

# Original $\mathbb{F}_1$ in emergent spacetime

Seyed Khaki\*

## Abstract

The existence of a quantum field theory over the “field with one element” was first addressed in 2012 by Bejleri and Marcolli, where it was shown that wonderful compactifications of the graph configuration spaces that appear in the calculation of Feynman integrals, as well as the moduli spaces of curves, admit a  $\mathbb{F}_1$  structure. Recently, we also examined some advantages of studying finite fields  $\mathbb{F}_q$ , wherein  $\mathbb{F}_1$  represents the fundamental string with the Planck length which plays a fundamental role in the model. Such a role was briefly described, by comparing the similarity between the collapse of the spacetime concept probing the scales below the Planck length and the mathematical collapse of the ‘field’ concept at  $q = 1$ . In this letter, we elaborate more on this role by explaining how Kapranov and Smirnov’s perspective based on the Iwasawa theory is a perfect mathematical fit for the string theory. Particularly, their work suggests that the existence of  $\mathbb{F}_1$  alone is sufficient to create other fields  $\mathbb{F}_q$  emergent as its extensions, which exactly reflects the postulate of the string theory (where various vibrational modes of the fundamental string manifest as other fields). As support, a couple of evidence are provided that illustrate the physical importance of the Weyl group (known to be a reductive group over  $\mathbb{F}_1$ ), and explain why the calculation of the amplitudes in the “amplitudes=combinatorial geometry” program (initiated by Arkani-Hamed et al. in 2013) exhibits a combinatorial nature and simplifies the exponentially growing ( $\mathcal{O}(4^n)$ ) calculations of the Feynmann diagrams to a polynomially growing order ( $\mathcal{O}(n^2)$ ) in the kinematic space of scattering data for the scalar theory with cubic interactions.

## 1 Introduction

Inspired by [1], the existence of a quantum field theory (QFT) over  $\mathbb{F}_1$ , the “field with one element”, was first studied in 2012 by Bejleri and Marcolli [2], where they explored the potential presence of an additional  $\mathbb{F}_1$  structure for particular classes of algebraic varieties which innately emerge within the perturbative QFT. In particular, they showed that wonderful compactifications of the graph configuration spaces, that appear in the calculation of Feynman integrals in position space, as well as the moduli spaces of curves, admit a  $\mathbb{F}_1$  structure. Unfortunately, such a bright idea was not adequately tracked. The second outstanding research avenue addressing a likely connection of  $\mathbb{F}_1$  with physics is the studies of endomotive, presented in [3] as a means of representing the formation of quantum statistical mechanical systems (QSMS) tied to number theory through arithmetic data, which began with the introduction of Bost-Connes (BC) system in [4]. Remarkably, Connes,

---

\*Academy of Fundamental Studies, Munich 80637, Germany  
Email: [s.khaki@afundas.com](mailto:s.khaki@afundas.com)

Consani, and Marcolli announced in [5] that the algebra and the endomotive of the QSMS of BC innately emerge by extension of scalars from  $\mathbb{F}_1$  to rational numbers. Further, in [6], a Galois field quantum mechanics whose  $q = 1$  limit becomes the classical mechanics is assessed.

On the other hand, by a different approach, we recently examined some advantages of studying finite fields  $\mathbb{F}_q$  in [7], where at  $q = 1$  limit, the string length  $\ell_s = q \times \ell_{Planck}$  becomes the Planck length representing the fundamental string, which, as its name suggests, plays a 'fundamental role' in the model. Such a role was briefly described by comparing the similarity between the collapse of the spacetime concept probing the scales below the Planck length and the mathematical collapse of the 'field' concept at  $q = 1$ . In this letter, we elaborate more on this role in the next part, by proposing that Kapranov and Smirnov's perspective [8] based on the Iwasawa theory is a promising mathematical fit for the string theory. Afterward, we present a brief tailored introduction to  $\mathbb{F}_1$  to ensure the necessary grasp of its 'simplification' and 'original' role. In the last section, we provide a couple of evidence that illustrates the physical importance of the Weyl group (known to be reductive group over  $\mathbb{F}_1$ ) and explain quantitatively why the calculation of the amplitudes in the "amplitudes=combinatorial geometry" program (initiated by Arkani-Hamed et al. in 2013 [9]) exhibits a combinatorial nature and simplifies the exponentially growing calculations of the Feynmann diagrams to a polynomially growing order in the kinematic space of the scattering data.

## 1.1 Proposal

Inspired by a strong analogy between number fields and curves over finite fields, in 1959, Iwasawa founded his  $Z_p$ -extensions theory [10], which investigates the growth of arithmetic objects in towers of number fields. In 1995, based on the Iwasawa theory that adjoining the roots of unity is analogous to producing extensions of a base field, Kapranov and Smirnov [8] formed the profound perspective that the set of the  $n$ -th roots can be introduced as  $\mathbb{F}_{1^n}$ , the field extensions of  $\mathbb{F}_1$ . In this manner,  $\mathbb{F}_q$  is an algebra of dimension  $\frac{q-1}{n}$  over  $\mathbb{F}_{1^n}$ , implying that all results of linear algebra over  $\mathbb{F}_q$  must be reproduced by replacing the equivalent extension of the field with one element  $\mathbb{F}_{1^{q-1}}$ . This suggests that only the existence of  $\mathbb{F}_1$  is sufficient to create other fields emergent as its extensions, which exactly reflects the postulate of the string theory where various vibrational modes of the fundamental string manifest as other fields/particles. Thus, *we propose that Kapranov and Smirnov's mathematical development is a perfect match for the string theory, where only one original field exists and its extensions manifest as other fields.*

## 1.2 Original field

In this part, we briefly introduce  $\mathbb{F}_1$  emphasizing its 'original' and 'simplification' role (for a general overview see [11]). Although based on the standard definition of the 'field',  $\mathbb{F}_2$  is the smallest field, and a field with only one element does not exist, mathematicians had an idea of the meaning of projective geometry over that field. One could recognize  $\mathbb{F}_1$  as a field-like algebraic object that behaves like a finite field with characteristic one if that field could exist. A more precise definition is that  $\mathbb{F}_1$  is the multiplicative monoid  $\{0,1\}$  without the additive structure which addresses the  $\mathbb{F}_1$  geometry as "non-additive geometry" (see [12] for an overview of several geometries over  $\mathbb{F}_1$ ). Studying geometry over  $\mathbb{F}_1$  indeed uncovers the minimum amount of information (i.e., the most

fundamental piece of data) about an object that enables examination of its geometric characteristics. This is also evident in the discovery of  $\mathbb{F}_1$  in 1956 which Tits uncovered in [13] (and Steinberg in [14]), where he observed that projective geometries over  $\mathbb{F}_q$  contain a meaningful analog at the minimal  $q = 1$  limit. As an instance, over  $\mathbb{F}_1$ , the symmetric group  $S_n$  on  $n$  elements behaves as if it were a linear group,

$$GL_n(\mathbb{F}_q) \xrightarrow{q \rightarrow 1} S_n \quad (1)$$

Here, the  $q = 1$  limit unveils the combinatorial skeleton/core of the Chevalley group  $GL_n(\mathbb{F}_q)$ . So, over  $\mathbb{F}_1$ , vector spaces (the objects on which  $GL_n(\mathbb{F}_q)$  acts) reduce/descend to pointed sets (the objects on which  $S_n$  acts) which one may imagine as **origin**. In this sense,  $\mathbb{F}_1$  is the 'original' base (i.e., a terminal object lying below the integers) for an "absolute point"  $\text{Spec } \mathbb{F}_1$  over which any algebraic scheme would sit. Such an original role in Tits' perspective includes many schemes of finite type over the integers that can be characterized by a polynomial with integer coefficients, serving as a counting function that counts  $\mathbb{F}_q$  points of the scheme for all prime power values of  $q$ . Instances contain affine spaces, projective spaces, and Grassmannians.

$$\#\mathbb{A}^n(\mathbb{F}_q) = q^n \quad \#\mathbb{P}^{n-1}(\mathbb{F}_q) = [n]_q \quad \#Gr(k, n)(\mathbb{F}_q) = \begin{bmatrix} n \\ k \end{bmatrix}_q \quad (2)$$

where  $[n]_q = q^{n-1} + \dots + q + 1$  is the Gauss number,  $[n]_q! = \prod_{i=1}^n [i]_q$  is the Gauss factorial, and  $\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[n]_q!}{[k]_q! [n-k]_q!}$  is the Gauss binomial. At  $q = 1$ , the number of points will be minimized to  $\mathbb{F}_1$ -rational points

$$\#\mathbb{A}^n(\mathbb{F}_1) = 1 \quad \#\mathbb{P}^{n-1}(\mathbb{F}_1) = n \quad \#Gr(k, n)(\mathbb{F}_1) = \binom{n}{k} \quad (3)$$

Now, let us view an example directly sourced from [11] to comprehend how  $\mathbb{F}_1$  simplifies/reduces the information to its minimal meaningful level.

**Example** We illustrate the ideas of Tits in the example of  $GL_3(F_q)$ . In this case, we consider the two Grassmannians  $Gr(1, F_q^3)$  and  $Gr(2, F_q^3)$ , whose points corresponds to the points and lines in  $P^2(F_q)$ , respectively. The incidence relation consists of pairs of a point  $P$  and a line  $L$  such that  $P \in L$ . In the case  $q = 2$ , we calculate  $\#Gr(1, F_2^3) = \begin{bmatrix} 3 \\ 1 \end{bmatrix}_2 = 7$  and  $\#Gr(2, F_2^3) = \begin{bmatrix} 3 \\ 2 \end{bmatrix}_2 = 7$ . Moreover, note that every line contains  $q + 1 = 3$  points and that every point is contained in  $q + 1 = 3$  lines. The corresponding geometry is illustrated on the left hand side of Figure 1, where the dots correspond to the points in  $P^2(F_q)$ , the circles correspond to the lines in  $P^2(F_q)$ , and an edge between a dot and a circle indicates that the corresponding point is contained in the corresponding line. Considering the limit  $q \rightarrow 1$  yields the geometry for  $S_3$  that consists of the sets  $\Sigma(1, 3) = \{\{1\}, \{2\}, \{3\}\}$  and  $\Sigma(2, 3) = \{\{1, 2\}, \{1, 3\}, \{2, 3\}\}$ . Note that every point of this geometry, i.e., an one element subset of  $\{1, 2, 3\}$ , is contained in  $q + 1 = 2$  lines, i.e., a 2-subset of  $\{1, 2, 3\}$ . Similarly, every line contains  $q + 1 = 2$  points. The corresponding geometry is depicted on the right hand side of Figure 1.




Figure 1: The geometry of  $GL_3(\mathbb{F}_q)$  for  $q = 2$  and its  $q = 1$  limit

## 2 Supports and implications

### 2.1 Weyl group

In his bold paper [13], Tits extends the analogy between  $S_n$  and  $GL_n(\mathbb{F}_q)$  to an analogy between the  $\mathbb{F}_q$ -rational points of a Chevalley group scheme  $G$  and its Weyl group  $W$  such that  $W = G(\mathbb{F}_1)$ . That is,  $\mathbb{F}_1$  unveils the Weyl group which is the core or essence of the main group. He indicates that, in the  $q = 1$  limit, the finite geometry attached to  $G(\mathbb{F}_q)$  will be the geometry of the Coxeter group  $W$ . In this way,  $\lim_{q \rightarrow 1} |G(\mathbb{F}_q)| / (q-1)^{n_w}$  should count the number of elements of the Weyl group (where  $n_w$  zeros of  $G(\mathbb{F}_q)$  at  $q = 1$  are removed). E.g., for the exceptional Chevalley group  $E_8(\mathbb{F}_q)$  we have  $|E_8(\mathbb{F}_q)| = q^{120}(q^{30} - 1)(q^{24} - 1)(q^{20} - 1)(q^{18} - 1)(q^{14} - 1)(q^{12} - 1)(q^8 - 1)(q^2 - 1)$ , and therefore,  $|W(E_8(\mathbb{F}_q))| = \lim_{q \rightarrow 1} \frac{|E_8(\mathbb{F}_q)|}{(q-1)^{20}} = 30 \times 24 \times 20 \times 18 \times 14 \times 12 \times 8 \times 2 = 696\,729\,600$ . This is the order we considered in [7] to calculate the energy scale corresponding to the  $E_8$ , which here reveals its interpretation over  $\mathbb{F}_1$ . Over  $\mathbb{F}_2$ , the order blow up to  $|E_8(\mathbb{F}_2)| \approx 3.3 \times 10^{74}$ . Hence,

$$|E_8(\mathbb{F}_2)| \approx 3.3 \times 10^{74} \xrightarrow{q \rightarrow 1} |E_8(\mathbb{F}_1)| \approx 6.9 \times 10^8 \quad (4)$$

Now, let us cite the physical importance of the Weyl group. As established in [15] (and followed in [16]), for a particle symmetry described by a Lie group  $G$ , not only the Weyl group  $W(G)$  canonically acts in all zero weight spaces of  $G$  (and therefore on observables), but also its action gives various physical relations among which are those inferred from the  $G$ -transformation properties of observables. Notably, it is shown that in the case of finite-dimensional representations, there exists a canonical action of the Weyl groups on observables such that many physical relations are a consequence of the Weyl group action rather than the action of the original group of root systems on observables. Thus, *we claim that such physical importance originates from the fact that the Weyl group is a reductive group over  $\mathbb{F}_1$ .*

### 2.2 "Amplitudes=Combinatorial Geometry" (ACG) program

In 2013, Arkani-Hamed et al. started the ACG program, first by introducing the "amplituhedron" in [9], based merely on elementary combinatorial and geometric structures in the kinematic space of the scattering data that 'simplifies' the conventional calculation of the amplitudes to simpler counting problems without referencing to bulk spacetimes and Hilbert space. In the simplest theory of colored scalar particles - the  $\text{Tr}\phi^3$  theory - considered in the ACG program, the  $n$ -point tree

amplitude is typically a sum over  $C_{n-2} = \frac{1}{n-1} \binom{2n-4}{n-2}$  number of Feynman diagrams (where  $C_n$  is the  $n^{\text{th}}$  Catalan number) which is of the size  $\mathcal{O}(4^n)$  (cf. [17] Outlook section, and [18] Eq. 1.1). On the other hand, their approach contains curve integrals of an action built from  $\mathcal{O}(n^2)$  piecewise linear functions, originating from the kinematic space for  $n$  massless momenta whose dimensionality is (cf. [19] Eq. 2.3)

$$\binom{n}{2} - n = \frac{n(n-3)}{2} \quad (5)$$

where  $\binom{n}{2}$  is the number of Lorentz invariant dot products of momenta  $p_i \cdot p_j$  and  $-n$  comes from the momentum conservation constraints. In this context, therefore, the 'simplification' means reducing the order  $\mathcal{O}(4^n)$  of the Feynman diagrams method which grows exponentially with the number of particles to the order  $\mathcal{O}(n^2)$  of the ACG method which grows as a polynomial with the number of particles (cf. [17] Outlook section). *We claim that such a simplification is similar to the familiar reduction mentioned above in Eq. (1), Figure 1, and Eq. (4). That is,*

$$\text{Feynman method: amplitudes } (\mathbb{F}_2) \xrightarrow{q \rightarrow 1} \text{ACG method: amplitudes } (\mathbb{F}_1)$$

Now, let us quantitatively assess this claim. Over  $\mathbb{F}_q$ , the dimensionality of kinematic space is the  $q$ -analog of Eq. (5) (for  $q$ -analogs compare Eq. (2) and (3))

$$\left[ \begin{matrix} n \\ 2 \end{matrix} \right]_q - [n]_q = \frac{[n]_q \times [n-1]_q}{[2]_q} - [n]_q \quad (6)$$

This yields  $\frac{(q^n-1)/(q-1) \times (q^{n-1}-1)/(q-1)}{1+q} - (q^n-1)/(q-1)$  which is  $\mathcal{O}(q^{2n})$ . Therefore, at  $q=2$ , it grows as  $\mathcal{O}(4^n)$ , and at  $q=1$ , it becomes Eq. (5) which is  $\mathcal{O}(n^2)$ . Notably, while the exponential behavior shows up for all  $q \geq 2$ , the polynomial behavior emerges only at  $q=1$ <sup>1</sup>.

Furthermore, it is worth mentioning that in [20], Arkani-Hamed et al. recently highlighted the unified stringy picture too as they generalized their results to the stringy  $\text{Tr}\phi^3$  amplitudes and stated "There is a unique shift of the kinematic data that preserves the zeros, and this shift is precisely the one that unifies colored scalars, pions, and gluons into a *single object*". They also remark "there is a beautiful reason for the universality of these zeros, which turn out to be deeply related to a surprising relation between these colored theories revealed upon understanding a unified "stringy" descriptions of all these amplitudes. These stringy generalizations inherit all the zeros and factorization patterns of the field theory amplitudes and in fact generalize them to infinite new families of zero/factorization patterns. They also allow us to see that amplitudes for colored scalars, pions, and gluons are all given by a single function, expanded about different points in the kinematic space."

In sum, this letter concludes that not only the quantum gravity in the emergent spacetime should enjoy  $\mathbb{F}_1$  and its geometry, but also a fundamental theory of nature should embody finite combinatorial properties emerging from the number theory.

---

<sup>1</sup>This may have some implications in computer science

## References

- [1] Oliver Schnetz. Quantum field theory over  $\mathbb{F}_q$ . *arXiv preprint arXiv:0909.0905*, 2009.
- [2] Dori Bejleri and Matilde Marcolli. Quantum field theory over  $\mathbb{F}_1$ . *Journal of Geometry and Physics*, 69:40–59, 2013.
- [3] Alain Connes, Caterina Consani, and Matilde Marcolli. Noncommutative geometry and motives: The thermodynamics of endomotives. *Advances in Mathematics*, 214(2):761–831, 2007.
- [4] Jean-Benoit Bost and Alain Connes. Hecke algebras, type III factors and phase transitions with spontaneous symmetry breaking in number theory. *Selecta Mathematica*, 1:411–457, 1995.
- [5] Alain Connes, Caterina Consani, and Matilde Marcolli. Fun with  $\mathbb{F}_1$ . *Journal of Number Theory*, 129(6):1532–1561, 2009.
- [6] Lay Nam Chang, Zachary Lewis, Djordje Minic, and Tatsu Takeuchi. Quantum  $\mathbb{F}_{un}$ : the  $q=1$  limit of galois field quantum mechanics, projective geometry and the field with one element. *Journal of Physics A: Mathematical and Theoretical*, 47(40):405304, 2014.
- [7] Seyed Khaki. An examination of the de sitter space dual to the monster conformal field theory. *arXiv preprint arXiv:2309.09762*, 2023.
- [8] M. Kapranov and A. Smirnov. Cohomology determinants and reciprocity laws: number field case. *Preprint Series 1912, Inst. für Experimentelle Mathematik, Essen*, 1995. Available online.
- [9] Nima Arkani-Hamed and Jaroslav Trnka. The amplituhedron. *Journal of High Energy Physics*, 2014(10):1–33, 2014.
- [10] Kenkichi Iwasawa. On  $\Gamma$ -extensions of algebraic number fields. *Bull. Am. Math. Soc.*, 65:183–226, 1959.
- [11] Oliver Lorscheid.  $\mathbb{F}_1$  for everyone. *Jahresbericht der Deutschen Mathematiker-Vereinigung*, 120(2):83–116, 2018.
- [12] J López Pena and Oliver Lorscheid. Mapping  $\mathbb{F}_1$ -land: An overview of geometries over the field with one element. *Noncommutative geometry, arithmetic, and related topics*, pages 241–265, 2011.
- [13] Jacques Tits. Sur les analogues algébriques des groupes semi-simples complexes. In *Colloque d’algèbre supérieure, tenue à Bruxelles du*, volume 19, pages 261–289, 1956.
- [14] Robert Steinberg. A geometric approach to the representations of the full linear group over a

- galois field. *Transactions of the American Mathematical Society*, 71(2):274–282, 1951.
- [15] Grzegorz Cieciura and Igor Szczyrba. Physical relations and the Weyl group. *Journal of mathematical physics*, 28(9):2115–2127, 1987.
- [16] Grzegorz Cieciura and Igor Szczyrba. Weyl group and tensor operators for hadron-type  $SU(n)$  representations. *Journal of mathematical physics*, 28(9):2128–2141, 1987.
- [17] N Arkani-Hamed, H Frost, G Salvatori, PG Plamondon, and H Thomas. All loop scattering as a counting problem. *arXiv preprint arXiv:2309.15913*, 2023.
- [18] Nima Arkani-Hamed, Hadleigh Frost, Giulio Salvatori, Pierre-Guy Plamondon, and Hugh Thomas. All loop scattering for all multiplicity. *arXiv preprint arXiv:2311.09284*, 2023.
- [19] Nima Arkani-Hamed, Yuntao Bai, Song He, and Gongwang Yan. Scattering forms and the positive geometry of kinematics, color and the worldsheet. *Journal of High Energy Physics*, 2018(5):1–78, 2018.
- [20] Nima Arkani-Hamed, Qu Cao, Jin Dong, Carolina Figueiredo, and Song He. Hidden zeros for particle/string amplitudes and the unity of colored scalars, pions and gluons. *arXiv preprint arXiv:2312.16282*, 2023.