LONG CYCLES IN LINEAR THRESHOLDING SYSTEMS

ANNA LADDACH AND MICHAEL SHAPIRO

INTRODUCTION

Linear thresholding systems have been studied as models of neural activation [1] and more recently as models of cell-intrinsic gene regulation [2]. More generally, they represent dynamics on a set of N objects which can be either on or off. That is, they are maps

$$f = f_{(J,\theta)} : \mathcal{P}_N = \{0,1\}^N \subset \mathbb{R}^N \to \{0,1\}^N$$
$$f(x) = \theta(Jx)$$

where J is a linear map and θ is a thresholding function

$$\theta(x_1, \dots, x_N) = (y_1, \dots, y_N)$$
$$y_i = \begin{cases} 0 & \text{if } x_i \le \theta_i \\ 1 & \text{if } x_i > \theta_i \end{cases}$$

We will restrict to the case in which each θ_i is positive. Since there are 2^N elements of \mathcal{P}_N , this provides an upper bound on the length of any cycle for a linear thresholding system in dimension N. Under the assumption that the θ_i are positive, 0 is a fixed point and this reduces this bound to $2^N - 1$. Here we exhibit systems of arbitrarily large N with cycles of length greater than $e^{\sqrt{N}}$. Along the way, we introduce a direct product on linear thresholding systems.

BACKGROUND AND DEFINITIONS

We record here some basic definitions and results. **Definition 1.**

- We say two linear thresholding systems (J,θ) and (J',θ') are equivalent and write $(J,\theta) \sim (J',\theta')$ if $f_{(J,\theta)} = f_{(J',\theta')}$.
- We say that (J,θ) is generic if for each $x \in \mathcal{P}$, and each coordinate i, $(Jx)_i \neq \theta_i$.
- We will say that (J, θ) is robust if there is an open set U_J around J and an open set U_{θ} around θ such that for each $(J', \theta') \in U_J \times U_{\theta}, (J, \theta) \sim (J', \theta')$

The following are not hard to see: **Proposition 1.**

- (J, θ) is robust if and only if it is generic.
- Every linear thresholding system is equivalent to a generic system.
- Every linear thresholding system is equivalent under a linear change of coordinates to a system in which the θ_i are independent of *i*. Further, their value can be taken to be any positive number.

Date: 22 November 2023.

We will take our thresholding value to be $\theta = \frac{1}{2}$. We will be interested in cases where J permutes \mathcal{P} , and here thresholding acts as the identity.

The dynamics of a function f on a finite set X can be seen as a directed graph $\Gamma(X, E)$ where the edges E are the pairs $\{(x, f(x)) \mid x \in X\}$. Each connected component of Γ consists of a recurrent cycle C (possibly of length 1) possibly decorated with directed trees T_i . Each T_i is directed towards its root, which is a vertex of C. The vertices on the cycles are *recurrent*, the others are *transient*.

LONG CYCLES

We start by defining a direct product on linear thresholding systems.

Definition 2. Given two systems, (J, θ) and (J', θ) where the first has N genes and the second has N' genes, the product system, $(J, \theta) \oplus (J', \theta)$ is the system (\bar{J}, θ) on N + N' genes where

$$ar{J} = \begin{bmatrix} J & 0 \\ 0 & J' \end{bmatrix}.$$

Proposition 2. The dynamics of the product system are the product of the dynamics of the two systems. That is, let f and f' be the functions of the two systems and \overline{f} be the function of the product system. Let x and x' be elements of $\mathcal{P}(1, \ldots, N)$ and $\mathcal{P}(1, \ldots, N')$. Then

$$\bar{f}\left(\begin{bmatrix} x\\x'\end{bmatrix}\right) = \begin{bmatrix} f(x)\\f'(x')\end{bmatrix}$$

Corollary 1. Suppose x_1, \ldots, x_r is a cycle of (J, θ) of length r and x'_1, \ldots, x'_s is a cycle of (J', θ) of length s. Then together, they form a cycle of length $\operatorname{lcm}(r, s)$ in the product system.

We now take a moment to consider the orbits of (J_p, θ) where $J = J_p$ is the matrix that cyclically permutes the bases of \mathbb{R}^p , p is prime and $\theta = \frac{1}{2}$. Observe that in this case the thresholding operation is superfluous since it acts as the identity on $J\mathcal{P}$. Note that J generates the cyclic group \mathbb{Z}_p and thereby induces an action of \mathbb{Z}_p on \mathcal{P} .

Lemma 1.

- (J, θ) has no transient states.
- The number of coordinates which are one is constant on each orbit.
- Each cycle of (J, θ) has length 1 or p.
- There are two cycles of length 1.
- There are $(2^p 2)/p$ cycles of length p.

Proof.

- Since $J^p = I$, $J^p x = x$ for each $x \in \mathcal{P}$.
- Each $x \in \mathcal{P}$ is a set coordinates which are one. J acts to cyclically permute this set. It follows that the number of coordinates which are one is constant along the orbit.
- The length of each orbit divides *p*. Since *p* is prime, it follows that this is necessarily 1 or *p*.
- The points (0,...,0) and (1,...,1) are fixed by J. On the other hand, if x is neither of these, x has two (cyclically) adjacent coordinates which are not equal. Hence Jx ≠ x, so the orbit of x has length p.

 $\mathbf{2}$

• This follows directly from the previous statement.

Corollary 2. Let p_1, \ldots, p_k be a sequence of distinct primes, $N = \sum p_i$, $P = \prod p_i$. Then there is a system (J, θ) on N genes with a cycle of length P.

Note that the sum of the first *n* primes grows as $\frac{1}{2}n^2 \log(n)$ [3] while the product of the first *n* primes grows as $e^{(1+o(1))n \log(n)}$ [4]. Thus, for arbitrarily large values *N* there are systems on *N* genes with cycles of length greater than $e^{\sqrt{N}}$.

In fact, (J, θ) is rich in long cycles. If $S \subset \{1, \ldots, k\}$, then there are

$$\prod_{i \in S} (2^{p_i} - 2)/p_i$$

orbits of length $\prod_{i \in S} p_i$.

Our definition is restricted to the case where the two systems have the same thresholding function θ . However up to equivalence, they can be given the same θ and the equivalence class of the product system depends only on the equivalence class of the factors. Thus, up to equivalence, the product is defined on arbitrary pairs of systems.

References

- A C C Coolen, P Sollich, and R Kuehn. Theory of neural information processing systems. Oxford University Press, 2005.
- [2] Ryan Hannam, Alessia Annibale, and Reimer Kühn. Cell reprogramming modelled as transitions in a hierarchy of cell cycles. *Journal of Physics A Mathematical General*, 50(42):425601, October 2017.
- [3] E. Landau. Handbuch der Lehre von der Verteilung der Primzahlen, volume 1. Teubner, Leipzig und Berlin, 1909.
- [4] Sebastian Martin Ruiz. A result on prime numbers. Math. Gaz., July 1997.

FRANCIS CRICK INSTITUTE, LONDON

FRANCIS CRICK INSTITUTE, LONDON