

One Dimensional Fixed Point Interactions and the Resolvent Algebra - Simple Remarks

Antonio Moscato*

Dipartimento di Matematica, Università di Trento
Via Sommarive 14, I-38123 Povo (Trento), Italy

Abstract

In this paper, the resolvent algebra $\mathcal{R}(\mathbb{R}^2, \sigma)$ stability under dynamics induced by the symbolic Hamiltonians

- $H = -\frac{d^2}{dx^2} + \alpha\delta(x - x_0)$, $\alpha \in \mathbb{R} \setminus \{0\}$, $x_0 \in \mathbb{R}$,
- $H = -\frac{d^2}{dx^2} + \sum_{i=1}^N \alpha_i \delta(x - x_i)$, $\alpha_i \in \mathbb{R} \setminus \{0\}$, $x_i \in \mathbb{R} : x_i \neq x_j$, $i, j \in \{1, \dots, N\}$,
- $H = -\frac{d^2}{dx^2} + \sum_{i=1}^{\infty} \alpha_i \delta(x - x_i)$, $\{\alpha_i\}_i \in l^1(\mathbb{N})$, $x_i \in \mathbb{R} : x_i \neq x_j$, $i, j \in \mathbb{N}$

is proved: if π_S is the Schrödinger representation of $\mathcal{R}(\mathbb{R}^2, \sigma)$ on $L^2(\mathbb{R})$, $[e^{itH} \pi_S(a) e^{-itH}] \in \pi_S[\mathcal{R}(\mathbb{R}^2, \sigma)]$ holds for all $a \in \mathcal{R}(\mathbb{R}^2, \sigma)$ and $(H - i\lambda\mathbb{1})^{-1} \in \pi_S[\mathcal{R}(\mathbb{R}^2, \sigma)]$, for all $\lambda \in \mathbb{R} \setminus \{0\}$. Results from [1], [7] and [9] are used for the purpose.

1 Introduction

In 2008, Buchholz and Grundling introduced a novel C^* -algebra, the resolvent algebra ([1]), to model (bosonic) quantum mechanical systems, aiming to overcome some of the drawbacks afflicting the historically famous Weyl algebra, typically employed for the purpose. In details, given a symplectic vector space (X, σ) , the dynamics of the Weyl algebra $CCR(X, \sigma)$ defined by symplectic transformations of X correspond to the dynamics induced by quadratic Hamiltonians, hence excluding many physically interesting situations; as a matter of fact, the following result holds.

Proposition ([2]): Let $(H_\lambda = H_0 + \lambda V, \mathcal{D}_{H_\lambda})$ be a self-adjoint operator on $L^2(\mathbb{R})$, where H_0 is the free Hamiltonian. If $V \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$, $\forall \lambda \in \mathbb{R} \setminus \{0\}, t \in \mathbb{R}$, the automorphism $\alpha_t^\lambda(\cdot) = e^{itH_\lambda}(\cdot)e^{-itH_\lambda}$ of $\mathfrak{B}(L^2(\mathbb{R}))$ is not an automorphism of $CCR(\mathbb{R}^2, \sigma)$ unless V is null. ■

Moreover, in regular representations, natural observables as bounded functions of the Hamiltonian do not belong to $CCR(X, \sigma)$. Resolvent algebras $\mathcal{R}(X, \sigma)$, on the other hand, have proved to be not affected by similar drawbacks in different interesting situations ([3], [4]); a crucial role for this to happen is its non-simple nature ([1], [5]).

*antonio.moscato@unitn.it

This paper takes a step forward in this direction: it shows that, for the case of a single, non-relativistic spinless particle, the resolvent algebra $\mathcal{R}(\mathbb{R}^2, \sigma)$, on the one hand, can accommodate dynamics induced by fixed located Dirac delta-like potentials, on the other hand, the corresponding Hamiltonians are affiliated to it, respectively meaning that, if H is the whatever point-like Hamiltonian of the case, $e^{itH} \pi_S(a) e^{-itH} \in \pi_S[\mathcal{R}(\mathbb{R}^2, \sigma)]$, for all $a \in \mathcal{R}(\mathbb{R}^2, \sigma)$ and $(H - i\lambda \mathbb{1})^{-1} \in \pi_S[\mathcal{R}(\mathbb{R}^2, \sigma)]$, for all $\lambda \in \mathbb{R} \setminus \{0\}$, where π_S is the Schrödinger representation of $\mathcal{R}(\mathbb{R}^2, \sigma)$ on $L^2(\mathbb{R})$.

The reason to focus on delta interactions lies in their ubiquity in applications: they represent good candidates to approximate unknown, short-scaled interactions. An extensive mathematical literature has been dedicated to such singular potential: [6], for example, provides a very well-known illustrative reference.

The paper is articulated as follows: section 2 recaps the necessary structural results about resolvent algebras, while section 3 first briefly recalls the content of [7], then proves what above anticipated, with H alternatively given by the symbols

- $H = -\frac{d^2}{dx^2} + \alpha \delta(x - x_0)$, $\alpha \in \mathbb{R} \setminus \{0\}$, $x_0 \in \mathbb{R}$,
- $H = -\frac{d^2}{dx^2} + \sum_{i=1}^N \alpha_i \delta(x - x_i)$, $\alpha_i \in \mathbb{R} \setminus \{0\}$, $x_i \in \mathbb{R} : x_i \neq x_j$, $i, j \in \{1, \dots, N\}$,
- $H = -\frac{d^2}{dx^2} + \sum_{i=1}^{\infty} \alpha_i \delta(x - x_i)$, $\{\alpha_i\}_i \in l^1(\mathbb{N})$, $x_i \in \mathbb{R} : x_i \neq x_j$, $i, j \in \mathbb{N}$.

2 The Resolvent Algebra

Definition 2.1. Let (X, σ) be a symplectic space and $\mathfrak{R} \doteq \{R(\lambda, f) \mid \lambda \in \mathbb{R} \setminus \{0\}, f \in X\}$ a set of symbols. Let L be the following list of relations.

1. $R(\lambda, 0) = -\frac{i}{\lambda} \mathbb{1}$
2. $R(\lambda, f)^* = R(-\lambda, f)$
3. $\nu R(\nu\lambda, \nu f) = R(\lambda, f)$
4. $R(\lambda, f) - R(\mu, f) = i(\mu - \lambda) R(\lambda, f) R(\mu, f)$
5. $[R(\lambda, f), R(\mu, g)] = i\sigma(f, g) R(\lambda, f) R(\mu, g)^2 R(\lambda, f)$
6. $R(\lambda, f) R(\mu, g) = R(\lambda + \mu, f + g) \left[R(\lambda, f) + R(\mu, g) + i\sigma(f, g) R(\lambda, f)^2 R(\mu, g) \right]$

with $\lambda, \mu, \nu \in \mathbb{R} \setminus \{0\}$, $f, g \in X$ and $\lambda + \mu \neq 0$ at 6.. Denoted by $\tilde{\mathcal{R}}_0$ the unital $*$ -algebra freely generated by \mathfrak{R} and the two-sided ideal \mathcal{I}_L generated by the given relations, the **unital pre-resolvent $*$ -algebra** $\mathcal{R}_0 \doteq \tilde{\mathcal{R}}_0 / \mathcal{I}_L$ is defined. \square

Remark 2.1. Turning \mathcal{R}_0 into a C^* -algebra amounts in defining a C^* -norm on it. The following result is preparatory. \square

Proposition 2.1. Let (X, σ) be a symplectic space and let \mathcal{R}_0 be as in definition 2.1.

1. If \mathcal{H} is a Hilbert space and $\pi_0 : \mathcal{R}_0 \rightarrow \mathcal{B}(\mathcal{H})$ is a (bounded) $*$ -representation of \mathcal{R}_0 , then $\left\| \pi_0 [R(\lambda, f)] \right\| \leq |\lambda|^{-1}$. In other words, for all $a \in \mathcal{R}_0$, there exists $c_a \geq 0$, only depending on a , such that $\|\pi(a)\| \leq c_a$, for all the bounded Hilbert space representations π of \mathcal{R}_0 .
2. Let ω be a positive functional of \mathcal{R}_0 , i.e. $\omega : \mathcal{R}_0 \rightarrow \mathbb{C}$ such that $\omega(a^*a) \geq 0$ for all $a \in \mathcal{R}_0$. Then, the GNS-construction yields a cyclic $*$ -representation of \mathcal{R}_0 , denoted by π_ω , consisting of bounded Hilbert space operators. ■

Remark 2.2. The importance of the foregoing result lies in the following fact: a $*$ -algebra can have unbounded representations, but, as long as the GNS-construction is concerned, the corresponding representation is bounded. □

Definition 2.2. Let \mathcal{R}_0 over (X, σ) be as in Definition 2.1. Its universal representation $\pi_u : \mathcal{R}_0 \rightarrow \mathcal{B}(\mathcal{H})$ is given by

$$\pi_u(a) \doteq \bigoplus \{ \pi_\omega(a) \mid \omega \in \mathfrak{S} \} \quad \text{and} \quad \|a\|_u \doteq \|\pi_u(a)\| \equiv \sup_{\omega \in \mathfrak{S}} \|\pi_\omega(a)\|$$

denotes the corresponding enveloping C^* -norm, where \mathfrak{S} is the set of all positive and normalized¹ (algebraic) states over \mathcal{R}_0 . The completion of $\mathcal{R}_0 / \ker \|\cdot\|_u$ with respect to $\|\cdot\|_u$ is denoted by $\mathcal{R}(X, \sigma)$ and defines the **resolvent algebra** over (X, σ) . □

Remark 2.3. Because of Proposition 2.1(1), $\|\cdot\|_u$ is well-defined, because of $\|\pi_\omega(a)\| \leq c_a$, for all $\omega \in \mathfrak{S}$, hence

$$\sup_{\omega \in \mathfrak{S}} \|\pi_\omega(a)\| \leq c_a < \infty.$$

□

Remark 2.4. It is reported that, given $f \in X$, the map

$$R(\cdot, f) : \lambda \in \mathbb{R} \setminus \{0\} \mapsto R(\lambda, f) \in \mathcal{R}(X, \sigma)$$

is analytic. Such a circumstance allows for the possibility of having complex-valued λ arguments for $R(\lambda, f)$ as long as these are not purely imaginary. Differently said, $\mathcal{R}(X, \sigma)$ contains $R(z, f)$, with $z \in \mathbb{C} \setminus i\mathbb{R}$ too. □

Definition 2.3. Let \mathcal{H} be a Hilbert space and (X, σ) be a symplectic vector space. Given $S \subseteq X$, a representation $\pi : \mathcal{R}(X, \sigma) \rightarrow \mathcal{B}(\mathcal{H})$ is said **regular on S** if and only if $\ker \left\{ \pi [R(1, f)] \right\} = \{0\}$ for all $f \in S$. Consequently, a state ω of $\mathcal{R}(X, \sigma)$ is said **regular on S** if and only if its GNS-representation is regular on S . A representation (state) is simply said **regular** if it is regular on X . The set of all regular representations of $\mathcal{R}(X, \sigma)$ on \mathcal{H} is denoted by $\text{reg}(\mathcal{R}(X, \sigma), \mathcal{H})$, while the set of all regular states by $\mathfrak{S}_r(\mathcal{R}(X, \sigma))$. □

Remark 2.5. The importance of the notion of regular representation stems from the fact that, given the Hilbert space \mathcal{H} together with the regular representation $\pi : \mathcal{R}(X, \sigma) \rightarrow \mathcal{B}(\mathcal{H})$, for all $f \in X$, one can define the linear operator $(\phi_\pi(f), \mathcal{D}_{\phi_\pi(f)})$, where

$$\begin{aligned} \mathcal{D}_{\phi_\pi}(f) &= \text{Ran} [R(1, f)] \\ \phi_\pi(f) &= R(1, f)^{-1} + i\mathbb{1}. \end{aligned}$$

Such an operator is proved to be self-adjoint and its definition is independent on the choice of $\lambda \equiv 1$. □

¹i.e. $\omega(\mathbb{1}) = 1$

Proposition 2.2. *Let (X, σ) be a symplectic vector space and $\mathcal{R}(X, \sigma)$ be the corresponding resolvent algebra. Every regular representation of $\mathcal{R}(X, \sigma)$ is faithful. ■*

Proposition 2.3. *Let \mathcal{H} be a Hilbert space and (X, σ) be a symplectic vector space. Given a regular representation $\pi \in \text{reg}(\mathcal{R}(X, \sigma), \mathcal{H})$, the mapping*

$$\tilde{\pi} : \delta_f \in CCR(X, \sigma) \longmapsto \tilde{\pi}(\delta_f) \doteq e^{i\phi_\pi(f)} \in \mathcal{B}(\mathcal{H})$$

results in a regular representation of the Weyl algebra over (X, σ) ; particularly,

$$\pi \in \text{reg}(\mathcal{R}(X, \sigma), \mathcal{H}) \longmapsto \tilde{\pi} \in \text{reg}(CCR(X, \sigma), \mathcal{H})$$

is a bijection preserving irreducibility and direct sums; its inverse is defined by

$$\pi[R(\lambda, f)] \doteq -i \int_0^{\eta\infty} e^{-\lambda t} \tilde{\pi}(\delta_{-tf}) dt, \quad \eta \equiv \text{sign}(\lambda)$$

where the integral is understood in the strong operator topology. ■

Remark 2.6. *The foregoing result allows to state that, as long as (X, σ) is finite-dimensional, the Stone-von Neumann theorem holds for the resolvent algebra too, i.e. $\mathcal{R}(X, \sigma)$ admits a unique (up to unitary equivalence) irreducible, regular representation, the Schrödinger representation. □*

Proposition 2.4. *Let (X, σ) be a finite-dimensional symplectic space and $\pi_0 : \mathcal{R}(X, \sigma) \longrightarrow \mathcal{B}(\mathcal{H}_0)$ be an irreducible regular representation of $\mathcal{R}(X, \sigma)$. Then, there exists a unique closed two-sided ideal \mathcal{K} of $\mathcal{R}(X, \sigma)$ isomorphic to the algebra of compact operators $\mathcal{B}_\infty(\mathcal{H}_0)$, such that $\pi_0(\mathcal{K}) = \mathcal{B}_\infty(\mathcal{H}_0) \subset \mathcal{B}(\mathcal{H}_0)$. ■*

2.1 Dynamics on $\mathcal{R}(X, \sigma)$

Definition 2.4. Let $(X, \sigma) \equiv (\mathbb{R}^{2N}, \sigma)$, $N \in \mathbb{N}$ be the standard symplectic vector space and $\pi_S : \mathcal{R}(X, \sigma) \longrightarrow \mathcal{B}(L^2(\mathbb{R}^N))$ be the Schrödinger representation of $\mathcal{R}(\mathbb{R}^{2N}, \sigma)$. A (H, \mathcal{D}_H) self-adjoint Hamiltonian on $L^2(\mathbb{R}^N)$ **induces a dynamics on $\mathcal{R}(X, \sigma)$** if and only if

$$e^{itH} \left\{ \pi_S[\mathcal{R}(X, \sigma)] \right\} e^{-itH} \subseteq \pi_S[\mathcal{R}(X, \sigma)], \quad \forall t \in \mathbb{R}.$$

□

Remark 2.7. *Such a definition is well-posed, because of the regularity, hence the injectivity, of the Schrödinger representation, together with the fact that, because of the Stone-von Neumann theorem, such a representation is unique up to unitary equivalence. □*

Proposition 2.5. *Let $\mathcal{R}(\mathbb{R}^2, \sigma)$ be and $\pi_S : \mathcal{R}(\mathbb{R}^2, \sigma) \longrightarrow \mathcal{B}(L^2(\mathbb{R}))$ be the corresponding Schrödinger representation. Given $V \in C_0(\mathbb{R})$, the self-adjoint Hamiltonian $(H \equiv H_0 + V, \mathcal{D}_H)$ induces a dynamics on $\mathcal{R}(\mathbb{R}^2, \sigma)$. Moreover $R_H(\lambda) = (H - i\lambda\mathbf{1})^{-1} \in \pi_S[\mathcal{R}(\mathbb{R}^2, \sigma)]$ for all $\lambda \in \mathbb{R} \setminus \{0\}$, i.e. H is affiliated to $\mathcal{R}(\mathbb{R}^2, \sigma)$. ■*

3 Point Interactions

Proposition 2.5 guarantees that, for one-dimensional quantum mechanical systems, Schrödinger Hamiltonians with $C_0(\mathbb{R})$ potentials do induce dynamics on $\mathcal{R}(\mathbb{R}^2, \sigma)$. This section, on the other hand, addresses the stability problem of $\mathcal{R}(\mathbb{R}^2, \sigma)$ under the action of symbolic Hamiltonians² as

$$H = -\frac{d^2}{dx^2} + \sum_{i=1}^N \alpha_i \delta(x - x_i), \quad (1)$$

with $N \in \mathbb{N} \cup \{\infty\}$, $x_i \in \mathbb{R} : x_i \neq x_j, \forall i, j$, $\alpha_i \in \mathbb{R} \setminus \{0\}, \forall i$, clearly not of Schrödinger type. Definition 2.4 requires showing

$$e^{itH} \pi_S(a) e^{-itH} \in \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right], \quad \forall a \in \mathcal{R}(\mathbb{R}^2, \sigma), \forall t \in \mathbb{R}, \quad (2)$$

hence the first issue to be dealt with is the explicit construction of e^{-itH} , $t \in \mathbb{R}$ given H as in (1). [7] is extremely useful for the announced purpose; concretely, by observing that

$$\begin{aligned} \Gamma_V(t) &= \mathbb{1} + \sum_{n \in \mathbb{N}} (-i)^n \int_0^t dt_n \cdots \int_0^{t_2} dt_1 V(t_1) \cdots V(t_n) \\ &= \mathbb{1} + \sum_{n \in \mathbb{N}} (-i)^n \int_0^t dt_n \cdots \left\{ \int_0^{t_3} dt_2 \left[\int_0^{t_2} dt_1 V(t_1) \right] V(t_2) \cdots \right\} V(t_n) \\ &= \mathbb{1} + \sum_{n \in \mathbb{N}} (-i)^n \int_0^t dt_n \Gamma_{V, (n-1)}(t_n) V(t_n) \equiv \mathbb{1} + \sum_{n \in \mathbb{N}} (-i)^n \Gamma_{V, (n)}(t), \end{aligned}$$

for all $t \in \mathbb{R}$, i.e. by considering

$$\Gamma_{V, (n)}(t) = \int_0^t dt_n \Gamma_{V, (n-1)}(t_n) V(t_n), \quad t \in \mathbb{R}, n \in \mathbb{N} \quad (3)$$

with $\Gamma_{V, (0)}(t) = \mathbb{1}$ for all $t \in \mathbb{R}$, $\Gamma_V(t)$ can be explicitly built out of the Fourier domain counterparts of $\Gamma_{V, (n)}(t)$, in turn defined by the integral kernels

$$K_{t, (1)}(p, q) = [\mathfrak{F} \Gamma_{V, (1)}(t) \mathfrak{F}^{-1}](p, q) = \left[\frac{e^{it(p^2 - q^2)} - 1}{i(p^2 - q^2)} \right] \frac{\tilde{V}(p - q)}{\sqrt{2\pi}} \quad (4)$$

$$K_{t, (n)}(p, q) = [\mathfrak{F} \Gamma_{V, (n)}(t) \mathfrak{F}^{-1}](p, q) = \int_0^t dt_n \int_{\mathbb{R}} dz_{n-1} K_{t_n, (n-1)}(p, z_{n-1}) e^{it_n(z_{n-1}^2 - q^2)} \frac{\tilde{V}(z_{n-1} - q)}{\sqrt{2\pi}}, \quad (5)$$

$t, p, q \in \mathbb{R}, n \in \mathbb{N} : n \geq 2$, where \mathfrak{F} is the Fourier-Plancherel operator. V is then allowed to be a distribution on \mathbb{R} whose Fourier transform \tilde{V} is a L^∞ function such that $\tilde{V}(p) = \tilde{V}(-p)$, $p \in \mathbb{R}$. Further, [7] ensures that each $K_{t, (n)}$ is a bounded operator on $L^2(\mathbb{R})$ for all $t \in \mathbb{R}$ and that the Dyson series $\sum_{n \in \mathbb{N}} K_{t, (n)}$ converges in the uniform norm topology; set, then, $K(t) = \mathbb{1} + \sum_{n \in \mathbb{N}} K_{t, (n)}$,

$$U(t) = e^{-itH_0} [\mathfrak{F}^{-1} K(t) \mathfrak{F}], \quad t \in \mathbb{R}$$

gives the rigorous unitary time evolution operator of a system governed by the symbolic Hamiltonian $H = H_0 + V$. \square

²In this section, $H_0 = -\frac{d^2}{dx^2}$ is assumed.

3.1 One Fixed-Center Point Interaction

One spinless particle undergoing a unique point interaction placed in a fixed location of the real line is considered. Its formal Hamiltonian is

$$H = -\frac{d^2}{dx^2} + \alpha\delta(x - x_0), \quad x_0 \in \mathbb{R}, \quad (6)$$

where $\alpha \in \mathbb{R} \setminus \{0\}$ is the coupling constant and $x_0 \in \mathbb{R}$ is the δ -location. Given $V = \alpha\delta(\cdot - x_0)$, one has

- $\tilde{V}(p) = \frac{\alpha}{(\sqrt{2\pi})} e^{-ipx_0}$, $p \in \mathbb{R} \implies \tilde{V} \in L^\infty(\mathbb{R})$,
- $\overline{\tilde{V}(p)} = \frac{\alpha}{(\sqrt{2\pi})} e^{-ipx_0} \equiv \frac{\alpha}{(\sqrt{2\pi})} e^{-i(-p)x_0} \equiv \tilde{V}(-p)$, $p \in \mathbb{R}$.

Consequently,

$$K_{t,(1)}^{(\alpha)}(p, q) = \frac{\alpha}{2\pi} \left[\frac{e^{it(p^2 - q^2)} - 1}{i(p^2 - q^2)} \right] e^{-i(p-q)x_0} \quad (7)$$

$$K_{t,(n)}^{(\alpha)}(p, q) = \frac{\alpha}{2\pi} \int_0^t dt_n \int_{\mathbb{R}} dz_{n-1} K_{t_n, (n-1)}^{(\alpha)}(p - z_{n-1}) e^{it_n(z_{n-1}^2 - q^2)} e^{-i(z_{n-1} - q)x_0}, \quad n \in \mathbb{N} \quad (8)$$

allow to build $\Gamma_V(t) \equiv \Gamma_\alpha(t)$ as described, hence the unitary time evolution operator $U_\alpha(t) = e^{-itH_0} \Gamma_\alpha(t)$ corresponding to (6), for all $t \in \mathbb{R}$. \square

Remark 3.1. *Endowed with the unitary time evolution operator, the resolvent algebra $\mathcal{R}(\mathbb{R}^2, \sigma)$ stability remains to be proved; the following strategy is adopted: given a non-negative smooth function of compact support W^3 , introducing W_ϵ as*

$$W_\epsilon : x \in \mathbb{R} \mapsto W_\epsilon(x) \doteq \frac{1}{\epsilon} W\left(\frac{x}{\epsilon}\right) \in \mathbb{R}, \quad \epsilon > 0,$$

along with the Schrödinger Hamiltonian ($H_\epsilon = H_0 + \alpha W_\epsilon$, \mathcal{D}_{H_0}), Proposition 2.5 allows to claim that $\exp(-itH_\epsilon) \in \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$ for all $t \in \mathbb{R}$. Therefore, should

$$\left\| U_\alpha(t) - e^{-itH_\epsilon} \right\|_{\mathcal{B}(L^2(\mathbb{R}))} \xrightarrow[\epsilon \downarrow 0]{} 0 \quad (9)$$

hold, the stability of $\mathcal{R}(\mathbb{R}^2, \sigma)$ would easily follow; in fact, for all $a \in \mathcal{R}(\mathbb{R}^2, \sigma)$, $t \in \mathbb{R}$,

$$\begin{aligned} & \left\| U_\alpha(t)^* \pi_S(a) U_\alpha(t) - e^{itH_\epsilon} \pi_S(a) e^{-itH_\epsilon} \right\| = \\ & = \left\| U_\alpha(t)^* \pi_S(a) U_\alpha(t) - e^{itH_\epsilon} \pi_S(a) U_\alpha(t) + e^{itH_\epsilon} \pi_S(a) U_\alpha(t) - e^{itH_\epsilon} \pi_S(a) e^{-itH_\epsilon} \right\| \leq \\ & \leq \left\| U_\alpha(t)^* - e^{itH_\epsilon} \right\| \left\| \pi_S(a) \right\| + \left\| \pi_S(a) \right\| \left\| U_\alpha(t) - e^{-itH_\epsilon} \right\| \xrightarrow[\epsilon \downarrow 0]{} 0. \end{aligned}$$

\square

³It does not harm generality assuming $\int_{\mathbb{R}} W = 1$

Remark 3.2. 1. $\{W_\epsilon\}_{\epsilon>0}$ converges to δ in $\mathcal{D}'(\mathbb{R})$ as $\epsilon \rightarrow 0$; in fact, given a whatever compactly supported real smooth function f on \mathbb{R} ,

$$\begin{aligned} \lim_{\epsilon \downarrow 0} \int_{\mathbb{R}} \left[\frac{1}{\epsilon} W \left(\frac{x - x_0}{\epsilon} \right) \right] f(x) dx &= \int_{\mathbb{R}} W(x) \left[\lim_{\epsilon \downarrow 0} f(\epsilon x + x_0) \right] dx = \left(\int_{\mathbb{R}} W \right) f(x_0) \equiv \\ &\equiv \int_{\mathbb{R}} [\delta(x - x_0)] f(x) dx, \end{aligned}$$

by using the Lebesgue dominated convergence theorem and $\int_{\mathbb{R}} W = 1$.

2. $\left\{ \mathcal{F} [W_\epsilon(\cdot - x_0)] \right\}_\epsilon^4$ is point-wise convergent to $\left[\left(\sqrt{2\pi} \right)^{-1} e^{-i(\cdot)x_0} \right]$ as $\epsilon \downarrow 0$ for all $x_0 \in \mathbb{R}$ and there exists $M \in \mathbb{R}^+$ such that $|\tilde{W}_\epsilon(p)| \leq M$, for all ϵ and p : straightforwardly,

$$\lim_{\epsilon \downarrow 0} \mathcal{F} [W_\epsilon(\cdot - x_0)](p) \equiv \lim_{\epsilon \downarrow 0} \tilde{W}_\epsilon(p) = \lim_{\epsilon \downarrow 0} \frac{e^{-ipx_0}}{\sqrt{2\pi}} \int_{\mathbb{R}} W(x) e^{-i(\epsilon p)x} dx = \frac{e^{-ipx_0}}{\sqrt{2\pi}}, \quad \forall p \in \mathbb{R},$$

by the Lebesgue dominated convergence theorem. Clearly

$$|\tilde{W}_\epsilon(p)| \leq \frac{1}{\sqrt{2\pi}} \equiv M.$$

□

Proposition 3.1. Let $\alpha \in \mathbb{R} \setminus \{0\}$ be and $W \in C_c^\infty(\mathbb{R})$ as in Remark 3.1. For all $t \in \mathbb{R}$,

$$\left\| U_\alpha(t) - e^{-itH_\epsilon} \right\|_{\mathcal{B}(L^2(\mathbb{R}))} \xrightarrow{\epsilon \downarrow 0} 0.$$

holds.

Proof. Set

$$K_{t,(1)}^{(\epsilon)}(p, q) = \frac{\alpha}{\sqrt{2\pi}} \left[\frac{e^{it(p^2 - q^2)} - 1}{i(p^2 - q^2)} \right] \tilde{W}_\epsilon(p - q) \quad (10)$$

$$K_{t,(n)}^{(\epsilon)}(p, q) = \frac{\alpha}{\sqrt{2\pi}} \int_0^t dt_n \int_{\mathbb{R}} dz_{n-1} K_{t_n, (n-1)}^{(\epsilon)}(p, z_{n-1}) e^{it_n(z_{n-1}^2 - q^2)} \tilde{W}_\epsilon(z_{n-1} - q), \quad n \in \mathbb{N} \quad (11)$$

it results

$$\begin{aligned} &\left\| U_\alpha(t) - e^{-itH_\epsilon} \right\| = \\ &= \left\| e^{-itH_0} \left[\mathbb{1} + \sum_{n \in \mathbb{N}} i^n \mathfrak{F}^{-1} K_{t,(n)}^{(\alpha)} \mathfrak{F} \right] - e^{-itH_0} \left[\mathbb{1} + \sum_{n \in \mathbb{N}} i^n \mathfrak{F}^{-1} K_{t,(n)}^{(\epsilon)} \mathfrak{F} \right] \right\| \leq \sum_{n \in \mathbb{N}} \left\| K_{t,(n)}^{(\alpha)} - K_{t,(n)}^{(\epsilon)} \right\|, \end{aligned}$$

meaning that proving the claim amounts in showing

$$\left\| K_{t,(n)}^{(\alpha)} - K_{t,(n)}^{(\epsilon)} \right\| \xrightarrow{\epsilon \downarrow 0} 0, \quad \forall n \in \mathbb{N}, \forall t \in \mathbb{R}.$$

⁴ \mathcal{F} denotes the L^1 -Fourier transform operator.

[7] thm. 3.4 allows for

$$\left\| K_{t,(n)}^{(\alpha)} - K_{t,(n)}^{(\epsilon)} \right\| \leq \left\{ \left(\sup_{p \in \mathbb{R}} \int_{\mathbb{R}} \left| K_{t,(n)}^{(\alpha)}(p, q) - K_{t,(n)}^{(\epsilon)}(p, q) \right| dq \right) \left(\sup_{p \in \mathbb{R}} \int_{\mathbb{R}} \left| K_{t,(n)}^{(\alpha)*}(p, q) - K_{t,(n)}^{(\epsilon)*}(p, q) \right| dq \right) \right\}^{\frac{1}{2}} < \infty, \quad n \in \mathbb{N}$$

hence the *induction principle* is going to be used.

$$\boxed{k = 1}$$

$$\lim_{\epsilon \downarrow 0} \int_{\mathbb{R}} \left| K_{t,(1)}^{(\alpha)}(p, q) - K_{t,(1)}^{(\epsilon)}(p, q) \right| dq = \frac{|\alpha|}{\sqrt{2\pi}} \lim_{\epsilon \downarrow 0} \int_{\mathbb{R}} \left| \frac{e^{it(p^2-q^2)} - 1}{p^2 - q^2} \right| \left| \frac{e^{-i(p-q)x_0}}{\sqrt{2\pi}} - \tilde{W}_\epsilon(p-q) \right| dq$$

is intended to be studied. By observing that

$$\frac{|\alpha|}{\sqrt{2\pi}} \int_{\mathbb{R}} \left| \frac{e^{it(p^2-q^2)} - 1}{p^2 - q^2} \right| \left| \frac{e^{-i(p-q)x_0}}{\sqrt{2\pi}} - \tilde{W}_\epsilon(p-q) \right| dq \leq \sqrt{\frac{2}{\pi}} |\alpha| M \int_{\mathbb{R}} \left| \frac{e^{it(p^2-q^2)} - 1}{p^2 - q^2} \right| dq < \infty$$

because of [7] thm. 2.3, remark 3.2 and the dominated convergence theorem allow for

$$\int_{\mathbb{R}} \left| \frac{e^{it(p^2-q^2)} - 1}{p^2 - q^2} \right| \left\{ \lim_{\epsilon \downarrow 0} \left| \frac{e^{-i(p-q)x_0}}{\sqrt{2\pi}} - \tilde{W}_\epsilon(p-q) \right| \right\} dq = 0$$

Since $K_{t,(1)} = K_{t,(1)}^*$,

$$\lim_{\epsilon \downarrow 0} \int_{\mathbb{R}} \left| K_{t,(1)}^{*(\alpha)}(p, q) - K_{t,(1)}^{*(\epsilon)}(p, q) \right| dq = 0$$

holds all the same.

$\boxed{k = n}$ It is assumed the statement holds for $k \leq n - 1$.

$$K_{t,(n)}^{(\alpha)}(p, q) - K_{t,(n)}^{(\epsilon)}(p, q) = \tag{12}$$

$$= \frac{\alpha}{\sqrt{2\pi}} \int_0^t dt_n \int_{\mathbb{R}} dz_{n-1} e^{it_n(z_{n-1}^2 - q^2)} K_{t_n, (n-1)}^{(\alpha)}(p, z_{n-1}) \left[\frac{e^{-i(z_{n-1}-q)x_0}}{\sqrt{2\pi}} - \tilde{W}_\epsilon(z_{n-1}-q) \right] + \tag{13}$$

$$+ \frac{\alpha}{\sqrt{2\pi}} \int_0^t dt_n \int_{\mathbb{R}} dz_{n-1} e^{it_n(z_{n-1}^2 - q^2)} \left[K_{t_n, (n-1)}^{(\alpha)}(p, z_{n-1}) - K_{t_n, (n-1)}^{(\epsilon)}(p, z_{n-1}) \right] \tilde{W}_\epsilon(z_{n-1}-q). \tag{14}$$

A priori, the foregoing integrals are *double* integrals; to use them as *iterated*, Fubini theorem hypotheses have to be ascertained.

$$\begin{aligned} \int_0^t \int_{\mathbb{R}} \left| K_{t_n, (n-1)}^{(\alpha)}(p, z_{n-1}) \right| \left| \frac{e^{-i(z_{n-1}-q)x_0}}{\sqrt{2\pi}} - \tilde{W}_\epsilon(z_{n-1}-q) \right| dz_{n-1} dt_n &\leq \\ &\leq 2M \int_0^t \int_{\mathbb{R}} \left| K_{t_n, (n-1)}^{(\alpha)}(p, z_{n-1}) \right| dz_{n-1} dt_n < \infty \end{aligned}$$

by being $K_{t,(n)}^{(\alpha)} \in \mathfrak{B}(L^2(\mathbb{R}))$, $\forall n \in \mathbb{N}$, $\forall t \in \mathbb{R}$. Fubini also holds for both (14) and the adjoint case, as can be readily verified. Then, to compute

$$\lim_{\epsilon \downarrow 0} \int_{\mathbb{R}} \left| K_{t,(n)}^{(\alpha)}(p, q) - K_{t,(n)}^{(\epsilon)}(p, q) \right| dq,$$

the dominated convergence theorem hypotheses need to be checked out. Therefore

$$\begin{aligned} & \left| \int_0^t \int_{\mathbb{R}} K_{t_n,(n-1)}^{(\alpha)}(p, z_{n-1}) \left[\frac{e^{-i(z_{n-1}-q)x_0}}{\sqrt{2\pi}} - \tilde{W}_\epsilon(z_{n-1}-q) \right] e^{it_n(z_{n-1}^2-q^2)} dz_{n-1} dt_n \right| \leq \\ & \leq \left| \int_{\mathbb{R}} \left[\frac{e^{-i(z_{n-1}-q)x_0}}{\sqrt{2\pi}} - \tilde{W}_\epsilon(z_{n-1}-q) \right] \int_0^t K_{t_n,(n-1)}^{(\alpha)}(p, z_{n-1}) e^{it_n(z_{n-1}^2-q^2)} dt_n dz_{n-1} \right| \leq \\ & \leq 2M \int_{\mathbb{R}} \left| \int_0^t K_{t_n,(n-1)}^{(\alpha)}(p, z_{n-1}) e^{it_n(z_{n-1}^2-q^2)} dt_n \right| dz_{n-1} \equiv \\ & \equiv (2M) \tilde{K}_{t,(n)}^{(\alpha)}(p, q), \end{aligned}$$

i.e.

$$\begin{aligned} & \int_{\mathbb{R}} \left| \int_0^t \int_{\mathbb{R}} K_{t_n,(n-1)}^{(\alpha)}(p, z_{n-1}) \left[\frac{e^{-i(z_{n-1}-q)x_0}}{\sqrt{2\pi}} - \tilde{W}_\epsilon(z_{n-1}-q) \right] e^{it_n(z_{n-1}^2-q^2)} dz_{n-1} dt_n \right| dq \leq \\ & \leq 2M \int_{\mathbb{R}} \tilde{K}_{t,(n)}^{(\alpha)}(p, q) dq \leq 2M \left[\sup_{p \in \mathbb{R}} \int_{\mathbb{R}} \tilde{K}_{t,(n)}^{(\alpha)}(p, q) dq \right] < \infty, \end{aligned}$$

the estimate holding because of [7] thm. 3.4. Hence, concerning (13),

$$\begin{aligned} & \lim_{\epsilon \downarrow 0} \int_{\mathbb{R}} \left| \int_0^t \int_{\mathbb{R}} K_{t_n,(n-1)}^{(\alpha)}(p, z_{n-1}) \left[\frac{e^{-i(z_{n-1}-q)x_0}}{\sqrt{2\pi}} - \tilde{W}_\epsilon(z_{n-1}-q) \right] e^{it_n(z_{n-1}^2-q^2)} dz_{n-1} dt_n \right| dq \leq \\ & \leq \int_{\mathbb{R}} \int_0^t \int_{\mathbb{R}} \left| K_{t_n,(n-1)}^{(\alpha)}(p, z_{n-1}) \right| \left[\lim_{\epsilon \downarrow 0} \left| \frac{e^{-i(z_{n-1}-q)x_0}}{\sqrt{2\pi}} - \tilde{W}_\epsilon(z_{n-1}-q) \right| \right] dz_{n-1} dt_n dq = 0. \end{aligned}$$

On the other hand, regarding (14), the inductive hypothesis gives

$$\begin{aligned} & \lim_{\epsilon \downarrow 0} \int_{\mathbb{R}} \left| \int_0^t \int_{\mathbb{R}} \left[K_{t_n,(n-1)}^{(\alpha)}(p, z_{n-1}) - K_{t_n,(n-1)}^{(\epsilon)}(p, z_{n-1}) \right] \tilde{W}_\epsilon(z_{n-1}-q) e^{it_n(z_{n-1}^2-q^2)} dz_{n-1} dt_n \right| dq \leq \\ & \leq M \int_{\mathbb{R}} \int_0^t \left\{ \lim_{\epsilon \downarrow 0} \int_{\mathbb{R}} \left| K_{t_n,(n-1)}^{(\alpha)}(p, z_{n-1}) - K_{t_n,(n-1)}^{(\epsilon)}(p, z_{n-1}) \right| dq \right\} dt_n dz_{n-1} = 0. \end{aligned}$$

By proceeding analogously for the adjoint relations, the Schur test gives

$$\left\| K_{t,(n)}^{(\alpha)} - K_{t,(n)}^{(\epsilon)} \right\| \xrightarrow{\epsilon \downarrow 0} 0, \quad \forall t \in \mathbb{R}, \quad \forall n \in \mathbb{N}.$$

■

Proposition 3.2. *What follows holds.*

1. $U_\alpha^*(t) \pi_S(a) U_\alpha(t) \in \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$ for all $a \in \mathcal{R}(\mathbb{R}^2, \sigma)$;
2. Denoted by $(H_\alpha, \mathcal{D}_{H_\alpha})$ the self-adjoint operator on $L^2(\mathbb{R})$ generating to the one parameter family of strongly continuous unitary operators $\{U_\alpha(t)\}_{t \in \mathbb{R}}$, $(H_\alpha, \mathcal{D}_{H_\alpha})$ is affiliated to $\mathcal{R}(\mathbb{R}^2, \sigma)$;
3. The map $\alpha_t : a \in \mathcal{R}(\mathbb{R}^2, \sigma) \mapsto \alpha_t(a) \in \mathcal{R}(\mathbb{R}^2, \sigma)$, with

$$\alpha_t(a) \doteq \pi_S^{-1} \left[e^{itH_\alpha} \pi_S(a) e^{-itH_\alpha} \right],$$

results in an automorphism of $\mathcal{R}(\mathbb{R}^2, \sigma)$ for all $t \in \mathbb{R}$.

Proof. 1. Directly from Proposition 3.1 and remark 3.1.

2. It is a very well known fact that norm dynamical convergence⁵ implies⁶ norm resolvent convergence, therefore

$$\left\| U_\alpha(t) - e^{-itH_\epsilon} \right\| \xrightarrow{\epsilon \downarrow 0} 0, \forall t \in \mathbb{R} \implies \left\| (H_\alpha - i\lambda \mathbb{1})^{-1} - (H_\epsilon - i\lambda \mathbb{1})^{-1} \right\| \xrightarrow{\epsilon \downarrow 0} 0, \forall \lambda \in \mathbb{R} \setminus \{0\}.$$

Proposition 2.5 states that $(H_\epsilon - i\lambda \mathbb{1})^{-1} \in \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$ for all $\lambda \in \mathbb{R} \setminus \{0\}$ and $\epsilon > 0$.

Since $\pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$ is closed with respect to the uniform norm topology, the affiliation of $(H_\alpha, \mathcal{D}_{H_\alpha})$ results.

3. Given $t \in \mathbb{R}$, the map

$$a \in \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right] \mapsto e^{itH_\alpha} a e^{-itH_\alpha} \in \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right] \quad (15)$$

is surely injective, being isometric. On the other hand, given $b \in \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$, because of proposition 3.1, $e^{-itH_\alpha} b e^{itH_\alpha} \equiv d \in \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$, hence

$$e^{itH_\alpha} d e^{-itH_\alpha} = b,$$

allowing to conclude that (15) is surjective. The same map is obviously a homomorphism; finally, being $\pi_S : \mathcal{R}(\mathbb{R}^2, \sigma) \longrightarrow \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$ an isomorphism, the result follows. ■

5

Definition 3.1. Let \mathcal{H} be a complex Hilbert space. Given self-adjoint operators (A_n, \mathcal{D}_{A_n}) , (A, \mathcal{D}_A) , A_n is **norm dynamically convergent** to A if and only if, for all $t \in \mathbb{R}$, $\{e^{itA_n}\}_n$ converges to e^{itA} with respect to the $\mathcal{B}(\mathcal{H})$ norm. □

⁶See [8], thm. 10.1.16.

3.2 Many Fixed-Centers Point Interactions

3.2.1 Finitely Many Fixed-Centers Point Interactions

Focus is set on the symbolic Hamiltonian

$$H = H_0 + \sum_{i=1}^N \alpha_i \delta(x - x_i), \quad (16)$$

with $N \in \mathbb{N}$, coupling constants $\alpha_i \in \mathbb{R} \setminus \{0\}$ and fixed-centers location $x_i \in \mathbb{R} : x_i \neq x_j$. By setting $(\alpha) \equiv (\alpha_1, \dots, \alpha_N) \in \mathbb{R}^N$ and $V = \sum_{i=1}^N \alpha_i \delta(\cdot - x_i)$,

1.

$$\tilde{V}(p) = \int_{\mathbb{R}} \left[\sum_{m=1}^N \alpha_m \delta(x - x_m) \right] e^{-ipx} \frac{dx}{\sqrt{2\pi}} = \sum_{m=1}^N \frac{\alpha_m}{\sqrt{2\pi}} e^{-ipx_m},$$

i.e. $\tilde{V} \in L^\infty(\mathbb{R})$ and

2.

$$\tilde{V}(u) = \sum_{m=1}^N \frac{\alpha_m}{\sqrt{2\pi}} e^{-ipx_m} = \sum_{m=1}^N \frac{\alpha_m}{\sqrt{2\pi}} e^{i(-p)x_m} = \overline{\sum_{m=1}^N \frac{\alpha_m}{\sqrt{2\pi}} e^{-i(-p)x_m}} = \overline{\tilde{V}(-p)},$$

for all $p \in \mathbb{R}$.

Consequently, one legitimately relies on

$$K_{t,(1)}^{(\alpha)}(p, q) = \left[\frac{e^{it(p^2 - q^2)} - 1}{i(p^2 - q^2)} \right] \left[\sum_{m=1}^N \frac{\alpha_m}{\sqrt{2\pi}} e^{-i(p-q)x_m} \right] \equiv \sum_{m=1}^N K_{t,(1)}^{(\alpha),m}(p, q), \quad (17)$$

$$K_{t,(n)}^{(\alpha)}(p, q) = \int_0^t \int_{\mathbb{R}} K_{t_n, (n-1)}^{(\alpha)}(p, z_{n-1}) \left[\sum_{m=1}^N \frac{\alpha_m}{\sqrt{2\pi}} e^{-i(z_{n-1}-q)x_m} \right] e^{it_n(z_n^2 - q^2)} dz_{n-1} dt_n = \quad (18)$$

$$\equiv \sum_{m=1}^N K_{t,(n)}^{(\alpha),m}(p, q), \quad (19)$$

for all $t, p, q \in \mathbb{R}$, to build $\Gamma_{(\alpha)}(t)$ up, hence the unitary time evolution operator $U_{(\alpha)}(t)$, $t \in \mathbb{R}$.

Proposition 3.3. *Given $N \in \mathbb{N}$, let $\alpha_1, \dots, \alpha_N \in \mathbb{R} \setminus \{0\}$ and non-negative smooth functions of compact support $W_1, \dots, W_N \in C_c^\infty(\mathbb{R})$ such that $\int_{\mathbb{R}} W_i = 1$ be. Considered the Schrödinger Hamiltonians*

$$H_\epsilon = H_0 + \sum_{i=1}^N \alpha_i W_{\epsilon,i} \equiv H_0 + W_\epsilon, \quad \epsilon > 0,$$

where $W_{\epsilon,i}(x) = \epsilon^{-1} W_i(x/\epsilon)$, $x \in \mathbb{R}$, $i \in \{1, \dots, N\}$, for all $t \in \mathbb{R}$,

$$\left\| U_{(\alpha)}(t) - e^{-itH_\epsilon} \right\|_{\mathcal{B}(L^2(\mathbb{R}))} \xrightarrow{\epsilon \downarrow 0} 0.$$

Proof. By using

$$K_{t,(1)}^{(\epsilon)}(p, q) = \left[\frac{e^{it(p^2 - q^2)} - 1}{i(p^2 - q^2)} \right] \frac{\tilde{W}_\epsilon(p - q)}{\sqrt{2\pi}} \equiv \sum_{m=1}^N K_{t,(1)}^{(\epsilon), m}(p, q) \quad (20)$$

$$K_{t,(n)}^{(\epsilon)}(p, q) = \int_0^t \int_{\mathbb{R}} K_{t_n, (n-1)}^{(\epsilon)}(p, z_{n-1}) \frac{\tilde{W}_\epsilon(z_{n-1} - q)}{\sqrt{2\pi}} e^{it_n(z_{n-1}^2 - q^2)} dz_{n-1} dt_n = \sum_{m=1}^N K_{t,(n)}^{(\epsilon), m}(p, q) \quad (21)$$

for all $t, p, q \in \mathbb{R}$, to build $\exp(-itH_\epsilon)$ up, $t \in \mathbb{R}$, one then has

$$\begin{aligned} \left\| U_{(\alpha)}(t) - e^{-itH_\epsilon} \right\| &\leq \sum_{n \in \mathbb{N}} \left\| K_{n,t,s}^{(\alpha)} - K_{n,t,s}^{(\epsilon)} \right\| \leq (\text{by Schur test}) \\ &\leq \sum_{n \in \mathbb{N}} \left\{ \left[\sup_p \int_{\mathbb{R}} \left| K_{t,(n)}^{(\alpha)}(p, q) - K_{t,(n)}^{(\epsilon)}(p, q) \right| dq \right] \left[\sup_p \int_{\mathbb{R}} \left| K_{t,(n)}^{(\alpha),*}(p, q) - K_{t,(n)}^{(\epsilon),*}(p, q) \right| dq \right] \right\}^{\frac{1}{2}}. \end{aligned}$$

Therefore, by observing that

$$\int_{\mathbb{R}} \left| K_{t,(n)}^{(\alpha)}(p, q) - K_{t,(n)}^{(\epsilon)}(p, q) \right| dq = \int_{\mathbb{R}} \left| \sum_{m=1}^N \left[K_{t,(n)}^{(\alpha), m}(p, q) - K_{t,(n)}^{(\epsilon), m}(p, q) \right] \right| dq \leq \quad (22)$$

$$\leq \sum_{m=1}^N \int_{\mathbb{R}} \left| K_{t,(n)}^{(\alpha), m}(p, q) - K_{t,(n)}^{(\epsilon), m}(p, q) \right| dq, \quad (23)$$

the result is proved as in Proposition 3.1. ■

Proposition 3.4. *What follows holds.*

1. $U_{(\alpha)}^*(t) \pi_S(a) U_{(\alpha)}(t) \in \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$ for all $a \in \mathcal{R}(\mathbb{R}^2, \sigma)$;
2. Denoted by $\left(H_{(\alpha)}, \mathcal{D}_{H_{(\alpha)}} \right)$ the self-adjoint operator on $L^2(\mathbb{R})$ generating the one parameter family of strongly continuous unitary operators $\{U_{(\alpha)}(t)\}_{t \in \mathbb{R}}$, $\left(H_{(\alpha)}, \mathcal{D}_{H_{(\alpha)}} \right)$ is affiliated to $\mathcal{R}(\mathbb{R}^2, \sigma)$;
3. The map $\alpha_t : a \in \mathcal{R}(\mathbb{R}^2, \sigma) \mapsto \alpha_t(a) \in \mathcal{R}(\mathbb{R}^2, \sigma)$, with

$$\alpha_t(a) \doteq \pi_S^{-1} \left[e^{itH_{(\alpha)}} \pi_S(a) e^{-itH_{(\alpha)}} \right],$$

results in an automorphism of $\mathcal{R}(\mathbb{R}^2, \sigma)$ for all $t \in \mathbb{R}$.

Proof. The proof closely mimics the one of Proposition 3.2. ■

3.2.2 Countably Many Fixed-Centers Point Interactions

Given $\{\alpha_i\}_{i \in \mathbb{N}} \in l^1(\mathbb{N}) \setminus \{0\}$ the symbolic Hamiltonian

$$H = -\frac{d^2}{dx^2} + \sum_{i=1}^{\infty} \alpha_i \delta(x - x_i),$$

with $\{x_i\}_{i \in \mathbb{N}} \subset \mathbb{R}$ such that $x_i \neq x_j$, for all $i \neq j$, is finally considered. Set $V = \sum_{i=1}^{\infty} \alpha_i \delta(\cdot - x_i)$,

1. $\tilde{V}(p) = \frac{1}{\sqrt{2\pi}} \sum_{m \in \mathbb{N}} \alpha_m e^{-ipx_m} = \overline{\frac{1}{\sqrt{2\pi}} \sum_{m \in \mathbb{N}} \alpha_m e^{ipx_m}} = \overline{\tilde{V}(-p)}$, $\forall p \in \mathbb{R}$ and
2. $\tilde{V} \in L^\infty(\mathbb{R})$,

therefore $\Gamma_{\{\alpha_i\}}(t)$ can be obtained via (4), (5).

Proposition 3.5. *What follows holds.*

1. For all $t \in \mathbb{R}$, the unitary time evolution operator $U_{\{\alpha_i\}}(t) \doteq e^{-itH_0} \Gamma_{\{\alpha_i\}}(t)$ belongs to $\pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$. Moreover, denoted by $\left(H_{\{\alpha_i\}}, \mathcal{D}_{H_{\{\alpha_i\}}} \right)$ the self-adjoint operator on $L^2(\mathbb{R})$ generating the one parameter family of unitary operators $\{U_{\{\alpha_i\}}(t)\}_{t \in \mathbb{R}}$, it is affiliated to $\mathcal{R}(\mathbb{R}^2, \sigma)$.
2. $U_{\{\alpha_i\}}(t)^* \pi_S(a) U_{\{\alpha_i\}}(t) \in \pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$ for all $a \in \mathcal{R}(\mathbb{R}^2, \sigma)$, $t \in \mathbb{R}$.
3. The map

$$\alpha_t : a \in \mathcal{R}(\mathbb{R}^2, \sigma) \longmapsto \alpha_t(a) = \pi_S^{-1} \left[U_{\{\alpha_i\}}(t)^* \pi_S(a) U_{\{\alpha_i\}}(t) \right] \in \mathcal{R}(\mathbb{R}^2, \sigma)$$

is an automorphism of $\mathcal{R}(\mathbb{R}^2, \sigma)$.

Proof. Concerning 1., the result follows from [9] prop. 2, Proposition 3.3 and the fact that norm dynamical convergence implies norm resolvent convergence. 2. and 3. are proved as in Proposition 3.2. ■

Remark 3.3. *The non-trivial ideal structure of the resolvent algebra has already proved to be fundamental for the possibility of accommodating non-trivial quantum dynamics. The same feature is also of primary importance for the following final result to hold.* □

Proposition 3.6. *Let \mathfrak{K}_0 be the C^* -subalgebra of $\pi_S \left[\mathcal{R}(\mathbb{R}^2, \sigma) \right]$ generated by $\mathcal{B}_\infty(L^2(\mathbb{R}))$ and the identity operator. $(\mathcal{K}_0 \equiv \pi_S^{-1}(\mathfrak{K}_0), \mathbb{R}, \beta)$, where*

$$\beta : t \in \mathbb{R} \longmapsto \beta_t \in \text{Aut}(\mathcal{K}_0)$$

and

$$\beta_t : a \in \mathcal{K}_0 \longmapsto \beta_t(a) \doteq \pi_S^{-1} \left[U(t)^* \pi_S(a) U(t) \right] \in \mathcal{K}_0,$$

$U(t) \in \mathcal{B}(L^2(\mathbb{R}))$ propagating fixed point interactions, is a C^* -dynamical system.

Proof. First of all, it is observed that Proposition 2.4 allows for $\mathcal{B}_\infty(L^2(\mathbb{R}))$ to be contained in $\pi_S[\mathcal{R}(\mathbb{R}^2, \sigma)]$; then, for all $t_0 \in \mathbb{R}$, $\|U(t)^*U(t) - U(t_0)^*U(t_0)\| \xrightarrow[t \rightarrow t_0]{} 0$. On the other hand, given $\psi, \varphi \in L^2(\mathbb{R})$, let the finite rank operator $T = \langle \psi, \cdot \rangle \varphi$ be. Fixed again $t_0 \in \mathbb{R}$, it results

$$\|U(t)^*T - U(t_0)^*T\| \leq \|\psi\| \|U(t)^*\varphi - U(t_0)^*\varphi\| \xrightarrow[t \rightarrow t_0]{} 0.$$

Analogously,

$$\|TU(t) - TU(t_0)\| \leq \|\varphi\| \|U(t)^*\psi - U(t_0)^*\psi\| \xrightarrow[t \rightarrow t_0]{} 0,$$

therefore

$$\|U(t)^*TU(t) - U(t_0)^*TU(t_0)\| \leq \|U(t)^*T - U(t_0)^*T\| + \|TU(t) - TU(t_0)\| \xrightarrow[t \rightarrow t_0]{} 0.$$

Linearity, density and continuity arguments prove the statement. ■

4 Conclusions

This paper shows that resolvent algebras can accommodate dynamics induced by self-adjoint Hamiltonians describing a single non-relativistic spinless particle undergoing one up to countably many different fixed point interactions located on the real line. Such a result is a one of a kind result, since, apart from Buchholz investigations, it deals with the open problem of establishing which physical systems the resolvent algebra formalism can manage. Moreover, the non-simple nature of such an algebra contributes in singling out a C^* -subalgebra of $\pi_S[\mathcal{R}(\mathbb{R}^2, \sigma)]$ constituting a C^* -dynamical system.

Acknowledgements

A. M. is grateful to professors Romeo Brunetti and Detlev Buchholz for precious comments, ideas and suggestions, allowing for this work to exist as it is as well as to the colleagues Andrea Moro, Matteo Crispino, Daniele Volpe and Dario De Stefano for their help with linguistics and debugging.

References

- [1] D. Buchholz, H. Grundling - The Resolvent Algebra: A New Approach to Canonical Quantum Systems, *Jour. Func. An.* 254, 2725-2779 (2008);
- [2] M. Fannes, A. Verbeure - On the Time Evolution Automorphisms of the CCR-Algebra for Quantum Mechanics, *Commun. Math. Phys.* 35, 257-264 (1974);
- [3] D. Buchholz - The resolvent algebra of non-relativistic Bose fields: Observables, Dynamics and States, *Commun. Math. Phys.*, 363, 949-981 (2018);
- [4] D. Buchholz - The Resolvent Algebra for Oscillating Lattice Systems, *Commun. Math. Phys.*, 353, 691-716 (2017);
- [5] D. Buchholz - The Resolvent Algebra: Ideals and Dimensions, *Jour. Func. An.* 266, 3286-3302 (2014);

- [6] S. Albeverio, F. Gesztesy, R. Heigh-Krohn, H. Holden - Solvable Models in Quantum Mechanics, AMS Chelsea Publishing, 2005;
- [7] R. J. Hughes, I. E. Segal - Singular Perturbation in the Interaction Representation, Jour. Func. An. 38, 71-98 (1980);
- [8] C. R. de Oliveira - Intermediate Spectral Theory and Quantum Dynamics, Birkhäuser, 2009;
- [9] R. J. Hughes, M. A. Kon - Norm group convergence for singular Schrödinger operators, Annales de l'I. H. P., sec. A tome 54 n.2 (1991), p. 179-198.