

Fine-tuning of the quasi-bound K^-pp state

N.V. Shevchenko

Received: date / Accepted: date

Abstract Characteristics of the quasi-bound state in the K^-pp system strongly depend on the model of antikaon-nucleon interaction and weakly - on the nucleon-nucleon potential. In the present paper, dynamically exact Faddeev-type calculations with coupled $\bar{K}NN$ and $\pi\Sigma N$ were performed using different models of the ΣN and πN interactions to study the influence of these "less important" interaction on the three-body result. In addition, dynamically exact three-body Faddeev-type AGS calculations with three coupled particle channels $\bar{K}NN - \pi\Sigma N - \pi\Lambda N$ were performed.

Keywords Few-body exotic systems · AGS equations · Antikaon-nucleon interaction

1 Introduction

The attractive nature of $\bar{K}N$ interaction has stimulated theoretical and experimental searches for K^- bound states in different systems. In particular, many theoretical calculations devoted to the lightest possible system $\bar{K}NN$ were performed using different methods and inputs. All of them agree that a quasi-bound state in the K^-pp system exists, but they yield quite diverse binding energies and widths. The experimental situation is also unsettled: several evidences for the K^-pp state were reported by several experimental groups, but the estimated binding energies and especially decay widths of such states differ from each other and are far from all theoretical predictions. The most recent E15 experiment at J-PARC [1] reported the first clear signal of the $\bar{K}NN$ quasi-bound state with binding energy $42 \pm 3(\text{stat.})_{-4}^{+3}(\text{syst.})$ MeV, and width $100 \pm 7(\text{stat.})_{-9}^{+19}(\text{syst.})$ MeV. The measured binding energy of the state is comparable with some of the theoretical results, while the width is much larger than the theoretical predictions.

The work was supported by the THEIA, STRONG-2020 Project funded by the EU Framework Programme for Research and Innovation, Horizon 2020

N.V. Shevchenko
Nuclear Physics Institute, 25068 Řež, Czech Republic
E-mail: shevchenko@ujf.cas.cz

We studied the quasi-bound state in the K^-pp system solving Faddeev-type dynamically exact three-body Alt-Grassberger-Sandhas (AGS) equations [3] with coupled $\bar{K}NN - \pi\Sigma N$ channels with different input. The most recent results of our calculations can be found in [2]. In particular, three models of the $\bar{K}N$ interaction were used, and calculations with two of them give K^-pp binding energies close to the experimental values [1]. However, the experimental width measured by the E15 experiment at J-PARC is much larger than our predictions. In our previous calculations we demonstrated that the antikaon-nucleon interaction plays the main role in the description of the K^-pp system. The binding energy and width of the quasi-bound state in the system strongly depend on it. On the contrary, dependence on the nucleon-nucleon potential is weak. The present calculations aim to check, what else could change the theoretical predictions for the characteristics of the K^-pp quasi-bound state. To do this we studied the dependence of the binding energy and width of the quasi-bound state on the $\Sigma N - \Lambda N$ interaction models and used different versions of the πN potential, excluded from the previous calculations. In addition, we performed for the first time dynamically exact three-body Faddeev-type AGS calculations with three coupled channels $\bar{K}NN$, $\pi\Sigma N$, and $\pi\Lambda N$. For this sake new $\bar{K}N - \pi\Sigma - \pi\Lambda$ potentials were constructed and used.

The paper is organized as follows. The three-body AGS equations with two coupled $\bar{K}NN - \pi\Sigma N$ channels are described in Section 2, the section after it is devoted to the two-body interaction models used in the calculations. Section 4 contains description of the AGS equations with three coupled $\bar{K}NN - \pi\Sigma N - \pi\Lambda N$ channels and the corresponding $\bar{K}N - \pi\Sigma - \pi\Lambda$ antikaon-nucleon potentials. The results of the three-body calculations are separated at those obtained with two coupled $\bar{K}NN - \pi\Sigma N$ (Section 5) and three coupled $\bar{K}NN - \pi\Sigma N - \pi\Lambda N$ channels (Section 6). The paper is finished with the Summary.

2 Three-body AGS equations with two coupled $\bar{K}NN - \pi\Sigma N$ channels

The quasi-bound state in the K^-pp system was studied using dynamically exact Faddeev-type AGS equations [3]. The system of three-body equations was written for separable potentials $V_i^{\alpha\beta}$ leading to separable T -matrices $T_i^{\alpha\beta}$

$$V_i^{\alpha\beta} = \lambda_i^{\alpha\beta} |g_i^\alpha\rangle \langle g_i^\beta| \quad \rightarrow \quad T_i^{\alpha\beta} = |g_i^\alpha\rangle \tau_i^{\alpha\beta} \langle g_i^\beta|, \quad (1)$$

where one-term separable potentials are assumed for simplicity. The AGS equations in the operator form

$$X_{ij}^{\alpha\beta}(z) = \delta_{\alpha\beta} Z_{ij}^\alpha + \sum_{k=1}^3 \sum_{\gamma=1}^3 Z_{ik}^\alpha \tau_k^{\alpha\gamma} X_{kj}^{\gamma\beta} \quad (2)$$

contain three-body transition $X_{ij}^{\alpha\beta}$ and kernel Z_{ij}^α operators. Faddeev indices $i, j, k = 1, 2, 3$ in Eq.(2) simultaneously denote a pair of particles and the corresponding spectator (which is the third particle). Since antikaon-nucleon interaction is coupled to $\pi\Sigma$ channel through $\Lambda(1405)$ resonance, the corresponding potentials with coupled $\bar{K}N$ and $\pi\Sigma$ channels were used. The second, $\pi\Sigma$ channel was included into the three-body equations Eq.(2) directly, due to this additional indices α, β, γ denoting a particle channel

$$\alpha = 1 : |\bar{K}_1 N_2 N_3\rangle, \quad \alpha = 2 : |\pi_1 \Sigma_2 N_3\rangle, \quad \alpha = 3 : |\pi_1 N_2 \Sigma_3\rangle \quad (3)$$

were introduced. The system of operator equations (2) was written down in momentum representation and antisymmetrized due to identical fermions in the $\bar{K}NN$ channel. The corresponding system of integral equations was solved numerically. The three-body system with spin $S^{(3)} = 0$ and isospin $I^{(3)} = 1/2$ was studied.

3 Two-body input

The input for the three-body Faddeev-type AGS equations Eq.(2) are two-body T -matrices Eq.(1), describing two-body interactions between all pairs of particles in all particle channels. Since searching for quasibound states assumes working at low energies, and low-energy interactions are satisfactorily described by s waves, we used orbital momentum $l = 0$ for all the two-body interactions.

3.1 Antikaon-nucleon interaction

Three our antikaon-nucleon interaction models, constructed before, were used in the present calculations. Two of them are phenomenological potentials $V_{\bar{K}N}^{1,\text{SIDD}}$ and $V_{\bar{K}N}^{2,\text{SIDD}}$ with one- or two-pole structure of the $\Lambda(1405)$ resonance correspondingly [4]. The phenomenological potentials couple two channels: $\bar{K}N$ and $\pi\Sigma$. The third, a chirally motivated potential $V_{\bar{K}N}^{\text{Chiral}}$ [5] with two-pole structure of the $\Lambda(1405)$ resonance describes three coupled channels: $\bar{K}N$, $\pi\Sigma$, and $\pi\Lambda$.

All three potentials with the same level of accuracy reproduce low-energy experimental data on K^-p system. In particular, they reproduce $1s$ level shift and width of kaonic hydrogen caused by strong interaction and measured by SIDHARTA collaboration [6]. The $1s$ level of the atom was calculated by us directly, without using some approximate Deser-type formula: Lippmann-Schwinger equation for the strong $\bar{K}N - \pi\Sigma(-\pi\Lambda)$ plus Coulomb K^-p potentials was solved. The experimental cross-sections of the elastic $K^-p \rightarrow K^-p$ and inelastic $K^-p \rightarrow MB$ reactions [7, 8, 9, 10, 11, 12] are also reproduced by all three of our potentials. Accurately measured threshold branching ratios γ , R_c and R_n [13, 14] are reproduced by the chirally motivated model $V_{\bar{K}N}^{\text{Chiral}}$ directly. The phenomenological potentials $V_{\bar{K}N}^{1,\text{SIDD}}$ and $V_{\bar{K}N}^{2,\text{SIDD}}$ reproduce experimental value of γ ratio directly, while the measured R_c and R_n were used for construction of an artificial $R_{\pi\Sigma}$ branching ratio, reproduced by the potentials [4]. The reason is the absence of the directly included $\pi\Lambda$ channel in the phenomenological antikaon-nucleon potentials. Finally, all three $\bar{K}N$ interaction models reveal a bump in $\pi\Sigma$ elastic cross-sections near the $\Lambda(1405)$ resonance mass $M_{\Lambda(1405)}^{PDG} = 1405.1_{-1.0}^{+1.3}$ MeV, according to the Particle Data Group [15].

3.2 Nucleon-nucleon interaction

Two-term Separable New potential (TSN) of nucleon-nucleon interaction was constructed in [2]. It has a form

$$V_{NN}^{\text{TSN}}(k, k') = \sum_{m=1}^2 g_m(k) \lambda_m g_m(k') \quad (4)$$

with form-factors

$$g_m(k) = \sum_{n=1}^3 \frac{\gamma_{mn}}{(\beta_{mn})^2 + k^2} \quad \text{for } m = 1, 2. \quad (5)$$

and parameters λ_m , γ_{mn} , β_{mn} fitted to phase shifts of Argonne V18 potential [16]. The potential reproduces triplet and singlet scattering lengths a and gives effective ranges r_{eff}

$$a_{np}^{\text{TSN}} = -5.400 \text{ fm}, \quad r_{\text{eff},np}^{\text{TSN}} = 1.744 \text{ fm}, \quad (6)$$

$$a_{pp}^{\text{TSN}} = 16.325 \text{ fm}, \quad r_{\text{eff},pp}^{\text{TSN}} = 2.792 \text{ fm}. \quad (7)$$

Deuteron binding energy given by the V_{NN}^{TSN} is $E_{\text{deu}} = 2.22 \text{ MeV}$.

3.3 $\Sigma N - \Lambda N$ interaction models

A simple separable model of the $\Sigma N - \Lambda N$ potential was used in our previous calculations [2]

$$V_{I,S}^{\Sigma N}(k, k') = \lambda_{I,S}^{\Sigma N} g_{I,S}^{\Sigma N}(k) g_{I,S}^{\Sigma N}(k'), \quad g_{I,S}^{\Sigma N}(k) = \frac{1}{(k^2 + \beta_{I,S}^{\Sigma N})^2} \quad (8)$$

with parameters $\lambda_{I,S}^{\Sigma N}$ and $\beta_{I,S}^{\Sigma N}$ fitted to not very accurate experimental cross-sections [17, 18, 19, 20, 21]. The interaction strongly depends on the two-body isospin. The ΣN interaction model in isospin $I = 3/2$ state is a one-channel one, it has real strength parameters $\lambda_{I,S}^{\Sigma N}$. Isospin $I = 1/2$ state ΣN system is coupled to the ΛN channel, therefore a coupled-channel $\Sigma N - \Lambda N$ potential with real parameters $\lambda_{I,S}^{\Sigma N}$ was constructed. Since the lowest $\pi \Lambda N$ channel is not directly included in the three-body calculations with two coupled $\bar{K} NN - \pi \Sigma N$ channels, the exact optical version of the coupled-channel potential $V_{I=1/2,S}^{\Sigma N}$ was used. The strength parameters $\lambda_{I=1/2,S}^{\Sigma N}$ of the exact optical potential turn to energy dependent functions. On the contrary, the three-body calculations with three coupled $\bar{K} NN - \pi \Sigma N - \pi \Lambda N$ channels were performed with the original version of the $I = 1/2$ potential with coupled $\Sigma N - \Lambda N$ channels.

New fits of the $\Sigma N - \Lambda N$ potential parameters Eq.(8) were performed to study dependence of the $K^- pp$ pole position on the $\Sigma N - \Lambda N$ interaction. Two new sets are spin-dependent and spin-independent ones fitted to the experimental cross-sections [17, 18, 19, 20, 21]. Two other sets were additionally fitted to scattering lengths of the different charge states of ΣN and ΛN systems taken from an "advanced" potential [22]. All together four new sets of parameters Eq.(8) were obtained for spin-dependent (denoted as "SDep") or spin-independent ("SInd") potentials with or without fits to scattering length ("ScL" or "noScL").

3.4 πN potential

It is known that s -wave πN interaction is weak. Due to this the πN potential $V^{\pi N}$ was neglected in our previous calculations of the $\bar{K} NN$ systems [2]. To check

dependence of the three-body results on the pion nucleon interaction, a one-term separable πN potential was constructed

$$V_I^{\pi N}(k, k') = \lambda_I^{\pi N} g_I^{\pi N}(k) g_I^{\pi N}(k'), \quad g_I^{\pi N}(k) = \frac{1}{(k^2 + \beta_I^{\pi N})^2}. \quad (9)$$

First, four sets of $\lambda_I^{\pi N}$ and $\beta_I^{\pi N}$ parameters (denoted as "1a", "1b", "2a", and "2b") were arbitrarily chosen with the only condition that there are no bound states in the πN system:

$$\beta_{1/2}^{\pi N} = \beta_{3/2}^{\pi N} = 3 \text{ fm}^{-1}, \quad \lambda_{1/2}^{\pi N} = \lambda_{3/2}^{\pi N} = -0.1 \text{ (1a)}, \quad = 0.1 \text{ (1b)}, \quad (10)$$

$$\beta_{1/2}^{\pi N} = \beta_{3/2}^{\pi N} = 1.5 \text{ fm}^{-1}, \quad \lambda_{1/2}^{\pi N} = \lambda_{3/2}^{\pi N} = -0.03 \text{ (2a)}, \quad = 0.03 \text{ (2b)}. \quad (11)$$

Two more sets of the strength parameters $\lambda_I^{\pi N}$ were fitted to the isospin $I = 1/2$ and $I = 3/2$ scattering lengths extracted from experimental data [23]

$$a_{\pi N, I=1/2}^{\text{Exp}} = 0.26 \text{ fm}, \quad a_{\pi N, I=3/2}^{\text{Exp}} = -0.11 \text{ fm}, \quad (12)$$

while the corresponding range parameters $\beta_I^{\pi N}$ were chosen by hand. These two sets of πN potential parameters Eq.(9) are denoted as "ScL-1" and "ScL-2":

$$\beta_{I=1/2}^{\pi N} = \beta_{3/2}^{\pi N} = 3 \text{ fm}^{-1}, \quad \lambda_{1/2}^{\pi N} = -0.627 \text{ fm}, \quad \lambda_{3/2}^{\pi N} = 0.456 \text{ fm} \text{ (ScL - 1)} \quad (13)$$

$$\beta_{I=1/2}^{\pi N} = \beta_{3/2}^{\pi N} = 1.5 \text{ fm}^{-1}, \quad \lambda_{1/2}^{\pi N} = -0.046 \text{ fm}, \quad \lambda_{3/2}^{\pi N} = 0.026 \text{ fm} \text{ (ScL - 2)} \quad (14)$$

4 Three-body Faddeev-type AGS equations and antikaon-nucleon potentials with three coupled particle channels

It was written in Sect.2 that due to the coupling of the $\bar{K}N$ system with $\pi\Sigma$ channel, the three-body Faddeev-type AGS equations were solved for two coupled particle channels: $\bar{K}NN$ and $\pi\Sigma N$. However, antikaon-nucleon interaction is coupled to the lowest $\pi\Lambda$ channel as well. If we include the third, $\pi\Lambda$ channel into three-body equations directly, the sum on γ index goes up to 5:

$$X_{ij}^{\alpha\beta}(z) = \delta_{\alpha\beta} Z_{ij}^{\alpha} + \sum_{k=1}^3 \sum_{\gamma=1}^5 Z_{ik}^{\alpha} \tau_k^{\alpha\gamma} X_{kj}^{\gamma\beta} \quad (15)$$

since now there are five particle channels

$$\alpha = 1 : |\bar{K}_1 N_2 N_3\rangle, \quad \alpha = 2 : |\pi_1 \Sigma_2 N_3\rangle, \quad \alpha = 3 : |\pi_1 N_2 \Sigma_3\rangle, \quad (16)$$

$$\alpha = 4 : |\pi_1 \Lambda_2 N_3\rangle, \quad \alpha = 5 : |\pi_1 N_2 \Lambda_3\rangle. \quad (17)$$

Antikaon-nucleon potentials with three coupled $\bar{K}N - \pi\Sigma - \pi\Lambda$ channels should be used in the system of integral equations derived from Eq.(15). The previously used chirally motivated antikaon-nucleon potential already couples all three channels. We refitted parameters of the potential to have strong poles corresponding to the $\Lambda(1405)$ resonance closer to other chiral models.

The $\pi\Lambda$ channel was taken in our phenomenological $V^{1,\text{SIDD}}$, $V^{2,\text{SIDD}}$ potentials into account indirectly through imaginary part of the complex strength parameter $\lambda_{I=1}^{11}$. We constructed new versions of the phenomenological potentials

directly coupling all three particle channels $\bar{K}N$, $\pi\Sigma$, and $\pi\Lambda$. As before, the $V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{1,\text{SIDD}}$ interaction model give one pole corresponding to the $\Lambda(1405)$ resonance, while the $V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{2,\text{SIDD}}$ potential lead to two strong poles for the resonance. All three new antikaon-nucleon $\bar{K}N - \pi\Sigma - \pi\Lambda$ potentials reproduce $1s$ level shift and width of kaonic hydrogen and low energy cross-sections of $K^-p \rightarrow K^-p$ and $K^-p \rightarrow MB$ reactions. They also reproduce two recently measured accurate data on $K^-p \rightarrow \pi^0\Sigma^0$ and $K^-p \rightarrow \pi^0\Lambda$ cross-sections [24]. In contrast to the previous versions, the new phenomenological potentials directly reproduce all three threshold branching ratios γ , R_c , and R_n . The same does the chirally motivated model with the new set of parameters. Physical characteristics of the three new antikaon-nucleon potentials are shown in Table 1.

	$V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{1,\text{SIDD}}$	$V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{2,\text{SIDD}}$	$V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{\text{Chiral}}$	Exp
ΔE_{1s}	-322.6	-323.5	-311.6	$-283 \pm 36 \pm 6$ [6]
Γ_{1s}	645.4	633.8	605.8	$541 \pm 89 \pm 22$ [6]
E_1	$1429.5 - i 35.0$	$1430.9 - i 41.6$	$1429.6 - i 33.2$	–
E_2	–	$1380.4 - i 79.9$	$1367.8 - i 66.5$	–
γ	2.35	2.36	2.36	2.36 ± 0.04 [13, 14]
R_c	0.666	0.664	0.664	0.664 ± 0.011 [13, 14]
R_n	0.190	0.189	0.190	0.189 ± 0.015 [13, 14]
a_{K^-p}	$-0.77 + i 0.97$	$-0.78 + i 0.95$	$-0.75 + i 0.90$	–

Table 1 Physical characteristics of the three $\bar{K}N - \pi\Sigma - \pi\Lambda$ potentials: $1s$ level shift ΔE_{1s} and width Γ_{1s} , strong poles E_1 and E_2 , threshold branching ratios γ , R_c , R_n . The K^-p scattering length a_{K^-p} was calculated with physical masses in all channels and Coulomb interaction in K^-p channel. Experimental data are also shown.

5 Results obtained with two coupled $\bar{K}NN - \pi\Sigma N$ channels

The results of the three-body $\bar{K}NN - \pi\Sigma N$ calculations with the new $\Sigma N - \Lambda N$ potentials are shown in Table 2. The three models of the antikaon-nucleon interaction were used: one- $V_{\bar{K}N}^{1,\text{SIDD}}$ and two-pole $V_{\bar{K}N}^{2,\text{SIDD}}$ phenomenological potentials and the chirally motivated potential $V_{\bar{K}N}^{\text{Chiral}}$. The πN interaction was switched off, as in our previous calculations. The first line contains our previous results [2] obtained with the older $\Sigma N - \Lambda N$ interaction model, denoted as $V_{\Sigma N-\Lambda N}^{\text{Prev}}$. The binding energies of the K^-pp quasi-bound state B_{K^-pp} and its widths Γ_{K^-pp} were calculated with the four new sets of parameters of the hyperon-nucleon interaction model: spin-dependent ("SDep") or spin-independent ("SInd") ones with ("ScL") or without ("noScL") fit to hyperon-nucleon scattering lengths from the "advanced" potential [22].

It is seen from Table 2 that the dependence of the three-body results on the hyperon-nucleon potential can be quite strong, especially when the phenomenological $\bar{K}N - \pi\Sigma$ potentials are used. Influence of the $V_{\Sigma N-\Lambda N}$ being used together

with the chirally motivated $\bar{K}N - \pi\Sigma(-\pi\Lambda)$ potential is much weaker. Surprisingly, not only width of the K^-pp state is influenced by the $\Sigma N - \Lambda N$ interaction, but its binding energy also can be changed sufficiently. Probably it is caused by the fact, that some of the four hyperon-nucleon potentials are characterized by very strong attraction in the $\Sigma N/\Lambda N$ pairs.

The most reliable hyperon-nucleon model among the four is the spin-dependent potential $V_{\Sigma N - \Lambda N}^{\text{ScL,SDep}}$ with parameters fitted to experimental cross-sections and scattering lengths. The resulting binding energies calculated with all three antikaon-nucleon potentials turned out to be smaller than the initial ones calculated with the previous $V_{\Sigma N - \Lambda N}^{\text{Prev}}$. The difference is much larger when the phenomenological models of $V_{\bar{K}N - \pi\Sigma}$ potential are used. The widths are changed slightly.

Dependence of the results of the three-body calculations with two coupled $\bar{K}NN - \pi\Sigma N$ channels on pion-nucleon interaction is shown in Table 3. Binding energy B_{K^-pp} and width Γ_{K^-pp} of the K^-pp quasi-bound state obtained using six new πN potentials are compared with the previously calculated results with switched off πN potential, shown at the first line. All old and new results were evaluated using the best hyperon-nucleon $V_{\Sigma N - \Lambda N}^{\text{ScL,SDep}}$ potential. The four versions of the πN potential with arbitrary parameters $V_{\pi N}^{(1a)}$, $V_{\pi N}^{(1b)}$, $V_{\pi N}^{(2a)}$, and $V_{\pi N}^{(2b)}$ lead to negligible changes in the binding energy obtained with every of the three $\bar{K}N$ potentials. The dependence of the three-body widths on the πN interaction model is more visible, especially when one of the phenomenological antikaon-nucleon potentials is used. Four πN interaction models change the width by few MeV. Changes in the three-body results are stronger when the πN models fitted to scattering lengths $V_{\pi N}^{\text{ScL}-1}$ and $V_{\pi N}^{\text{ScL}-2}$ are used. However, the dependence of the K^-pp quasi-bound state binding energy and width on the πN interaction models is quite weak, it is much weaker than on the hyperon-nucleon interaction.

Comparing the first line of Table 2, containing the previous results from [2], with the two last lines of Table 3, we see the cumulative effect of the "less important" pion-nucleon and hyperon-nucleon interactions. It is seen that the results of the three-body calculations with two coupled $\bar{K}NN - \pi\Sigma N$ channels evaluated

	$V_{\bar{K}N - \pi\Sigma}^{1,\text{SIDD}}$		$V_{\bar{K}N - \pi\Sigma}^{2,\text{SIDD}}$		$V_{\bar{K}N - \pi\Sigma(-\pi\Lambda)}^{\text{Chiral}}$	
	B_{K^-pp}	Γ_{K^-pp}	B_{K^-pp}	Γ_{K^-pp}	B_{K^-pp}	Γ_{K^-pp}
$V_{\Sigma N - \Lambda N}^{\text{Prev}}$	52.2	67.1	46.6	51.2	29.4	46.4
$V_{\Sigma N - \Lambda N}^{\text{noScL,SInd}}$	63.5	96.5	52.2	61.0	28.3	49.5
$V_{\Sigma N - \Lambda N}^{\text{noScL,SDep}}$	52.5	67.0	46.1	49.8	29.6	46.8
$V_{\Sigma N - \Lambda N}^{\text{ScL,SInd}}$	38.1	48.9	35.4	41.1	29.5	39.3
$V_{\Sigma N - \Lambda N}^{\text{ScL,SDep}}$	34.3	65.6	34.7	53.4	27.9	42.8

Table 2 Dependence of the binding energy B_{K^-pp} (MeV) and width Γ_{K^-pp} (MeV) of the quasi-bound state in the K^-pp system on $\Sigma N - \Lambda N$ interaction models. Previous results evaluated with $V_{\Sigma N - \Lambda N}^{\text{Prev}}$ from [2] are also shown. All observables were obtained with switched off πN potential. Three-body calculations were performed with two coupled $\bar{K}NN - \pi\Sigma N$ channels.

with the chirally motivated model of antikaon-nucleon interaction are influenced by the $\Sigma N - \Lambda N$ and πN interactions slightly. These interactions in the lower channel are much more important together with the phenomenological models. Using of the accurate $V_{\Sigma N - \Lambda N}$ and $V_{\pi N}$ potentials lead to much smaller binding energies $B_{K^- pp}$ and larger widths $\Gamma_{K^- pp}$ of the $K^- pp$ system calculated with $V_{\bar{K}N}^{1,\text{SIDD}}$ and $V_{\bar{K}N}^{2,\text{SIDD}}$.

	$V_{\bar{K}N-\pi\Sigma}^{1,\text{SIDD}}$		$V_{\bar{K}N-\pi\Sigma}^{2,\text{SIDD}}$		$V_{\bar{K}N-\pi\Sigma(-\pi\Lambda)}^{\text{Chiral}}$	
	$B_{K^- pp}$	$\Gamma_{K^- pp}$	$B_{K^- pp}$	$\Gamma_{K^- pp}$	$B_{K^- pp}$	$\Gamma_{K^- pp}$
$V_{\pi N}^{\text{off}}$	34.3	65.6	34.7	53.4	27.9	42.8
$V_{\pi N}^{(1a)}$	34.6	66.4	34.9	54.2	27.7	43.3
$V_{\pi N}^{(1b)}$	34.1	64.8	34.4	52.7	28.0	42.4
$V_{\pi N}^{(2a)}$	34.6	69.5	35.1	56.6	27.1	43.9
$V_{\pi N}^{(2b)}$	33.9	62.3	34.1	50.8	28.4	41.6
$V_{\pi N}^{\text{ScL-1}}$	37.6	71.4	37.7	58.2	27.2	46.6
$V_{\pi N}^{\text{ScL-2}}$	35.7	70.5	35.9	57.4	27.0	44.6

Table 3 Dependence of the binding energy $B_{K^- pp}$ (MeV) and width $\Gamma_{K^- pp}$ (MeV) of the quasi-bound state in the $K^- pp$ system on πN interaction models. All observables were calculated with $V_{\Sigma N - \Lambda N}^{\text{ScL,SDep}}$ potential, including the previous ones from Table 2 with switched off pion-nucleon interaction $V_{\pi N}^{\text{off}}$. The three-body calculations were performed with two coupled $\bar{K}NN - \pi\Sigma N$ channels.

6 Results obtained with three coupled $\bar{K}NN - \pi\Sigma N - \pi\Lambda N$ channels

The results of the direct inclusion of the lowest $\pi\Sigma N$ channel in the $\bar{K}NN - \pi\Sigma N - \pi\Lambda N$ system are shown in Table 4. The calculations were performed with the newly constructed phenomenological potentials directly coupling all three particle channels $V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{1,\text{SIDD}}$ and $V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{2,\text{SIDD}}$ together with the chirally motivated potential $V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{\text{Chiral}}$ with refitted parameters. The new potentials are described in Section 4. The most accurate $V_{\Sigma N - \Lambda N}^{\text{ScL,SDep}}$ potential and the two versions of the πN interaction model $V_{\pi N}^{\text{ScL-1}}$, $V_{\pi N}^{\text{ScL-2}}$ were used in the calculations.

It is seen that the two-pole phenomenological and chiral potentials give approximately equal widths of the quasi-bound $K^- pp$ state, while the width obtained with the one-pole phenomenological potential is smaller. As before, the smallest $K^- pp$ binding energy is the result of using the chirally motivated potential, the difference between the one-pole and two-pole phenomenological results is now larger than before. Comparing the results of the three-body calculations with two coupled $\bar{K}NN - \pi\Sigma N$ in Table 3 and three coupled $\bar{K}NN - \pi\Sigma N - \pi\Lambda N$ channels in Table 4 we see large differences between the observables. However, it

is not clear, whether the difference is caused by the three-channel structure or by using different $\bar{K}N$ potentials leading to different two-body observables.

	$V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{1,\text{SIDD}}$		$V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{2,\text{SIDD}}$		$V_{\bar{K}N-\pi\Sigma-\pi\Lambda}^{\text{Chiral}}$	
	B_{K^-pp}	Γ_{K^-pp}	B_{K^-pp}	Γ_{K^-pp}	B_{K^-pp}	Γ_{K^-pp}
$V_{\pi N}^{\text{ScL}-1}$	34.5	52.0	42.9	60.4	26.8	59.6
$V_{\pi N}^{\text{ScL}-2}$	34.3	52.1	40.2	57.7	27.2	56.3

Table 4 Results of the three-body AGS calculations with three coupled $\bar{K}NN - \pi\Sigma N - \pi\Sigma\Lambda$ channels. Binding energy B_{K^-pp} (MeV) and width Γ_{K^-pp} (MeV) of the quasi-bound state in the K^-pp system were evaluated using $V_{\Sigma N-\Lambda N}^{\text{ScL},\text{SDep}}$ potential and two pion-nucleon interaction models $V_{\pi N}^{\text{ScL}-1}$ and $V_{\pi N}^{\text{ScL}-2}$.

7 Summary

Fine-tuning of the dynamically exact calculations of the binding energy and width of the K^-pp quasi-bound state was performed. Influence of the "less important" interactions in the lower $\pi\Sigma N$ channel was studied in three-body Faddeev-type calculations with two coupled $\bar{K}NN - \pi\Sigma N$ channels. It was found that the $\Sigma N - \Lambda N$ interaction model can change the three-body result quite strongly, while the dependence of the B_{K^-pp} and Γ_{K^-pp} on the πN potential is weaker. The new antikaon-nucleon potentials coupling three $\bar{K}N - \pi\Sigma - \pi\Lambda$ channels were constructed. The corresponding three-body calculations of the K^-pp quasi-bound state with three coupled $\bar{K}NN - \pi\Sigma N - \pi\Lambda N$ channels were performed. The three-body results evaluated with three coupled channels lead to different changes in binding energy and/or width in comparison with the calculations with two coupled channels depending on the particular model of antikaon-nucleon interaction. However, the reason for the differences: whether it is new potentials or the third coupled channel, - should be additionally studied.

References

1. T. Yamaga, T.: Experimental study of $\bar{K}NN$ and future experiments for kaonic nuclei. EPJ Web Conf. 271, 07001 (2022)
2. Shevchenko, N.V.: On a quasi-bound state in the K^-d system caused by strong interactions. Few. Body. Syst. 61, 27 (2020)
3. Alt, E.O., Grassberger, P., Sandhas, W.: Reduction of the three-particle collision problem to multi-channel two-particle Lippmann-Schwinger equations. Nucl. Phys. B 2, 167 (1967)
4. Shevchenko, N.V.: Near-threshold K^-d scattering and properties of kaonic deuterium. Nucl. Phys. A 890-891, 50-61 (2012)
5. Shevchenko, N.V., Révai, J.: Faddeev calculations of the $\bar{K}NN$ system with chirally-motivated $\bar{K}N$ interaction. I. Low-energy K^-d scattering and antikaonic deuterium. Phys. Rev. C 90, 034003 (2014)
6. Bazzi, M. *et al.* (SIDDHARTA Collaboration): A new measurement of kaonic hydrogen X-rays. Phys. Lett. B 704, 113 (2011)

7. Sakitt, M. et al.: Low-energy K^- -meson interactions in hydrogen. *Phys. Rev.* 139, B 719 (1965)
8. Kim, J.K.: Low-energy K^- -p interaction and interpretation of the 1405-MeV Y_0^* resonance as a $\bar{K}N$ bound state. *Phys. Rev. Lett.* 14, 29 (1965)
9. Kim, J.K.: Multichannel phase-shift analysis of $\bar{K}N$ interaction in the region 0 to 550 MeV/c. *Phys. Rev. Lett.* 19, 1074 (1967)
10. Kittel, W., Otter, G., Wacek, I.: The K^- proton charge exchange interactions at low energies and scattering lengths determination. *Phys. Lett.* 21, 349 (1966)
11. Ciborowski, J. et al.: Kaon scattering and charged Sigma hyperon production in K^-p interactions below 300 MeV/c. *J. Phys. G* 8, 13 (1982)
12. Evans, D. et al.: Charge-exchange scattering in K^-p interactions below 300 MeV/c. *J. Phys. G* 9, 885 (1983)
13. Tovee, D.N. et al.: Some properties of the charged Σ hyperons. *Nucl. Phys. B* 33, 493 (1971)
14. Nowak, R.J. et al.: Charged Σ hyperon production by K^- meson interactions at rest. *Nucl. Phys. B* 139, 61 (1978)
15. Workman, R.L. et al. (Particle Data Group): The Review of Particle Physics (2022). *Prog. Theor. Exp. Phys.* 2022, 083C01 (2022) and 2023 update.
16. Wiringa, R.B., Stoks, V.G.J., Schiavilla, R.: Accurate nucleon-nucleon potential with charge-independence breaking. *Phys. Rev. C* 51, 38 (1995)
17. Alexander, G. et al.: Study of the $\Lambda - N$ system in low-energy $\Lambda - p$ elastic scattering. *Phys. Rev.* 173, 1452 (1968)
18. Sechi-Zorn, B., Kehoe, B., Twitty, J., Burnstein, R.A.: Low-energy Λ -proton elastic scattering. *Phys. Rev.* 175, 1735 (1968)
19. Eisele, F. et al.: Elastic $\Sigma^\pm p$ scattering at low energies. *Phys. Lett. B* 37, 204 (1971)
20. Engelmann, R., Filthuth, H., Hepp, V., Kluge, E.: Inelastic Σ^-p -interactions at low momenta. *Phys. Lett.* 21, 587 (1966)
21. Hepp, V., Schleich, M.: A new determination of the capture ratio $r_c = \frac{\Sigma^-p \rightarrow \Sigma^0 n}{(\Sigma^-p \rightarrow \Sigma^0 n) + (\Sigma^-p \rightarrow \Lambda^0 n)}$, the Λ^0 -lifetime and the $\Sigma^- - \Lambda^0$ mass difference. *Z. Phys.* 214, 71 (1968)
22. J. Haidenbauer, J. et al.: Hyperon-nucleon interaction in chiral effective field theory at next-to-next-to-leading order. *arXiv nucl-th/2301.00722* (2022)
23. Hirtl, A. et al.: Redetermination of the strong-interaction width in pionic hydrogen. *Eur. Phys. J. A* 57, 70 (2021)
24. Piscicchia, K. et al.: First simultaneous $K^-p \rightarrow (\Sigma^0/\Lambda)\pi^0$ cross sections measurements at 98 MeV/c. *arXiv: nucl-ex/2210.10342* (2022)