Implications of the Weak Gravity Conjecture on Charge, Kinetic Mixing, the Photon Mass, and More

Fayez Abu-Ajamieh*

Centre for High Energy Physics, Indian Institute of Science, Bangalore 560012, India

Nobuchika Okada[†]

Department of Physics and Astronomy; University of Alabama; Tuscaloosa; Alabama 35487; USA

Sudhir K. Vempati[‡]

Centre for High Energy Physics, Indian Institute of Science, Bangalore 560012, India

We investigate several phenomenological implications of the Weak Gravity Conjecture (WGC). We find that the WGC implies that the SM neutrinos must be electrically neutral, that the electric charge in the SM must be quantized, and that the photon must be massless. In addition, we use the WGC to set lower bounds on the electric charge of milli-charged particles (mCP), the gauge coupling of several U(1) extensions of the SM, their kinetic mixing parameter with the SM $U(1)_{\rm EM}$, and the axion couplings to photons and fermions. We also set an upper bound on the lifetime of the proton.

I. INTRODUCTION

One of the fundamental questions in Nature is why gravity is the weakest force amongst the four fundamental interactions. The Weak Gravity Conjecture (WGC) [1] tries to address the question and has attracted a lot of attention in recent years. A major implication of the conjecture is that any U(1) gauge theory with a coupling g, must breakdown at a scale Λ below the Planck scale, such that $\Lambda \sim gM_{\rm Pl}$, where $M_{\rm Pl} = 2.4 \times 10^{18}$ GeV. More concretely, consider a U(1) gauge group with coupling g. The electric WGC implies that there must be a light charged particle with mass

$$m_{\rm el} \lesssim g_{\rm el} M_{\rm Pl},$$
 (1)

which also should hold for magnetic monopoles, i.e.

$$m_{\rm mag} \lesssim g_{\rm mag} M_{\rm Pl} \sim \frac{1}{g_{\rm el}} M_{\rm Pl}.$$
 (2)

As the monopole has a mass that is at least of the order of the magnetic field it generates, which is linearly divergent, then one has

$$m_{\rm mag} \sim \frac{\Lambda}{g_{\rm el}^2},$$
 (3)

plugging eq. (3) in eq. (2), we thus arrive at the magnetic WGC

$$\Lambda \lesssim g M_{\rm Pl}.$$
 (4)

This result implies that any U(1) gauge theory has a natural cutoff where it breaks down. For instance, for

 $U(1)_{\rm EM}$, the WGC suggests $\Lambda \sim 10^{17}$ GeV, close to the scale of the heterotic string. We shall see how eq. (4) arises naturally when applying the WGC in its simplest form.

A major motivation for this conjecture lies with the fact that the existence of a small charge g without a corresponding light mass m as dictated by eq. (1), means that black holes carrying such charges cannot evaporate via the Hawking radiation, which in turn will lead to the problematic issue of remnants [2].

The result in eq. (4) is quite powerful, as instead of predicting a cutoff scale for a U(1) theory, it can be used to set a lower bound on g if the cutoff scale Λ is known. In this paper, we shall show that the magnetic WGC has several deep phenomenological implications on various U(1) gauge groups, including the electric neutrality of neutrinos and the electric charge quantization of the SM; setting lower bounds on the electric charge of milli-Charged Particles (mCPs), the charges of U(1) extension of the SM and the kinetic mixing parameter between U(1) and $U(1)_{\rm EM}$; the masslessness of the photon, and setting lower bounds on the axion's couplings to photons and fermions and an upper bound on the proton's lifetime.

This paper is organized as follows: In Section II, we illustrate the application of the (magnetic) WGC and how to extract the cutoff scale Λ to be used with it. In Section III we investigate the various phenomenological implications of the WGC, and then we present our conclusions in Section IV. We relegate some technical details to the Appendix.

II. APPLYING THE WGC

Here we show how to apply the WGC to extract a lower limit on the coupling g of a U(1) gauge group. Consider the s-channel scattering $X\overline{X} \to X\overline{X}$. Gravi-

^{*} favezajamieh@iisc.ac.in

[†] okadan@ua.edu

[‡] vempati@iisc.ac.in



FIG. 1. Applying the WGC.

tational interaction proceeds through the exchange of a graviton, whereas the U(1) interaction proceeds through the exchange of the corresponding gauge boson. The WGC implies that the gravitational interaction should be weaker than the U(1) interaction, as shown in Figure 1. This implies that

$$|\mathcal{M}_{\text{grav}}(s)| \lesssim |\mathcal{M}_{\text{U}(1)}(s)|. \tag{5}$$

Since at high energy $\mathcal{M}_{\mathrm{U}(1)} \sim g^2$ and $\mathcal{M}_{\mathrm{grav}} \sim s/M_{\mathrm{Pl}}^2$, and if we set set $\sqrt{s} = \Lambda$ as the cutoff scale, then we immediately arrive at eq. (4).

Notice that if Λ is known, or at least if a lower bound on it is known, then one can set a lower bound on g. Strictly speaking, Λ is interpreted as the scale at which the corresponding QFT breaks down. In our calculation, we will be mainly concerned with the coupling of U(1) to SM fermions and photons. For the former, we can extract a lower limit on Λ from the results of LEP, whereas for the latter, we can use the results of Light-By-Light (LBL) scattering in the LHC.

A. The Scale of New Physics (NP) from LEP

LEP applied the technique presented in [3] on the differential cross-section of $e^+e^- \rightarrow l^+l^-, q^+q^ (l = e, \mu, \tau$ and q = u, d; in order to set limits on the scale of NP arising from the operator [4]

$$\mathcal{L}_{\text{eff}} = \frac{g^2}{(1+\delta)\Lambda_{\pm}^2} \sum_{i,j=L,R} \eta_{ij}(\bar{e}_i \gamma_\mu e_i)(\bar{f}_j \gamma^\mu f_j), \quad (6)$$

where $\delta = 1(0)$ for f = e $(f \neq e)$ respectively, $\eta_{ij} \pm 1, 0$, $\Lambda_+(\Lambda_-)$ for constructive (destructive) interference with the SM, and by convention $g^2/4\pi \to 1$.

Defining $\epsilon_{\pm} \equiv 1/\Lambda_{\pm}^2$ with $\epsilon = 0$ corresponding to the SM, the limits on ϵ_{\pm} were used to set a 95% C.L. limit on Λ_{\pm} of ~ 10 TeV. This scale is defined as the lower scale where QFT breaks down if the NP is heavy such that it can be integrated out, i.e., if the NP is of the decoupling type, then we can set $\Lambda \equiv \Lambda_{\text{LEP}} \simeq 10$ TeV. On the other hand, if the NP is light (e.g. axions), then this scale cannot be used as the NP in non-decoupling. Instead, we can drop the corresponding mass and set $\Lambda = \sqrt{s_{\text{LEP}}} = 209$ GeV as the lowest scale where QFT breaks down.

B. The Scale of NP from Light-By-Light (LBL) Scattering

The LEP bound extracted above cannot be used for certain processes when the scattering particles are photons. Instead, the experimental results from LBL scattering should be used. Here, we follow the treatment in [5, 6]. The effective Lagrangian for LBL scattering can be expressed as:

$$\mathcal{L}_{\text{eff}} \supset c_1 F_{\mu\nu} F^{\mu\nu} F_{\rho\sigma} F^{\rho\sigma} + c_2 F_{\mu\nu} F^{\nu\rho} F_{\rho\sigma} F^{\mu\sigma}, \qquad (7)$$

from which the differential cross-section of LBL scattering at LO is obtained

$$\frac{d\sigma}{d\Omega} = \frac{1}{16\pi^2 \hat{s}} (\hat{s}^2 + \hat{t}^2 + \hat{s}\hat{t})^2 (48c_1^2 + 11c_2^2 + 40c_1c_2), \quad (8)$$

where $\hat{s} = m_{\gamma\gamma}^2$ and $\hat{t} = -\frac{\hat{s}}{2}(1 - \cos\theta)$. Eq. (8) can be immediately integrated to yield the total cross-section

$$\sigma_{\gamma\gamma} = \frac{7\hat{s}^3}{40\pi} (48c_1^2 + 11c_2^2 + 40c_1c_2). \tag{9}$$

Notice that $c_{1,2}$ are model-dependent parameters that have the dimension of $1/(\text{mass})^4$. For example, for the Born-Infeld (BI) model, $c_1 = -1/32\beta^2$ and $c_2 = 1/8\beta^2$, where β is the BI scale parameter. To set a conservative limit, we follow the scaling of LEP above and set

$$c_{1,2} \equiv \frac{g_i^4}{\Lambda^4} \to \frac{(4\pi)^2}{\Lambda^4}.$$
 (10)

The latest experimental results of the LBL scattering from ATLAS are given by [7]

$$\sigma_{\gamma\gamma}^{\text{Exp}} = 70 \pm 24 \text{ (stat.)} \pm 17 \text{ (sys.) nb;}$$
 (11)

setting the theoretical uncertainty at 10 nb as in [5] and summing the uncertainties in quadrature, we get $\delta\sigma^{\text{Exp}} = 31.0644$ nb. Demanding that the cross-section in eq. (9) be less than twice this experimental uncertainty and using the scaling in eq. (10), we can estimate the scale of NP in LBL processes @ 2σ C.L. as

$$\Lambda_{\rm LBL} \gtrsim \left(\frac{22176\pi^3 \hat{s}^3}{10\delta\sigma^{\rm Exp}}\right)^{1/8} \simeq 7.8 \text{ TeV}, \qquad (12)$$

where we have used $\sqrt{\hat{s}} = m_{\gamma\gamma} = 5.02 \text{ TeV} [7].$

III. PHENOMENOLOGICAL IMPLICATIONS OF THE WGC

In this section, we discuss several phenomenological implications of the WGC based mainly on eqs. (4) and (5).

A. Bounds on the Electric Charge

The WGC can be used to set a lower bound on the electric charge. It is well-known that milli-charges can be introduced to the SM in a variety of ways, such as adding an $SU(2) \times SU(2)$ Dirac fermion with hypercharge 2ϵ , [8], an additional U(1) [9], allowing neutrinos to have millicharge [10, 11], or through introducing non-locality [12, 13]. For an electric milli-charge ϵe , the WGC in eq. (4) sets the lower bound $|\epsilon| \gtrsim \Lambda/(eM_{\rm Pl})$, which using $\Lambda_{\rm LEP} = 10$ TeV translates to $|\epsilon| \gtrsim 1.4 \times 10^{-14}$. On the other hand, globular cluster star cooling sets an upper bound of $|\epsilon| \leq 2 \times 10^{-14}$ [14]. Thus we have

$$1.4 \times 10^{-14} \lesssim |\epsilon| \lesssim 2 \times 10^{-14}.$$
 (13)

For the particular case of SM neutrinos, it can be shown that if they are Dirac fermions, then their charges must be equal [10, 11, 19]

$$Q_{\nu_e} = Q_{\nu_{\mu}} = Q_{\nu_{\tau}}.$$
 (14)

Matter neutrality sets an upper bound on their electric charge of [15–19]

$$|Q_{\nu_e}| \lesssim 3.2 \times 10^{-21} e,$$
 (15)

which in conjunction with the bound from the WGC, and in light of eq. (14) implies that all neutrinos must be electrically neutral. Since in the SM model, charge cannot de dequantized without inducing milli-charge in neutrinos [10, 11, 19], this also implies that in the SM, charge must be quantized.¹ Thus, we arrive at the following important result:

The WGC, in conjunction with experimental limits, implies that the electric charge in the SM is quantized.

Before we conclude this section, we point out that we can use eq. (2) to set a lower bound on the magnetic charge of neutrinos (if such a charge exists)

$$Q_{\nu_{\rm mag}} \gtrsim \frac{m_{\nu}}{M_{\rm Pl}} \simeq 4.2 \times 10^{-29},\tag{16}$$

where we've used $m_{\nu} = 0.1$ eV. On the other hand, an upper bound on the magnetic charge of neutrinos can be obtained from magnetars [22]. Both bounds imply

$$4.2 \times 10^{-29} \lesssim Q_{\nu_{\rm mag}} \lesssim 10^{-16}.$$
 (17)

B. Implication for Gauged U(1) Extensions

The WGC can also be used to set bounds on additional gauged U(1) interactions. In particular, a gauged U(1) has a corresponding gauge boson Z', and the process $e^+e^- \rightarrow Z' \rightarrow \overline{f}f$ yields an effective interaction identical to the one in eq. (6). Thus, the LEP results can be used in conjunction with the WGC to set lower bounds on their gauge coupling g'.

The only non-anomalous U(1) gauge group extensions are B - L, $L_{\mu} - L_e$, $L_{\tau} - L_e$ and $L_{\tau} - L_{\mu}$, and we can see that the LEP bound can be applied to the first 3, whereas the LEP bound cannot be used to bound the last as it doesn't interact with electrons at tree-level. The interaction Lagrangians can be expressed as

$$\mathcal{L}_{B-L} \supset g' Z'_{\mu} \Big(\frac{1}{3} (\overline{u} \gamma^{\mu} u + \overline{d} \gamma^{\mu} d) - \overline{e} \gamma^{\mu} e - \overline{\mu} \gamma^{\mu} \mu - \overline{\tau} \gamma^{\mu} \tau \Big),$$

$$\mathcal{L}_{\mu-e} \supset -g' Z'_{\mu} \Big(\overline{e} \gamma^{\mu} e - \overline{\mu} \gamma^{\mu} \mu \Big),$$

$$\mathcal{L}_{\tau-e} \supset -g' Z'_{\mu} \Big(\overline{e} \gamma^{\mu} e - \overline{\tau} \gamma^{\mu} \tau \Big).$$
(18)

For $m_{Z'} \gg \sqrt{s_{\text{LEP}}} = 209$ GeV, Z' can be integrated out to yield an operator identical to eq. (6) with $m_{Z'} = \Lambda_{\text{LEP}} = 10$ TeV. Thus, the WGC implies

$$\frac{s}{M_{\rm Pl}^2} \lesssim \frac{g'^2 s}{m_{Z'}^2},\tag{19}$$

which for $m_{Z'} = 10$ TeV yields the bound

$$g' \gtrsim 4.2 \times 10^{-15}$$
. (20)

On the other hand, if $m_{Z'} \ll \sqrt{s_{\text{LEP}}}$, then it can be neglected to yield an operator identical to eq. (6) with $\Lambda = \sqrt{s_{\text{LEP}}}$. In this case the WGC implies

$$g' \gtrsim \frac{\sqrt{s_{\rm LEP}}}{M_{\rm Pl}} \simeq 8.7 \times 10^{-17}.$$
 (21)

We can use the experimental bounds on the $g' - m_{Z'}$ parameter space to recast this bound to become a bound on $m_{Z'}$. For B - L, the limits in [23] can be used to translate the bound in eq. (21) to become

$$m_{Z'} \gtrsim 10^{-2} \text{ eV},$$
 (22)

which arises from the cooling of Horizontal Branch (HB) stars, whereas the latest bounds on $L_{\mu} - L_e$ and $L_{\tau} - L_e$ from [24] indicate that no lower bound on the mass can be set.

Another important result is that the bounds in eqs. (20) and (21) exclude the viability of an extra U(1) as a solution to the hierarchy problem. In [25], it was proposed to use an extra U(1) gauge group in conjunction with the WGC to furnish a solution to the hierarchy problem. The argument rested on the electric form of the WGC, which postulates the existence of a light particle with mass given by eq. (1). More specifically, if neutrinos are Dirac fermions, then they obtain their mass from the Higgs VEV

$$m_{\nu} \sim y_{\nu} v,$$
 (23)

¹ If neutrinos are Majorana fermions, then they must be neutral [20, 21], which also implies that charge is quantized in the SM.



FIG. 2. $e^+e^- \rightarrow \overline{f}f$ through kinetic mixing.

which according to the WGC should be $m_{\nu} \leq g M_{\rm Pl}$ with g being the coupling of some extra U(1) gauge group. This implies that

$$\frac{v}{M_{\rm Pl}} \lesssim \frac{g}{y_{\nu}},\tag{24}$$

and a large hierarchy between v and $M_{\rm Pl}$ can be obtained if $g \ll y_{\nu}$. However, if we use $y_{\nu} \sim 10^{-12}$, then generating the hierarchy between the EW scale and the Planck scale would require

$$g \sim \frac{(10^{-12})(246)}{2.4 \times 10^{18}} \sim 10^{-28},$$
 (25)

which is many orders of magnitudes below the limits in eqs. (20) and (21). This excludes the viability of this solution to the hierarchy problem.²

C. Bounds on Kinetic Mixing

In addition to setting bounds on the gauge coupling of any additional U(1) gauge group, the WGC can also set bounds on the kinetic mixing parameter of an extra U(1) with the SM $U(1)_{\text{EM}}$. In general, the kinetic term of $U(1) \times U(1)_{\text{EM}}$ can be written as

$$\mathcal{L}_{\rm kin} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{4} X_{\mu\nu} X^{\mu\nu} - \frac{1}{2} \chi F_{\mu\nu} X^{\mu\nu}.$$
 (26)

To apply the WGC with the LEP bound, we need two insertions of the kinetic mixing, as shown in Figure (2). Like the case with Z', the amplitude and bound depends on whether the gauge boson X is light or heavy.

For $m_X \ll \sqrt{s_{\text{LEP}}}$, the amplitude of this process is given by $\mathcal{M} \sim e^2 \chi^2$, which according to the WGC sets the following lower bound on the kinetic mixing parameter

$$\chi \gtrsim \frac{\sqrt{s_{\rm LEP}}}{eM_{\rm Pl}} \simeq 2.9 \times 10^{-16}.$$
 (27)

Other experimental bounds on the kinetic mixing parameters can be found in [26]. For gauge boson masses up

to ~ 10 keV , the strongest bounds arise from the cooling of Red Giant (RG) stars, which is equal to ~ 6×10^{-14} . Thus, we can set the following bound

$$2.9 \times 10^{-16} \lesssim \chi \lesssim 6 \times 10^{-14}$$
 for $m_X \lesssim 10$ keV. (28)

On the other hand, for $m_X \gg \sqrt{s_{\text{LEP}}}$, the amplitude is give by $\mathcal{M} \sim e^2 \chi^2 s / m_X^2$, which according to the WGC sets the following lower bound on the kinetic mixing parameter

$$\chi \gtrsim \frac{\Lambda}{eM_{\rm Pl}} \simeq 1.4 \times 10^{-14},$$
 (29)

where $\Lambda = m_X = 10$ TeV. For $m_X = 10$ TeV, there is no upper bond on χ . For an earlier analysis using magnetic form of the WGC, please see [27].

D. Bound on Axion Couplings

The WGC can also be used to set lower bounds on several axion couplings. The interaction Lagrangian of axions with photons and fermions can be written as

$$\mathcal{L}_{\rm int} = -\frac{1}{4} g_{a\gamma} a F_{\mu\nu} \tilde{F}^{\mu\nu} + \frac{g_{af}}{2m_f} (\partial_\mu a) (\overline{f} \gamma^\mu \gamma^5 f), \quad (30)$$

where $\tilde{F}^{\mu\nu} = \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}$. The WGC can be used to set bounds on $g_{a\gamma}$ by comparing the amplitude of the LBL scattering through the exchange of an axion with that through the exchange of a graviton. The former is given by $\mathcal{M}_{\text{LBL}(a)} \sim g_{a\gamma}^2 s$, whereas the latter is given by $\mathcal{M}_{\text{LBL}(G)} \sim s/M_{\text{Pl}}^2$. This implies that

$$g_{g\gamma} \gtrsim \frac{1}{M_{\rm Pl}},$$
 (31)

which is reminiscent of the Axion Weak Gravity Conjecture [28–41]. Things are more interesting when we consider the axion's coupling to fermions. If we consider the process $e^+e^- \rightarrow a \rightarrow \overline{f}f$, then we can use the WGC in conjunction with the LEP results to set a lower bound on the couplings g_{af} . Notice here that since axions are light, then the cutoff scale we use is $\Lambda = \sqrt{s_{\text{LEP}}}$. The amplitude of the process is given by $\mathcal{M} \sim g_{ae}g_{af}$, which implies that

$$g_{ae}g_{af} \gtrsim \frac{s_{\rm LEP}^2}{M_{\rm Pl}^2}.$$
(32)

For the particular case of f = e, this bound reads $g_{ae} \gtrsim 8.7 \times 10^{-17}$. This, together with the bound from cooling of RG stars [42] implies that

$$8.7 \times 10^{-17} \lesssim g_{ae} \lesssim 1.3 \times 10^{-13}.$$
 (33)

For $f = \mu, \tau, u, d$, the lower limit on g_{ae} can be used in eq. (32) to set a conservative lower bound on g_{af}

$$g_{af} \gtrsim \frac{(209)^2}{(2.4 \times 10^{18})^2 (8.7 \times 10^{-17})} \simeq 8.7 \times 10^{-17}.$$
 (34)

² Although the bounds in eqs. (20) and (21) do not hold for the gauge group $L_{\tau} - L_{\mu}$ since the LEP bound does not apply here, using any reasonably low QED cutoff, (even ~ 1 GeV), would exclude the viability of this gauge group as well as a solution to the hierarchy problem.

This bound also applies to the axion's coupling to protons and neutrons, since the LEP bound also applies to $e^+e^- \rightarrow \overline{u}u/\overline{d}d$. Combined with the upper limits on g_{ap} and g_{an} from the cooling of supernovae [43], the bounds read

$$8.7 \times 10^{-17} \lesssim g_{ap} \lesssim 1.3 \times 10^{-9},$$
 (35)

$$8.7 \times 10^{-17} \lesssim g_{an} \lesssim 1.5 \times 10^{-9}.$$
 (36)

E. Implications of the WGC on the Mass of the Photon

Although in the SM the photon is massless, it is possible to give it a small mass while still remaining within experimental bounds. In general, a massless U(1) gauge boson can acquire mass in a gauge-invariant way either via the Stueckelberg mechanism or the Higgs mechanism,³ and for the latter case, the photon can be Higgsed either by the SM Higgs or by a Beyond the SM (BSM) Higgs. Below, we discuss the implication of the WGC for each case individually.

1. Photon Mass via the Stueckelberg Mechanism

The photon can acquire mass via the gauge-invariant Stueckelberg action

$$\mathcal{L}_{\text{Stueck}} = -\frac{1}{2} F^{\dagger}_{\mu\nu} F^{\mu\nu} - (\partial^{\mu} A^{\dagger}_{\mu} + mB^{\dagger}) (\partial^{\nu} A_{\nu} + mB) + m^2 \Big(A^{\dagger}_{\mu} - \frac{1}{m} \partial_{\mu} B^{\dagger} \Big) \Big(A^{\mu} - \frac{1}{m} \partial^{\mu} B \Big).$$
(37)

A simple calculation shows that the photon will acquire the following mass [44, 45]

$$m_A^2 = \frac{1}{8} (g^2 + g'^2) v^2 \Big[1 + \epsilon - \sqrt{1 - 2\epsilon \Big(\frac{g^2 - g'^2}{g^2 + g'^2}\Big) + \epsilon^2} \Big],$$

$$\simeq m^2 \cos^2 \theta_W, \qquad (38)$$

where θ_W is the Weinberg angle, and

$$\epsilon = \frac{4m^2}{v^2(g^2 + g'^2)} \ll 1. \tag{39}$$

A consequence of the photon acquiring a Stueckelberg mass is that it will interact with neutrinos at tree-level, i.e., neutrinos will acquire an electric charge

$$Q_{\nu} = -\frac{1}{2}g'\cos\theta_W + \frac{1}{2}g\sin\theta_W \simeq \frac{g\sin\theta_W m_A^2}{2M_W^2},\quad(40)$$

where M_W is the mass of the W boson. Given that the upper limit on the photon mass reads [46–48]

$$m_A \lesssim 1.8 \times 10^{-14} \text{ eV},^4$$
 (41)

it is easy to show that the corresponding induced neutrino charge is excluded by the limit in eq. (15). Put differently, we have shown in Section III A that the WGC implies that neutrinos must be electrically neutral. Therefore, this implies that the possibility of the photon acquiring mass via the Stueckelberg is excluded.

2. Photon Mass via the SM Higgs

It is also possible for the photon to acquire mass via the SM Higgs mechanism.⁵ In this case, the photon can obtain mass via the kinetic term of the Higgs

$$\mathcal{L}_{\rm SM}^A \supset (D_\mu H)^{\dagger} (D^\mu H), \tag{42}$$

where $D_{\mu} = \partial_{\mu} + ieQA_{\mu}$. Inserting the Higgs doublet in the unitary gauge, we can write Lagrangian explicitly as

$$\mathcal{L}_{\rm SM}^A \supset \frac{1}{2} \Big(e^2 Q^2 v^2 + 2e^2 Q^2 v h + e^2 Q^2 v^2 h^2 \Big) A_\mu A^\mu,$$
(43)

and we see now that the photon acquires a mass $m_A = eQv$. Moreover, tree-level cubic and quartic interactions between the Higgs and the photon are induced.

In order to use the WGC to set a lower limit on the photon mass via the SM Higgs, we can utilize the LBL scattering via the exchange of a Higgs and compare it to that via the exchange of a graviton. For $\sqrt{s} \gg m_h$, the former yields $\mathcal{M}_{\text{LBL}} \sim m_A^4/v^2 s$, which when compared to the graviton exchange yields the limit

$$m_A \gtrsim \Lambda_{\rm LBL} \sqrt{\frac{v}{M_{\rm Pl}}} \simeq 79 \ {\rm keV},$$
 (44)

which is clearly excluded. Thus, we conclude that the WGC excludes the possibility of a photon mass via the SM Higgs.

3. Photon Mass via a BSM Higgs

It is also possible for the photon to acquire a mass via a BSM Higgs ϕ . This case is identical to the SM case, except now the VEV v_{ϕ} is a free parameter.⁶

³ It is possible for the photon to acquire mass through the Proca action, however, this action is not gauge-invariant.

⁴ This limit is model-independent. A stronger limit of $\sim 10^{-18}$ eV can be obtained with some assumptions [49, 50].

⁵ In fact, the Higgs mechanism was originally introduced as a mechanism to give mass to the photon. In this letter, we do not concern ourselves with how Elecroweak Symmetry Breaking (EWSB) is modified when the photon is Higgsed.

⁶ Obviously a BSM Higgs will mix with the SM Higgs at the level of the potential, however, we can set the mixing to be very small to avoid any experimental constraints, since we are only interested in giving mass to the photon.

Applying the WGC yields a condition similar to eq. (44) with $v_{\rm SM} \rightarrow v_{\phi}$. As such, it is not possible to set a limit on the photon mass directly. However, one can recast eq. (44) to set a lower bound on the coupling of ϕ to photons by setting

$$\frac{2m_A^2}{v_\phi} \equiv g_{\phi\gamma} \gtrsim \frac{2\Lambda_{\rm LBL}}{M_{\rm Pl}} \simeq 5.1 \times 10^{-11} \text{ GeV.}$$
(45)

On the other hand, one can set an upper bound on $g_{\phi\gamma}$ from the cooling of HB stars (see the Appendix)

$$g_{\phi\gamma} \lesssim 2 \times 10^{-38} \text{ GeV},$$
 (46)

which clearly excludes the possibility of a photon mass via a BSM Higgs as well. Thus, the results of this section show that:

The WGC, in conjunction with the experimental limits, implies that the photon is strictly massless.

F. Implications of the WGC on the Proton Lifetime

It is well-known that Grand Unified Theories (GUTs) [51–53] lead to proton decay [54–60] $p^+ \rightarrow e^+ \pi^0$ due to the exchange of a leptoquark X. The lifetime of the proton decay is given by

$$\tau_{p^+ \to e^+ \pi^0} \sim \frac{M_X^4}{m_p^5},$$
(47)

where M_X is the mass of the leptoquark and m_p is the mass of the proton. Since the X is the gauge boson, the WGC can be applied here as well. The WGC (eq. (1)) implies that the upper bound of the mass of X is $M_{\rm Pl}$. This translates into an upper bound of the lifetime of the proton

$$\tau_{p^+ \to e^+ \pi^0} \lesssim \frac{M_{\rm Pl}^4}{m_p^5} \sim 10^{42} \text{ yr.}$$
 (48)

Unfortunately, this bound is much weaker than the current bound of ~ 10^{36} yr for non-SUSY SU(5) and ~ 10^{39} yr for SUSY SU(5) that is calculated in [61, 62].

IV. CONCLUSIONS

In this paper, we have shown that the WGC, in conjunction with the relevant experimental limits, can have many important phenomenological implications, including the electric neutrality of neutrinos, charge quantization of the SM, and the masslessness of photons. In addition, we have used the WGC to set lower limits on the electric charge of mCPs, the coupling of several BSM U(1) gauge groups and their kinetic mixing parameter with the SM U(1) gauge group, and the coupling of the axion to photons and fermions. These limits can be used in synergy with the limits from experiment to define the viable range of these couplings that is bound both from above and below. Other set of complementarity bounds can come from theoretical constraints like positivity conditions in the presence of gravity[63].

The profound implications of the WGC provide motivation for further efforts study it in more depth, including trying to prove or disprove it.

ACKNOWLEDGMENTS

The work of FA is supported by the C.V. Raman fellowship from CHEP at IISc. The work on NO is supported in part by the United States Department of Energy.

Appendix A: Setting an Upper Bound on $g_{\phi\gamma}$ from the Cooling of HB Stars

Here we show how we obtain the bound in eq. (46) from the cooling HB stars. But before doing so, let's try to set a limit on m_{ϕ} . First, notice that we can use the upper limit on the photon mass in (eq. 41) with eq. (44) to set an upper bound on the BSM VEV

$$v_{\phi} \lesssim 1.3 \times 10^{-26} \text{ eV},$$
 (A1)

where we have used Λ_{LBL} . If we assume that the quartic coupling of ϕ is λ_{ϕ} , such that $m_{\phi} = \sqrt{2\lambda_{\phi}}v_{\phi}$, then we can also set an upper bound on m_{ϕ} by assuming that λ_{ϕ} saturates the perturbativity limit of 4π . Thus we obtain

$$m_{\phi} \lesssim 6.4 \times 10^{-26} \text{ eV}.$$
 (A2)

Thus, we see that any BSM Higgs that gives mass of the photon must be extremely light, justifying the assumption of small mixing with the SM Higgs. The dominant process in HB cooling is the Primakoff (or inverse Primakoff) production shown in Figure 3.



FIG. 3. Primakoff production.

Assuming that the electron is non-relativistic and neglecting its recoil energy, both of which are reasonable assumptions given the conditions of the HB medium and the smallness of m_{ϕ} , the cross-section of the Primakoff production is given by

$$\sigma(\omega) \simeq \frac{\alpha g_{\phi\gamma}^2}{16m_e^2 \omega^2} \left[\log\left(\frac{4\omega^2}{m_A^2 + m_\phi^2}\right) + \frac{2m_e^2}{m_A^2 + m_\phi^2} \right], \quad (A3)$$

where ω is the photon energy. The energy loss rate per unit volume can be expressed as [64]

$$Q = 2n_e \int \frac{d^3\vec{k}}{(2\pi)^3} \frac{\omega\sigma(\omega)}{e^{\omega/T} - 1},$$
 (A4)

where n_e is the electron number density and T is the temperature of the star. To set a conservative limit on $g_{\phi\gamma}$, we assume m_A to be the largest possible value allowed by experiment (saturating the upper limit in eq. (41)). This implies that $m_{\phi} \ll m_A$, so we drop m_{ϕ} in our calculation. Thus, plugging eq. (A3) in eq. (A4), the upper bound on $g_{\phi\gamma}$ can be expressed as

$$g_{\phi\gamma} \lesssim \sqrt{\frac{8\pi^2 m_e^2 m_u \dot{\varepsilon}_{\max}}{\alpha y_e I(\omega)}},$$
 (A5)

where m_u is the atomic mass unit, $y_e = n_e m_u / \rho$ is the electron fraction in the star, $\dot{\varepsilon}_{\text{max}} = Q / \rho$ is the maximum energy loss rate per unit mass, ρ is the density of the star, and the integral $I(\omega)$ is given by

$$I(\omega) = \int_0^\infty d\omega \left(\frac{\omega}{e^{\omega/T} - 1}\right) \left[\log\left(\frac{2\omega}{m_A}\right) + \frac{m_e^2}{m_A^2}\right].$$
 (A6)

For HB stars, $\dot{\varepsilon}_{\rm max} = 10 \text{ erg s}^{-1} \text{ g}^{-1}$. Using $T = 10^8 \text{ K}$, $\rho = 10^4 \text{ g cm}^{-3}$ and $y_e = 0.5$; and evaluating the integral in eq. (A6) numerically, we arrive at the upper bound:

$$g_{\phi\gamma} \lesssim 2 \times 10^{-38} \text{ GeV.}$$
 (A7)

This bound, however, is only valid in the free streaming regime, i.e., when ϕ in the Primakoff process, or the photon in the inverse Primakoff process, stream freely out of the star. To verify the validity of this assumption, we need to check both the decay length and the mean free path.

To find the decay length, let's assume that ϕ only couples to the photon at tree-level. Given the upper limit on the mass of ϕ , we can see that it can only decay to a pair of photons if $m_{\phi} \geq 2m_A$. Otherwise ϕ would be stable. Assuming that ϕ can decay to a pair of photons, the decay width is given by

$$\Gamma_{\phi} = \frac{1}{\tau_{\phi}} = \frac{2\lambda_{\phi}m_A^4}{\pi m_{\phi}^3},\tag{A8}$$

where we have assumed ϕ has a quartic coupling λ_{ϕ} . The decay length is given by

$$L_{\phi} = \tau_{\phi} \beta \gamma c, \tag{A9}$$

where $\gamma = 1/\sqrt{1-\beta^2} = E_{\phi}/m_{\phi}$. Assuming that ϕ is relativistic, its average energy is given by

$$\langle E_{\phi} \rangle = \frac{\pi^4}{30\xi(3)}T \simeq 2.701T.$$
 (A10)

To get a conservative limit, we set $\lambda = 4\pi$ and use the upper limit on m_{ϕ} . This yields a lower limit on the decay length of ~ $O(10^{43})$ Km, which is many orders of magnitude larger than the typical radius of an HB star of ~ 10^9 Km. Thus, ϕ does not decay inside the star.

Next, we evaluate the mean free path. The mfp can be expressed as

$$mfp = \frac{1}{n_e \sigma(\omega)}.$$
 (A11)

Using $\langle \omega \rangle \simeq 2.701T$, we find that mph $\simeq 4 \times 10^{18}$ Km, which is also many orders of magnitude larger than radius of an HB star, thereby justifying our free streaming assumption.

- N. Arkani-Hamed, L. Motl, A. Nicolis and C. Vafa, "The String landscape, black holes and gravity as the weakest force," JHEP 06, 060 (2007) hep-th/0601001.
- [2] L. Susskind, "Trouble for remnants," hep-th/9501106.
- [3] E. Eichten, K. D. Lane and M. E. Peskin, "New Tests for Quark and Lepton Substructure," Phys. Rev. Lett. 50, 811-814 (1983)
- S. Schael *et al.* [ALEPH, DELPHI, L3, OPAL and LEP Electroweak], "Electroweak Measurements in Electron-Positron Collisions at W-Boson-Pair Energies at LEP," Phys. Rept. 532, 119-244 (2013) hep-ex/1302.3415.
- [5] J. Ellis, N. E. Mavromatos and T. You, "Lightby-Light Scattering Constraint on Born-Infeld Theory," Phys. Rev. Lett. 118, no.26, 261802 (2017) hep-ph/1703.08450.
- [6] J. Ellis, N. E. Mavromatos, P. Roloff and T. You, "Lightby-light scattering at future e⁺e⁻ colliders," Eur. Phys. J. C 82, no.7, 634 (2022) hep-ph/2203.17111.

- [7] M. Aaboud *et al.* [ATLAS], "Evidence for light-by-light scattering in heavy-ion collisions with the ATLAS detector at the LHC," Nature Phys. **13**, no.9, 852-858 (2017) hep-ex/1702.01625.
- [8] L. B. Okun, M. B. Voloshin and V. I. Zakharov, "ELEC-TRICAL NEUTRALITY OF ATOMS AND GRAND UNIFICATION MODELS," Phys. Lett. B 138, 115-120 (1984)
- [9] B. Holdom, "Two U(1)'s and Epsilon Charge Shifts," Phys. Lett. B 166, 196-198 (1986)
- [10] R. Foot, G. C. Joshi, H. Lew and R. R. Volkas, "Charge quantization in the standard model and some of its extensions," Mod. Phys. Lett. A 5, 2721-2732 (1990)
- [11] R. Foot, H. Lew and R. R. Volkas, "Electric charge quantization," J. Phys. G 19, 361-372 (1993) [erratum: J. Phys. G 19, 1067 (1993)] hep-ph/9209259.
- [12] A. Capolupo, G. Lambiase and A. Quaranta, "Muon g-2 anomaly and non-locality," Phys. Lett. B 829, 137128

(2022) hep-ph/2206.06037.

- [13] F. Abu-Ajamieh, P. Chattopadhyay, A. Ghoshal and N. Okada, "Anomalies in String-inspired Non-local Extensions of QED," hep-th/2307.01589.
- [14] S. Davidson, S. Hannestad and G. Raffelt, "Updated bounds on millicharged particles," JHEP 05, 003 (2000) hep-ph/0001179.
- [15] M. Marinelli and G. Morpurgo, "The Electric Neutrality of Matter: A Summary," Phys. Lett. B 137, 439-442 (1984)
- [16] G. G. Raffelt, "Limits on neutrino electromagnetic properties: An update," Phys. Rept. 320, 319-327 (1999)
- [17] C. Giunti and A. Studenikin, "Neutrino electromagnetic interactions: a window to new physics," Rev. Mod. Phys. 87, 531 (2015) hep-ph/1403.6344.
- [18] G. Bressi, G. Carugno, F. Della Valle, G. Galeazzi, G. Ruoso and G. Sartori, "Testing the neutrality of matter by acoustic means in a spherical resonator," Phys. Rev. A 83, no.5, 052101-1-052101-14 (2011) physics.atom-ph/1102.2766.
- [19] A. Das, D. Ghosh, C. Giunti and A. Thalapillil, "Neutrino charge constraints from scattering to the weak gravity conjecture to neutron stars," Phys. Rev. D 102, no.11, 115009 (2020) hep-ph/2005.12304.
- [20] K. S. Babu and R. N. Mohapatra, "Is There a Connection Between Quantization of Electric Charge and a Majorana Neutrino?," Phys. Rev. Lett. 63, 938 (1989)
- [21] K. S. Babu and R. N. Mohapatra, "Quantization of Electric Charge From Anomaly Constraints and a Majorana Neutrino," Phys. Rev. D 41, 271 (1990)
- [22] A. Hook and J. Huang, "Bounding millimagnetically charged particles with magnetars," Phys. Rev. D 96, no.5, 055010 (2017) hep-ph/1705.01107.
- [23] J. Heeck, "Unbroken B L symmetry," Phys. Lett. B 739, 256-262 (2014) hep-ph/1408.6845.
- [24] M. Bauer, P. Foldenauer and J. Jaeckel, "Hunting All the Hidden Photons," JHEP 07, 094 (2018) hep-ph/1803.05466.
- [25] N. Craig, I. Garcia Garcia and S. Koren, "The Weak Scale from Weak Gravity," JHEP 09, 081 (2019) hep-ph/1904.08426.
- [26] M. Ahlers, J. Jaeckel, J. Redondo and A. Ringwald, "Probing Hidden Sector Photons through the Higgs Window," Phys. Rev. D 78, 075005 (2008) hep-ph/0807.4143.
- [27] K. Benakli, C. Branchina and G. Lafforgue-Marmet, Eur. Phys. J. C 80 (2020) no.12, 1118 doi:10.1140/epjc/s10052-020-08691-4 [arXiv:2007.02655 [hep-ph]].
- [28] T. Rudelius, "Constraints on Axion Inflation from the Weak Gravity Conjecture," JCAP 09, 020 (2015) hep-th/1503.00795.
- [29] M. Montero, A. M. Uranga and I. Valenzuela, "Transplanckian axions!?," JHEP 08, 032 (2015) hep-th/1503.03886.
- [30] J. Brown, W. Cottrell, G. Shiu and P. Soler, "Fencing in the Swampland: Quantum Gravity Constraints on Large Field Inflation," JHEP 10, 023 (2015) hep-th/1503.04783.
- [31] B. Heidenreich, M. Reece and T. Rudelius, "Weak Gravity Strongly Constrains Large-Field Axion Inflation," JHEP 12, 108 (2015) hep-th/1506.03447.
- [32] A. de la Fuente, P. Saraswat and R. Sundrum, "Natural Inflation and Quantum Gravity," Phys. Rev. Lett. 114,

no.15, 151303 (2015) hep-th/1412.3457.

- [33] A. Hebecker, P. Mangat, F. Rompineve and L. T. Witkowski, "Winding out of the Swamp: Evading the Weak Gravity Conjecture with F-term Winding Inflation?," Phys. Lett. B 748, 455-462 (2015) hep-th/1503.07912.
- [34] T. C. Bachlechner, C. Long and L. McAllister, "Planckian Axions and the Weak Gravity Conjecture," JHEP 01, 091 (2016) hep-th/1503.07853.
- [35] T. Rudelius, "On the Possibility of Large Axion Moduli Spaces," JCAP 04, 049 (2015) hep-th/1409.5793.
- [36] D. Junghans, "Large-Field Inflation with Multiple Axions and the Weak Gravity Conjecture," JHEP 02, 128 (2016) hep-th/1504.03566.
- [37] K. Kooner, S. Parameswaran and I. Zavala, "Warping the Weak Gravity Conjecture," Phys. Lett. B 759, 402-409 (2016) hep-th/1509.07049.
- [38] L. E. Ibanez, M. Montero, A. Uranga and I. Valenzuela, "Relaxion Monodromy and the Weak Gravity Conjecture," JHEP 04, 020 (2016) hep-th/1512.00025.
- [39] A. Hebecker, F. Rompineve and A. Westphal, "Axion Monodromy and the Weak Gravity Conjecture," JHEP 04, 157 (2016) hep-th/1512.03768.
- [40] A. Hebecker and P. Henkenjohann, "Gauge and gravitational instantons: From 3-forms and fermions to Weak Gravity and flat axion potentials," JHEP 09, 038 (2019) hep-th/1906.07728.
- [41] T. Daus, A. Hebecker, S. Leonhardt and J. March-Russell, "Towards a Swampland Global Symmetry Conjecture using weak gravity," Nucl. Phys. B 960, 115167 (2020) hep-th/2002.02456.
- [42] F. Capozzi and G. Raffelt, "Axion and neutrino bounds improved with new calibrations of the tip of the red-giant branch using geometric distance determinations," Phys. Rev. D 102, no.8, 083007 (2020) astro-ph.SR/2007.03694.
- [43] M. Buschmann, C. Dessert, J. W. Foster, A. J. Long and B. R. Safdi, "Upper Limit on the QCD Axion Mass from Isolated Neutron Star Cooling," Phys. Rev. Lett. 128, no.9, 091102 (2022) hep-ph/2111.09892.
- [44] S. V. Kuzmin and D. G. C. McKeon, "Stueckelberg mass in the Glashow-Weinberg-Salam model," Mod. Phys. Lett. A 16, 747-753 (2001)
- [45] H. Ruegg and M. Ruiz-Altaba, "The Stueckelberg field," Int. J. Mod. Phys. A 19, 3265-3348 (2004) hep-th/0304245.
- [46] X. F. Wu, S. B. Zhang, H. Gao, J. J. Wei, Y. C. Zou, W. H. Lei, B. Zhang, Z. G. Dai and P. Mészáros, "Constraints on the Photon Mass with Fast Radio Bursts," Astrophys. J. Lett. 822, no.1, L15 (2016) astro-ph.HE/1602.07835.
- [47] L. Bonetti, J. Ellis, N. E. Mavromatos, A. S. Sakharov, E. K. G. Sarkisyan-Grinbaum and A. D. A. M. Spallicci, "Photon Mass Limits from Fast Radio Bursts," Phys. Lett. B 757, 548-552 (2016) astro-ph.HE/1602.09135.
- [48] L. Bonetti, J. Ellis, N. E. Mavromatos, A. S. Sakharov, E. K. Sarkisyan-Grinbaum and A. D. A. M. Spallicci, "FRB 121102 Casts New Light on the Photon Mass," Phys. Lett. B 768, 326-329 (2017) astro-ph.HE/1701.03097.
- [49] A. S. Goldhaber and M. M. Nieto, "Photon and Graviton Mass Limits," Rev. Mod. Phys. 82, 939-979 (2010) hep-ph/0809.1003.
- [50] L. C. Tu, J. Luo and G. T. Gillies, "The mass of the

photon," Rept. Prog. Phys. 68, 77-130 (2005)

- [51] J. C. Pati and A. Salam, "Unified Lepton-Hadron Symmetry and a Gauge Theory of the Basic Interactions," Phys. Rev. D 8, 1240-1251 (1973)
- [52] H. Georgi and S. L. Glashow, "Unified weak and electromagnetic interactions without neutral currents," Phys. Rev. Lett. 28, 1494 (1972)
- [53] H. Fritzsch and P. Minkowski, "Unified Interactions of Leptons and Hadrons," Annals Phys. 93, 193-266 (1975)
- [54] S. Weinberg, "Baryon and Lepton Nonconserving Processes," Phys. Rev. Lett. 43, 1566-1570 (1979)
- [55] F. Wilczek and A. Zee, "Operator Analysis of Nucleon Decay," Phys. Rev. Lett. 43, 1571-1573 (1979) doi:10.1103/PhysRevLett.43.1571
- [56] S. Weinberg, "Varieties of Baryon and Lepton Nonconservation," Phys. Rev. D 22, 1694 (1980)
- [57] S. Weinberg, "Supersymmetry at Ordinary Energies. 1. Masses and Conservation Laws," Phys. Rev. D 26, 287 (1982)
- [58] N. Sakai and T. Yanagida, "Proton Decay in a Class of Supersymmetric Grand Unified Models," Nucl. Phys. B 197, 533 (1982)

- [59] S. Dimopoulos, S. Raby and F. Wilczek, "Proton Decay in Supersymmetric Models," Phys. Lett. B 112, 133 (1982)
- [60] J. R. Ellis, D. V. Nanopoulos and S. Rudaz, "GUTs 3: SUSY GUTs 2," Nucl. Phys. B 202, 43-62 (1982)
- [61] I. Dorsner, P. Fileviez Perez and R. Gonzalez Felipe, "Phenomenological and cosmological aspects of a minimal GUT scenario," Nucl. Phys. B 747, 312-327 (2006) hep-ph/hep-ph/0512068.
- [62] I. Dorsner, P. Fileviez Perez and G. Rodrigo, "Fermion masses and the UV cutoff of the minimal realistic SU(5)," Phys. Rev. D 75, 125007 (2007) hep-ph/hep-ph/0607208.
- [63] K. Aoki, T. Q. Loc, T. Noumi and J. Tokuda, Phys. Rev. Lett. **127** (2021) no.9, 091602 doi:10.1103/PhysRevLett.127.091602 [arXiv:2104.09682 [hep-th]];

T. Q. Loc, [arXiv:2312.09132 [hep-th]].

[64] G. G. Raffelt, "Stars as laboratories for fundamental physics: The astrophysics of neutrinos, axions, and other weakly interacting particles," 1996, ISBN 978-0-226-70272-8