On Mössbauer rotor effect, clock synchronization and third postulate of relativity

 February 17, 2024

 Christian Corda

 UNY Polytechnic Institute, 13502 Utica, New York, USA; Istituto Livi, 59100 Prato, Prato, USA; Istituto Livi, 59100 Prato, Prato, USA; Istituto Livi, 59100 Prato, Prato, USA; Istica Science Centre, Adarsh Nagar, Hyderabad 500063, India. E-mail:

 Spirla Science Centre, Adarsh Nagar, Hyderabad 500063, India. E-mail:

 Cordac.galilei@gmail.com

 Data

 The Mössbauer rotor effect recently gained a renewed interest due to the discovery and explanation of an additional effect of clock synchronization which has been missed for about 50 years, i.e. starting from a famous book of Pauli, till some more recent experimental analyses. The theoretical explanation of such an additional effect of clock synchronization can be calculated in another way via the third postulate of relativity.

 In the general relativistic framework, Einstein grasped that the gravitational field can be represented via

In the general relativistic framework, Einstein grasped that the gravitational field can be represented via space-time curvature in a paper which analysed the rotating frame, verbatim [1]:

"The following important argument also speaks in favor of a more relativistic interpretation. The centrifugal force which acts under given conditions of a body is determined precisely by the same natural constant that also gives its action in a gravitational field. In fact we have no means to distinguish a centrifugal field from a gravitational field. We thus always measure as the weight of the body on the surface of the earth the superposed action of both fields, named above, and we cannot separate their actions. In this manner the point of view to interpret the rotating system K' as at rest, and the centrifugal field as a gravitational field, gains justification by all means. This interpretation is reminiscent of the original (more special) relativity where the pondermotively acting force, upon an electrically charged mass which moves in a magnetic field, is the action of the electric field which is found at the location of the mass as seen by the reference system at rest with the moving mass."

Such an interpretation enabled various general relativistic treatments of Mössbauer rotor [2-10] and Sagnac effects [11], via the Einstein Equivalence Principle (EEP) which states the equivalence between gravitation and inertia [4–11]. The EEP indeed includes the rotating reference frame [4–10]. The discovery and explanation of a new effect, neglected for over 50 years since Pauli's famous book on relativity [2], has led to a recent new interest in the Mössbauer rotor effect. The correct theoretical explanation of this new effect, which is added to the traditionally known one, is due to a series of works in which the analysis was carried out with a general relativistic treatment [4–10]. In this letter this additional effect of clock synchronization will be calculated in a simple way via the third postulate of relativity.

This effect is due to the German physicist R. Mössbauer in 1958 [12], and involves gamma rays' resonant and recoil-free emission and absorption, without loss of energy, by atomic nuclei bound in a solid. The particular case of the *Mössbauer rotor effect* (see Figure 1), shows an absorber orbiting around the radiation's source. In Figure 1 it is assumed that the z - axis is perpendicular to the plane of the apparatus within the circumference and that the same apparatus rotates around such a z - axis having constant angular velocity ω . The final detector in the right of the Figure is assumed at rest. The idea is to measure the so-called *transverse Doppler effect* via the fractional energy shift for the resonant absorber [3–10], because the transverse Doppler effect involves a relative energy shift between emission and absorption lines through its motion.

Kündig developed a fundamental Mössbauer rotor experiment in 1963 [3]. The shift of the 14.4-keV Mössbauer absorption line of Fe^{57} was measured in function of the angular velocity of the rotor [3]. The result appeared to be consistent with the theory of relativity with a sensitivity of 1.1%. Kündig also discussed potential systematic errors [3]. Kündig's experiment has been recently investigated by an experimental group [13, 14]. In the first work [13], the original data of Kündig's paper have been reanalysed. Because of the strangeness of the results in [13], the cited experimental group realized an independent Mössbauer rotor experiment [14]. In [13], it was shown that mistakes were present in Kündig's original data. After correcting those mistakes, the Authors found a higher value of the fractional energy shift [13]

$$\frac{\Delta E}{E} = -k\frac{v^2}{c^2},\tag{0.1}$$



Figure 1 The scheme of the Mössbauer rotor experiment in this Figure is adapted from [14]. One assumes that the z-axis is perpendicular to the plane of the Figure and that the apparatus within the circumference rotates around such a z-axis having constant angular velocity ω . The final detector in the right of the Figure results at rest.

with $k = 0.596 \pm 0.006$ where Kündig [3], Pauli [2] and others found k = 0.5 in agreement with the relativity theory. This was a strange issue, but here it will be shown that a correct relativistic analysis can solve it. We recall that the Authors of [13] found that: i) the deviation of the coefficient k Eq. (0.1) from 0.5 is much higher than the measuring error (about 20 times); ii) that deviation did not arise from some kinds of disturbing factors like rotor vibrations because the excellent methodology applied by Kündig [3] surely excluded those potential disturbing factors. In [13] it was also emphasized that also the experiment in [15] appeared to be consistent with k > 0.5. Remarkably, in the experiment in [15] much more data than similar Mössbauer rotor experiments analysed in [16–19] have been involved. Based on the strange results in [13], the experimental group decided to realize a new experiment [14], where neither the scheme of Kündig's experiment [3], nor the schemes of the other previously cited experiments [15–19] have been followed. This permitted them to get independent information, with respect to previous experiments, about the real value of k in Eq. (0.1). The final result of the value of k was $k = 0.68 \pm 0.03$ [14]. Hence, the experiment [14] confirmed that the coefficient k in Eq. (0.1) substantially exceeds 0.5. One can find the scheme of this Mössbauer rotor experiment in Figure 1, see [14] for details.

The puzzle has been solved by us in [4–8]. The key point is that in previous works in the literature [2, 3, 13–19] an important effect of clock synchronization of has not been considered for about 50 years, starting from the aforementioned book by Pauli on relativity [2], up to the recent experiments [13, 14].

The Langevin transformation in cylindrical coordinates is about moving from a Lorentz-Minkowski coordinate system, centered in the source of the apparatus in Figure 1, having the z - axis perpendicular to the plane of the Figure, to a second coordinate system rotating around the z - axis (Langevin frame). In the following the coordinates and the proper time of the Lorentz-Minkowski frame will be labelled with the subscript L, while the coordinates and the proper time of the observer in the Langevin frame will be labelled with the subscript R (R stays for rotating). The Lorentz-Minkowski frame is the laboratory frame, and the metric is [4–8, 20]

$$ds^{2} = c^{2}dt_{L}^{2} - dr_{L}^{2} - r_{L}^{2}d\phi_{L}^{2} - dz_{L}^{2}.$$
(0.2)

The Langevin transformation generates the Langevin reference frame $\{t_R, r_R, \phi_R, z_R\}$, having constant angular velocity ω around the z - axis. In details, one sets [4–8, 20]

$$t_L = t_R \quad r_L = r_R \quad \phi_L = \phi_R + \omega t_R \qquad z_L = z_R \ . \tag{0.3}$$

This leads to Langevin metric for the Langevin frame [4–8, 20]

$$ds^{2} = \left(1 - \frac{r_{R}^{2}\omega^{2}}{c^{2}}\right)c^{2}dt_{R}^{2} - 2\omega r_{R}^{2}d\phi_{R}dt_{R} - dr_{R}^{2} - r_{R}^{2}d\phi_{R}^{2} - dz_{R}^{2}.$$
(0.4)

Considering the EEP, the line element (0.4) is interpreted in the Einsteinian sense in terms of a stationary gravitational field [4–8, 20]. Technical details of the last sentence can be found in paragraph 89 of [20]. The EEP means that the inertial force experienced by a Langevin observer is interpreted in terms of a gravitational "force" [4–8]. The first well known effect arises from the "gravitational redshift" [4–8]. This effect is well known in previous literature [3–10, 12-19]. Kündig's standard interpretation [3] argues that the Langevin observer concludes that his clock is slowed down by the "gravitational potential". Hence, the clock of the observer in the laboratory must be faster than the clock of the Langevin observer. This is the so-called gravitational redshift". Then, the fractional energy shift in the laboratory results [3–10, 12-19]

$$\frac{E_2 - E_1}{E_1} = \frac{\triangle E_1}{E_1} = \simeq -\frac{1}{2} \frac{v^2}{c^2}.$$
(0.5)

Hence, one gets $k_1 = \frac{1}{2}$ as the contribution to k from the first effect [3–10, 12-19].

The key point for needing an additional effect is the following. The computation that permits to obtain Eq. (0.5) is usually performed in the Langevin reference frame. [4–8]. But the final detector is in motion with

respect to the Langevin observer |4-8|, see Figure 1. Thus, the clock of the fixed observer in the Lorentz-Minkowski laboratory frame is *not* synchronized with the clock of the Langevin observer. This implies that an additional effect must contribute to the total effect [4–8]. This effect of clock synchronization was not considered in previous literature on the subject [2, 3, 13–19]. The general relativistic approach in [7, 8] permitted us to explain this second effect in a complete way. Here we show that this additional effect can be obtained in another way via the third postulate of relativity [21], which states that, if one considers a physical system (the Langevin observer in the current case) moving through a flat space-time, than there is at any moment a local inertial system such that, in it, the system is at rest (the Langevin reference frame is locally inertial). In that case, at any instant, the coordinates and state of the system can be Lorentz transformed to the other system through some Lorentz transformation. This means that if one considers the frame of reference of the rotating Langevin observer, where the observer is at rest, the coordinates and state of the system can be Lorentz transformed to the Lorentz-Minkowski frame of reference of the laboratory, which moves with respect to the frame of reference of the Langevin observer. Relative to the Lorentz-Minkowski frame of reference of the laboratory, the instantaneous velocity of the rotating Langevin observer is $\vec{v}(t_L)$ with magnitude $|\vec{v}| = v$ bounded by the speed of light c, so that $0 \le v < c$. Here the time t_L is the time as measured in the Lorentz-Minkowski frame of reference of the laboratory, which corresponds with the proper time of the laboratory τ_{L} , it is not the time measured in the frame of reference of the rotating Langevin observer. The rotating Langevin frame of reference and the Lorentz-Minkowski frame of reference of the laboratory are Lorentz connected via the instantaneous Lorentz factor [21]

$$\gamma \equiv \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}.\tag{0.6}$$

Then, one gets via the definition of the instantaneous Lorentz factor

$$\frac{d\tau_L}{d\tau_R} = \gamma, \tag{0.7}$$

where τ_R is the proper time measured in the frame of reference of the rotating Langevin observer. Hence, there is a time dilation between the two frames of references given by

$$d\tau_L = \gamma d\tau_R,\tag{0.8}$$

which means that the proper time between two ticks as measured in the frame in which the clock is moving

(the Lorentz-Minkowski frame of reference of the laboratory), is longer than the proper time between these ticks as measured in the rest frame of the clock (the frame of reference of the rotating Langevin observer). In order to compute γ one must compute v. For a radial coordinate r_L it is obviously $v = \omega r$. Thus, one gets:

$$d\tau_R = \frac{d\tau_L}{\gamma} = \sqrt{1 - \frac{v^2}{c^2}} d\tau_L = \sqrt{1 - \frac{\omega^2 r_L^2}{c^2}} d\tau_L,$$
 (0.9)

which means that the clock of the rotating Langevin observer and the clock of the Lorentz-Minkowski observer in the laboratory measure the same proper time **if and only if** $r_L = r_R = 0$, i.e. only in the origin of the two reference frames. Thus, clock synchronization is necessary in all the other points. Hence, the contribution must be calculated along the whole trajectory of photons. Thus, one must find r_L as a function of the time τ_L in Eq. (0.9) for the light rays. Clearly, as r_L and τ_L refer to the Lorentz-Minkowski frame of reference of the laboratory it is obviously $r_L = c\tau_L$. Then, Eq. (0.9) becomes

$$d\tau_R = \sqrt{1 - \omega^2 \tau_L^2} d\tau_L. \tag{0.10}$$

One can approximate Eq. (0.10) with [7]

$$d\tau_R \simeq \left(1 - \frac{1}{2}\omega^2 \tau_L^2 + \dots\right) d\tau_L. \tag{0.11}$$

Eq. (0.11) takes into account the second effect of order $\frac{v^2}{c^2}$ to time dilation, which, integrated along the entire trajectory of the photons, gives

$$\Delta \tau_{RT} = \int_0^{\tau_{LT}} \left(1 - \frac{1}{2} \omega^2 \tau_L^2 \right) d\tau_L - \tau_{LT} = -\frac{1}{6} \tau_L^3 \omega^2 = -\frac{1}{6} r_{LT} \frac{v^2}{c^2}.$$
 (0.12)

The subscript T in the quantities in Eq. (0.12) stands for "trajectory". $\Delta \tau_{RT}$ represents the difference between the proper time that has been measured by the rotating Langevin observer and the proper time that has been measured by the fixed Lorentz-Minkowski observer and r_{LT} is the radial coordinate of the detector. Then, the additional effect of clock synchronization (at order $\frac{v^2}{c^2}$) to the energy shift is

$$z \equiv \frac{\Delta \tau_{RT}}{\tau_{LT}} = \frac{\Delta E_2}{E_1} = -k_2 \frac{v^2}{c^2} = -\frac{1}{6} \frac{v^2}{c^2}.$$
 (0.13)

This means $k_2 = \frac{1}{6}$.

Therefore, Eqs. (0.5) and (0.13) permit to obtain the total fractional energy shift as

$$\frac{\triangle E}{E_1} = \frac{\triangle E_1}{E_1} + \frac{\triangle E_2}{E_1} \simeq -\frac{2}{3} \frac{v^2}{c^2}.$$
(0.14)

This theoretical result is completely consistent with the experimental result $k = 0.68 \pm 0.03$ in [14] and with previous analyses in [4–10, 22], but in this letter it has been obtained via a new, different analysis, which uses the third postulate of relativity.

Here are some additional considerations. After our original discovery of the additional effect of clock synchronization in [4], this result has been independently confirmed by other Authors [9, 10, 22]. The experimental group that reanalysed the original data of Kündig [13] and realized the new experiment [14] also realized a second experiment with some additional collaborators [23]. In this new experiment they found the result $k = 0.69 \pm 0.02$ [23], which is still consistent with the theoretical prediction of Eq. (0.14). For the sake of completeness, we stress the important analogy between the effect discussed here and the use of relativity theory in Global Positioning Systems (GPS) [24]. The additional term $-\frac{1}{6}$ in Eq. (0.13) is similar to the correction that one has to consider in GPS when one accounts for the difference between the time measured in a frame co-rotating with the Earth geoid and the time measured in a non-rotating Earth centered frame, which is locally inertial [4–8] (and also the difference between the proper time of an observer at the surface of the Earth and at infinity). In fact, by simply considering the redshift due to the Earth's gravitational field, but neglecting the effect of the Earth's rotation, GPS cannot work, see [4–8] for details.

Finally, the attentive reader may be interested in some other useful papers regarding the topic covered in this letter, which are Refs. [25–27].

Conclusion remarks

Based on its renewed interest, due to the discovery and explanation of an additional effect of clock synchronization which has been missed for about 50 years, in this letter we addressed the theoretical explanation of such an additional effect via a novel approach founded on the third postulate of relativity by finding complete consistence with the experimental results in [14, 23] and with previuos theoretical approaches in [4–10, 22].

Data availability statement

This manuscript has no associated data.

References

- A. Einstein, *The Collected Papers*, Volume 6: The Berlin Years: Writings, 1914-1917 (English translation supplement) Pages 31-32.
- [2] W. Pauli, *Theory of Relativity*, Pergamon Press, London (1958).
- [3] W. Kündig, Phys. Rev. **129**, 2371 (1963).
- [4] C. Corda, Ann. Phys. **355**, 360 (2015).
- [5] C. Corda, Ann. Phys. **368**, 258 (2016).
- [6] C. Corda, Int. Journ. Mod. Phys. D 27, 1847016 (2018).
- [7] C. Corda, Found Phys **52**, 42 (2022).
- [8] C. Corda, Int. Journ. Mod. Phys. D 28, 1950131 (2019).
- [9] G. Iovane, E. Benedetto, Ann. Phys. **403**, 106 (2019).
- [10] E. Benedetto, A. Briscione and G. Iovane, EPL **132**, 49001 (2020)
- [11] E. Benedetto, F. Feleppa, I. Licata, H. Moradpour, C. Corda, Eur. Phys. J. C 79, 187 (2019).
- [12] R. L. Mössbauer, Zeitschrift für Physik A (in German) 151, 124 (1958).
- [13] A. L. Kholmetskii, T. Yarman and O. V. Missevitch, Phys. Scr. 77, 035302 (2008).
- [14] A. L. Kholmetskii, T. Yarman, O.V. Missevitch and B. I. Rogozev, Phys. Scr. 79, 065007 (2009).
- [15] D. C. Champeney, G. R. Isaak and A. M. Khan, Proc. Phys. Soc. 85, 583 (1965).
- [16] H. J. Hay, in Proc. 2nd Conf. Mössbauer Effect, ed A Schoen and D M T Compton (New York: Wiley) p 225 (1962).
- [17] T. E. Granshaw and H. J. Hay, in Proc. Int. School of Physics, 'Enrico Fermi' (New York: Academic) p 220 (1963).
- [18] D. C. Champeney and P. B. Moon, Proc. Phys. Soc. 77, 350 (1961).
- [19] H. J. Hay et al, Phys. Rev. Lett. 4, 165 (1960).

- [20] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*. 2nd edition, Pergamon Press, (1962).
- [21] H. Goldstein, *Classical Mechanics* (2nd ed., Addison Wesley 1980).
- [22] J. Foukzon, E. R. Men'kova, Ann. Phys. **413**, 168047 (2020).
- [23] T. Yarman, A. L. Kholmetskii, M. Arik, B. Akkus, Y. Oktem, L. A. Susam, O. V. Missevitch, Can. J. Phys. 94, 780 (2016).
- [24] N. Ashby, Liv. Rev. Rel. 6, 1 (2003).
- [25] E. Benedetto, F. Feleppa, G. Iovane, E. Laserra, Int. J. Geom. Meth. Mod. Phys. 17, 2050128 (2020).
- [26] E. Benedetto, F. Feleppa, B. Sersante, Afr. Math. **31**, 793 (2020).
- [27] E. Benedetto, A. Feoli, Eur. Phys. J. Plus 133, 53 (2018).