

MAGNETIC DIRAC SYSTEMS: VIOLATION OF BULK-EDGE CORRESPONDENCE IN THE ZIGZAG LIMIT

J.-M. BARBAROUX, H. D. CORNEAN, L. LE TREUST, N. RAYMOND,
AND E. STOCKMEYER

ABSTRACT. We consider a Dirac operator with constant magnetic field defined on a half-plane with boundary conditions that interpolate between infinite mass and zigzag. By a detailed study of the energy dispersion curves we show that the infinite mass case generically captures the profile of these curves, which undergoes a continuous pointwise deformation into the topologically different zigzag profile. Moreover, these results are applied to the bulk-edge correspondence. In particular, by means of a counterexample, we show that this correspondence does not always hold true in the zigzag case.

CONTENTS

1. Introduction and main results	1
1.1. The energy dispersion curves	4
1.1.1. Setting	4
1.1.2. Main results for the energy dispersion curves	6
2. Edge conductance formula	9
3. Energy dispersion curves	12
3.1. Preliminaries	12
3.2. A characterization of the eigenvalues for the non-zigzag case	13
4. The auxiliary quadratic forms	15
4.1. Study of $\alpha \mapsto \nu_n(\alpha, \xi)$	15
4.2. Study of $\xi \mapsto \nu_n^\pm(\alpha, \xi)$	17
5. Proofs for the energy dispersion curves	18
5.1. Proof of Theorems 1.7 and 1.8	19
Appendix A. Lemma on trace class operators	20
Acknowledgments	21
References	21

1. INTRODUCTION AND MAIN RESULTS

Bulk-edge correspondence is a notion that arises in the study of certain condensed matter physics systems possessing non-trivial topology. It establishes a connection between the bulk properties of a material (its interior or bulk region) and its edge or boundary properties. This correspondence may be given through an equation that links a Chern number, that depends only on the bulk operator, and an expression involving the edge-states localized close to the boundary; eventually yielding the

so-called topological quantization of edge currents [13]. Due to the integer (or topological) nature of the Chern number these relations are very stable under smooth changes of the material parameters and therefore its importance for potential applications. In this context, Dirac Hamiltonians are prominent examples that exhibit interesting phenomena. They are used to model various materials, among them, graphene and topological insulators [14].

In this article we investigate the bulk-edge correspondence for a two-dimensional Dirac Hamiltonian with a constant magnetic field defined on a half-plane. This model was recently considered in [6], where bulk-edge correspondence was shown to hold, provided infinite-mass boundary conditions along the edge are imposed. Our motivation is to investigate the validity of these results when any fixed admissible local boundary condition is allowed. To this end we define a family of Dirac Hamiltonians \mathcal{D}_γ , where $\gamma \in \mathbb{R} \cup \{+\infty\}$ characterizes the boundary conditions, in particular, $\gamma = \pm 1$ corresponds to infinite mass.

Our main result Theorem 1.2 indicates that bulk-edge correspondence for this model still holds, provided the boundary conditions are not zigzag (*i.e.* $\gamma \notin \{0, +\infty\}$). Moreover, Theorem 1.2 shows that this correspondence is violated for certain energies when zigzag boundary conditions are imposed. In order to show Theorem 1.2 certain knowledge on the energy dispersion curves associated to \mathcal{D}_γ is helpful. In this work we present a fairly detailed analysis of them extending the results of [3] to any local boundary condition. We complement our analysis with numerical illustrations of the energy dispersion curves for different values of the boundary parameter and the magnetic field.

Let us now turn to define our model. We consider a magnetic Dirac system on the half-plane denoted by $\mathbb{R}_+^2 = \{(x_1, x_2) \in \mathbb{R}^2 : x_2 > 0\}$ in the presence of an orthogonal magnetic field whose component in the x_3 direction is given by $b \in \mathbb{R} \setminus \{0\}$. The corresponding Hamiltonian acts on functions in $L^2(\mathbb{R}_+^2, \mathbb{C}^2)$ as

$$\sigma \cdot (-i\nabla - \mathbf{A}) = \begin{pmatrix} 0 & -i\partial_1 - \partial_2 + bx_2 \\ -i\partial_1 + \partial_2 + bx_2 & 0 \end{pmatrix}. \quad (1.1)$$

Here \mathbf{A} refers to a vector potential associated with the magnetic field *i.e.* $\text{rot}\mathbf{A} = b\mathbf{e}_3$. We choose

$$\mathbf{A} = (-bx_2, 0).$$

We recall that

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The study of the magnetic Dirac operator on the half-plane with infinite mass boundary conditions was recently carried forward in [3]. Here we consider general local conditions at the edge interpolating between zigzag and infinite mass. More precisely: Let $\gamma \in \mathbb{R} \cup \{+\infty\}$, then we consider the self-adjoint realization $\mathcal{D}_\gamma \equiv \mathcal{D}_\gamma(b)$ acting as (1.1) on a subset of functions $\psi = (\psi_1, \psi_2) \in L^2(\mathbb{R}_+^2, \mathbb{C}^2)$ satisfying, for all $x_1 \in \mathbb{R}$,

$$\begin{cases} \psi_2(x_1, 0) = \gamma\psi_1(x_1, 0) & \text{if } \gamma \in \mathbb{R}, \\ \psi_1(x_1, 0) = 0 & \text{if } \gamma = +\infty. \end{cases} \quad (1.2)$$

The two cases $\gamma \in \{0, +\infty\}$ are called zigzag, while $\gamma = \pm 1$ corresponds to infinite mass boundary conditions (see Remark 1.3 below).

Remark 1.1. The domains of self-adjointness of the operators \mathcal{D}_γ are already known: See [3, Section 1C] for the zigzag cases and [3, Theorem 1.15] for the infinite mass; the latter result can be easily adapted for the non-zigzag cases. For the essential self-adjointness of \mathcal{D}_γ on the class of Schwartz functions with infinite mass boundary conditions see [6, Proposition 1.1], and for the general case see [1].

We denote by X_1 the operator of multiplication with x_1 , and by J_1 the current density operator, we have

$$J_1 = -i[\mathcal{D}_\gamma, X_1] = -\sigma_1.$$

Recall that the Landau Hamiltonian $\mathcal{D}_{bulk}(b)$ acts on the whole plane as in (1.1) and its spectrum consists of the Landau levels given by $\{\pm\sqrt{2n|b|}, n \in \mathbb{N}_0\}$.

We say that $F : \mathbb{R} \rightarrow \mathbb{R}$ is equal to $a \in \mathbb{R}$ near $x_0 \in \mathbb{R}$ if there exists an open interval I around x_0 where $F(x) = a, x \in I$, holds.

Our main result is the following.

Theorem 1.2. *Let $b > 0$ and let $\chi = \mathbb{1}_{(0,1)}$ be the indicator function of the interval $(0, 1)$. Let $\gamma \in \mathbb{R} \cup \{+\infty\}$. Let $F \in C_0^2(\mathbb{R})$ be such that it equals 1 near $n \geq 1$ Landau levels, and 0 near the others. Then, the operator $\chi(X_1)J_1F'(\mathcal{D}_\gamma)$ is trace class and the edge Hall conductance is given by*

$$2\pi \text{Tr}(\chi(X_1)J_1F'(\mathcal{D}_\gamma)) = \begin{cases} n-1 & \text{if } \gamma = 0 \text{ and } F \text{ equals 1 near } 0, \\ n+1 & \text{if } \gamma = +\infty \text{ and } F \text{ equals 1 near } 0, \\ n & \text{otherwise.} \end{cases} \quad (1.3)$$

Comments:

- (1) The fact that the left hand side of this formula can be interpreted as an edge conductance is explained for instance in [10].
- (2) Let us make the connection with the bulk-edge correspondence. Let F and n be as in Theorem 1.2 and define the orthogonal projection $P_n = F(\mathcal{D}_{bulk}(b))$. In our case, P_n contains exactly n bulk Landau levels and one can show that its Chern number equals n , which encapsulates the non-trivial topology of the bulk projection (see *e.g.* [6]). On the other hand, since F' equals 0 near the Landau levels, $F'(\mathcal{D}_\gamma)$ only selects edge-states.
- (3) If $b < 0$, the Chern number of P_n becomes $-n$, and the right hand side of formula (1.3) must be multiplied by -1 .
- (4) Our third alternative in (1.3) indicates that the bulk-edge correspondence should hold for all non-zigzag conditions, a result which in our case is confirmed by brute force, i.e. by direct computation and comparison with the bulk Chern number. The general proof of this fact, under more general conditions than purely constant magnetic field, will be considered in [1]. At least for $\gamma = 1$, this has been shown to be the case [6]; for Schrödinger-like operators see [7].
- (5) If we work with zigzag boundary conditions, and if the zero-energy bulk Landau level belongs to the projection P_n , then one of the first two alternatives in (1.3) occurs. Thus the bulk-edge correspondence does not hold

in this case. Such an anomaly has been previously observed in other continuous models such as shallow-water waves [11, 18], and for "regularized" non-magnetic Dirac-like operators [18]. The latter are in fact second order differential operators, with boundary conditions that are incompatible with first-order self-adjoint differential operators.

Remark 1.3 (On the boundary condition). General local boundary conditions for Dirac operators are usually written as $(-i\sigma_3(\sigma \cdot \mathbf{n}) \cos \eta + \sigma_3 \sin \eta)\psi = \psi$, on $\partial\mathbb{R}_+^2$, where $\eta \in [-\frac{\pi}{2}, \frac{3\pi}{2})$ (see *e.g.* [5] and [4, 3]). In the present situation, we have $\mathbf{n} = -e_2$ and thus

$$(\sigma_1 \cos \eta + \sigma_3 \sin \eta)\psi = \psi, \quad \text{on } \partial\mathbb{R}_+^2.$$

We obtain (1.2) by setting $\gamma = \frac{\cos \eta}{1 + \sin \eta} = \tan(\frac{\pi}{4} - \frac{\eta}{2})$ with the convention $\gamma = +\infty$ in the case $\eta = -\pi/2$ *i.e.* $\psi_1 = 0$ on $\partial\mathbb{R}_+^2$.

1.1. The energy dispersion curves. Our proof of Theorem 1.2 requires certain knowledge on the energy dispersion curves and their corresponding eigenfunctions. In what follows we present a description of these curves for different values of the boundary parameter γ . The main technical issue here is the lack of semi-boundedness of \mathcal{D}_γ . This can however be treated using appropriate variational methods [12, 9, 16]; we follow the approach proposed in [3]. Before presenting our main results in this context we establish the basic framework.

1.1.1. Setting. In view of the translation invariance in the x_1 -direction, we may use the partial Fourier transform to represent $\mathcal{D}_\gamma(b)$ as a family of fiber operators $\mathcal{D}_{\gamma,\xi}(b) \equiv \mathcal{D}_{\gamma,\xi}$, with $\xi \in \mathbb{R}$. Indeed, we have (see *e.g.* [3])

$$\mathcal{D}_\gamma = \int_{\mathbb{R}}^{\oplus} \mathcal{D}_{\gamma,\xi} d\xi,$$

where the $1d$ magnetic Dirac operator $\mathcal{D}_{\gamma,\xi}$ acts as

$$\begin{pmatrix} 0 & d_\xi(b) \\ d_\xi^\dagger(b) & 0 \end{pmatrix} \equiv \begin{pmatrix} 0 & d_\xi \\ d_\xi^\dagger & 0 \end{pmatrix},$$

with $d_\xi = \xi - \partial_2 + bx_2$ and $d_\xi^\dagger = \xi + \partial_2 + bx_2$. Its domain is given for $\gamma \notin \{0, +\infty\}$ as

$$\text{Dom}(\mathcal{D}_{\gamma,\xi}) = \{\psi \in H^1(\mathbb{R}_+, \mathbb{C}^2) : x_2\psi \in L^2(\mathbb{R}_+) \text{ and } \psi_2(0) = \gamma\psi_1(0)\}. \quad (1.4)$$

As for the zigzag cases, by denoting

$$B^1(\mathbb{R}_+) = \{\psi \in H^1(\mathbb{R}_+, \mathbb{C}^2) : x_2\psi \in L^2(\mathbb{R}_+)\}$$

we have

$$\text{Dom}(\mathcal{D}_{0,\xi}) = \{u \in L^2(\mathbb{R}_+) : d_\xi^\dagger u \in L^2(\mathbb{R}_+)\} \times H_0^1(\mathbb{R}_+) \cap B^1(\mathbb{R}_+),$$

$$\text{Dom}(\mathcal{D}_{\infty,\xi}) = H_0^1(\mathbb{R}_+) \cap B^1(\mathbb{R}_+) \times \{u \in L^2(\mathbb{R}_+) : d_\xi u \in L^2(\mathbb{R}_+)\}.$$

We have that (*cf.* [3]), for any $\gamma \in \mathbb{R} \cup \{+\infty\}$ and $\xi \in \mathbb{R}$, the operator $\mathcal{D}_{\gamma,\xi}$ is self-adjoint and has compact resolvent (for $\gamma \neq \{0, +\infty\}$ it follows directly from the compact embedding of H^1 in L^2). We write the spectrum of $\mathcal{D}_{\gamma,\xi}$ as the set $\{-\vartheta_j^-(\gamma, \xi) : j \in \mathbb{N}\} \cup \{\vartheta_j^+(\gamma, \xi) : j \in \mathbb{N}\}$ such that

$$\dots \leq -\vartheta_2^-(\gamma, \xi) \leq -\vartheta_1^-(\gamma, \xi) < 0 \leq \vartheta_1^+(\gamma, \xi) \leq \vartheta_2^+(\gamma, \xi) \leq \dots \quad (1.5)$$

For a given boundary condition $\gamma \in \mathbb{R} \cup \{+\infty\}$ the map $\mathbb{R} \ni \xi \mapsto \vartheta_n^\pm(\gamma, \xi)$ defines the energy dispersion relation.

The following two propositions are shown in Section 3.1.

Proposition 1.4. *Let $\gamma \in \mathbb{R} \cup \{+\infty\}$, $\xi \in \mathbb{R}$, $b \neq 0$. We have*

- (i) *For all $n \geq 1$, the eigenvalues $\vartheta_n^\pm(\gamma, \xi)$ are simple.*
- (ii) *For all $n \geq 1$, $\xi \mapsto \vartheta_n^\pm(\gamma, \xi)$ and $\gamma \mapsto \vartheta_n^\pm(\gamma, \xi)$ are real-analytic.*
- (iii) *0 belongs to the spectrum of $\mathcal{D}_{\gamma, \xi}$ iff $\gamma = 0$, in case $b > 0$, or $\gamma = +\infty$, in case $b < 0$.*

In order to study the zigzag case we introduce further the 1d fibers of a magnetic Dirichlet Pauli operator $\mathcal{H}_\xi(b)$ for $b > 0$. It acts as $-\partial_2^2 + (\xi + bx_2)^2 + b$, with

$$\text{Dom}(\mathcal{H}_\xi(b)) = \{\psi \in H^2(\mathbb{R}_+; \mathbb{C}) : x_2^2 \psi \in L^2(\mathbb{R}_+), \psi(0) = 0\}.$$


It is well-known [8] that $\mathcal{H}_\xi(b)$ is self-adjoint with compact resolvent and that its spectrum consists on simple eigenvalues $(\nu_n^{\text{Dir}}(b, \xi))_{n \in \mathbb{N}}$ with

$$2b < \nu_1^{\text{Dir}}(b, \xi) < \nu_2^{\text{Dir}}(b, \xi) < \dots$$

The following statements are well-known (see [17] and [2, 3]). We specialize in the case $b > 0$ since otherwise one can use the charge conjugation symmetry described in Remark 1.9 below.

Proposition 1.5. *Consider the case $\gamma \in \{0, +\infty\}$. Then, for all $\xi \in \mathbb{R}$ and $b \neq 0$, the spectrum of $\mathcal{D}_{\gamma, \xi}$ is symmetric with respect to 0. Moreover, for $b > 0$, we have*

$$\begin{aligned} \vartheta_n^+(+\infty, \xi) &= \sqrt{\nu_n^{\text{Dir}}(b, \xi) - 2b} \quad (n \geq 1), \\ \vartheta_1^+(0, \xi) &= 0, \quad \text{and} \quad \vartheta_n^+(0, \xi) = \sqrt{\nu_{n-1}^{\text{Dir}}(b, \xi)} \quad (n \geq 2). \end{aligned}$$



(a) Case $\gamma = 0$: Here 0 is an eigenvalue of \mathcal{D}_γ

(b) Case $\gamma = \infty$: Here 0 is not an eigenvalue. Dashed lines correspond to the Landau levels of the bulk operator.

Figure 1. Dispersion curves for the zigzag boundary conditions

Remark 1.6. Recall [8] that, for $n \geq 1$, the function $\nu_n^{\text{Dir}}(b, \cdot)$ is increasing and

$$\lim_{\xi \rightarrow -\infty} \nu_n^{\text{Dir}}(b, \xi) = 2nb, \quad \lim_{\xi \rightarrow +\infty} \nu_n^{\text{Dir}}(b, \xi) = +\infty.$$

Figures 1a and 1b give the dispersion curves of the zigzag Dirac operators and on the left of each figure, their spectrum. All illustrations presented in this article are obtained thanks to standard finite difference schemes, inverse power and Newton-like methods.

1.1.2. *Main results for the energy dispersion curves.* In view of the symmetry it is enough to consider the cases of positive magnetic field and non-negative boundary parameter i.e. the case $(b, \gamma) \in (0, +\infty) \times [0, +\infty]$ (see Remark 1.9).

The following theorem gives a description of the dispersion curves when $\gamma \in (0, +\infty)$ and generalizes the result obtained in [3] for $\gamma = 1$.

Theorem 1.7. *Let $\gamma \in (0, +\infty)$, $\xi \in \mathbb{R}$, $b > 0$. The spectrum of $\mathcal{D}_{\gamma, \xi}(b)$ can be described as follows. Let $n \geq 1$.*

(i) *The function $\vartheta_n^+(\gamma, \cdot)$ is increasing and*

$$\lim_{\xi \rightarrow -\infty} \vartheta_n^+(\gamma, \xi) = \sqrt{2(n-1)b}, \quad \lim_{\xi \rightarrow +\infty} \vartheta_n^+(\gamma, \xi) = +\infty.$$

(ii) *The function $\vartheta_n^-(\gamma, \cdot)$ has a unique critical point, which is a non-degenerate minimum, and*

$$\lim_{\xi \rightarrow -\infty} \vartheta_n^-(\gamma, \xi) = \sqrt{2nb}, \quad \lim_{\xi \rightarrow +\infty} \vartheta_n^-(\gamma, \xi) = +\infty. \quad (1.6)$$




Figure 2. The dispersion curves of $\mathcal{D}_{\gamma, \xi}$ for $\gamma = b = 1$

Figure 2 gives the dispersion curves of the infinite mass Dirac magnetic operator with a special focus on the global maxima of the negative dispersion curves. Here $a_0 \in (0, \sqrt{2})$, was introduced in [3], it is the minimum of $\vartheta_1^-(1, \cdot)$, i.e., it is the size of the spectral gap of the Dirac operator with infinite mass boundary condition. The spectral gap of \mathcal{D}_γ as a function of γ can be read off from Figure 3.




Figure 3. The value of the maximal negative energy of the full operator as a function of γ for $b = 1$.




Figure 4. The location of the critical point of the first negative dispersion curve for $b = 1$. The red bullet refers to $\gamma = 1$.

Our next result describes the dispersion curves as functions of γ and their zigzag limits.

Theorem 1.8. *Let $b > 0$. The families of functions $[0, +\infty) \ni \gamma \mapsto \vartheta_n^+(\gamma, \cdot)$ and $[0, +\infty) \ni \gamma \mapsto -\vartheta_n^-(\gamma, \cdot)$ are increasing with γ . Moreover, for all $n \geq 1$ and all $\xi \in \mathbb{R}$, we have*

$$\lim_{\gamma \rightarrow 0} \vartheta_n^\pm(\gamma, \xi) = \vartheta_n^\pm(0, \xi), \quad \lim_{\gamma \rightarrow +\infty} \vartheta_n^\pm(\gamma, \xi) = \vartheta_n^\pm(+\infty, \xi).$$

In Figure 5 we present various pictures of the dispersion curves with varying $\gamma \in (0, +\infty)$. Moreover, Figure 6 illustrates the action of the symmetries, described in the following.

Remark 1.9 (Symmetries). In view of the underlying symmetries it is enough to study the dispersion curves when $(b, \gamma) \in (0, +\infty) \times [0, +\infty]$. Indeed, in order to also consider $\gamma < 0$, we notice that

$$\sigma_3 \mathcal{D}_\gamma(b) = -\mathcal{D}_{-\gamma}(b) \sigma_3.$$




Figure 5. The dispersion curves for $b = 1$ and various γ

Moreover, for $b < 0$ we used the charge conjugation $C\psi = \sigma_1 \bar{\psi}$ which turns the boundary conditions into $\psi_2 = \gamma^{-1}\psi_1$ and hence

$$C\mathcal{D}_\gamma(b) = -\mathcal{D}_{\gamma^{-1}}(-b)C.$$

For the fiber operators this leads to:

$$C\mathcal{D}_{\gamma,\xi}(b) = -\mathcal{D}_{\gamma^{-1},-\xi}(-b)C, \quad \sigma_3 \mathcal{D}_{\gamma,\xi}(b) = -\mathcal{D}_{-\gamma,\xi}(b)\sigma_3. \quad (1.7)$$

In particular, we obtain the dispersion curves when $b < 0$ from the curves when $b > 0$ by changing γ into γ^{-1} and ξ into $-\xi$.




Figure 6. Action of the symmetries on the dispersion curves.

Organization of this article. In Section 2 we prove Theorem 1.2. Sections 3, 4, and 5 are devoted to the description of the energy dispersion relations $\vartheta_n^\pm(\gamma, \xi)$. In Section 3 we prove Propositions 1.4 and 1.5. In addition, we state in Theorem 3.5 a fixed-point characterization of $\vartheta_n^\pm(\gamma, \xi)$ in terms of a family $(\nu_n^\pm(\alpha, \xi))_{\alpha>0}$ of the eigenvalues of certain magnetic Schrödinger-like operators with Robin boundary conditions. We give a proof of this characterization in Section 5. In Section 4 we investigate the fundamental mapping properties of $\nu_n^\pm(\alpha, \xi)$ for $\alpha > 0$ and $\xi \in \mathbb{R}$. Finally, in Section 5, we apply Theorem 3.5, together with the analysis of Section 4, to prove Theorems 1.7 and 1.8.

2. EDGE CONDUCTANCE FORMULA

In this section we prove Theorem 1.2. In doing so we use various results on the energy dispersion curves which are stated in Section 1.1 and proved in the next sections.

We let, for all $j \geq 1$,

$$\lambda_j(\xi) = \vartheta_j^+(\gamma, \xi),$$

and, for all $j \leq -1$,

$$\lambda_j(\xi) = -\vartheta_j^-(\gamma, \xi).$$

Let $(\Psi_{\xi,j})_{\xi \in \mathbb{R}}$ be an analytic family of normalized eigenfunctions of $\mathcal{D}_{\gamma,\xi}$ associated with $\lambda_j(\xi)$ (see Proposition 1.4).

In view of the asymptotic behavior of $\lambda_j(\xi)$ as $\xi \rightarrow \pm\infty$ – stated in Theorem 1.7 (for the non-zigzag case) and Proposition 1.5 and Remark 1.6 (for the zigzag case) – the proof of Theorem 1.2 reduces to showing the following result.

Proposition 2.1. *We let $\chi = \mathbb{1}_{(0,1)}$ and consider $\gamma \in \mathbb{R} \cup \{+\infty\}$. Let us consider a function $f \in C_0^1(\mathbb{R})$ being zero near the Landau levels. Then, the operator $\chi(X_1)\sigma_1 f(\mathcal{D}_\gamma)$ is trace class and there exists a finite $J \subset \mathbb{Z} \setminus \{0\}$ such that*

$$2\pi \operatorname{Tr}(\chi(X_1)\sigma_1 f(\mathcal{D}_\gamma)) = \sum_{j \in J} \int_{\mathbb{R}} f(\lambda_j(\xi)) \lambda_j'(\xi) d\xi.$$

In particular, if $n \in \mathbb{N}$ and $F \in C_0^2(\mathbb{R})$ equals 1 near n Landau levels and 0 near the others, we have

$$-2i\pi \operatorname{Tr}(\chi(X_1)[\mathcal{D}_\gamma, X_1]F'(\mathcal{D}_\gamma)) = \sum_{j \in J} \left(F(\lambda_j(-\infty)) - F(\lambda_j(+\infty)) \right). \quad (2.1)$$

Let us first state two useful elementary results. The first one is a direct consequence of Proposition 1.4 and Theorem 1.7.

Lemma 2.2. *Let us consider a function $f \in C_0(\mathbb{R})$ being zero near the Landau levels $\{\pm\sqrt{2n}, n \in \mathbb{N}\}$. Then, there exists a finite $J \subset \mathbb{Z} \setminus \{0\}$ such that for all $k \notin J$, $f \circ \lambda_k = 0$ and for all $j \in J$, the functions $\xi \mapsto f(\lambda_j(\xi))$ have compact supports.*

The following result might be elementary. We give its proof for the reader's convenience.

Lemma 2.3. *Consider an operator T on $L^2(\mathbb{R}_+^2)$ given as a Bochner integral (on a finite interval) of a continuous family of rank one operators*

$$T = \int_a^b |\psi(\xi)\rangle\langle\phi(\xi)| d\xi. \quad (2.2)$$

Assume further that the trace norm $\|\psi(\xi)\| \|\phi(\xi)\|$ of the above integrand is uniformly bounded on $[a, b]$. Then, T is trace class and

$$\mathrm{Tr}(T) = \int_a^b \mathrm{Tr} |\psi(\xi)\rangle\langle\phi(\xi)| d\xi = \int_a^b \langle\phi(\xi), \psi(\xi)\rangle d\xi. \quad (2.3)$$

Proof. The integral in (2.2) can be seen as a limit of a Riemann sum T_n , which a priori only converges in the operator norm topology. The integrand is a rank-one trace class operator, with a trace norm which is uniformly bounded in ξ on $[a, b]$. Hence, the trace norm of the T_n 's is uniformly bounded in n . By Lemma A.1 from the Appendix we see that T , which a priori is only a compact operator, is actually trace class.

Let $\{f_j\}_{j \geq 1}$ be any orthonormal system. Then for all $N \geq 1$ we have

$$\sum_{j=1}^N \langle f_j, T f_j \rangle = \int_a^b \sum_{j=1}^N \langle f_j, \psi(\xi) \rangle \langle \phi(\xi), f_j \rangle d\xi. \quad (2.4)$$

Using Cauchy-Schwarz and Bessel inequalities we get for every N :

$$\left| \sum_{j=1}^N \langle f_j, \psi(\xi) \rangle \langle \phi(\xi), f_j \rangle \right| \leq \sqrt{\sum_{j=1}^N |\langle f_j, \psi(\xi) \rangle|^2} \sqrt{\sum_{j=1}^N |\langle f_j, \phi(\xi) \rangle|^2} \leq \|\psi(\xi)\| \|\phi(\xi)\|.$$

By Lebesgue's dominated convergence theorem we can take $N \rightarrow \infty$ in (2.4) to get (2.3). \square

Proof of Proposition 2.1. Let us first investigate the integral kernel of $f(\mathcal{D}_\gamma)$ for some $f \in C_0^1(\mathbb{R})$. We can write

$$\left(f(\mathcal{D}_\gamma)\psi \right)(x) = \int_{\mathbb{R}_+^2} K_f(x, x') \psi(x') dx',$$

with

$$K_f(x, x') = \frac{1}{2\pi} \int_{\mathbb{R}} d\xi e^{i(x_1 - x'_1)\xi} k_f(\xi, x_2, x'_2),$$

where

$$k_f(\xi, x_2, x'_2) = \sum_{j \in J} f(\lambda_j(\xi)) |\Psi_{\xi, j}(x_2)\rangle \langle \Psi_{\xi, j}(x'_2)|,$$

and $J \subset \mathbb{Z} \setminus \{0\}$ is the finite set from Lemma 2.2. The technical issue here is that, even if we multiply by $\chi \equiv \chi(X_1)$ from the left we can not directly apply Lemma 2.3 since the function $(x_1, x_2) \mapsto e^{ix_1\xi} \Psi_{\xi, j}(x_2)$ is not square integrable on \mathbb{R}_+^2 . However, we observe that

$$\sigma_1 \chi f(\mathcal{D}_\gamma) = \sigma_1 \chi (1 + iX_1) f(\mathcal{D}_\gamma) (1 + iX_1)^{-1} + \sigma_1 \chi (1 + iX_1) [(1 + iX_1)^{-1}, f(\mathcal{D}_\gamma)]. \quad (2.5)$$

The first operator above can be written as

$$\begin{aligned} & \sigma_1 \chi (1 + iX_1) f(\mathcal{D}_\gamma) (1 + iX_1)^{-1} \\ &= \frac{1}{2\pi} \sum_{j \in J} \int_{\mathbb{R}} f(\lambda_j(\xi)) |(1 + iX_1) \chi e^{iX_1 \xi} \Psi_{\xi, j}\rangle \langle (1 - iX_1)^{-1} e^{iX_1 \xi} \Psi_{\xi, j}| d\xi. \end{aligned} \quad (2.6)$$

The second operator has a commutator term $[\cdot, \cdot]$ which can be explicitly computed using the following identity: We get, by doing partial integration,

$$\begin{aligned} 2\pi K_f(x, x') (1 + i x'_1) &= \int_{\mathbb{R}_\xi} d\xi \left((1 + i x_1 - \partial_\xi) e^{i(x_1 - x'_1)\xi} \right) k_f(\xi, x_2, x'_2) \\ &= 2\pi (1 + i x_1) K_f(x, x') + \int_{\mathbb{R}} d\xi e^{i(x_1 - x'_1)\xi} \partial_\xi k_f(\xi, x_2, x'_2). \end{aligned}$$

Therefore, we get as operators on $L^2(\mathbb{R}_+^2)$

$$\begin{aligned} & \sigma_1 \chi (1 + iX_1) [(1 + iX_1)^{-1}, f(\mathcal{D}_\gamma)] \\ &= \frac{1}{2\pi} \sum_{j \in J} \int_{\mathbb{R}} f'(\lambda_j(\xi)) \lambda'_j(\xi) |\sigma_1 \chi e^{iX_1 \xi} \Psi_{\xi, j}\rangle \langle (1 - iX_1)^{-1} e^{iX_1 \xi} \Psi_{\xi, j}| d\xi \\ &+ \frac{1}{2\pi} \sum_{j \in J} \int_{\mathbb{R}} f(\lambda_j(\xi)) |\sigma_1 \chi e^{iX_1 \xi} (\partial_\xi \Psi_{\xi, j})\rangle \langle (1 - iX_1)^{-1} e^{iX_1 \xi} \Psi_{\xi, j}| d\xi \\ &+ \frac{1}{2\pi} \sum_{j \in J} \int_{\mathbb{R}} f(\lambda_j(\xi)) |\sigma_1 \chi e^{iX_1 \xi} \Psi_{\xi, j}\rangle \langle (1 - iX_1)^{-1} e^{iX_1 \xi} (\partial_\xi \Psi_{\xi, j})| d\xi. \end{aligned} \quad (2.7)$$

Notice that thanks to Lemma 2.2 the integrals above take place on a finite interval. Therefore, each of the four terms appearing in (2.6) and (2.7) can be seen as Bochner integrals involving rank one operators in $L^2(\mathbb{R}_+^2)$ whose trace is uniformly bounded on compact sets. Hence, Lemma 2.3 can be applied to each of the terms involved in (2.5). In particular, as a finite sum of trace class operators, $\sigma_1 \chi f(\mathcal{D}_\gamma)$ is trace class. A quick computation using Lemma 2.3 for each term in (2.7) gives

$$\begin{aligned} & \text{Tr} \left(\sigma_1 \chi (1 + iX_1) [(1 + iX_1)^{-1}, f(\mathcal{D}_\gamma)] \right) \\ &= \frac{1}{2\pi} \sum_{j \in J} \int_{\mathbb{R}} \partial_\xi \left(f(\lambda_j(\xi)) \langle \sigma_1 \chi \Psi_{\xi, j}, (1 - iX_1)^{-1} \Psi_{\xi, j} \rangle \right) d\xi = 0, \end{aligned}$$

where in the last step we used that the term $\partial_\xi(\dots)$ has compact support in ξ . Thus, we get

$$\begin{aligned} & \text{Tr}(\chi \sigma_1 f(\mathcal{D}_\gamma)) = \text{Tr}(\sigma_1 \chi (1 + iX_1) f(\mathcal{D}_\gamma) (1 + iX_1)^{-1}) \\ &= \frac{1}{2\pi} \sum_{j \in J} \int_{\mathbb{R}} f(\lambda_j(\xi)) \langle \sigma_1 \Psi_{\xi, j}, \Psi_{\xi, j} \rangle_{L^2(\mathbb{R}_+, \mathbb{C}^2)} d\xi. \end{aligned}$$

Now the conclusion follows since $\langle \sigma_1 \Psi_{\xi, j}, \Psi_{\xi, j} \rangle_{L^2(\mathbb{R}_+, \mathbb{C}^2)} = \langle \partial_\xi(\mathcal{D}_\gamma, \xi) \Psi_{\xi, j}, \Psi_{\xi, j} \rangle_{L^2(\mathbb{R}_+, \mathbb{C}^2)}$ which, by the Feynman-Hellmann theorem, equals $\lambda'_j(\xi)$. In particular, we get (2.1) by writing $F' = f$ and integrating in ξ . \square

3. ENERGY DISPERSION CURVES

We start this section by showing Propositions 1.4 and 1.5. They state the basic properties of the solutions of the eigenvalue problem, for $\xi \in \mathbb{R}$ and $(b, \gamma) \in (0, +\infty) \times [0, +\infty]$

$$\mathcal{D}_{\gamma, \xi} u = \lambda u. \quad (3.1)$$

We show that these solutions are related to a Schrödinger-like problem with Robin boundary conditions. For zigzag boundary conditions this property is already clear from Proposition 1.5. For $\gamma \in (0, +\infty)$ we establish this relation in Lemma 3.4 below.

Moreover, we present in Theorem 3.5 a characterization of the eigenvalues $\vartheta_n^\pm(\gamma, \xi)$ in terms of a fixed-point problem that runs along a family of eigenvalues $(\nu^\pm(\alpha, \xi))_{\alpha > 0}$ of certain Schrödinger-like operators.

3.1. Preliminaries. Let us investigate some preliminary facts. (Throughout this paragraph we assume $b > 0$.) The eigenvalue equation (3.1) can be rewritten as

$$d_\xi^\dagger u_1 = \lambda u_2, \quad d_\xi u_2 = \lambda u_1.$$

Then, we have $d_\xi d_\xi^\dagger u_1 = \lambda^2 u_1$ and $d_\xi^\dagger d_\xi u_2 = \lambda^2 u_2$. Moreover, from the classical theory of ODEs, we see that u_1 and u_2 are smooth on $[0, +\infty)$. Since $u_2(0) = \gamma u_1(0)$ (or $u_1(0) = 0$ when $\gamma = +\infty$) we obtain Robin-type boundary conditions for u_1 and u_2 , separately. Thus, (3.1) implies

$$d_\xi d_\xi^\dagger u_1 = \lambda^2 u_1 \quad (d_\xi^\dagger u_1(0) = \gamma \lambda u_1(0)), \quad (3.2)$$

$$d_\xi^\dagger d_\xi u_2 = \lambda^2 u_2 \quad (d_\xi u_2(0) = \frac{\lambda}{\gamma} u_2(0)). \quad (3.3)$$

Proof of Proposition 1.4. Let us consider the eigenvalue equations (3.2) and (3.3). From the standard theory of initial value problems, we see that u_j belongs to a space of dimension at most 1. Therefore, $\dim \ker(\mathcal{D}_{\gamma, \xi} - \lambda) \leq 1$. This proves the simplicity of the non-zero eigenvalues.

Let us now discuss the existence of zero modes. For $\lambda = 0$, we have $d_\xi u_2 = 0$ so that u_2 is proportional to $e^{\frac{1}{2b}(\xi + bx_2)^2}$, which is not in $L^2(\mathbb{R}_+)$ implying that $u_2 = 0$ holds. Moreover, we also check that $\gamma u_1(0) = 0$. Using $d_\xi^\dagger u_1 = 0$ we see that u_1 is proportional to $e^{-\frac{1}{2b}(\xi + bx_2)^2}$, which belongs to $L^2(\mathbb{R}_+)$ but it does not vanish at $x_2 = 0$. Therefore, we find that $u_1 = 0$ unless $\gamma = 0$.

The family $(\mathcal{D}_{\gamma, \xi})_{\xi \in \mathbb{R}}$ being analytic of type (A) (in the Kato sense, see [15]), the simplicity of the eigenvalues implies their analyticity. \square

Next we discuss the zigzag operators *i.e.* the cases $\gamma \in \{0, +\infty\}$.

Proof of Proposition 1.5. The fact that the eigenvalues are symmetric with respect to zero follows from (1.7), hence, we may look at the non-negative ones only.

Let us consider the case $\gamma = 0$ *i.e.* $u_2(0) = 0$. As we have just seen, we have a zero mode and, with our convention, we have $\vartheta_1^+(0, \xi) = 0$. Let us describe the non-zero eigenvalues. Let λ be a positive eigenvalue and u a corresponding eigenfunction. In view of (3.2), we see that u_2 cannot be 0 and it is an eigenfunction of

$$d_\xi^\dagger d_\xi = -\partial_2^2 + (\xi + bx_2)^2 + b,$$

with Dirichlet condition. In particular, λ^2 belongs to the spectrum of $d_\xi^\dagger d_\xi$ with Dirichlet condition. Conversely, if $\mu > 0$ is an eigenvalue of this operator, we write $d_\xi^\dagger d_\xi v = \mu v$ with $v(0) = 0$ and we let $u = \mu^{-\frac{1}{2}} d_\xi v$ ($\neq 0$) and we have

$$d_\xi^\dagger u = \sqrt{\mu} v, \quad d_\xi v = \sqrt{\mu} u, \quad v(0) = 0,$$

which means that $\sqrt{\mu}$ is an eigenvalue of $\mathcal{D}_{0,\xi}$.

The case $\gamma = +\infty$ is quite similar although we have no zero modes. Now, u_1 is an eigenfunction (with eigenvalue λ^2) of

$$d_\xi d_\xi^\dagger = -\partial_2^2 + (\xi + bx_2)^2 - b,$$

with Dirichlet condition. Conversely, consider an eigenvalue $\mu > 0$ of this operator. Proceeding as before, we write $d_\xi d_\xi^\dagger u = \mu u$ with $u(0) = 0$ and let $v = \mu^{-\frac{1}{2}} d_\xi^\dagger u$ ($\neq 0$) to get that $\sqrt{\mu}$ is an eigenvalue of $\mathcal{D}_{\infty,\xi}$. \square

3.2. A characterization of the eigenvalues for the non-zigzag case. We consider $\gamma \in (0, +\infty)$. Let $\lambda \neq 0$ be an eigenvalue of $\mathcal{D}_{\gamma,\xi}(b)$. Multiplying (3.2) by u_1 and integrating by parts yields

$$\langle d_\xi d_\xi^\dagger u_1, u_1 \rangle = \|d_\xi^\dagger u_1\|^2 + d_\xi^\dagger u_1(0) u_1(0) = \|d_\xi^\dagger u_1\|^2 + \lambda u_2(0) u_1(0) = \|d_\xi^\dagger u_1\|^2 + \lambda \gamma u_1^2(0).$$

Moreover, proceeding analogously for the second component in (3.3) we get

$$\langle d_\xi^\dagger d_\xi u_2, u_2 \rangle = \|d_\xi u_2\|^2 - d_\xi u_2(0) u_2(0) = \|d_\xi u_2\|^2 - \lambda u_1(0) u_2(0) = \|d_\xi u_2\|^2 - \frac{\lambda}{\gamma} u_1^2(0).$$

This suggests to introduce the following family of quadratic forms.

Definition 3.1. Let $\alpha > 0$. We define the auxiliary quadratic forms, for $u \in B^1(\mathbb{R}_+)$, as

$$\begin{aligned} q_{b,\alpha,\xi}^+(u) &= \|d_\xi^\dagger u\|^2 + \alpha u^2(0), \\ q_{b,\alpha,\xi}^-(u) &= \|d_\xi u\|^2 + \alpha u^2(0). \end{aligned} \tag{3.4}$$

They are both non-negative and closed. We denote by $\mathfrak{h}_{\alpha,\xi}^\pm$ the corresponding self-adjoint Schrödinger operators.

Remark 3.2. By Friedrichs' extension theorem we have that $\text{Dom}(\mathfrak{h}_{\alpha,\xi}^\pm) \subset B^1(\mathbb{R}_+)$ and for $u^\pm \in \text{Dom}(\mathfrak{h}_{\alpha,\xi}^\pm)$

$$\mathfrak{h}_{\alpha,\xi}^+ u^+ = d_\xi d_\xi^\dagger u^+, \quad d_\xi^\dagger u^+(0) = \alpha u^+(0), \tag{3.5}$$

$$\mathfrak{h}_{\alpha,\xi}^- u^- = d_\xi^\dagger d_\xi u^-, \quad d_\xi u^-(0) = -\alpha u^-(0). \tag{3.6}$$

Remark 3.3. Integration by parts yields

$$\begin{aligned} q_{b,\alpha,\xi}^+(u) &= \|u'\|^2 + \|(\xi + bx_2)u\|^2 - b\|u\|^2 + (\alpha - \xi)u^2(0), \\ q_{b,\alpha,\xi}^-(u) &= \|u'\|^2 + \|(\xi + bx_2)u\|^2 + b\|u\|^2 + (\alpha + \xi)u^2(0). \end{aligned} \tag{3.7}$$

We also observe that

$$q_{b,\alpha,\xi}^- = q_{-b,\alpha,-\xi}^+, \tag{3.8}$$

which reflects the first relation in (1.7). In what follows, we drop the reference to b in the notation.

In relation to our problem we see that, for $u = (u_1, u_2)$ an eigenfunction of $\mathcal{D}_{\gamma,\xi}$, we have

$$\begin{aligned} q_{\lambda\gamma,\xi}^+(u_1) &= \langle u_1, d_\xi d_\xi^\dagger u_1 \rangle = \lambda^2 \|u_1\|^2, \quad \text{for } \lambda > 0 \quad \text{and,} \\ q_{-\lambda\gamma^{-1},\xi}^-(u_2) &= \langle u_2, d_\xi^\dagger d_\xi u_2 \rangle = \lambda^2 \|u_2\|^2, \quad \text{for } \lambda < 0. \end{aligned}$$

Next, we describe a bijection existing between the kernels of $\mathcal{D}_{\gamma,\xi} - \lambda$ and $\mathfrak{h}_{\gamma\lambda,\xi}^+ - \lambda^2$ provided $\gamma\lambda > 0$. If $\gamma\lambda < 0$ analogous statements can be obtained for $\mathfrak{h}_{-\lambda/\gamma,\xi}^- - \lambda^2$. The following lemma is a straightforward adaptation of [3, Proposition 2.9]. We recall its proof for the convenience of the reader and we emphasize that it does not require sign assumptions on b and γ .

Lemma 3.4. *Let $(b, \gamma) \in \mathbb{R}^2$, $\xi \in \mathbb{R}$. Then, for any $\lambda \in \mathbb{R} \setminus \{0\}$ with $\gamma\lambda > 0$, the map*

$$\mathcal{J} : \ker(\mathfrak{h}_{\gamma\lambda,\xi}^+ - \lambda^2) \ni u \mapsto (u, \lambda^{-1} d_\xi^\dagger u) \in \ker(\mathcal{D}_{\gamma,\xi} - \lambda)$$

is well-defined and it is an isomorphism.

Proof. First, let $u \in \ker(\mathfrak{h}_{\gamma\lambda,\xi}^+ - \lambda^2)$ and $v \in \text{Dom}(\mathcal{D}_{\gamma,\xi} - \lambda)$. We have

$$\langle \mathcal{J}(u), (\mathcal{D}_{\gamma,\xi} - \lambda)v \rangle = \langle u, d_\xi v_2 - \lambda v_1 \rangle + \lambda^{-1} \langle d_\xi^\dagger u, d_\xi^\dagger v_1 - \lambda v_2 \rangle,$$

so that, by integrating by parts,

$$\langle \mathcal{J}(u), (\mathcal{D}_{\gamma,\xi} - \lambda)v \rangle = \lambda^{-1} (q_{\gamma\lambda,\xi}^+(u, v_1) - \lambda^2 \langle u, v_1 \rangle) = 0,$$

where we used that $u \in \ker(\mathfrak{h}_{\gamma\lambda,\xi}^+ - \lambda^2)$. Thus, \mathcal{J} is well defined. It is also injective from the very definition. For the surjectivity, we consider $(u_1, u_2) \in \ker(\mathcal{D}_{\gamma,\xi} - \lambda)$. We have

$$d_\xi u_2 = \lambda u_1, \quad d_\xi^\dagger u_1 = \lambda u_2.$$

We only have to check that $u_1 \in \ker(\mathfrak{h}_{\gamma\lambda,\xi}^+ - \lambda^2)$. Take $v \in B^1(\mathbb{R}_+)$ and notice that

$$\begin{aligned} q_{\gamma\lambda,\xi}^+(u_1, v) - \lambda^2 \langle u_1, v \rangle &= \langle d_\xi^\dagger u_1, d_\xi^\dagger v \rangle + \gamma \lambda u_1(0)v(0) - \lambda^2 \langle u_1, v \rangle \\ &= \lambda \langle u_2, d_\xi^\dagger v \rangle + \gamma \lambda u_1(0)v(0) - \lambda^2 \langle u_1, v \rangle \\ &= \lambda \langle d_\xi u_2, v \rangle - \lambda^2 \langle u_1, v \rangle \\ &= 0. \end{aligned}$$

This finishes the proof. \square

Let us now turn to the characterization. Since the family $(q_{\alpha,\xi}^\pm)_{(\alpha,\xi) \in \mathbb{R}_+ \times \mathbb{R}}$ is analytic on the common domain $B^1(\mathbb{R}_+)$ the eigenvalues of $\mathfrak{h}_{\alpha,\xi}^\pm$ (which are all simple) are also real analytic with respect to α and to ξ . We denote them by $\nu_n^\pm(\alpha, \xi)$ so that

$$0 \leq \nu_1^\pm(\alpha, \xi) < \nu_2^\pm(\alpha, \xi) < \dots$$

The following result completely characterizes positive and negative eigenvalues of $\mathcal{D}_{\gamma,\xi}$ in terms of $\alpha \mapsto \nu_n^\pm(\alpha, \xi)$. Recall the notation in (1.5).

Theorem 3.5. *Let $b \neq 0$ and $\gamma \in (0, +\infty)$. The equation*

$$\nu_n^+(\gamma\lambda, \xi) = \lambda^2 \tag{3.9}$$

has a unique positive solution $\lambda = \vartheta_n^+(\gamma, \xi)$. Moreover, the equation

$$\nu_n^-(\gamma^{-1}\lambda, \xi) = \lambda^2 \tag{3.10}$$

has a unique positive solution $\lambda = \vartheta_n^-(\gamma, \xi)$.

The proof of this theorem uses Lemma 3.4 and requires the analysis of the auxiliary quadratic forms performed in Section 4; we postpone it to Section 5.

4. THE AUXILIARY QUADRATIC FORMS

In this section we perform a detailed study of the auxiliary quadratic forms from Definition 3.4. We restrict the analysis to the case in which $(b, \gamma) \in (0, +\infty) \times (0, +\infty)$.

For $\alpha > 0$ and $\xi \in \mathbb{R}$ consider the eigenvalue problems (recall Remark 3.2)

$$\mathfrak{h}_{\alpha, \xi}^{\pm} u_{\alpha, \xi}^{\pm} = \nu^{\pm}(\alpha, \xi) u_{\alpha, \xi}^{\pm}. \quad (4.1)$$

Most of the following results can be traced back to [3] (notice, however, the different convention for $q_{\alpha, \xi}^-$). For the sake of completeness we present a concise argument.

4.1. Study of $\alpha \mapsto \nu_n(\alpha, \xi)$.

Lemma 4.1. *Let $\nu^{\pm}(\alpha, \xi)$ be an eigenvalue as in (4.1). Then, the function $\alpha \mapsto \nu^{\pm}(\alpha, \xi)$ is increasing and*

$$\partial_{\alpha} \nu^{\pm}(\alpha, \xi) = (u_{\alpha, \xi}^{\pm}(0))^2 > 0. \quad (4.2)$$

Proof. We only argue for the + case. To simplify notation, we denote the corresponding normalized solution of (4.1) as u ($\equiv u_{\alpha, \xi}^+$) and we drop the reference to ξ and α when not relevant.

Let us first observe that for any smooth function g on $[0, \infty)$ we have, integrating by parts,

$$\begin{aligned} \langle u, (dd^{\dagger} - \nu)g \rangle &= \langle d^{\dagger}u, d^{\dagger}g \rangle + u(0)(d^{\dagger}g)(0) - \nu \langle u, g \rangle \\ &= u(0)(d^{\dagger}g)(0) - (d^{\dagger}u)(0)g(0). \end{aligned} \quad (4.3)$$

In view of the smoothness of u with respect to α and x , we see that $d^{\dagger} \partial_{\alpha} u = \partial_{\alpha} d^{\dagger} u$. Hence, since $d^{\dagger}u(0) = \alpha u(0)$, we get that

$$(d^{\dagger} \partial_{\alpha} u)(0) = u(0) + \alpha \partial_{\alpha} u(0). \quad (4.4)$$

Taking derivative with respect to α in the eigenvalue equation we get

$$(dd^{\dagger} - \nu) \partial_{\alpha} u = (\partial_{\alpha} \nu) u.$$

After multiplying by u and integrating we use (4.3) to get

$$\partial_{\alpha} \nu = \langle u, dd^{\dagger} \partial_{\alpha} u \rangle - \nu \langle u, \partial_{\alpha} u \rangle = u(0)(d^{\dagger} \partial_{\alpha} u)(0) - (d^{\dagger}u)(0)(\partial_{\alpha} u)(0).$$

The conclusion follows using (3.5) and (4.4). \square

Lemma 4.2. *For all $n \geq 1$, we have*

$$\lim_{\alpha \rightarrow 0} \nu_n^-(\alpha, \xi) = \nu_n^-(0, \xi) = \nu_n^{\text{Dir}}(-b, -\xi) \quad (4.5)$$

$$\lim_{\alpha \rightarrow +\infty} \nu_n^-(\alpha, \xi) = \nu_n^{\text{Dir}}(b, \xi). \quad (4.6)$$

Proof. The first equality follows by analyticity. Then, we have $q_{0,\xi}^-(u) = \|d_\xi u\|^2$. The corresponding operator $\mathfrak{h}_{0,\xi}^-$ has no zero mode. Now, if ν is a positive eigenvalue, we have

$$d_\xi^\dagger d_\xi u = \nu u, \quad d_\xi u(0) = 0.$$

Letting $v = d_\xi u$, we get $d_\xi d_\xi^\dagger v = \nu v$ with $v(0) = 0$. This shows that ν belongs to the spectrum of the Dirichlet realization of $d_\xi d_\xi^\dagger$. Conversely, if $\nu > 0$ is an eigenvalue of $d_\xi d_\xi^\dagger$ associated with the eigenfunction v , we have

$$d_\xi^\dagger d_\xi u = \nu u, \quad u = d_\xi^\dagger v, \quad d_\xi u(0) = 0.$$

We can check that $u \neq 0$ (unless $v = 0$). Thus, ν belongs to the spectrum of the operator $\mathfrak{h}_{0,\xi}^-$.

When $\alpha \rightarrow +\infty$, we are in a singular regime. By using that $H_0^1(\mathbb{R}_+) \cap B^1(\mathbb{R}_+) \subset B^1(\mathbb{R}_+)$ and the min-max principle, we see that

$$\nu_n^-(\alpha, \xi) \leq \nu_n^{\text{Dir}}(b, \xi).$$

Conversely, let us consider

$$E_n(\alpha, \xi) := \text{span}_{1 \leq k \leq n} u_{\alpha,\xi,k}^-.$$

We notice that, for all $u \in E_n(\alpha, \xi)$,

$$(\alpha + \xi)u^2(0) \leq q_{\alpha,\xi}^-(u) \leq \nu_n^-(\alpha, \xi)\|u\|^2 \leq \nu_n^{\text{Dir}}(b, \xi)\|u\|^2,$$

so that, for α large enough,

$$u^2(0) \leq \frac{\nu_n^{\text{Dir}}(b, \xi)}{\alpha + \xi}\|u\|^2 = \mathcal{O}(\alpha^{-1})\|u\|^2.$$

Then, we also notice that

$$\|u'\|^2 + \|(\xi + bx_2)u\|^2 + b\|u\|^2 \leq q_{\alpha,\xi}^-(u) \leq \nu_n^{\text{Dir}}(b, \xi)\|u\|^2.$$

Let us consider a smooth cutoff function χ with compact support equal to 1 near 0. The function

$$\tilde{u}(x_2) = u(x_2) - u(0)\chi(x_2)$$

satisfies the Dirichlet boundary condition. We notice that

$$(1 - C\alpha^{-\frac{1}{2}})\|u\| \leq \|\tilde{u}\| \leq (1 + C\alpha^{-\frac{1}{2}})\|u\|.$$

This tells us that, when u runs over $E_n(\alpha, \xi)$, \tilde{u} also runs over a space of dimension n .

In the same way, we get

$$\|u'\|^2 + \|(\xi + bx_2)u\|^2 + b\|u\|^2 \geq (1 - C\alpha^{-\frac{1}{2}}) (\|\tilde{u}'\|^2 + \|(\xi + bx_2)\tilde{u}\|^2 + b\|\tilde{u}\|^2).$$

We deduce that

$$\|\tilde{u}'\|^2 + \|(\xi + bx_2)\tilde{u}\|^2 + b\|\tilde{u}\|^2 \leq (1 + C\alpha^{-\frac{1}{2}})\nu_n^-(\alpha, \xi)\|\tilde{u}\|^2.$$

Using the min-max principle, we infer that

$$\nu_n^{\text{Dir}}(b, \xi) \leq (1 + C\alpha^{-\frac{1}{2}})\nu_n^-(\alpha, \xi),$$

and the result follows. \square

Lemma 4.3. *For $n \geq 1$, we have*

$$\lim_{\alpha \rightarrow 0} \nu_n^+(\alpha, \xi) = \nu_n^+(0, \xi) = \begin{cases} 0 & n = 1 \\ \nu_{n-1}^{\text{Dir}}(b, \xi) & n \geq 2 \end{cases}, \quad (4.7)$$

$$\lim_{\alpha \rightarrow +\infty} \nu_n^+(\alpha, \xi) = \nu_n^{\text{Dir}}(-b, -\xi). \quad (4.8)$$

Proof. The proof is similar to that of Lemma 4.2. We have $q_{0,\xi}^+(u) = \|d_\xi^\dagger u\|^2$. In particular, 0 is an eigenvalue associated with $x_2 \mapsto e^{-\frac{1}{b}(\xi+bx_2)^2}$. So, $\nu_1^+(0, \xi) = 0$. Then, let us consider a positive eigenvalue ν . We have

$$d_\xi d_\xi^\dagger u = \nu u, \quad d_\xi^\dagger u(0) = 0.$$

This implies that

$$d_\xi^\dagger d_\xi v = \nu v, \quad \text{with } v = d_\xi^\dagger u \neq 0, \text{ and } v(0) = 0.$$

Conversely, if v is an eigenfunction of the Dirichlet realization of $d_\xi^\dagger d_\xi$ with eigenvalue ν , we have

$$d_\xi d_\xi^\dagger u = \nu u, \quad \text{with } u = d_\xi v, \quad \text{and } d_\xi^\dagger u(0) = 0.$$

The argument to show the limit in (4.8) follows the same lines as the proof of (4.6). \square

4.2. Study of $\xi \mapsto \nu_n^\pm(\alpha, \xi)$.

Lemma 4.4. *Let $\nu^\pm(\alpha, \xi)$ be an eigenvalue as in (4.1). Then, we have*

$$\partial_\xi \nu^\pm(\alpha, \xi) = \frac{1}{b} (\nu^\pm(\alpha, \xi) + \alpha^2 \mp 2\alpha\xi) (u_{\alpha,\xi}^\pm(0))^2. \quad (4.9)$$

In addition, if ξ_α is a critical point of $\xi \mapsto \nu(\alpha, \xi)$, we have

$$\partial_\xi^2 \nu^\pm(\alpha, \xi_\alpha) = \mp \alpha \frac{2}{b} (u_{\alpha,\xi}^\pm(0))^2. \quad (4.10)$$

In particular, $\nu^\pm(\alpha, \cdot)$ has at most one critical point. This critical point can only be a local maximum for $\nu^+(\alpha, \cdot)$ and a local minimum for ν^- .

Proof. We give again the proof only for the + case. We use the notation from the proof of the previous Lemma 4.1. (We also replace x_2 by t in the notation.)

Observe that since $\partial_\xi d^\dagger u = d^\dagger \partial_\xi u + u$, we get

$$(d^\dagger \partial_\xi u)(0) = \alpha(\partial_\xi u)(0) - u(0). \quad (4.11)$$

By differentiating (4.1) with respect to ξ we get

$$(dd^\dagger - \nu)\partial_\xi u = [\partial_\xi \nu - 2(\xi + bt)]u.$$

Hence, (4.3) and (4.11) yield

$$\partial_\xi \nu = \langle u, 2(\xi + bt)u \rangle - u(0)^2. \quad (4.12)$$

In addition, integrating by parts and using (4.1), we calculate

$$\begin{aligned}
\langle u, 2(\xi + bt)u \rangle &= \frac{1}{b} \int_{\mathbb{R}_+} u(t)^2 \partial_t (\xi + bt)^2 dt \\
&= -\frac{\xi^2}{b} u(0)^2 - \frac{2}{b} \int_{\mathbb{R}_+} u'(t) (\xi + bt)^2 u(t) dt \\
&= -\frac{\xi^2}{b} u(0)^2 - \frac{2}{b} \int_{\mathbb{R}_+} u'(t) (\nu + b + \partial_t^2) u(t) dt \\
&= -\frac{\xi^2}{b} u(0)^2 - \frac{1}{b} \int_{\mathbb{R}_+} \partial_t [((\nu + b)u(t)^2 + u'(t)^2)] dt.
\end{aligned}$$

Using that $u'(0) = (\alpha - \xi)u(0)$ we readily obtain (4.9). Hence, if a critical point ξ_α exists, it satisfies $\nu^+(\alpha, \xi_\alpha) + \alpha^2 - 2\alpha\xi_\alpha = 0$. Taking the derivative of (4.9) with respect to ξ and evaluating at ξ_α we obtain (4.10). \square

With the help of the perturbation theory, we get the following (see [3, Lemma 4.14]).

Lemma 4.5. *We have*

$$\lim_{\xi \rightarrow -\infty} \nu_n^+(\alpha, \xi) = (2n - 2)b, \quad \lim_{\xi \rightarrow +\infty} \nu_n^+(\alpha, \xi) = +\infty,$$

and

$$\lim_{\xi \rightarrow -\infty} \nu_n^-(\alpha, \xi) = 2nb, \quad \lim_{\xi \rightarrow +\infty} \nu_n^-(\alpha, \xi) = +\infty.$$

This allows us to show the following.

Lemma 4.6. *The function $\nu^+(\alpha, \cdot)$ has no critical points. Moreover, $\nu^-(\alpha, \cdot)$ has a unique critical point, which is a global minimum.*

Proof. Since $\nu^+(\alpha, \cdot) > 0$, from (4.9) we see that it is increasing on $(-\infty, 0)$. If it has a (unique) critical point for some $\xi_\alpha > 0$, it must be a non-degenerate global maximum. This contradicts the limit at $\xi = +\infty$, hence $\nu^+(\alpha, \cdot)$ is increasing on \mathbb{R} .

Now assume that $\nu^-(\alpha, \cdot)$ does not have critical points. From (4.9) we must have $\nu^-(\alpha, \xi) + \alpha^2 + 2\alpha\xi < 0$ for all ξ (since it is the case for $\xi \rightarrow -\infty$). But this would imply that $\nu^-(\alpha, \cdot)$ is decreasing on \mathbb{R} , which contradicts its limit at $\xi \rightarrow +\infty$. \square

5. PROOFS FOR THE ENERGY DISPERSION CURVES

In this section we start by proving the characterization described in Theorem 3.5. Next, we apply that result to show Theorems 1.7 and 1.8.

Proof of Theorem 3.5. It is enough to deal with the positive eigenvalues of $\mathcal{D}_{\gamma, \xi}(b)$. Indeed, due to the charge conjugation (1.7), $\vartheta_n^-(\gamma, \xi)$ is the n -th positive eigenvalue of $\mathcal{D}_{\gamma^{-1}, -\xi}(-b)$. Thus, if the characterization (3.9) is established, $\vartheta_n^-(\gamma, \xi)$ is the unique positive solution of $\nu_n^+(-b, \gamma^{-1}\lambda, -\xi) = \lambda^2$ or equivalently of (3.10) (here we use (3.8)).

Let us now prove that (3.9) has exactly one positive solution. Remember that $\gamma > 0$. We let

$$f(\lambda) = \nu_n^+(\gamma\lambda, \xi) - \lambda^2,$$

and notice that $\lim_{\lambda \rightarrow +\infty} f(\lambda) = -\infty$, $f(0) \geq 0$, and $f'(0) > 0$ (see (4.2)). Thus (3.9) has at least one positive solution. If E is such a solution, we have $f(E) = 0$ and we notice that

$$f'(E) = \gamma \partial_\alpha \nu_n^+(\gamma E, \xi) - 2E = \gamma [u_{\gamma E, \xi}(0)]^2 - 2E.$$

To get the sign of $f'(E)$, we consider the polynomial of degree two given by

$$P(\lambda) = q_{\gamma \lambda, \xi}^+(u_{\gamma E, \xi}) - \lambda^2.$$

Because $P(0) \geq 0$, $P(-\infty) = -\infty$, and $P(E) = f(E) = 0$ with $E > 0$, the polynomial must have two roots of opposite sign. Thus, $f'(E) = P'(E) < 0$. This shows that (3.9) has at most one positive solution and thus exactly one, which is denoted by $E_n(\gamma, \xi)$.

In fact, $(E_n)_{n \geq 1}$ is increasing. Indeed,

$$0 = \nu_{n+1}^+(\gamma E_{n+1}, \xi) - E_{n+1}^2 > \nu_n^+(\gamma E_{n+1}, \xi) - E_{n+1}^2 = f(E_{n+1}),$$

which implies that $E_n < E_{n+1}$.

For all $n \geq 1$, due to Lemma 3.4, $E_n(\gamma, \xi)$ is a positive eigenvalue of $\mathcal{D}_{\gamma, \xi}$. This tells us that

$$\mathcal{A} : \mathbb{N}^* \ni n \mapsto E_n(\gamma, \xi) \in \text{spec}(\mathcal{D}_{\gamma, \xi}) \cap \mathbb{R}_+$$

is well-defined (and it is injective).

We now show that the map is surjective. For all $n \geq 1$, Lemma 3.4 implies that $\mathfrak{h}_{\gamma \vartheta_n^+(\gamma, \xi), \xi}^+ - (\vartheta_n^+(\gamma, \xi))^2$ has a non-zero kernel. This means that, for some $m \geq 1$, we have

$$\nu_m^+(\gamma \vartheta_n^+(\gamma, \xi), \xi) = (\vartheta_n^+(\gamma, \xi))^2,$$

and thus $\vartheta_n^+(\gamma, \xi) = E_m(\gamma, \xi)$. This implies that \mathcal{A} is bijective, hence $E_n(\gamma, \xi) = \vartheta_n^+(\gamma, \xi)$ for all $n \geq 1$. \square

5.1. Proof of Theorems 1.7 and 1.8. In what follows, in order to ease the readability, we drop the reference to the index n in the notation. In view of Theorem 3.5 we have

$$\nu^+(\gamma \vartheta^+, \xi) = (\vartheta^+)^2$$

and, due to the analyticity and the chain rule, the derivative with respect to ξ gives (with $\alpha \equiv \gamma \vartheta^+$)

$$(\gamma \partial_\alpha \nu^+(\gamma \vartheta^+, \xi) - 2\vartheta^+) \partial_\xi \vartheta^+ + \partial_\xi \nu^+(\gamma \vartheta^+, \xi) = 0, \quad (5.1)$$

and differentiating with respect to γ yields:

$$(\gamma \partial_\alpha \nu^+(\gamma \vartheta^+, \xi) - 2\vartheta^+) \partial_\gamma \vartheta^+ + \partial_\alpha \nu^+(\gamma \vartheta^+, \xi) \vartheta^+ = 0. \quad (5.2)$$

We saw in the proof of Theorem 3.5 that

$$\gamma \partial_\alpha \nu^+(\gamma \vartheta^+, \xi) - 2\vartheta^+ < 0.$$

In the $+$ case, we see that $\vartheta^+(\gamma, \cdot)$ has no critical points and is increasing. We also see that $\gamma \mapsto \vartheta^+(\gamma, \xi)$ is increasing (by using (4.2)), which proves the monotonicity in γ of ϑ^+ announced in Theorem 1.8.

In the $-$ case, by performing the same derivatives on (3.10), we have

$$(\gamma^{-1} \partial_\alpha \nu^-(\gamma^{-1} \vartheta^-, \xi) - 2\vartheta^-) \partial_\xi \vartheta^- + \partial_\xi \nu^-(\gamma^{-1} \vartheta^-, \xi) = 0, \quad (5.3)$$

and

$$(\gamma^{-1} \partial_\alpha \nu^-(\gamma^{-1} \vartheta^-, \xi) - 2\vartheta^-) \partial_\gamma \vartheta^- - \gamma^{-2} \partial_\alpha \nu^-(\gamma^{-1} \vartheta^-, \xi) \vartheta^- = 0. \quad (5.4)$$

We still have $\gamma^{-1}\partial_\alpha\nu^-(\gamma^{-1}\vartheta^-, \xi) - 2\vartheta^- < 0$. In particular, $\gamma \mapsto \vartheta^-(\gamma, \xi)$ is decreasing.

If ξ_γ is a critical point of $\vartheta^-(\gamma, \cdot)$, then we have

$$\partial_\xi\nu^-(\gamma^{-1}\vartheta^-(\gamma, \xi_\gamma), \xi_\gamma) = 0.$$

Remark 5.1. We recall Lemma 4.4 and we have

$$\nu^-(\alpha, \xi_\gamma) + \alpha^2 + 2\alpha\xi_\gamma = 0, \quad \text{with } \alpha = \gamma^{-1}\vartheta^-(\gamma, \xi_\gamma).$$

Hence, $\vartheta^-(\gamma, \xi_\gamma) = -\frac{2\gamma}{\gamma^2+1}\xi_\gamma$ and $\xi_\gamma = \xi_\alpha$ by the uniqueness of the critical point.

Being a non-degenerate minimum, it is necessary that $\partial_\xi^2\nu^-(\gamma^{-1}\vartheta^-(\gamma, \xi_\gamma), \xi_\gamma) > 0$ and, by taking one more derivative in ξ of (5.3), we see that $\partial_\xi^2\vartheta^-(\gamma, \xi_\gamma) > 0$. Therefore, all the critical points of $\vartheta^-(\gamma, \cdot)$ are local non-degenerate minima and thus there is at most one such point. If there is no critical point, we have, for all ξ ,

$$\partial_\xi\vartheta^-(\gamma, \xi) = C \left((1 + \gamma^{-2})\vartheta^-(\gamma, \xi) + 2\gamma^{-1}\xi \right) \neq 0, \quad C > 0.$$

Let us assume for the moment that (1.6) is true; we will prove that later on. If ξ is sufficiently negative, then the left-hand side of the above expression is negative, so it must remain negative for all ξ . This implies that $\vartheta^-(\gamma, \xi)$ must be bounded from above, contradicting the limit $\xi \rightarrow +\infty$ in (1.6). This ends the analysis of critical points announced in Theorem 1.7.

It remains to explain why (1.6) holds. We only consider the limit $\xi \rightarrow -\infty$. We recall Lemma 4.5. Let us fix $\varepsilon > 0$ and define $\lambda_1 = \sqrt{2nb - \varepsilon}$ and $\lambda_2 = \sqrt{2nb + \varepsilon}$. Then there exists $\xi(\varepsilon) < 0$ such that for all $\xi < \xi(\varepsilon)$ we have

$$\nu^-(\gamma^{-1}\lambda_1, \xi) - \lambda_1^2 > 0 \quad \text{and} \quad \nu^-(\gamma^{-1}\lambda_2, \xi) - \lambda_2^2 < 0.$$

This implies that

$$\lambda_1 < \vartheta^-(\gamma, \xi) < \lambda_2, \quad \forall \xi < \xi(\varepsilon).$$

The limit $\xi \rightarrow +\infty$ can be analyzed similarly (as well as the limits for the ϑ^+). This ends the proof of Theorem 1.7.

It remains to discuss the limits in Theorem 1.8. Let us consider $\vartheta_n^+(\gamma, \xi)$. Take $\varepsilon > 0$ and $\lambda = \vartheta_n(0, \xi) - \varepsilon$ (for $n \geq 2$). We have $\nu^+(\gamma\lambda, \xi) - \lambda^2 > 0$ for γ small enough since $\nu^+(0, \xi) = (\vartheta^+(0, \xi))^2$. Thus, $\vartheta^+(0, \xi) - \varepsilon < \vartheta^+(\gamma, \xi)$. In the same way, we get $\vartheta^+(\gamma, \xi) < \vartheta^+(0, \xi) + \varepsilon$. This proves the first limit in Theorem 1.8.

Next, we consider the limit $\gamma \rightarrow +\infty$. We take $\lambda = \vartheta^+(+\infty, \xi) - \varepsilon$. We have $\nu^+(\gamma\lambda, \xi) - \lambda^2 > 0$ for γ large enough since $\nu^+(+\infty, \xi) = (\vartheta^+(+\infty, \xi))^2$. Thus, $\vartheta^+(+\infty, \xi) - \varepsilon < \vartheta^+(\gamma, \xi)$. We easily get the upper bound $\vartheta^+(\gamma, \xi) < \vartheta^+(+\infty, \xi) + \varepsilon$. The case of $\vartheta^-(\gamma, \xi)$ is similar.

APPENDIX A. LEMMA ON TRACE CLASS OPERATORS

Lemma A.1. *Let $\{T_n\}_{n \geq 1}$ be a sequence of trace class operators on some separable Hilbert space, having the property that their trace norms are uniformly bounded, i.e. $\sup_{n \geq 1} \|T_n\|_1 = c < \infty$. Assume that T_n converges to T in the operator norm topology. Then T is trace class.*

Proof. Since the T_n 's are compact operators, T is also compact and admits a singular value decomposition (SVD), i.e. there exist two orthonormal systems $\{f_j\}_{j \geq 1}$ and $\{g_j\}_{j \geq 1}$, together with a set of non-increasing singular values $s_j \geq 0$ such that

$$T = \sum_{j \geq 1} s_j |f_j\rangle\langle g_j|.$$

T is trace class if $\sum_{j \geq 1} s_j < \infty$. We will show that for every $N \geq 1$ we have $\sum_{j=1}^N s_j \leq c$. Let us introduce the SVD of each T_n as

$$T_n = \sum_{j \geq 1} s_j^{(n)} |f_j^{(n)}\rangle\langle g_j^{(n)}|, \quad \|T_n\|_1 = \sum_{j \geq 1} s_j^{(n)} \leq c.$$

Then

$$\sum_{k=1}^N s_k = \lim_{n \rightarrow \infty} \sum_{k=1}^N \langle f_k, T_n g_k \rangle = \lim_{n \rightarrow \infty} \sum_{k=1}^N \sum_{j \geq 1} s_j^{(n)} \langle f_k, f_j^{(n)} \rangle \langle g_j^{(n)}, g_k \rangle.$$

Using Bessel's inequality for the orthonormal systems f_k and g_k we have

$$\left| \sum_{k=1}^N \langle f_k, f_j^{(n)} \rangle \langle g_j^{(n)}, g_k \rangle \right| \leq \sqrt{\sum_k |\langle f_k, f_j^{(n)} \rangle|^2} \sqrt{\sum_k |\langle g_k, g_j^{(n)} \rangle|^2} \leq \|f_j^{(n)}\| \|g_j^{(n)}\| = 1$$

which holds for all $j, n, N \geq 1$. This implies

$$\left| \sum_{k=1}^N \sum_{j \geq 1} s_j^{(n)} \langle f_k, f_j^{(n)} \rangle \langle g_j^{(n)}, g_k \rangle \right| \leq \sum_{j \geq 1} s_j^{(n)} = \|T_n\|_1 \leq c,$$

hence $\sum_{j=1}^{\infty} s_j < \infty$ and T is trace class. □

Acknowledgments. This work was partially conducted within the France 2030 framework programme, the Centre Henri Lebesgue ANR-11-LABX-0020-01. The authors thank T. Ourmières-Bonafos for useful discussions. They are also very grateful to the CIRM (and its staff) where this work was initiated. E.S. acknowledge support from Fondecyt (ANID, Chile) through the grant # 123-1539. This work has been partially supported by CNRS International Research Project Spectral Analysis of Dirac Operators – SPEDO. H.C. acknowledge support from the Independent Research Fund Denmark–Natural Sciences, grant DFF-10.46540/2032-00005B.

REFERENCES

- [1] J.-M. Barbaroux, H. Cornean, L. Le Treust, N. Raymond, and E. Stockmeyer. Bulk-edge correspondence. *In preparation*, 2024.
- [2] J.-M. Barbaroux, L. Le Treust, N. Raymond, and E. Stockmeyer. On the semiclassical spectrum of the Dirichlet–Pauli operator. *Journal of the European Mathematical Society*, 23(10):3279–3321, 2021.
- [3] J.-M. Barbaroux, L. Le Treust, N. Raymond, and E. Stockmeyer. On the Dirac bag model in strong magnetic fields. *Pure and Applied Analysis*, 2023.
- [4] R. D. Benguria, S. , E. Stockmeyer, and H. Van Den Bosch. Self-Adjointness of Two-Dimensional Dirac Operators on Domains. *Annales Henri Poincaré*, pages 1–13, 2017.
- [5] M. V. Berry and R. J. Mondragon. Neutrino billiards: time-reversal symmetry-breaking without magnetic fields. *Proc. Roy. Soc. London Ser. A*, 412(1842):53–74, 1987.

- [6] H. D. Cornean, M. Moscolari, and K. S. Sørensen. Bulk–edge correspondence for unbounded Dirac–Landau operators. *Journal of Mathematical Physics*, 64(2):021902, feb 2023.
- [7] H. D. Cornean, M. Moscolari, and S. Teufel. General bulk-edge correspondence at positive temperature, 2022.
- [8] S. De Bièvre and J. V. Pulé. Propagating edge states for a magnetic hamiltonian. In *Mathematical Physics Electronic Journal: (Print Version) Volumes 5 and 6*, pages 39–55. World Scientific, 2002.
- [9] J. Dolbeault, M. J. Esteban, and E. Séré. On the eigenvalues of operators with gaps. Application to Dirac operators. *J. Funct. Anal.*, 174(1):208–226, 2000.
- [10] P. Elbau and G. M. Graf. Equality of bulk and edge Hall conductance revisited. *Commun. Math. Phys.*, 229(3):415–432, 2002.
- [11] G. M. Graf, H. Jud, and C. Tauber. Topology in shallow-water waves: a violation of bulk-edge correspondence. *Communications in Mathematical Physics*, 383:731–761, 2021.
- [12] M. Griesemer and H. Siedentop. A minimax principle for the eigenvalues in spectral gaps. *J. London Math. Soc. (2)*, 60(2):490–500, 1999.
- [13] J. Kellendonk and H. Schulz-Baldes. Quantization of edge currents for continuous magnetic operators. *Journal of Functional Analysis*, 209(2):388–413, 2004.
- [14] R. S. K. Mong and V. Shivamoggi. Edge states and the bulk-boundary correspondence in Dirac Hamiltonians. *Phys. Rev. B*, 83:125109, Mar 2011.
- [15] M. Reed and B. Simon. *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press, New York, 1978.
- [16] L. Schimmer, J. P. Solovej, and S. Tokus. Friedrichs extension and min–max principle for operators with a gap. 21(2):327–357, 2020.
- [17] K. M. Schmidt. A remark on boundary value problems for the Dirac operator. *Quart. J. Math. Oxford Ser. (2)*, 46(184):509–516, 1995.
- [18] C. Tauber, P. Delpierre, and A. Venaille. Anomalous bulk-edge correspondence in continuous media. *Phys. Rev. Res.*, 2:013147, Feb 2020.

(J.-M. Barbaroux) AIX MARSEILLE UNIV, UNIVERSITÉ DE TOULON, CNRS, CPT, MARSEILLE, FRANCE

Email address: barbarou@univ-tln.fr

(H. D. Cornean) DEPARTMENT OF MATHEMATICAL SCIENCES, AALBORG UNIVERSITY, SKJERNVEJ 4A, 9220 AALBORG, DENMARK

Email address: cornean@math.aau.dk

(L. Le Treust) AIX MARSEILLE UNIV, CNRS, CENTRALE MARSEILLE, I2M, MARSEILLE, FRANCE

Email address: loic.le-treust@univ-amu.fr

(N. Raymond) UNIV ANGERS, CNRS, LAREMA, INSTITUT UNIVERSITAIRE DE FRANCE, SFR MATHSTIC, F-49000 ANGERS, FRANCE

Email address: nicolas.raymond@univ-angers.fr

(E. Stockmeyer) INSTITUTO DE FÍSICA, PONTIFICIA UNIVERSIDAD CATÓLICA DE CHILE, VICUÑA MACKENNA 4860, SANTIAGO 7820436, CHILE.

Email address: stock@fis.puc.cl