

Bi-Hamiltonian in Semiflexible Polymer as Strongly Coupled System

Heeyuen Koh*

*Soft Foundry Institute, Seoul National University,
1 Gwanak-ro, Gwanak-gu, Seoul, 08826, South Korea*

Shigeo Maruyama

Mechanical Engineering Dept., The University of Tokyo, Hongo 3-7-1, Bunkyo, Tokyo, Japan

(Dated: May 21, 2024)

When many body system in Classical Mechanics is described using bi-Hamiltonian for its group of particles which forms a coarse grained bead, the damping energy from the heat bath perturbs the motion of a particle which is along the trajectory of multiple Hamiltonian system simultaneously. As managing the coarse grained modeling of the suspended semiflexible polymer like single walled carbon nanotubes(SWCNT) in the vacuum, the authors discovered that such interconnection between Hamiltonian systems as sharing a heat bath builds a strongly coupled system. The consequence of this interconnection is revealed certain form of cross correlated momentum in coarse grained system as well as in the original atomic scales. In this paper, we adapt the framework of Stochastic Thermodynamics to describe the cross correlation of bi-Hamiltonian system which replicates the macroscopic behaviour of SWCNT under the Smoluchowski picture for cross correlation between Hamiltonian systems. The numerical experiments using collision between tubes confirm the derivation and the justification of the usage of the heat diffusion to compensate the abnormality from cross correlated momentum.

I. INTRODUCTION

As of Zwanzig's nonlinear generalized Langevin equation[1], the memory effect could be the term that explains the intercorrelation between the target system and the heat bath in the equation of motion. From the integration of correlated trajectory, the memory effect offers the amount of action that is involved with the interconnection between the heat bath and the system of interest as the Hamiltonian of mean force[2–4] proposes the involved energy amount. If the coordinate system of a bead entangled with conservative force evolves in a Lagrangian perspective, the reason for the memory effect is quite evident. Yet, it is hard to specify the cause of the interaction mechanism or characteristics of the inter-connection that differentiates the system of interest whether this coupling between the system and the heat bath is weak enough so that the interconnection can be negligible or whether it is engineerable to enhance the functionality.

When the system is simplified into two Hamiltonians with coarse grained variables, which describe the motion for each degree of freedom(DOF) condition in an adiabatic environment, such as SWCNT or other quasi-one dimensional systems, the relation between each Hamiltonian system is exactly that of the system versus heat bath. Recently, Koh et al.[5, 6] found that coarse grained system like SWCNT and dsDNA that is composed of two independent variables like bond length and angle has a cross-correlated momentum between two variables during the evolution process, and they form a memory effect to each other which is strong enough to affect the trajectory of motion from Mori Zwanzig formalism[5].

In the case of SWCNT[5] and DNA[6], the diffusion of cross correlated energy between translational and rotational deformation adjusted in their equation of motion to resolve the discrepancy between CGMD and molecular dynamic simulation. The macroscopic motion of coarse grained bi-Hamiltonian system which is composed of the simple beads system to replicate the SWCNT's nonlinear bending motion[7], is almost perfectly synchronized with the atomic scale simulation result. Inspired by the energetics of Langevin equation following Sekimoto's frameworks[8], we can presume the stochastic nature of the equation of motion modified with the diffusion term. The diffusion is applied as the physics underpinning a quantifiable energetic form that resolved a memory effect in the Markovian regime without concrete phenomenological ground. Nonetheless, additional derivation is required to effectively address the cross correlated states using the diffusion term.

The heat diffusion damping which resolved the alternation of conservative force pathway is the result of fast variable dynamics which is not included in the coarse grained system, but it could be included as the Hamiltonian of mean force[2, 3, 9] defined for coarse grained system. The quantification of the Hamiltonian of mean force which is either of heat or work[10, 11] has raised several arguments, especially on its measurability and localization[11–13]. The coarse grained system in Markovian regime with the assumption of the usage of heat diffusion term could be validated with overdamping spring mass system[12], or through the energetics represented by Fokker-Plank equation[9] as its evolution equation. Nonetheless, if the projection of fast dynamics to the slow ones is related to the heat diffusion, it should be revealed with the irregularity of entropy along the structure with the entropy definition which needs to be de-

* hy_koh@snu.ac.kr

fined from memory effect or Hamiltonian of mean force for ensembles. Such express should be involved with the evolution on the fast dynamics. Therefore, new frameworks to distinguish thermal fluctuation of the molecular system upon Thermodynamics[14] need to be thoroughly investigated.

In this paper, strongly coupled two hamiltonian systems from the memory effect is analyzed using Stochastic Thermodynamics and its evolution is proved with numerical experiment. Applying Stochastic Thermodynamics to crarify cross correlation momentum of SWCNT in atomic scale trajectory explicates the intramolecular thermal energy mechanism as well as the justification of the usage of heat diffusion term in the equation of motion in coarse-grained molecular dynamcis (CGMD), which corrects the memory effect from cross correlation. In section II. Theoretical modeling is derived based on the framework by Jarzynski[14]. The data provided by molecular dynamics and coarse grained molecular dynamics is explained in section III. Simulation. The meaning of derivation which is confirmed from Simulation is described in IV Results and Discussion. The summary and further works are included in V. Conclusion.

II. THEORETICAL MODELING

A. System definition

Let a quasi-one dimensional discretely be a connected system like SWCNT with the potential energy ϕ with N_α atoms. The Hamiltonian is

$$\begin{aligned} H &= K + U, \\ K &= \sum_{\alpha}^N \sum_i^{N_\alpha} \frac{p_{\alpha i}^2}{2m_{\alpha i}}, \\ U &= \frac{1}{2} \sum_{\alpha\beta} \sum_{i,j} \phi(|r_{\alpha i} - r_{\beta j}|). \end{aligned} \quad (1)$$

α and β are the indices of a coarse grained particle. i and j are the indices for each atom. ϕ is the potential energy function between atoms. Following the expression for the Hamiltonian system and the phase space density definition by Kinjo and Hyodo[15], the description by coordinate and momenta on the center of mass of a group of atoms becomes as below:

$$\hat{R}_\alpha \equiv \frac{\sum_i m_{\alpha i} \hat{r}_\alpha}{M_\alpha}, \quad (2)$$

$$\hat{P}_\alpha \equiv \sum_i \hat{p}_{\alpha i}, \quad (3)$$

$$M_\alpha \equiv \sum_i m_{\alpha i} \quad (4)$$

\hat{R}_α and \hat{P}_α are the coordinate of the COM and the momentum of the CG particle. M_α is the mass. The phase

space density for each scale of the system should be as below:

$$\begin{aligned} f(\hat{\Gamma}(t); \Gamma) &\equiv \delta(\hat{\Gamma}(t) - \Gamma) \\ &\equiv \prod_{\alpha, i} \delta[\hat{r}_{\alpha i} - r_{\alpha i}] \delta[\hat{p}_{\alpha i} - p_{\alpha i}], \end{aligned} \quad (5)$$

$$\begin{aligned} f(\hat{\Gamma}_s(t); \Gamma_s) &\equiv \delta(\hat{\Gamma}_s(t); \Gamma_s) \\ &\equiv \prod_{\alpha} \delta[\hat{R}_\alpha - R_{\alpha i}] \delta[\hat{P}_\alpha - P_\alpha]. \end{aligned} \quad (6)$$

Here, the trajectory of the atoms and CG particles at time t are $\hat{\Gamma}(t) \equiv \{\hat{r}_{\alpha i}(t), \hat{p}_{\alpha, i}(t)\}$, and $\hat{\Gamma}_s(t) \equiv \{\hat{R}_\alpha(t), \hat{P}_\alpha(t)\}$, respectively.

which is composed of the set of the group of atoms and each group can be regarded as being connected by length and bending deformation energy, i.e. potential energy which are noted as ℓ and θ , respectively.

$$H_{CG} = H_\ell + H_\theta, \quad (7)$$

$$H_\ell = \frac{1}{2} \frac{\mathbf{P}_\ell^T \mathbf{P}_\ell}{m} + \Phi_\ell, \quad (8)$$

$$H_\theta = \frac{1}{2} \frac{\mathbf{P}_\theta^T \mathbf{P}_\theta}{I} + \Phi_\theta. \quad (9)$$

Here, H_ℓ and H_θ with the potential energy Φ_ℓ and Φ_θ , respectively. The radial deformation is supposed to be canceled out during averaging as shown in Figure 1A-b. Each CG particle is connected with bond length and angle potential energy functions as Figure 1B, and each of them belongs to independent Hamiltonian system.

B. Cross correlated momentum between Hamiltonians

When the coordinate system of a bead in a coarse grained system, defined by the bond length and angle deformation is on evolution as shown in Figure 1C-> Figure 2, such evolution slightly alters the unit axes at t and $t + \Delta t$ so that the bond length-wise deformation at t and that from the angle at $t + \Delta t$ are not orthogonal like bond and angle deformation does at the same moment. The momentum of i th CG particle is composed with the update from each Hamiltonian as below:

$$\begin{aligned} \hat{P}_i &= P_{\ell i} \hat{e}_{P_\ell}(t) + P_{\theta i} \hat{e}_{P_\theta}(t), \\ &= (P_\ell^i + P_\theta^i) \cdot \hat{e}_{\hat{P}_i}(t). \end{aligned} \quad (10)$$

Here, $\hat{e}_{P_\ell}(t)$ and $\hat{e}_{P_\theta}(t)$ are the unit vector on each momentum from H_ℓ and H_θ , respectively. $\hat{e}_{\hat{P}_i}(t)$ is the unit vector long the momentum direction. The direction of all unit vectors is decided by the coordinate of CG particles connected by the bond and angle so the unit vectors should be time-varying. The 3 DOF in the suspended system as shown in Fig. 1A makes the unit vector to be varying on time, therefore, the unit vector at $t + \Delta t$ is




FIG. 1. A. a. Schematic figure of suspended SWCNT and the system of the interest in blue shade, b. The cartoon of the coarse grained system from suspended SWCNT with two independent potential energy which builds bi-Hamiltonian system. B. The system of interest with deformation of elongation ($\Delta\ell$) and bending ($\Delta\theta$). C. Evolution of the deformation along the bond length and angle direction at t and $t + \Delta t$. The coordinate system is also developed along the macroscopic motion of the tube.

slightly disturbed from the previous state as below:

$$\hat{\mathbf{e}}_{\hat{P}_\theta}(t + \Delta t) = \hat{\mathbf{e}}_{\hat{P}_\theta}(t) + \eta \hat{\mathbf{e}}_{\hat{P}_\ell}(t), \quad (11)$$

$$\hat{\mathbf{e}}_{\hat{P}_\ell}(t + \Delta t) = \hat{\mathbf{e}}_{\hat{P}_\ell}(t) + \eta' \hat{\mathbf{e}}_{\hat{P}_\theta}(t). \quad (12)$$

η and η' are arbitrary given parameters. The canonical equation of Hamiltonian H_{CG} along the given bond length and angle simultaneously, therefore, perturbs the trajectory, $\dot{R}_\alpha \rightarrow \dot{R}_\alpha + \dot{R}'_\alpha$ as shown in Fig. 1C. Such perturbation affects the least of action in the modified Hamilton's principle as below:

$$\delta I_\sigma = \delta \int_{t_i}^{t_f} (\dot{R}_{\alpha\sigma} \hat{P}_{\alpha\sigma} - H_\sigma(\hat{R}_{\alpha\sigma}, \dot{R}_{\alpha\sigma}, t)) \neq 0. \quad (13)$$

σ is either of ℓ or θ . Lets suppose there is a perturbation term from fast variables as $H_\sigma + H'_\sigma$ which satisfies $\delta I = 0$. The evolution of f and f_s along the perturbed trajectory can be

$$\left(\frac{d}{dt} \right)_\Gamma f_s = \sum_\alpha \sum_i \left\{ \frac{\partial H_\sigma}{\partial \hat{r}_{\alpha i}} \cdot \frac{\partial}{\partial \hat{p}_{\alpha i}} - \frac{\partial H_\sigma}{\partial \hat{p}_{\alpha i}} \cdot \frac{\partial}{\partial \hat{r}_{\alpha i}} \right\} f, \quad (14)$$

$$\equiv i\mathcal{L}_s f_s + i\mathcal{L}' f_s \quad (15)$$

with

$$\hat{F}_\alpha = - \sum_{i=1}^{N_\alpha} \frac{\partial U}{\partial \hat{r}_{\alpha i}}, \quad (16)$$

$$i\mathcal{L}_s = - \sum_\alpha \left\{ \hat{F}_\alpha \cdot \frac{\partial}{\partial P_\alpha} + \frac{\hat{P}_\alpha}{M_\alpha} \cdot \frac{\partial}{\partial R_\alpha} \right\}. \quad (17)$$

Here, the evolution of f_s which is from f is proceeded separately as $i\mathcal{L}_s$ and $i\mathcal{L}'$ for coarse grained particle and fast variable dynamics effect. We are going to deal $i\mathcal{L}'$ in more detail in the next section. The perturbation H'_σ is from H_ℓ when σ is θ and vice versa.

In Mori-Zwanzig formalism, the projection operator divides the dynamic variable $g(\hat{\Gamma}) = g_P(\hat{\Gamma}) + g_Q(\hat{\Gamma})$ according to the base set $f_S(\Gamma_S(t_0); \Gamma_S)$, and adjusts the variable $g_P(\hat{\Gamma})$ which is from f_s with the evolution of $g_Q(\hat{\Gamma})$ [15]. The perturbation from $g_Q(\hat{\Gamma})$ is, therefore, as shown in Figure 2A-a. But in the case of bi-Hamiltonian system which is coupled with the memory effect, it is the landscape of the potential energy that is distorted by the perturbation during evolution as shown in Figure 2A-b. It is the reason for the additional operator $i\mathcal{L}'$ is essential. The two Hamiltonians for each orthogonal variable coexist, then the perturbation from each other can be understood as Figure 2B.

C. Jarzynski's frameworks for cross correlation

Due to the adiabatic environment, the probability of density of the fluctuation from a finite bath [16] starts from the very beginning of the definition as below:

$$\rho(z, \lambda) = \frac{\Omega_B(E_{tot} - H(z, \lambda))}{\Omega_{tot}(E_{tot})}, \quad (18)$$

$$\lim_{C \rightarrow \infty} \rho_C(z; T, \lambda) = \frac{e^{-H/T}}{Z(T, \lambda)}. \quad (19)$$

Here, $z = (\mathbf{q}, \mathbf{p})$ is the set of state variable, and λ is controllable parameter. The density of the state of the bath and the total system are $\Omega_B(E)$ and $\Omega_{tot}(E)$, respectively. The form of the probability density function, ρ in Eq. (18) becomes as Eq. (19) when the specific heat $C \rightarrow 0$ as $dn \rightarrow \infty$ where the DOF of the system is d and the number of particles is n .

A more specific condition for $C \rightarrow 0$ is derived in Appendix B to overcome the limit of the number of states as




FIG. 2. A. Dissipation and random force effect on harmonic oscillator(left) and the deformed harmonic oscillator by cross correlation in bi-Hamiltonian system(right) from Eq. (13). B. Effect of cross correlation on each Hamiltonian system H_ℓ and H_θ . The potential energy curvature is deformed and affected by another Hamiltonian system as well as its momentum. The red and yellow dot has interconnectivity from cross correlation and its reconciliation.

the system defined in the adiabatic condition. The result of derivation proves the ready-known condition $dn \rightarrow \infty$, but it also shows the finite amount of dn from the number of carbon atoms in SWCNT can afford the condition for $C \rightarrow 0$ so that we adopt the framework of Jarzynski to handle the Hamiltonian perturbation as below:

$$H = h_s(x_j; \lambda) + H_{\mathcal{E}}(y_j) + h_{SE}(x_j, y_j) + PV_{\mathcal{E}}. \quad (20)$$

The state variable x_j is for the system of interest and then $y_j = (\mathbf{q} - \mathbf{q}_j, \mathbf{p} - \mathbf{p}_j)$. $h_s(x_j; \lambda)$ is the Hamiltonian of the system of the interest which is either of H_ℓ or H_θ . $H_{\mathcal{E}}(y_j)$ is that of the thermal environment, therefore, it becomes the rest of the SWCNT tube system. $h_{SE}(x_j, y_j)$ is the energy related to both the system and the environment. P is the pressure of thermal environment and $V_{\mathcal{E}}$ is the volume of the total system in the reference[14]. The isothermal-isobaric ensemble of the system becomes as below:

$$\begin{aligned} \pi_{\lambda NPT}^{eq}(\zeta) &= \frac{1}{\mathcal{Y}_A} e^{-\beta H_{S+E}(z; \lambda) + PV_{\mathcal{E}}}, \\ &= \frac{1}{\mathcal{Y}_A} e^{-\beta H}, \end{aligned} \quad (21)$$

$$\mathcal{Y}_A = \int d\zeta e^{-\beta H} = \int dx e^{-\beta h_s(x; \lambda)} \mathcal{Z}_x^{\mathcal{E}} = \mathcal{Z}_x \mathcal{Z}_0^{\mathcal{E}}.$$

Here, H_{S+E} is $h_s(x_j; \lambda) + H_{\mathcal{E}}(y_j) + h_{SE}(x_j, y_j)$, and z is (x_j, y_j) . Then, the ensemble of the system of interest can be derived as below:

$$\begin{aligned} \rho(t) &= \pi_{\lambda NPT}^{eq} = \int dy_j \pi^{eq}(x_j, y_j), \\ &= \frac{1}{\mathcal{Z}_\lambda} e^{-\beta(h_s(x_j; \lambda) + \phi(x_j))}, \end{aligned} \quad (22)$$

$$\begin{aligned} \phi(x_j) &= \phi(x_j; N, P, T), \\ &= -\beta^{-1} \frac{\int dy_j \exp[-\beta(H_{\mathcal{E}}(y_j) + h_{SE} + PV_{\mathcal{E}})]}{\int dy \exp[-\beta(H_{\mathcal{E}}(y_j) + PV_{\mathcal{E}})]}. \end{aligned} \quad (23)$$

where

$$\mathcal{Z}_\lambda(N, P, T) = \int dx_j e^{-\beta(h_s + \phi(x_j))}. \quad (24)$$

$x_j = (\mathbf{q}_j, \mathbf{p}_j)$ are separately described for each system along $j = \theta$ or ℓ . θ represents the angle and ℓ represent the bond length. When the state variable x_j belongs to H_ℓ then x_θ becomes part of $y_j = (\mathbf{q} - \mathbf{q}_j, \mathbf{p} - \mathbf{p}_j)$. The concept of separation of x_j which is causing the heat bath variable no longer to be a complete orthogonality with the state variable of y_j has explained in Appendix A.

As $\phi(x_j)$ is the perturbation of energy from the interconnection with the thermal environment, it includes, therefore, the energy of conformation change of the system of interest by the thermal environment. For the ergodicity based on the modified Hamilton's principle in Eq. (13), ϕ can be regarded as the perturbation of Hamiltonian to satisfy the least action when its amount is equivalent to $\hat{R}_{\sigma-1} P_\sigma$ with $\sigma = \ell$ or θ . $1 - \sigma$ indicates θ or ℓ , respectively. ϕ is, therefore, the deformation caused by cross correlation of the bi-Hamiltonian system and it should be the function of \hat{R} . Note that the system of the interest is a part of SWCNT which is connected to the rest of the tubes, and we are considering the motion either of bending or elongation. The thermal environment \mathcal{E} for the Hamiltonian H_θ , is therefore, H_ℓ and connected tubes on both ends of the system.

In the next subsection, the evolution of the ensemble modified with ϕ , which is introduced as the operator $i\mathcal{L}'$ is discussed.

D. Probability Density Function in Smoluchowski equation

For resolving the discrepancy of cross correlation between Hamiltonian systems, it is supposed that the rearrangement of atoms in the target system is not inducing any motion of the group of atoms. The evolution of state variable from probability density function is, then governed by Smoluchowski equation as below:

$$\xi \frac{\partial \rho}{\partial t} = \frac{\partial}{\partial x} (U' \rho) + \frac{1}{\beta} \frac{\partial^2 \rho}{\partial x^2}. \quad (25)$$

Here, ξ is damping coefficient. U is the potential function. ρ , the probability density function can be derived from Jarzynski's frameworks as the ensemble for the target system:

$$\rho = p^{eq} = \frac{e^{-\beta[u_s + P v_s]}}{\int dx e^{-\beta[u_s + P v_s]}} = \frac{A}{Z}. \quad (26)$$

A is $e^{-\beta[u_s + Pv_s]}$, and Z is $\int dx e^{-\beta[u_s + Pv_s]}$. The LHS divided by ξ becomes:

$$\frac{\partial \rho}{\partial t} = \frac{A}{Z} \left(-\beta \frac{\partial}{\partial t} (u_s + Pv_s) \right). \quad (27)$$

The first term of RHS of Eq. (25) is:

$$\frac{\partial}{\partial x} (U' \rho) = \frac{\partial^2 U}{\partial x^2} \rho + U' \frac{\partial \rho}{\partial x} \quad (28)$$

$$\frac{\partial \rho}{\partial x} = -\frac{A^2}{Z^2} - \beta \frac{A}{Z} \frac{\partial}{\partial x} (u_x + Pv_s). \quad (29)$$

The second term of RHS of Eq.(25) is as below:

$$\frac{1}{\beta} \frac{\partial^2 f}{\partial x^2} = \frac{1}{\beta} \frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} \left(\frac{A}{Z} \right) \right), \quad (30)$$

$$= 2 \frac{A^3}{Z^3} + \frac{A^2}{Z^2} \frac{\partial^2 Z}{\partial x^2} + \frac{1}{Z} \frac{\partial^2 A}{\partial x^2}. \quad (31)$$

To make the assumption of $\frac{A}{Z} \ll 1$, we need the restriction on $|\frac{\partial A}{\partial x}| < 1/\beta = kT$. The restrain indicates that the possible number of state should be varied in range of kT between its neighbors. Otherwise $\frac{A}{Z} \ll 1$ is not satisfied so that the high order of $\frac{A}{Z}$ can not be eliminated. The full derivation of Eq.(25) is as below:

$$\begin{aligned} & \xi \frac{A}{Z} \frac{\partial}{\partial t} (u_s + Pv_s), \\ = & \frac{\partial^2 U}{\partial x^2} \frac{A}{Z} - \frac{1}{\beta} \left(-\beta \frac{A}{Z} \frac{\partial^2}{\partial x^2} (u_s + Pv_s) \right), \\ & \frac{\partial}{\partial t} (u_s + Pv_s) = \frac{1}{\xi} \frac{\partial^2}{\partial x^2} (pV_s + u_s). \end{aligned} \quad (32)$$

E. Evolution of Cross Correlation under Smoluchowski picture

In the atomic scale system, we presume that the small scale motion of the atoms, which is averaged as zero, resolves the cross correlated condition. Since such a condition has no momentum, the evolution equation in for CG scale particles becomes Smoluchowski equation. The probability density function has its governing equation as below:

$$\partial_t \rho = L_C \rho + L_D \rho + L_S \rho. \quad (33)$$


L_C is the Liouville operator with conservative force from potential energy. L_D is the Fokker-Plank operator for the dissipative and random force. Lastly, L_S is Smoluchowski operator which has its evolution as below:

$$L_S \rho = \nabla \cdot (\rho \nabla U) + T \nabla^2 \rho. \quad (34)$$

The substitution of Eq. (23) to Smoluchowski eq. is as follows in equilibrium case:

$$\Delta u_s + P \Delta v_s = \frac{1}{\xi} \frac{\partial^2}{\partial x^2} (E_{ke} + Pv_s) = 0. \quad (35)$$

A



B




FIG. 3. A. The snapshot of the MD simulation, B. The normal mode decomposition results of 1st bending mode(upper), x,y and z direction of n th mode, respectively.

The derivation is in Appendix. A. Intuitively, Eq. (35) should be zero in equilibrium condition to maintain the harmonic oscillator, and such condition can be from $\frac{\partial^2}{\partial x^2} E_{ke}$ when equivalent to $-\frac{\partial^2}{\partial x^2} Pv_s$. This indicates the boundary condition of the target system and how the target system is interconnected with geometrically contacted heat bath. In the case of SWCNT in vacuum, the geometrical heat bath is the tube segment at both sides of the target system.

$$\Delta u_s + P \Delta v_s = \frac{\partial^2}{\partial x^2} (-U_{def} + E_{ke} + Pv_s). \quad (36)$$

When the system has external stimuli which provoke deformation, The first term in RHS of Eq. (36) remained with unresolved deformation energy, U_{def} , which is deviated from the potential energy in equilibrium state as part of p^{eq} . When the system is close to equilibrium from external stimuli, RHS of Eq. (36) becomes zero. According to Eq. (36), it is E_{ke} and Pv_s which compensate the $-U_{def}$. To confirm this process of thermalization of U_{def} , the molecular dynamics simulation is conducted.




FIG. 4. Normal mode decomposition of the result from CGMD with the same configuration in Fig. 3A., A. 1st bending mode amplitude, B. N th mode along x and y direction.

III. RESULTS

To examine the response of the bi-Hamiltonian from Eq. (36), the collision between SWCNT is conducted using molecular dynamic(MD) simulation. Two (5,5) SWCNTs with 20 nm length are arranged with 90 degrees to be intercepted at the middle of the tube during collision. To invoke the collision, one of the tube is artificially bent with 5 nm depth and released to make a collision with another tube underneath the bent SWCNT as shown in Fig. 3A.

The consequence of the deformation caused by collision which draws a bending curvature on the tube is far larger than the curvature of bending from equilibrium. The trajectory of the SWCNT which is affected by the collision is analyzed after the post processing of averaging the displacement from a ring, which has 10 atoms shaping the circumference as shown in Fig. 1B. The angle from the deformation of collision on each bead is irregular, $\Delta\theta_i \neq \Delta\theta_j, i \neq j$, and it brings the perturbation $Pv_s = \eta\Delta_t\theta\Delta_t\ell$ for H_ℓ . Through the diffusion term, the double derivative $\frac{\partial^2}{\partial x^2}$ along the $x = q_\ell, p_\ell$ variable, offers the information of evolution along time variable. Among the modes that the enthalpy $U_{def} + u_s + Pv_s$, the highest wavevector will be emphasized. The expected response from the collision, therefore, maximize the shorted wavelength mode, $k = \frac{\pi}{L/N}x$ where L is the length of the tube, N is the total number of the node and x is the location of the node due to the $\frac{\partial^2}{\partial x^2}$.

The level of the amplitude analyzed by projecting $\exp(ikX_i), k = \frac{\pi}{L/N}X_i$ where X_i is the coordinate of each simple bead along y axis into simple beads system from the trajectory data of the collided tube is as shown in Fig. 3B. L is the total length of the tube and N is the number of the bead which is connected in a series as shown in Fig. 1A-b. In this numerical experiment, we have 200

simple beads which is from 10 atoms in a ring of 20nm SWCNT which has 2000 atoms. The effect of the double partial derivative is confirmed through the amplitude of $k = \frac{\pi}{L/N}$ is spiked up along the collision from another tube.

As another validation, the collision simulation using CGMD with heat diffusion damping is conducted. The CGMD that we adapted has its governing equation as the equation of motion derived from the reference[5] whose thermalization is considering the cross correlation with heat diffusion term so that the response from the external collision governed by the diffusion term in EOM is observed. The result is in Fig. 4.

IV. DISCUSSION

$\Delta u_s + P\Delta v_s$ is equivalent to heat q and work w given to the system. According to the definition of the dependency of variable x and λ , each term in Eq. (35) becomes as below:

$$\mathcal{E} = h_s + Pv_s, \quad (37)$$

$$q = \int dt \dot{x} \frac{\partial h}{\partial x} \approx \frac{\partial^2}{\partial x^2} E_{ke}, \quad (38)$$

$$w = \int dt \dot{\lambda} \frac{\partial h}{\partial \lambda} \approx \frac{\partial^2}{\partial x^2} Pv_s. \quad (39)$$

Here, \mathcal{E} is the enthalpy of the system, and q and w are heat and work applied to the system, respectively. Note that the evolution caused by Pv_s of the double partial derivative that acts as work in the system resolves the perturbation of Hamiltonian of coarse grained model, which is induced from the heat, the dynamics of the fast variable caused by another Hamiltonian.

According to VI Partial Molar Representation in Jarzynski's framework[14], a certain amount of heat is induced when the ϕ is the function of temperature as below:

$$\phi_T = T \frac{\partial \phi}{\partial T}, \quad (40)$$

$$\bar{q} = q - \Delta\phi_T, \quad (41)$$

$$\bar{s} = s - \frac{\langle \phi_T \rangle^{eq}}{T}. \quad (42)$$

The temperature dependence of Pv_s which is equivalent to ϕ and the cross correlation term $\dot{R}_{\sigma-1}P_\sigma$ is clear since the temporally defined kinetic energy of the system of interest is varying due to its macroscopic motion from given DOFs which is > 1 . The fluctuation from $\dot{R}_{\sigma-1}P_\sigma$ is dependent on the macroscopic deformation during its bending motion so that we can conjecture that $\frac{\partial \phi}{\partial T} \neq 0$ and $\frac{\partial \phi}{\partial x} \neq 0$. The irregularity of $\dot{R}_{\sigma-1}P_\sigma$ along the tube, which is from the asymmetry of the deformation from bending, can cause the unequal value of \bar{s} so that this justifies the diffusion energy from as the source of heat caused by ϕ .

The description provided by the framework of Stochastic Thermodynamics [14] justifies the perturbation of Hamiltonian Pv_s can be regarded as the cross correlation term in atomic scale system. Applying Smoluchowski equation for modified PDF from the perturbation shows the evolution of enthalpy of the target system is managed from double partial derivatives. The existence of the double partial derivatives is discussed from two independent studies. First, the trajectory from the numerical experiment of collision between two SWCNTs is analyzed using normal mode analysis. The response of n th mode of the simple beads system is well shown. The similar response is observed in its replication using heat diffusion damping CGMD.

V. CONCLUSION

In this paper, we tried to localize the scope of the observation and description of Stochastic Thermodynamics description into the part of the target molecular system further from the framework suggested by Jarzynski. The pixelization of the observation may disagree with the probability density function defined upon $N \rightarrow \infty$. Yet, the system of interest which is a Hamiltonian of coarse grained model for the semiflexible polymer like SWCNT is in the criteria defined by Finite Bath fluctuation theorem that assures the quantifiable condition for the partition function of Jarzynski frameworks.

When there are multiple Hamiltonian systems in a coarse grained model, the trajectory of the system is decided by the superposition of the information from each Hamiltonian. The disturbances of the trajectory caused by the co-existing Hamiltonian system which causes the perturbation of the conservative energy are well explained in the main text and the reference by Koh et al. [5]. The discrepancy of coarse grained model from the atomic scale energy conservation implies that there is a hidden process that is not counted as the motion of the group of atoms that forms a CG particle. Smoluchowski equation is adapted as the evolution equation of such, and the derivation using the pdf based on Jarzynski's frameworks proves the usage of heat diffusion for abnormally formed cross correlated momentum between two Hamiltonian systems manages the CGMD simulation to be a conservative system. The result of derivation is demonstrated as the normal mode decomposition data of SWCNT collision experiment in molecular dynamics simulation.

As adapting the heat diffusion process as the damping term in EOM, the simple bead system having the same initial velocity and displacement data of SWCNT collision simulation in atomic scale shows a similar impact response in normal mode decomposition. From these findings, we could conclude that the diffusion process from fast dynamics is resolving the discrepancy of the coarse grained modeling using multiple Hamiltonians.

ACKNOWLEDGMENTS

The simulation in this manuscript was conducted under the supervision of Prof. Maruyama at the University of Tokyo. The theoretical approach is solely responsible for Heeyuen Koh, Dr. or Eng.(heeyuen.koh@gmail.com) This research is supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education (NRF-2022R1I1A1A01063582). Its computational resources are from National Supercomputing Center with supercomputing resources including technical support (KSC-2020-CRE-0345). There are no conflicts to declare.

Appendix A: Fast dynamics in cross correlated momentum

The momentum \vec{p}_i of the i th atom is defined with the Cartesian coordinates as $\vec{p}_i = \vec{p}_{i\tau} + \vec{p}_{ir} + \vec{p}_{i\ell}$. Each term represents the component of the momentum along the axis defined in the system of interest. \hat{e} is the unit vector along the tube length.

$$\vec{p}_i = \vec{p}_{i,\theta} + \vec{p}_{i,\ell} + \vec{p}_{i,\Delta}. \quad (\text{A1})$$

$\vec{p}_{i,\theta}$ and $\vec{p}_{i,\ell}$ are the components of momentum belonging to elongation and bending deformation. $\vec{p}_{i,\Delta}$ is remained level of momentum, which is not included in neither of projections.

When the target system is described using two independent Hamiltonian systems for bond length deformation and angle deformation with N coarse grained particles, we are going to regard the system of interest is represented by a CG particle. Then during the evolution of each Hamiltonian system, the damping for each momentum $\vec{p}_{i,\theta}$ and $\vec{p}_{i,\ell}$ are caused by $\vec{p}_{\ell,damp}$ and $\vec{p}_{\theta,damp}$ is $\vec{p}_{\ell,damp} = \vec{p}_{i,\theta} + \vec{p}_{i,\Delta}$ and $\vec{p}_{\theta,damp} = \vec{p}_{i,\ell} + \vec{p}_{i,\Delta}$ Since $\vec{p}_{\ell,damp}$ and $\vec{p}_{\theta,damp}$ is shared by $\vec{p}_{i,\Delta}$, we have a certain correlation in atomic scale between two independent Hamiltonians, and it is not - .

The existence of this kind of correlation can resolve the abnormal discrepancy observed in coarse grained system which is defined as bi-Hamiltonian as shown in Fig. 1A-b. The damping momentum for the set of $p_j = j = \theta$ or ℓ is paired with the perturbation of potential energy U_ℓ and U_θ as below:

$$U_\ell = U_{\ell 0} + U'_\theta, \quad (\text{A2})$$

$$U_\theta = U_{\theta 0} + U'_\ell. \quad (\text{A3})$$

$U_{\ell 0}$ and $U_{\theta 0}$ is the potential function of bond length and angle when SWCNT is modeled as simple beads system. When the tube is thermalized, the influence of damping momentum perturbs the harmonic potential energy well from Eq. (13). The existence of such alternation has well known in Classical Mechanics since the elongation of the tube disturbs the neutral axis of the Euler beam and vice versa from the imperfect bending symmetry also known as Timoshenko beam theory.

Appendix B: Finite Bath Fluctuation Theorem for SWCNT coarse graining

Lets suppose we have a Hamiltonian of atomic system in the heat bath as below:

$$H_B(\{\vec{p}_i\}, \{\vec{q}_i\}) = \sum_i \frac{\vec{p}_i^2}{2m} + \sum V(|\vec{q}_i - \vec{q}_j|). \quad (\text{B1})$$

The heat capacity of the bath is defined by the energy of the bath E_B and the microcanonical temperature T_B as below:

$$C(E_B) = \left(\frac{\partial}{\partial E_B} T_B(E_B) \right)^{-1}, \quad (\text{B2})$$

$$T_B = \frac{\Phi_B(E_B)}{\Omega_B(E_B)}. \quad (\text{B3})$$

For $C \rightarrow 0$

$$\frac{\partial}{\partial E_B} T_B(E_B) = 1 - \Phi_B \cdot \frac{1}{\Omega_B^2} \cdot \frac{\partial^2 \Phi_B(E_B)}{\partial E_B^2} \rightarrow 0 \quad (\text{B4})$$

when the bath density of the state and the phase space volume Φ_B satisfies

$$\frac{\Omega_B^2}{\Phi_B} \approx \frac{\partial^2 \Phi_B(E_B)}{\partial E_B^2}. \quad (\text{B5})$$

According to Campsi et al, the phase space volume can be defined under the assumption that the phase volume of \vec{q}_i is independent of E_B as below:

$$\Phi_B(E_B)_0 = \int \prod_{i=1}^n d\vec{q}_i \int \prod_{i=1}^n d\vec{p}_i \times \theta \left(E_B - \sum_i \frac{\vec{p}_i^2}{2m} - \sum V(|\vec{q}_i - \vec{q}_j|) \right), \quad (\text{B6})$$

$$= A_{dn} (2m)^{dn/2} E_B^{dn/2} V^n \quad (\text{B7})$$

where $A_{dn} = \pi^{N/2} \Gamma(N/2 + 1)$ and $V^n = \int \prod_{i=1}^N d\vec{q}_i$ for ideal gas of the hard sphere particles in volume V which is fixed. $\theta(x)$ is a heavyside step function.

Although the system of interest in the main text may composed of hard spheres, it is connected as a single strand so that the term $\int \prod_{i=1}^N d\vec{q}_i$ is no longer independent to E_B . Instead of independent of E_B , we can bring the interdependent condition between kinetic energy and potential energy that the sum of those quantities should be always E_B so that the value of E_B in Eq. (B6) can

be divided as E_a and $E_B - E_a$ as below:

$$\begin{aligned} \Phi_B(E_B) &= \int \prod_{i=1}^n d\hat{R}_{\alpha i} \int \prod_{i=1}^n d\hat{P}_{\alpha} \times \\ &\theta \left(E_B - \sum_i \frac{\vec{p}_i^2}{2m} - \sum V(|\vec{q}_i - \vec{q}_j|) \right), \quad (\text{B8}) \\ &= \prod dE_a \delta(\hat{E}_a - E_a) \times \\ &\int \prod_{i=1}^n d\hat{P}_{\alpha i} \theta \left((E_B - E_a) - \sum_i \frac{\vec{p}_i^2}{2m} \right) \times \\ &\int \prod_{i=1}^n d\hat{R}_{\alpha} \theta \left(E_a - \sum V(|\vec{q}_i - \vec{q}_j|) \right) \quad (\text{B9}) \end{aligned}$$

Then the result of Eq. (B8) becomes

$$\Phi_B(E_B) = \prod dE_a \delta(\hat{E}_a - E_a) A_{dn}^2 (m/k)^{dn/2} \times (E_b - E_a)^{dn/2} E_a^{dn/2}, \quad (\text{B10})$$

when $V(|\vec{q}_i - \vec{q}_j|)$ is supposed to be a harmonic potential function. The assumption on E_a to divide the potential and kinetic energy is valid even the function V is not harmonic. From this result of Eq. (B10), the density of states of the bath Ω_B , $\frac{\Omega_B^2}{E_B}$ and $\frac{\partial^2 \Phi_B}{\partial E_B^2}$ which are the terms in Eq. (B4) can be as follows:

$$\begin{aligned} \Omega_B(E_B) &= \frac{\partial \Phi_B(E_B)}{\partial E_B} \\ &= dn/2 \cdot \prod \delta(\hat{E}_a - E_a) \times \\ &A_{dn}^2 (m/k)^{dn/2} (E_b - E_a)^{dn/2-1} a^{dn/2}, \quad (\text{B11}) \end{aligned}$$

$$\begin{aligned} \frac{\Omega_B^2}{E_B} &= (dn/2)^2 \prod \delta(\hat{E}_a - E_a) \times \\ &A_{dn}^2 (m/k)^{dn/2} (E_b - E_a)^{dn/2-2} E_a^{dn/2}, \quad (\text{B12}) \end{aligned}$$

$$\begin{aligned} \frac{\partial^2 \Phi_B}{\partial E_B^2} &= dn/2 \cdot (d/2 - 1) \times \\ &\prod \delta(\hat{E}_a - E_a) A_{dn}^2 (m/k)^{dn/2} (E_b - E_a)^{dn/2-2} a^{dn/2}. \quad (\text{B13}) \end{aligned}$$

Therefore,

$$\frac{\partial}{\partial E_B} \left(T_B(E_B) \right) = 1 - \frac{\Phi_B}{\Omega_B^2} \cdot \frac{\partial^2 \Phi_B(E_B)}{\partial E_B^2} \approx 0 \quad (\text{B14})$$

when

$$\begin{aligned} \frac{\partial \Phi_B}{\partial E_B} \cdot \frac{\Phi_B}{\Omega_B^2} &= \frac{dn - 1}{\frac{dn}{2}} \rightarrow 1 \\ &\text{as } dn \rightarrow \infty. \quad (\text{B15}) \end{aligned}$$

Appendix C: Simulation Condition

The AiREBO potential energy function[17] is used for carbon atoms in the SWCNT. Both tubes are arranged at 300 K with 1 ns of thermalization using Langevin thermostat[18] and 1 ns of relaxation. The artificial bending is applied after the relaxation. Fixation at both ends has zero kelvin for a unit cell of the tube. Lammmps packaged[19] is adapted. The sampling is conducted for

each 1 ps during total simulation time which is 25 ns.

The same arrangement is prepared using coarse grained molecular dynamics model. The heat diffusion damping which considers the abnormal cross correlation between two independent Hamiltonian in simple beads system is adapted. More detail is in the reference[5]. The initial configuration of the simple beads system is from the molecular dynamic simulation by averaging the location of the beads and its velocity for each unit cell which becomes a bead.

-
- [1] R. Zwanzig, *Journal of Statistical Physics* **9**, 215 (1973).
 - [2] J. G. Kirkwood, *J. Chem. Phys.* **3**, 300 (1935).
 - [3] M. Campisi, P. Talkner, and P. Hänggi, *Physical Review Letters* **102** (2009), 10.1103/PhysRevLett.102.210401.
 - [4] P. Talkner and P. Hänggi, *Reviews of Modern Physics* **92** (2020), 10.1103/RevModPhys.92.041002.
 - [5] H. Koh, S. Chiashi, J. Shiomi, and S. Maruyama, *Scientific Reports* **11** (2021), 10.1038/s41598-020-79200-6.
 - [6] H. Koh, J. Y. Lee, and J. G. Lee, (2023).
 - [7] H. Koh, J. J. Cannon, T. Shiga, J. Shiomi, S. Chiashi, and S. Maruyama, *Physical Review B* **92**, 024306 (2015).
 - [8] K. Sekimoto, *Progress of Theoretical Physics Supplement* **130** (1998), 10.1143/PTPS.130.17/1842313.
 - [9] U. Seifert, *Reports on Progress in Physics* **75** (2012), 10.1088/0034-4885/75/12/126001.
 - [10] M. Ding and X. Xing, *Physical Review Research* **5** (2023), 10.1103/PhysRevResearch.5.013193.
 - [11] P. Talkner and P. Hänggi, *Physical Review E* **94** (2016), 10.1103/PhysRevE.94.022143.
 - [12] P. Strasberg and M. Esposito, *Physical Review E* **95** (2017), 10.1103/PhysRevE.95.062101.
 - [13] P. Strasberg and M. Esposito, *Physical Review E* **101** (2020), 10.1103/PhysRevE.101.050101.
 - [14] C. Jarzynski, *Physical Review X* **7** (2017), 10.1103/PhysRevX.7.011008.
 - [15] T. Kinjo and S.-a. Hyodo, *Phys. Rev. E* **75**, 051109 (2007).
 - [16] M. Campisi, P. Talkner, and P. Hänggi, *Phys. Rev. E* **80**, 031145 (2009).
 - [17] S. J. Stuart, A. B. Tutein, and J. A. Harrison, *Journal of Chemical Physics* **112**, 6472 (2000).
 - [18] T. Schneider and E. Stoll, *Physical Review B* **17**, 1302 (1978).
 - [19] A. P. Thompson, H. M. Aktulga, R. Berger, D. S. Bolintineanu, W. M. Brown, P. S. Crozier, P. J. in 't Veld, A. Kohlmeyer, S. G. Moore, T. D. Nguyen, R. Shan, M. J. Stevens, J. Tranchida, C. Trott, and S. J. Plimpton, *Computer Physics Communications* **271** (2022), 10.1016/j.cpc.2021.108171.