

Charged gravastar model in Rastall theory of gravity

Debadri Bhattacharjee¹ and Pradip Kumar Chattopadhyay²

^{1,2}IUCAA Centre for Astronomy Research and Development (ICARD), Department of Physics,
Cooch Behar Panchanan Barma University, Vivekananda Street, District: Cooch Behar,
Pin: 736101, West Bengal, India

E-mail: debadriwork@gmail.com, pkc_76@rediffmail.com

Abstract. Gravastars are considered as one of the prime exotic type compact objects which may be found at the end state of gravitational collapse of massive stars with a view to resolve the complexities that are pertinent in case of a black hole [1]-[3]. In this paper, we analyse the role of charge on the possible formation of isotropic spherically symmetric gravastar configuration in the framework of Rastall gravity. Gravastar contains three distinct layers *viz.* i) Interior region, ii) Thin shell and iii) Exterior region. The interior region is characterised by the equation of state $p = -\rho$ that defines the repulsive outward pressure in radial direction at all points on the thin shell. The thin shell, contains ultra-relativistic stiff fluid which is denoted by the equation of state $p = \rho$ following Zel'dovich's criteria [4, 5] for cold baryonic universe, can withstand the repulsive pressure exerted by the interior region. The exterior region is the vacuum space-time represented by the Reissner-Nordström solution. In view of the above specifications, we construct and analyse a charged gravastar model in Rastall theory of gravity which represents several salient features. The basic physical attributes, *viz.* proper length, energy, entropy and equation of state parameter of the shell are investigated. In this model, it is interesting to note that for large value of the radius of hyper-surface (R) the EoS parameter of the thin shell corresponds to dark energy EoS with $\mathcal{W}(R) \rightarrow -1$. However, for small value of R the EoS parameter $\mathcal{W}(R) \rightarrow 0$, defines a dust shell. The stability of the model is ensured through the study of gravitational surface redshift and maximisation of shell entropy within the framework of Rastall theory of gravity.

Contents

1	Introduction	1
2	The Field Equations in Rastall Gravity	4
3	Gravastar Model	5
3.1	Interior Region	6
3.2	Thin Shell	6
3.3	Exterior Region	7
4	Junction Condition	8
5	Boundary Condition	9
6	Important features of gravastar in this model	9
6.1	Proper Length of the shell	10
6.2	Energy of the shell	10
6.3	Entropy of the shell	10
6.4	EoS parameter	12
7	Stability Analysis	12
7.1	Surface Redshift	12
7.2	Entropy maximisation	13
8	Conclusion	14

1 Introduction

Compact astrophysical objects mark the end state of stellar evolution. When a stellar body is unable to counter-balance its own gravitational collapse though the generation of thermal pressure, a compact object is created. The theory and evolution of compact objects attract the attention of astrophysicists as this terminal point of stellar collapse creates extreme pressure and density regimes to test different relevant theories within the context of general theory of relativity. It is conjectured that if a massive stellar body goes through supernova explosion then depending on the mass of the remnant, it either ends as a neutron star or a black hole. In 1916, Schwarzschild provided the first black hole solution from the Einstein Field Equation (henceforth EFE) in vacuum and this is termed as the most important solution of a classical charged as well as uncharged black hole. But Schwarzschild solution has two fundamental drawbacks, *viz.*, i) existence of central singularity and ii) presence of event horizon. The central singularity ($r \rightarrow 0$) is an irremovable singularity which cannot be removed by some coordinate transformation and it is called the dynamical singularity. Whereas the singularity at the Schwarzschild radius $r = r_s = \frac{2GM}{c^2}$ which in case of a black hole constitutes the event horizon at $r = 2M$ (in a system of units $G = c = 1$) is termed as the kinematic or coordinate singularity poses fundamental problematic situations. In the quantum mechanical aspect, the presence of event horizon shows that near this region the energy of a photon diverges and there are no a-priori parameters to control this deformity in the region. We know that, Hawking

temperature is inversely proportional to the mass of the black hole ($T = \frac{\hbar}{8\pi k_B GM}$) [6] which in turn shows that black holes being in thermal equilibrium with their own hawking radiation have negative specific heat which makes them unstable under thermodynamic fluctuations. Again, black holes have gigantic amount of entropy and the presence of information paradox which is another viable condition to seek for an alternative approach. Mazur and Mottola [1]-[3] extended the idea of gravitational Bose-Einstein condensate (GBEC) to hypothesize a new model of cold, dark compact object named Gravitational Vacuum Stars or abbreviated as Gravastars. The basic model of gravastar contained five layers which was then reduced to a three layer model by Visser and Wiltshire [7]. It is now widely accepted that gravastars have a three layered structure, namely the interior region, the exterior region and the intermediate thin shell separating the interior and exterior spaces. Three regions are specified by three distinct equations of states (henceforth EoS). The interior region being in de-Sitter condensate phase is characterised by positive pressure and negative energy density ($p = -\rho$) which is equivalent to the positive cosmological constant that Einstein instigated in order to modify the static universe model based on Mach's principle. The presence of negative energy density shows that the interior region exerts repulsive outward pressure at all point on the thin shell. The thin shell is hypothesized to contain ultra-relativistic stiff fluid with an EoS $p = \rho$, which is consistent with Zel'dovich criterion of EoS in the context of cold baryonic universe. Within the context of causality, it is noted that the velocity of sound and velocity of light are equal in this is region. The shell is thin but it has a finite thickness. The exterior region is the vacuum flat space-time as indicated by the EoS $p = \rho = 0$.

Mazur and Mottola [1]-[3] in their work hypothesized that gravastars are an alternative manifestation of black holes and they studied the stability of the model through the concept of entropy maximization. Visser and Wiltshire [7] investigated the dynamical stability of gravastar model against spherically symmetric perturbations of the matter or gravity field. In pursuing the stability, they developed a mathematical structure, compatible with the Mazur-Mottola hypothesis, that describes the stability of the model through some realistic values of EoS parameter. Cattoen et al. [8] imposed the pressure anisotropy on the formation gravastars. They claimed that gravastars cannot be perfect fluids and the inclusion of pressure anisotropy could be useful in supporting gravastars with high compactness. Considering associated the range of parameters, Carter [9] investigated the existence of thin shell and stability of gravastars having Reissner-Nordström [10, 11] exterior. He also explored the dependence of EoS parameter on the possible gravastar formation. Regarding the theoretical modeling of gravastars in presence of electromagnetic field, Horvat et al. [12] introduced two models. Using suitable matching conditions they obtained the stability of gravastars through dominant energy condition. In their approach, they studied the effects of electromagnetic field on the EoS parameter, velocity of sound and surface redshift. In the framework of conformal motion, Usmani et al. [13] proposed a charged gravastar model considering the Reissner-Nordström exterior and the model was further generalized to higher dimensional approach by Bhar [14]. Rahaman et al. [15] studied the viable configuration of charged gravastars in (2+1)-dimensional space-time. In a new approach, Bhattacharjee and Chattopadhyay [16] studied the formation of charged gravastars in a generalized cylindrically symmetric space-time. Considering charged shell configuration, Chan and Silva [18] studied the effect of charge on the stability of charged gravastars. In the framework of non-commutative geometry Övgün et al. [19] constructed charged thin shell gravastars.

An interesting approach to understand the structure formation and evolution of the universe is through the light of modified gravity theories. Though Einstein's general theory of

relativity is the most promising theory to unveil the mysteries of the universe, the observational evidences in favour of the notion of accelerating universe, dark energy and dark matter pose some serious challenges to the theoretical predictions of general relativity [20]-[31]. The importance and need of incorporating modified gravity has been elaborately discussed in [32]. The generalization of Einstein-Hilbert action leads to the formalism of various alternative theories of gravity, *viz.* $f(R)$ (R is the Ricci scalar) [33, 34], $f(\mathcal{T})$ (\mathcal{T} is the torsion scalar) [35], $f(R, \square R, T)$ (\square is the de Alembert's operator, and T is the trace of energy-momentum tensor) [36]-[38], $f(R, T)$ [39] etc. A large number of works have been done on gravastars in the skeleton of modified gravity theories [40]-[45]. In comparison to general theory of relativity, the interesting feature about these theories is the non-conservation of energy-momentum tensor which leads to the non-minimal coupling of geometry and matter fields. This intriguing aspect of null covariant divergence is the main motivation for us to model a charged gravastar system in Rastall theory of gravity as introduced by P. Rastall [46, 47]. In this approach, the covariant derivative of energy-momentum tensor is proportional to the divergence of Ricci scalar curvature which is expressed as $\nabla_\nu T^\nu_\mu = \eta \mathfrak{R}_{,\mu}$.

Rastall theory provides a straightforward and tractable form for the EFE which yields interesting features in astrophysical and cosmological aspects. In the recent past, Rastall theory has been found to be successful in the formalism of black hole solutions [48], charged and rotating NUT black holes [49], rotating black holes [50], Kerr-Newman-AdS black holes [51], black holes inspired by non-commutative geometry [52], nonlinear charged black holes [53] etc.. Rastall theory has also been important in the compact star modeling. In a general approach, Mota et al. [54] described the effects of Rastall gravity on compact astrophysical objects. Using the Krori and Barua metric ansatz [55] Abbas and Shahzad [56] studied the anisotropic solutions of compact stars in Rastall theory. In another work, Abbas and Shahzad [57] explored the same formalism by varying the cosmological constant. Neutron stars in the framework of Rastall gravity was studied by Oliveira et al. [58]. The effect of Rastall gravity on the basic characteristic features i.e. mass, radius, sound velocity etc. of PSR J0740+6620 was studied in Ref. [59]. Considering perfect fluid spheres, the impact of Rastall parameter was investigated in Ref. [60]. In literature, there exists numerous number of studies that depict the impact of Rastall theory on the astrophysical as well as cosmological aspect [61]-[64].

The arrangement of this paper is done as follows. In section 2, we have set up the Einstein-Maxwell field equations along with the notion of conservation of energy-momentum tensor in the framework of Rastall gravity. In section 3, we derive the solutions of the field equations in the three distinct layers of gravastar, *viz.*, interior region, thin shell and exterior vacuum space-time for the characteristic EoS of the three regions. In this section, we have also obtained the expression for the pressure as well as matter density of the thin shell. Smooth matching of the interior and exterior solutions at the hyper-surface ($r = R$) is ensured in the section 4. Section 5 deals with the boundary conditions, i.e., the matching of the interior and thin shell solutions at the interior boundary ($r = r_1$) and thin shell and exterior solutions at the exterior boundary ($r = r_2$) to determine the numerical values of the necessary constants. Some fundamental characteristics of a gravastar namely proper length, energy and entropy of the thin shell, the EoS parameter of the hyper-surface along with the impact of charge on these attributes are studied in the section 6. In section 7, we have analysed the stability of the model on the basis of surface redshift parameter (Z_s) and the consideration of entropy maximisation. And finally in section 8 we have discussed the major findings of the paper.

2 The Field Equations in Rastall Gravity

We begin with a short introduction of Rastall theory as introduced by P. Rastall [46, 47]. In the original idea [47], Rastall modified the Einstein theory of gravity by considering the non-vanishing divergence of energy-momentum tensor which is proportional to the covariant derivative of Ricci Scalar expressed as,

$$\nabla_\nu T_\mu^\nu = \beta_\mu, \quad (2.1)$$

where β sustains its form in curved space-time but vanishes in a flat geometry. We note that $\beta_\mu = \eta \mathfrak{R}_{,\mu}$, where η is a constant, $\mathfrak{R} = g^{\mu\nu} \mathfrak{R}_{\mu\nu}$ is the Ricci Scalar and T_μ^ν is energy-momentum tensor expressed as $T_\mu^\nu = (T_\mu^\nu)_m + (T_\mu^\nu)_q$, where the subscripts define the matter and electromagnetic (charge) sectors respectively. In the context of Rastall theory, the consideration of a non-minimal coupling between matter field and geometry leads to the modification of Einstein's tensor written as:

$$G_{\mu\nu} + k\eta g_{\mu\nu} \mathfrak{R} = kT_{\mu\nu}. \quad (2.2)$$

The relation connecting Ricci scalar and energy-momentum tensor is modified as:

$$\mathfrak{R}_{\mu\nu} + (k\eta - \frac{1}{2})g_{\mu\nu} \mathfrak{R} = kT_{\mu\nu}. \quad (2.3)$$

In Rastall gravity k plays the role of gravitational coupling parameter. Contracting (2.3) one obtains, $\mathfrak{R} = \frac{kT_\mu^\mu}{4k\eta - 1}$. In the present case, we exclude $k\eta = \frac{1}{4}$ as it leads to $T_\mu^\mu = 0$ and $\mathfrak{R} = 0$ which in turn reduces to Einstein theory with $\mathfrak{R}_{\mu\nu} = 0$. We consider $\zeta = k\eta$ as the dimensionless Rastall parameter. Therefore, in the units of $G = c = 1$, the gravitational coupling parameter k and constant η are expressed as:

$$k = 8\pi \frac{4\zeta - 1}{6\zeta - 1}, \quad (2.4)$$

and

$$\eta = \frac{\zeta(6\zeta - 1)}{8\pi(4\zeta - 1)}. \quad (2.5)$$

It is evident from Eq. (2.4) that for $\zeta = 0$ k reduces to Einstein gravitational constant. From Eqs. (2.4) and (2.5) we can see that for $\zeta = \frac{1}{6}$, k diverges and again we exclude $\zeta = \frac{1}{4}$ for Rastall gravity properties. In relativistic units, $G = c = 1$ the field equations in Rastall theory of gravity take the form:

$$G_{\mu\nu} + \zeta g_{\mu\nu} \mathfrak{R} = 8\pi T_{\mu\nu} \left(\frac{4\zeta - 1}{6\zeta - 1} \right). \quad (2.6)$$

The static spherically symmetric space-time is expressed as:

$$ds^2 = -e^{2\nu(r)} dt^2 + e^{2\lambda(r)} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (2.7)$$

In eq. (2.6), $T_{\mu\nu} = (T_{\mu\nu})_m + (T_{\mu\nu})_q$, is the composite energy-momentum tensor where the subscripts depict the matter sector and electromagnetic(charge) distribution respectively. The perfect fluid consideration of the interior matter sector is expressed in the following form:

$$(T_{\mu\nu})_m = (p + \rho)u_\mu u_\nu + pg_{\mu\nu}, \quad (2.8)$$

where ρ is the energy density, p is the isotropic pressure and u^μ ($u^\mu u_\mu = -1$) is the four-velocity of the perfect fluid in consideration. On the other hand, the electromagnetic(charge) distribution takes the form:

$$(T_{\mu\nu})_q = \frac{1}{4\pi}(F_\mu^\sigma F_{\nu\sigma} - \frac{1}{4}g_{\mu\nu}F_{\sigma\beta}F^{\sigma\beta}). \quad (2.9)$$

Here, $F_{\mu\nu}$ is the Faraday-Maxwell tensor defined by the form, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, where $A_{\mu\nu}$ is the electromagnetic four potential. The corresponding Maxwell electromagnetic field equations are

$$(\sqrt{-g}F^{\mu\nu})_{,\nu} = 4\pi J^\mu \sqrt{-g} \quad , \quad F_{[\mu\nu,\delta]} = 0. \quad (2.10)$$

Here, J^i is the electric four current obeying $J^i = \sigma u^i$ where σ is the charge density and the corresponding electric field intensity E is defined through the charge parameter q as,

$$E = \frac{q}{4\pi r^2}. \quad (2.11)$$

Within the framework of Rastall gravity, eqs. (2.6)-(2.9) fabricates the following set of tractable field equations:

$$\frac{2\lambda'e^{-2\lambda}}{r} + \frac{(1-e^{-2\lambda})}{r^2} + \zeta e^{-2\lambda} \left[2\nu'' + 2(\nu')^2 - 2\lambda'\nu' + \frac{4}{r}(\nu' - \lambda') + \frac{2}{r^2}(1-e^{2\lambda}) \right] = \frac{(4\zeta - 1)}{6\zeta - 1} \left(8\pi\rho + \frac{q^2}{r^4} \right), \quad (2.12)$$

$$\frac{2\nu'e^{-2\lambda}}{r} - \frac{(1-e^{-2\lambda})}{r^2} - \zeta e^{-2\lambda} \left[2\nu'' + 2(\nu')^2 - 2\lambda'\nu' + \frac{4}{r}(\nu' - \lambda') + \frac{2}{r^2}(1-e^{2\lambda}) \right] = \frac{(4\zeta - 1)}{6\zeta - 1} \left(8\pi p - \frac{q^2}{r^4} \right), \quad (2.13)$$

$$e^{-2\lambda} \left[\nu'' + (\nu')^2 - \lambda'\nu' + \frac{\nu' - \lambda'}{r} \right] - \zeta e^{-2\lambda} \left[2\nu'' + 2(\nu')^2 - 2\lambda'\nu' + \frac{4}{r}(\nu' - \lambda') + \frac{2}{r^2}(1-e^{2\lambda}) \right] = \frac{(4\zeta - 1)}{6\zeta - 1} \left(8\pi p + \frac{q^2}{r^4} \right). \quad (2.14)$$

Here, ‘ \prime ’ denotes the differentiation with respect to the radial parameter r and we have considered the formalism in the geometrised unit of $G = c = 1$. The conservation of energy-momentum tensor in Rastall theory leads to

$$p' + (p + \rho)\nu' - \frac{\zeta}{4\zeta - 1}(3p' - \rho') = 0. \quad (2.15)$$

It is interesting to note that, in case of $\zeta \rightarrow 0$ one retains the Einstein theory of gravity.

3 Gravastar Model

In the preceding section we will study the structure of gravastar in Rastall gravity regime.

3.1 Interior Region

The interior region of a gravastar has been hypothesised to be a vacuum de-Sitter space characterised by the EoS, $p = -\rho$ [1]-[3]. The negative or repulsive pressure in this region is responsible for the radial outward force on all point on the shell which counterbalances the inward gravitational collapse. On the other hand, the negative energy density in this region is equivalent to the positive cosmological constant. This particular choice of EoS is also termed as the ‘degenerate vacuum’ or ‘ ρ -vacuum’ and it is a prime example of dark energy EoS. Using the EoS in eq. (2.15) we obtain, $p = -\rho = -\rho_c$, where ρ_c is a constant implying the constant interior density. Applying the EoS in eqs. (2.12) to (2.14), we obtain the form of metric potential $\lambda(r)$ as:

$$e^{-2\lambda(r)} = 1 + \frac{1}{3}c_1 r^2 - \frac{k}{8\pi} \frac{q^2}{r^2} + \frac{c_2}{r}, \quad (3.1)$$

where c_1 , c_2 are the integration constants. To ensure the regularity of the solution at the center, one can demand $c_2 = 0$. Therefore, eq. (3.1) reduces to

$$e^{-2\lambda(r)} = 1 + \frac{1}{3}c_1 r^2 - \frac{k}{8\pi} \frac{q^2}{r^2}. \quad (3.2)$$

From eqs. (2.13) and (2.14) we get:

$$e^{2\nu(r)} = c_3 \left(1 + \frac{1}{3}c_1 r^2 - \frac{k}{8\pi} \frac{q^2}{r^2} \right), \quad (3.3)$$

where c_3 is an integration constant. It is to be noted that eqs. (3.2) and (3.3) are non-singular solutions. Therefore, the problem of central singularity in the interior region of the gravastar is removed.

The active gravitational mass of the interior regions is obtained as:

$$M_{active} = \int_0^{r_1} \left(\rho + \frac{q^2}{8\pi r^4} \right) dr = \frac{4\pi\rho_c r_1^3}{3} - \frac{q^2}{2r_1}. \quad (3.4)$$

3.2 Thin Shell

The intermediate thin shell contains ultra-relativistic fluid obeying the EoS $p = \rho$. In the backdrop of cold baryonic universe, Zel’dovich [4, 5] originated the idea of this type of fluid and it is also termed as ‘stiff fluid’. In the present work, we may argue that this kind of approximation may be taken into consideration due to the thermal fluctuations with zero chemical potential or due to the conservation of number density of gravitational quanta at zero temperature. Several researchers [65]-[70] have used this stiff fluid to investigate different cosmological as well as astrophysical scenarios. The non-vacuum thin shell approximation along with the stiff fluid EoS makes it a difficult task to obtain the exact solutions for this region. However, analytical solutions can be obtained using the thin shell approximation i.e. $0 < e^{-2\lambda} \ll 1$. Following the thin shell approximation along with the EoS and eqs. (2.12)-(2.14), we can obtain the following set of equations:

$$\frac{2\lambda' e^{-2\lambda}}{r} + \frac{1}{r^2} + \zeta e^{-2\lambda} \left[-2\lambda'\nu' - \frac{4\lambda'}{r} \right] - \frac{2\zeta}{r^2} = \frac{(4\zeta - 1)}{6\zeta - 1} \left(8\pi\rho + \frac{q^2}{r^4} \right), \quad (3.5)$$

$$-\frac{1}{r^2} - \zeta e^{-2\lambda} \left[-2\lambda'\nu' - \frac{4\lambda'}{r} \right] + \frac{2\zeta}{r^2} = \frac{(4\zeta - 1)}{6\zeta - 1} \left(8\pi p - \frac{q^2}{r^4} \right), \quad (3.6)$$

$$e^{-2\lambda} \left[-\lambda'\nu' - \frac{\lambda'}{r} \right] - \zeta e^{-2\lambda} \left[-2\lambda'\nu' - \frac{4\lambda'}{r} \right] + \frac{2\zeta}{r^2} = \frac{(4\zeta - 1)}{6\zeta - 1} (8\pi p + \frac{q^2}{r^4}). \quad (3.7)$$

From eqs. (3.6) and (3.7) we get

$$\nu' = -\frac{1}{r} - \frac{e^{2\lambda}(r^2(1 - 6\zeta) + q^2(8\zeta - 2))}{\lambda'r^4(6\zeta - 1)}. \quad (3.8)$$

Using EoS $p = \rho$ in eqs. (3.5), (3.6) and (3.8) we obtain

$$e^{-2\lambda} = \frac{(4\zeta - 1)^2}{(6\zeta - 1)(2\zeta - 1)} \frac{q^2}{r^2} + 2 \frac{(4\zeta - 1)}{2\zeta - 1} \ln(r) + c_4. \quad (3.9)$$

Using eq. (3.9) in eq. (3.8) we get

$$\nu = \frac{(6 - 16\zeta) \ln(r) + (2\zeta - 1) \ln[q^2(1 - 4\zeta) + r^2(6\zeta - 1)]}{(8\zeta - 2)} + c_5, \quad (3.10)$$

where c_4 and c_5 are integration constants. Using eq. (3.10) in eq. (2.15) we obtain the matter density as well as pressure of the thin shell as:

$$p = \rho = \rho_0 e^{\frac{2(8\zeta - 3) \ln(r) + (1 - 2\zeta) \ln[q^2(1 - 4\zeta) + r^2(6\zeta - 1)]}{(2\zeta - 1)}}. \quad (3.11)$$

The variation of the matter density as well as pressure with the charge and shell thickness is shown in figure 1. It is to be noted here that the matter density or pressure increases towards the outer edge of the shell and it decreases with the increment of charge.




Figure 1. Variation of matter density or pressure with radius r (km).

3.3 Exterior Region

The exterior region for gravastar obeys the EoS $p = \rho = 0$ which describes a vacuum space-time. Coupling the EoS with eqs. (2.12) and (2.13) we obtain

$$\lambda' + \nu' = 0. \quad (3.12)$$

In the above consideration, we use the line element for the exterior region to be designated by the well known Reissner-Nordström [10, 11] metric expressed as:

$$ds^2 = -\left(1 - \frac{2M}{r} + \frac{q^2}{r^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{q^2}{r^2}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (3.13)$$

where M is the total mass and q is the charge of the gravitating system.

4 Junction Condition

The structure of a charged gravastar contains three layers. The interior, intermediate thin shell and the exterior region where the thin shell separates the the interior and exterior geometry. According to the fundamental junction condition, the smooth matching of the interior and exterior metric at the hyper-surface $r = R$ is ensured by Israel criterion [71, 72]. It is to be noted here that even though the metric coefficients are continuous at the interface, their derivatives are not necessarily continuous. We consider Darmois-Israel junction condition [71]-[73] to compute the surface stresses at the junction interface. Lanczos equation [74]-[77] expresses the intrinsic surface energy tensor $S_{\gamma\beta}$ in the following form:

$$S_{\beta}^{\gamma} = -\frac{1}{8\pi}(K_{\beta}^{\gamma} - \delta_{\beta}^{\gamma}K_{\kappa}^{\kappa}), \quad (4.1)$$

here, $K_{\gamma\beta} = K_{\gamma\beta}^{+} - K_{\gamma\beta}^{-}$, where (+) and (-) sign indicate the value of $K_{\gamma\beta}$ at the exterior and interior interfaces respectively. Now the form of second fundamental $K_{\gamma\beta}^{\pm}$ is given below:

$$K_{\gamma\beta}^{\pm} = -\eta_{\tau}^{\pm} \left(\frac{\partial^2 X_{\tau}}{\partial \zeta^{\gamma} \partial \zeta^{\beta}} + \Gamma_{\alpha\beta}^{\tau} \frac{\partial X_{\alpha}}{\partial \zeta^{\gamma}} \frac{\partial X_{\beta}}{\partial \zeta^{\beta}} \right). \quad (4.2)$$

The double sided normal on the surface is defined as,

$$\eta_{\tau}^{\pm} = \pm \left| g^{\alpha\beta} \frac{\partial f}{\partial x^{\alpha}} \frac{\partial f}{\partial x^{\beta}} \right|^{-\frac{1}{2}} \frac{\partial f(r)}{\partial x^{\tau}}, \quad (4.3)$$

with $\eta^{\tau}\eta_{\tau} = 1$. Following Lanczos equation [74]-[77], the surface stress-energy tensor at the boundary of the interface is defined as, $S_{\gamma\beta} = \text{diag}(-\Sigma, \mathfrak{P}, \mathfrak{P}, \mathfrak{P})$, where Σ and \mathfrak{P} are the surface energy density and the surface pressure respectively and they are expressed as:

$$\Sigma = -\frac{1}{4\pi R} \left(\sqrt{f(r)} \right)_{-}^{+}, \quad (4.4)$$

$$\mathfrak{P} = -\frac{\Sigma}{2} + \frac{1}{16\pi} \left(\frac{f'(r)}{\sqrt{f(r)}} \right)_{-}^{+}, \quad (4.5)$$

where, $f(r)_{\pm}^{\pm}$ represents the g_{rr} components of the exterior and interior regions respectively. Using eqs. (3.2) and (3.13), eqs. (4.4) and (4.5) reduces to,

$$\Sigma = \frac{1}{4\pi R} \left[\sqrt{1 + \frac{c_1 R^2}{3} + \frac{kq^2}{8\pi R^2}} - \sqrt{1 - \frac{2M}{R} + \frac{q^2}{R^2}} \right], \quad (4.6)$$

$$\mathfrak{P} = \frac{1}{8\pi R} \left[\frac{1 - \frac{M}{R}}{\sqrt{1 - \frac{2M}{R} + \frac{q^2}{R^2}}} - \frac{1 + \frac{2}{3}c_1 R^2}{\sqrt{1 + \frac{1}{3}c_1 R^2 - \frac{kq^2}{8\pi R^2}}} \right]. \quad (4.7)$$

The mass of the gravastar shell is obtained as:

$$M_{shell} = 4\pi R^2 \Sigma = R \left(\sqrt{1 + \frac{c_1 R^2}{3} + \frac{kq^2}{8\pi R^2}} - \sqrt{1 - \frac{2M}{R} + \frac{q^2}{R^2}} \right). \quad (4.8)$$

The total mass of the gravastar in terms of the mass of the shell can be written as:

$$M = \frac{4M_{shell} \sqrt{6\pi(24\pi R^2 + 8c_1 \pi R^4 - 3kq^2)} - 8c_1 \pi R^4 + 24\pi q^2 + 3kq^2 - 24\pi M_{shell}^2}{48\pi R}. \quad (4.9)$$

5 Boundary Condition

Analysis of the physical behaviour of this model of charged gravastar depends on the constants arising through mathematical formalism. In this section, we analyse the boundary condition by the matching the interior region and the thin shell at the interface $r = r_1$ and similarly matching the thin shell and the exterior region at the boundary $r = r_2$. By the matching the solutions we obtain the numerical values of the necessary constants namely c_1, c_3, c_4 and c_5 . Here we make use of eq. (4.6) and the concept that at the boundary $r = r_2$ the surface energy density vanishes [78].

(i) At $r = r_1$ we match the interior and thin shell solutions:

$$1 + \frac{1}{3}c_1r_1^2 - \frac{k}{8\pi} \frac{q^2}{r_1^2} = \frac{(4\zeta - 1)^2}{(6\zeta - 1)(2\zeta - 1)} \frac{q^2}{r_1^2} + 2 \frac{(4\zeta - 1)}{2\zeta - 1} \ln(r_1) + c_4. \quad (5.1)$$

$$c_3 \left(1 + \frac{1}{3}c_1r_1^2 - \frac{k}{8\pi} \frac{q^2}{r_1^2} \right) = e^{2 \left(\frac{(6-16\zeta)\ln(r_1) + (2\zeta-1)\ln[q^2(1-4\zeta) + r_1^2(6\zeta-1)]}{(8\zeta-2)} + c_5 \right)}. \quad (5.2)$$

(ii) At $r = r_2$ we match the thin shell and exterior solutions:

$$\frac{(4\zeta - 1)^2}{(6\zeta - 1)(2\zeta - 1)} \frac{q^2}{r_2^2} + 2 \frac{(4\zeta - 1)}{2\zeta - 1} \ln(r_2) + c_4 = \left(1 - \frac{2M}{r_2} + \frac{q^2}{r_2^2} \right). \quad (5.3)$$

$$e^{2 \left(\frac{(6-16\zeta)\ln(r_2) + (2\zeta-1)\ln[q^2(1-4\zeta) + r_2^2(6\zeta-1)]}{(8\zeta-2)} + c_5 \right)} = \left(1 - \frac{2M}{r_2} + \frac{q^2}{r_2^2} \right). \quad (5.4)$$

We consider interior radius $r_1 = 10$ Km [78] and accounting for the infinitesimally small thickness of the shell, we take the range of outer radius as $r_2 = 10.001 - 10.009$ Km. within the characteristic limit of Rastall theory, we consider the value of Rastall parameter as $\zeta = 0.23$. The prediction [78] that the surface energy density vanishes at the outer radius of the gravastar leads to the numerical evaluation of the necessary constants. Here, we have taken the mass of the gravastar $M = 2.75 M_\odot$ and the notion of $\frac{2M}{r} < \frac{8}{9}$ is suitably obeyed. It is also noted that this arbitrary combination of radii and mass provide a unique solution as the amount of charge is varied. We tabulate the numerical values of all the constants in tab 1.

q	c_1	c_4	c_5
0	-0.0243	-0.4925	20.275
2	-0.0233	-0.451	20.345
4	-0.0205	-0.3275	20.476
6	-0.0158	-0.122	20.567

Table 1. Numerical evaluation of necessary constants with the variation of charge q.

6 Important features of gravastar in this model

The finite slice of thickness of the thin shell, the extreme high density ($\gtrsim 10^{15} g/cm^3$) in this region and through the use of table 1, we have analysed the physical attributes of a gravastar, *viz*, proper length, energy, entropy and the EoS parameter of the shell.

6.1 Proper Length of the shell

The thickness of the thin shell is infinitesimally small ($\epsilon = r_2 - r_1 \ll 1$) [1]-[3] and it is the partition between the interior and the exterior region. Using eq. (3.9) the expression of the proper length is obtained as:

$$\ell = \int_{r_1}^{r_2} e^\lambda dr = \int_{r_1}^{r_2} \frac{dr}{\sqrt{\frac{(4\zeta-1)^2}{(6\zeta-1)(2\zeta-1)} \frac{q^2}{r^2} + 2 \frac{(4\zeta-1)}{(2\zeta-1)} \ln(r) + c_4}}. \quad (6.1)$$

Using eq. (6.1) and table 1, we graphically represent the proper length of the shell with shell thickness (ϵ). It is noted here that as the thickness of the shell increases the proper length




Figure 2. Variation of proper length of the shell ℓ (Km) against shell thickness ϵ (Km).

also increases. On the other hand, the increment of charge results in a decrease in the proper length. Similar results are also obtained in Refs. [17, 41].

6.2 Energy of the shell

The shell region is characterised by the EoS $p = \rho$, which is the special class of barotropic EoS $p = \omega\rho$, with $\omega = 1$. Maur-Mottola in their model hypothesised that the thin shell contains ultra-relativistic stiff fluid which satisfies Zel'dovich's criteria [4, 5]. The energy contained in case of a charged gravastar is expressed as:

$$\mathcal{E} = \int_{r_1}^{r_2} 4\pi r^2 \rho dr. \quad (6.2)$$

Using eq. (3.11) in eq. (6.2) we obtain the mathematical form of the energy contained in the shell. Due to the complexity of the equation, we have opted for the graphical representation of energy of a charged gravastar. From figure 3, it is evident that more charged fluid is found towards the outer edge of the shell. Moreover, the increase in charge results in a decrease in energy. Similar result has been obtained in the study of Bhar and Rej [41].

6.3 Entropy of the shell

Entropy is a measure of disorderness in mechanical systems. According to the Mazur-Mottola model [1]-[3], the entropy of the vacuum interior is zero and the thin shell is solely responsible




Figure 3. Variation of energy of the shell \mathcal{E} (Km) against shell thickness ϵ (Km).

for the entropy considerations. It is evaluated on the basis of the entropy function of the form:

$$S = 4\pi \int_{r_1}^{r_2} \mathfrak{s}(r) r^2 e^\alpha dr. \quad (6.3)$$

Here $\mathfrak{s}(r) = \frac{\alpha^2 k_B^2 T(r)}{4\pi \hbar^2} = \frac{\alpha k_B}{\hbar} \sqrt{\frac{p}{2\pi}}$ is the entropy density corresponding to a local specific temperature $T(r)$. α is a dimensionless constant which is consider $\alpha = 1$ without any loss of generality, k_B is the Boltzmann constant, $\hbar = \frac{h}{2\pi}$ is the Planck constant. Using Eq. (3.11) in eq. (6.3) we obtain the total entropy in the Planckian units ($\hbar = k_B = 1$) as:

$$S = 2\sqrt{2\pi} \int_{r_1}^{r_2} r^2 \sqrt{\frac{\rho_0 e^{\frac{2(8\zeta-3)\ln(r)+(1-2\zeta)\ln[q^2(1-4\zeta)+r^2(6\zeta-1)]}{(2\zeta-1)}}}{\sqrt{\frac{(4\zeta-1)^2}{(6\zeta-1)(2\zeta-1)} \frac{q^2}{r^2} + 2\frac{(4\zeta-1)}{(2\zeta-1)} \ln(r) + c_4}}} dr. \quad (6.4)$$

The variation of entropy of the thin shell with the variation of charge and shell thickness is shown in figure 4. A linear relationship between entropy and shell thickness is obtained and




Figure 4. Variation of entropy of the shell S against shell thickness ϵ (Km).

it is also noted that as the charge increases, the entropy decreases. Usmani et al. [13] showed that the entropy of a charged gravastar depends on the thickness of the shell which is found to be true in our model too. On the other hand, similar nature of result is also obtained in Yusuf et al. [17].

6.4 EoS parameter

The EoS parameter at the hyper-surface $r = R$ is expressed as:

$$\mathcal{W} = \frac{\mathfrak{P}}{\Sigma}. \quad (6.5)$$

Substituting the values of Σ and \mathfrak{P} from eqs. (4.6) and (4.7) in eq. (6.5), we get

$$\mathcal{W}(R) = \frac{\left[\frac{1 - \frac{M}{R}}{\sqrt{1 - \frac{2M}{R} + \frac{q^2}{R^2}}} - \frac{1 + \frac{2}{3}c_1 R^2}{\sqrt{1 + \frac{1}{3}c_1 R^2 - \frac{kq^2}{8\pi R^2}}} \right]}{2 \left[\sqrt{1 + \frac{c_1 R^2}{3} + \frac{kq^2}{8\pi R^2}} - \sqrt{1 - \frac{2M}{R} + \frac{q^2}{R^2}} \right]}. \quad (6.6)$$

To obtain a real solution of $\mathcal{W}(R)$, we impose the restriction $\frac{2M}{R} - \frac{q^2}{R^2} < 1$ or $2M < (R + \frac{q^2}{R})$ which is in essence satisfied by eq. (3.13). The sign of $\mathcal{W}(R)$ depends on the sign of numerator or denominator. Interestingly, for large value of R , the EoS steps into the dark energy EoS with $\mathcal{W}(R) \rightarrow -1$. On the other had, for small value of R and suitable combination of R and c_2 , $\mathcal{W}(R) \rightarrow 0$, which defines a dust shell.

7 Stability Analysis

This section is devoted to the stability analysis of the present theoretical model. The two relevant methods employed to study the stability of gravastar in this model are i) Surface Redshift and ii) Entropy maximisation.

7.1 Surface Redshift

Investigation of gravitational surface redshift can be considered as a key ingredient to analyse the stability of a gravastar. In fact, this study can be useful to obtain information regarding the detection of gravastars. The surface redshift is expressed as $Z_s = \frac{\lambda_e - \lambda_o}{\lambda_e}$, where λ_o and λ_e are the observed and emitted signals respectively. Buchdahl asserted that for a static, isotropic perfect fluid distribution the the value of surface redshift should not exceed 2, i.e., $Z_s < 2$ [79]-[81]. Barraco and Hamity [82] further stated that $Z_s < 2$ is also a valid condition in absence of cosmological constant. But Böhmer and Harko [81] showed that the presence of anisotropy in the stellar configuration extends the surface redshift to $Z_s \leq 5$. Ivanov in his study of anisotropic stars [83] further extended the limit of surface redshift from $Z_s < 3.84$ to $Z_s \leq 5.211$. DeBenedictis et al. [84] examined the stability of gravastars through axial perturbations and they also studied the surface redshift parameter in gravastars. Through their study that have found conformity in the surface redshift consideration with the above discussion.

The surface redshift is expressed as:

$$Z_s = -1 + \frac{1}{\sqrt{g_{tt}}}. \quad (7.1)$$

From eq. (3.10) and $g_{tt} = e^{2\nu}$ we obtain:

$$Z_s = -1 + \frac{1}{\sqrt{e^{2\left(\frac{(6-16\zeta)\ln(r)+(2\zeta-1)\ln[q^2(1-4\zeta)+r^2(6\zeta-1)]}{8\zeta-2} + c_5\right)}}}. \quad (7.2)$$

We have represented the radial variation of surface redshift in figure 5. We have found that the Z_s increases with the shell thickness and decreases with the increment of charge. In light




Figure 5. Radial variation of surface redshift (Z_s).

of the above discussion we can claim that our present model of charged gravastar is a stable and physically viable.

7.2 Entropy maximisation

In this section we have used the same method as employed by Mazur and Mottola in their original work [1]-[3] to check the stability criteria of gravastar model. To ensure a stable model we have applied the procedure of maximisation of entropy functional with respect to the mass function $M(r)$ and checked the sign of the second variation of the entropy functional. The first variation must vanish at the thin shell boundaries at $r = r_1$ and $r = r_2$, i.e., $\delta S = 0$. The entropy functional is written as:

$$S = \frac{\sqrt{2}\alpha k_B}{2\hbar} \int_{r_1}^{r_2} r dr \left(\frac{dM}{dr}\right)^{\frac{1}{2}} \frac{1}{\sqrt{g_{rr}}}, \quad (7.3)$$

where, α is a dimensionless constant, k_B is Boltzmann constant, $\hbar = \frac{h}{2\pi}$ is Planck constant and $g_{rr} = 1 - \frac{2M}{r} + \frac{q^2}{r^2}$. Using eqs. (3.9), (3.13) and (5.1) we can obtain the mass $M(r)$ in Rastall gravity in the form:

$$M = -r \left[\frac{(4\zeta - 1)^2}{2(2\zeta - 1)(6\zeta - 1)} q^2 \left(\frac{1}{r^2} - \frac{1}{r_1^2}\right) + \frac{(4\zeta - 1)}{(2\zeta - 1)} \ln\left(\frac{r}{r_1}\right) - \frac{q^2}{r^2} - \frac{kq^2}{16\pi r_1^2} + \frac{c_1 r^2}{6} \right]. \quad (7.4)$$

Therefore, following Refs.[1]-[3] we can obtain the second variation of entropy functional in the form:

$$\delta^2 S = \frac{\sqrt{2}\alpha k_B}{2\hbar} \int_{r_1}^{r_2} r dr \left(\frac{dM}{dr}\right)^{-\frac{3}{2}} \frac{1}{\sqrt{g_{rr}}} \left[- \left(\frac{d(\delta M)}{dr}\right)^2 + \frac{(\delta M)^2}{g^2 r^2} \left(\frac{dM}{dr}\right) \left(1 + \frac{dM}{dr}\right) \right]. \quad (7.5)$$

Here we have conveniently considered a linear combination of $M(r)$ as $\delta M = \chi\eta$, where η vanishes at the boundaries. Plugging this consideration in eq. (7.5) and integrating by parts keeping $\delta M = 0$ at the apex of the shell we obtain:

$$\delta^2 S = -\frac{\sqrt{2}\alpha k_B}{2\hbar} \int_{r_1}^{r_2} r dr \left(\frac{dM}{dr}\right)^{-\frac{3}{2}} \frac{1}{\sqrt{g_{rr}}} \chi^2 \left[\frac{d\psi}{dr}\right]^2 < 0. \quad (7.6)$$

It is evident that the entropy function in Rastall gravity attains a maximal value for all the radial variations which vanish at the extreme points of the thin shell of the charged gravastar. Therefore, we can conclusively say that the perturbations in the ultra-relativistic fluid in the region of thin shell further decreases the entropy of the region which in turn proves that our model of charged gravastar is stable against small perturbations having fixed end points. Fundamentally, the employment of modified gravity theory does not affect the stability of a charged gravastar.

8 Conclusion

In this paper we have investigated the gravastar formalism in the framework of Rastall theory of gravity in spherically symmetric space-time and studied the effect of charge in the formulation of gravastar. We have obtained a new class of exact, singularity free and physically acceptable solutions for three layers of an isotropic, charged gravastar and we have noted the interesting features appearing in the fabrication of gravastar due to the incorporation of Rastall theory. The following prominent gravastar attributes are listed below:

- **Interior solution:** Using the EoS $p = -\rho$ and the conservation of energy-momentum tensor equation eq. (2.15) it is found that the matter density as well as pressure is constant in the interior region. In this region, we have obtained singularity free solutions as described in eqs. (3.2) and (3.3). Also, we have determined the active gravitational mass contained in the interior region of a charged gravastar.
- **EoS of the thin shell:** Eq. (3.11) shows the behaviour of matter density as well as pressure of the thin shell region and it is illustrated graphically in figure 1. The figure depicts the presence of denser fluid towards the outer edge of the shell.
- **Junction Condition:** The interior and exterior solutions are matched at the smooth hypersurface $r = R$ following Israel condition [71, 72]. Through the matching we have obtained the intrinsic surface energy tensor at given in eq. (4.1). The formulation of Lanczos equation [74]-[77] yields the surface energy density and surface pressure which is used to express the thin shell mass in terms of the total mass of gravastar as given in eq.(4.8).
- **Proper length of the thin shell:** Figure 2 shows the radial variation of proper length of the thin shell of a charged gravastar. It is evident that the proper length increases linearly with shell thickness and the increment of charge results in a decrease of length.
- **Energy of the shell:** Figure 3 shows the radial variation of the energy of the charged fluid contained within the shell. It is found that with the increase of shell thickness more charged fluid is found and when the charge increases the energy decreases which in turn illustrates the non-repulsive nature of energy in the shell.

- **Entropy:** In figure 4 we find the increasing nature of the entropy of a charged gravastar. It is also found the more charged fluid contains less entropy.
- **EoS parameter:** The formulation of the EoS parameter of the shell region shows that for large values of radius, the EoS enters dark energy regime ($\mathcal{W}(R) \rightarrow -1$). On the other had, for small value of R and suitable choice of parameters c_2 and R , the EoS parameter $\mathcal{W}(R) \rightarrow 0$, which defines a dust shell configuration.
- **Surface Redshift:** The stability of the model is ensured through the consideration of gravitational surface redshift which is found to obey the Buchdahl limit [79] for isotropic stellar configuration.
- **Entropy Maximisation:** The achievement of the condition of maximisation of entropy shows that our model of charged gravastar is stable under small radial perturbations. It also obtained that the entropy function attains a maximum value when radial perturbations vanish at the end point of the shell.

From the entire discussion above, we can assert that charge has a significant effect on the formation and basic properties of gravastar and charged gravastar model may exist in Rastall theory of gravity. One of the most noteworthy feature of this model is that for $q = 0$, we recover the results which are similar in form as obtained by Ghosh et al. [78]. We have found that the use of Rastall theory modifies the form of basic properties of a charged gravastar from those obtained in Einstein gravity. Interestingly, the insertion of $\zeta = 0$ in the theory retains Einstein's GR formalism. Finally, it may rightly be said that our model represents a stable and viable stellar configuration preserving all the characteristic features of a stable charged gravastar model within the fabric of Rastall theory.

Acknowledgments

DB is thankful to DST for providing the fellowship vide no: DST/INSPIRE/03/2022/000001.

References

- [1] P. O. Mazur, E. Mottola, *Gravitational Condensate Stars: An Alternative to Black Holes*, Report number: LA-UR-01-5067 (2001) [[gr-qc/0109035](#)].
- [2] P. O. Mazur, E. Mottola, *Gravitational vacuum condensate stars*, *Proc. Natl. Acad. Sci. USA* **101** (2004) 9545 [[gr-qc/0407075](#)].
- [3] P. O. Mazur and E. Mottola, *Gravitational Condensate Stars: An Alternative to Black Holes*, *Universe* **9** (2023) 88.
- [4] Y. B. Zel'dovich, *The Equation of State at Ultrahigh Densities and Its Relativistic Limitations*, *Sov. Phys. JETP* **14** (1962) 5, pg. 1143.
- [5] Y.B. Zel'dovich, *A hypothesis, unifying the structure and the entropy of the Universe*, *Mon. Not. R. Astron. Soc.* **160** (1972) 1P.
- [6] S. W. Hawking, *Black hole expansions?*, *Nature* **248** (1974) 30.
- [7] M. Visser and Wiltshire D. L., *Stable gravastars - an alternative to black holes?*, *Class. Quantum Gravity* **21** (2004) 1135 [[gr-qc/0310107](#)].
- [8] C. Cattoen, T. Faber, and M. Visser, *Gravastars must have anisotropic pressures*, *Class. Quantum Gravity* **22** (2005) 4189 [[gr-qc/0505137](#)].

- [9] B.M.N. Carter, *Stable gravastars with generalised exteriors*, *Class. Quantum Gravity* **22** (2005) 4551, [[gr-qc/0509087](#)].
- [10] H. Reissner, *Über die Eigengravitation des elektrischen Feldes nach der Einsteinschen Theorie*, *Ann. Phys.* **50** (1916) 106.
- [11] G. Nordström, *On the Energy of the Gravitation field in Einstein's Theory*, *Verhandl. Koninkl. Ned. Akad. Wetenschap., Afdel. Natuurk.* **26** (1918) 1201.
- [12] D. Horvat, S. Ilijic and A. Marunovic, *Electrically charged gravastar configurations*, *Class. Quantum Gravity* **26** (2009) 025003, [[gr-qc/0807.2051](#)].
- [13] A. A. Usmani, Farook Rahaman, Sk. A. Rakib, Saibal Ray, K. K. Nandi, Peter K. F. Kuhfittig, and Z. Hasan, *Charged gravastars admitting conformal motion*, *Phys. Lett. B* **701** (2011) 388, [[gr-qc/1012.5605](#)].
- [14] P. Bhar, *Higher dimensional charged gravastar admitting conformal motion*, *Astrophys. Space Sci.* **354** (2014) 457.
- [15] F. Rahaman, A. A. Usmani, S. Ray, and S. Islam, *The (2+1)-dimensional charged gravastars*, *Phys. Lett. B* **717** 1 (2012).
- [16] D. Bhattacharjee and P.K. Chattopadhyay, *Stable charged gravastar model in cylindrically symmetric space-time*, *Phys. Scr.* **98** (2023) 085013 [[gr-qc/2303.02592](#)].
- [17] Z. Yousaf, K. Bamba, M. Z. Bhatti and U. Ghafoor, *Charged gravastars in modified gravity*, *Phys. Rev. D* **100** (2019) 024062.
- [18] R. Chan and M.F.A. da Silva, *How the charge can affect the formation of gravastars*, *JCAP* **07** (2010) 29, [[gr-qc/1005.3703](#)].
- [19] A. Övgün, A. Banerjee and K. Jusufi, *Charged thin-shell gravastars in noncommutative geometry*, *Eur. Phys. J. C* **77** (2017) 566.
- [20] A.G. Riess et al., *Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant*, *Astron. J.* **116** (1998) 1009 [[astro-ph/9805201](#)].
- [21] A.G Riess et al., *New Hubble Space Telescope Discoveries of Type Ia Supernovae at $z > 1$: Narrowing Constraints on the Early Behavior of Dark Energy*, *Astrophys. J.* **659** (2007) 98 [[astro-ph/0611572](#)].
- [22] S. Perlmutter et al., *Measurements of Ω and Λ from 42 High-Redshift Supernovae*, *Astrophys. J.* **517** (1999) 565.
- [23] P. de Bernardis et al., *A flat Universe from high-resolution maps of the cosmic microwave background radiation*, *Nature (London)* **404** (2000) 955 [[astro-ph/0004404](#)].
- [24] S. Hanany et al., *MAXIMA-1: A Measurement of the Cosmic Microwave Background Anisotropy on angular scales of 10 arcminutes to 5 degrees*, *Astrophys. J.* **545** (2000) L5 [[astro-ph/0005123](#)].
- [25] P.J.E. Peebles and B. Ratra, *The cosmological constant and dark energy*, *Rev. Mod. Phys.* **75** (2003) 559 [[astro-ph/0207347](#)].
- [26] T. Padmanabhan, *Cosmological Constant - the Weight of the Vacuum*, *Phys. Rep.* **380** (2003) 235 [[hep-th/0212290](#)].
- [27] V. Sahni and A. Starobinsky, *The Case for a Positive Cosmological Λ -term*, *Int. J. Mod. Phys. D* **09** (2000) 373 [[astro-ph/9904398](#)].
- [28] T. Clifton, P. G. Ferreira, A. Padilla, and C. Skordis, *Modified gravity and cosmology*, *Phys. Rep.* **513** 1 (2012) [[astro-ph/1106.2476](#)].
- [29] R. Amanullah et al., *Spectra and Light Curves of Six Type Ia Supernovae at $0.511 < z < 1.12$ and the Union2 Compilation*, *Astrophys. J.* **716** (2010) 712 [[astro-ph/1004.1711](#)].

- [30] E. Komatsu et al., *Seven-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation*, *Astrophys. J. Suppl.* **192** (2011) 18 [[astro-ph/1001.4538](#)].
- [31] K. Tegmark et al., *Cosmological parameters from SDSS and WMAP*, *Phys. Rev. D* **69** (2004) 103501 [[astro-ph/0310723](#)].
- [32] S. Nojiri and S. D. Odintsov, *INTRODUCTION TO MODIFIED GRAVITY AND GRAVITATIONAL ALTERNATIVE FOR DARK ENERGY*, *Int. J. Geom. Methods Mod. Phys.* **04** (2007) 115 [[hep-th/0601213](#)].
- [33] S. Capozziello, *Curvature Quintessence*, *Int. J. Mod. Phys. D* **11** (2002) 483 [[gr-qc/0201033](#)].
- [34] E. Elizalde, S. Nojiri, S. D. Odintsov, L. Sebastiani, and S. Zerbini, *Nonsingular exponential gravity: A simple theory for early- and late-time accelerated expansion*, *Phys. Rev. D* **83** (2011) 086006.
- [35] K. Bamba, C.-Q. Geng, C.-C. Lee, and L.-W. Luo, *Equation of state for dark energy in $f(T)$ gravity*, *JCAP* **01** (2011) 021 [[astro-ph.CO/1011.0508](#)].
- [36] M.J.S. Houndjo, M. E. Rodrigues, N. S. Mazhari, D. Momeni, and R. Myrzakulov, *Higher-derivative $f(R, \square R, T)$ theories of gravity* *Int. J. Mod. Phys. D* **26** (2017) 1750024 [[gr-qc/1601.00196](#)].
- [37] Z. Yousaf, M. Sharif, M. Ilyas, and M. Z. Bhatti, *Energy conditions in higher derivative $f(R, \square R, T)$ gravity*, *Int. J. Geom. Methods Mod. Phys.* **15** (2018) 1850146.
- [38] M. Ilyas, Z. Yousaf, and M. Z. Bhatti, *Bounds on higher derivative $f(R, \square R, T)$ models from energy conditions*, *Mod. Phys. Lett. A* **34** (2019) 1950082 [[gr-qc/1901.04308](#)].
- [39] T. Harko, F.S.N. Lobo, S. Nojiri, and S.D. Odintsov, *Phys. Rev. D* **84** (2011) 024020 [[gr-qc/1104.2669](#)].
- [40] A. Das, S. Ghosh, B.K. Guha, S. Das, F. Rahaman, S. Ray, *Gravastars in $f(R, T)$ gravity*, *Phys. Rev. D* **95** (2017) 124011 [[gr-qc/1702.08873](#)].
- [41] P. Bhar and P. Rej, *Stable and self consistent charged gravastar model within the framework of $f(R, T)$ gravity*, *Eur. Phys. J. C*, **81** (2021) 763 [[gr-qc/2108.06989](#)].
- [42] S. Ghosh, A.D. Kanfon, A. Das, M.J.S. Houndjo, I. G. Salako and S. Ray, *Gravastars in $f(T, T)$ gravity*, *Int. J. of Mod. Phys. A* **35** (2020) 2050017 [[gr-qc/2108.06989](#)].
- [43] A. Das, S. Ghosh, D. Deb, B.K. Guha, F. Rahaman and S. Ray, *Study of gravastars under $f(T)$ gravity*, *Nuc. Phys. B* **954** (2020) 114986 [[gr-qc/2003.10842](#)].
- [44] R. Sengupta, S. Ghosh, S. Ray, B. Mishra and S. K. Tripathy, *Gravastar in the framework of braneworld gravity*, *Phys. Rev. D* **102** (2020) 024037 [[gr-qc/2004.01519](#)].
- [45] S. Banerjee, S. Ghosh, N. Paul, F. Rahaman, *Study of gravastars in Finslerian geometry*, *Euro. Phys. J Plus* **135** (2020) 185.
- [46] P. Rastall, *Generalization of the Einstein Theory*, *Phys. Rev. D* **6** (1972) 3357.
- [47] P. Rastall, *A Theory of Gravity*, *Can. J. Phys.* **54** (1976) 66.
- [48] Y. Heydarzade, H. Moradpour and F. Darabi, *Black Hole Solutions in Rastall Theory*, *Can. J. Phys.* **95** (2017) 1253 [[gr-qc/1610.03881](#)].
- [49] H.L. Parihadi, M.F.A.R. Sakti, G. Hikmawan and F.P. Zen, *Dynamics of Charged and Rotating NUT Black Holes in Rastall Gravity*, *Int. J. Mod. Phys. D* **29** (2020) 2050021 [[gr-qc/1908.09629](#)].
- [50] R. Kumar and S.G. Ghosh, *Rotating black hole in Rastall theory*, *Eur. Phys. J. C* **78** (2018) 750 [[gr-qc/1711.08256](#)].
- [51] Z. Xu and J. Wang, *Kerr-Newman-AdS Black Hole Surrounded by Perfect Fluid Matter in Rastall Gravity*, *Eur. Phys. J. C* **78** (2018) 513 [[gr-qc/1711.04542](#)].

- [52] M.S. Ma and R. Zhao, *Noncommutative geometry inspired black holes in Rastall gravity*, *Eur. Phys. J. C* **77** (2017) 629 [[hep-th/1706.08054](#)].
- [53] G. Gergess and L. Nashed, *Nonlinear Charged Black Hole Solution in Rastall Gravity*, *Universe* **8** (2022) 510.
- [54] C.E. Mota et al., *Generalized Rastall's gravity and its effects on compact objects*, *Int. J. Mod. Phys. D* **31** (2022) 2250023 [[gr-qc/2007.01968](#)].
- [55] K.D. Krori and J. Barua, *A singularity-free solution for a charged fluid sphere in general relativity*, *J. Phys. A: Math. Gen.* **8** (1975) 508.
- [56] G. Abbas and M.R. Shahzad, *Comparative analysis of Einstein gravity and Rastall gravity for the compact objects*, *Chinese J. Phys.* **63** (2020) 1.
- [57] G. Abbas and M.R. Shahzad, *Models of anisotropic compact stars in the Rastall theory of gravity*, *Astrophys Space Sci.* **364** (2019) 50.
- [58] A.M. Oliveira, H.E.S. Velten, J.C. Fabris and L. Casarini, *Neutron stars in Rastall gravity*, *Phys. Rev. D* **92** (2015) 044020 [[gr-qc/1506.00567](#)].
- [59] W.E. Hanafy, *Impact of Rastall gravity on mass, radius and sound speed of the pulsar PSR J0740+6620*, *Astrophys. J* **940** (2022) 51 [[astro-ph.HE/2209.10287](#)].
- [60] S. Hansraj, A. Banerjee and P. Channuie, *Impact of the Rastall parameter on perfect fluid spheres*, *Ann. Phys.* **400** (2019) 320 [[gr-qc/1805.00003](#)].
- [61] M. Capone, V.F. Cardone and M.L. Ruggiero, *The possibility of an accelerating cosmology in Rastall's theory*, *J. Phys. Conf. Ser.* **222** (2010) 012012.
- [62] C.E.M. Batista, M.H. Daouda, J.C. Fabris, O.F. Piattella and D.C. Rodrigues, *Rastall cosmology and the LambdaCDM model*, *Phys. Rev. D* **85** (2012) 084008 [[astro-ph/1112.4141](#)].
- [63] P. Bhar, F. Tello-Ortiz, Ángel Rincón, Y. Gomez-Leyton, *Study on anisotropic stars in the framework of Rastall gravity*, *Astrophys. Space Sci.* **365** (2020) 145.
- [64] G. Mustafa, S. Waheed, M. Zubair and T.C. Xia, *Non-commutative wormholes exhibiting conformal motion in Rastall gravity*, *Chinese J. Phys.* **65** (2020) 163.
- [65] M.S. Madsen, J.P. Mimoso, J.A. Butcher, G.F.R. Ellis, *Evolution of the density parameter in inflationary cosmology reexamined*, *Phys. Rev. D* **46** (1992) 1399.
- [66] B.J. Carr, *The Primordial black hole mass spectrum*, *Astrophys. J.* **201** (1975) 1.
- [67] I. Chakraborty, A. Pradhan, *LRS BIANCHI I MODELS WITH TIME-VARYING GRAVITATIONAL AND COSMOLOGICAL CONSTANTS*, *Gravit. Cosmol.* **7** (2001) 55.
- [68] T. Buchert, *On Average Properties of Inhomogeneous Fluids in General Relativity: Perfect Fluid Cosmologies*, *Gen. Relativ. Gravit.* **33** (2001) 1381 [[gr-qc/0102049](#)].
- [69] L.P. Linares, M. Malheiro, S. Ray, *THE IMPORTANCE OF THE RELATIVISTIC CORRECTIONS IN HYPERON STARS*, *Int. J. Mod. Phys. D* **13** (2004) 1355.
- [70] T.M. Braje, R.W. Romani, *RX J1856A3754: EVIDENCE FOR A STIFF EQUATION OF STATE*, *Astrophys. J.* **580** (2002) 1043 [[astro-ph/0208069v1](#)].
- [71] W. Israel, *Singular hypersurfaces and thin shell in general relativity*, *Nuo. Cim. B* **44** (1966) 1.
- [72] W. Israel, *erratum-ibid.* **48** (1967) 463.
- [73] G. Darrois, *Memorial des Sciences Mathematiques XXV, Fascicule XXV, Gauthier-Villars, Paris, France, Chap. V* (1927).
- [74] K. Lanczos, *Ann. Phys. (Berlin)* **379** (1924) 518.
- [75] N. Sen, *Ann. Phys. (Leipzig)* **378** (1924) 365.

- [76] G.P. Perry, R.B. Mann, *Traversable wormholes in $(2 + 1)$ dimensions*, *Gen. Relativ. Gravit.* **24** (1992) 305 [[gr-qc/0307034](#)].
- [77] P. Musgrave and K. Lake, *Junctions and thin shells in general relativity using computer algebra: I. The Darmois - Israel formalism*, *Class. Quantum Gravity* **13** (1996) 1885 [[gr-qc/9510052](#)].
- [78] S. Ghosh, S. Dey, A. Das, A. Chanda and B. C. Paul, *Study of Gravastars in Rastall Gravity*, *JCAP* **07** (2021) 004.
- [79] H. A. Buchdahl, *General Relativistic Fluid Spheres*, *Phys. Rev.* **116** (1959) 1027.
- [80] N. Straumann, *General Relativity and Relativistic Astrophysics*, Springer, Berlin (1984).
- [81] C. G. Böhmer and T. Harko, *Bounds on the basic physical parameters for anisotropic compact general relativistic objects*, *Class. Quantum Gravity* **23** (2006) 6479 [[gr-qc/0609061](#)].
- [82] D.E. Barraco and V.H. Hamity, *Maximum mass of a spherically symmetric isotropic star*, *Phys. Rev. D* **65** (2002) 124028.
- [83] B.V. Ivanov, *Maximum bounds on the surface redshift of anisotropic stars*, *Phys. Rev. D* **65** (2002) 104011 [[gr-qc/0201090v2](#)].
- [84] A. DeBenedictis et al., *Gravastar Solutions with Continuous Pressures and Equation of State*, *Class. Quantum Gravity* **23** (2006) 2303 [[gr-qc/0511097](#)].