Is equivalence principle valid for quantum gravitational field?

Baocheng Zhang1, [∗](#page-0-0)

¹ School of Mathematics and Physics, China University of Geosciences, Wuhan μ 3007 μ , China

Entanglement can be generated through the gravitational interaction between two massive bodies that are initially in a product state. This shows that the gravitational field is quantum. When the third massive body is introduced and the gravitational interaction only between the third body with either one of the former two bodies is considered, we find that no entanglement is generated between the former two bodies up to the monopole approximation, even though the considered gravitational interaction is quantum. This resembles the behavior of two accelerating two-level atoms that is usually regarded as the Unruh-DeWitt detectors. By linking the acceleration to that generated by the gravitational field, we show that the equivalence principle is still valid even though the gravitational field is quantum.

PACS numbers:

I. INTRODUCTION

Einstein equivalence principle (EEP) states that all effects of a uniform gravitational field are identical to the effects of a uniformly acceleration of a coordinate system [\[1\]](#page-3-0). Thus, one cannot distinguish a gravitational field from a uniformly accelerating frame of reference by making any measurements in a small enough region of spacetime. EEP is at the heart of the theory of General Relativity. Initially, Einstein called it the "equivalence hypothesis" when he fully comprehended the importance of equivalence between gravitation and inertia. In 1908, he elevated its name to the principle level as the "equivalence principle" after he realized that this "equivalence hypothesis" could be used as a heuristic tool to build a physically satisfactory relativistic theory of gravitation [\[2,](#page-3-1) [3\]](#page-3-2). In the past 100 years, there have been many experiments to test the validity of EEP through the table-based or space-based implementation [\[4\]](#page-3-3). No evidence of violation are found until now, but interest in such experiments still remains high due to its significance.

The EEP was constructed based on the classical theory in which the considered elements were the classical massive objects and the classical gravitational field. In recent years, however, some investigations [\[5](#page-3-4), [6](#page-3-5)] attempted to involve the quantum objects that may have internal degrees of freedom instead of the classical objects to discuss whether EEP is still valid in such a case. These studies aimed to construct a general equivalence principle that is also proper for quantum objects, by which it is expected to search for the possibility or the clues of constructing the theory of quantum gravity. Some experiments using neutron [\[7](#page-3-6), [8](#page-3-7)] or atomic interferometry [\[9,](#page-3-8) [10](#page-3-9)] has tested such ideas, and no violation is found until now. Nevertheless, it is hardly discussed whether the EEP is valid when the classical gravitational field is substituted by the quantum gravitational field. The reason is that no one knows how to describe the quantum gravitational

field properly due to the absence of the complete theory about quantum gravity. However, there exists some methods to test gravity's quantum behavior but don't require any particular theory of quantum gravity in advance, as pointed out in recent works [\[11,](#page-3-10) [12](#page-3-11)]. Based on this, here we will study the question whether the EEP is valid for the quantum gravitational field.

II. GRAVITATIONAL FIELD

According to the thought from Ref. [\[11](#page-3-10), [12\]](#page-3-11) and as stated in Ref. [\[13](#page-3-12)], if gravity follows quantum theory, it should set into a superposition of many states at once when it interacts with a mass that is also behaving in this way. A second mass could be used as a probe to pick up that quantum state. Measuring the probe's state could determine whether it has been superposed, thus proving whether gravity exhibits quantum behavior. This proposed theoretically the methods for the experimental confirmation of the assumption that the gravitational field is quantum by the result that either it is in a superposed state or it generates entanglement between two massive bodies. We will continue along this line. For simplicity, two quantum states are considered to be complete for the description of the gravitational field in this paper. At first, we provide a way that has the same evolution essentially as before [\[11,](#page-3-10) [12\]](#page-3-11), that is,

$$
\left|m_{1}\right\rangle\left|m_{1}'\right\rangle \rightarrow \alpha_{1}\left|m_{1}\right\rangle\left|m_{1}'\right\rangle+\sqrt{P}\alpha_{2}\left|m_{2}\right\rangle\left|m_{2}'\right\rangle, \tag{1}
$$

where two massive bodies separated by a distance d are distinguished by the primed and unprimed labels. As stated in Ref. [\[13\]](#page-3-12), the quantum behavior of the gravitational field can be described by analogy from the massive body that produced it. Thus, the quantum states $|m_1\rangle$ and $|m_2\rangle$ for one massive body (or for the gravitational field it produces) can be regarded as energy eigenstates like that usually used in the quantum theory, and m_1 or m_2 is the mass-like parameter related to the energy of the corresponding state by the Einstein mass-energy relation [\[1](#page-3-0)]. The primed labels have the same meaning

[∗]Electronic address: zhangbaocheng@cug.edu.cn

for another massive body. $\alpha_1 \sim \exp[-i\frac{Gm_1m_1'}{d} \frac{t}{\hbar}]$ and $\alpha_2 \sim \exp[-i\frac{Gm_2m'_2}{d}\frac{t}{\hbar}]$ represent the phases derived from the gravitational interaction. P represents the probability for the change of the state from $|m_1\rangle |m'_1\rangle$ to $|m_2\rangle |m'_2\rangle$ through the gravitational interaction between them. Evidently, the initial product state becomes entangled after the interaction, which shows that the gravitational interaction is quantum. This is a simple presentation for the earlier proposals [\[11](#page-3-10)[–14\]](#page-3-13) in which the interaction is exerted at such a product state $\left(\frac{|m_1\rangle+|m_2\rangle}{\sqrt{2}}\right)\left(\frac{|m'_1\rangle+|m'_2\rangle}{\sqrt{2}}\right)$. It is easy to recover the earlier results with the evolution presented in Eq. [\(1\)](#page-0-1) up to a normalization factor.

Now we consider a gravitational field G generated by the Earth. It can be described by two quantum states $|M_1\rangle$ and $|M_2\rangle$ simply as stated above. An atom is placed into the field G at height h to "feel" the interaction. The atom is regarded as a point-like two-level quantum system with the ground $|g\rangle$ and excited $|e\rangle$ states. According to Eq. [\(1\)](#page-0-1), the coupling between the atom and the field G is expressed as

$$
U^G |g\rangle |M_1\rangle = C_0 \left(\alpha_1^G |g\rangle |M_1\rangle + \sqrt{P_{ge}^G} \alpha_2^G |e\rangle |M_2\rangle \right),
$$

\n
$$
U^G |e\rangle |M_1\rangle = C_1 \left(\beta_1^G |e\rangle |M_1\rangle + \sqrt{P_{eg}^G} \beta_2^G |g\rangle |M_2\rangle \right),
$$
\n(2)

where $C_{0,1}$ is the state normalization factor, and U^G represents the interaction between the atom and the gravitational field. The interaction derives from the quantization of the Hamiltonian $H_{int}^G = -\frac{1}{2}h_{\mu\nu}T^{\mu\nu}$ where $T^{\mu\nu}$ is the stress-energy tensor and $h_{\mu\nu}$ is the perturbation of the metric tensor away from the flat (Minkowski) spacetime. In the earlier proposals [\[11,](#page-3-10) [12\]](#page-3-11), the quantum states of gravitational field were described as spatially localized states, i.e. the path states separated by up and down in Ref. [\[11](#page-3-10)] and that separated by left and right in Ref. [\[12\]](#page-3-11). Here the states $|M_1\rangle$ and $|M_2\rangle$ are essentially the same with the earlier usage, but it might be better to be understood roughly as the energy eigenstates of the gravitational field. For example, the atom absorbs the energy from the field G , which would lead to the change of the energy or the state of the field G. Because the mass of the Earth is much larger than the atom, the quantum states corresponding to its gravitational field are dense in this sense that $\frac{|M_2 - M_1|}{M_1} \ll \frac{|m_e - m_g|}{m_g}$. Moreover, the states $|M_1\rangle$ and $|M_2\rangle$ can be orthogonal only if the quantum gravitational field has a complete set of state vectors. As stated above, we consider here that the number of state vectors for the set is two for simplify. The parameters α_1^G , α_2^G , β_1^G , and β_2^G can be obtained according to the Eq. [\(1\)](#page-0-1). P_{ge}^{G} (P_{eg}^{G}) represents the transition probability from ground (exited) to exited (ground) state for the atom interacting with the field G.

When two atoms A and B with the initial product state $|q\rangle |e\rangle$ are placed into the gravitational field $|M_1\rangle$ at the same height but different sites, through the interaction

with the gravitational field G , the state becomes

$$
\begin{split} \left| \Psi_{f}^{G} \right\rangle &= \left(U^{G} \left| g \right\rangle \left| M_{1} \right\rangle \right) \left(U^{G} \left| e \right\rangle \left| M_{1} \right\rangle \right) \\ &= C_{0} C_{1} \left[\alpha_{1}^{G} \beta_{1}^{G} \left| g \right\rangle \left| e \right\rangle \left| M_{1} \right\rangle \left| M_{1} \right\rangle \\ &+ \sqrt{P_{eg}^{G}} \alpha_{1}^{G} \beta_{2}^{G} \left| g \right\rangle \left| g \right\rangle \left| M_{1} \right\rangle \left| M_{2} \right\rangle \\ &+ \sqrt{P_{ge}^{G}} \alpha_{2}^{G} \beta_{1}^{G} \left| e \right\rangle \left| e \right\rangle \left| M_{1} \right\rangle \left| M_{2} \right\rangle \\ &+ \sqrt{P_{ge}^{G}} \sqrt{P_{eg}^{G}} \alpha_{2}^{G} \beta_{2}^{G} \left| e \right\rangle \left| g \right\rangle \left| M_{1} \right\rangle \left| M_{2} \right\rangle \end{split} \tag{3}
$$

Tracing out the quantum states of the gravitational field, we obtain the final state for the two atoms as

$$
\rho_f^G = \rho_A^G \otimes \rho_B^G,\tag{4}
$$

where $\rho_A = |C_0|^2 (|\alpha_1^G|)$ $\left| \right| 2 \left| g \right\rangle \left\langle g \right| + P_{ge}^G \left| \alpha_2^G \right|$ $2 |e\rangle \langle e|$ and $\rho_B = |C_1|^2 \left(\left| \beta_1^G \right| \right)$ $\langle e| + P_{eg}^G |\beta_2^G|$ $2|g\rangle\langle g|$. It shows that no entanglement is created when the two atoms are coupled to the same gravitational field. This is obtained by ignoring the gravitational interaction between two atoms, since the interaction is much smaller than that between the atom and the Earth. But in the earlier proposals [\[11](#page-3-10), [12\]](#page-3-11), the generation of entanglement is due to the gravitational interaction between two atoms. This is embodied here in the generation of entanglement between the atom and the Earth, which guarantees that the gravitational field G is quantum. Moreover, when the two atoms are placed in the different height, the similar result can be obtained.

III. ACCELERATION

In order to investigate the EEP, we turn to the effects in an accelerating frame of reference. It is widely accepted by now that an observer with the uniform acceleration a in the Minkowski vacuum would feel a thermal bath of particles at the temperature $T_U = \hbar a/2\pi c k_B$. This is just the well-known Unruh effect [\[15\]](#page-3-14). It has been digested and extended to many different situations (see the review [\[16\]](#page-3-15) and references therein). In one famous application to the Unruh-DeWitt (UDW) detector [\[17\]](#page-3-16), it is found that a quantum system consisting of a detector uniformly accelerating in Minkowski vacuum can sense the thermal emission and, thus, cause the coupling of the quantum system with the thermal field. In what follow, we will make the discussion using the UDW model.

Starting with the consideration that constitutes a scalar field ϕ interacting with a point-like two-level quantum system, or an atom as used in the discussion above. The interaction Hamiltonian for this $(1 + 1)$ -dimension model can be modeled by the interaction Hamiltonian, $H_I = \lambda \mu(\tau) \phi(x(\tau))$, with λ the coupling strength. τ is the atom's proper time along its trajectory $x(\tau)$, $\mu(\tau)$ is the atom's monopole momentum, and $\phi(x(\tau))$ is the scalar field related to the vacuum. For an atom accelerating in the Minkowski vacuum, the evolution of the total quantum state is determined perturbatively by the unitary operator which up to first order is given by, $U = I + U^{(1)} + O(\lambda^2) = I - i \int d\tau H(\tau) + O(\lambda^2)$. Within the first-order approximation and in the interaction picture, this evolution is described by [\[18](#page-3-17), [19](#page-3-18)]

$$
U|g\rangle|0\rangle = D_0 (|g\rangle|0\rangle + \eta_0|e\rangle|1_k\rangle),
$$

$$
U|e\rangle|0\rangle = D_1 (|e\rangle|0\rangle + \eta_1|g\rangle|1_k\rangle),
$$
 (5)

where k denotes the mode of the $(1 + 1)$ -dimension scalar field with (bosonic) annihilation (creation) operator a_k (a_k^{\dagger}) , $a_k |0\rangle = 0$ and $a_k^{\dagger} |0\rangle = |1_k\rangle$. $D_{0,1}$ is the state normalization factor. η_0 and η_1 are related to the excitation and deexcitation probability of the atom, i.e. $P_{ge} = \sum_{k} | \langle e, 1_k | U^{(1)} | g, 0 \rangle |$ 2 = $|\eta_{0}|^2$ and P_{eg} = $\sum_{k} |\langle g, 1_k | U^{(1)} | e, 0 \rangle|$ $^2 = |\eta_1|^2$ (the relative phases are absorbed into the parameters η_0 and η_1 whose concrete form refers to Ref. [\[16,](#page-3-15) [18\]](#page-3-17)). $t(\tau) = \frac{c}{a} \sinh(\frac{a\tau}{c})$ and $x(\tau) = \frac{c^2}{a} \left(\cosh(\frac{a\tau}{c}) - 1 \right)$ is the trajectory of the accelerating atom with acceleration a. Approximately, $t \approx \tau$ and $x \simeq \frac{1}{2} a \tau^2$ indicates the trajectory of uniformly accelerated rectilinear motion.

When two atoms A and B with the initial product state $|q\rangle |e\rangle$ are accelerated with the same acceleration, and the initial product state then evolves into

$$
\begin{aligned} \left| \Psi_f \right\rangle &= D_0 D_1(\left| g \right\rangle \left| e \right\rangle \left| 0 \right\rangle \left| 0 \right\rangle + \eta_1 \left| g \right\rangle \left| g \right\rangle \left| 0 \right\rangle \left| 1_k \right\rangle \\ &+ \eta_0 \left| e \right\rangle \left| e \right\rangle \left| 1_k \right\rangle \left| 0 \right\rangle + \eta_0 \eta_1 \left| e \right\rangle \left| g \right\rangle \left| 1_k \right\rangle \left| 1_k \right\rangle. \end{aligned} \tag{6}
$$

Again, the reduced two-atom density matrix is obtained by tracing out the scalar field modes

$$
\rho_f = \rho_A \otimes \rho_B,\tag{7}
$$

It shows that no entanglement is created. This is the same as the result [\(4\)](#page-1-0) obtained from the quantum gravitational field. Moreover, when the two atoms assume different accelerations, the final quantum state is found to take a similar form to that Eq. [\(7\)](#page-2-0) for two atoms with the same acceleration.

IV. EQUIVALENCE BETWEEN GRAVITATION AND ACCELERATION

Then, let's see how to relate the gravitational field with the acceleration. As well-known, the accelerating frame of reference can be expressed with the Rindler metric $ds^2 = \rho^2 d\xi^2 - d\rho^2$ [\[16](#page-3-15)]. It might be easier to understand this by using the associated coordinates $z = \rho - \frac{c^2}{a}$ $\frac{a}{a}$ and $\tau = \frac{\rho}{c}\xi$, in terms of which the Rindler metric becomes [\[20\]](#page-3-19)

$$
ds^{2} = -\left(1 + \frac{az}{c^{2}}\right)^{2}c^{2}d\tau^{2} + dz^{2},
$$
 (8)

where τ represents the proper time, and $a = \frac{c^2}{a}$ $\frac{e^2}{\rho}$ (at the point $z = 0$) represents the acceleration. This coordinates can also be interpreted with the static gravitational field where $g = -a$ is the acceleration of free fall at that point with respect to the chosen body at rest. In the Newtonian approximation, when the potential $\phi \ll c^2$, one has

$$
ds^{2} = -\left(1 + \frac{2\phi}{c^{2}}\right)^{2} c^{2} d\tau^{2} + \left(1 - \frac{2\phi}{c^{2}}\right) dz^{2}
$$

$$
\simeq -\left(1 + \frac{\phi}{c^{2}}\right)^{2} c^{2} d\tau^{2} + dz^{2}, \tag{9}
$$

up to a conformal factor in the last step. From this, it can be seen that the metric [\(8\)](#page-2-1) corresponds to the case of a homogeneous gravitational field (in the paper, we consider the atoms rested in the specific points of the gravitational field, and thus for the small enough regime around the points, the field can be regarded as homogeneous) in the Newtonian approximation with the potential $\phi = -gz = az$ which provides the relation between the description of accelerated atoms and the atoms rested in the gravitational field.

For the accelerated atom, the transition ratio can be calculated as $\frac{P_{ge}}{P_{eg}} \simeq \exp[-\frac{\Delta E}{k_B T_U}] = \exp[-\frac{2\pi c \Delta E}{a}]$ for a long enough time, where ΔE is the energy gap between the ground $|q\rangle$ and excited $|e\rangle$ states of the atom. It indicates the thermal response of a particle detector evaluated by the detailed-balance condition obeyed by KMS states [\[21\]](#page-3-20). Similarly, the transition ratio caused by the gravitational field can be calculated as $\frac{P_{ge}^G}{P_{eq}^G} = \exp[\frac{2\pi c\Delta E}{g}],$ which derives from the consideration [\[20\]](#page-3-19) that a detector at rest in a static homogeneous gravitational field at point at which the acceleration of free fall is $q = -a$ behaves in the case of interaction with the quantized field in the Hartle-Hawking vacuum state [\[22\]](#page-3-21). The Hartle-Hawking vacuum state is related to the thermal equilibrium state in the free falling frame of reference in the static gravitational field, while the Boulware vacuum [\[23](#page-3-22)] is seen for observers rested in the static gravitational field. This confirms that the EEP is still valid when the gravitational field is regarded as quantum, although the calculation is made in the monopole approximation both for the gravity and the acceleration. When the higher expansion is considered, entanglement could be generated according to the process stated above, which is similar to the mechanism of entanglement harvesting [\[24\]](#page-3-23) in the acceleration.

Notably, the connection between acceleration and gravity relies on the transition probability of a single atom. However, our validation of the EEP is grounded in the entanglement of two atoms. This extension holds true as, within the considered approximation, verifying the EEP necessitates that the two atoms experience identical gravitational acceleration and are subjected to the same acceleration in a flat spacetime background. The entanglement under discussion is intricately associated with the transition probability of each atom, allowing the confirmation of the EEP through the alteration of entanglement in distinct scenarios.

V. CONCLUSION

In conclusion, the EEP is discussed under the background of quantum gravitational field. The quantum property of the gravitational field is confirmed by generating entanglement between the source (Earth) and the objects (atoms). When two atoms are placed in the gravitational field and interact with the field respectively, we show that no entanglement is generated between the two atoms. This conclusion is obtained by taking the interaction in the monopole approximation and ignoring the interaction between two atoms since the mass of the atom is much smaller than that of the Earth. On the other hand, when two atoms are accelerated in the Minkowski vacuum, no entanglement is generated either under the interaction of the first-order approximation. Thus, by linking the acceleration to the static homogeneous (at

least locally) gravitational field, we show that the EEP is still valid for the quantum gravitational field up to the monopole approximation. More general confirmation for the EEP might require more knowledge about the vacuum and the high-order interaction for the strong gravitational field [\[25](#page-3-24)], and more refined quantum criteria for the gravitational field [\[26](#page-3-25)].

VI. ACKNOWLEDGEMENTS

This work is supported by National Natural Science Foundation of China (NSFC) with Grant No. 12375057, and the Fundamental Research Funds for the Central Universities, China University of Geosciences (Wuhan) with No. G1323523064.

- [1] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravita*tion (W. H. Freeman and Company, San Francisco, USA, 1973).
- [2] A. Einstein, Jahrbuch Radiaoktiv. 4, 411 (1908).
- [3] A. Einstein, Ann. Phys. Lpz., 35, 898 (1911).
- [4] C. M. Will, Living Rev. Relativity, 9, 3 (2006).
- $[5]$ M. Zych and C. Brukner, Nature Phys 14, 1027–1031 (2018).
- [6] C. Anastopoulos and B. L. Hu, Class. Quantum Grav. 35, 035011 (2018).
- [7] R. Colella, A. Overhauser, and S. Werner, Phys. Rev. Lett. 34, 1472–1474 (1975).
- [8] U. Bonse and T. Wroblewski, Phys. Rev. Lett. 51, 1401 (1983).
- [9] S. Fray, C. A. Diez, T. W. Hänsch, and M. Weitz, Phys. Rev. Lett. 93, 240404 (2004).
- [10] S. Herrmann, H. Dittus, C. Lämerzahl, et al., Classical and Quantum Gravity 29, 184003 (2012).
- [11] C. Marletto and V. Vedral, Phys. Rev. Lett. 119, 240402 (2017).
- [12] Sougato Bose, Anupam Mazumdar, Gavin W. Morley, et al., Phys. Rev. Lett. 119, 240401 (2017).
- [13] C. Marletto and V. Vedral, Nature 547, 156 (2017).
- [14] C. Marletto, V. Vedral, and D. Deutsch, New J. Phys.

20, 083011 (2018).

- [15] W. G. Unruh, Phys. Rev. D 14, 870 (1976). [16] L. C. B. Crispino, A. Higuchi, and G. E. A. Matsas, Rev.
- Mod. Phys. 80, 787 (2008). [17] B. S. DeWitt, S. Hawking, and W. Israel, *General Rela*tivity: An Einstein Centenary Survey (Cambridge University Press Cambridge, England, 1979).
- [18] W. Brenna, R. B. Mann, and E. Martín-Martínez, Phys. Lett. B 757, 307 (2016).
- [19] T. Li, B. Zhang and L. You, Phys. Rev. D 97, 045005 (2018).
- [20] V. L. Ginzburg and V. P. Frolov, Sov. Phys. Usp. 30 1073 (1987).
- [21] S. Takagi, Prog. Theor. Phys. Suppl. 88, 1 (1986).
- [22] J. B. Hartle and S. W. Hawking, Phys. Rev. D 13, 2188 (1976).
- [23] D. G. Boulware, Phys. Rev. D 11, 1404 (1975).
- [24] G. Salton, R. B. Mann, and N. C. Menicucci, New J. Phys. 17, 035001 (2015).
- [25] D. Singleton and S. Wilburn, Phys. Rev. Lett. 107, 081102 (2011).
- [26] M. J W Hall and M. Reginatto, J. Phys. A: Math. Theor. 51, 085303 (2018).