A VERTEX-SKIPPING PROPERTY FOR ALMOST-MINIMIZERS OF THE RELATIVE PERIMETER IN CONVEX SETS

GIAN PAOLO LEONARDI AND GIACOMO VIANELLO

ABSTRACT. Given a convex domain $\Omega \subset \mathbb{R}^3$ and an almost-minimizer E of the relative perimeter in Ω , we prove that the closure of $\partial E \cap \Omega$ does not contain vertices of Ω .

1. INTRODUCTION

Given a convex open set $\Omega \subset \mathbb{R}^n$, $n \geq 2$, we aim to study the boundary behavior of local almostminimizers of the relative perimeter in Ω near certain singular points of $\partial\Omega$. More specifically, we take a measurable set $E \subset \Omega$ and, for any open set $A \subset \mathbb{R}^n$, we define the relative perimeter of E in Ω restricted to A as

$$P_{\Omega}(E;A) := P(E;\Omega \cap A),$$

with the short form $P_{\Omega}(E) := P(E; \Omega)$ when $A \supset \Omega$, and with P(E; B) denoting the standard perimeter of E in B à la De Giorgi. We then say that E is a local almost-minimizer of P_{Ω} if, for any $x \in \overline{\Omega}$ there exists $r_x > 0$ such that, for any $0 < r < r_x$ and any measurable subset F of Ω with $F\Delta E \subset B_r(x)$, one has

$$P_{\Omega}(E; B_r(x)) \le P_{\Omega}(F; B_r(x)) + |F\Delta E|^{\frac{n-1}{n}} \psi_{\Omega}(E; x, r), \qquad (1.1)$$

for a suitable function $\psi_{\Omega}(E; x, r)$ such that $\lim_{r \to 0^+} \psi_{\Omega}(E; x, r) = 0$. In particular, E is a perimeter minimizer in $B_r(x)$ if and only if $\psi_{\Omega}(E; x, r) = 0$.

Similar notions of almost-minimality are well-known in the literature, and the most relevant ones satisfy condition (1.1). For instance, the lambda-minimality, which characterizes solutions to isoperimetric and prescribed mean curvature problems, implies (1.1) by choosing $\psi_{\Omega}(E; x, r) = \Lambda r \omega_n^{1/n}$. It is worth recalling that De Giorgi's interior regularity for perimeter minimizers is wellknown to hold up to $C^{1,\alpha}$ -regularity for almost-minimizers in the sense of Tamanini [30, 31], which in our case corresponds to (1.1) combined with the extra summability of $r^{-1}\psi(E; x, r)$ on some interval $(0, r_0)$, for any $x \in \Omega$ and with $r_0 > 0$ possibly depending on x.

²⁰²⁰ Mathematics Subject Classification. Primary: 49Q05. Secondary: 49Q10.

Key words and phrases. perimeter, almost-minimizers, stability, capillarity.

G.P.Leonardi has been partially supported by: PRIN 2017TEXA3H "Gradient flows, Optimal Transport and Metric Measure Structures"; PRIN 2022PJ9EFL "Geometric Measure Theory: Structure of Singular Measures, Regularity Theory and Applications in the Calculus of Variations" (financed by European Union - Next Generation EU, Mission 4, Component 2 - CUP:E53D23005860006); Grant PID2020-118180GB-I00 "Geometric Variational Problems". Giacomo Vianello has been supported by GNAMPA (INdAM) Project 2023: "Esistenza e proprietà fini di forme ottime". The authors wish to thank the anonymous referees for very helpful comments on the first version of the paper, and in particular for pointing out a remarkable consequence of our main result in combination with [10, Theorem 1.1].

Further regularity results are known in the context of integral currents and varifolds, where instead of almost-minimality hypotheses one considers summability assumptions on the generalized mean curvature of the (associated) varifold [1,2]. In particular, the problem of boundary regularity for integral k-varifolds with free boundary and mean curvature in L^p with p > k, has been first tackled in [15].

To state our main result, we need to introduce the notion of vertex of Ω . Given $x_0 \in \partial \Omega$ we define the (open) tangent cone to Ω at x_0 as

$$T_{x_0}\Omega = \lim_{t \to +\infty} t(\Omega - x_0) = \bigcup_{t > 0} t(\Omega - x_0).$$

We say that x_0 is a vertex of Ω if $T_{x_0}\Omega$ does not contain lines. Notice that this is equivalent to require that, up to isometries, $T_{x_0}\Omega$ cannot be written as $\mathbb{R} \times C$ for some convex cone C in \mathbb{R}^{n-1} . That said, our main result is the following

Theorem 1.1 (Vertex-skipping). Let $\Omega \subset \mathbb{R}^3$ be a open, convex set. Let E be a local almostminimizer of the relative perimeter in Ω . Then $\overline{\partial E \cap \Omega}$ does not contain vertices of Ω .

Theorem 1.1 provides a partial answer to the broader question whether the internal boundary of an almost-minimizer of the relative perimeter in Ω can reach singular points of $\partial\Omega$. The restriction to dimension n = 3 is not purely technical. Indeed, in the forthcoming paper [21], we show that in dimension n = 4 the intersection of the half-space

$$E = \{ (x_1, \dots, x_4) \in \mathbb{R}^4 : x_1 < 0 \}$$

with the cone

$$C_{\lambda} = \{(x_1, \dots, x_4) \in \mathbb{R}^4 : x_4 > \lambda \sqrt{x_1^2 + x_2^2 + x_3^2}\}$$

is strictly stable with respect to a suitable class of compact variations up to the boundary of C_{λ} , as soon as $\lambda > 0$ is small enough. Even though the global minimality of E for the relative perimeter in C_{λ} for small λ (with respect to *all* compact variations) remains an open question, the strict stability of E in C_{λ} supports the conjecture that Theorem 1.1 is false in dimension $n \geq 4$.

In the more general context of capillarity, it is well-known that when Ω is of class $C^{1,1}$, a capillary surface interface in Ω meets $\partial\Omega$ according to Young's law [32], i.e., forming a contact angle $\beta = \arccos \gamma$. Here $\gamma \in [-1, 1]$ is the wetting coefficient appearing in the capillary energy (without bulk terms)

$$P_{\Omega}(E) + \gamma P(E; \partial \Omega),$$

so that one obtains the relative perimeter in the special case $\gamma = 0$. The rigorous derivation of Young's law is based on a boundary regularity result for local minimizers of the capillary energy, which in turn requires $\partial\Omega$ to be sufficiently smooth [9]. However, the study of capillarity in nonsmooth containers is not only as relevant in applications, as it is Young's law for smooth ones (see [12]), but also one of the first historical testimonies of scientific research around this natural phenomenon. Indeed, one of the first studies on the behavior of capillary surfaces in wedges was conducted by Taylor about a century before Young's and Laplace's works in the context of smooth containers (see [12]). The interest for the capillarity phenomenon in containers with wedge- or corner-type singularities emerges in various works, like in Concus-Finn [7], Lancaster [18], Chen-Finn-Miersemann [5], Lancaster-Siegel [17], Tamanini [29]. We also mention that, in the same spirit, the first author in collaboration with G. Saracco studied in [20] the boundary behavior of non-parametric solutions of the prescribed mean curvature equation on a weakly-regular domain A, with application to the capillarity for perfectly wetting fluids ($\gamma = -1$) that partially fill the cylinder $A \times \mathbb{R}$.

We now consider the case of the relative perimeter in convex domains, and recall some relevant results in this context. In [3], Bokowski and Sperner obtained the characterization of minimizers for the relative isoperimetric problem in Euclidean balls, as well as a relative isoperimetric inequality in a convex domain with a non-sharp constant explicitly written in terms of a Bonnesen-type asymmetry index. Later on, Lions and Pacella [23] showed the optimal isoperimetric inequality

$$P(E;\mathcal{C}) \ge P(B_E;\mathcal{C})$$

in a convex cone \mathcal{C} with a vertex at the origin, for all measurable $E \subset \mathcal{C}$ with $|E| < +\infty$, where B_E denotes the Euclidean ball centered at the origin and such that $|B_E \cap \mathcal{C}| = |E|$. The same inequality has been later strengthened by Figalli and Indrei [11] who proved its sharp quantitative form using optimal transport. We also mention the relative isoperimetric problem in Ω such that $\mathbb{R}^n \setminus \Omega$ is convex. This problem was first tackled by Choe, Ghomi and Ritoré [6], who proved the optimal inequality

$$P_{\Omega}(E) \ge \frac{n\omega_n^{\frac{1}{n}}}{2} |E \cap \Omega|^{(n-1)/n}$$

plus the characterization of the equality case, under the assumption $\partial \Omega \in C^2$. More recently, this result has been extended by Fusco and Morini [13] to the non-smooth case. Concerning the properties of minimizers, Sternberg and Zumbrun proved in [28] that the internal boundary of any local minimizer of the relative perimeter in a convex domain with $C^{2,\alpha}$ boundary is either connected, or a union of parallel planes meeting $\partial \Omega$ orthogonally. Such a property follows quite directly from the concavity of the isoperimetric profile, which is another key result obtained in the very same paper. See also [16], [25], [27], [19] for extensions to general convex sets.

The study of critical points of the relative perimeter is closely connected with the classification of free-boundary minimal surfaces. The internal boundary of a local minimizer of the relative perimeter without volume constraints is indeed a free-boundary minimal surface. The study of free-boundary minimal surfaces, initiated by seminal works of Courant [8] and Lewy [22], has been carried out by various authors in the past (see e.g. Nitsche's book [26] for an updated list of classical references, up to 1975) and still represents a very active research area (see [4] for a comprehensive overview of more recent advances in the field). However, the majority of classification results is usually obtained on specific domains - or ambient manifolds - with smooth boundary, while the case of non-smooth domains is typically neglected due to the technical difficulties in implementing variation arguments in presence of boundary singularities. It is, however, worth mentioning the very recent, Allard-type ε -regularity result for free-boundary minimal surfaces, recently proved by Edelen and Li in [10] for locally-almost-polyhedral domains. In this regard, Theorem 1.1 can be easily extended to free-boundary, locally (almost) areaminimizing 2-currents in 3-dimensional convex domains. Moreover, as observed by one of the anonymous referees of the paper, if we combine our result with [10, Theorem 1.1], we can prove that the singular set of any such current in a polyhedral domain is empty, which improves [10, Theorem 1.2].

2. Preliminaries

2.1. Notation. For this work, unless otherwise specified, Ω denotes a open, convex subset of \mathbb{R}^n . Given $A \subset \mathbb{R}^n$ open and $F \subset \Omega$ measurable, we define the relative perimeter of F in A as

$$P_{\Omega}(F;A) := P(F;A \cap \Omega),$$

where

$$P(F;A) = \sup\left\{\int_{A\cap F} \operatorname{div} g(x) \, dx : g \in C_c^1(A;\mathbb{R}^n), \ |g| \le 1\right\}$$

It is convenient to introduce the minimality gap of E in A (relative to Ω) as

$$\Psi_{\Omega}(E;A) = P_{\Omega}(E;A) - \inf \left\{ P_{\Omega}(F;A) : F\Delta E \subset A \cap \overline{\Omega} \right\}.$$

We observe that, if E is a local almost-minimizer of P_{Ω} then, for all $x \in \overline{\Omega}$ and for all $0 < r < r_x$,

$$\Psi_{\Omega}(E; B_r(x)) \le \omega_n^{\frac{n-1}{n}} r^{n-1} \psi_{\Omega}(E; x, r) \,. \tag{2.1}$$

We denote the characteristic function of F by $\mathbf{1}_F$. We denote by Df the distributional gradient of $f \in BV(A)$, and identify it as usual with a vector-valued Radon measure on A. Given a vectorvalued Radon measure $\mu = (\mu_1, ..., \mu_p) : \mathcal{B}(\mathbb{R}^n) \to \mathbb{R}^p$, we denote by $|\mu|$ its total variation. Let $u = (u_1, ..., u_p) : \mathbb{R}^n \to \mathbb{R}^p, u \in L^1_{loc}(\mu)$, then we denote by $u \cdot \mu$ the Radon measure defined by

$$(u \cdot \mu)E := \int_E u \cdot d\mu = \sum_{q=1}^p \int_E u_q d\mu_q$$

Let $S \subset \mathbb{R}^n$ and t > 0, we define $S_t := t^{-1}S$. Consequently, the tangent cone to a convex set Ω at $x_0 \in \overline{\Omega}$ is denoted by $T_{x_0}\Omega$ and defined as the (set-theoretic, loc-Hausdorff, L^1_{loc}) limit of $(\Omega - x_0)_t$ as $t \to 0$. We shall often assume that $x_0 = 0$ and abbreviate $\Omega_0 = T_0\Omega$.

Finally, we say that a convex set $C \subset \mathbb{R}^n$ is a wedge provided there exist $x_0 \in \mathbb{R}^n$ and two linearly independent unit-normal vectors $w_1, w_2 \in \partial B_1$ such that

$$C = \{x \in \mathbb{R}^n : (x - x_0) \cdot w_1 \le 0, (x - x_0) \cdot w_2 \le 0\}.$$

We note that a wedge C is a cone with respect to all points of the n-2 dimensional affine subspace

$$\sigma_C = \{x_0 + v : \langle v, w_i \rangle = 0 \ \forall i = 1, 2\},\$$

that we call the spine of C.

2.2. Boundary density estimates. Here we establish perimeter and volume density estimates for almost-minimizers at a boundary point for Ω , and we provide the full proof for the reader's convenience. We highlight that, in what follows, Ω is required to be just Lipschitz (i.e., the convexity of Ω is not needed).

Lemma 2.1. Let $\Omega \subset \mathbb{R}^n$ be an open set with Lipschitz boundary. Let E be an almost-minimizer in Ω . Up to a translation, we assume that $0 \in \partial \Omega$ and that $P_{\Omega}(E; B_r) > 0$ for all r > 0. Then, there exist a constant $C \geq 1$ and a radius $\overline{r} > 0$, both depending on 0, such that

$$C^{-1}r^{n-1} \le P_{\Omega}(E; B_r) \le Cr^{n-1}$$
(2.2)

$$\min\left(|E \cap B_r \cap \Omega|, \ |(B_r \cap \Omega) \setminus E|\right) \ge C^{-1}r^n, \tag{2.3}$$

for all $0 < r < \overline{r}$.

Proof. We start proving (2.3). Given $0 < r < r_0$ we set

$$m(r) := |B_r \cap \Omega \cap E|, \quad \mu(r) := |B_r \cap \Omega \setminus E|.$$

Both m and μ are non-decreasing, thus differentiable for almost all r > 0. By [24, Example 13.3], for almost all r > 0, we have

$$m'(r) = \mathcal{H}^{n-1}(E \cap \partial B_r \cap \Omega), \quad \mu'(r) = \mathcal{H}^{n-1}(\partial B_r \cap \Omega \setminus E).$$

Since the one-parameter family of rescaled domains $D_r := r^{-1}(\Omega \cap B_r)$, $0 < r \leq 1$ is precompact in the class of Lipschitz and connected domains with respect to the L^1 -convergence, there exists a constant $\overline{C} > 0$ such that the following, relative isoperimetric inequality holds:

$$P_{\Omega}(E; B_r) \ge \overline{C} \min\{m(r), \mu(r)\}^{\frac{n-1}{n}}, \qquad (2.4)$$

for all 0 < r < 1. Set 0 < t < r and define the competitor

$$F_t = \begin{cases} E \cup B_t \cap \Omega & \text{if } m(t) > \mu(t), \\ E \setminus B_t \cap \Omega & \text{otherwise.} \end{cases}$$

We note that in the first case $F_t \Delta E = B_t \cap \Omega \setminus E$, while in the second case $F_t \Delta E = B_t \cap \Omega \cap E$. In any case, we have $F_t \Delta E \subset B_r \cap \Omega$. Thus, by the almost-minimality of E in Ω , and for almost all 0 < t < r, we infer that either

$$P_{\Omega}(E; B_r) \leq P_{\Omega}(F_t; B_r) + \mu(t)^{\frac{n-1}{n}} \psi(r)$$

$$\leq P_{\Omega}(E; B_r \setminus \overline{B_t}) + \mathcal{H}^{n-1}(\partial B_t \cap \Omega \setminus E) + \mu(r)^{\frac{n-1}{n}} \psi(r) ,$$
(2.5)

or

$$P_{\Omega}(E; B_r) \leq P_{\Omega}(F_t; B_r) + m(t)^{\frac{n-1}{n}} \psi(r)$$

$$\leq P_{\Omega}(E; B_r \setminus \overline{B_t}) + \mathcal{H}^{n-1}(\partial B_t \cap \Omega \cap E) + m(r)^{\frac{n-1}{n}} \psi(r).$$
(2.6)

where $\psi(r) := \psi_{\Omega}(E; 0, r)$. Taking the limit as $t \nearrow r$ in (2.5) and (2.6), and using (2.4), we deduce that, if $m(r) > \mu(r)$, then for almost all $0 < r < r_0$ we have

$$\mu'(r) + \mu(r)^{\frac{n-1}{n}}\psi(r) \ge \overline{C}\mu(r)^{\frac{n-1}{n}},$$

while otherwise we have

$$m'(r) + m(r)^{\frac{n-1}{n}}\psi(r) \ge \overline{C}m(r)^{\frac{n-1}{n}}$$

Therefore, calling $s(r) := \min\{m(r), \mu(r)\}$ and owing to the infinitesimality of $\psi(r)$ as $r \to 0$, we obtain

$$\frac{s'(r)}{s(r)^{\frac{n-1}{n}}} \ge C,$$

for $0 < r < \overline{r}$, for some $C, \overline{r} > 0$. Integrating this inequality on the interval (0, r) we obtain (2.3). Then, the first inequality in (2.2) follows from (2.3) and (2.4). Finally, the second inequality in (2.2) follows from the observation that, taking the limit as $t \nearrow r$ in (2.5) and possibly redefining \overline{r} and C, we have

$$P_{\Omega}(E; B_r) \leq \mathcal{H}(\partial B_r \cap \Omega \setminus E) + \mu(r)^{\frac{n-1}{n}} \psi(r) \leq Cr^{n-1},$$

for every $0 < r < \overline{r}$.

2.3. Blow-up limits. We now show that a sequence of dilations of an almost-minimizer E in Ω converge, up to subsequences, to a minimizer of the relative perimeter in the tangent cone Ω_0 .

Lemma 2.2. Let E be a almost-minimizer in Ω , and assume $0 \in \partial \Omega$ and $P_{\Omega}(E; B_r) > 0$ for all r > 0. Then, there exist a sequence $t_j \searrow 0$ and a measurable set $E_0 \subset \Omega_0$ such that $E_{t_j} \to E_0$ in L^1_{loc} and E_0 is a perimeter-minimizer in Ω_0 , namely

$$\Psi_{\Omega_0}(E_0; B_R) = 0 \qquad for \ any \ R > 0.$$

Proof. First, we fix R > 0 and prove that there exist $t_0, C > 0$ such that

$$P(E_t; B_R) \le CR^{n-1} \quad \forall 0 < t < t_0.$$
 (2.7)

In what follows, for more simplicity, we will write C to denote a constant that might change from one line to another. To prove (2.7), we note that

$$P(\Omega_t; B_R) = t^{1-n} P(\Omega; B_{Rt}) \le CR^{n-1}$$
(2.8)

since $\partial \Omega$ is Lipschitz. Then owing to (2.2) and (2.8), and assuming $t < t_0 := \min(1, \bar{r}/R)$, we obtain

$$P(E_t; B_R) \le P_{\Omega_t}(E_t; B_R) + P(\Omega_t; B_R) = t^{1-n} (P_{\Omega}(E; B_{tR}) + P(\Omega; B_{Rt})) \le C R^{n-1},$$

which proves (2.7). Consequently, we obtain the global perimeter bound

ŀ

$$P(E_t) \le P(E_t; B_R) + P(\Omega_t; B_R) \le CR^{n-1},$$

hence any blow-up sequence E_{t_j} admits a (not relabeled) subsequence converging in $L^1(B_R)$ to a limit set E_0 . By a standard diagonal argument, one can prove the existence of a subsequence and a limit set, still denoted respectively as E_{t_j} and E_0 , such that $E_{t_j} \to E_0$ in $L^1_{loc}(\mathbb{R}^n)$. Since the corresponding sequence of rescaled domains Ω_{t_j} converges to the tangent cone Ω_0 in $L^1_{loc}(\mathbb{R}^n)$, we infer that $E_0 \subset \Omega_0$.

Now we have to show that $\Psi_{\Omega_0}(E_0; B_R) = 0$, for all R > 0. To do so, we claim that

$$\Psi_{\Omega_0}(E_0; B_R) \le \liminf_{j \to \infty} \Psi_{\Omega_{t_j}}(E_{t_j}; B_R) \,. \tag{2.9}$$

Indeed, let $F_0 \subset \Omega_0$ be such that

$$F_0 \Delta E_0 \subset B_R \cap \overline{\Omega_0}$$
.

By well-known properties of the trace of a BV function, for all j, for a.e. $0 < \rho < R$ and \mathcal{H}^{n-1} -a.e. $x \in \partial B_{\rho}$, we have

$$\mathbf{1}_{E_{t_j}}(x) = \operatorname{Tr}^{\pm}(E_{t_j}, \partial B_{\rho})(x) \quad \text{and} \quad \mathbf{1}_{E_0}(x) = \operatorname{Tr}^{\pm}(E_0, \partial B_{\rho})(x),$$
(2.10)

where $\operatorname{Tr}^{\pm}(E, \partial B_{\rho})$ denotes the inner (+) or outer (-) trace of $\mathbf{1}_{E}$ on ∂B_{ρ} . By the L^{1}_{loc} convergence of $E_{t_{j}}$ to E_{0} , we can choose $\rho < R$ with the above property, and such that

$$F_0 \Delta E_0 \subset B_\rho \cap \overline{\Omega_0}$$
 and $\lim_{j \to \infty} \int_{\partial B_\rho \cap \Omega_0} |\mathbf{1}_{E_{t_j}} - \mathbf{1}_{E_0}| d\mathcal{H}^{n-1} = 0.$ (2.11)

For any j we define

$$F_j := \left[(F_0 \cap B_\rho) \cup \left(E_{t_j} \cap (B_R \setminus B_\rho) \right) \right] \cap \Omega_{t_j}.$$

By construction, and thanks to (2.11), we have

$$F_j \Delta E_{t_j} \subset \subset B_R \cap \overline{\Omega_{t_j}}$$

and

$$P(F_j; B_R \cap \Omega_{t_j}) = P(F_0; B_\rho \cap \Omega_{t_j}) + P(E_{t_j}; (B_R \setminus \overline{B_\rho}) \cap \Omega_{t_j}) + \int_{\partial B_\rho \cap \Omega_{t_j}} |\mathbf{1}_{E_{t_j}} - \mathbf{1}_{E_0}| d\mathcal{H}^{n-1},$$

hence owing to (2.10) we infer $P(E_{t_j}; \partial B_\rho \cap \Omega_{t_j}) = 0$, and obtain

$$\Psi_{\Omega_{t_j}}(E_{t_j}; B_R) \ge P(E_{t_j}; B_R \cap \Omega_{t_j}) - P(F_j; B_R \cap \Omega_{t_j})$$

= $P(E_{t_j}; B_\rho \cap \Omega_{t_j}) - P(F_0; B_\rho \cap \Omega_{t_j}) - \int_{\partial B_\rho \cap \Omega_{t_j}} |\mathbf{1}_{E_{t_j}} - \mathbf{1}_{E_0}| d\mathcal{H}^{n-1}$
 $\ge P(E_{t_j}; B_\rho \cap \Omega_{t_j}) - P(F_0; B_R \cap \Omega_0) - \int_{\partial B_\rho \cap \Omega_{t_j}} |\mathbf{1}_{E_{t_j}} - \mathbf{1}_{E_0}| d\mathcal{H}^{n-1}.$ (2.12)

Given $\varepsilon > 0$, we can select j_0 and possibly update the choice of $0 < \rho < R$ in such a way that all the previous requirements are still satisfied, and moreover we have

$$P(E_0; B_\rho \cap \Omega_{t_{j_0}}) \ge P(E_0; B_R \cap \Omega_0) - \varepsilon.$$
(2.13)

Now by (2.12), we infer that for $j \ge j_0$

$$\Psi_{\Omega_{t_j}}(E_{t_j}; B_R) \ge P(E_{t_j}; B_\rho \cap \Omega_{t_{j_0}}) - P(F_0; B_R \cap \Omega_0) - \int_{\partial B_\rho \cap \Omega_{t_j}} |\mathbf{1}_{E_{t_j}} - \mathbf{1}_{E_0}| d\mathcal{H}^{n-1}$$
(2.14)

Taking the lim inf in (2.14), and exploiting (2.11), (2.13) together with the lower-semicontinuity of the perimeter, we deduce that

$$\liminf_{j \to \infty} \Psi_{\Omega_{t_j}}(E_{t_j}; B_R) \ge P(E_0; B_\rho \cap \Omega_{t_{j_0}}) - P(F_0; B_R \cap \Omega_0)$$
$$\ge P(E_0; B_R \cap \Omega_0) - P(F_0; B_R \cap \Omega_0) - \varepsilon,$$

which proves (2.9) by the arbitrary choice of ε and F_0 . Finally, by (2.1) we obtain

$$\Psi_{\Omega_{t_j}}(E_{t_j}; B_R) = \frac{1}{t_j^{n-1}} \Psi_{\Omega}(E; B_{t_j R}) \le \omega_n^{1-\frac{1}{n}} R^{n-1} \psi_{\Omega}(E; 0, t_j R) \longrightarrow 0,$$

which concludes the proof.

2.4. Boundary monotonicity formula in Lipschitz cones. We shall need the following conical deviation estimate for a set G minimizing the relative perimeter in a Lipschitz cone Ω_0 .

Theorem 2.3. Let G be a finite perimeter set in Ω_0 , and let $\mathcal{B}_r := \Omega_0 \cap B_r$. Then

$$\left\{ \left| \frac{x}{|x|^{n}} \cdot D\mathbf{1}_{G} \right| (\mathcal{B}_{r} \setminus \mathcal{B}_{\rho}) \right\}^{2} \leq 2 |x|^{1-n} |D\mathbf{1}_{G}| (\mathcal{B}_{r} \setminus \mathcal{B}_{\rho}) \qquad (2.15)$$

$$\left\{ r^{1-n} |D\mathbf{1}_{G}| (\mathcal{B}_{r}) - \rho^{1-n} |D\mathbf{1}_{G}| (\mathcal{B}_{\rho}) + (n-1) \int_{\rho}^{r} t^{-n} \Psi_{\Omega_{0}}(G; B_{t}) dt \right\},$$

for almost every $0 < \rho < r$.

Theorem 2.3 is well-known under the assumption that the origin is an internal point, and $B_r \subset \Omega$, see [14], [24]. When the origin is a vertex of Ω_0 , the classical proof is adapted without much effort, and we provide it here for the reader's convenience.

Remark 2.4. We note that, if G is a perimeter minimizer in Ω_0 , then $\Psi_{\Omega_0}(E; B_r) = 0$, hence by (2.15) one obtains that $r^{1-n}P_{\Omega_0}(G, B_r)$ is non-decreasing w.r.t. r.

Proof of Theorem 2.3. We follow the proof of [14, Lemma 5.8]. First, we take $f \in \mathcal{C}^{\infty}(\mathcal{B}_R) \cap BV(\mathcal{B}_R)$ and prove that, for almost all $0 < \rho < r < R$, one has

$$\left\{ \left| \frac{x}{|x|^{n}} \cdot Df \right| (\mathcal{B}_{r} \setminus \mathcal{B}_{\rho}) \right\}^{2} \leq 2 |x|^{1-n} \cdot |Df| (\mathcal{B}_{r} \setminus \mathcal{B}_{\rho}) \cdot \left\{ r^{1-n} |Df| (\mathcal{B}_{r}) - \rho^{1-n} |Df| (\mathcal{B}_{\rho}) + (n-1) \int_{\rho}^{r} t^{-n} \Psi_{\Omega_{0}}(f;t) dt \right\},$$
(2.16)

where

$$\Psi_{\Omega_0}(f;t) \equiv |Df|(\mathcal{B}_t) - \inf\{|Dg|(\mathcal{B}_t): g \in BV(\mathcal{B}_t), \operatorname{spt}(g-f) \subset \mathcal{B}_t \cap \overline{\Omega_0}\}.$$
(2.17)
For $0 < t < R$, let

 $01 \ 0 < t < 1t, 1et$

$$f_t(x) = \begin{cases} f(x) & \text{if } t < |x| < R\\ f\left(t\frac{x}{|x|}\right) & \text{if } 0 < |x| < t. \end{cases}$$

Standard computations yield

$$\int_{\mathcal{B}_t} |Df_t| dx = \frac{t}{n-1} \int_{\partial B_t \cap \Omega_0} |Df| \left\{ 1 - \frac{\langle x, Df \rangle^2}{|x|^2 |Df|^2} \right\}^{1/2} d\mathcal{H}^{n-1}.$$

By definition of $\Psi_{\Omega_0}(f;t)$, we have

$$\begin{split} \int_{\mathcal{B}_t} |Df| dx - \Psi_{\Omega_0}(f;t) &\leq \int_{\mathcal{B}_t} |Df_t| dx \\ &= \frac{t}{n-1} \int_{\partial B_t \cap \Omega_0} |Df| \left\{ 1 - \frac{\langle x, Df \rangle^2}{|x|^2 |Df|^2} \right\}^{1/2} d\mathcal{H}^{n-1} \\ &\leq \frac{t}{n-1} \int_{\partial B_t \cap \Omega_0} |Df| d\mathcal{H}^{n-1} - \frac{t}{2(n-1)} \int_{\partial B_t \cap \Omega_0} \frac{\langle x, Df \rangle^2}{|x|^2 |Df|} d\mathcal{H}^{n-1} \,. \end{split}$$

Multiplying both sides by $(n-1)t^{-n}$, we get

$$\frac{t^{1-n}}{2} \int_{\partial B_t \cap \Omega_0} \frac{\langle x, Df \rangle^2}{|x|^2 |Df|} d\mathcal{H}^{n-1} \leq t^{1-n} \int_{\partial B_t \cap \Omega_0} |Df| d\mathcal{H}^{n-1} - (n-1) t^{-n} \int_{\mathcal{B}_t} |Df| dx + (n-1) t^{-n} \Psi_{\Omega_0}(f;t) \\
= \frac{d}{dt} \left(t^{1-n} \int_{\mathcal{B}_t} |Df| dx \right) + (n-1) t^{-n} \Psi_{\Omega_0}(f;t),$$
(2.18)

and, by integrating (2.18) with respect to t between ρ and r, we obtain

$$\frac{1}{2} \int_{\mathcal{B}_r \setminus \mathcal{B}_\rho} \frac{\langle x, Df \rangle^2}{|x|^{n+1} |Df|} dx \le r^{1-n} \int_{\mathcal{B}_r} |Df| dx - \rho^{1-n} \int_{\mathcal{B}_\rho} |Df| dx + (n-1) \int_{\rho}^r t^{-n} \Psi_{\Omega_0}(f; t) dt \,.$$
(2.19)

By Hölder's inequality, we find that

$$\left\{\int_{\mathcal{B}_r \setminus \mathcal{B}_\rho} \frac{|\langle x, Df \rangle|}{|x|^n} dx\right\}^2 \le \left\{\int_{\mathcal{B}_r \setminus \mathcal{B}_\rho} |x|^{1-n} |Df| dx\right\} \left\{\int_{\mathcal{B}_r \setminus \mathcal{B}_\rho} \frac{\langle x, Df \rangle^2}{|x|^{n+1} |Df|} dx\right\} \,. \tag{2.20}$$

Then, the combination of (2.19) and (2.20) gives the desired estimate (2.16). Finally, we can approximate χ_E by a sequence of smooth BV functions and pass to the limit in (2.16), thus getting (2.15).

3. FLATNESS OF THE BLOW-UP OF AN ALMOST-MINIMIZER

Thanks to Lemma 2.2, we know that any blow-up sequence of the almost-minimizer E in Ω , with respect to $0 \in \partial \Omega$, converges to a minimizer E_0 in the tangent cone Ω_0 , up to subsequences. Applying Lemma 2.2 to E_0 , we deduce the existence of a sequence $s_j \searrow 0$ and a set E_{00} minimizing the relative perimeter in Ω_0 , such that

$$(E_0)_{s_i} \longrightarrow E_{00} \quad \text{in } L^1_{loc}(\Omega_0).$$
 (3.1)

Actually, up to a diagonal argument, one can even show that there exists a suitable sequence $t_j \searrow 0$ such that $E_{t_j} \to E_{00}$ in $L^1_{loc}(\mathbb{R}^n)$. A consequence of Theorem 2.3 is the following

Proposition 3.1. The set E_{00} is a perimeter-minimizing cone with a vertex at the origin.

Proof. The argument requires a slight variant of the proof of [14, Theorem 9.3]. Let $s_j \searrow 0$ be such that $(E_0)_{s_j}$ converges to E_{00} in $L^1_{loc}(\mathbb{R}^n)$. For t > 0, we let $p(t) := t^{1-n} P_{\Omega_0}(E_0, B_t)$ and note that, by Remark 2.4, p is non-decreasing in t and, for all R > 0, we have

$$p(tR) = R^{1-n} P_{\Omega_0}((E_0)_t, B_R)$$

Consequently, as the perimeter measure of $(E_0)_{s_j}$ weakly-* converges to that of E_{00} by a wellknown property of perimeter almost-minimizers (see, e.g., [30]), we deduce that, for almost every R > 0,

$$R^{1-n}P_{\Omega_0}(E_{00}, B_R) = \lim_{i \to \infty} p(s_j R).$$
(3.2)

At the same time, the monotonicity of p(t) implies that

$$\lim_{j\to\infty} p(s_j R) = \lim_{t\to 0^+} p(t) \,,$$

hence the limit does not depend on R. Then by approximating a generic radius R by means of sequences of smaller/larger radii for which (3.2) is satisfied, one easily obtains that

$$R^{1-n}P_{\Omega_0}(E_{00}, B_R) = P_{\Omega_0}(E_{00}, B_1)$$
 for all $R > 0$.

We can now apply (2.15) and get

$$|x \cdot D\mathbf{1}_{E_{00}}| (B_r \setminus B_{\rho}) = 0 \quad \text{for all } 0 < \rho < r.$$

This implies that

$$\langle x, \nu_{E_{00}}(x) \rangle = 0$$
 for \mathcal{H}^{n-1} -a.e. $x \in \partial^* E_{00}$

By [24, Proposition 28.8], we conclude that E_{00} is, up to null sets, a cone with vertex at the origin.

3.1. Characterization of the conical minimizer in \mathbb{R}^3 . Starting from an almost-minimizer E in Ω which satisfies volume density estimates at the origin, and applying Lemma 2.2 at most twice, we have obtained a conical minimizer E_{00} of the relative perimeter in the tangent cone Ω_0 . The next theorem shows that, in dimension n = 3, $\partial E_{00} \cap \Omega_0$ coincides with a convex angle contained in a 2-plane through the origin, that meets $\partial \Omega_0$ orthogonally.

Theorem 3.2. Let n = 3 and E_{00} be the conical minimizer obtained in the previous subsection. Then $\partial E_{00} \cap \Omega_0$ coincides with a 2-plane intersected with Ω_0 , that meets $\partial \Omega_0 \setminus \{0\}$ orthogonally.

Proof. Set $F = E_{00}$ for brevity, then the proof is accomplished by showing that there exists exactly one geodesic arc $\gamma \subset \partial B_1 \cap \Omega_0$ such that

$$\partial F \cap \partial B_1 \cap \Omega_0 = \gamma \,, \tag{3.3}$$

and γ meets $\partial \Omega_0 \cap \partial B_1$ orthogonally. We split the proof into some steps.

Step 1. We claim that $\partial F \cap \partial B_1 \cap \Omega_0$ is made of countably-many (open) geodesic arcs γ_i , $i \in \mathbb{N}$, such that

$$\gamma_i \cap \gamma_j = \emptyset$$
, for $i \neq j$, $\bigcup_i \gamma_i = \partial F \cap \partial B_1 \cap \Omega_0$.

By interior regularity, $\partial F \cap \Omega_0$ is smooth, and its outer normal vector ν_F is orthogonal to the radial directions (recall that F is a cone with vertex at the origin). Hence, ∂F intersects transversally $\partial B_1 \cap \Omega_0$ along smooth curves γ_i that cannot cross each other. Since F has locally-finite perimeter, the family of these curves is at most countable. Let us now show that γ_i is a geodesic arc, for all i. With a slight abuse of notation, we identify γ_i with its arc-legth parametrization defined on the interval $(0, L_i)$, where L_i is the length of the curve. The connected component of ∂F that intersects ∂B_1 along γ_i can be then parametrized through

$$\sigma_i(s,t) := s \,\gamma_i(t) \,, \, \text{for } s > 0, \, t \in (0, L_i).$$
(3.4)

We can choose the parametrization $\gamma_i(t)$ in such a way that $\nu_F(\sigma_i(s,t)) = \gamma_i(t) \times \gamma'_i(t)$, for all s > 0. Exploiting (3.4), and using $\operatorname{div}_{\partial F} \nu_F = 0$ by the minimality of F, we infer

$$0 = \operatorname{div}_{\partial F} \nu_F(\sigma_i(s, t)) \tag{3.5}$$

$$= \frac{d}{dt} (\gamma_i(t) \times \gamma'_i(t)) \cdot \gamma'_i(t)$$

= $-\gamma_i(t) \times \gamma'_i(t) \cdot \gamma''_i(t)$, for all t. (3.6)

Since we also have $\gamma'_i(t) \cdot \gamma''_i(t) = 0$ by the choice of the arc-length parametrization, and observing that $\{\gamma'_i(t), \gamma_i(t) \times \gamma'_i(t)\}$ is an orthonormal basis for the tangent space to ∂B_1 at $\gamma(t)$, we conclude that $\gamma''_i(t)$ is orthogonal to the tangent space to ∂B_1 at $\gamma(t)$, which is precisely the definition of geodesic arc.

Step 2. We prove that γ_i meets $\partial \Omega_0$ orthogonally at its endpoints. More precisely, if p is an endpoint of γ_i , then Ω_0 admits a unique supporting plane at p, hence the outer unit normal vector $\nu_0(p)$ to $\partial \Omega_0$ at p is well-defined, and moreover if we denote by ν_i the constant unit outer normal to the connected component of ∂F containing γ_i , we have

$$\nu_i \cdot \nu_0(p) = 0. (3.7)$$

Let us first prove that Ω_0 admits a unique supporting plane at p. Up to a rotation, we may assume that p = (0, 0, 1), hence it follows that $\nu_i \cdot e_3 = 0$. Owing to Lemma 3.3 below, we can

11

find a sequence $t_j \searrow 0$ such that $\Omega_0^{p,t_j} := t_j^{-1}(\Omega_0 - p)$ locally converge to a cylinder of type $C \times \mathbb{R}$, and $F^{p,t_j} := t_j^{-1}(F-p)$ locally converge to a cylinder that can be written as $G \times \mathbb{R}$, where $G \subset C$. Moreover, both C, and G are cones with respect to 0 in the plane z = 0, and G is perimeter-minimizing in C. By convexity of C, up to a further rotation, we can assume that

$$C = \{(x_1, x_2, 0) : x_2 > \lambda |x_1|\},\$$

for some $\lambda \geq 0$. Clearly, $\lambda = 0$ if and only if p admits a unique supporting plane for Ω_0 . The only possibility is then that $\partial G \cap C$ is made of finitely many half-lines $L_1, ..., L_k$ of the form $L_j = \{t v_j : t \geq 0\}$, for some unit vectors v_j . Up to relabeling, we can assume

$$0 < v_1 \cdot e_2 \leq v_j \cdot e_2$$
 for all j

If either $\lambda > 0$ or k > 1, we could replace an initial portion of L_1 with a projection segment onto the closest side of C, which strictly decreases the perimeter and thus contradicts the fact that G is perimeter-minimizing in C. Hence we necessarily have $\lambda = 0$ and k = 1, i.e., there exists a unique supporting plane to Ω_0 at p with $\nu_0(p) = -e_2$, and moreover $v_1 = e_2$. This proves the claim and, additionally, shows that two different geodesic arcs cannot share a common endpoint.

Step 3. Finally, we prove that $\partial F \cap \partial B_1 \cap \Omega_0$ is made of exactly one geodesic arc.

Suppose by contradiction that there exist two geodesic arcs $\gamma_1 \neq \gamma_2$ contained in $\partial F \cap \partial B_1 \cap \Omega_0$. By the previous steps we know that

$$\overline{\gamma_1} \cap \overline{\gamma_2} = \emptyset \,. \tag{3.8}$$

For i = 1, 2 we denote by Π_i the plane through the origin that contains γ_i , and by p_i , q_i the boundary points of γ_i (see Figure 1).

Then, we consider the point $N_i \in \partial B_1$ such that γ_i is contained in the equator with north pole N_i and south pole $S_i = -N_i$, and denote by μ_{p_i}, μ_{q_i} the corresponding meridians connecting N_i with S_i and passing through p_i and q_i , respectively. These meridians bound a region of ∂B_1 , that we denote as Σ_i , which satisfies

$$\Omega_0 \cap \partial B_1 \subset \Sigma_i, \qquad i = 1, 2.$$

Incidentally, thanks to the previous step, Σ_i is also obtained by intersecting the sphere ∂B_1 with the wedge W_i given by the intersection of the two supporting half-spaces to Ω_0 at p_i and q_i , respectively. Since in particular W_i is a convex cone, it is immediate to check that $\Sigma_i = W_i \cap \partial B_1$ is geodesically convex. Moreover, using the fact that the angle formed by the vectors p_i, q_i is strictly smaller than π (recall that the origin is an isolated vertex for Ω_0) we infer that the internal angle formed by the two geodesic sides of Σ_i , i.e. the meridians μ_{p_i} and μ_{q_i} , at N_i (or S_i) is strictly smaller than π .

Now, observe that the closure of Σ_i is the union of two closed geodesic triangles T_{N_i}, T_{S_i} with vertices p_i, q_i, N_i and p_i, q_i, S_i , respectively. Since in particular $\gamma_2 \subset \Omega_0 \cap \partial B_1 \subset \Sigma_1$, and γ_2 has a strictly positive distance from γ_1 , we must have that either $\gamma_2 \subset T_{N_1}$ or $\gamma_2 \subset T_{S_1}$. Without loss of generality, we assume $\gamma_2 \subset T_{N_1}$.

Now, we set $\tilde{p}_2 = \Pi_2 \cap \mu_{p_1}$ and $\tilde{q}_2 = \Pi_2 \cap \mu_{q_1}$, and denote by $\tilde{\gamma}_2$ the geodesic connecting \tilde{p}_2 and \tilde{q}_2 , and by $\tilde{\Sigma}_2$ the associated geodesically convex region bounded by the meridians $\mu_{\tilde{p}_2}, \mu_{\tilde{q}_2}$ meeting at poles $\tilde{N}_2 = N_2$ and $\tilde{S}_2 = S_2$. Clearly we have $\gamma_2 \subset \tilde{\gamma}_2$ and thus $\Sigma_2 \subset \tilde{\Sigma}_2$. Moreover,



FIGURE 1. A possible geometric configuration in the contradiction argument for the proof of Step 3.

we have

$\widetilde{p}_2, \widetilde{q}_2 \in T_{N_1} \setminus \{p_1, q_1, N_1\}.$

Indeed, the geodesic $\tilde{\gamma}_2$ cannot intersect γ_1 , hence it is contained in T_{N_1} and its closure is disjoint from γ_1 because $\Pi_2 \cap \overline{\gamma_1} = \overline{\gamma_2} \cap \overline{\gamma_1} = \emptyset$, by (3.8); moreover, if we had $\tilde{p}_2 = N_1$ (or $\tilde{q}_2 = N_1$) we would conclude that $\mu_{q_1} \subset \Pi_2$ (respectively, $\mu_{p_1} \subset \Pi_2$), but this is impossible because Π_2 and $\overline{\gamma_1}$ are disjoint.

Now, consider the geodesic quadrilateral D determined by the four points $p_1, \tilde{p}_2, \tilde{q}_2, q_1$. By the previous argument, $D \subset T_{N_1}$. Denote by $\alpha_1, \tilde{\alpha}_2, \tilde{\beta}_2, \beta_1$ the angles formed by the pairs of geodesic sides meeting at the respective vertices. Then, consider the two geodesic triangles $R_1 = p_1 \tilde{p}_2 q_1$ and $R_2 = \tilde{p}_2 \tilde{q}_2 q_1$. Call $\tilde{\alpha}_{2,1}$ the internal angle to R_1 at \tilde{p}_2 , and $\tilde{\alpha}_{2,2}$ the internal angle to R_2 at \tilde{p}_2 . Similarly, call $\beta_{1,1}$ the internal angle to R_1 at q_1 , and $\beta_{1,2}$ the internal angle to R_2 at q_1 . We thus have $\tilde{\alpha}_2 = \tilde{\alpha}_{2,1} + \tilde{\alpha}_{2,2}, \beta_1 = \beta_{1,1} + \beta_{1,2}$, and therefore we deduce

$$\alpha_1 = \beta_1 = \pi/2 \,, \tag{3.9}$$

$$\max(\widetilde{\alpha}_2, \beta_2) \le \pi/2. \tag{3.10}$$

Indeed, (3.9) follows from the orthogonality of γ_1 with the meridians μ_{p_1} and μ_{q_1} , while (3.10) follows from the fact that the quadrilateral D is contained in one of the two geodesic triangles $\widetilde{T}_{N_2} = \widetilde{p}_2 \widetilde{q}_2 N_2$, $\widetilde{T}_{S_2} = \widetilde{p}_2 \widetilde{q}_2 S_2$ (indeed, by a symmetric argument, we have either $\gamma_1 \subset \widetilde{T}_{N_2}$ or $\gamma_1 \subset \widetilde{T}_{S_2}$) and we know by construction that the internal angles to \widetilde{T}_{N_2} (or \widetilde{T}_{S_2}) at \widetilde{p}_2 and at \widetilde{q}_2 are both equal to $\pi/2$.

Now we notice that R_1 must be a non-degenerate geodesic triangle, because it possesses an internal angle at p_1 measuring $\alpha_1 = \pi/2$, and the other two vertices do not coincide. Thus we have that the sum of the internal angles of R_1 satisfies

$$\alpha_1 + \tilde{\alpha}_{2,1} + \beta_{1,1} > \pi \,. \tag{3.11}$$

At the same time, the sum of the internal angles of R_2 is not smaller than π :

$$\widetilde{\alpha}_{2,2} + \widetilde{\beta}_2 + \beta_{1,2} \ge \pi \,. \tag{3.12}$$

By combining (3.9), (3.10), (3.11) and (3.12), we reach the contradiction

$$2\pi < (\alpha_1 + \widetilde{\alpha}_{2,1} + \beta_{1,1}) + (\widetilde{\alpha}_{2,2} + \widetilde{\beta}_2 + \beta_{1,2}) = \alpha_1 + \widetilde{\alpha}_2 + \widetilde{\beta}_2 + \beta_1 \le 2\pi$$

and this completes the proof of the theorem.

Next we state a slight variant of the classical Federer's Reduction Lemma, which is used in the proof of Theorem 3.2. In what follows, by "cone" we shall always mean a cone with respect to the origin.

Lemma 3.3. Let $K \subset \mathbb{R}^n$ be a convex cone, $C \subset K$ be a minimizing cone for the relative perimeter. Let $x_0 \in \partial C \cap \partial K \setminus \{0\}$. For t > 0, we set

$$C_t := x_0 + \frac{C - x_0}{t}, \quad K_t := x_0 + \frac{K - x_0}{t}.$$

Then there exist a sequence $t_j \searrow 0$ and two sets C_0 , K_0 such that

$$C_j := C_{t_j} \longrightarrow C_0, \quad K_j := K_{t_j} \longrightarrow K_0 \tag{3.13}$$

in L^1_{loc} -topology, $C_0 \subset K_0$ and C_0 , K_0 are cylinders with axis coinciding with the line joining 0 to x_0 . Moreover, the sets $C'_0 := C_0 \cap x_0^{\perp}$ and $K'_0 := K_0 \cap x_0^{\perp}$ are cones with respect to 0 in the hyperplane x_0^{\perp} , and C'_0 is perimeter-minimizing in K'_0 .

The proof of this lemma is omitted, as it can be obtained via a slight modification of [14, Proposition 9.9].

4. Proof of the main result

We now dispose of all the necessary tools to prove Theorem 1.1. We argue by contradiction. By the results of the previous section, via a blow-up argument we can restrict the proof to the case of a domain Ω_0 being a convex cone with vertex at the origin, and of a minimizer given by the intersection with Ω_0 of a half-space whose boundary plane passes through the origin and meets $\partial \Omega_0$ orthogonally.

Before giving the proof of Theorem 1.1, we introduce a class of convex cones that will play a key role in the first part of the proof.

Definition 4.1. We say that a convex cone $C \subset \mathbb{R}^3$ is a pyramid provided there exist two wedges W_1, W_2 with orthogonally incident spines such that $C = W_1 \cap W_2$.

We note that a pyramid $C \subset \mathbb{R}^3$ is always a cone with vertex at the point V_0 in which the spines of the wedges intersect each other. Moreover, up to a rotation and a translation, there exist a, b > 0 such that

$$C = C_{a,b} := \{ x \in \mathbb{R}^3 : x_3 \ge \max\{a|x_1|, b|x_2|\} \}.$$

Proof (of Theorem 1.1). The proof is split into two steps. In the first, we show that a suitable plane through the vertex of a *pyramid*, i.e., of a cone over a rectangle, cannot be area-minimizing in the pyramid itself. In the second, we employ a "packing-box" technique that allows us to reduce the case of a general convex cone to that of a suitably associated pyramid.

Step 1. Consider a pyramid cone $C_{a,b}$. We want to show that the plane

$$\pi_0 = \{ x \in \mathbb{R}^3 : x_1 = 0 \}$$

is not locally area-minimizing in $C_{a,b}$. To do so, we build a family of competitors which improve the area of π_0 in $C_{a,b}$. For $\varepsilon \ge 0$, let $\pi_{\varepsilon} = \{x \in \mathbb{R}^3 : x_1 = \varepsilon\}$ and define

$$R_{\varepsilon} := C_{a,b} \cap \{ x \in \pi_{\varepsilon} : x_3 \le 1 \}, \qquad A_{\varepsilon} := \mathcal{H}^2(R_{\varepsilon})$$

We note that R_0 is a triangle in the plane π_0 , and that R_{ε} is a trapezium in the plane π_{ε} whenever $0 < \varepsilon < 1$. Moreover, up to translations, R_{ε} is obtained from R_0 by removing a triangle of area $\frac{a^2 \varepsilon^2}{b}$, so that we have

$$A_{\varepsilon} = A_0 - \frac{a^2}{b} \varepsilon^2 \,. \tag{4.1}$$

The idea is now to connect the trapezium R_{ε} with $\pi_0 \cap C_{a,b}$ in order to obtain a local variation of $\pi_0 \cap C_{a,b}$. We formulate the problem in the following way. Let h > 0 and let T_h be the trapezium defined as

$$T_h = C_{a,b} \cap \{x \in \pi_0 : 1 \le x_3 \le 1 + h\}.$$

We immediately note that

$$\mathcal{H}^2(T_h) = \frac{h(2+h)}{b} \,. \tag{4.2}$$

We look for those smooth functions ϕ_h defined on the segment $\{1 \le x_3 \le 1 + h\}$ satisfying the following conditions:

$$\phi_h(1) = 1, \qquad \phi_h(1+h) = 0.$$
 (4.3)

We observe that, looking at ϕ_h as a function of both variables x_2 and x_3 defined in T_h , the ruled surface

$$G_{\varepsilon}(\phi_h) = \{ (\varepsilon \phi_h(x_3), x_2, x_3) : (x_2, x_3) \in T_h \}$$

connects R_{ε} with $\pi_0 \cap C_{a,b}$. By suitably choosing h and the map ϕ_h , we claim that

$$A_{\varepsilon} + \mathcal{H}^2(G_{\varepsilon}(\phi_h)) < A_0 + \mathcal{H}^2(T_h).$$
(4.4)

Using (4.1), (4.2), and the area formula, (4.4) turns out to be equivalent to

$$\iint_{T_h} \sqrt{1 + \varepsilon^2 |\phi_h'(x_3)|^2} dx_2 dx_3 < \frac{a^2}{b} \varepsilon^2 + \frac{h(2+h)}{b}.$$
(4.5)

In order to guarantee (4.5), we only need to impose that the second-order derivative at 0 of the left-hand side is strictly smaller than the same derivative of the right-hand side. Differentiating both sides and applying Dominated Convergence, it suffices to choose h and ϕ_h so that

$$\int_{1}^{1+h} t \,\phi_h'(t)^2 dt < a^2 \,. \tag{4.6}$$

Then, for $\alpha > 0$, we choose

$$\phi_h = \phi_{h,\alpha}(t) := \frac{(1+h)^{\alpha}t^{-\alpha} - 1}{(1+h)^{\alpha} - 1}$$

We observe that $\phi_{h,\alpha}$ fulfills (4.3). Taking $\phi_h = \phi_{h,\alpha}$, condition (4.6) becomes

$$\frac{\alpha}{2} \frac{(1+h)^{\alpha} + 1}{(1+h)^{\alpha} - 1} = \int_{1}^{1+h} t \,\phi_{h,\alpha}'(t)^2 dt < a^2 \,. \tag{4.7}$$

As $h \to +\infty$, the term on the left-hand side of (4.7) tends to $\frac{\alpha}{2}$, hence it is enough to choose $\alpha < 2a^2$ and h large enough to enforce (4.7). This ultimately proves (4.4) and shows that π_0 cannot be area-minimizing in $C_{a,b}$.

Step 2. Let now Ω_0 be a generic convex cone with vertex at the origin. Thanks to Theorem 3.2, and up to rotations, we may suppose that the boundary of the minimizer E_{00} is the intersection of the plane π_0 with Ω_0 , hence there exists b > 0 such that

$$\partial E_{00} \cap \Omega_0 = \{(0, x_2, x_3) : x_3 \ge b |x_2|\}.$$

Now, by Theorem 3.2 we have

$$\Omega_0 \subset W_1 := \{ x \in \mathbb{R}^3 : x_3 \ge b |x_2| \}.$$

Since the origin is an isolated vertex for Ω_0 , it is not possible that $\partial \Omega_0$ contains the whole line $\{(t, 0, 0) : t \in \mathbb{R}\}$, hence there must exist a > 0 such that the pyramid $C_{a,b}$ verifies either

$$\Omega_0 \cap \{x_1 \ge 0\} \subset C_{a,b} \cap \{x_1 \ge 0\}$$
(4.8)

or

$$\Omega_0 \cap \{x_1 \le 0\} \subset C_{a,b} \cap \{x_1 \le 0\}.$$
(4.9)

We can assume for instance that (4.8) holds true, otherwise we simply flip the argument. We take $\varepsilon > 0$ and set

$$\hat{R}_{\varepsilon} := \Omega_0 \cap \left\{ x \in \pi_{\varepsilon} : \ x_3 \le 1 \right\}, \qquad \hat{A}_{\varepsilon} := \mathcal{H}^2(\hat{R}_{\varepsilon}) \,.$$

With the choice of suitable values h and α , we already know that the connection map $\phi_{h,\alpha}$ constructed in the proof of Theorem 3.2 satisfies

$$A_{\varepsilon} + \mathcal{H}^2(G_{\varepsilon}(\phi_{h,\alpha})) < A_0 + \mathcal{H}^2(T_h)$$

whenever ε is small enough. Finally, we observe that

$$\hat{A}_{\varepsilon} + \mathcal{H}^2(G_{\varepsilon}(\phi_{h,\alpha}) \cap \Omega_0) \le A_{\varepsilon} + \mathcal{H}^2(G_{\varepsilon}(\phi_{h,\alpha}))$$
$$\hat{A}(0) = A(0),$$

which shows that π_0 cannot be a minimizer in Ω_0 . This concludes the proof of Theorem 1.1.

References

- [1] William K Allard. On the first variation of a varifold. Annals of mathematics, 95(3):417-491, 1972.
- [2] William K Allard. On the first variation of a varifold: boundary behavior. Annals of Mathematics, 101(3):418–446, 1975.
- [3] Jürgen Bokowski and Emanuel Sperner, Jr. Zerlegung konvexer Körper durch minimale Trennflächen. J. Reine Angew. Math., 311/312:80–100, 1979.
- [4] Alessandro Carlotto. Free boundary minimal surfaces: a survey of recent results. Rendiconto dell'Accademia delle Scienze Fisiche e Matematiche, 86(1):103–121, 12 2019.
- [5] Jin Tzu Chen, Robert Finn, and Erich Miersemann. Capillary surfaces in wedge domains. Pacific J. Math., 236(2):283–305, 2008.
- [6] Jaigyoung Choe, Mohammad Ghomi, and Manuel Ritoré. The relative isoperimetric inequality outside convex domains in Rⁿ. Calc. Var. Partial Differential Equations, 29(4):421–429, 2007.
- [7] Paul Concus and Robert Finn. On the behavior of a capillary surface in a wedge. Proc. Natl. Acad. Sci. USA, 63(2):292–299, 1969.
- [8] R. Courant. On Plateau's problem with free boundaries. Proc. Nat. Acad. Sci. U.S.A., 31:242–246, 1945.
- [9] G. De Philippis and F. Maggi. Regularity of free boundaries in anisotropic capillarity problems and the validity of Young's law. Arch. Ration. Mech. Anal., 216(2):473–568, 2015.

- [10] Nicholas Edelen and Chao Li. Regularity of free boundary minimal surfaces in locally polyhedral domains. Comm. Pure Appl. Math., 75(5):970–1031, 2022.
- [11] A. Figalli and E. Indrei. A sharp stability result for the relative isoperimetric inequality inside convex cones. J. Geom. Anal., 23(2):938–969, 2013.
- [12] Robert Finn. Equilibrium capillary surfaces, volume 284 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, New York, 1986.
- [13] N. Fusco and M. Morini. Total positive curvature and the equality case in the relative isoperimetric inequality outside convex domains. *Calc. Var. Partial Differential Equations*, 62(3):Paper No. 102, 32, 2023.
- [14] Enrico Giusti. Minimal surfaces and functions of bounded variation, volume 80 of Monographs in Mathematics. Birkhäuser Verlag, Basel, 1984.
- [15] Michael Grüter and Jürgen Jost. Allard type regularity results for varifolds with free boundaries. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 13(1):129–169, 1986.
- [16] Ernst Kuwert. Note on the isoperimetric profile of a convex body. In Geometric analysis and nonlinear partial differential equations, pages 195–200. Springer, Berlin, 2003.
- [17] K. E. Lancaster and D. Siegel. Behavior of a bounded non-parametric H-surface near a reentrant corner. Z. Anal. Anwendungen, 15(4):819–850, 1996.
- [18] Kirk E. Lancaster. Remarks on the behavior of nonparametric capillary surfaces at corners. Pacific J. Math., 258(2):369–392, 2012.
- [19] Gian Paolo Leonardi, Manuel Ritoré, and Efstratios Vernadakis. Isoperimetric inequalities in unbounded convex bodies. Mem. Amer. Math. Soc., 276(1354):1–86, 2022.
- [20] Gian Paolo Leonardi and Giorgio Saracco. The prescribed mean curvature equation in weakly regular domains. NoDEA Nonlinear Differential Equations Appl., 25(2):Paper No. 9, 29, 2018.
- [21] Gian Paolo Leonardi and Giacomo Vianello. Stability of axial 3-planes in 4-dimensional circular cones of large aperture. *In preparation*.
- [22] Hans Lewy. On minimal surfaces with partially free boundary. Comm. Pure Appl. Math., 4:1–13, 1951.
- [23] Pierre-Louis Lions and Filomena Pacella. Isoperimetric inequalities for convex cones. Proc. Amer. Math. Soc., 109(2):477–485, 1990.
- [24] Francesco Maggi. Sets of finite perimeter and geometric variational problems, volume 135 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2012. An introduction to geometric measure theory.
- [25] Emanuel Milman. On the role of convexity in isoperimetry, spectral gap and concentration. Invent. Math., 177(1):1–43, 2009.
- [26] Johannes C. C. Nitsche. Lectures on minimal surfaces. Vol. 1. Cambridge University Press, Cambridge, 1989. Introduction, fundamentals, geometry and basic boundary value problems, Translated from the German by Jerry M. Feinberg, With a German foreword.
- [27] Manuel Ritoré and Efstratios Vernadakis. Isoperimetric inequalities in Euclidean convex bodies. Trans. Amer. Math. Soc., 367(7):4983–5014, 2015.
- [28] Peter Sternberg and Kevin Zumbrun. On the connectivity of boundaries of sets minimizing perimeter subject to a volume constraint. Comm. Anal. Geom., 7(1):199–220, 1999.
- [29] Italo Tamanini. Il problema della capillarità su domini non regolari. Rend. Sem. Mat. Univ. Padova, 56:169– 191, 1976.
- [30] Italo Tamanini. Boundaries of Caccioppoli sets with Hölder-continuous normal vector. J. Reine Angew. Math., 334:27–39, 1982.
- [31] Italo Tamanini. Regularity results for almost minimal oriented hypersurfaces in \mathbb{R}^n . Quaderni del Dipartimento di Matematica dell'Università di Lecce, 1984.
- [32] Thomas Young. III. An essay on the cohesion of fluids. Philosophical transactions of the royal society of London, (95):65–87, 1805.

DIPARTIMENTO DI MATEMATICA, VIA SOMMARIVE 14, IT-38123 POVO - TRENTO (ITALY) *Email address*: gianpaolo.leonardi@unitn.it

DIPARTIMENTO DI MATEMATICA, VIA SOMMARIVE 14, IT-38123 POVO - TRENTO (ITALY) *Email address:* giacomo.vianello-1@unitn.it