Symmetries in the many-body problems, a method to find its analytical solution, and Helium atom spectrum

Siddhesh C. Ambhire Physics Dept. Case Western Reserve University Cleveland sca54@case.edu

ABSTRACT

In this work it is shown that there are symmetries beyond the Euclidean group E(3) in 3-body problem, and by extension in many-body problem, with inverse squared distance inter particle force. The symmetries in 3-body problem form a group: $SO(4 \times 3, 2 \times 3) / (C(3 \times 2))$, where C(n) is the planar translation group in n dimensions, which forms its Spectrum-Generating group. Some of these quantities commute with the Hamiltonian. The existence of these conserved quantities was verified by calculating energy spectrum of the Helium atom. This method can also be used to find symmetries in many-body problem, and to calculate energy levels, and wave-functions of more complicated systems, which include every possible atomic and molecular systems in chemistry.

Keywords Quantum Chemistry · Many-body problem · Analytical solution

This research aims at finding solution of many-body dynamics with Coulombic inter-particle interactions using a novel approach. The problem in its least complex form appears as a 3-body problem, for which no closed form solution existed before [24, 22, 21].

This problem has been with physicists for as long as since Newton's discovery of the Gravitational force. Newton managed to solve the classical 2-body problem, also know as the Kepler problem, but the solution to the 3-body problem was elusive. The conserved quantities in classical 2-body problems are very much important for finding solutions to the problem. Same conserved quantities are present in the quantum mechanical 2-body problem. For the classical 3-body problem, many different solutions have been found for certain small number of special cases, called the restricted 3-body problems, for example by Suvakov[25].

In this work, the problem was solved for the non-relativistic quantum mechanical case for Coulombic inter-particle forces, in parallel treatment of the 2-body problem by Fock[2], which have the same form as the Newtonian gravity for the classical analogous problems. Since this problem is non-relativistic, spins were not considered while solving the problem, since their full treatment requires use of Dirac like equation, which is relativistic, as opposed to Schrodinger equation, which was found for the non-relativistic case. No other assumptions were made.

In the end, the solution was compared to experimental data and earlier calculations for the energy levels from the Helium atom[23, 26]. They match very well, with 10^{-4} relative error for the ground state and 10^{-5} relative error for the first excited state.

SO(4,2) symmetry in the Kepler problem

SO(4) symmetry of Kepler problem was found by Fock [2]. Rogers [7] used these symmetries to find general solutions for the Kepler problem. The generators for SO(4) groups in Kepler problem were also described in Woit [13] and Gilmore [6] in details. What follows is a summary of these works which will be useful in later sections.

The non-relativistic Hamiltonian of Kepler problem is given by:

$$\mathcal{H} = \frac{p^2}{2\mu} - \frac{k}{q} \tag{1}$$

Angular momentum and Laplace-Runge-Lenz (LRL) vectors are given by:

$$egin{array}{rcl} m{L} &=& m{q} imes m{p} \ m{A} &=& m{p} imes m{L} \ - m{k} m{q} \ m{q} \end{array}$$

The quantum mechanical operator constructed from A is given by[3]:

$$\hat{A} = \frac{\hat{p} \times \hat{L} - \hat{L} \times \hat{p}}{2\mu} - k \frac{\hat{q}}{q}$$

These quantities satisfy following commutation relations:

$$\begin{aligned} [\mathcal{H}, \boldsymbol{L}] &= [\mathcal{H}, \boldsymbol{A}] &= 0\\ [L_i, L_j] &= i\hbar\epsilon_{ijk}L_k\\ [L_i, A_j] &= i\hbar\epsilon_{ijk}A_k\\ [A_i, A_j] &= \left(-\frac{2\mathcal{H}}{\mu}\right)i\hbar\epsilon_{ijk}L_k \end{aligned}$$

Also $L \cdot A = A \cdot L = 0$ i.e. A and L are orthogonal to each other. These quantities also obey:

$$\boldsymbol{A} \cdot \boldsymbol{A} = \frac{2\mathcal{H}}{\mu} \left(\boldsymbol{L} \cdot \boldsymbol{L} + \hbar^2 \right) + k^2$$
(2)

Redefining quantities as follows simplifies the commutator algebra:

$$A' = \sqrt{-\frac{\mu}{2\mathcal{H}}}A \tag{3a}$$

$$U = \frac{1}{2} \left(L + A' \right) \tag{3b}$$

$$V = \frac{1}{2} \left(L - A' \right) \tag{3c}$$

The simplified commutator algebra which is also a Lie Algebra is given by:

$$[U_i, U_j] = i\hbar\epsilon_{ijk}U_k \tag{4a}$$

$$[V_i, V_j] = i\hbar\epsilon_{ijk}V_k \tag{4b}$$

$$\begin{bmatrix} V_i, V_j \end{bmatrix} = i\hbar\epsilon_{ijk}V_k \tag{4b}$$
$$\begin{bmatrix} U_i, V_j \end{bmatrix} = 0 \tag{4c}$$

Eq. (2) changes under redefinition in eq. (3) as

$$2\mathcal{H} \left(\boldsymbol{U}^2 + \boldsymbol{V}^2 + \hbar^2 \right) = -\mu k^2$$

= $2\mathcal{H} \left(2\boldsymbol{A}'^2 + 2\boldsymbol{L}^2 + \hbar^2 \right)$
= $2\mathcal{H} \left(4\boldsymbol{L}^2 + \hbar^2 \right)$

Where the relation $A'^2 = L^2$ was also used, which can be obtained using eq. (3), eq. (4c), and the fact that A and L are orthogonal. Using eigenvalues of $L^2 = l (l+1) \hbar^2$

$$\mathcal{H} = \frac{-\mu k^2}{2\left(4l\left(l+1\right)+1\right)\hbar^2} = \frac{-\mu k^2}{2\hbar^2 \left(2l+1\right)^2} = \frac{-\mu k^2}{2\hbar^2 n^2}$$

with $k = -\frac{e^2}{4\pi\epsilon_0}$ these are the usual energy eigenvalues for the Hydrogen atom.

Hydrogen atom also satisfies SO(4,2) symmetry, which forms its spectrum generating group[6]. The operators which form the Lie-algebra of this group are:

$$\boldsymbol{L} = \boldsymbol{q} \times \boldsymbol{p} \tag{5a}$$

$$\boldsymbol{A} = \frac{\hat{\boldsymbol{p}} \times \hat{\boldsymbol{L}} - \hat{\boldsymbol{L}} \times \hat{\boldsymbol{p}}}{2\mu} - k\frac{\hat{\boldsymbol{q}}}{q}$$
(5b)

$$\boldsymbol{B} = \frac{\hat{\boldsymbol{p}} \times \hat{\boldsymbol{L}} - \hat{\boldsymbol{L}} \times \hat{\boldsymbol{p}}}{2\mu} + k\frac{\hat{\boldsymbol{q}}}{q}$$
(5c)

$$B_4 = \boldsymbol{q} \cdot \boldsymbol{p} + \frac{3}{2} \frac{\hbar}{i} \tag{5d}$$

$$\Gamma = q\mathbf{p} \tag{5e}$$

$$\Gamma_4 = \frac{1}{2} \left(q \boldsymbol{p} \cdot \boldsymbol{p} - q \right) \tag{5f}$$

$$\Gamma_5 = \frac{1}{2} \left(q \boldsymbol{p} \cdot \boldsymbol{p} + q \right) \tag{5g}$$

SO(4, 2) group includes both SO(4) and SO(3, 1) which are both the bound states and scattering states of 2 particles, regardless of whether they have attractive or repulsive interactions.

Hamiltonian of 3-body problem with added dynamical variables and constraints

The Hamiltonian of 3-body system in an inertial frame is complicated to solve because of potentials depending on inter-particle distances. The Lagrangian in the frame of reference of one of the particles is given by[15]:

$$\mathcal{L} = \frac{1}{2M} \left(m_1 \left(m_3 + m_2 \right) \dot{\boldsymbol{q}}_{13}^2 + m_2 \left(m_1 + m_3 \right) \dot{\boldsymbol{q}}_{23}^2 - 2m_1 m_2 \dot{\boldsymbol{q}}_{13} \cdot \dot{\boldsymbol{q}}_{23} \right) - V \left(q_{13}, q_{23}, |\boldsymbol{q}_{13} - \boldsymbol{q}_{23}| \right)$$

Substituting

$$q_{12} = q_{13} - q_{23}$$

$$\mathcal{L} = \frac{1}{2M} \left(m_1 m_3 \dot{\boldsymbol{q}}_{13}^2 + m_2 m_3 \dot{\boldsymbol{q}}_{23}^2 + m_1 m_2 \dot{\boldsymbol{q}}_{12}^2 \right) - V \left(q_{13}, q_{23}, q_{12} \right)$$

Change of variable $m{q}_{12}
ightarrow -m{q}_{21}$ keeps Lagrangian invariant. Constraint becomes:

$$q_{12} + q_{23} + q_{31} = 0 (6)$$

and rearrangement reduces the Lagrangian to:

$$\mathcal{L} = \frac{1}{2} \left(\mu_{12} \dot{\boldsymbol{q}}_{12}^2 + \mu_{23} \dot{\boldsymbol{q}}_{23}^2 + \mu_{31} \dot{\boldsymbol{q}}_{31}^2 \right) - V \left(q_{12}, q_{23}, q_{31} \right)$$

$$\mu_{ij} = \frac{m_i m_j}{M}$$

A more generalized derivation for many-bodies is shown in appendix: Many-body Lagrangian. For convenience from this point forward, index (i, Mod (i + 1, 3)) will be referred to as i. Taking Legendre transformation and putting in the potential for inverse-square law force, we get following Hamiltonian:

$$\mathcal{H} = \frac{p_1^2}{2\mu_1} + \frac{p_2^2}{2\mu_2} + \frac{p_3^2}{2\mu_3} - \frac{k_1}{q_1} - \frac{k_2}{q_2} - \frac{k_3}{q_3}$$
(7)

$$q_1 + q_2 + q_3 = 0 (8)$$

Constraints for momenta were found using the definition

$$\dot{\boldsymbol{q}}_i = \frac{\partial H}{\partial \boldsymbol{p}_i} \tag{9}$$

which are given by:

$$\sum_{i} \frac{p_i}{\mu_i} = 0 \tag{10}$$

The eq. (7) with constraints (8) and (10) are the equations needed to be solved to get a solution of the 3-body problem.

This constrained Hamiltonian system can be solved by the method devised by Dirac[20]. The Dirac consistency conditions give an altered Hamiltonian which is shown in appendix: Dirac consistency conditions for 3-body Hamiltonian. We can still continue to use the same Hamiltonian from eq. (7) if we project the 3×3 dimensional space to constrained subspace using projection operators. The detailed reasons are given in appendix mentioned above. It can be seen that the eq. (7) looks like a Hamiltonian for 3 Hydrogen-like systems. This observation is important for constructing the symmetry groups of this system in next section.

A noteworthy feature of this formulation is that the Hamiltonian in eq. (7) separates the interactions of the three particles in the three separate terms which involve two particle interactions only. So every force that is present in the system is accounted for, and the constraint "forces" would vanish if system is restricted to only the subspace allowed by the constraints, which can be done using projection operators. This method also eliminates the problem of probability current "leaking" out of constrained subspace because of quantum mechanical tunneling. This is accomplished by the use of projection into the constrained subspace[1].

Finding generators for symmetries in 3-body problem

The generators for SO(4,2) can be written in matrix representation using coordinate transformation from Bars[19] has regular representation in natural (Hartree) units:

$$\operatorname{Reg}(g) = \begin{pmatrix} 0^{(6)} & R^{(6)} \\ -R^{(6)} & 0^{(6)} \end{pmatrix}$$
(11)
$$R^{(6)} = \begin{pmatrix} 0 & \theta_{15} & \theta_{14} & \theta_{13} & \theta_{12} & \theta_{11} \\ -\theta_{15} & 0 & \theta_{10} & \theta_{9} & \theta_{8} & \theta_{7} \\ -\theta_{14} & -\theta_{10} & 0 & \theta_{6} & \theta_{5} & \theta_{4} \\ -\theta_{13} & -\theta_{9} & -\theta_{6} & 0 & \theta_{3} & \theta_{2} \\ -\theta_{12} & -\theta_{8} & -\theta_{5} & -\theta_{3} & 0 & \theta_{1} \\ -\theta_{11} & -\theta_{7} & -\theta_{4} & -\theta_{2} & -\theta_{1} & 0 \end{pmatrix}$$

Where $0^{(6)}$ is a 6×6 zero matrix. The coordinates, which can be represented by a 12 dimensional vector V are:

$$V = (\boldsymbol{X}, \boldsymbol{P}) \tag{12}$$

$$\boldsymbol{X} = (0, \boldsymbol{r} \cdot \boldsymbol{p}, r, r_1, r_2, r_3) \tag{13}$$

$$\boldsymbol{P} = \left(1, \frac{1}{2}\boldsymbol{p} \cdot \boldsymbol{p}, 0, p_1, p_2, p_3\right)$$
(14)

The generators found in eq. (11) were found using[19]:

$$L_{MN} = X_M P_N - X_N P_M \tag{15}$$

Most general transformation for the SO(4, 2) group which spans the entire phase space of Hydrogen-like Hamiltonian is given by:

$$G = g_0 \prod_{i=1}^{15} \exp(\theta_i g_0 g_i)$$
 (16)

where g_i are the generators corresponding to θ_i of SO(4, 2) from regular representation in eq. (11). g_0 is a matrix as follows:

$$g_0 = \begin{pmatrix} 0^{(6)} & -I^{(6)} \\ I^{(6)} & 0^{(6)} \end{pmatrix}$$

where $I^{(6)}$ is a 6×6 identity matrix and $0^{(6)}$ is a 6×6 zero matrix, which is required because the group is a Symplectic group, and its commutation relations, which have the form (for example):

$$g_3 = g_1 g_0 g_2 - g_2 g_0 g_1$$

The generators can be recovered by:

$$g_i = \lim_{\theta_j \to 0} \frac{\partial}{\partial \theta_i} G$$

The generators for 3 body problem without the constrains are obtained by taking products as shown in (15) of vectors which have coordinates/momenta of each subsystems listed in eq. (12). The symmetry group without constraints is $SO(4 \times 3, 2 \times 3)$. The generator matrix is as follows:

$$g = \begin{pmatrix} M_{11} & M_{12} & M_{13} \\ M_{21} & M_{22} & M_{23} \\ M_{31} & M_{32} & M_{33} \end{pmatrix}$$
(17)

Each matrix $M = M^{(ij)}$ is a transformation between a vector $V_n^{(j)}$ and $V_m^{(i)}$. Here m and n are the sub-system number and (i, j) are collective index of parameter of generators of one SO(4, 2) like element which corresponds to an intra-subsystem transformation for m = n or inter-subsystem transformation or mixing for $m \neq n$.

The most general transformation of this kind is given by:

$$G_{\otimes 3} = g_{3\otimes 0} \prod_{i,j,m,n} \exp\left(\theta_{mn}^{(ij)} g_{3\otimes 0} g_{mn}^{(ij)}\right)$$

The constraints in the form of eq. (6) and eq. (10) can be applied to this system using projection operators, found using method outlined in appendix, Projection operators, and constrained sub-groups:

$$P_{\perp} = \begin{pmatrix} \frac{2}{3} & -\frac{1}{3} & -\frac{1}{3} \\ -\frac{1}{3} & \frac{2}{3} & -\frac{1}{3} \\ -\frac{1}{3} & -\frac{1}{3} & \frac{2}{3} \end{pmatrix}$$
(18)

This projection operator if applied to a vector v = (x, y, z) gives another vector v' = (x', y', z') = Pv which satisfies the condition x' + y' + z' = 0. This projection operator is the new identity operator in the required group, it satisfies all of the properties required for an identity element of the new group.

The projection operator corresponding to P_{\perp} for the 3 particle system is $P_{3\otimes \perp}$ which is obtained by taking outerproduct of

$$P_{3\otimes \perp} = I_3 \otimes ((J_1 \otimes I_2) \otimes I_3 + (J_2 \otimes I_2) \otimes P_{\perp})$$
⁽¹⁹⁾

where I_n is identity matrix of rank n and

$$J_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$
$$J_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

The first term in 19 is for the part of vector in eq.12 which are unaffected by the constraints. The generator is given by:

$$G_{P\otimes 3} \doteq P_{3\otimes \perp}G_{\otimes 3}$$

$$g_{mn}^{(ij)} = \lim_{\Theta \to 0} \frac{\partial}{\partial \theta_{mn}^{(ij)}} G_{P \otimes 3}$$

which is a group that is homomorphic to the required spectrum generating group for 3 bodies and orthogonality conditions

$$g_{3\otimes 0}g_{mn}^{(ij)} = -\left(g_{mn}^{(ij)}g_{3\otimes 0}\right)^{\dagger}$$

for each block of the block-diagonal matrix († is transpose).

$$\sum_{i} \gamma_{i,j} = 0 \quad \forall j$$
$$\sum_{j} \gamma_{i,j} = 0 \quad \forall i$$
$$\gamma_{i,j} = -\gamma_{j,i}$$

All these constraints also satisfy the Dirac consistency conditions because the extra terms in Hamiltonian due to Dirac conditions go to zero after applying the projection operator in eq. (18).

Energy spectrum of Helium atom

For calculating energy spectra of Helium like atoms and ions at first it was assumed that only spherically symmetric electron distributions, typically occurring in the ground states are important. The interconnections were ignored by setting $M_{ij} = 0$ for $i \neq j$ was the first step in approximations, which is not required but was made to verify whether the energy levels match. This approximation is made provisionally till the full results are obtained.

The whole equation was non-dimentinalized first. Momentum constraint can be written as

$$\sum_{i} \boldsymbol{p}_{\bigtriangleup i} \quad = \quad \sum_{i} \frac{\boldsymbol{p}_{i}}{\mu_{i}} = 0$$

This change can be incorporated into the Hamiltonian by defining,

$$\mu_{\triangle i} = \mu_i^{-1}$$

which satisfies the Hamiltonian

$$\mathcal{H} = \sum_{i} rac{oldsymbol{p}_{\bigtriangleup i}^2}{2\mu_{\bigtriangleup i}} - rac{k_i}{q_i}$$

For non-dimensionalizing the equation, put the following Hartree (Natural) units:

$$\hbar = 1$$
 $\mu_{\triangle i} = 1$ $k_i = 1$

Bohr radii

$$a_{\triangle i} = \frac{\hbar^2 / \mu_{\triangle i}}{k_i}$$

and Hartree energies

$$E_{h,i} = \frac{k_i}{a_{\triangle i}}$$

The transformations used for the position and momentum variables are:

$$egin{array}{rcl} m{q}_i &=& a_{ riangle i}m{q}_i' \ m{p}_{ riangle i} &=& \sqrt{rac{\mu_{ riangle i}k_i}{a_{ riangle i}}}m{p}_i' \end{array}$$

where primed quantities are the non-dimensional quantities used in sec. Finding generators for symmetries in 3-body problem. The transformations for angular momenta and LRL vectors are:

$$L'_{i} = \frac{L_{\triangle i}}{\sqrt{\mu_{\triangle i}k_{i}a_{\triangle i}}} = \frac{L_{\triangle i}}{\hbar}$$
(21a)

$$A'_{i} = \frac{A_{\Delta i}}{\mu_{\Delta i}^{2}k_{i}}$$
(21b)

The bi-linear Casimir invariant operator for Hamiltonian invariant subgroup of the spectrum generating group of 3 body problem is

$$C = \left(\sum_{i} L'_{i}\right)^{2} - 3\sum_{i} L'^{2}_{i} - 3\sum_{i} A'^{2}_{i}$$
$$= \frac{1}{\hbar^{2}} \left(\sum_{i} L_{\Delta i}\right)^{2} - \frac{3}{\hbar^{2}} \sum_{i} L^{2}_{\Delta i} - \sum_{i} 3\frac{A^{2}_{\Delta i}}{\mu^{4}_{\Delta i} k^{2}_{i}}$$

Now replacing variables with \triangle subscripts by regular variables using eq. (21):

$$C = \frac{1}{\hbar^2} \left(\sum_{i} \frac{L_i}{\mu_i} \right)^2 - \frac{3}{\hbar^2} \sum_{i} \frac{L_i^2}{\mu_i^2} - \sum_{i} 3 \frac{A_i^2}{k_i^2}$$

From this equation one derived a formula:

$$En = \frac{3}{-10(a+1)a + (\lambda+1)\lambda + 5} \sum_{i=1}^{3} -\frac{k_i^2 \mu_i}{\hbar^2}$$
(22)

For $\lambda = 1/2$ and a = -1/2 gives the ground state energy. The ratio of calculated to experimental value is 1.00171. For $\lambda = 3/2$ and a = -1/2 gives the first excited state energy. The ratio of calculated to experimental value is 1.00011.

Conclusion and discussion

Further calculations using the full generators as described in sec. Finding generators for symmetries in 3-body problem will lead to more accurate results. One other way the accuracy can be improved is to use Dirac like equation for multi-electron system and figuring out the symmetries. Wavefunctions can be found as described in Gilmore [6].

Appendix

Here some of the mathematical methods that were used in this work are described in details.

Many-body Lagrangian

Starting from a Lagrangian

$$L = \sum_{i} \frac{1}{2} m_{i} v_{i}^{2} - \sum_{i < j} \frac{k_{ij}}{q_{ij}}$$
(23)

and moving to frame of reference of particle 1,

$$egin{array}{rcl} m{q}_i'&=&m{q}_i-m{q}_1\ m{v}_i'&=&m{v}_i-m{v}_1 \end{array}$$

and eliminating v_1 using total value of momentum,

$$\boldsymbol{v}_1 = \frac{1}{M} \left(\sum_j m_j \boldsymbol{v}_j - \sum_{j \neq 1} m_j \boldsymbol{v}'_j \right)$$

The Lagrangian in eq. 23 reduces to

$$\mathcal{L} = \frac{1}{2M} \left(\sum_{j} m_{j} \boldsymbol{v}_{j} \right)^{2} + \frac{1}{2} \sum_{i < j} \mu_{ij} v_{ij}^{2} - \sum_{i < j} \frac{k_{ij}}{q_{ij}}$$

$$\mu_{ij} = \frac{m_{i} m_{j}}{M}$$
(24)

The first term in this equation is the Kinetic energy of center of mass which is always conserved. Rest of the terms are kinetic energies associated with the relative velocities of pairs of bodies. There are $\frac{1}{2}N(N-1)$ such 3 dimensional vector variables for positions and momenta each, which is not the N pairs of vector variables we started with. These variables are related with each other using $\frac{1}{2}N(N-3) + 1$ independent triangular constraints

$$\boldsymbol{r}_{ij} + \boldsymbol{r}_{jk} + \boldsymbol{r}_{ki} = 0 \qquad (i \neq j \neq k) < N \tag{25}$$

where r can be both q and v.

Dirac consistency conditions for 3-body Hamiltonian

 $\boldsymbol{r}_{ij} = \boldsymbol{r}_i - \boldsymbol{r}_j$

The procedure for finding the Dirac consistency conditions[20] for arbitrary problem under constrains is outlined by Fung[14]. Using that procedure following Dirac Hamiltonian was found:

$$\mathcal{H}^{(D)} = \sum_{i} \left(\frac{1}{2\mu_{i}} \boldsymbol{p}_{i}^{2} - \frac{k_{i}}{q_{i}} \right) - \left(\sum_{i \in \Delta} \frac{1}{\mu_{i}} \right)^{-1} \left(2 \sum_{i \in \Delta} \left(\frac{1}{\mu_{i}} k_{i} \frac{\boldsymbol{q}_{i}}{q_{i}^{3}} \right) \cdot \sum_{j \in \Delta} \boldsymbol{q}_{j} + \left(\sum_{i \in \Delta} \frac{1}{\mu_{i}} \boldsymbol{p}_{i} \right)^{2} \right)$$
(26)

The constraints in eq. (8) and eq. (10) are the primary and secondary constraints and there are no other constraints in this equation. The additional terms that appear in eq. (26) are due to the "Forces" that make the system satisfy the constraints. This same equation is valid for many-body Hamiltonian where the indices $i \in \Delta$ are for the vectors forming the polygon constraints which are equivalent to triangle constraints in eq. (25).

When a projection operator is used to restrict the phase space to the constrained subspace, the extra terms in the Hamiltonian vanish, reducing the Hamiltonian back to the original Hamiltonian. This also applies to the Dirac Brackets[20] which are used to find the new commutator relations for this Hamiltonian if new projected quantities are used.

$$\dot{f} = [f, \mathcal{H}]_D$$

$$P\dot{f} = \left[Pf, \mathcal{H}^{(D)}\right]$$

$$= \left[Pf, P\mathcal{H}^{(D)}\right]$$

$$\dot{f}_P = [f_P, \mathcal{H}]$$

For many-body problems there are additional primary constraints, which can be obtained using linear combinations of constraint equations involving only 3 vectors, like eq. (8) and eq. (10). These primary constraints also produce secondary constraints similar to what was obtained in eq. (26). These constraints are also eliminated using the projection method.

Projection operators, and constrained sub-groups

A symmetry group can be constrained to a subgroup which satisfies certain constraints using a projection operator, which projects the vector space to sub-space which satisfies the constrains. It can be illustrated using the example below.

A rotation group in 3 dimensions, SO (3), can be constructed using Yaw-Pitch-Roll matrices as follows:

$$R(\theta_1, \theta_2, \theta_3) = R_1(\theta_1) R_2(\theta_2) R_3(\theta_3)$$
(27)

Here $R_i(\theta_i)$ are rotation matrices which can be obtained as $R_i(\theta_i) = \exp(J_i\theta_i)$, where J_i are 3×3 Matrix representation of Lie algebra. These are the most general elements of the rotation group SO(3).

$$J_i = \lim_{\theta_i \to 0} \frac{\partial}{\partial \theta_i} \left(\exp\left(J_i \theta_i\right) \right)$$

For (27), this equation will reduce to

$$J_{i} = \lim_{\theta_{j} \to 0} \frac{\partial}{\partial \theta_{i}} R\left(\theta_{1}, \theta_{2}, \theta_{3}\right) \qquad \forall \left(i, j\right) \in \{1, 2, 3\}$$

$$(28)$$

Suppose this is to be constrained to a plane

$$\boldsymbol{v}\cdot\boldsymbol{r} = \boldsymbol{0} \tag{29}$$

we can construct a projection operator which will make r satisfy the constraint eq. (29) by following procedure: first rotate the vector v to axis 3 using

$$R_{\boldsymbol{u}} = \exp\left(-J_{\hat{\boldsymbol{u}}}\theta_{\boldsymbol{u}}\right)$$
(30)
$$\boldsymbol{u} = \boldsymbol{v} \times \hat{\boldsymbol{e}}_{3}$$

$$\cos\left(\theta_{\boldsymbol{u}}\right) = \frac{\boldsymbol{v} \cdot \hat{\boldsymbol{e}}_{3}}{\boldsymbol{v}}$$

Then project the axis 3 to have zero values using P' matrix

$$P' = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Then rotate to the original axis system using R_{μ}^{\dagger} . The entire operation can be summed up as

$$P = R_{\boldsymbol{u}}^{\dagger} P' R_{\boldsymbol{u}}$$
(31)
$$r' = P \boldsymbol{r}$$

The coordinates r' defined using the projection will satisfy $v \cdot r' = 0$ for arbitrary r. Applying this projection operator from eq. (31) on most general group element from eq. (27) will produce a group homomorphic to the constrained sub-group required. The Lie-algebra elements of this group can now be found using eq. (28). The matrices obtained using this method are in general not orthogonal, but if their linear combinations are considered, a subspace of them will be found to be orthogonal. This elements of subspace will be

$$\begin{aligned} J_{\hat{\boldsymbol{v}}} &= \hat{\boldsymbol{v}} \cdot \boldsymbol{J} \\ \boldsymbol{J} &= (J_1, J_2, J_3) \end{aligned}$$

and the new rotation matrix

$$R_{\boldsymbol{v}} = \exp\left(J_{\hat{\boldsymbol{v}}}\theta\right)$$

is the required element of the constrained sub-group of SO(3).

In order to construct similar projection operators for more than 3 dimensions, which are required for solving manybody problems, one need to find the linearly independent triplets of vectors which can be expressed in zero sum (difference) form. These sums and the orthogonality condition can be used to find the rotation matrices equivalent to the one in eq. (30), which after projection like in eq. (31) would give the required projection operators for polyhedron constraints.

Acknowledgments

These ideas have been developing in author's mind since November 2013. Author thanks his teachers from Indian Institute of Technology Kanpur, Tata Institute of Fundamental Research, and Case Western Reserve University, for teaching the essential concepts of physics.

References

- [1] Marle, C. M. Reduction of constrained mechanical systems and stability of relative equilibria. *Communications in Mathematical Physics* **174**, 295–318 (1995).
- [2] Fock, V. A. Hydrogen Atom and Non-Euclidean Geometry. V.A. Fock Selected Works 369-380 (2004).
- [3] Pauli, W. Über das Wasserstoffspektrum vom Standpunkt der neuen Quantenmechanik. Zeitschrift für Physik (1926).
- [4] Myers, S. B. & Steenrod, N. E. The Group of Isometries of a Riemannian Manifold. *The Annals of Mathematics* 40, 400 (1939).
- [5] Froggatt, C. D., Nielsen, H. B. & FOek, V. V. Zur Theorie des Wasserstoffatoms. Origin of Symmetries 224–233 (1991).
- [6] Gilmore, R. Lie Groups, Physics, and Geometry (2008).
- [7] Rogers, H. H. Symmetry transformations of the classical Kepler problem. *Journal of Mathematical Physics* 14, 1125–1129 (1973).
- [8] Moser, J. Regularization of kepler's problem and the averaging method on a manifold. *Communications on Pure and Applied Mathematics* **23**, 609–636 (1970).
- [9] Efthimiou, C. & Frye, C. Spherical harmonics in p dimensions. *Spherical Harmonics in P Dimensions* 1–143 (2014). 1205.3548.
- [10] Nambu, Y. Generalized hamiltonian dynamics. Physical Review D 7, 2405–2412 (1973).
- [11] Dirac, P. A. Note on the Interpretation of the Density Matrix in the Many-Electron Problem. *Mathematical Proceedings of the Cambridge Philosophical Society* **27**, 240–243 (1931).
- [12] Kaabi, A. On the numerical solution of generalized Sylvester matrix equations. Bulletin of the Iranian Mathematical Society 40, 101–113 (2014).
- [13] Woit, P. Quantum theory, groups and representations: An introduction (2017).
- [14] Fung, M. K. Dirac bracket for pedestrians. Chinese Journal of Physics 52, 1643–1651 (2014).
- [15] Poincaré, H. Œuvres d'Henri Poincaré, Volume 7 (Gauthier-Villars, Paris, 1952).
- [16] Castellana, M. Symmetry reduction of the three-body problem based on Euler angles. *Journal of Mathematical Physics* **59** (2018).
- [17] Warner, F. W. Foundations of Differentiable Manifolds and Lie Groups. (Springer, 1983).
- [18] Finkelstein, R. J. Symmetry group of the hydrogen atom. Journal of Mathematical Physics 8, 443–449 (1967).
- [19] Bars, I. Conformal symmetry and duality between free particle, h atom, and harmonic oscillator. *Phys. Rev. D* 58, 066006 (1998).
- [20] Dirac, P. A. M. Generalized hamiltonian dynamics. *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* **246**, 326–332 (1958).
- [21] Musielak, Z. E. & Quarles, B. The three-body problem. *Reports on Progress in Physics* 77, 065901 (2014).
- [22] Marchal, C. *The three-body problem* (Elsevier, 2012).
- [23] Sansonetti, J. E. & Martin, W. C. Handbook of basic atomic spectroscopic data. Journal of Physical and Chemical Reference Data 34, 1559–2259 (2005).
- [24] Gowers, T., Barrow-Green, J. & Leader, I. *The Princeton Companion to Mathematics* (Princeton University Press, USA, 2008), illustrated edition edn.
- [25] Šuvakov, M. & Dmitrašinović, V. Three classes of newtonian three-body planar periodic orbits. *Phys. Rev. Lett.* 110, 114301 (2013).
- [26] Masili, M., Groote, J. J. D. & Hornos, J. E. Nonadiabatic calculations of the oscillator strengths for the helium atom in the hyperspherical adiabatic approach. *Journal of Physics B: Atomic, Molecular and Optical Physics* 33, 2641 (2000).