# Off-diagonal estimates of partial Bergman kernels on $S^1$ -symmetric Kähler manifolds

Ood Shabtai\*

January 30, 2024

#### Abstract

We establish local asymptotic estimates of partial Bergman kernels on closed,  $S^1$ -symmetric Kähler manifolds. The main result concerns the scaling asymptotics of partial Bergman kernels at generic off-diagonal points in which they are not negligible. The case of the two-dimensional sphere is discussed in detail.

# Contents

1	Introduction	<b>2</b>
	1.1 Main results	3
2	Rotation of the two-dimensional sphere	5
3	The microsupport of partial Bergman kernels	11
	3.1 Definition and basic properties	11
	3.2 Proof of Theorem 1.2	12
4	Fourier theory and partial Bergman kernels	<b>14</b>
	4.1 The Cauchy-Szegö projection on the circle	14
	4.2 Spectral projections and unitary representations of $S^1$	15
	4.3 Proof of Theorems 1.6, 1.7	16
5	Estimates of integrals involving the Bergman kernel	<b>21</b>
6	Concluding remarks	26

\*shabtai@imj-prg.fr

Partially supported by the DFG funded project SFB/TRR 191 (Project-ID 281071066-TRR 191), and the ANR-DFG project QuaSiDy (Project-ID ANR-21-CE40-0016).

## 1 Introduction

Let  $(M, \omega)$  be a connected closed Kähler manifold of complex dimension n. Assume that  $L \to M$  is a holomorphic Hermitian line bundle such that the curvature of its Chern connection equals  $-i\omega$ . The space of holomorphic sections of  $L^{\otimes k}$  is a finite-dimensional complex vector space  $\mathcal{H}_k$ , equipped with a natural inner product  $\langle \cdot, \cdot \rangle$ . The latter is obtained by integrating the fiberwise Hermitian product of  $L^{\otimes k}$  with respect to the Liouville measure  $\mu = \frac{|\omega^{\wedge n}|}{n!}$ . Assume that  $\varphi : S^1 \times M \to M$  is a holomorphic Hamiltonian  $S^1$ -action.

Assume that  $\varphi : S^1 \times M \to M$  is a holomorphic Hamiltonian  $S^1$ -action. Thus, there exists a smooth function  $H : M \to \mathbb{R}$  such that  $\varphi$  is the flow of its Hamiltonian vector field  $X_H$ . The "quantum counterpart" of H is the Hermitian operator  $\hat{H}_k \in \text{End}(\mathcal{H}_k)$  specified by<sup>1</sup>

$$\hat{H}_k = \mathcal{M}_H - \frac{i}{k} \nabla_{X_H}^{\otimes k},$$

where  $\nabla : C^{\infty}(M, L) \to C^{\infty}(M, T^*M \otimes L)$  is the Chern connection, and  $\mathcal{M}_H$  is the operator of multiplication by H.

Fix a regular value  $E \in H(M)$ , and consider the spectral projection

$$\Pi_{k,E} = \mathbb{1}_{[E,\infty)}(H_k).$$

The Schwartz kernel of  $\Pi_{k,E}$ , which we also denote by  $\Pi_{k,E}$  (by an abuse of notation), is termed a partial Bergman kernel<sup>2</sup> ([27, 28, 1, 7, 21, 22, 9, 10]). This terminology references the "full" Bergman kernel, which is the Schwartz kernel of the orthogonal projection  $\Pi_k : L^2(M, L^{\otimes k}) \to \mathcal{H}_k$ . The main goal of the present article is to study the asymptotic behaviour of the partial Bergman kernel  $\Pi_{k,E}$  as  $k \to \infty$ . We focus on the behaviour away from the diagonal  $\Delta_M \subset M \times M$ , since the asymptotic properties of  $\Pi_{k,E}$  on  $\Delta_M$  are already understood quite well. Specifically, it is shown (among other things) in [27, 28] that

$$\Pi_{k,E}(z,z) = \begin{cases} \Pi_k(z,z) + \mathcal{O}(k^{-\infty}) & \text{if } H(z) > E, \\ \Pi_k(z,z)(\frac{1}{2} + \mathcal{O}(k^{-\frac{1}{2}})) & \text{if } H(z) = E, \\ \mathcal{O}(k^{-\infty}) & \text{if } H(z) < E. \end{cases}$$
(1)

In this work, we formulate statements analogous to (1), but for  $\Pi_{k,E}(z,w)$ , where  $z \neq w$ . Notably,  $\Pi_{k,E}$  will be shown to "concentrate" about the set

$$\{(z,z) \mid H(z) \ge E\} \cup \{(z,\varphi(t,z)) \mid H(z) = E, \ t \in S^1\} \subset M \times M.$$

It is instructive to compare this type of behaviour with that of the Bergman kernel  $\Pi_k$  (or more generally, the kernels of Toeplitz operators<sup>3</sup>). The latter "concentrates" on  $\Delta_M$ , and additionally, it admits a full, uniform asymptotic expansion. We refer the reader to Sect. 6 for some further details in this context.

<sup>&</sup>lt;sup>1</sup>Since  $X_H$  generates a holomorphic circle action,  $\hat{H}_k$  preserves holomorphic sections.

<sup>&</sup>lt;sup>2</sup>The projection  $\Pi_{k,E}$  is a spectral projection, hence the partial Bergman kernel  $\Pi_{k,E}$  is sometimes called a *spectral* partial Bergman kernel.

<sup>&</sup>lt;sup>3</sup>That is, operators of the form  $\Pi_k \mathcal{M}_F : \mathcal{H}_k \to \mathcal{H}_k$ , where  $F \in C^{\infty}(M)$ .

The properties of  $\Pi_k$  have been studied thoroughly (e.g., [18, 3, 2, 8, 4, 26]), and have found many applications in various fields of mathematics; they also underlie the results presented here.

Perhaps most notably, Theorem 4 of [27] describes the leading-order scaling asymptotics of  $\Pi_{k,E}(z,z)$  in a  $\frac{C}{\sqrt{k}}$ -neighborhood of  $H^{-1}(E)$ . The main result of the present paper (Theorem 1.6) addresses the leading-order scaling asymptotics of  $\Pi_{k,E}$  away from the diagonal.

Finally, we refer the reader to [27] for a concise review of various properties of  $S^1$ -symmetric Kähler manifolds which are relevant in our context. There is some overlap between the present paper and [27]; however, to the best of our knowledge, the off-diagonal asymptotic behaviour of partial Bergman kernels corresponding to spectral projections of quantum observables has yet to be described in the research literature. A treatment of the on-diagonal asymptotic properties of partial Bergman kernels of the same type considered here, but without an assumption of  $S^1$ -symmetry, may be found in [28].

#### 1.1 Main results

The local behaviour of  $\Pi_{k,E}$  away from  $H^{-1}(E) \times H^{-1}(E)$  is straightforward to derive (up to an error of order  $\mathcal{O}(k^{-\infty})$ ) using existing tools from the theory of Toeplitz operators ([4]), and does not require the assumption that M is equipped with a holomorphic Hamiltonian  $S^1$ -action. In order to formulate the statements concisely, we use the following terminology (see also Sect. 3).

**Definition 1.1** ([4]). We say that a sequence  $A_k \in \text{End}(\mathcal{H}_k)$  is negligible at  $(z, w) \in M \times M$  if there exists a neighborhood  $\mathcal{N} \subset M \times M$  of (z, w) such that

$$\sup_{\mathcal{N}} |A_k| = \mathcal{O}(k^{-\infty}).$$

Here (by an abuse of notation),  $A_k \in C^{\infty} (M \times M, L^{\otimes k} \boxtimes (L^*)^{\otimes k})$  denotes the Schwartz kernel of  $A_k \in \text{End}(\mathcal{H}_k)$ , and its point-wise norm is defined using the Hermitian metric on  $L^{\otimes k} \boxtimes (L^*)^{\otimes k}$  which is induced from that of L.

**Theorem 1.2.** Let  $F \in C^{\infty}(M)$ , and denote  $\hat{F}_k = \prod_k \left( \mathcal{M}_F - \frac{i}{k} \nabla_{X_F}^{\otimes k} \right) \prod_k$ . Fix  $E \in [\min F, \max F]$ , not necessarily a regular value. Let  $z, w \in M$ . There exists a neighborhood  $\mathcal{N} \subset M \times M$  of (z, w) such that the following holds.

- 1. If F(z) > E or F(w) > E, then  $\mathbb{1}_{[E,\infty)}(\hat{F}_k) \Pi_k = \mathbb{1}_{(-\infty,E)}(\hat{F}_k)$  is negligible at (z,w).
- 2. If F(z) < E or F(w) < E, then  $\mathbb{1}_{[E,\infty)}(\hat{F}_k)$  is negligible at (z,w).

**Remark 1.3.** In particular, if  $z \neq w$  and  $z \notin F^{-1}(E)$  or  $w \notin F^{-1}(E)$ , then  $\mathbb{1}_{[E,\infty)}(\hat{F}_k)$  is negligible at (z,w).

**Remark 1.4.** In some cases (e.g., the settings of Lemma 4.6), the negligibility estimates provided by Theorem 1.2 can be improved, using suitable estimates of the Bergman kernel ([19]), to exponential decay estimates (cf. [27], Theorem 3).

Similarly, in some cases (e.g., the settings of Corollary 4.9), exponential decay estimates are valid for Schwartz kernels of orthogonal projections onto single eigenspaces (cf. [27], Theorem 1).

We note that Theorem 1.2 implies that the ranges of spectral projections corresponding to disjoint classical domains are "orthogonal up to a negligible error". More precisely,

**Corollary 1.5.** Let  $F, G \in C^{\infty}(M)$ . Assume that  $E_1 \in (\min F, \max F)$ , that  $E_2 \in (\min G, \max G)$ , and that  $\{F \ge E_1\} \cap \{G \ge E_2\} = \emptyset$ . Then

$$\|\mathbb{1}_{[E_1,\infty)}(\hat{F}_k)\mathbb{1}_{[E_2,\infty)}(\hat{G}_k)\|_{\mathrm{op}} = \mathcal{O}(k^{-\infty}).$$

Consequently, the algebras generated by the pairs  $\mathbb{1}_{[E_1,\infty)}(\hat{F}_k)$ ,  $\mathbb{1}_{[E_2,\infty)}(\hat{G}_k)$  are "asymptotically trivial" (cf. [24]).

The main result of the present article describes the scaling asymptotics of  $\Pi_{k,E}$  near  $z_0, w_0 \in H^{-1}(E), z_0 \neq w_0$ . Throughout, we fix the normalization min H = 0 (note that H is defined up to a constant). There exists  $N \geq 1$  and an open dense subset  $M_N \subset M$  such that for all  $z \in M_N$ , the stabilizer group of z is the finite subgroup  $S_N^1 \subset S^1$  of order N. Accordingly, the spectrum of  $\hat{H}_k$  equals<sup>4</sup>  $H(M) \cap \frac{N}{k}\mathbb{Z}$ . Let  $\{\psi_a\}_{a \in \mathbb{R}}$  denote the gradient flow of H, and  $\varphi_t(z) = \varphi(t, z)$  (where  $t \in S^1 = \mathbb{R}/2\pi\mathbb{Z}$ ).

**Theorem 1.6.** Let  $\mathcal{O}_{z_0} \subset M$  denote the  $S^1$ -orbit of  $z_0 \in H^{-1}(E)$ . Choose a local non-vanishing invariant section  $s_{inv}$  of L in a neighborhood of  $\mathcal{O}_{z_0}$ . Denote

$$\Pi_{k,E}(z,w) = \mathcal{K}_{k,E}(z,w)S_k(z,w),$$

where  $S_k(z,w) = \sigma^{\otimes k}(z) \otimes (\sigma^*(w))^{\otimes k}$ , with  $\sigma = \frac{s_{\text{inv}}}{|s_{\text{inv}}|}$ .

1. If  $w_0 \notin \mathcal{O}_{z_0}$ , then  $\Pi_{k,E}$  is negligible at  $(z_0, w_0)$ .

2. If  $z_0 \in M_N$  and  $t \in S^1$  satisfies  $\varphi_t(z_0) \neq z_0$ , then

$$\left(\frac{2\pi}{k}\right)^{n} \mathcal{K}_{k,E}\left(\psi_{\frac{a}{\sqrt{k}}}(z_{0}), \psi_{\frac{b}{\sqrt{k}}} \circ \varphi_{t}(z_{0})\right) = e^{-\frac{(a^{2}+b^{2})\|X_{H}(z_{0})\|^{2}}{2}} e^{-iN\left\lceil\frac{kE}{N}\right\rceil t} \frac{N\left(1-i\cot\left(\frac{Nt}{2}\right)\right)}{2\|X_{H}(z_{0})\|\sqrt{\pi k}} + C_{k}(a,b,t).$$

Here,

$$C_k(a,b,t) = \frac{C_{k,1}(a,b,t)}{k} + \frac{C_{k,2}(a,b,t)}{k^{\frac{3}{2}}}$$

where  $C_{k,1}(a, b, t)$ ,  $C_{k,2}(a, b, t)$  satisfy that for any bounded  $B \subset \mathbb{R}$  and closed interval  $\mathcal{I} \subset S^1 \setminus S_N^1$  there exist  $c_{B,\mathcal{I},1}, c_{B,\mathcal{I},2} > 0$  such that

$$|C_{k,1}(a,b,t)| \le (|a|+|b|)c_{B,\mathcal{I},1}, \ |C_{k,2}(a,b,t)| \le c_{B,\mathcal{I},2}$$

for all  $a, b \in B, t \in \mathcal{I}$ .

<sup>&</sup>lt;sup>4</sup>The case N > 1 is reduced, in Lemma 4.4, to the case N = 1, which is covered by [12].

Let  $V_k(t) = e^{ikt\hat{H}_k}$  denote the quantum evolution defined by  $\hat{H}_k$ . The proof of Theorem 1.6 is essentially an adaptation of the arguments used in [23]; it relies on basic methods from the theory of Fourier series, applied in the context of the unitary representation  $V_k : S^1 \to \text{End}(\mathcal{H}_k)$ . A key role is played by the *periodic Hilbert transform* ([15]), which is a classical singular integral operator on  $L^2(S^1)$ . We also use an estimate of the Schwartz kernels of projections onto single eigenspaces of  $\hat{H}_k$  (the so-called *equivariant Bergman kernels*). The estimate is obtained by essentially the same method that was used in [27] to describe the on-diagonal scaling asymptotics of equivariant Bergman kernels.

**Theorem 1.7.** Let  $\lambda_k$  be an eigenvalue of  $\hat{H}_k$  such that  $|\lambda_k - E| = \mathcal{O}(k^{-1})$ . Let  $\prod_{k,\lambda_k}^{\text{eq}}$  be the orthogonal projection onto the corresponding eigenspace of  $\hat{H}_k$ . Fix  $z_0, w_0 \in M$ , and denote

$$\Pi_{k,\lambda_k}^{\mathrm{eq}}(z,w) = \mathcal{K}_{k,\lambda_k}^{\mathrm{eq}}(z,w) S_k(z,w),$$

where  $S_k(z, w)$  is as specified in Theorem 1.6.

- 1. If  $H(z_0) \neq E$  or  $H(w_0) \neq E$  or  $w_0 \notin \mathcal{O}_{z_0}$ , then  $\Pi_{k,\lambda_k}^{eq}$  is negligible at  $(z_0, w_0)$ .
- 2. If  $z_0 \in M_N$ , then

$$\left(\frac{2\pi}{k}\right)^{n} \mathcal{K}_{k,\lambda_{k}}^{\mathrm{eq}} \left(\psi_{\frac{a}{\sqrt{k}}}(z_{0}), \psi_{\frac{b}{\sqrt{k}}} \circ \varphi_{t}(z_{0})\right) = e^{-\frac{(a^{2}+b^{2})\|X_{H}(z_{0})\|^{2}}{2}} e^{-ik\lambda_{k}t} \frac{N}{\|X_{H}(z_{0})\|\sqrt{\pi k}} + C_{k}^{\mathrm{eq}}(a,b,t).$$

Here,

$$C_k^{\text{eq}}(a, b, t) = \frac{C_{k,1}^{\text{eq}}(a, b, t)}{k} + \frac{C_{k,2}^{\text{eq}}(a, b, t)}{k^{\frac{3}{2}}}$$

where  $C_{k,1}^{\text{eq}}(a, b, t)$ ,  $C_{k,2}^{\text{eq}}(a, b, t)$  satisfy that for any bounded  $B \subset \mathbb{R}$  and closed interval  $\mathcal{I} \subset S^1$  there exist  $c_{B,\mathcal{I},1}^{\text{eq}}, c_{B,\mathcal{I},2}^{\text{eq}} > 0$  such that

$$|C_{k,1}^{\rm eq}(a,b,t)| \le (|a|+|b|)c_{B,\mathcal{I},1}^{\rm eq}, \ |C_{k,2}^{\rm eq}(a,b,t)| \le c_{B,\mathcal{I},2}^{\rm eq}$$

for all  $a, b \in B, t \in \mathcal{I}$ .

## 2 Rotation of the two-dimensional sphere

In this section, we illustrate the results of Sect. 1.1 in the example of the action of  $S^1$ , by rotations, on the two-dimensional sphere. The latter is identified (via the stereographic projection from the north pole) with the complex projective line  $\mathbb{C}P^1$  and equipped with the Fubini-Study form such that the total area equals  $2\pi$ . For conciseness, we focus on the case N = 1, and a = b = 0. The Hamiltonian vector field of the function

$$H([z]) = \frac{|z_0|^2}{|z_0|^2 + |z_1|^2}, \ z = (z_0, z_1) \in \mathbb{C}^2 \setminus \{0\},\$$

is specified (on  $\{[z_0:z_1] \in \mathbb{C}P^1 \mid z_1 \neq 0\}$ ) by

$$X_H([\zeta:1]) = i \left( \zeta \partial_{\zeta} - \bar{\zeta} \partial_{\bar{\zeta}} \right),$$

and it generates the Hamiltonian flow

$$\varphi_t([z]) = [U_t z], \ U_t = \begin{pmatrix} e^{it} & 0\\ 0 & 1 \end{pmatrix}.$$

Clearly,  $\varphi: S^1 \times \mathbb{C}P^1 \to \mathbb{C}P^1$  is a holomorphic circle action.

**Remark 2.1.** On  $S^2 = \{x \in \mathbb{R}^3 \mid |x| = 1\}$ , it holds that

$$H(x_1, x_2, x_3) = \frac{1}{2}(x_3 + 1),$$

and  $\varphi_t$  is the rotation by angle t about the  $x_3$  axis.

Let  $L = \mathcal{O}(1)$  be the dual of the tautological line bundle on  $\mathbb{C}P^1$ . Then L is a prequantum line bundle, and we let  $\mathcal{H}_k = H^0(\mathbb{C}P^1, L^{\otimes k})$  be the space of holomorphic sections of  $L^{\otimes k}$ . The quantum counterpart of H is given by (5)

$$\hat{H}_k = \prod_k \mathcal{M}_{H-\frac{1}{2k}\Delta H} \prod_k,$$

and in our case,  $\Delta H = -4H + 2$ , which means that

$$\hat{H}_k = \frac{k+2}{k} \Pi_k \mathcal{M}_H \Pi_k - \frac{1}{k} \operatorname{Id}_{\mathcal{H}_k}.$$

As is common, we identify  $\mathcal{H}_k$  with the space of bivariate homogeneous polynomials of degree k, denoted by  $\mathbb{C}_k[z_0, z_1]$ , and equipped with a suitable inner product so that the set  $\{s_{k,l} \mid l = 0, 1, ..., k\}$ , where

$$s_{k,l} = \sqrt{\frac{(k+1)\binom{k}{l}}{2\pi}} z_0^l z_1^{k-l}, \qquad (2)$$

forms an orthonormal basis of  $\mathcal{H}_k$ . In fact, this is an eigenbasis of  $\hat{H}_k$  (cf. [16], Example 5.2.4), with

$$\hat{H}_k(s_{k,l}) = \lambda_{k,l} s_{k,l}, \ \lambda_{k,l} = \frac{l}{k}.$$

In particular,  $s_{inv} = s_{1,0}$  is an invariant section of  $L \to \mathbb{C}P^1$ .

We can estimate the leading order of  $s_{k,l}$  using only basic tools, as follows. Note that H([z]) = E if and only if  $[z] = \left[\sqrt{\frac{E}{1-E}}e^{i\theta}:1\right]$  for some  $\theta \in [0, 2\pi]$ . **Lemma 2.2.** As in Theorems 1.6, 1.7, consider the local normalized section  $\sigma = \frac{s_{\text{inv}}}{|s_{\text{inv}}|}$  on  $\{[z_0:z_1] \in \mathbb{C}P^1 \mid z_1 \neq 0\}$ , and write

$$s_{k,l}([z]) = \kappa_{k,l}([z])\sigma([z])^{\otimes k}.$$

Let  $E \in (0,1)$  and  $\left|\frac{l_k}{k} - E\right| = O(k^{-1})$ . Then, applying Stirling's approximation formula, we see that

$$\kappa_{k,l_k}([z]) = \begin{cases} \mathcal{O}(k^{-\infty}) & \text{if } H([z]) \neq E, \\ \frac{k^{\frac{1}{4}}}{(2\pi)^{\frac{3}{4}}} \frac{e^{il_k\theta}}{(E(1-E))^{\frac{1}{4}}} + \mathcal{O}(k^{-\frac{3}{4}}) & \text{if } [z] = \left[\sqrt{\frac{E}{1-E}}e^{i\theta} : 1\right]. \end{cases}$$

*Proof.* Stirling's approximation formula produces

$$\binom{k}{l_k} = \frac{1}{\sqrt{2\pi}} \sqrt{\frac{k}{l_k(k-l_k)}} \frac{k^k}{l_k^{l_k}(k-l_k)^{k-l_k}} (1 + \mathcal{O}(k^{-1})).$$

Thus, the pointwise norm of  $s_{k,l_k}$  is given by

$$\left| s_{k,l_k} ([\zeta:1]) \right|_{[\zeta:1]}^2 = \frac{k+1}{(2\pi)^{\frac{3}{2}}} \sqrt{\frac{k}{l_k(k-l_k)}} \left( \left( \left(\frac{k}{l_k} - 1\right) |\zeta|^2 \right)^{\frac{l_k}{k}} \frac{1}{1 - \frac{l_k}{k}} \frac{1}{1 + |\zeta|^2} \right)^k (1 + \mathcal{O}(k^{-1})).$$

Now, note that

$$\left(\left(\frac{k}{l_k} - 1\right)|\zeta|^2\right)^{\frac{l_k}{k}} \frac{1}{1 - \frac{l_k}{k}} \frac{1}{1 + |\zeta|^2} = \left(\left(\frac{1}{E} - 1\right)|\zeta|^2\right)^E \frac{1}{1 - E} \frac{1}{1 + |\zeta|^2} + \mathcal{O}(k^{-1}).$$

We readily verify that

$$a(E,\zeta) = \left(\left(\frac{1}{E} - 1\right)|\zeta|^2\right)^E \frac{1}{1 - E} \frac{1}{1 + |\zeta|^2} = 1$$

if and only if  $|\zeta|^2 = \frac{E}{1-E}$ , that is, if and only if  $H([\zeta : 1]) = E$ , and otherwise  $0 \le a(E, \zeta) < 1$ . Thus,

$$\left|\kappa_{k,l_k}([z])\right| = \left|s_{k,l_k}([\zeta:1])\right|_{[\zeta:1]} = \mathcal{O}(k^{-\infty})$$

whenever  $H([z]) \neq E$ .

Assume that  $H([\zeta : 1]) = E$ , so that  $\zeta = \sqrt{\frac{E}{1-E}}e^{i\theta}$  for some  $\theta \in [0, 2\pi]$ . Then

$$\kappa_{k,l_k}([\zeta:1]) = \sqrt{\frac{k+1}{2\pi}} \sqrt{\binom{k}{l_k}} E^{l_k} (1-E)^{k-l_k} e^{il_k\theta},$$

where

$$\binom{k}{l_k} E^{l_k} (1-E)^{k-l_k} = \frac{1}{\sqrt{2\pi}} \sqrt{\frac{k}{l_k(k-l_k)}} \left(\frac{kE}{l_k}\right)^{l_k} \left(\frac{k(1-E)}{k-l_k}\right)^{k-l_k} \left(1+\mathcal{O}(k^{-1})\right),$$

and it is readily verified that

$$\sqrt{\frac{k}{l_k(k-l_k)}} = \frac{1}{\sqrt{kE(1-E)}} + \mathcal{O}(k^{-\frac{3}{2}}).$$

Writing  $\frac{l_k}{k} = E + \frac{c_k}{k}$  with  $c_k = \mathcal{O}(1)$ , we obtain

$$b_k = \left(\frac{k}{l_k}E\right)^{l_k} \left(\frac{k}{k-l_k}(1-E)\right)^{k-l_k} = \left(1-\frac{c_k}{l_k}\right)^{l_k} \left(1+\frac{c_k}{k-l_k}\right)^{k-l_k}$$

If  $c_k = 0$ , then  $b_k = 1$ . If  $c_k \neq 0$ , noting that  $\left| e - \left( 1 + \frac{1}{x} \right)^x \right| = \mathcal{O}(x^{-1})$  as  $x \to \infty$ , we conclude that  $b_k = 1 + \mathcal{O}(k^{-1})$ . Thus,

$$\binom{k}{l_k} E^{l_k} (1-E)^{k-l_k} = \frac{1}{\sqrt{2\pi}} \frac{1}{\sqrt{kE(1-E)}} (1+\mathcal{O}(k^{-1})),$$

which implies that

$$\kappa_{k,l_k}([z]) = \frac{k^{\frac{1}{4}}}{(2\pi)^{\frac{3}{4}}} \frac{e^{il_k\theta}}{(E(1-E))^{\frac{1}{4}}} + \mathcal{O}(k^{-\frac{3}{4}}).$$

The equivariant Bergman kernel corresponding to the eigenvalue  $\frac{l}{k}$  of  $\hat{H}_k$  is specified by

$$\Pi_{k,l}^{\rm eq}([z],[w]) = s_{k,l}([z]) \otimes s_{k,l}^*([w]).$$
(3)

Hence, Lemma 2.2 provides an estimate of  $\Pi_{k,l}^{\text{eq}}$  which can be used to verify Theorem 1.7 (one could similarly verify the cases N > 1 or  $a \neq 0$  or  $b \neq 0$ ).

**Corollary 2.3.** As in Theorem 1.7, let  $S_k([z], [w]) = \sigma([z])^{\otimes k} \otimes (\sigma^*([w]))^{\otimes k}$ , and write

$$\Pi_{k,l_k}^{\text{eq}}([z],[w]) = \mathcal{K}_{k,l_k}^{\text{eq}}([z],[w])S_k([z],[w]).$$

If we substitute the estimate of Lemma 2.2 in (3), then

$$\begin{aligned} & \mathcal{K}_{k,l_{k}}^{\mathrm{eq}}([z],[w]) = \\ \left\{ \begin{array}{l} \mathcal{O}(k^{-\infty}) & \text{if } H([z]) \neq E \text{ or } H([w]) \neq E, \\ \frac{k^{\frac{1}{2}}}{(2\pi)^{\frac{3}{2}}} \frac{e^{-il_{k}t_{0}}}{\sqrt{E(1-E)}} + \mathcal{O}(k^{-\frac{1}{2}}) & \text{if } [z] = \left[\sqrt{\frac{E}{1-E}}e^{i\theta}:1\right], \ [w] = [U_{t_{0}}z]. \end{aligned} \right. \end{aligned}$$

Clearly, the case  $H([z]) \neq E$  or  $H([w]) \neq E$  is in accordance with Theorem 1.7. Otherwise,  $\zeta = \sqrt{\frac{E}{1-E}} e^{i\theta}$  for some  $\theta \in [0, 2\pi]$ , hence

$$||X_H([\zeta:1])||^2 = \frac{2|\zeta|^2}{(1+|\zeta|^2)^2} = 2E(1-E).$$

Thus, Theorem 1.7 produces the estimate

j

$$\mathcal{K}_{k,l_k}^{\text{eq}}([z], [U_{t_0}z]) = \frac{k}{2\pi} \frac{e^{-il_k t_0}}{\sqrt{2E(1-E)}\sqrt{\pi k}} + \mathcal{O}(k^{-\frac{1}{2}}),$$

which, as expected, is identical to the estimate coming from Lemma 2.2.



Figure 1: A plot of  $|\mathcal{K}_{80,0.5}^{eq}([1:1], [\zeta:1])|$ . The circle  $\{|\zeta| = 1\} = \{H = 0.5\}$  is the orbit of [1:1].

The partial Bergman kernel associated with a regular value  $E \in (0, 1)$  of H is specified by (noting the identification  $\mathcal{H}_k \simeq \mathbb{C}_k[z_0, z_1]$ )

$$\Pi_{k,E} = \frac{k+1}{2\pi} \sum_{\frac{l}{k} \ge E} \binom{k}{l} (z_0 \bar{w}_0)^l (z_1 \bar{w}_1)^{k-l}.$$

This partial binomial sum is difficult to estimate directly. Denote

$$\Pi_{k,E}([z], [w]) = \mathcal{K}_{k,E}([z], [w])S_k([z], [w])$$

The following image illustrates the behaviour of  $\mathcal{K}_{k,E}([z], [w])$  (as expected, the image appears to be in accordance with the estimate provided by Theorem 1.6).



Figure 2: A plot of  $|\mathcal{K}_{80,0.5}([1:1],[\zeta:1])|$ . The circle  $\{|\zeta|=1\} = \{H=0.5\}$  is the orbit of [1:1], and  $\zeta = 1$  corresponds to the point ([1:1],[1:1]) on the diagonal  $\Delta_{\mathbb{C}P^1} \subset \mathbb{C}P^1 \times \mathbb{C}P^1$ .

Consider  $[z] \in H^{-1}(E)$ , and denote

$$\operatorname{Er}_{k}(t_{0}, E, [z]) = \left| \mathcal{K}_{k, E}([z], [U_{t_{0}}z]) - \mathcal{K}_{k, E, \operatorname{approx}}([z], [U_{t_{0}}z]) \right|,$$

where

$$\mathcal{K}_{k,E,\mathrm{approx}}([z],[U_{t_0}z]) = \frac{k}{4\pi} \frac{e^{-i\lceil kE\rceil t_0}}{\sqrt{2E(1-E)}\sqrt{\pi k}} \left(1 - i\cot\left(\frac{t_0}{2}\right)\right)$$

is the leading term in the estimate of Theorem 1.6. The next images illustrate the behaviour of the error as k grows.



Figure 3: A plot of  $\operatorname{Er}_k\left(\frac{\pi}{2}, \frac{1}{2}, [1:1]\right)$  as a function of k (top), and (bottom) a plot of log  $\operatorname{Er}_k\left(\frac{\pi}{2}, \frac{1}{2}, [1:1]\right)$  (in blue) and  $-1.5 - 0.5 \log(k)$  (in red) as functions of  $\log(k)$ . The slope of -0.5 is in accordance with the fact that the sub-leading term in the estimate provided by Theorem 1.6 is (in this case) of order  $k^{-0.5}$ .

## 3 The microsupport of partial Bergman kernels

#### **3.1** Definition and basic properties

The notion of microsupport of an admissible sequence of holomorphic sections is specified in [4], as follows. Let  $(s_k)_{k\in\mathbb{N}}$  denote a sequence such that  $s_k \in \mathcal{H}_k$ . We say that  $(s_k)_{k\in\mathbb{N}}$  is admissible if there exists N > 0 such that  $||s_k|| = \mathcal{O}(k^N)$ . An admissible sequence  $(s_k)_{k\in\mathbb{N}}$  is called *negligible at*  $z_0 \in M$  if there exists a neighborhood V of  $z_0$  such that  $\sup_{z\in V} |s_k(z)|_z = \mathcal{O}(k^{-\infty})$ , where  $|\cdot|_z$  is the norm on  $L_z^{\otimes k}$  induced by the Hermitian product.

**Definition 3.1.** The microsupport of an admissible sequence  $(s_k)_{k \in \mathbb{N}}$  is the set

 $MS(s_k) \subset M$  specified by

$$MS(s_k) = M \setminus \{ z \in M \mid (s_k)_{k \in \mathbb{N}} \text{ is negligible at } z \}.$$

The notion of microsupport is also applicable to sequences  $(T_k)_{k\in\mathbb{N}}$  with  $T_k \in \operatorname{End}(\mathcal{H}_k)$ . To this end, equip  $M \times M$  with the symplectic form  $\pi_1^* \omega - \pi_2^* \omega$ , where  $\pi_1, \pi_2$  are the natural projections on the left and right factors. Then  $\operatorname{MS}(T_k) \subset M \times M$  is defined to be the microsupport of the sequence of Schwartz kernels of  $T_k$ , which are holomorphic sections of  $L^{\otimes k} \boxtimes (L^*)^{\otimes k} \to M \times M$ .

**Example 3.2.** ([4]) Assume that  $T_k = \prod_k \mathcal{M}_{G_k} \prod_k$ , where  $G_k \in C^{\infty}(M)$  admits the expansion  $G_k = \sum_{l=0}^{\infty} g_l k^{-l}$  in the  $C^{\infty}$  topology. Then  $\mathrm{MS}(T_k) \subset \Delta_M$ , where  $\Delta_M \subset M \times M$  is the diagonal. Viewed as a subset of M,

$$MS(T_k) = \bigcup_{l \ge 0} \operatorname{supp}(g_l).$$
(4)

#### 3.2 Proof of Theorem 1.2

The operator  $\hat{F}_k$  admits the representation ([25], [16], Proposition 8.1.3)

$$\hat{F}_k = \Pi_k \mathcal{M}_{F - \frac{1}{2k}\Delta F} \Pi_k,\tag{5}$$

where  $\Delta$  is the Laplacian defined by the Kähler metric of M.

**Lemma 3.3.** Let  $\varepsilon > 0$ . Let  $\psi_{\varepsilon} : \mathbb{R} \to [0,1]$  be a smooth function such that  $\psi_{\varepsilon}(t) = 0$  for all  $t \leq E - \varepsilon$ ,  $\psi_{\varepsilon}(t) = 1$  for all  $t \geq E$ . Write  $\hat{\Psi}_{k,\varepsilon} = \psi_{\varepsilon}(\hat{F}_k)$ . Then  $\mathrm{MS}(\hat{\Psi}_{k,\varepsilon}) \subset \{F \geq E - \varepsilon\}$ .

*Proof.* There exists ([4], Proposition 12)  $\Psi_{k,\varepsilon} \in C^{\infty}(M)$ , admitting an expansion

$$\Psi_{k,\varepsilon} = \sum_{l=0}^{\infty} k^{-l} \psi_{l,\varepsilon}$$

in the  $C^{\infty}$  topology, such that  $\psi_{\varepsilon}(\hat{F}_k) = \prod_k \mathcal{M}_{\Psi_{k,\varepsilon}} \prod_k + \mathcal{O}(k^{-\infty})$ . If  $F(z) < E - \varepsilon$ , then  $\psi_{\varepsilon}$  vanishes in a neighborhood of F(z), hence (using [4], p27) there exists a neighborhood V of z such that  $\psi_{l,\varepsilon}|_V \equiv 0$  for all  $l \ge 0$ . Thus, in light of (4),  $z \notin MS(\psi_{\varepsilon}(\hat{F}_k))$ .

The first item of Theorem 1.2 now readily follows.

**Corollary 3.4.** The microsupport of  $\mathbb{1}_{[E,\infty)}(\hat{F}_k)$  satisfies

$$\mathrm{MS}(\mathbb{1}_{[E,\infty)}(\hat{F}_k)) \subset \{F \ge E\} \times \{F \ge E\}.$$

The same holds for  $\mathbb{1}_{(E,\infty)}(\hat{F}_k)$ .

*Proof.* We prove for  $A_k = \mathbb{1}_{[E,\infty)}(\hat{F}_k)$ , but the proof for  $\mathbb{1}_{(E,\infty)}(\hat{F}_k)$  is identical. Let  $\psi_{\varepsilon}$  be as in Lemma 3.3, and write  $\hat{\Psi}_{k,\varepsilon} = \psi_{\varepsilon}(\hat{F}_k)$ . Then

$$A_k = \bar{\Psi}_{k,\varepsilon} A_k = A_k \bar{\Psi}_{k,\varepsilon}$$

hence (using [4], p24)

$$\begin{split} \mathrm{MS}(A_k) &= \mathrm{MS}(\hat{\Psi}_{k,\varepsilon}A_k) \\ &\subset \{(z_1, z_3) \mid \exists z_2 \in M \text{ such that } (z_1, z_2) \in \mathrm{MS}(\hat{\Psi}_{k,\varepsilon}) \text{ and } (z_2, z_3) \in \mathrm{MS}(A_k)\} \\ &\subset \mathrm{MS}(\hat{\Psi}_{k,\varepsilon}) \times M, \end{split}$$

where by a slight abuse of notation, in the second line  $MS(\hat{\Psi}_{k,\varepsilon})$  is viewed as a subset of  $M \times M$ , and in the third line it is viewed as a subset of M. Repeating this argument,

$$\mathrm{MS}(A_k) = \mathrm{MS}(A_k \hat{\Psi}_{k,\varepsilon}) \subset M \times \mathrm{MS}(\hat{\Psi}_{k,\varepsilon}),$$

therefore (noting Lemma 3.3)

$$MS(A_k) \subset \left(MS(\hat{\Psi}_{k,\varepsilon}) \times M\right) \cap \left(M \times MS(\hat{\Psi}_{k,\varepsilon})\right)$$
  
= MS( $\hat{\Psi}_{k,\varepsilon}$ ) × MS( $\hat{\Psi}_{k,\varepsilon}$ ) ⊂ {F ≥ E -  $\varepsilon$ } × {F ≥ E -  $\varepsilon$ }.

Since  $\varepsilon > 0$  is arbitrary, we obtain the required.

The second item of Theorem 1.2 immediately follows from Corollary 3.4.

**Corollary 3.5.** Replacing F with G = -F and E with -E, if G(z) < -E or G(w) < -E then there exists a neighborhood  $\mathcal{N} \subset M \times M$  of (z, w) such that

$$\sup_{\mathcal{N}} \left| \mathbb{1}_{(-E,\infty)}(\hat{G}_k) \right| = \mathcal{O}(k^{-\infty}),$$

where

$$\mathbb{1}_{(-E,\infty)}(\hat{G}_k) = \Pi_k - \mathbb{1}_{(-\infty,-E]}(\hat{G}_k) = \Pi_k - \mathbb{1}_{[E,\infty)}(\hat{F}_k),$$

that is,  $\sup_{\mathcal{N}} \left| \Pi_k - \mathbb{1}_{[E,\infty)}(\hat{F}_k) \right| = \mathcal{O}(k^{-\infty}).$ 

Finally,

Lemma 3.6. Corollary 3.4 implies Corollary 1.5.

*Proof.* We keep the notations used in the formulation of Corollary 1.5, and also write

$$\Pi_{k,1} = \mathbb{1}_{[E_1,\infty)}(F_k), \ \Pi_{k,2} = \mathbb{1}_{[E_2,\infty)}(G_k).$$

Then using Corollary 3.4,

 $MS(\Pi_{k,1}\Pi_{k,2})$ 

 $\subset \{(z_1, z_3) \mid \exists z_2 \in M \text{ such that } (z_1, z_2) \in \mathrm{MS}(\Pi_{k,1}) \text{ and } (z_2, z_3) \in \mathrm{MS}(\Pi_{k,2}) \} \\ \subset \{(z_1, z_3) \mid \exists z_2 \in M \text{ such that } z_2 \in \{F \ge E_1\} \cap \{G \ge E_2\} \} = \emptyset.$ 

Thus  $\Pi_{k,1}\Pi_{k,2}$  is a negligible sequence, which implies ([4], Remark 5) that  $\|\Pi_{k,1}\Pi_{k,2}\|_{\text{op}} = \mathcal{O}(k^{-\infty}).$ 

The proof of the first part of the first item of Theorem 1.7 is essentially identical to that of Theorem 1.2. Namely,

**Lemma 3.7.** Let  $\lambda_k$  be a sequence of eigenvalues of  $\hat{F}_k$  with  $|\lambda_k - E| = \mathcal{O}(k^{-1})$ , where  $E \in F(M)$ . Let  $\Pi_{k,\lambda_k,F}$  be the orthogonal projection onto the eigenspace associated with  $\lambda_k$ . Then  $\mathrm{MS}(\Pi_{k,\lambda_k,F}) \subset \{F = E\} \times \{F = E\}$ .

*Proof.* Given  $\varepsilon > 0$  we consider a smooth function  $\psi_{\varepsilon} : \mathbb{R} \to [0, 1]$  such that  $\psi_{\varepsilon}(t) = 1$  whenever  $|t - E| < \frac{\varepsilon}{2}$  and  $\psi_{\varepsilon}(t) = 0$  whenever  $|t - E| \ge \varepsilon$ . Write  $\hat{\Psi}_{k,\varepsilon} = \psi_{\varepsilon}(\hat{F}_k)$ , and note that exactly as in Lemma 3.3,

$$\mathrm{MS}(\hat{\Psi}_{k,\varepsilon}) \subset \{ |F - E| \le \varepsilon \} \times \{ |F - E| \le \varepsilon \}.$$

For every sufficiently large k, it holds that  $\hat{\Psi}_{k,\varepsilon}\Pi_{k,\lambda_k,F} = \Pi_{k,\lambda_k,F}\hat{\Psi}_{k,\varepsilon} = \Pi_{k,\lambda_k,F}$ . Thus, exactly as in Corollary 3.4, we obtain that

$$\mathrm{MS}(\Pi_{k,\lambda_k,F}) \subset \mathrm{MS}(\hat{\Psi}_{k,\varepsilon}) \times \mathrm{MS}(\psi_{\varepsilon}(\hat{F}_k)) \subset \{|F-E| \leq \varepsilon\} \times \{|F-E| \leq \varepsilon\}.$$

Since  $\varepsilon$  is arbitrary,  $MS(\Pi_{k,\lambda_k}) \subset \{F = E\} \times \{F = E\}.$ 

## 4 Fourier theory and partial Bergman kernels

#### 4.1 The Cauchy-Szegö projection on the circle

Let  $\mathbb{R}/2\pi\mathbb{Z} = S^1 \subset \mathbb{C}$  denote the unit circle. Let  $\hat{g}(p) = \langle g, e_p \rangle_{L^2(S^1)}$  denote the *p*-th Fourier coefficient of  $g \in L^1(S^1)$ , where  $p \in \mathbb{Z}$  and  $e_p(t) = e^{ipt}$ . The *Cauchy-Szegö projection*  $\Pi_{S^1} : L^2(S^1) \to H^2(S^1)$  is the orthogonal projection on the Hardy space

$$H^{2}(S^{1}) = \left\{ g \in L^{2}(S^{1}) \mid \hat{g}(p) = 0 \,\,\forall p < 0 \right\},$$

and it admits the formula

$$\Pi_{S^1}(g)(t) = \lim_{r \to 1^-} \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{g(t-s)}{1 - re^{is}} ds$$

The periodic Hilbert transform  $\mathcal{H}_{S^1}$ :  $L^2(S^1) \to L^2(S^1)$  is specified by  $\mathcal{H}_{S^1}(e_p) = -i \operatorname{sgn}(p) e_p$ , or equivalently

$$\mathcal{H}_{S^1}(g)(t) = \lim_{\varepsilon \to 0^+} \frac{1}{2\pi} \int_{\varepsilon \le |t-s| \le \pi} g(s) \cot\left(\frac{t-s}{2}\right) ds.$$

**Corollary 4.1.** The Cauchy-Szegö projection acts by  $e_p(t) \mapsto \mathbb{1}_{[0,\infty)}(p)e_p(t)$ , hence it can be expressed in terms of the periodic Hilbert transform, as follows:

$$\Pi_{S^1}(g) = \frac{1}{2} \left( i \mathcal{H}_{S^1}(g) + g + \hat{g}(0) \right).$$

#### 4.2 Spectral projections and unitary representations of $S^1$

Let  $\mathcal{H}$  denote a finite dimensional complex Hilbert space, and write dim  $\mathcal{H} = d$ . Let  $A \in \text{End}(\mathcal{H})$  be a Hermitian operator. Denote

$$U_A(t) = e^{itA},$$

and assume that  $t \mapsto U_A(t)$  is  $2\pi$ -periodic.

Let  $v_1, ..., v_d$  be an orthonormal eigenbasis of A such that  $Av_m = p_m v_m$ , with  $p_1 \leq p_2 \leq ... \leq p_d$ . Then  $p_m \in \mathbb{Z}$ , and we can write

$$A = \sum_{m=1}^d p_m v_m \otimes v_m^*, \ U_A(t) = \sum_{m=1}^d e^{itp_m} v_m \otimes v_m^*,$$

where  $v_m^*$  is the dual vector of  $v_m$ .

**Corollary 4.2.** The spectral projection  $\Pi_A = \mathbb{1}_{[0,\infty)}(A)$  can be written in the following form:

$$\Pi_A = (\Pi_{S^1}(U_A))(0) = \lim_{r \to 1^-} \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{U_A(s)}{1 - re^{is}} ds.$$

In terms of the Hilbert transform,

$$\Pi_A = \frac{1}{2} \left( i \mathcal{H}_{S^1}(U_A)(0) + \mathrm{Id}_{\mathcal{H}} + \widehat{U}_A(0) \right), \tag{6}$$

where

$$\widehat{U}_A(0) = \frac{1}{2\pi} \int_{-\pi}^{\pi} U_A(t) dt$$

and

$$\mathcal{H}_{S^1}(U_A)(0) = \frac{1}{2\pi} \int_0^\pi \left( U_A(-t) - U_A(t) \right) \cot\left(\frac{t}{2}\right) dt.$$

More generally, given  $p, p_0 \in \mathbb{Z}$  and  $g \in L^2(S^1)$ , recall that

$$\widehat{e_{-p_0}g}(p) = \langle e_{-p_0}g, e_p \rangle_{L^2(S^1)} = \langle g, e_{p+p_0} \rangle_{L^2(S^1)} = \widehat{g}(p+p_0)$$

This relation allows us to extend formula (6) to  $\Pi_{A,E} = \mathbb{1}_{[E,\infty)}(A), E \in \mathbb{R}$ . Corollary 4.3. Let  $E \in \mathbb{R}$ . Denote  $U_{A,E} = e_{-\lceil E \rceil} U_A$ . Then

$$\Pi_{A,E} = (\Pi_{S^1}(U_{A,E}))(0) = \frac{1}{2} \left( i\mathcal{H}_{S^1}(U_{A,E})(0) + \mathrm{Id}_{\mathcal{H}} + \widehat{U}_{A,E}(0) \right),$$
(7)

where

$$\widehat{U}_{A,E}(0) = \frac{1}{2\pi} \int_{-\pi}^{\pi} U_{A,E}(t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} U_A(t) e^{-i\lceil E\rceil t} dt$$

and

$$\mathcal{H}_{S^1}(U_{A,E})(0) = \frac{1}{2\pi} \int_0^\pi \left( U_A(-t)e^{i\lceil E\rceil t} - U_A(t)e^{-i\lceil E\rceil t} \right) \cot\left(\frac{t}{2}\right) dt.$$

#### 4.3 Proof of Theorems 1.6, 1.7

The proof of Theorem 1.6 relies on formula (7), which is also applicable to the Schwartz kernels of the operators involved. Thus, we obtain a representation of  $\Pi_{k,E}(z,w)$  in terms of the Bergman kernel and the Schwartz kernel of the quantum propagator defined by  $\hat{H}_k$ . The well-known asymptotic properties of the Bergman kernel, with the help of the stationary phase lemma, produce the desired estimate. Notably, we can assume without loss of generality that the circle action is effective (Lemma 4.4), and then the third term in (7) is the equivariant Bergman projection associated with the eigenvalue  $\frac{\lceil kE \rceil}{k}$  of  $\hat{H}_k$ . Accordingly, the proof of Theorem 1.6 relies on the second item of Theorem 1.7. The latter is established by similar arguments, namely, the stationary phase lemma is used together with the integral representation of equivariant Bergman kernels as "Fourier coefficients" of the full Bergman kernel.

Recall that there exists  $N \geq 1$  and an open dense subset  $M_N \subset M$  such that the stabilizer group of every  $z \in M_N$  is the subgroup  $S_N^1 \subset S^1$  of order N. We assume throughout that N = 1 (i.e., that the circle action is effective); this is justified by the following lemma.

**Lemma 4.4.** It suffices to prove Theorems 1.6, 1.7 in the case N = 1.

*Proof.* Assume that N > 1. If  $t \in S_N^1$ , then  $\varphi_t|_{M_N} = \text{Id}$ , hence by continuity  $\varphi_t = \text{Id}$ . It follows that  $t \mapsto \varphi_{N,t} = \varphi_{\frac{t}{N}}, t \in \mathbb{R}/2\pi\mathbb{Z}$ , is an effective holomorphic circle action, generated by the Hamiltonian  $H_N = \frac{1}{N}H$ . Writing  $\hat{H}_{N,k} = \frac{1}{N}\hat{H}_k$ ,

$$\Pi_{k,E} = \mathbb{1}_{[E,\infty)}(\hat{H}_k) = \mathbb{1}_{[\frac{E}{N},\infty)}\left(\hat{H}_{N,k}\right).$$

Thus, the case N > 1 is reduced to the case N = 1. Similarly,

$$\Pi_{k,\lambda_k}^{\mathrm{eq}} = \mathbb{1}_{\{\lambda_k\}}(\hat{H}_k) = \mathbb{1}_{\{\frac{\lambda_k}{N}\}}(\hat{H}_{N,k}).$$

Recall that  $E \in H(M)$  is a regular value of H, and assume that  $z \in H^{-1}(E)$ . Denote  $V_k(t) = e^{ikt\hat{H}_k}$ , and consider the map

$$V_k(z,w): S^1 \to L_z^{\otimes k} \otimes (L_w^*)^{\otimes k}$$

specified by  $t \mapsto V_k(t)(z, w)$ , where the latter is the Schwartz kernel of  $V_k(t)$ . Denote  $E_k = \frac{\lceil kE \rceil}{k}$  and  $V_{k,E_k}(t)(z,w) = e^{-ikE_kt}V_k(t)(z,w)$ . According to (7),

$$\Pi_{k,E}(z,w) = \frac{1}{2} \left( i \mathcal{H}_{S^1} \left( V_{k,E_k}(z,w) \right)(0) + \Pi_k(z,w) + V_{k,E_k}(z,w)(0) \right), \quad (8)$$

where

$$\widehat{V_{k,E_{k}}(z,w)}(0) = \frac{1}{2\pi} \int_{0}^{2\pi} V_{k,E_{k}}(t)(z,w)dt = \Pi_{k,E_{k}}^{\text{eq}}(z,w)$$

is the equivariant Bergman kernel associated with the eigenvalue  $E_k$  of  $H_k$ . The Schwartz kernel of  $V_{k,E_k}(t)$  admits the following useful formula.

**Lemma 4.5.** The Schwartz kernel of  $V_{k,E_k}(t)$  is specified by

$$V_{k,E_k}(t)(z,w) = c_{k,E_k}(z,t)\mathcal{T}_{k,t}\left(\Pi_k(\varphi_t(z),w)\right)$$

where  $\mathcal{T}_{k,t}: L_{\varphi_t(z)}^{\otimes k} \otimes (L^*)_w^{\otimes k} \to L_z^{\otimes k} \otimes (L^*)_w^{\otimes k}$  is the parallel transport along the curve  $t' \mapsto (\varphi_{t-t'}(z), w), t' \in [0, t]$ , and  $c_{k, E_k}(z, t) = e^{ik(H(z) - E_k)t} = e^{ik(E - E_k)t}$ .

*Proof.* As shown in [16], Proposition 8.2.1, the action of  $V_k(t)$  is specified by

$$V_k(t)s(z) = e^{iktH(z)}\tau_{k,t}s(\varphi_t(z)), \ s \in \mathcal{H}_k,$$

where  $\tau_{k,t}: L_{\varphi_t(z)}^{\otimes k} \to L_z^{\otimes k}$  is the parallel transport along the curve  $t' \mapsto \varphi_{t-t'}(z)$ ,  $t' \in [0, t]$ . Thus, considering any orthonormal basis  $s_1, \ldots, s_{d_k}$  of  $\mathcal{H}_k$ , we find that the Schwartz kernel of  $V_k(t)$  is given by

$$V_k(t)(z,w) = \sum_{m=1}^{d_k} (V_k(t)s_m)(z) \otimes s_m^*(w)$$
  
=  $e^{iktH(z)} \sum_{m=1}^{d_k} (\tau_{k,t}s_m)(\varphi_t(z)) \otimes s_m^*(w)$   
=  $e^{iktH(z)} (\tau_{k,t} \otimes \mathrm{Id}) \left( \sum_{m=1}^{d_k} s_m(\varphi_t(z)) \otimes s_m^*(w) \right)$   
=  $e^{iktH(z)} \mathcal{T}_{k,t} (\Pi_k(\varphi_t(z),w)).$ 

Multiplying both sides by  $e^{-ikE_kt}$ , we obtain the required.

The microsupport  $MS(\Pi_k)$  of  $\Pi_k$  equals the diagonal  $\Delta_M \subset M \times M$ . In particular, this implies (see [4], Proposition 8, or [19], Theorem 1) that for every  $\varepsilon > 0$  and N > 0 there exists  $C_{\varepsilon,N} > 0$  such that for every  $z, w \in M$ , if  $dist(z, w) \ge \varepsilon$  then

$$|\Pi_k(z,w)| < C_{\varepsilon,N} k^{-N},\tag{9}$$

and that for every vector field Y on  $M \times M$  there exists  $C_{Y,\varepsilon,N}$  such that for every  $z, w \in M$ , if  $dist(z, w) \geq \varepsilon$  then

$$|(\tilde{\nabla}_k)_Y \Pi_k(z, w)| < C_{Y,\varepsilon,N} k^{-N}.$$
(10)

Here,  $\tilde{\nabla}_k$  is the connection on  $L^{\otimes k} \boxtimes (L^*)^{\otimes k}$  induced by  $\nabla$ . The first part of Theorem 1.6 readily follows from these estimates.

**Lemma 4.6.** If  $z \in M$  and  $w \in M \setminus \mathcal{O}_z$ , then  $(z, w) \notin MS(\Pi_{k,E})$ .

*Proof.* Let  $\mathcal{N}_0 \subset M \times M$  denote a neighborhood of (z, w) such that for every  $(z', w') \in \mathcal{N}_0$  and  $t \in [-\pi, \pi]$ ,

$$\operatorname{dist}(\varphi_t(z'), w') > \frac{\varepsilon}{2}$$

for some  $\varepsilon > 0$ . We wish to prove that

$$\sup_{\mathcal{N}_0} |\Pi_{k,E}| = \mathcal{O}(k^{-\infty}),$$

which would mean (by definition) that  $(z, w) \notin MS(\Pi_{k,E})$ . Let

$$\mathcal{N} = \{ (\varphi_t(z'), w') \mid (z', w') \in \mathcal{N}_0, \ t \in [-\pi, \pi] \}$$

Looking at the representation (8) in light of (9), we immediately note that the second term satisfies

$$\sup_{\mathcal{N}_0} |\Pi_k| = \mathcal{O}(k^{-\infty}).$$

Similarly, estimate (9) implies that the term  $V_{k,E_k}(z',w')(0) = \prod_{k,E_k}^{eq}(z',w')$ , which is the equivariant Bergman kernel associated with the eigenvalue  $E_k$ , satisfies

$$\sup_{(z',w')\in\mathcal{N}_0} \left| \Pi_{k,E_k}^{\mathrm{eq}}(z',w') \right| \le \sup_{\mathcal{N}_0} \frac{1}{2\pi} \int_{-\pi}^{\pi} \left| \Pi_k\left(\varphi_t(z'),w'\right) \right| dt = \mathcal{O}(k^{-\infty}).$$
(11)

Finally, consider the first term in (8). Let  $V'_{k,E_k}(u)(z',w') = \frac{d}{dt}(V_k(z',w'))(u)$ . Note that  $V_{k,E_k}(0)(z',w') = \prod_k(z',w')$ . If  $t \in (0,\pi)$ , then by the mean-value theorem,

$$|V_{k,E_k}(t)(z',w') - \Pi_k(z',w')| \le t \sup_{u \in [0,t]} |V'_{k,E_k}(u)(z',w')|,$$

where (in light of Lemma 4.5)

$$V'_{k,E_{k}}(u)(z',w') = c_{k,E_{k}}(u)\mathcal{T}_{k,u}\left(ik(E-E_{k})\Pi_{k}(\varphi_{u}(z'),w') + (\tilde{\nabla}_{k})_{Y_{H}}(\Pi_{k})(\varphi_{u}(z'),w')\right),$$

with  $Y_H = (X_H, 0)$ .

Applying (9) and (10), we see that

$$\begin{aligned} \left| V_{k,E_k}'(u)(z',w') \right| &\leq k |E - E_k| |\Pi_k(\varphi_u(z'),w')| + |(\tilde{\nabla}_k)_{Y_H} \Pi_k(\varphi_u(z'),w')| \\ &\leq \sup_{\mathcal{N}} |\Pi_k| + \sup_{\mathcal{N}} |(\tilde{\nabla}_k)_{Y_H} \Pi_k| = \mathcal{O}(k^{-\infty}). \end{aligned}$$
(12)

Thus,

$$\sup_{\mathcal{N}_0 \times (0,\pi)} \frac{1}{t} |V_{k,E_k}(t)(z',w') - \Pi_k(z',w')| = \mathcal{O}(k^{-\infty}),$$

which implies (since if  $t \approx 0$ , then  $\cot\left(\frac{t}{2}\right) \approx \frac{2}{t}$ ) that

$$\sup_{\mathcal{N}_0} |\mathcal{H}_{S^1}(V_{k,E_k}(z',w'))(0)| = \mathcal{O}(k^{-\infty}).$$

We proceed with the proof of the second item of Theorem 1.6, relying on formula (8). Fix  $a, b \in \mathbb{R}$ . The behaviour of the third term in (8) is specified in Theorem 1.7, hence in what follows we focus on  $\mathcal{H}_{S^1}(V_{k,E_k}(z_{a_k,0}, z_{b_k,t_0}))(0)$ , where we denote  $a_k = \frac{a}{\sqrt{k}}$ ,  $b_k = \frac{b}{\sqrt{k}}$ , and for any  $\alpha \in \mathbb{R}$ ,  $t \in S^1$ ,

$$z_{\alpha} = \psi_{\alpha}(z_0), \ z_{\alpha,t} = \varphi_t(z_{\alpha}).$$

**Lemma 4.7.** Fix  $z_0 \in H^{-1}(E) \cap M_1$ . Let  $\mathcal{I} \subset S^1 \setminus \{0\}$  be a closed interval. Then for every fixed  $\delta > 0$  small enough, if  $t_0 \in \mathcal{I}$  then

$$\mathcal{H}_{S^1}(V_{k,E_k}(z_{a_k}, z_{b_k,t_0}))(0) = -\frac{1}{2\pi} \int_{t_0-\delta}^{t_0+\delta} V_{k,E_k}(t)(z_{a_k}, z_{b_k,t_0}) \cot\left(\frac{t}{2}\right) dt + r_k(a, b, t_0)$$

Here,  $r_k(a, b, t_0) = \mathcal{O}(k^{-\infty})$  uniformly, in the following sense: for every bounded set  $B \subset \mathbb{R}$  and N > 0, there exists  $c_{B,N} > 0$  such that  $|r_k(a, b, t_0)| \leq c_{B,N}k^{-N}$  for every  $a, b \in B$ ,  $t_0 \in \mathcal{I}$ .

*Proof.* Assume that  $\mathcal{I} = [t', t''] \subset (0, 2\pi)$ . Let  $\delta > 0$  be small enough so that

$$([0,\delta] \cup [2\pi - \delta, 2\pi]) \cap [t' - \delta, t'' + \delta] = \emptyset.$$

There exists  $\varepsilon > 0$  such that for every  $t \in [0, 2\pi]$  and  $t_0 \in \mathcal{I}$ , if  $|t - t_0| \geq \delta$ then  $\operatorname{dist}(z_{0,t}, z_{0,t_0}) > \varepsilon$ . Let  $B \subset \mathbb{R}$  be bounded. There exists  $k_0 \in \mathbb{N}$  such that for every  $k \geq k_0$ , for every  $a, b \in B$  and for every  $t \in [0, 2\pi]$ ,  $t_0 \in \mathcal{I}$  such that  $|t - t_0| \geq \delta$ , it holds that

$$\operatorname{dist}(z_{a_k,t}, z_{b_k,t_0}) > \frac{\varepsilon}{2}.$$
(13)

Thus, the mean-value theorem together with a suitable version of estimate (12) imply that

$$\sup_{(0,\delta]} \left| (V_{k,E_k}(t)(z_{a_k}, z_{b_k,t_0}) - V_{k,E_k}(-t)(z_{a_k}, z_{b_k,t_0})) \cot\left(\frac{t}{2}\right) \right| = \mathcal{O}(k^{-\infty}),$$

and (in light of (13), (10)) the estimate is uniform for  $a, b \in B$  and  $t_0 \in \mathcal{I}$ . Thus,

$$\begin{split} \int_{0}^{\pi} \left( V_{k,E_{k}}(t)(z_{a_{k}}, z_{b_{k},t_{0}}) - V_{k,E_{k}}(-t)(z_{a_{k}}, z_{b_{k},t_{0}}) \right) \cot\left(\frac{t}{2}\right) dt &= \\ \int_{\delta}^{\pi} \left( V_{k,E_{k}}(t)(z_{a_{k}}, z_{b_{k},t_{0}}) - V_{k,E_{k}}(-t)(z_{a_{k}}, z_{b_{k},t_{0}}) \right) \cot\left(\frac{t}{2}\right) dt + \mathcal{O}(k^{-\infty}) \\ &= \int_{\delta}^{2\pi-\delta} V_{k,E_{k}}(t)(z_{a_{k}}, z_{b_{k},t_{0}}) \cot\left(\frac{t}{2}\right) dt + \mathcal{O}(k^{-\infty}) \end{split}$$

where (in light of (13), (9)) the remainder estimate is uniform for  $a, b \in B$  and  $t_0 \in \mathcal{I}$ . Finally,

$$\int_{\delta}^{t_0-\delta} V_{k,E_k}(t)(z_{a_k}, z_{b_k,t_0}) \cot\left(\frac{t}{2}\right) dt = \mathcal{O}(k^{-\infty}),$$

and again, by (13), (9), the estimate is uniform for  $a, b \in B$ ,  $t_0 \in \mathcal{I}$ . Similarly for the integration over  $[t_0 + \delta, 2\pi - \delta]$ .

**Corollary 4.8.** Let  $S_k(z, w) = \sigma(z)^{\otimes k} \otimes (\sigma_{inv}^*(w))^{\otimes k}$ , where  $\sigma = \frac{s_{inv}}{|s_{inv}|}$ , with  $s_{inv}$  a local non-vanishing invariant section of L in a neighborhood of  $\mathcal{O}_{z_0}$ . Note that  $||X_H||$  is constant in  $\mathcal{O}_{z_0}$ . In light of the previous lemma and (17), we conclude that

$$-\left(\frac{2\pi}{k}\right)^{n}\mathcal{H}_{S^{1}}(V_{k,E_{k}}(z_{a_{k}},z_{b_{k},t_{0}}))(0) = \\ \left(e^{-\frac{(a^{2}+b^{2})}{2}\|X_{H}(z_{0})\|^{2}}e^{-ikE_{k}t_{0}}\frac{\cot\left(\frac{t_{0}}{2}\right)}{\|X_{H}(z_{0})\|\sqrt{\pi k}} + C_{k}(a,b,t_{0})\right)S_{k}(z_{a_{k}},z_{b_{k},t_{0}}).$$

Here,  $C_k(a, b, t_0) = \frac{C_k^v(a, b, t_0)}{2\pi}$ , where  $C_k^v(a, b, t)$  is as specified in Corollary 5.2, with  $v(t) = \cot\left(\frac{t}{2}\right)$ .

In order to complete the proof of Theorem 1.6, it remains to establish the off-diagonal estimate for the equivariant Bergman kernel  $\Pi_{k,E_k}$ , as formulated in Theorem 1.7. Hence, we now turn to address the proof of the latter. Assume that  $\lambda_k$  is an eigenvalue of  $\hat{H}_k$  such that  $|\lambda_k - E| = \mathcal{O}(k^{-1})$ . Let

$$\Pi_{k,\lambda_k}^{\rm eq} = \frac{1}{2\pi} \int_0^{2\pi} V_k(t) e^{-ik\lambda_k t} dt$$

denote the orthogonal projection on the eigenspace associated with  $\lambda_k$ .

**Corollary 4.9.** Estimate (11) in the proof of Lemma 4.6 is also valid for  $\Pi_{k,\lambda_k}^{\text{eq}}$ . Namely, if  $z \in M$  and  $w \in M \setminus \mathcal{O}_z$ , then  $(z, w) \notin \text{MS}(\Pi_{k,\lambda_k}^{\text{eq}})$ . Together with Lemma 3.7, we obtain the first item of Theorem 1.7.

**Corollary 4.10.** Let  $z_0 \in H^{-1}(E) \cap M_1$  and  $\mathcal{I} \subset S^1$  be a closed interval. The same argument as in the proof of Lemma 4.7 implies that for every fixed  $\delta > 0$  small enough, if  $t_0 \in \mathcal{I}$  then

$$\Pi_{k,\lambda_k}^{\rm eq}(z_{a_k}, z_{b_k,t_0}) = \int_{t_0-\delta}^{t_0+\delta} V_k(t)(z_{a_k}, z_{b_k,t_0})e^{-ik\lambda_k t}dt + r_k^{\rm eq}(a, b, t_0)dt$$

Here,  $r_k^{\text{eq}}(a, b, t_0) = \mathcal{O}(k^{-\infty})$  uniformly, as before: for every  $B \subset \mathbb{R}$  bounded and N > 0, there exists  $c_{B,N} > 0$  such that  $|r_k^{\text{eq}}(a, b, t_0)| \leq c_{B,N}k^{-N}$  for every  $a, b \in B, t_0 \in \mathcal{I}$ .

Thus, by (17),

$$\left(\frac{2\pi}{k}\right)^{n} \Pi_{k,\lambda_{k}}^{\mathrm{eq}}(z_{a_{k}}, z_{b_{k},t_{0}}) = \left(e^{-\frac{(a^{2}+b^{2})}{2}\|X_{H}(z_{0})\|^{2}}e^{-ik\lambda_{k}t_{0}}\frac{1}{\|X_{H}(z_{0})\|\sqrt{\pi k}} + C_{k}^{\mathrm{eq}}(a, b, t_{0})\right)S_{k}(z_{a_{k}}, z_{b_{k},t_{0}}),$$

where  $C_k^{\text{eq}}(a, b, t_0) = \frac{C_k^v(a, b, t_0)}{2\pi}$ , with  $C_k^v(a, b, t)$  as specified in Corollary 5.2, for v = 1 (also,  $S_k(z, w)$  is as specified in Corollary 4.8).

# 5 Estimates of integrals involving the Bergman kernel

We begin by noting that  $\Pi_k$  admits the representation ([4], [16], Theorem 7.2.1, [14], Theorem 4.11)

$$\Pi_k(z,w) = \left(\frac{k}{2\pi}\right)^n u(z,w,k) \mathcal{E}^{\otimes k}(z,w) + R_k(z,w),$$
(14)

where  $\mathcal{E} \in C^{\infty}(M \times M, L \boxtimes L^*)$ ,  $u(\cdot, \cdot, k) \in C^{\infty}(M \times M)$  is real valued, and  $R_k$  have the following properties.

- The section  $\mathcal{E}$  satisfies  $|\mathcal{E}(z,w)| < 1$  for  $z \neq w$ , and  $\mathcal{E}(z,z) = 1$ .
- The function  $u(\cdot, \cdot, k)$  is real-valued, and  $u(\cdot, \cdot, k) \sim \sum_{l=0}^{\infty} k^{-l} u_l(\cdot, \cdot)$  in the  $C^{\infty}$ -topology<sup>6</sup>, with  $u_0(z, z) \equiv 1$ .
- $R_k = \mathcal{O}(k^{-\infty})$  uniformly in (z, w).

Recall that  $\{\varphi_t\}_{t\in[0,2\pi]}$  and  $\{\psi_a\}_{a\in\mathbb{R}}$  denote the Hamiltonian flow and gradient flow associated with H. The two flows commute, and define a  $\mathbb{C}^*$ -action on M(the gradient flow also consists of biholomorphisms). As before, given  $z_0 \in M$ , if  $\alpha \in \mathbb{R}$  and  $t \in S^1$ , then we denote

$$z_{\alpha} = \psi_{\alpha}(z_0), \ z_{\alpha,t} = \varphi_t(z_{\alpha}).$$

**Lemma 5.1.** Fix  $z_0 \in M$ . Let  $s_{inv}$  be a local invariant holomorphic section of L in a neighborhood of  $\mathcal{O}_{z_0}$ . Let  $S(z, w) = \sigma(z) \otimes \sigma^*(w)$ , where  $\sigma = \frac{s_{inv}}{|s_{inv}|}$ . For  $\delta > 0$  small enough, define

$$g_{z_0}: (-\delta, \delta) \times (-\delta, \delta) \times (-\delta, \delta) \times S^1 \to \mathbb{C}$$

by

$$\mathcal{E}(z_{a,t+t'}, z_{b,t'}) = e^{ig_{z_0}(t, a, b, t')} S(z_{a,t+t'}, z_{b,t'}).$$
(15)

Let  $t_0 \in S^1$ . Then

$$\begin{split} g_{z_0}(0,t_0) &= 0, \\ \partial_t g_{z_0}(0,t_0) &= H(z_0), \partial_a g_{z_0}(0,t_0) = \partial_b g_{z_0}(0,t_0) = 0, \\ \partial_t^2 g_{z_0}(0,t_0) &= \partial_a^2 g_{z_0}(0,t_0) = \partial_b^2 g_{z_0}(0,t_0) = -\partial_a \partial_b g_{z_0}(0,t_0) = \frac{i}{2} \|X_H\|^2, \\ \partial_a \partial_t g_{z_0}(0,t_0) &= \partial_b \partial_t g_{z_0}(0,t_0) = \frac{1}{2} \|X_H\|^2. \end{split}$$

<sup>5</sup>Here, we use the Hermitian metric to make the identification  $L_z \otimes L_z^* \simeq \mathbb{C}$ .

<sup>&</sup>lt;sup>6</sup>This means that for every  $N \ge 0$ , the function  $u(\cdot, \cdot, k) - \sum_{l=0}^{N} k^{z_l} u_l(\cdot, \cdot)$  and all its derivatives are uniformly  $\mathcal{O}(k^{-(N+1)})$ .

Proof. First,  $\mathcal{E}(\varphi_{t_0}(z_0), \varphi_{t_0}(z_0)) = S(\varphi_{t_0}(z_0), \varphi_{t_0}(z_0))$ , hence  $g_{z_0}(0, t_0) = 0$ . Let  $\tilde{\nabla}$  denote the connection on  $L \boxtimes L^* \to M \times M$  induced from  $\nabla$ . Then

$$\begin{split} \tilde{\nabla}_{(X_{H},0)} \mathcal{E}(z_{a,t+t_{0}}, z_{b,t_{0}}) &= \\ & i\partial_{t}g_{z_{0}}(t, a, b, t_{0})\mathcal{E}(z_{a,t+t_{0}}, z_{b,t_{0}}) \\ &+ e^{ig_{z_{0}}(t, a, b, t_{0})}\tilde{\nabla}_{(X_{H},0)}S(z_{a,t+t_{0}}, z_{b,t_{0}}) \\ &= i\left(\partial_{t}g_{z_{0}}(t, a, b, t_{0}) - H(z_{a})\right)\mathcal{E}(z_{a,t+t_{0}}, z_{b,t_{0}}), \end{split}$$

where we used the invariance of  $s_{inv}$ , and the fact that  $\varphi_{t_0}$  preserves level sets of H.

Let  $\alpha_{\mathcal{E}}$  be the 1-form defined in a neighborhood of the diagonal  $\Delta_M \subset M \times M$ by the equation

$$\nabla \mathcal{E} = -i\alpha_{\mathcal{E}} \otimes \mathcal{E}.$$
(16)

Then

$$i(\partial_t g_{z_0}(t, a, b, t_0) - H(z_a)) = -i\alpha_{\mathcal{E}}(X_H, 0)(z_{a, t+t_0}, z_{b, t_0})$$

that is,

$$\partial_t g_{z_0}(t, a, b, t_0) = H(z_a) - \alpha_{\mathcal{E}}(X_H, 0)(z_{a, t+t_0}, z_{b, t_0})$$

However,  $\alpha_{\mathcal{E}}$  vanishes on  $\Delta_M$  ([16], Lemma 7.1.3), hence  $\partial_t g_{z_0}(0, t_0) = H(z_0)$ . Similarly (noting that  $\nabla_{\text{grad } H} \sigma = 0$ ),

$$\partial_a g_{z_0}(t, a, b, t_0) = -\alpha_{\mathcal{E}}(\operatorname{grad} H, 0)(z_{a,t+t_0}, z_{b,t_0}),$$
  
$$\partial_b g_{z_0}(t, a, b, t_0) = -\alpha_{\mathcal{E}}(0, \operatorname{grad} H)(z_{a,t+t_0}, z_{b,t_0}),$$

and since  $\alpha_{\mathcal{E}}$  vanishes on  $\Delta_M$ , we obtain  $\partial_a g_{z_0}(0, t_0) = \partial_b g_{z_0}(0, t_0) = 0$ . Next, we note that

$$\begin{aligned} \partial_t^2 g_{z_0} &= -\mathcal{L}_{(X_H,0)} \alpha_{\mathcal{E}}(X_H,0), \ \partial_a^2 g_{z_0} &= -\mathcal{L}_{(\operatorname{grad} H,0)} \alpha_{\mathcal{E}}(\operatorname{grad} H,0), \\ \partial_b^2 g_{z_0} &= -\mathcal{L}_{(0,\operatorname{grad} H)} \alpha_{\mathcal{E}}(0,\operatorname{grad} H), \partial_a \partial_b g_{z_0} &= -\mathcal{L}_{(\operatorname{grad} H,0)} \alpha_{\mathcal{E}}(0,\operatorname{grad} H), \\ \partial_t \partial_a g_{z_0} &= -\mathcal{L}_{(X_H,0)} \alpha_{\mathcal{E}}(\operatorname{grad} H,0), \ \partial_t \partial_b g_{z_0} &= -\mathcal{L}_{(X_H,0)} \alpha_{\mathcal{E}}(0,\operatorname{grad} H). \end{aligned}$$

Now, as shown in [16], Lemma 7.1.3, there exists a smooth section  $B_{\mathcal{E}}$  of the bundle  $T^*(M \times M) \otimes T^*(M \times M) \otimes \mathbb{C} \to \Delta_M$  such that for vector fields X, Y on  $M \times M$ ,

$$\mathcal{L}_X(\alpha_{\mathcal{E}}(Y)) = B_{\mathcal{E}}(Y,Y) = \tilde{\omega}(q(Y_H),Y_H)$$

on  $\Delta_M$ . Here, q is the projection from  $T_{(w,w)}(M \times M) \otimes \mathbb{C}$  onto  $T_{(w,w)}^{0,1}(M \times M)$ with kernel  $T_{(w,w)}\Delta_M \otimes \mathbb{C}$ , and  $\tilde{\omega} = \pi_1^* \omega - \pi_2^* \omega$  is the Kähler form on  $M \times M$ . We can compute q(X, 0) and q(0, X) explicitly, and obtain

$$q(X,0) = \frac{1}{2} \left( X + ijX, -X + ijX \right), \ q(0,X) = \frac{1}{2} \left( -X + ijX, X + ijX \right)$$

where j is the complex structure on M. Thus, a straightforward computation produces

$$\partial_t^2 g_{z_0}(0, t_0) = -\tilde{\omega}(q(X_H, 0), (X_H, 0)) = -\frac{1}{2}\omega(ijX_H, X_H) = \frac{i}{2} ||X_H||^2,$$

and noting that grad  $H = -jX_H$ ,

$$\partial_t \partial_a g_{z_0}(0, t_0) = -\tilde{\omega}(q(X_H, 0), (\text{grad } H, 0)) = -\frac{1}{2}\omega(X_H, -jX_H) = \frac{1}{2}||X_H||.$$

The rest of the cases are computed in the same way.

Using Lemma 5.1, we can apply the stationary phase approximation in order to estimate integrals (along  $S^1$ -trajectories) which involve the Bergman kernel. In light of Lemma 4.4, we assume that N = 1 (i.e., there exists an open dense  $M_1 \subset M$  such that the stabilizer group of every  $z \in M_1$  is trivial).

**Corollary 5.2.** Let  $\lambda_k$  be an eigenvalue of  $\hat{H}_k$  such that  $|\lambda_k - E| = \mathcal{O}(k^{-1})$ , where  $E \in H(M)$  is a regular value. Fix  $z_0 \in H^{-1}(E) \cap M_1$ . Let  $\mathcal{I} \subset S^1$ be a closed interval. Let v be a smooth function, compactly supported in a neighborhood of  $\mathcal{I}$ , such that  $v|_{\mathcal{I}} \neq 0$ . Fix  $a, b \in \mathbb{R}$ . For  $\delta > 0$  small enough and  $t_0 \in \mathcal{I}$ , denote

$$I_{k,\delta}^{v,\lambda_k}(\alpha,\beta,t_0) = \left(\frac{2\pi}{k}\right)^n \int_{t_0-\delta}^{t_0+\delta} v(t) V_{k,\lambda_k}(t)(z_\alpha,z_{\beta,t_0}) dt,$$

where  $V_{k,\lambda_k}(t) = e^{-ik\lambda_k t} V_k(t)$ . Write  $a_k = \frac{a}{\sqrt{k}}$ ,  $b_k = \frac{b}{\sqrt{k}}$ . Then

$$I_{k,\delta}^{\nu,\lambda_k}(a_k, b_k, t_0) = \mathcal{K}_k^{\nu,\lambda_k}(a, b, t_0) S_k(z_{a_k}, z_{b_k, t_0}),$$
(17)

with  $S_k$  as specified in Theorem 1.6, and

$$\mathcal{K}_{k}^{v,\lambda_{k}}(a,b,t_{0}) = e^{-\frac{(a^{2}+b^{2})\|X_{H}(z_{0})\|^{2}}{2}}e^{-ik\lambda_{k}t_{0}}\frac{2v(t_{0})}{\|X_{H}(z_{0})\|}\sqrt{\frac{\pi}{k}} + C_{k}^{v}(a,b,t_{0}).$$

Here,

$$C_k^v(a,b,t) = \frac{C_{k,1}^v(a,b,t)}{k} + \frac{C_{k,2}^v(a,b,t)}{k^{\frac{3}{2}}}$$

where  $C_{k,1}^v(a, b, t)$ ,  $C_{k,2}^v(a, b, t)$  satisfy that for any bounded set  $B \subset \mathbb{R}$  there exist  $c_{B,1}^v, c_{B,2}^v > 0$  such that

$$|C_{k,1}^{v}(a,b,t)| \le (|a|+|b|)c_{B,1}^{v}, \ |C_{k,2}^{v}(a,b,t)| \le c_{B,2}^{v}$$

for all  $a, b \in B$  and  $t \in \mathcal{I}$ .

Proof. According to Lemma 4.5,

$$V_{k,\lambda_k}(t)(z_{\alpha}, z_{\beta,t_0}) = c_{k,\lambda_k}(z_{\alpha}, t)\mathcal{T}_{k,t}\Pi_k(z_{\alpha,t}, z_{\beta,t_0}),$$

where  $c_{k,\lambda_k}(z,t) = e^{ik(H(z)-\lambda_k)t}$ . Then using (14),

$$\mathcal{T}_{k,t}\Pi_k(z_{\alpha,t}, z_{\beta,t_0}) = \left(\frac{k}{2\pi}\right)^n u(z_{\alpha,t}, z_{\beta,t_0}, k) \mathcal{T}_{k,t} \left(\mathcal{E}^{\otimes k}(z_{\alpha,t}, z_{\beta,t_0})\right).$$

The invariance of  $s_{inv}$  implies that  $\mathcal{T}_{k,t}S_k(\varphi_t(z), w) = e^{-ikH(z)t}S_k(z, w)$ . Thus, in the notations of Lemma 5.1,

$$\mathcal{T}_{k,t}\left(\mathcal{E}^{\otimes k}(z_{\alpha,t}, z_{\beta,t_0})\right) = e^{ik(g_{z_0}(t-t_0, \alpha, \beta, t_0) - H(z_\alpha)t)} S_k(z_\alpha, z_{\beta,t_0}).$$

Hence,

$$I_{k,\delta}^{v,\lambda_k}(a_k,b_k,t_0) = \left(\int_{t_0-\delta}^{t_0+\delta} \tilde{v}_k(t,a_k,b_k,t_0)e^{ikg_{z_0}(t-t_0,a_k,b_k,t_0)}dt\right)S_k(z_{a_k},z_{b_k,t_0}),$$

with

$$\tilde{v}_k(t,\alpha,\beta,t_0) = v(t)e^{-ik\lambda_k t}u(z_{\alpha,t},z_{\beta,t_0},k).$$

Next, applying a change of variables,

$$\int_{t_0-\delta}^{t_0+\delta} \tilde{v}_k(t,a_k,b_k,t_0) e^{ikg_{z_0}(t-t_0,a_k,b_k,t_0)} dt$$
  
=  $e^{-ik\lambda_k t_0} \int_{-\delta}^{\delta} v_k(t,a_k,b_k,t_0) e^{ikf_{z_0}(t,a_k,b_k,t_0)} dt$ 

where (noting that  $H(z_0) = E$ )

$$v_k(t, \alpha, \beta, t_0) = v(t+t_0)e^{ik(E-\lambda_k)t}u(z_{\alpha,t+t_0}, z_{\beta,t_0}, k),$$
  
$$f_{z_0}(t, \alpha, \beta, t') = g_{z_0}(t, \alpha, \beta, t') - Et.$$

Now,  $\partial_t f_{z_0}(0, t_0) = 0$  and  $\partial_t^2 f_{z_0}(0, t_0) \neq 0$  by Lemma 5.1, and we can assume without loss of generality that t = 0 is the unique such point in  $(-\delta, \delta)$ . Also, the derivatives of  $v_k$  are bounded as  $k \to \infty$  (since  $|E - \lambda_k| = \mathcal{O}(k^{-1})$ ). Hence, we can apply the stationary phase lemma for a complex valued phase ([13], Theorem 7.7.12), and obtain

$$\int_{-\delta}^{\delta} v_k(t, a_k, b_k, t_0) e^{ikf_{z_0}(t, a_k, b_k, t_0)} dt$$
  
=  $\left(\frac{2\pi i}{k \left(\partial_t^2 f_{z_0}\right)^0 (a_k, b_k, t_0)}\right)^{\frac{1}{2}} e^{ikf_{z_0}^0 (a_k, b_k, t_0)} v_k^0(a_k, b_k, t_0) + \mathcal{O}(k^{-\frac{3}{2}}).$ 

Here, for a function F(t, a, b, t'), the notation  $F^0(a, b, t')$  stands for a function of a, b, t' only which belongs to the same residue class as F modulo the ideal generated by  $\partial_t F$  (i.e.,  $F - F^0 = G \partial_t F$  for some function G). Also, the square root is defined such that its real part is non-negative. Finally, the  $\mathcal{O}(k^{-\frac{3}{2}})$ estimate is uniform for a, b in bounded sets and  $t_0 \in \mathcal{I}$ .

The derivatives of  $v_k^0$  are bounded as  $k \to \infty$ , hence (using Taylor's theorem)

$$\begin{aligned} v_k^0(a_k, b_k, t_0) &= v_k^0(0, t_0) + \frac{c_0(k, a, b, t_0)}{\sqrt{k}} \\ &= v_k(0, t_0) + \frac{c_0(k, a, b, t_0)}{\sqrt{k}} \\ &= v(t_0) + \frac{c_0(k, a, b, t_0)}{\sqrt{k}} + \mathcal{O}(k^{-1}) \end{aligned}$$

for some  $c_0(k, a, b, t_0)$  satisfying that for every bounded set  $B \subset \mathbb{R}$  there exists  $c_{0,B} > 0$  such that  $|c_0(k, a, b, t_0)| \leq (|a| + |b|)c_{0,B}$  for all  $a, b \in B$ ,  $t_0 \in \mathcal{I}$ . Also, the  $\mathcal{O}(k^{-1})$  estimate is uniform for  $a, b \in B$  and  $t_0 \in \mathcal{I}$ .

Similarly,

$$\left( \frac{2\pi i}{k \left(\partial_t^2 f_{z_0}\right)^0 (a_k, b_k, t_0)} \right)^{\frac{1}{2}} = \left( \frac{2\pi i}{k \left(\partial_t^2 f_{z_0}\right)^0 (0, t_0)} \right)^{\frac{1}{2}} + \frac{c_1(k, a, b, t_0)}{k}$$
$$= \frac{2}{\|X_H\|} \sqrt{\frac{\pi}{k}} + \frac{c_1(k, a, b, t_0)}{k}.$$

Again,  $c_1(k, a, b, t_0)$  satisfies that for every bounded set  $B \subset \mathbb{R}$  there exists  $c_{1,B} > 0$  such that  $|c_1(k, a, b, t_0)| \leq (|a| + |b|)c_{1,B}$  for all  $a, b \in B$ ,  $t_0 \in \mathcal{I}$ .

Finally, we may choose (see [13], 7.7.16, and the succeeding paragraph)

$$f_{z_0}^0(a,b,t') = f_{z_0}(0,a,b,t') - q(0,a,b,t')T(a,b,t')^2,$$

where T such that  $T(0, t_0) = 0$  for every  $t_0 \in \mathcal{I}$ , and q is some smooth function. It follows that

$$\begin{split} f_{z_0}^0(0,t_0) &= f_{z_0}(0,t_0) = 0, \\ \partial_a f_{z_0}^0(0,t_0) &= \partial_a f_{z_0}(0,t_0) = 0 = \partial_b f_{z_0}(0,t_0) = \partial_b f_{z_0}^0(0,t_0). \end{split}$$

Also,

$$\begin{aligned} \partial_a^2 f_{z_0}^0(0,t_0) &= \partial_a^2 f_{z_0}(0,t_0) - 2q(0,t_0)(\partial_a T(0,t_0))^2, \\ \partial_a \partial_b f_{z_0}^0(0,t_0) &= \partial_a \partial_b f_{z_0}(0,t_0) - 2q(0,t_0)\partial_b T(0,t_0)\partial_a T(0,t_0), \\ \partial_b^2 f_{z_0}^0(0,t_0) &= \partial_b^2 f_{z_0}(0,t_0) - 2q(0,t_0)(\partial_b T(0,t_0))^2, \end{aligned}$$

with

$$\begin{split} &2q(0,t_0) = \partial_t^2 f_{z_0}(0,t_0), \\ &\partial_a T(0,t_0) = -\frac{\partial_a \partial_t f_{z_0}(0,t_0)}{\partial_t^2 f_{z_0}(0,t_0)}, \ \partial_b T(0,t_0) = -\frac{\partial_b \partial_t f_{z_0}(0,t_0)}{\partial_t^2 f_{z_0}(0,t_0)} \end{split}$$

Thus, noting Lemma 5.1, a straightforward computation produces

$$\begin{split} f^{0}_{z_{0}}(a_{k},b_{k},t_{0}) &= f^{0}_{z_{0}}(0,t_{0}) + a_{k}\partial_{a}f^{0}_{z_{0}}(0,t_{0}) + b_{k}\partial_{b}f^{0}_{z_{0}}(0,t_{0}) \\ &+ \frac{a_{k}^{2}}{2}\partial_{a}^{2}f^{0}_{z_{0}}(0,t_{0}) + a_{k}b_{k}\partial_{a}\partial_{b}f^{0}_{z_{0}}(0,t_{0}) \\ &+ \frac{b_{k}^{2}}{2}\partial_{b}^{2}f^{0}_{z_{0}}(0,t_{0}) + \frac{c_{2}(k,a,b,t_{0})}{k^{\frac{3}{2}}} \\ &= \frac{i}{2k}(a^{2}+b^{2})\|X_{H}\|^{2} + \frac{c_{2}(k,a,b,t_{0})}{k^{\frac{3}{2}}}. \end{split}$$

As before,  $c_2(k, a, b, t_0)$  satisfies that for every bounded set  $B \subset \mathbb{R}$  there exists  $c_{2,B} > 0$  such that  $|c_2(k, a, b, t_0)| \leq (|a|+|b|)c_{2,B}$  for all  $a, b \in B, t_0 \in \mathcal{I}$ . Putting everything together, we obtain the required.

## 6 Concluding remarks

The spectral projections addressed in Theorem 1.6 (the main result of this work) are analogues of so-called *Melrose-Uhlmann projections* [11], which are defined in the framework of pseudodifferential quantization. Melrose-Uhlmann projections on  $S^1$ -symmetric manifolds are closely related to the quantization of symplectic cuts ([17, 11]). More generally, they are instances of operators whose Schwartz kernels are *distributions of Melrose-Uhlmann type* ([20]); these are distributions associated with (suitable) pairs of intersecting Lagrangian submanifolds, and they admit a symbol calculus. It would be interesting to study whether an analogous theory exists in the framework of geometric quantization.

Finally, we note that our study of partial Bergman kernels has two obvious shortcomings. First, the main result (Theorem 1.6) only addresses the case of Hamiltonians generating holomorphic circle actions; we except that more robust methods would make it possible to establish a version of Theorem 1.6 which is valid in greater generality (cf. [28]). Second, the results presented in this work are all local; it would be desirable to obtain a uniform description of partial Bergman kernels (it is not obvious that such a description exists), as has been done for Schwartz kernels of Toeplitz operators and of quantum propagators ([5, 6]).

#### Acknowledgements

This research has been supported by the DFG funded project SFB/TRR 191 (Project-ID 281071066-TRR 191) "Symplectic Structures in Geometry, Algebra and Dynamics", and the ANR-DFG project QuaSiDy (Project-ID ANR-21-CE40-0016) "Quantization, Singularities, and Holomorphic Dynamics". I wish to express my sincere gratitude to the ANR and to the DFG.

I wish to thank X. Ma and G. Marinescu for many helpful discussions and for their valuable comments on an earlier version of this work. I also wish to thank L. Charles, C.-Y. Hsiao, N. Savale and A. Uribe for interesting and useful discussions on the topics addressed in this work.

### References

- R. Berman, Bergman kernels and equilibrium measures for line bundles over projective manifolds, Am. J. Math. 131, 1485-1524 (2009)
- [2] R. Berman, B. Berndtsson, J. Sjöstrand, A direct approach to Bergman kernel asymptotics for positive line bundles, Ark. Mat. 46 (2), 197-217 (2008)
- [3] D. Catlin, The Bergman kernel and a theorem of Tian, in: Analysis and Geometry in Several Complex Variables (Katata, 1997), Trends Math., Boston
   : Birkhäuser Boston, 1-23, 1999.
- [4] L. Charles, Berezin-Toeplitz Operators, a Semi-Classical Approach, Commun. Math. Phys. 239, 1-28 (2003)

- [5] L. Charles, Quasimodes and Bohr-Sommerfeld conditions for the Toeplitz operators. Comm. Partial Differential Equations 28, 1527-1566 (2003)
- [6] L. Charles, Y. Le Floch, Quantum propagation for Berezin-Toeplitz operators, arXiv:2009.05279 (2020)
- [7] D. Coman, G. Marinescu, On the first order asymptotics of partial Bergman kernels, Ann. Fac. Sci. Toulouse Math. 26(6), 1193-1210 (2017)
- [8] X. Dai, K. Liu, X. Ma, On the asymptotic expansion of Bergman kernel, J. Diff. Geom. 72 (1), 1-41 (2006)
- [9] S. Finski, Semiclassical Ohsawa-Takegoshi extension theorem and asymptotics of the orthogonal Bergman kernel, arXiv:2109.06851
- [10] S. Finski, The asymptotics of the optimal holomorphic extensions of holomorphic jets along submanifolds, arXiv:2207.02761
- [11] V. Guillemin, E. Lerman, Melrose-Uhlmann projectors, the metaplectic representation and symplectic cuts, J. Diff. Geom 61, 365-396 (2002)
- [12] V. Guillemin, S. Sternberg, Geometric quantization and multiplicities of group representations, Invent. Math. 67, no. 3, 515538 (1982)
- [13] L. Hörmander, The Analysis of Linear Partial Differential Operators I: Distribution Theory and Fourier Analysis, Classics in Mathematics, Springer Berlin Heidelberg, 2003.
- [14] C.-Y. Hsiao, G. Marinescu, Asymptotics of spectral function of lower energy forms and Bergman kernel of semi-positive and big line bundles, Commun. Anal. Geom. 22, 1-108 (2014)
- [15] F. W. King, Hilbert Transforms, volume 125 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2009.
- [16] Y. Le Floch, A Brief Introduction to Berezin-Toeplitz Operators on Compact Kähler Manifolds, CRM Short Courses, Springer International Publishing, 2014.
- [17] E. Lerman, Symplectic cuts, Math. Res. Lett. 2, 247-258 (1995)
- [18] X. Ma, G. Marinescu, Holomorphic Morse inequalities and Bergman kernels, Progress in Mathematics, vol. 254, Birkhäuser Boston Inc, Boston, 2007.
- [19] X. Ma, G. Marinescu, Exponential estimate for the asymptotics of Bergman kernels, Math. Ann. 362, no. 3-4, 1327-1347 (2015)
- [20] R.B. Melrose, G. Uhlmann, Lagrangian intersections and the Cauchy problem, Comm. Pure Appl. Math. 32, 483-519 (1979)
- [21] F. Pokorny, M. Singer, Toric partial density functions and stability of toric varieties, Math. Ann. 358 (3-4), 879-923 (2014)

- [22] J. Ross, M. Singer, Asymptotics of partial density functions for divisors, J. Geom. Anal. 27(3), 1803-1854 (2017)
- [23] O. Shabtai, Commutators of spectral projections of spin operators, J. Lie Theory 31(3), 599-624 (2020)
- [24] O. Shabtai, On polynomials in spectral projections of spin operators, Lett. Math. Phys 111, 119 (2021)
- [25] G.M. Tuynman, Quantization: Towards a comparison between methods. J. Math. Phys. 28(12), 2829-2840 (1987)
- [27] S. Zelditch, P. Zhou, Interface asymptotics of partial Bergman kernels on S<sup>1</sup>-symmetric Kähler manifolds, J. Symplectic Geom **17** (3), 793-856 (2019)
- [28] S. Zeditch, P. Zhou, Central limit theorem for spectral partial Bergman kernels. Geom. Topol. 23(4), 1961-2004 (2019)

Ood Shabtai

UNIVERSITÉ PARIS CITÉ CNRS, IMJ-PRG, BÂTIMENT SOPHIE GERMAIN, UFR DE MATHÉMATIQUES CASE 7012, 75205 PARIS CEDEX 13 FRANCE *E-mail address*: shabtai@imj-prg.fr

AND

UNIVERSITÉ DE LILLE LABORATOIRE DE MATHÉMATIQUES PAUL PAINLEVÉ CNRS U.M.R. 8524 59655 VILLENEUVE D'ASCQ CEDEX FRANCE