

# ‘Anti-Gravity’ inside a Menger Sponge

Karl Svozil<sup>1,\*</sup>

<sup>1</sup>*Institute for Theoretical Physics, TU Wien, Wiedner Hauptstrasse 8-10/136, 1040 Vienna, Austria*  
(Dated: January 30, 2024)

This speculative argument proposes that within a physical, ponderable model of space-time, anti-gravity may naturally arise. The central concept involves a paradigm shift from ‘adding stuff’ to ‘thinning out,’ wherein the local geometry of space-time, as perceived by embedded observers, plays a crucial role. In this context, the equivalence principle holds, emphasizing the focus on the intrinsic structure of space-time rather than interactions among individual particles or quanta.

Keywords: fractal gravity

## I. INTUITIVE IMAGES ON FRACTAL GRAVITY

In this speculative paper we try to answer the following question: How would an embedded [1], intrinsic [2] observer experience a fractal substratum which supports motion? In particular, what would be the intrinsic physics of a Menger Sponge?

The Menger Sponge is a three-(exterior)dimensional generalization of the two-(exterior)dimensional Sierpinski carpet, which in turn is an extension of a one-(exterior)dimensional Cantor set. It is generated by beginning with a cube, and dividing every face of the cube into nine squares, then removing or ‘cutting out’ the smaller cube in the middle of each face, as well as removing the smaller cube in the center of the original cube; and by recursively repeating these steps for each of the remaining smaller cubes.

One issue that comes to mind (among many others) is the apparent ‘discontinuity’ of the ‘disjoint’, although path-connected, parts of this construction—which nevertheless forms a set theoretic continuum. Stated pointedly: how can an embedded, intrinsic observer ‘experience’ continuous motion if the underlying substratum is disjointed, disconnected, discontinuous and discrete? We propose that this is an old question that was already put forward by the Eleatics in Zeno’s ‘arrow paradox’. We will not propose any ‘(re)solution’ of this paradox here, but just observe that the canonical information of evolution or ‘movement’ is not only about space but about the (information encoded in the) velocity (momentum/inertia).

Another issue is the possible operational constructions [3] of entities such as dimensional [4] and metric spaces allowing the definition of metrics and distances [5, 6]. In the context of fractals, this may be seen as the projection of fractal objects [7, 8] in terms of dimensional shadowing [9].

What follows will be a generalization of relativity on geometric structures such as fractals. Thereby, these geometric structures can be interpreted by abstract forms of ‘ether’. Relativity did not abandon the ether, but rather redefined it as a property of space-time itself. According to Einstein [10, 11], ether is not the same as the classical ‘mechanical, ponderable’ Hertzian ether that was assumed to be a mechanical substrate consisting of microscopic constituents that served

as a medium for electromagnetic waves. While Maxwell and Lorentz had already succeeded in eliminating certain mechanical attributes associated with the Hertzian ether, they still retained its immobility. The ether of Einstein is not an immobile mechanical substance or a kind of ponderable entity, but a geometrical structure, devoid of any definite state of (mechanical) motion, that can be curved by the presence of matter and energy. Therefore, one may say that relativity did not abandon the ether, but transformed it into a more abstract and general concept.

It is most instructive to consider Einstein’s metaphor for this situation: Consider waves on the surface of water. Two distinct aspects can be described in this process. Firstly, one can observe how the wavelike interface between water and air changes over time. Alternatively, using small floating bodies, one can track the shifting positions of individual water particles over time. However, without any means to operationalize such floating entities to trace fluid particle motion fundamentally, and if only the temporally changing position of the space occupied by water were noticeable, intrinsic, embedded observers would have no reason to assume that water consists of moving particles. Nevertheless, we could still refer to it as a medium—or ether. It is this nonmechanical ether, devoid of any (operational) microphysical property that Einstein promotes for the propagation of electromagnetic waves. In particular, one has to abstain ascribing a state of motion to (constituents of) the ether. Moreover, in Einstein’s general theory of relativity, space possesses physical qualities, implying the existence of a certain type of ether. Without this ether, there would be no light propagation, measures, clocks, or spatial-temporal distances. However, this ether differs from ponderable media, lacking traceable parts through time (such as floating bodies on water) and not subject to the concept of motion.

Quantum field theory challenged this view insofar as it ascribed a certain ‘ponderability’ to the vacuum state. In 1948, Casimir and Polder suggested that London-Van der Waals interactions could give rise to forces between electrically neutral objects, particularly between a neutral atom and a conducting plane, or two neutral atoms [12], or between parallel conducting planes [13]. Such dispersion forces have an alternative characterization in terms of quantum (vacuum) fluctuations, using the methods of zero-point energy [14–18].

Independently, Dirac [19] pointed out that the quantum vacuum “is no longer a trivial state, but needs elaborate mathe-

\* [svozil@tuwien.ac.at](mailto:svozil@tuwien.ac.at); <http://tph.tuwien.ac.at/~svozil>

matics for its description.” (In 1982, I had the opportunity to inquire personally with Dirac about this issue. At that time, Dirac was emphasizing that all he cared about was the electron equation.)

Subsequently, Sakharov observed that in general relativity, spacetime action is postulated to depend on curvature, resulting in ‘metrical elasticity’ that generates forces resisting space curvature [20–25]. He suggests that this action can be linked to changes in quantum vacuum fluctuations when space is curved, viewing metrical elasticity as a level displacement effect. Beyond the Casimir-Polder forces, Sakharov proposes that gravitation—that is, the apparent curvature of space-time—arises from the elasticity of the vacuum, originating from the quantum vacuum fluctuations of interacting electromagnetic and other fields. Therefore, gravity is a secondary, ‘emergent’ interaction, akin to how hydrodynamics or continuum elasticity theory emerges from molecular physics [26]. Indeed, recent indirect [27] methods, as well as direct comparisons [28] of the propagation speeds of electromagnetic and gravitational waves, consistently indicate that this speed is identical to the speed of light in a vacuum, suggesting a common origin.

From the onset the elasticity theory of solids has been linked to the formalism of the general theory of relativity [29, 30]. A geometrical approach to the theory of structural defects in solids by (four-dimensional) continuum mechanics [31–47] effectively [48] encodes defects in terms of elastoplasticity. This formalism is tensor based. The operational, intrinsic viewpoint of embedded observers is emphasized. For instance, Kröner states [38] that “lengths are measured and atoms identified by counting lattice steps in the three crystallographic directions, then applying Pythagoras’ theorem  $ds^2 = g_{kl} dx^k dx^l$ , where  $ds$  is the distance of two atoms with relative position  $dx^k$ . ...  $ds$  ... is not the distance obtained by an external observer by means of a constant scale, but is, rather, the distance found by an internal observer with the help of the counting procedure.”

The aforementioned analogy between general relativity and the elasticity theory of solids has led to speculations that dark matter is a solid [49]. Kleinert even suggested that a general relativity-type ‘crystal gravity’ can be derived for a ‘world crystal’ with defects [40, 50–52]. In this analogy, the conserved defect tensor can be identified with the Einstein curvature tensor. The fourth (time) dimension enters because of the dynamics: the movement of defects and the change of the crystal’s plastic state [41].

For the sake of demonstrating the analogy between geometry and solid-state defects [30], suppose you are given a flat paper representing a two-dimensional spatial manifold, with a circle on it. If you cut out a section of the paper and fold the remaining paper into a cone, this folding introduces curvature, thereby reducing the circle’s circumference. We can also expect that any geodesics (for a flat paper, it is just a straight line) between two points (remaining and not inside the section that has been cut out) will shrink with decreasing opening angle of the cone.

This folding operation takes place in, or requires, a third dimension. Any attempt to force or crush the paper cone onto

a surface confined to a flat two-dimensional plane will exert a force on the paper cone, eventually tearing apart the paper fabric due to its flatness.

However, suppose that the two-dimensional surface is capable of adapting to the shape of the cone. Confinement by flatness can, for instance, be overcome by injecting or inserting additional ‘surface stuff’ into the two-dimensional surface, which might thereby reproduce the cone shape to a certain degree (without a perfect cone dip). This ‘added stuff’ is akin to a disclination in solid-state physics.

This metaphor can be perceived from an extrinsic, three-dimensional perspective, as we did at first. But for flatlanders [53] embedded in ‘cone-world,’ they would experience a need to explain their world in terms of nonflat, curved geometry.

In the next step, let us invert the question, as well as the role of the cone and the two-dimensional surface: So far, we have considered *adding* stuff to an existing flat manifold to resolve confinement. What about removing stuff from this manifold, thereby creating vacancies? In particular, consider the case of the Sierpinski carpet or the Menger Sponge. What kind of intrinsic operational phenomenology would embedded observers experience?

An intuition can be obtained from Kröner’s aforementioned observation that, embedded observers measure lengths ‘by counting lattice steps’. Indeed, if we need to ‘add’ stuff, like in the case to accommodate the cone, then we need to ‘subtract’ stuff to adapt a fractal object like Sierpinski carpet. In the former case intrinsic observers take ‘longer’ to traverse a region inhabited by a cone, whereas in the latter case they seem to take a ‘shortcut’.

## II. SEMI-QUANTITATIVE ANALYSIS

In what follows we shall approximate space-time metrics of fractals—or rather, approximations of fractals where scaling has not been performed in the limit—and then construct the associated Ricci tensor, which is a measure of how a volume in a ‘curved’ space differs from a volume in Euclidean space. We shall argue that, unlike normal matter and normal energy, this gives rise to negative curvature, which is a signature for ‘anti-gravity’ in the sense that it effectively describes a repulsive force.

When talking about anti-gravity it needs to be acknowledged that the formalism of general relativity does not outrightly disallow it [54]: The Einstein equations  $G_{ij} = \kappa T_{ij}$ , where  $G_{ij}$  denotes the Einstein curvature tensor,  $T_{ij}$  the stress-energy tensor (the source of the gravitational field), and  $\kappa$  is the gravitational constant, connect energy (density) to (local) geometry. Indeed, it is only after making some (physically motivated) assumptions regarding the stress-energy tensor  $T_{ij}$ , that the geometry of space-time  $G_{ij}$  reflects an ‘attractive gravitational force’. If  $T_{ij}$  is ‘exotic’ this could give rise to all sorts of ‘strange’ geometries  $G$ , including ‘anti-gravity’. It is not totally unreasonable to speculate that quantum field theory, in particular, vacuum fluctuations (beyond repulsive Casimir or London-Van der Waals forces that appear not to be

mediated by geometry [55]), may provide such ‘exotic’ states of the vacuum or matter (density) [56–58].

The metric or geodesic distance  $d_F(p, q)$  between two points  $p, q$  on a fractal set (or geodesic mask [59, Chapter 7]) of point defects  $F$  such as the Cantor set, the Sierpinski carpet or the Menger Sponge is the minimum of the length  $L$  of the (therefore, ‘shortest’) path or chain [60, 61]  $P = (p_1, p_2, \dots, p_l)$  joining  $p$  and  $q$  and included in  $F$ , that is,  $d_F(p, q) = \min \{L(P) | p_1 = p, p_l = q, \text{ and } P \subset F\}$ .

At that point it should be emphasized that we are not interested in ‘extrinsic’ observables such the average distance [62], or the geodesics in the Sierpinski Carpet and Menger Sponge [60, 61, 63–65]. We will be interested in intrinsic metrics for imbedded observers. To this end we shall combine ‘crystal space-time’ [41, Equation (1.6)] of vacancies with density (vacancy per cell of a space crystal  $F$ )  $N_v$ , as well as interstitial (additional ‘stuff’) density  $N_i$  with an idealized metric tensor  $g_{ij}$  in its most elementary diagonal form [39, Equations (3,4)]

$$g = \text{diag} \left[ -1, (1 - N_v + N_i)^{\frac{2}{3}}, (1 - N_v + N_i)^{\frac{2}{3}}, (1 - N_v + N_i)^{\frac{2}{3}} \right]. \quad (1)$$

The exponent  $2/3$  can be motivated by noticing that the volume of an infinitesimal parallelepiped can be written in terms of the modulus of the Jacobian determinant, which in turn can be written as the square root (the positive inverse of the square) of the modulus of the determinant of the metric tensor. Therefore, integration over these infinitesimal parallelepipeds yield an intrinsic volume that is proportional to the density  $1 - N_v + N_i$ .

At the lowest scale resolution, for the Menger Sponge,  $N_i = 0$  and  $N_v = 20/3^3 \approx 0.74$ . This vacancy density gets larger as the resolution increases, and in the limit approaches unity: ‘almost all stuff is lacking’. Let us, for the sake of the argument, suppose that we are not dealing with ‘added interstitial stuff’, so from now on,  $N_i = 0$ .

If the metric is not modelled by a uniform continuum of vacancies as in (1) but by a ‘one off’ localized vacancy, we could, in spherical coordinates  $\{t, r, \theta, \phi\}$ , make the Ansatz

$$g = \text{diag} \left[ -1, (1 - N_v)^{\frac{2}{3}} e^{-r^2}, r^2, r^2 \sin^2 \theta \right]. \quad (2)$$

The metric tensor  $g_{ij}$  in Equation (2) determines the geometry, and also the Einstein curvature scalar  $G = g^{ij} G_{ij}$  via the Ricci tensor and the Ricci scalar. This computation yields the Ricci scalar  $R$

$$G = R = - \frac{2e^{-r^2} \left[ (1 - N_v)^{\frac{2}{3}} e^{r^2} + 2r^2 - 1 \right]}{(1 - N_v)^{\frac{2}{3}} r^2}, \quad (3)$$

which, for small  $r \ll 1$ , is of the order of

$$G = O \left[ \frac{1 - (1 - N_v)^{\frac{2}{3}}}{(1 - N_v)^{\frac{2}{3}} r^2} \right]. \quad (4)$$

With respect to a frame aligned with the motion of perfect fluids or massive scalar fields the stress-energy tensor  $T^{ij}$  is of the form

$$T = \begin{pmatrix} \rho & 0 & 0 & 0 \\ 0 & p & 0 & 0 \\ 0 & 0 & p & 0 \\ 0 & 0 & 0 & p \end{pmatrix}, \quad (5)$$

and thus

$$T^i_j = g_{kj} T^{ik} \equiv \begin{pmatrix} -\rho & 0 & 0 & 0 \\ 0 & (1 - N_v)^{\frac{2}{3}} p & 0 & 0 \\ 0 & 0 & (1 - N_v)^{\frac{2}{3}} p & 0 \\ 0 & 0 & 0 & (1 - N_v)^{\frac{2}{3}} p \end{pmatrix}, \quad (6)$$

where  $\rho$  is a density and  $p$  is a pressure. Tracing this out, and inserting both  $G$  and  $T = T^i_i$  into the (traced-out) Einstein equation yields

$$T = -\rho + 3(1 - N_v)^{\frac{2}{3}} p = \frac{1}{\kappa} O \left[ \frac{1 - (1 - N_v)^{\frac{2}{3}}}{(1 - N_v)^{\frac{2}{3}} r^2} \right]. \quad (7)$$

For a particle of mass  $m$  at rest,  $\rho$  can be identified with  $m$  and  $p$  vanishes, so that Equation (7), with  $(1 - N_v)^{\frac{2}{3}} \approx 1 - 2N_v/3$ , reduces to

$$m = -\frac{1}{\kappa} O \left[ \frac{1 - (1 - N_v)^{\frac{2}{3}}}{(1 - N_v)^{\frac{2}{3}} r^2} \right] \approx -\frac{1}{\kappa} O \left[ \frac{N_v}{(1 - \frac{2}{3} N_v) r^2} \right], \quad (8)$$

which is negative for  $0 < N_v < 1$ .

### III. DISCUSSION

This estimation does not propose any physically feasible method to induce vacancies in the vacuum. As mentioned earlier, considerations related to Casimir-type quantum field effects may be relevant in this context.

Furthermore, the analysis has not leveraged the Menger Sponge’s analytic structure in the limit where, metaphorically speaking, space becomes ‘thinned out’ to an extent that renders it transparent to motion. In such a scenario, distances measured in terms of Lebesgue measures may approach vanishing values.

Nevertheless, the argument presented here aims to suggest that, in a physical—even ponderable, to use Einstein’s terminology of the time—model or representation of space-time, anti-gravity may emerge quite naturally. This conceptualization revolves around the notion of ‘thinning out’ rather than ‘adding stuff’. Consequently, the equivalence principle remains applicable, as we are not referring to interactions among individual particles or quanta, but rather to the local geometry of space-time as inherently perceived by embedded observers.

### ACKNOWLEDGMENTS

The research of K. Svozil was funded in whole, or in part, by the Austrian Science Fund (FWF), Project No. I 4579-N.

- [1] T. Toffoli, in *Applied General Systems Research: Recent Developments and Trends*, edited by G. J. Klir (Plenum Press, Springer US, New York, London, and Boston, MA, 1978), pp. 395–400, ISBN 978-1-4757-0555-3, URL [https://doi.org/10.1007/978-1-4757-0555-3\\_29](https://doi.org/10.1007/978-1-4757-0555-3_29).
- [2] K. Svozil, in *Inside versus Outside*, edited by H. Atmanspacher and G. J. Dalenoort (Springer, Berlin Heidelberg, 1994), vol. 63 of *Springer Series in Synergetics*, pp. 273–288, ISBN 978-3-642-48649-4, URL [https://doi.org/10.1007/978-3-642-48647-0\\_15](https://doi.org/10.1007/978-3-642-48647-0_15).
- [3] A. Zeilinger and K. Svozil, *Physical Review Letters* **54**, 2553 (1985), URL <https://doi.org/10.1103/PhysRevLett.54.2553>.
- [4] K. Menger, *The American Mathematical Monthly* **50**, 2 (1943), URL <https://doi.org/10.1080/00029890.1943.11991313>.
- [5] R. E. Hodel, in *Topology Conference*, edited by R. F. Dickman and P. Fletcher (Springer Berlin Heidelberg, Berlin, Heidelberg, 1974), pp. 120–136, ISBN 978-3-540-37948-5, URL <https://doi.org/10.1007/BFb0064018>.
- [6] J.-i. Nagata, *A survey of metrization theory* (Cambridge University Press, Cambridge, 1985), pp. 113–126, London Mathematical Society Lecture Note Series, URL <https://doi.org/10.1017/CBO9781107359925.005>.
- [7] J. M. Marstrand, *Proceedings of the London Mathematical Society* **s3-4**, 257 (1954), URL <https://doi.org/10.1112/plms/s3-4.1.257>.
- [8] K. J. Falconer, J. M. Fraser, and X. Jin, *Sixty years of fractal projections* (Birkhäuser Basel, 2015), vol. 70 of *Progress in Probability*, pp. 3–25, ISBN 978-3-319-18659-7, 978-3-319-18660-3, URL [https://doi.org/10.1007/978-3-319-18660-3\\_1](https://doi.org/10.1007/978-3-319-18660-3_1).
- [9] K. Svozil, *Journal of Physics A: Mathematical and General* **19**, L1125 (1986), URL <https://doi.org/10.1088/0305-4470/19/18/002>.
- [10] A. Einstein, *Äther und Relativitätstheorie. Rede gehalten am 5. Mai 1920 an der Reichs-Universität Leiden* (Springer, Berlin, 1920), URL <https://doi.org/10.1007/978-3-642-64927-1>.
- [11] A. Einstein, *Eiher and the Theory of Relativity* (Springer Dordrecht, Dordrecht, Netherlands, 2007), vol. 250 of *Boston Studies in the Philosophy and History of Science*, pp. 1537–1542, URL [https://doi.org/10.1007/978-1-4020-4000-9\\_34](https://doi.org/10.1007/978-1-4020-4000-9_34).
- [12] H. B. G. Casimir and D. Polder, *Physical Review* **73**, 360 (1948), URL <https://doi.org/10.1103/PhysRev.73.360>.
- [13] H. B. G. Casimir, *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen* **51**, 793 (1948), URL <https://dwc.knaw.nl/DL/publications/PU00018547.pdf>.
- [14] H. B. G. Casimir, *Journal de Chimie Physique* **46**, 407 (1949), ISSN 0021-7689, URL <https://doi.org/10.1051/jcp/1949460407>.
- [15] E. M. Lifshitz, *Journal of Experimental and Theoretical Physics (JETP)* **2**, 73 (1956), Russian original - ZhETF, Vol. 29, No. 1, p. 94, January 1956, URL [http://jetp.ras.ru/cgi-bin/dn/e\\_002\\_01\\_0073.pdf](http://jetp.ras.ru/cgi-bin/dn/e_002_01_0073.pdf).
- [16] T. A. Roman, *Physical Review D* **33**, 3526 (1986), URL <https://doi.org/10.1103/physrevd.33.3526>.
- [17] P. W. Milonni, *The Quantum Vacuum. An Introduction to Quantum Electrodynamics* (Academic Press, San Diego, 1994), ISBN 978-0-08-057149-2, URL <https://doi.org/10.1016/C2009-0-21295-5>.
- [18] P. W. Milonni, *An introduction to quantum optics and quantum fluctuations* (Oxford University Press, Oxford, United Kingdom, 2019), ISBN 978-0-19-921561-4, URL <https://doi.org/10.1093/oso/9780199215614.001.0001>.
- [19] P. A. M. Dirac, *Nature* **168**, 906 (1951), URL <https://doi.org/10.1038/168906a0>.
- [20] A. D. Sakharov, *General Relativity and Gravitation* **32**, 365 (2000), ISSN 0001-7701, translated from *Doklady Akademii Nauk SSSR*, vol. 177, No. 1, pp. 70-71, November 1967, URL <https://doi.org/10.1023/A:1001947813563>.
- [21] H. E. Puthoff, *Physical Review A* **39**, 2333 (1989), URL <https://doi.org/10.1103/physreva.39.2333>.
- [22] B. Haisch, A. Rueda, and H. E. Puthoff, *Physical Review A* **49**, 678 (1994), URL <https://doi.org/10.1103/PhysRevA.49.678>.
- [23] B. Haisch, A. Rueda, and Y. Dobyms, *Annalen der Physik* **10**, 393 (2001), arXiv:gr-qc/0009036, URL [https://doi.org/10.1002/1521-3889\(200105\)10:5<393::](https://doi.org/10.1002/1521-3889(200105)10:5<393::)
- [24] A. Rueda and B. Haisch, *Annalen der Physik* **14**, 479 (2005), <https://onlinelibrary.wiley.com/doi/pdf/10.1002/andp.200510147>, URL <https://doi.org/10.1002/andp.200510147>.
- [25] E. W. Davis, V. L. Teofilo, B. Haisch, H. E. Puthoff, L. J. Nickisch, A. Rueda, and D. C. Cole, *AIP Conference Proceedings* **813**, 1390 (2006), URL <https://doi.org/10.1063/1.2169324>.
- [26] M. Visser, *Modern Physics Letters A* **17**, 977 (2002), URL <https://doi.org/10.1142/S0217732302006886>.
- [27] N. Cornish, D. Blas, and G. Nardini, *Physical Review Letters* **119**, 161102 (2017), URL <https://doi.org/10.1103/PhysRevLett.119.161102>.
- [28] B. P. Abbott and et al, *The Astrophysical Journal Letters* **848**, L13 (2017), URL <https://doi.org/10.3847/2041-8213/aa920c>.
- [29] H. Schaefer, *Zeitschrift für angewandte Mathematik und Mechanik* (Journal of Applied Mathematics and Mechanics) **33**, 356 (1953), vorgetragen auf der Jahrestagung der GaMM in Braunschweig 1952, URL <https://doi.org/10.1002/zamm.19530331006>.
- [30] J. Zaanen, F. Balm, and A. J. Beekman, *SciPost physics* **13** (2022), arXiv:2109.11325, URL <https://doi.org/10.21468/scipostphys.13.2.039>.
- [31] E. Kröner, *Kontinuumstheorie der Versetzungen und Eigenspannungen*, vol. 5 of *Ergebnisse der angewandten Mathematik* (Springer-Verlag OHG, Berlin, Heidelberg, Germany, 1958), ISBN 978-3-540-02261-9, URL <https://archive.org/details/kontinuumstheori0000dre>.
- [32] E. Kröner, *Archive for Rational Mechanics and Analysis* **4**, 273 (1959), URL <https://doi.org/10.1007/bf00281393>.
- [33] A. M. Kosevich, *Journal of Experimental and Theoretical Physics (JETP)* **15**, 108 (1962), Russian original - J. Exptl. Theoret. Phys. (U.S.S.R.) (ZhETF), Vol. 42, No. 1, p. 152-162, January 1962, URL <http://www.ippt.pan.pl/repository/open/o6097.pdf>.
- [34] L. Turski, *The Bulletin of the Polish Academy of Sciences Technical Sciences* **14**, 289 (1966).
- [35] E. Kröner, *International Journal of Solids and Structures* **3**, 731 (1967), URL [https://doi.org/10.1016/0020-7683\(67\)90049-2](https://doi.org/10.1016/0020-7683(67)90049-2).

- [36] E. Kröner and K. Anthony, *Annual Review of Materials Science* **5**, 43 (1975), URL <https://doi.org/10.1146/annurev.ms.05.080175.000375>.
- [37] E. Kossecka and R. de Wit, *Archives of Mechanics* **29**, 633 (1977), URL <http://www.ippt.pan.pl/repository/open/o6097.1544>.
- [38] E. Kröner, **21**, 747 (1985), URL [https://doi.org/10.1016/0020-7683\(85\)90077-0](https://doi.org/10.1016/0020-7683(85)90077-0).
- [39] E. Kröner, *International Journal of Theoretical Physics* **29**, 1219 (1990), URL <https://doi.org/10.1007/bf00672933>.
- [40] E. Kröner, *International Journal of Solids and Structures* **38**, 1115 (2001), URL [https://doi.org/10.1016/s0020-7683\(00\)00077-9](https://doi.org/10.1016/s0020-7683(00)00077-9).
- [41] S.-i. Amari, *RAAG Memoirs of the Unifying Study of Basic Problems in Engineering and Physical Sciences by Means of Geometry IV, Division D*, 142 (1968), URL <https://bsi-ni.brain.riken.jp/database/file/475833.pdf>.
- [42] H. Günther, *physica status solidi (b)* **49**, 551 (1972), URL <https://doi.org/10.1002/pssb.2220490217>.
- [43] H. Günther, *Gerlands Beiträge zur Geophysik* **88**, 23 (1979).
- [44] H. Günther, *International Journal of Engineering Science* **19**, 1799 (1981), URL [https://doi.org/10.1016/0020-7225\(81\)90170-1](https://doi.org/10.1016/0020-7225(81)90170-1).
- [45] H. Günther, *Annalen der Physik* **495**, 220 (1983), URL <https://doi.org/10.1002/andp.19834950406>.
- [46] A. Golebiewska-Lasota, *International Journal of Engineering Science* **17**, 329 (1979), URL [https://doi.org/10.1016/0020-7225\(79\)90095-8](https://doi.org/10.1016/0020-7225(79)90095-8).
- [47] A. Golebiewska-Lasota and D. G. B. Edelen, *International Journal of Engineering Science* **17**, 335 (1979), URL [https://doi.org/10.1016/0020-7225\(79\)90096-x](https://doi.org/10.1016/0020-7225(79)90096-x).
- [48] P. W. Anderson, *Science* **177**, 393 (1972), URL <https://doi.org/10.1126/science.177.4047.393>.
- [49] M. Bucher and D. Spergel, *Physical Review D* **60**, 043505 (1999), URL <https://link.aps.org/doi/10.1103/PhysRevD.60.043505>.
- [50] H. Kleinert, *Annalen der Physik* **499**, 117 (1987), URL <https://doi.org/10.1002/andp.19874990206>.
- [51] H. Kleinert, *General Relativity and Gravitation* **32**, 769 (2000), URL <https://doi.org/10.1023/a:1001962922592>.
- [52] H. Kleinert and J. Zaanen, *Physics Letters* **324**, 361 (2004), URL <https://doi.org/10.1016/j.physleta.2004.03.048>.
- [53] E. Abbott, *Flatland: a romance of many dimensions. Second and Revised Edition* (Van Nostrand Reinhold Company, 1884), URL <http://www.gutenberg.org/etext/201andhttp://www.geon>.
- [54] P. Martín-Moruno and M. Visser, *Classical and Semi-Classical Energy Conditions* (Springer International Publishing, Cham, Switzerland, 2017), pp. 193–213, arXiv:1702.05915, URL [https://doi.org/10.1007/978-3-319-55182-1\\_9](https://doi.org/10.1007/978-3-319-55182-1_9).
- [55] J. N. Munday, F. Capasso, and V. A. Parsegian, *Nature* **457**, 170 (2009), URL <https://doi.org/10.1038/nature07610>.
- [56] J. D. Bekenstein, *Physical review* **88** (2013), URL <https://doi.org/10.1103/physrevd.88.125005>.
- [57] E.-A. Kontou and K. Sanders, *Classical and Quantum Gravity* **37**, 193001 (2020), URL <https://doi.org/10.1088/1361-6382/ab8fcf>.
- [58] A. Costa and G. E. A. Matsas, *Physical Review D* **105** (2022), URL <https://doi.org/10.1103/physrevd.105.085016>.
- [59] P. Soille, *Morphological image analysis. Principles and Applications* (Springer Berlin Heidelberg, Berlin, Heidelberg, 2004), 2nd ed., ISBN 978-3-540-42988-3, 978-3-642-07696-1, 978-3-662-05088-0, URL <https://doi.org/10.1007/978-3-662-05088-0>.
- [60] J. Kigami, *Geometry and analysis of metric spaces via weighted partitions* (Springer Nature Switzerland AG, Cham, Switzerland, 2020), URL <https://doi.org/10.1007/978-3-030-54154-5>.
- [61] Q. Gu, K. Lau, H. Qiu, and H.-J. Ruan, *Science China Mathematics* **66**, 907 (2023), URL <https://doi.org/10.1007/s11425-021-1989-3>.
- [62] A. M. Hinz and A. Schief, *Probability Theory and Related Fields* **87**, 129 (1990), URL <https://doi.org/10.1007/bf01217750>.
- [63] E. Berkove and D. A. Smith, *Fractals* **28**, 2050120 (2020), URL <https://doi.org/10.1142/s0218348x20501200>.
- [64] L. L. Cristea, *Quaestiones Mathematicae* **28**, 251 (2005), URL <https://doi.org/10.2989/16073600509486126>.
- [65] J. Kigami, *Analysis on Fractals*, vol. 143 of *Cambridge Tracts in Mathematics* (Cambridge University Press, 2001), ISBN 0521793211, 9780521793216, URL <https://doi.org/10.1017/CBO9780511470943>.