# Resting-State fingerprints of Acceptance and Reappraisal. The role of Sensorimotor, Executive and Affective networks

Parisa Ahmadi Ghomroudi\*1, Roma Siugzdaite<sup>2</sup>, Irene Messina<sup>3</sup>, Alessandro Grecucci<sup>1,4</sup>

<sup>1</sup>DiPSCo – Department of Psychology and Cognitive Sciences, University of Trento, Rovereto, Italy

<sup>2</sup>Department of Experimental Psychology, Faculty of Psychology and Pedagogical Sciences, Ghent University, Ghent, Belgium

<sup>3</sup>Universitas Mercatorum, Rome, Italy

<sup>4</sup>CISMed – Center for Medical Sciences, University of Trento, Trento, Italy

## **Corresponding author:**

Parisa Ahmadi Ghomroudi

Department of Psychology and Cognitive Sciences,

University of Trento,

Corso Bettini, 84, 38068,

Rovereto, Italy

E-mail: p.ahmadighomroudi@unitn.it

Tel. +39 04 64 808302

*Keywords* Acceptance, Reappraisal, Emotion Regulation, resting state, functional connectivity.

#### Abstract

Acceptance and Reappraisal are considered adaptive emotion regulation strategies. While previous studies have explored the neural underpinnings of these strategies using task-based fMRI and sMRI, a gap exists in the literature concerning resting-state functional brain networks contributions to these abilities, especially for what concerns Acceptance. Another intriguing question is whether these strategies rely on similar or different neural mechanisms. Building on the well-known improved emotion regulation and increased cognitive flexibility of individuals who rely on acceptance, we expected to find decreased activity inside the Affective network and increased activity inside the Executive and Sensorimotor networks to be predicted of acceptance. We also expect that these networks may be associated at least in part with Reappraisal, indicating a common mechanism behind different strategies. To test these hypotheses, we conducted a functional connectivity analysis of resting-state data from 134 individuals (95 females; mean age:  $30.09 \pm 12.87$  years, mean education:  $12.62 \pm 1.41$ years). To assess acceptance and reappraisal abilities, we used the Cognitive Emotion Regulation Questionnaire (CERQ) and a group-ICA unsupervised machine learning approach to identify resting-state networks. Subsequently, we conducted backward regression to predict acceptance and reappraisal abilities. As expected, results indicated that acceptance was predicted by decreased Affective, and increased Executive, , and Sensorimotor networks, while reappraisal was predicted by an increase in the Sensorimotor network only. Notably, these findings suggest both distinct and overlapping brain contributions to acceptance and reappraisal strategies, with the Sensorimotor network potentially serving as a core common mechanism. These results not only align with previous findings but also expand upon them,

demonstrating the complex interplay of cognitive, affective, and sensory abilities in emotion regulation.

## Introduction

The ability to regulate emotions is considered fundamental to mental health and well-being and difficulties in regulating emotions have been associated with a wide range of psychological conditions<sup>1–5</sup> For example, anxiety, depression, and personality disorders have all been linked to emotion dysregulation<sup>6,7</sup>. Due to the prevalence of emotion regulation challenges across various psychological disorders, clinicians have started integrating different emotion regulation strategies into their therapeutic approaches <sup>4,8–10</sup>

Acceptance is characterized by open curiosity towards ongoing mental and sensory experiences <sup>11,12</sup>. It is considered a fundamental concept in third-wave behavioral therapies <sup>13,14</sup> and experiential-dynamic approaches<sup>4,5,15,16</sup>. Within these frameworks, acceptance is defined as "the active and ware embrace of private experiences without unnecessary attempts to change their frequency or form" <sup>17</sup>. Reappraisal and acceptance are considered two highly effective strategies frequently used in psychotherapy <sup>5,18,19</sup>. Reappraisal refers to a voluntary effort to reinterpret the significance of a situation to change its emotional effect <sup>20</sup>. This process is defined as "construing a potentially emotion-eliciting situation in non-emotional terms"<sup>21</sup>. Reappraisal is an antecedent-focused regulation strategy that alters emotion before the complete onset of emotional response.

Reappraisal and acceptance are commonly regarded as adaptive strategies due to their positive associations with well-being and mental health<sup>22,23</sup>. In a study conducted by Dan-Glauser and Gross<sup>24</sup>, when compared to no regulation, acceptance was found to lead to an increase in positive emotions and decrease in respiration rate. Additionally, reappraisal is believed to be negatively correlated with anxiety<sup>25–27</sup>. Uchida et al<sup>28</sup> showed individuals who were more successful in reappraisal had lower levels of trait anxiety and experienced more

positive emotions in their daily lives. Hofmann et al<sup>29</sup> reported effectiveness for both strategies in reducing heart rate when compared to suppression. Goldin et al<sup>12</sup> found no difference in respiration rate and skin conductance between both strategies but observed a higher heart rate during reappraisal compared to acceptance.

For what concerns the neural bases of acceptance, just a few task-based fMRI studies inquired into its nature. Traditional models of emotion regulation are based on top-down control processes <sup>30</sup>. However, neuroimaging studies exploring the neural correlates of acceptance show inconsistent findings. Some of these studies align with traditional models by demonstrating prefrontal activations within the dorsal attention network during acceptance <sup>12,31</sup>. Other studies report either less activity in prefrontal areas or activations that are more medially located compared to traditional strategies<sup>32–34</sup>. Furthermore, several studies reveal that neural correlates of acceptance may depend on bottom-up mechanisms and occur without the involvement of prefrontal cortical areas<sup>35–39</sup>. In an effort to provide a synthetic view of the neural contributions to acceptance, Messina et al<sup>39</sup> (2021) performed a meta-analysis of 13 fMRI studies that revealed a consistent association between acceptance and decreased brain activity in emotion related regions such as the posterior cingulate cortex (PCC)/precuneus, insula and limbic subcortical regions such as the thalamus and the parahippocampal gyrus, regardless of the control condition. In another study, Sezer et al <sup>40</sup> showed that mindfulness, an ability strongly correlated with acceptance, correlated with increased functional connectivity between the posterior cingulate cortex, a central part of the default mode network (DMN), and the dorsolateral prefrontal cortex, potentially enhancing attention control. They further found that mindfulness is associated with increased connectivity in areas implicated in pain relief and self-awareness, pointing to the multifaceted nature of these cognitive strategies. Of note, the studies included in these meta-analyses did not include information on the individual differences in the abilities to use acceptance, as they were taskbased fMRI studies. Besides task based functional studies, to our knowledge, only one study tried to understand the abilities in acceptance abilities measured via dedicated questionnaires<sup>41</sup>. In this study, a data fusion machine learning approach was used to identify joint gray and white matter contributions to high and low acceptance abilities. Results revealed individuals with higher acceptance ability showed reduced gray and white matter concentrations within the default mode network, particularly in posterior and anterior midline structures, anterior temporal regions, the angular gyrus, the dorsal anterior cingulate cortex (dACC), and the right insula. Additionally, these individuals exhibited increased gray and white matter concentrations in the cognitive and attention networks, especially within prefrontal and superior parietal regions. However, this study was limited to the structural properties of the brain and the question of whether similar networks, but at a functional level, may contribute to acceptance remain unaddressed.

Regarding reappraisal, Buhle et al<sup>42</sup> conducted a meta-analysis of 48 task-related fMRI studies and reported that both downregulation and upregulation of emotion are associated with increased activation in the bilateral dorsolateral and ventrolateral prefrontal cortex (dIPFC, vIPFC), dorsal anterior cingulate cortex (dACC), supplementary motor area (SMA), and the inferior/superior parietal cortex. Beside task-based fMRI studies of reappraisal, a few studies inquired into the nature of reappraisal abilities and how they can be predicted by resting state and structural networks. For example, Uchida et al<sup>28</sup> found that effective reappraisal correlates with decreased connectivity between the right amygdala and both the medial prefrontal cortex (DLPFC) and posterior visual regions during resting-state functional connectivity. Additionally, Morawetz et al<sup>43</sup> reported that the ability to reappraise is linked to stronger functional connectivity between the ventrolateral prefrontal cortex and the amygdala. In another study, Zanella et al<sup>44</sup> found that cognitive and positive reappraisal

were predicted by sensorimotor networks. Of note, positive reappraisal, differently from cognitive reappraisal can be seen as a hybrid form of reappraisal and acceptance, for which someone can attach a positive meaning to the event in terms of personal growth <sup>45</sup>. Further, from a structural point of view, Ghomroudi et  $al^{46}$  applied machine learning methods to gray matter structural data and found that a temporo-parahippocampal-orbitofrontal network, which includes regions such as the Thalamus, Parahippocampal Gyrus, Superior Temporal Gyrus, Lentiform Nucleus, Uncus, and Cerebellar Tonsil, predicts the use of reappraisal. These studies point toward the direction of different mechanisms behind acceptance and reappraisal with a possible partial overlap in insula and frontal regions of the brain such as dorsolateral prefrontal cortex, but also in regions of the sensorimotor network<sup>44</sup>. To test this hypothesis, a recent meta-analysis of 42 fMRI studies conducted by Monachesi et al<sup>47</sup>, was run and it was found that reappraisal was associated with increased activity in the superior frontal gyrus and the middle frontal gyrus, while reducing activity in sublobar regions like the globus pallidus and putamen. Conversely, acceptance was associated with increased activity in the claustrum and decreased activity in various limbic structures, including the posterior cingulate cortex (PCC)/precuneus, the parahippocampal gyrus, and the thalamus (pulvinar). Interestingly, the conjunction analysis revealed that both acceptance and reappraisal engage the VLPFC and the insula, indicating shared neural pathways in these emotional regulation strategies.

It was suggested that emotion regulation may rely on a core inhibitory (VLPFC) / sensorial awareness (insula) and on strategy-specific top-down and bottom-up processes distinct for different strategies. Although this study suggests different and only partially overlapping mechanisms behind acceptance and reappraisal usage (task-based fMRI studies), we still do not know whether the same applies to the resting state networks associated with the ability to apply these strategies. The aim of this study is to test for the first time the possibility to detect

resting state macro networks contributions to acceptance and reappraisal abilities, and to test the hypothesis that they rely on different mechanisms with a common core behind them. If demonstrated, this result may support the model of common and specific neural mechanisms of emotion regulation that expand the previous simplistic dual-routes models<sup>5</sup>.

Of note, in this study we used different approach to the previous resting-state studies on emotion regulation. Resting-state functional connectivity has been mainly estimated using seed-based correlation<sup>48</sup> in the field of emotion<sup>28,49</sup>. Seed-based functional connectivity analysis, also known as ROI-based functional connectivity, involves identifying brain regions that exhibit correlations with the activity in a predefined seed region. This analysis calculates cross-correlations between the time-series of the seed region and the rest of the brain. Seedbased analysis requires the a priori selection of seed regions. One advantage of seed analysis is its simplicity and intuitive interpretation. However, a drawback is its sensitivity to seed selection, as changing the seed region can significantly alter the results, making it susceptible to bias. The ICA method<sup>50,51</sup> is a whole-brain, model-free method that provides a more datadriven approach to quantifying functional connectivity. ICA is a computational approach that decomposes BOLD fMRI signal time courses from the entire brain into spatially and temporally independent components. Several resting-state networks typically emerge from ICA analysis in resting-state fMRI studies, including the default mode network, auditory network, salience network, executive control network, medial visual network, lateral visual network, sensorimotor cortex, dorsal visual stream (frontoparietal attention network), basal ganglia network, limbic network, and precuneus network. Unlike seed-based analysis, which relies on a priori assumptions and the selection of regions of interest, ICA is a data-driven method and can be executed without predefined assumptions, except for specifying the number of independent components to identify. In this study, resting-state macro-networks

were determined using an ICA approach. To the best of our knowledge, all previous research on emotion regulation involving resting-state analysis has used seed-based analysis<sup>28,49</sup>. Based on previous research<sup>12,31,39</sup>, we expect modulation in BOLD temporal variability across the following networks to predict acceptance ability. Firstly, we predict a decrease in BOLD temporal variability within a network that includes subcortical regions such as the thalamus and the hippocampus/parahippocampal region, which can be considered part of an affective network. Furthermore, an increase in BOLD temporal variability is expected within the executive network, encompassing cognitive control regions such as the DLPFC, VLPFC, and the dorsal anterior cingulate cortex. Additionally, based on Zanella et  $al^{44}$  resting state study and Monachesi et al<sup>52</sup> meta-analysis, we expect increased BOLD temporal variability within the somatosensory network, particularly within the insula and supplementary motor cortex (SMA), to predict acceptance ability. Building upon previous research<sup>44,53</sup>, we predict that increased temporal variability within the sensorimotor network is predictive of reappraisal. We hypothesize that regions such as the precentral and postcentral gyrus, along with the supplementary motor area, which play a pivotal role in the execution of regulatory functions, form a network predictive of reappraisal. Based on the meta-analysis conducted by Monachesi et al<sup>52</sup>, we expect that the sensorimotor network, particularly regions such as the insula, may act as a fundamental network underlying both reappraisal and acceptance strategies.

#### Results

### Macro networks contributions to acceptance

The Backward Multiple Regression analysis returned a significant winning model (F(4, 130) = 10.603,  $R^2 = 0.246$ , p < 0.001) in which the BOLD variability across four networks, IC2 ( $\beta$ 

= -60.85, p = 0.001), IC11 ( $\beta$  = -67.89, p = 0.001), IC13 ( $\beta$  = 80.16, p < 0.001), and IC18 ( $\beta$  = 24.42, p = 0.022) predicted acceptance ability, thereby yielding an overall significant model. The corresponding regression equation was as follows: Acceptance ability = 8.765 - 60.858 IC2 - 67.899 IC11 + 80.167 IC13 + 24.429 IC18. The identified ICs encompass clusters of both cortical and subcortical regions at cluster statistical significance level of p < 0.05 (pFDR corrected) and at the voxel significant level p < 0.001 (pFDR corrected). These networks correspond to well-established resting state networks, specifically: IC2: Associated with the affective system, IC11: Corresponding to the Executive network, IC13: Representative of the Sensorimotor network and IC18: Aligned partly with the language network.

- - - -

Please insert Figure 1

- - - -

- - - -

Please insert Figure 2

- - - -

- - - -

Please insert Figure 3

- - - -

- - - -

Please insert Figure 4

- - - -

- - - -

Please insert Table 1

- - - -

- - - -

Please insert Table 2

- - - -

#### Macro networks contributions to Reappraisal

Backward Multiple Regression analysis returned a significant winning model (F (1, 133) = 28.59,  $R^2 = 0.421$ , p < 0.001) in which the BOLD variability of IC20 ( $\beta = 0.339$ , p < 0.001), predicted reappraisal usage. The corresponding regression equation was as follows: Reappraisal usage = 6.757 + 0.339 IC20. IC20 encompasses a cluster of sensory motor regions at cluster statistical significance level of p < 0.05 (pFDR corrected) and at the voxel significant level p < 0.001 (pFDR corrected). this network correspond to the well-established Sensory motor resting-state network. The level of variability observed in a specific brain region is positively correlated with the degree of functional connectivity<sup>54</sup>. In other words, when a brain area is highly interconnected with other brain regions, it tends to exhibit increased variability in its BOLD signal. Greater BOLD variability corresponds to an increased frequency of both reappraisal and acceptance usage. Conversely, lower BOLD variability corresponds to a decreased frequency of both acceptance and reappraisal usage.

- - - -

Please insert Figure 5

- - - -

- - - -

Please insert Table 3

- - - -

#### Discussion

In this study, we applied group-ICA, an unsupervised machine learning technique, to identify the resting state networks that predcit acceptance and reappraisal, and eventually the commonalities between the two strategies. The findings indicated that modulations in four BOLD temporal variability networks were predictive of acceptance, while one network was predictive of reappraisal. Specifically, reduced BOLD temporal variability in the Affective (IC2) and Executive (IC11) networks, and increased BOLD temporal variability in the Sensorimotor (IC13) and part of the language (IC18) networks, were associated with predicting acceptance. Further, the ability to reappraise was predicted by an increase of variability inside the Sensorimotor network (IC20). Of note the sensorimotor network was associated with both acceptance and reappraisal, demonstrating a commonality between the two.

The results indicated that decreased BOLD temporal variability in IC2 network is associated with acceptance ability. Notably, several regions within the IC2 brain area overlap with the

affective network, which is crucial for emotional processing. Key regions in IC2 encompass the Parahippocampal Gyrus, Hippocampus, Amygdala, and Thalamus. Previous studies <sup>38,39,52</sup> have found a correlation between reduced activity in the Parahippocampal Gyrus and acceptance. The Parahippocampal Gyrus (PHG) is significant in the early and automatic assessment of emotional significance during emotion regulation<sup>55</sup>. Additionally, decreased connectivity in the Parahippocampal Gyrus has been noted during mindfulness and meditation practices<sup>56</sup>. This suggests that diminished PHG activity during acceptance may reflect a reduced impact of emotional events on the individual, potentially influencing memory associations or the retrieval of stimuli<sup>52,57</sup>. The thalamus, characterized as a critical relay hub in the brain for processing sensory information<sup>58</sup>. Its deactivation might signify a sensory filtering process, leading to enhanced openness and a non-judgmental attitude inherent in acceptance<sup>59</sup>. Furthermore, the Thalamus, along with the Hippocampus and Parahippocampal Gyrus, contributes to regulation efficacy<sup>39,60,61</sup>. Overall these results align with the observation that individuals who rely on acceptance have a better regulation of emotions<sup>62,63</sup>.

The findings showed a reduction in BOLD temporal variability within the frontoparietal network, also name Central Executive network (IC11), which correlates with the acceptance ability. This network includes the Superior Frontal Gyrus and the Middle Frontal Gyrus. This result is aligned<sup>37,64</sup>, the decreased activity in Middle Frontal Gyrus might indicate an enhanced capacity to maintain attention and regulate impulses during acceptance<sup>64</sup>. This result confirms that individuals who practice acceptance may display increased cognitive abilities and enhanced flexibility. Moreover, they demonstrate improved attentional focus to execute attention tasks and better impulse control<sup>41,65.</sup>

The results revealed that increased BOLD temporal variability in Somatosensory networks (IC13) is associated with the acceptance ability. This network consists of Precentral Gyrus

and Postcentral Gyrus. The postcentral gyrus and the insula are associated with interoceptive awareness<sup>66–68</sup>. In addition, staying in the present moment a key in mindfulness, is linked to a brain network that includes the thalamus, insula, and sensorimotor regions such as post and precentral gyrus<sup>69,70</sup>. This result is consistent with the fact that individuals using acceptance may have a coherent perception of their emotional states, fostering a non-judgmental attitude towards emotional experiences. Moreover, it highlights the significance of interoceptive awareness, emphasizing the importance of remaining present, a fundamental aspect of acceptance<sup>39,41,64</sup>.

Last but not least, results revealed that increased BOLD temporal variability within IC18, recognized by CONN as the language networks, which include the Ventrolateral Prefrontal Cortex (VLPFC) and the insula, is associated with the ability to accept. The VLPFC plays a role in various functions, such as response selection and inhibition, as well as language processing<sup>71–73</sup>. Several studies suggest that both the insula and the VLPFC are consistently linked to effective emotion regulation across different strategies, including acceptance<sup>39,61,74</sup>. The insula plays a critical role in integrating sensory information from both internal and external sources, helps in the formation of an awareness of the body's emotional state and labeling of the emotion<sup>75,76</sup>, and was previously found in meta-analyses on this topic <sup>39</sup>. In addition to language-related regions, this network includes the dorsolateral prefrontal cortex (DLPFC) and the anterior cingulate cortex (ACC), which are more associated with control mechanisms. The DLPFC is a crucial area in the central executive network, playing significant roles in attention, decision-making, working memory, and cognitive control<sup>77–79</sup>. Moreover, the dorsal anterior cingulate cortex (dACC) is related to the regulation of cognitive processes<sup>80–82</sup> and, is also linked to increased connectivity in mindful individuals<sup>70,83</sup>. A few studies examining resting state connectivity indicate that DLPFC and VLPFC are effectively

interconnected during emotion regulation<sup>43,84</sup>. Increased activity in executive network regions such as the DLPFC, VLPFC, and ACC, along with decreased activity in affective network regions like the amygdala, supports a dual-process model<sup>85–87</sup>.

The result indicates the increase BOLD variability in another somatosensory network (IC 20) is predictive of reappraisal ability. The somatosensory network including regions such as Precentral Gyrus, Postcentral Gyrus, Supplementary Motor Area, Insula. Somatosensory cortex is involved in the recognition of emotions, the understanding of the emotional states of others<sup>88</sup>. Picó-Pérez et al<sup>52 53</sup> revealed the significant role of the Supplementary Motor Area (SMA) in cognitive reappraisal. The key role of SMA is top-down inhibitory control over the amygdala, whose role is in the initial processing and linking of sensory and affective input<sup>89</sup>. The insula plays a crucial role in integrating sensory information from both internal and external environments. This integration is essential for forming a coherent and conscious representation of one's internal emotional state<sup>63,76</sup> Moreover, functional connectivity of the left insula, SMA is associated with frequency of use of reappraisal<sup>53</sup>. The dorsal anterior cingulate cortex (dACC) and superior parietal cortex are linked to goal-oriented attention<sup>90,91</sup>. Specifically, superior parietal cortex is involved in detecting salience and directing attention <sup>92</sup> and the dACC plays a role in control allocation<sup>93</sup> The results of this study suggest that the Sensorimotor network functions as the common core network underlying both acceptance and reappraisal strategies. Considering that emotion regulation involves paying attention to and being aware of one's emotional state, it is plausible to associate this process with the awareness of bodily states<sup>94</sup>. The somatosensory cortex is pivotal in emotional processing, the generation of emotional states, and interoceptive awareness<sup>95–97</sup> Since regulating emotions requires an awareness of both emotional and bodily states, it follows that increased interoceptive awareness could enhance emotion regulation. This enhancement could be through the improved detection of early bodily reactions to emotional stimuli <sup>94</sup>. Therefore,

due to the role of the somatosensory network in awareness, it could be a common core in emotion regulation, regardless of the specific strategy used.

To conclude, this study aimed to identify specific resting state functional brain networks that are predictive of acceptance and reappraisal capabilities. Acceptance and reappraisal are both recognized as effective emotion regulation strategies, as highlighted in several research studies<sup>5,23,98,99</sup>. These strategies are frequently used in psychotherapy<sup>18</sup>. While numerous studies have explored the brain mechanisms behind acceptance and reappraisal in task-based fMRI and sMRI techniques, there remains a lack of understanding about the resting state functional brain networks that underlie these abilities. Our findings indicate that Acceptance was predicted by the Executive, the Affective and the Sensorimotor networks, whereas reappraisal was predicted by the Sensorimotor network. These findings not only are inline but also extend previous findings, revealing interaction among cognitive, emotional, and sensory processes in emotion regulation. Additionally, our results suggest unique and shared neural contributions to acceptance and reappraisal, with the Sensorimotor network appearing to be a crucial shared element in both strategies.

### Method

#### **Participants**

Participants included in this study comprised 134 (95 female) native German speaker individuals, with a mean age of  $30.09 \pm 12.87$  and an average of  $12.62 \pm 1.41$  years of education. The participants' mean acceptance score was  $7.06 \pm 2.85$ , and the mean reappraisal score was  $6.78 \pm 2.84$ . The data was drawn from "Leipzig study for mind-body-emotion interactions" (OpenNeuro database, accession number ds000221) (LEMON). The data collection was conducted at the Max Planck Institute for Human Cognitive and Brain Sciences (MPI CBS) in Leipzig <sup>100</sup>The inclusion criteria were, absence of cardiovascular, psychiatric, neurological disorders, and malignant diseases, as well as the non-use of certain medications. Individuals reporting drug or excessive alcohol use were excluded. Participants provided written informed consent and agreed to anonymous data sharing. Compensation was provided upon completion of all assessments. A power analysis conducted in R aimed to determine the required sample size for the multiple regression analysis. This analysis was conducted based on the following parameters: a number of predictors of 20, an effect size of 0.2 (Cohen's d), a significance level set at 0.05, and a desired statistical power (power) of 0.85. The result of this analysis yielded a recommended sample size of 133 participants.

### **Behavioural data**

The German version of Cognitive Emotion Regulation Questionnaire (CERQ)<sup>45,101</sup> was included in this study. This questionnaire assesses nine cognitive coping strategies, which encompass self-blame, acceptance, rumination, positive refocusing, refocus on planning, positive reappraisal, putting into perspective, catastrophizing, and blaming others. The German version of the questionnaire comprises of 27 items, with each strategy measured through three questions. Participants responded using 5-point Likert-type scale ranging from 1 (almost never) to 5 (almost always). In this study investigation of functional connectivity associated with Acceptance and positive reappraisal scales were mainly focused.

#### **Image Acquisition**

Structural and functional MRI imaging was conducted using a 3 Tesla scanner (MAGNETOM Verio, Siemens Healthcare GmbH, Erlangen, Germany) equipped with a 32channel head coil. Throughout the MRI data acquisition process no significant maintenance or updates were carried out that could have impacted the data quality. Our analyses focused on a BOLD rs-fMRI scan using a T2-weighted multiband EPI sequence (TR = 1400 ms, TE = 30 ms, flip angle = 69°, echo spacing = 0.67 ms, number of volumes = 657, voxel size = 2.3 mm), with a total acquisition time of 15 minutes and 30 seconds. Additionally, T1-weighted structural volumes were obtained using the MP2RAGE sequence (TR = 5000 ms, TE = 2.92 ms, TI1 = 700 ms, TI2 = 2500 ms, FOV = 256 mm, voxel size = 1 mm isotropic). The acquisition of the structural volumes contained 176 slices acquired interspersed over a scanning duration of 8 minutes and 22 seconds. During the image acquisition, participants were instructed to maintain wakefulness, keep still, and gaze at a low-contrast fixation cross with their eyes open.

### Data analysis

#### **Pre-processing**

Data pre-processing was performed using CONN (version 2022), SPM 12, and the MATLAB Toolbox (version 2021b). Frist, the CONN's default pre-processing pipeline using SMP12's default parametere. This pipeline encompassed several stages: functional realignment and unwarping, translation and centering, conservative functional outlier detection, direct segmentation and normalization of functional data (1 mm resolution), translation and centering of structural data, segmentation and normalization of structural data (2.4 mm resolution), and lastly, spatial smoothing of functional and structural data using an 8 mm Gaussian kernel. Subsequently, the denoising phase was conducted. The aim of this phase is to pinpoint and eliminate confounding variables and artifacts from the estimated BOLD signal. These factors arise from three distinct sources: the BOLD signal originating from masks of white matter or cerebrospinal fluid, parameters and outliers defined during the pre-processing step, and an

estimation of the subjects' motion parameters. Following identification, these factors were

time series underwent temporal band-pass filtering within the 0.0008 Hz to infinity range, a standard procedure for resting-state connectivity analyses.

#### **Resting state Analysis**

For functional connectivity analysis in this study, the data-driven group-independent component approach (group-ICA) was performed using CONN. The CONN group-ICA consist of several steps: pre-conditioning variance normalization, concatenation of BOLD signal temporally, group-level dimensionality reduction, fast-ICA for spatial component estimation, and back-projection for individual spatial estimation. The analysis aimed to identify 20 independent components, aligning with earlier research adopting low model order analysis <sup>44,102,103</sup>. To differentiate noise components from underlying resting-state networks, each identified independent component (IC) underwent visual inspection and was compared with CONN's networks atlas via a spatial match-to-template function. This function gauged the overlap between individual IC's spatial maps and eight predefined brain networks (Default Mode, Sensorimotor, Visual, Salience, Dorsal Attention, Frontoparietal, Language, Cerebellar), as defined by CONN's ICA analyses of the HCP dataset (497 subjects). Subsequently, the temporal variability and frequency of each IC were quantified using CONN, calculated as the standard deviation of BOLD time-series. To control for Type I errors, cluster-size-based false discovery rate (FDR) correction was applied (p < 0.05, voxel thresholded at p < 0.001 within each analysis).

- - - -

#### Please insert Figure 6

- - - -

To determine which of the 20 identified ICs were predictive of the use of acceptance and reappraisal strategies, we examined the impact of each IC's BOLD signal variability on these emotion regulation strategies. This was achieved by conducting a Multiple Linear Regression model (Ordinary Least Squares) with a backward elimination approach. The analyses were conducted separately for the dependent variables of acceptance and reappraisal, incorporating gender as a categorical fixed factor to assess its influence within the regression framework.

## **Data Availability:**

The complete LEMON Data is accessible through Gesellschaft für wissenschaftliche

Datenverarbeitung mbH Göttingen (GWDG) at https://www.gwdg.de/. Both raw and

preprocessed data can be accessed via web browser (https://ftp.gwdg.de/pub/misc/MPI-

Leipzig\_Mind-Brain-Body-LEMON/) or through a fast FTP connection

(ftp://ftp.gwdg.de/pub/misc/MPI-Leipzig\_Mind-Brain-Body-LEMON/). In the event of any

future changes in the data's location, the dataset can be located using PID 21.11101/0000-

0007-C379-5 (e.g., http://hdl.handle.net/21.11101/0000-0007-C379-5).

## References

- 1. Kring, A. M. & Sloan, D. M. Emotion Regulation and Psychopathology: A Transdiagnostic Approach to Etiology and Treatment. (Guliford, New York, 2009).
- 2. Sheppes, G., Suri, G. & Gross, J. J. Emotion regulation and psychopathology. *Annu Rev Clin Psychol* **11**, 379–405 (2015).
- 3. Dadomo, H., Panzeri, M., Caponcello, D., Carmelita, A. & Grecucci, A. Schema therapy for emotional dysregulation in personality disorders: A review. *Current Opinion in Psychiatry* vol. 31 43–49 Preprint at https://doi.org/10.1097/YCO.00000000000380 (2018).
- 4. Frederickson, J. J., Messina, I. & Grecucci, A. Dysregulated anxiety and dysregulating defenses: Toward an emotion regulation informed dynamic psychotherapy. *Front Psychol* **9**, (2018).
- 5. Grecucci, A. *et al.* A dual route model for regulating emotions: Comparing models, techniques and biological mechanisms. *Front Psychol* **11**, 1–13 (2020).
- 6. Mennin, D. S., McLaughlin, K. A. & Flanagan, T. J. Emotion regulation deficits in generalized anxiety disorder, social anxiety disorder, and their co-occurrence. *J Anxiety Disord* **23**, (2009).

- 7. Schulze, L. *et al.* Neuronal correlates of cognitive reappraisal in borderline patients with affective instability. *Biol Psychiatry* **69**, (2011).
- 8. Leahy, R. L., Tirch, D. & Napolitano, L. A. *Emotion Regulation in Psychotherapy: A Practitioner's Guide.* (Guilford, 2011).
- 9. Messina, I., Sambin, M., Palmieri, A. & Viviani, R. Neural correlates of psychotherapy in anxiety and depression: a meta-analysis. *PLoS One* **8**, (2013).
- Dadomo, H., Panzeri, M., Caponcello, D., Carmelita, A. & Grecucci, A. Schema therapy for emotional dysregulation in personality disorders: A review. *Current Opinion in Psychiatry* vol. 31 43–49 Preprint at https://doi.org/10.1097/YCO.00000000000380 (2018).
- 11. Grecucci, A. *et al.* Baseline and strategic effects behind mindful emotion regulation: Behavioral and physiological investigation. *PLoS One* **10**, (2015).
- 12. Goldin, P. R., Moodie, C. A. & Gross, J. J. Acceptance versus reappraisal: Behavioral, autonomic, and neural effects. *Cogn Affect Behav Neurosci* **19**, 927–944 (2019).
- 13. Hayes, S. C. Integrative Behavioral Couples Therapy (IBCT; Jacobson & Christensen, 1996), and Mindfulness-Based Cognitive Therapy (MBCT. Borkovec & Roemer vol. 35 (2004).
- 14. Kahl, K. G., Winter, L., Schweiger, U. & Sipos, V. The third wave of cognitivebehavioural psychotherapies: concepts and efficacy. *Fortschr Neurol Psychiatr* **79**, 330–339 (2011).
- Messina, I., Sambin, M., Beschoner, P. & Viviani, R. Changing views of emotion regulation and neurobiological models of the mechanism of action of psychotherapy. *Cognitive, Affective and Behavioral Neuroscience* vol. 16 571– 587 Preprint at https://doi.org/10.3758/s13415-016-0440-5 (2016).
- 16. Messina, I., Grecucci, A., MAROGNA, C. & CALVO, V. Relational exposure and semantic processes as mechanisms of change in psychodynamic psychotherapy: Convergences between psychotherapy research and affective neuroscience. *TPM Test Psychom Methodol Appl Psychol* **27**, 43–56 (2020).
- Hayes, S. C., Pistorello, J. & Levin, M. E. Acceptance and Commitment Therapy as a Unified Model of Behavior Change. *Couns Psychol* 40, 976–1002 (2012).
- 18. Wolgast, M., Lundh, L. G. & Viborg, G. Cognitive reappraisal and acceptance: An experimental comparison of two emotion regulation strategies. *Behaviour Research and Therapy* **49**, 858–866 (2011).
- Spencer, S. D., Buchanan, J. A. & Masuda, A. Effects of Brief Acceptance and Cognitive Reappraisal Interventions on Experiential Avoidance in Socially Anxious Individuals: A Preliminary Investigation. *Behav Modif* 44, 841–864 (2020).
- 20. Gross, J. The Emerging Field of Emotion Regulation: An Integrative Review. *Review of General Psychology* **2**, 271–299 (1998).
- 21. Gross, J. Emotion regulation: Affective, cognitive, and social consequences. *Psychophysiology* **39**, (2002).
- 22. Aldao, A. & Nolen-Hoeksema, S. When are adaptive strategies most predictive of psychopathology? *J Abnorm Psychol* **121**, 276–281 (2012).
- 23. Aldao, A., Nolen-Hoeksema, S. & Schweizer, S. Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review* vol. 30 217–237 Preprint at https://doi.org/10.1016/j.cpr.2009.11.004 (2010).

- 24. Dan-Glauser, E. S. & Gross, J. J. The temporal dynamics of emotional acceptance: Experience, expression, and physiology. *Biol Psychol* **108**, 1–12 (2015).
- 25. Garland, E. L., Gaylord, S. A. & Fredrickson, B. L. Positive Reappraisal Mediates the Stress-Reductive Effects of Mindfulness: An Upward Spiral Process. *Mindfulness (N Y)* **2**, 59–67 (2011).
- 26. Desrosiers, A., Vine, V., Klemanski, D. H. & Nolen-Hoeksema, S. Mindfulness and emotion regulation in depression and anxiety: Common and distinct mechanisms of action. *Depress Anxiety* **30**, 654–661 (2013).
- 27. Peh, C. X. *et al.* Emotion regulation and emotional distress: The mediating role of hope on reappraisal and anxiety/depression in newly diagnosed cancer patients. *Psychooncology* 1191–1197 (2017).
- 28. Uchida, M. *et al.* Emotion regulation ability varies in relation to intrinsic functional brain architecture. *Soc Cogn Affect Neurosci* **10**, 1738–1748 (2014).
- 29. Hofmann, S. G., Heering, S., Sawyer, A. T. & Asnaani, A. How to handle anxiety: The effects of reappraisal, acceptance, and suppression strategies on anxious arousal. *Behaviour Research and Therapy* **47**, 389–394 (2009).
- Ochsner, K. N. & Gross, J. J. The cognitive control of emotion. *Trends in Cognitive Sciences* vol. 9 242–249 Preprint at https://doi.org/10.1016/j.tics.2005.03.010 (2005).
- 31. Lebois, L. A. M. *et al.* A shift in perspective: Decentering through mindful attention to imagined stressful events. *Neuropsychologia* **75**, 505–524 (2015).
- 32. Murakami, H. *et al.* Neural networks for mindfulness and emotion suppression. *PLoS One* **10**, (2015).
- 33. Smoski, M. J. *et al.* Neural indicators of emotion regulation via acceptance vs reappraisal in remitted major depressive disorder. *Soc Cogn Affect Neurosci* **10**, 1187–1194 (2014).
- Ellard, K. K., Barlow, D. H., Whitfield-Gabrieli, S., Gabrieli, J. D. E. & Deckersbach, T. Neural correlates of emotion acceptance vs worry or suppression in generalized anxiety disorder. *Soc Cogn Affect Neurosci* 12, 1009–1021 (2017).
- 35. Kross, E., Davidson, M., Weber, J. & Ochsner, K. Coping with Emotions Past: The Neural Bases of Regulating Affect Associated with Negative Autobiographical Memories. *Biol Psychiatry* **65**, 361–366 (2009).
- 36. Westbrook, C. *et al.* Mindful attention reduces neural and self-reported cueinduced craving in smokers. *Soc Cogn Affect Neurosci* **8**, 73–84 (2013).
- Kober, H., Buhle, J., Weber, J., Ochsner, K. N. & Wager, T. D. Let it be: Mindful acceptance down-regulates pain and negative emotion. Soc Cogn Affect Neurosci 14, 1147–1158 (2019).
- 38. Dixon, M. L. *et al.* Emotion Regulation in Social Anxiety Disorder: Reappraisal and Acceptance of Negative Self-beliefs. *Biol Psychiatry Cogn Neurosci Neuroimaging* **5**, 119–129 (2020).
- 39. Messina, I., Grecucci, A. & Viviani, R. Neurobiological models of emotion regulation: a meta-analysis of neuroimaging studies of acceptance as an emotion regulation strategy. *Soc Cogn Affect Neurosci* **16**, 257–267 (2021).
- 40. Sezer, I., Pizzagalli, D. A. & Sacchet, M. D. Resting-state fMRI functional connectivity and mindfulness in clinical and non-clinical contexts: A review and synthesis. *Neurosci Biobehav Rev* **135**, 104583 (2022).

- 41. Grecucci, A., Ghomroudi, P. A., Monachesi, B. & Messina, I. *The Neural Signature of Inner Peace: Morphometric Differences between High and Low Accepters.*
- 42. Buhle, J. T. *et al.* Cognitive reappraisal of emotion: A meta-analysis of human neuroimaging studies. *Cerebral Cortex* **24**, 2981–2990 (2014).
- 43. Morawetz, C., Bode, S., Baudewig, J., Kirilina, E. & Heekeren, H. R. Changes in Effective Connectivity Between Dorsal and Ventral Prefrontal Regions Moderate Emotion Regulation. *Cerebral Cortex* **26**, 1923–1937 (2016).
- 44. Zanella, F., Monachesi, B. & Grecucci, A. Resting-state BOLD temporal variability in sensorimotor and salience networks underlies trait emotional intelligence and explains differences in emotion regulation strategies. *Sci Rep* **12**, (2022).
- 45. Garnefski, N., Kraaij, V. & Spinhoven, P. *Negative Life Events, Cognitive Emotion Regulation and Emotional Problems.* www.elsevier.com/locate/paid (2001).
- 46. Ghomroudi, P. A., Scaltritti, M. & Grecucci, A. Decoding reappraisal and suppression from neural circuits: A combined supervised and unsupervised machine learning approach. *Cogn Affect Behav Neurosci* (2023) doi:10.3758/s13415-023-01076-6.
- 47. Monachesi, B., Grecucci, A., Ahmadi Ghomroudi, P. & Messina, I. Comparing reappraisal and acceptance strategies to understand the neural architecture of emotion regulation: a meta-analytic approach. *Frontiers in Psychology* vol. 14 Preprint at https://doi.org/10.3389/fpsyg.2023.1187092 (2023).
- 48. Biswal, B., Yetkin, F. Z., Haughton, V. M. & Hyde, J. S. Functional connectivity Echo-Planar MRI. *Magn Reson Med* **34**, 537–541 (1995).
- 49. Dörfel, D., Gärtner, A. & Scheffel, C. Resting State Cortico-Limbic Functional Connectivity and Dispositional Use of Emotion Regulation Strategies: A Replication and Extension Study. *Front Behav Neurosci* **14**, (2020).
- 50. Bell, A. J. & Sejnowski, T. J. An Information-Maximisation Approach t o Blind Separation and Blind Deconvolution. (1995).
- 51. Calhoun, V. D., Adali, T., Pearlson, G. D. & Pekar, J. J. A Method for Making Group Inferences from Functional MRI Data Using Independent Component Analysis. (2001) doi:10.1002/hbm.
- 52. Monachesi, B., Grecucci, A., Ahmadi Ghomroudi, P. & Messina, I. Comparing reappraisal and acceptance strategies to understand the neural architecture of emotion regulation: a meta-analytic approach. *Frontiers in Psychology* vol. 14 Preprint at https://doi.org/10.3389/fpsyg.2023.1187092 (2023).
- 53. Picó-Pérez, M. *et al.* Dispositional use of emotion regulation strategies and resting-state cortico-limbic functional connectivity. *Brain Imaging Behav* **12**, 1022–1031 (2018).
- 54. Garrett, D. D., Epp, S. M., Perry, A. & Lindenberger, U. Local temporal variability reflects functional integration in the human brain. *Neuroimage* **183**, (2018).
- 55. Phillips, M. L., Ladouceur, C. & Drevets, W. C. Automatic and Volontary Regulation of Emotion. *Mol Psychiatry* **13**, 829–857 (2008).
- 56. Hernández, S. E., Barros-Loscertales, A., Xiao, Y., González-Mora, J. L. & Rubia, K. Gray Matter and Functional Connectivity in Anterior Cingulate Cortex are Associated with the State of Mental Silence During Sahaja Yoga Meditation. *Neuroscience* 371, (2018).

- 57. Yang, L. Z. *et al.* Electrical stimulation reduces smokers' craving by modulating the coupling between dorsal lateral prefrontal cortex and parahippocampal gyrus. *Soc Cogn Affect Neurosci* **12**, 1296–1302 (2017).
- 58. Katz, B., Adams, R. D. & Victor, M. Principles of Neurology. *Journal of Neuro-Ophthalmology* **16**, (1996).
- 59. Zeidan, F. *et al.* Brain mechanisms supporting the modulation of pain by mindfulness meditation. *Journal of Neuroscience* **31**, 5540–5548 (2011).
- 60. Frank, D. W. *et al.* Emotion regulation: Quantitative meta-analysis of functional activation and deactivation. *Neuroscience and Biobehavioral Reviews* vol. 45 202–211 Preprint at https://doi.org/10.1016/j.neubiorev.2014.06.010 (2014).
- 61. Morawetz, C., Bode, S., Derntl, B. & Heekeren, H. R. The effect of strategies, goals and stimulus material on the neural mechanisms of emotion regulation: A meta-analysis of fMRI studies. *Neuroscience and Biobehavioral Reviews* vol. 72 111–128 Preprint at https://doi.org/10.1016/j.neubiorev.2016.11.014 (2017).
- 62. Grecucci, A., Giorgetta, C., Bonini, N. & Sanfey, A. G. Living emotions, avoiding emotions: Behavioral investigation of the regulation of socially driven emotions. *Front Psychol* **3**, 1–14 (2013).
- 63. Grecucci, A., Giorgetta, C., Van't Wout, M., Bonini, N. & Sanfey, A. G. Reappraising the ultimatum: An fMRI study of emotion regulation and decision making. *Cerebral Cortex* **23**, 399–410 (2013).
- 64. Kozasa, E. H. *et al.* Meditation training increases brain efficiency in an attention task. *Neuroimage* **59**, (2012).
- 65. Brefczynski-Lewis, J. A., Lutz, A., Schaefer, H. S., Levinson, D. B. & Davidson, R. J. Neural correlates of attentional expertise in long-term meditation practitioners. *Proc Natl Acad Sci U S A* **104**, (2007).
- 66. Craig, A. D. How do you feel? Interoception: the sense of the physiological condition of the body. *Nature Reviews Neuroscience volume 3*, *)Nature Reviews Neuroscience* **3**, 655–656 (2002).
- 67. Khalsa, S. S., Rudrauf, D., Feinstein, J. S. & Tranel, D. The pathways of interoceptive awareness. *Nat Neurosci* **12**, 1494–1496 (2009).
- 68. Min, J. *et al.* Emotion Downregulation Targets Interoceptive Brain Regions While Emotion Upregulation Targets Other Affective Brain Regions. *Journal of Neuroscience* **42**, 2973–2985 (2022).
- 69. Farb, N. A. S., Anderson, A. K. & Segal, Z. V. The mindful brain and emotion regulation in mood disorders. *Canadian Journal of Psychiatry* **57**, 70–77 (2012).
- 70. Parkinson, T. D., Kornelsen, J. & Smith, S. D. Trait mindfulness and functional connectivity in cognitive and attentional resting state networks. *Front Hum Neurosci* **13**, (2019).
- Ochsner, K. N., Silvers, J. A. & Buhle, J. T. Functional imaging studies of emotion regulation: a synthetic review and evolving model of the cognitive control of emotion. *Annals of the New York Academy of Sciences* vol. 1251 E1– E24 Preprint at https://doi.org/10.1111/j.1749-6632.2012.06751.x (2012).
- 72. Kohn, N. *et al.* Neural network of cognitive emotion regulation An ALE metaanalysis and MACM analysis. *Neuroimage* **87**, 345–355 (2014).
- 73. Messina, I., Bianco, S., Sambin, M. & Viviani, R. Executive and semantic processes in reappraisal of negative stimuli: Insights from a meta-analysis of neuroimaging studies. *Front Psychol* **6**, (2015).
- 74. Li, W. *et al.* Differential involvement of frontoparietal network and insula cortex in emotion regulation. *Neuropsychologia* **161**, (2021).

- 75. Craig, A. D. Emotional moments across time: A possible neural basis for time perception in the anterior insula. *Philosophical Transactions of the Royal Society B: Biological Sciences* vol. 364 1933–1942 Preprint at https://doi.org/10.1098/rstb.2009.0008 (2009).
- 76. Zaki, J., Davis, J. I. & Ochsner, K. N. Overlapping activity in anterior insula during interoception and emotional experience. *Neuroimage* **62**, (2012).
- 77. MacDonald, D. D., Ingersoll, C. G. & Berger, T. A. Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. *Arch Environ Contam Toxicol* **39**, 20–31 (2000).
- 78. Curtis, C. E. & D'Esposito, M. Persistent activity in the prefrontal cortex during working memory. *Trends Cogn Sci* **7**, 415–423 (2003).
- 79. Hare, T. A., Camerer, C. F. & Rangel, A. Valuation System. **324**, 646–649 (2009).
- 80. Bush, G., Luu, P. & Posner, M. I. Cognitive and emotional influences in anterior cingulate cortex. *Trends in Cognitive Sciences* vol. 4 Preprint at https://doi.org/10.1016/S1364-6613(00)01483-2 (2000).
- 81. Stevens, F. L., Hurley, R. A. & Taber, K. H. Anterior cingulate cortex: Unique role in cognition and emotion. *Journal of Neuropsychiatry and Clinical Neurosciences* 23, 121–125 (2011).
- 82. Piretti, L. *et al.* Impaired processing of conspecifics in Parkinson's disease. *Applied Neuropsychology:Adult* (2022) doi:10.1080/23279095.2022.2074299.
- 83. Bilevicius, E., Smith, S. D. & Kornelsen, J. Resting-State Network Functional Connectivity Patterns Associated with the Mindful Attention Awareness Scale. *Brain Connect* **8**, 40–48 (2018).
- 84. Goulas, A., Uylings, H. B. M. & Stiers, P. Unravelling the intrinsic functional organization of the human lateral frontal cortex: A parcellation scheme based on resting state fMRI. *Journal of Neuroscience* **32**, 10238–10252 (2012).
- 85. Ochsner, K. N. & Gross, J. J. The cognitive control of emotion. *Trends in Cognitive Sciences* vol. 9 242–249 Preprint at https://doi.org/10.1016/j.tics.2005.03.010 (2005).
- 86. Ochsner, K. N., Silvers, J. A. & Buhle, J. T. Functional imaging studies of emotion regulation: a synthetic review and evolving model of the cognitive control of emotion. *Annals of the New York Academy of Sciences* vol. 1251 E1– E24 Preprint at https://doi.org/10.1111/j.1749-6632.2012.06751.x (2012).
- 87. Messina, I., Bianco, S., Sambin, M. & Viviani, R. Executive and semantic processes in reappraisal of negative stimuli: Insights from a meta-analysis of neuroimaging studies. *Front Psychol* **6**, (2015).
- 88. Grecucci, A., Rubicondo, D., Siugzdaite, R., Surian, L. & Job, R. Uncovering the social deficits in the autistic brain. A source-based morphometric study. *Front Neurosci* **10**, (2016).
- 89. Bzdok, D., Laird, A. R., Zilles, K., Fox, P. T. & Eickhoff, S. B. An investigation of the structural, connectional, and functional subspecialization in the human amygdala. *Hum Brain Mapp* **34**, 3247–3266 (2013).
- 90. Cole, M. W. & Schneider, W. The cognitive control network: Integrated cortical regions with dissociable functions. *Neuroimage* **37**, (2007).
- Lückmann, H. C., Jacobs, H. I. L. & Sack, A. T. The cross-functional role of frontoparietal regions in cognition: Internal attention as the overarching mechanism. *Progress in Neurobiology* vol. 116 Preprint at https://doi.org/10.1016/j.pneurobio.2014.02.002 (2014).

- 92. Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven attention in the brain. *Nat Rev Neurosci* **3**, 201–215 (2002).
- 93. Shenhav, A., Botvinick, M. M. & Cohen, J. D. The expected value of control: An integrative theory of anterior cingulate cortex function. *Neuron* vol. 79 Preprint at https://doi.org/10.1016/j.neuron.2013.07.007 (2013).
- 94. Füstös, J., Gramann, K., Herbert, B. M. & Pollatos, O. On the embodiment of emotion regulation: Interoceptive awareness facilitates reappraisal. *Soc Cogn Affect Neurosci* **8**, 911–917 (2013).
- 95. Kropf, E., Syan, S. K., Minuzzi, L. & Frey, B. N. From anatomy to function: the role of the somatosensory cortex in emotional regulation. *Braz J Psychiatry* **41**, 261–269 (2019).
- 96. Critchley, H. D., Wiens, S., Rotshtein, P., Öhman, A. & Dolan, R. J. Neural systems supporting interoceptive awareness. *Nat Neurosci* **7**, 189–195 (2004).
- 97. Damasio, A. R. *et al.* Subcortical and cortical brain activity during the feeling of self-generated emotions. *Nat Neurosci* **3**, 1049–1056 (2000).
- 98. Kohl, A., Rief, W. & Glombiewski, J. A. How effective are acceptance strategies? A meta-analytic review of experimental results. *J Behav Ther Exp Psychiatry* **43**, 988–1001 (2012).
- 99. McRae, K. Cognitive emotion regulation: A review of theory and scientific findings. *Current Opinion in Behavioral Sciences* vol. 10 119–124 Preprint at https://doi.org/10.1016/j.cobeha.2016.06.004 (2016).
- Babayan, A. *et al.* Data descriptor: A mind-brain-body dataset of MRI, EEG, cognition, emotion, and peripheral physiology in young and old adults. *Sci Data* 6, (2019).
- 101. Loch, N., Hiller, W. & Witthöft, M. Der cognitive emotion regulation questionnaire (CERQ) erste teststatistische überprüfung einer deutschen adaption. *Z Klin Psychol Psychother* **40**, 94–106 (2011).
- 102. Elseoud, A. A. *et al.* Group-ICA model order highlights patterns of functional brain connectivity. *Front Syst Neurosci* (2011) doi:10.3389/fnsys.2011.00037.
- 103. Saviola, F. *et al.* Trait and state anxiety are mapped differently in the human brain. *Sci Rep* **10**, (2020).

## **Author Contributions**

P.AG. Contributed to the writing of the manuscript, and performed data analysis. R.S. Provided supervision. I.M. Reviewed the manuscript. A.G. Conceived the work, reviewed the manuscript, and provided supervision. All the co-authors have read and approved the content of the final version of the manuscript.

## **Competing interests**

The authors declare no competing interests.

Figure 1: Resting-state BOLD temporal variability predicting acceptance ability in the affective network (IC2)




Figure 2: Resting-state BOLD temporal variability predicting acceptance ability in the frontoparietal network (IC11)




Figure 3: Resting-state BOLD temporal variability predicting acceptance ability in the sensorimotor network (IC13)




Figure 4: Resting-state BOLD temporal variability predicting acceptance ability in the executive network (IC18)




Figure 5: Resting-state BOLD temporal variability predicting reappraisal ability in the sensorimotor network (IC20)




Figure 6 Schematic diagram of the methodology. First the resting state data was preprocessed. Then 20 independent components were extracted using an unsupervised machine learning Group ICA approach.




Table 1: Result of Backward Regression	n
--	---

Emotion Regulation Strategy	β	Р
	-60.85	0.001
Accentance	-67.89	0.001
Acceptance	80.16	< 0.001
	24.42	0.022
Reapprasial	0.339	< 0.001
	Acceptance	-60.85 -67.89 80.16 24.42

Beta ( $\beta$ ) and corrected p-value for the significant relationships between the BOLD temporal variability and both ER strategies.

Strategy	Network	ROI	Voxels	MNI coordinates (mm)
		L/R Temporal Fusiform Cortex	654	(+36,-24,-28), (+32,-2,-42), (-32,-4,-42), (-34,-28,- 26)
		L/R Inferior Temporal Gyrus	647	(+52,-20,-32), (+46,-2,-42), (-48,-6,-40), (-52,-20,- 30
		L/R Parahippocampal Gyrus	649	(+22,-8,-30), (-22,-10,-30)
	IC 2	L/R Hippocampus	668	(+26,-20,-14), (-24,-22,-14)
	10.2	L/R Middle Temporal Gyrus	377	(-56,-4,-24)
		L/R Amygdala	296	(+24,-4,-18), (-24,-6,-18)
		L/R Frontal Orbital Cortex	899	(-26,+22,-20), (+24,+22,-20)
		R Superior Temporal Gyrus	193	(+56,-2,-12)
		R Thalamus	444	(+14,-26,+2)
		L/R Angular Gyrus	724	(-50,-56,+34), (+52,-52,+38)
		L Supramarginal Gyrus	636	(-54,-46,+42)
		L/R Middle Temporal Gyrus	1212	(-62,-28,-12), (+64,-22,-14)
		L/R Superior Frontal Gyrus	1643	(-14,+30,+50), (+14,+30,+52)
		L/R Middle Frontal Gyrus	1513	(-36,+18,+46), (+38,+22,+46)
	IC 11	Paracingulate Gyrus Left	803	(-6,+44,+16)
		L/R Amygdala	327	(-22,-4,-18), (+24,-4,-18)
		Anterior Cingulate Gyrus	1359	(+0,+14,+30)
		L/R Putamen	761	(-24,+2,+0), (+24,+4,+0)
Acceptance		L Frontal Orbital CorteX	623	(-28,+24,-12)
		L Insular Cortex	875	(-36,+8,-2)
		L/R Precentral Gyrus	2219	(+50,-4,+40), (-48,-6,+40)
	IC13	L/R Postcentral Gyrus	1898	(+50,-20,+42), (-52,-20,+42)
		L/R Superior Temporal Gyrus	261	(-56,-4,-8), (-62,-28,+4), (+60,-24,+2)
		L/R Insular Cortex	1117	(+38,+0,+2), (-38,-4,+2)
		R/L Parietal Operculum Cortex	533	(+48,-28,+22), (-48,-32,+20)
		L/R Supramarginal Gyrus	678	(+58,-26,+38), (-58,-30,+34)
		L/R Caudate r	213	(+14,+2,+18), (-14,+2,+18)
		L/R Thalamus r	843	(+12,-20,+10), (-10,-20,+10)
		Inferior Frontal Gyrus	464	(+52,+16,+20)
		L/R Frontal Orbital Cortex	1356	(+30,+22,-16), (-30,+24,-16)
		R Middle Frontal Gyrus	1672	(+40,+20,+40)
	IC18	Precuneous	4016	(+0,-60,+44)
	1010	L/R Insular Cortex	649	(+38,+4,-2), (-38,+4,-2)
		L/R Precentral Gyrus	2493	(+36,-10,+52), (-34,-12,+50)
		Anterior Cingulate Gyrus	1097	(+0,+24,+20)
		R Posterior Parahippocampal Gyrus	274	(+22,-32,-16)
		R Middle Temporal Gyrus	1064	(+58,-50,+2)

## Table 2: Summary of the main neural results for acceptance

Strategy	Network	ROI	Voxels	MNI coordinates (mm)
Reappraisal	IC20	L/R Precentral Gyrus Left	2784	(-24,-18,+60), (+26,-18,+60)
		Postcentral Gyrus Left	2492	(-30,-32,+60), (+30,-30,+60)
		L/R Supplementary Motor Cortex	587	(+6,-4,+58), (-6,-2,+56)
		Precuneous	1059	(+8,-46,+48)
		L/R Superior Frontal Gyrus	699	(-14,+0,+66), (+16,-4,+68)
		Cingulate Gyrus AC/PC	468	(+0,-4,+42), (+4,-28,+42)
		L/R Insular Cortex	204	(+36,-18,+10), (-36,-20,+10)
		L/R Parietal Operculum Cortex	304	(+44,-26,+20), (-44,-30,+20)
		L/R Central Opercular Cortex	240	(-46,-18,+16), (+44,-14,+16)
		L/R Superior Parietal Lobule	469	(+22,-46,+66), (-24,-48,+64)
		R Central Opercular Cortex	155	(+44,-14,+16)
		R Heschl's Gyrus	204	(+46,-20,+8)

## Table 3: Summary of the main neural results for reappraisal