

A review of statistical models used to characterize species-habitat associations with animal movement data

Katie R.N. Florko^{1,*}, Ron R. Togunov^{2,3,4}, Rowenna Gryba^{1,5,6}, Evan Sidrow⁵, Steven H. Ferguson^{7,8}, David J. Yurkowski^{7,8}, Marie Auger-Méthé^{1,5}

¹Institute for the Oceans and Fisheries, University of British Columbia, Vancouver BC, Canada

²Department of Zoology, University of British Columbia, Vancouver BC, Canada

³Department of Mathematical Sciences, Norwegian University of Science and Technology, Trondheim, Norway

⁴Centre for Biodiversity Dynamics, Norwegian University of Science and Technology, Trondheim, Norway

⁵Department of Statistics, University of British Columbia, Vancouver BC, Canada

⁶Department of Geography, University of British Columbia, Vancouver BC, Canada

⁷Fisheries and Oceans Canada, Freshwater Institute, Winnipeg MB, Canada

⁸Department of Biological Sciences, University of Manitoba, Winnipeg MB, Canada

***Corresponding author:**

Katie R.N. Florko

Aquatic Ecosystem Research Laboratory

University of British Columbia

2202 Main Mall

Vancouver BC V6T 1Z4

Phone: 1-604-822-2731

Email: katieflorko@gmail.com

Abstract

1. Understanding species-habitat associations is fundamental to ecological sciences and for species conservation. Consequently, various statistical approaches have been designed to infer species-habitat associations. Due to their conceptual and mathematical differences, these methods can yield contrasting results.
2. In this paper, we describe and compare commonly used statistical models that relate animal movement data to environmental data. Specifically, we examined selection functions which include resource selection function [RSF] and step-selection function [SSF], as well as hidden Markov models (HMMs) and related methods such as state-space models. We demonstrate differences in assumptions of each method while highlighting advantages and limitations. Additionally, we provide guidance on selecting the most appropriate statistical method based on research objectives and intended inference. To illustrate the varying ecological insights derived from each statistical model, we apply them to the movement track of a single ringed seal (*Pusa hispida*) in a case study.
3. Through our case study, we demonstrate that each model yields varying ecological insights. For example, the RSF indicated *selection* of areas with high prey diversity, whereas the SSFs indicated *no discernable relationship* with prey diversity. Furthermore, the HMM reveals variable associations with prey diversity across different behaviors, for example, a positive relationship between prey diversity and a slow-movement behaviour. Notably, the three models identified different “important” areas.
4. This case study highlights the critical significance of selecting the appropriate model as an essential step in the process of identifying species-habitat relationships and specific areas of importance. Our comprehensive review provides the foundational information required

for making informed decisions when choosing the most suitable statistical methods to address specific questions, such as identifying expansive corridors or protected zones, understanding movement patterns, or studying behaviours. In addition, this study informs researchers with the necessary tools to apply these methods most effectively.

Keywords

animal movement, biologging, habitat selection, hidden Markov models, integrated step-selection functions, movement ecology, Poisson point process models, resource selection functions, ringed seal (*Pusa hispida*), step-selection functions, telemetry

1. Background

Understanding the relationships between animals' space use and their physical environment is a fundamental aspect of ecological research and is essential for species conservation (Andrewartha & Browning, 1961; Huey, 1991; Matthiopoulos et al., 2015). Movement data from biologging devices provide valuable information on animal space use (Hussey et al., 2015; Nathan et al., 2022), and researchers often use statistical models to relate the movement data (in terms of occurrence, movement, or behaviour) to, for example, indicators of resources (e.g., vegetation, Valeix et al. 2011), proxies of energy (e.g., terrain ruggedness, Sells et al. 2022), or perceived predation risk (e.g., canopy cover, Godvik et al. 2009). These models can provide insight on the environmental conditions associated with key ecological concepts such as home range, habitat selection, movement corridors, behaviour, and critical habitat (Morales et al., 2010; Nathan et al., 2022). Identifying and designating critical habitat is one of the main ways governments protect species from rapid and widespread habitat degradation and climate changes (e.g., under the

Environment Protection and Biodiversity Conservation (EPBC) Act in Australia, Species at Risk Act (SARA) in Canada, Wildlife and Countryside Act in the United Kingdom, Endangered Species Act (ESA) in the United States). Thus, it is essential that statistical models that use movement data to identify critical habitat are chosen, implemented, and interpreted properly.

Various models aim to link observed animal locations to environmental covariates, but each model is appropriate for specific research questions. Many of these models can be easily implemented using R packages (e.g., `amt`, Signer et al. 2019; `momentuHMM`, McClintock and Michelot 2018; see Joo et al. 2020 for a relevant review of packages), and their use is ubiquitous in animal movement literature (e.g., Alston et al. 2020; Chimienti et al. 2021). However, choosing an appropriate method is a complex and challenging aspect of movement modeling (Joo et al., 2020). For example, different models address ecological questions at different scales, from large-scale questions on important areas for species (e.g., first- or second-order selection at the species or home range scale, respectively), to smaller-scale movement- or behaviour-specific questions (e.g., third- or fourth-order selection at the habitat usage or actual food intake scale, respectively; Owen 1972; Johnson 1980; Nams 2013). Some methods are designed to understand habitat preference via selection functions (e.g., resource selection function [RSF], step-selection function [SSF], or integrated-SSF) and others are interested in understanding how habitat relates to discrete behavioural states (e.g., hidden Markov model [HMM]). As a result, consideration of scale and the behaviour of interest when choosing a model is imperative for meaningful interpretation.

Here, we describe and compare the mainstream models for linking animals' movement data to the environment: RSF, SSF, and HMM. Our objective is to help clarify the intended use and

mathematical underpinnings of each model to help ecologists properly choose between them. Our review is unique as we compare all three models. Reviews comparing selection functions (e.g., Hooten et al. 2020; Northrup et al. 2022) or selection functions with inhomogenous point process models (IPPs, e.g., Mercker et al. 2021) are available, but here we also included HMMs, a fundamentally different, but increasingly popular, model to relate animal movement data to environmental covariates.

Table 1. Terminology used in this paper.

Term	Definition
Habitat	The set of environmental covariates (biotic resources and abiotic conditions) that characterize the space an animal inhabits (Northrup et al., 2013)
Habitat selection	The process whereby individuals preferentially use, or occupy, a non-random set of available habitats (Morris, 2003)
Habitat selection function	A function proportional to the probability of selection of habitat (Northrup et al., 2013)
Habitat availability	The accessibility, prevalence, and procurability of physical and biological components of a habitat by animals (Wiens, 1984)
Habitat use	The exploitation of habitat to meet a biological need (Hall et al. 1997), in the RSF analysis, the presence of an animal at a

	location (Northrup et al., 2013)
--	----------------------------------

2. Description of statistical models

2.1 Resource selection functions (RSF)

A resource selection function (RSF) is a widely used function that relates available habitat to the *relative* probability of use by an animal (Boyce & McDonald, 1999; Fieberg et al., 2021; C. J. Johnson et al., 2006; Manly et al., 1993). RSFs originally assumed that the environmental conditions selected by an animal provides desirable resources, and that the probability of habitat selection (Table 1) is a function of predictor variables (e.g., environmental, geographical) representing a resource distribution on a landscape (Lele et al., 2013). RSFs are now often referred to as *habitat* selection functions or *analyses* as they can include covariates other than resources (e.g., covariates that describe habitats avoided by animals; Matthiopoulos et al. 2015).

When applied to movement data, RSFs model observed/available resource units. Observed animal locations are often referred to as “used” locations. The observed animal locations represent a sample of the habitat used by the animals (Fig. 1, Table 1), and are often compared to a sample of available locations randomly selected within an animal’s home range (e.g., the minimum convex polygon (MCP), or extent, of the observed locations, Fig. 1). RSFs are widely applied due to their ease of use (e.g., using the `amt` R package, Signer et al. 2019) and the general information on species-habitat relationships they provide (Northrup et al., 2022).




Figure 1. A) observed animal locations, B) the home range (here via minimum convex polygon), and C) available locations randomly sampled from within the minimum convex polygon.

The RSF, $w(\mathbf{x})$, essentially represents selection, and more formally represents the ratio between the use (i.e., utilization) distribution $F^U(\mathbf{x})$ (the frequency distribution of habitat covariates used by an animal) and available distribution $F^A(\mathbf{x})$ (the frequency distribution of habitat covariates assumed to be available) where x denotes habitat covariates. To estimate $w(\mathbf{x})$, we first define the use distribution, $F^U(\mathbf{x})$, as follows:

Equation 1

$$F^U(\mathbf{x}) = \frac{w(\mathbf{x})F^A(\mathbf{x})}{\int_{\mathbf{x}' \in \Omega} w(\mathbf{x}')F^A(\mathbf{x}')d\mathbf{x}'},$$

where in the denominator we integrate over the entire domain of the environment, Ω , to ensure that $F^U(\mathbf{x})$ is a valid probability distribution that integrates to one. Equation 1 shows that the use distribution is a function of both the available habitat (as defined by $F^A(\mathbf{x})$) and the selection function, $w(\mathbf{x})$, and that the model is parameterized in *environmental space* (Fithian & Hastie, 2012; Johnson et al., 2006; Lele & Keim, 2006; Northrup et al., 2013).

Rearranging terms in equation 1 shows how the RSF relates to the ratio between the use and available distributions (i.e., $w(\mathbf{x}) \propto F^U(\mathbf{x})/F^A(\mathbf{x})$). The RSF only provides an estimate of the *relative* selection probability, and not the *absolute* probability of selection, known as the resource selection probability function, RSPF, (Lele et al., 2013; Lele & Keim, 2006). Specifically, the RSF $w(\mathbf{x})$ encodes the relative strength with which an animal selects a given habitat with covariates \mathbf{x} . The RSF, $w(\mathbf{x})$, is typically defined in the exponential form:

Equation 2

$$w(\mathbf{x}) = \exp(\beta_1 x_1 + \beta_2 x_2 + \dots + \beta_k x_k),$$

where $\mathbf{x} = x_1, \dots, x_k$ denotes the values of k predictor habitat variables and β_1, \dots, β_k are the associated selection coefficients (Manly et al., 1993). The coefficients of the RSF (β_1, \dots, β_k) can be estimated using a binomial logistic regression where the response, $Y = \{y_1, \dots, y_n\}$ is a set of binary random variables representing observed ($y_i = 1$) and available ($y_i = 0$) resource units, and the predictor is the linear form of the RSF, $\ln(w(\mathbf{x}))$. As movement datasets only contains information on used habitat variables, available resource units, which are often erroneously treated as true absences, must be randomly sampled from F^A .

Alternative to using a logistic regression, RSFs can also be estimated using an inhomogeneous Poisson point process model (IPP). IPPs model the density of points in space and thus are ideal for modeling presence-only data resulting from movement datasets, because, unlike selection functions, they do not use available locations as pseudo-absence points (Warton & Shepherd, 2010). The point process likelihood can be evaluated using the `inlabru` R package

(Bachl et al., 2019). An IPP can be derived by considering movement locations as a realization of a space-time point process (Johnson et al., 2013) and is connected to an RSF that models the use distribution in *geographical space* rather than in *environmental space* (eqn 1):

Equation 3

$$F^U(s) = \frac{w(\mathbf{x}(s)) F^A(s)}{\int_{s' \in G} w(\mathbf{x}(s')) F^A(s') ds'}$$

where the selection function, $w(\mathbf{x}(s))$, is now a function of $\mathbf{x}(s)$, the habitat covariates \mathbf{x} at location s , and the available distribution, $F^A(s)$, is a function of location and is generally assumed to be constant (i.e., $F^A(s) = \frac{l}{|G|}$; Fieberg et al. 2021). In the denominator, we integrate over the geographical area available to the animal, G (Elith & Leathwick, 2009).

Instead of directly modeling $w(\mathbf{x}(s))$, the IPP models the density of points as the exponential of a linear function of spatial predictors – interpreted as the intensity function, $\lambda(s)$, in geographical space. We can parameterize $\lambda(s)$ similarly to $w(\mathbf{x}(s))$ from eqn 2:

Equation 4

$$\lambda(s) = \exp(\beta_0 + \beta_1 x_1(s) + \dots + \beta_k x_k(s)),$$

where s is the location in geographical space, $x_1(s), \dots, x_k(s)$ are k predictor habitat variables associated with location s , β_0 is an intercept term, and β_1, \dots, β_k , are the selection coefficients similar to eqn 2.

The *intercept* (β_0), when estimated, must be interpreted with care as it is often not biologically meaningful (i.e., it is the density of observations, but for movement data that may reflect frequency of observations; Fithian and Hastie 2012; Fieberg et al. 2021). As the number of available points sampled for the logistic regression grows to infinity, the *selection coefficients* (β_1, \dots, β_k) from eqn 2 (which are estimated using logistic regression), converge to the selection coefficients (β_1, \dots, β_k) from eqn 4 (which are estimated using an IPP, Warton and Shepherd 2010). As such, the selection coefficients of an RSF can be estimated by both methods.

Using movement data in a RSF can provide valuable information on animal relationships with environmental covariates. However, autocorrelation in movement data can introduce spatio-temporal autocorrelation that cannot be explained by habitat-related covariates. If not accounted for, this autocorrelation can result in underestimation of standard errors, potentially inflating Type I error rates (Nielsen et al., 2002; Northrup et al., 2013). This issue is particularly common when using high-resolution datasets (Hooten et al., 2013) and is expected to become more significant with the advancement of technology that enables increasingly fine-resolution movement data (Fieberg et al., 2010). Various approaches have been proposed to mitigate this problem, including data thinning (i.e., subsampling) as proposed by Hooten et al., (2013) (but see Alston et al., 2023 for a discussion on the effects of potential remaining pseudoreplication), correcting the standard errors (Koper & Manseau, 2012), and adding an autocorrelation term (e.g., intrinsic conditional autoregressive (ICAR) model, Johnson et al., 2013).

2.2 Step selection functions (SSF)

A step selection function (SSF) – also referred to as step selection *analysis* – builds on RSFs to estimate an animals’ resource selection with each selected location (Fortin et al., 2005; Rhodes et al., 2005; Thurfjell et al., 2014), providing a solution to the autocorrelation issues associated with RSFs. SSFs model the environmental conditions of observed sequential locations or ‘steps’, (i.e., the linear segment between two consecutive locations) relative to alternative random available steps or locations (drawn from a distribution of step lengths or ‘distance’ and turning angles) (Fig. 2, Forester et al., 2009; Fortin et al., 2005; Rhodes et al., 2005), and thereby allow for inference of step-scale resource selection.




Figure 2. Example of an animal movement track with linear “observed steps” between locations. Observed turn angle is relative to the trajectory from the preceding step. At each observed location, random “available” locations ($n = 3$; and their associated step lengths and turn angles) are generated.

There are many versions of SSFs, and in our review we use a commonly applied version often referred to as integrated-SSFs (Avgar et al., 2016) because it allows for environmental covariates to influence animal movement itself in addition to selection. Specifically, these SSFs model habitat selection and movement explicitly, by defining use distribution as the product of the habitat selection, w^* , and a selection-free movement kernel, ϕ (Avgar et al., 2016; Forester et al., 2009). The movement kernel is essentially equivalent to having the availability distribution of an RSF, $F^A(\mathbf{x})$, be temporally varying and it represents how the animal would move in the absence of habitat selection (Avgar et al., 2016; Forester et al., 2009). Thus, the probability density of an animal using location $s_{t+\Delta t}$ at time $t + \Delta t$ given that it was at location s_t at time t is modeled as:

Equation 5

$$F^U(s_{t+\Delta t} | s_t) = \frac{w^*(\mathbf{x}(s_{t+\Delta t}))\phi(s_{t+\Delta t}, s_t)}{\int_{s'_{t+\Delta t} \in G} w^*(\mathbf{x}(s'_{t+\Delta t}))\phi(s'_{t+\Delta t}, s_t) ds'_{t+\Delta t}},$$

where $w^*(\mathbf{x}(s_{t+\Delta t}))$ is the habitat selection function and $\phi(s_{t+\Delta t}, s_t)$ is the selection-free movement kernel (although this terminology is inconsistently used, see Michelot et al., 2023). As in eqn 1, the denominator integrates over the entire geographic domain G to normalize the use distribution. The parameters (β in the step-selection function and γ in the movement kernel) are dependent on Δt , thus, observed locations are typically recorded at a constant time interval.

These SSFs are often fitted using a conditional logistic regression. Including interactions between movement characteristics (including step length and turning angle) and environmental covariates relaxes the assumption that observed movement attributes (i.e., velocities and directional persistence) are independent of resource selection (Avgar et al., 2016; Forester et al.,

2009). Like RSFs, SSFs can also be fit using IPP or using Monte Carlo integration techniques (Michelot et al., 2023).

The selection-function lineage of modeling has evolved to consider the influence of scales on habitat relationships. However, choosing the number and the spatial extent of the availability sample is an important consideration and can affect the coefficient estimates and subsequently the ecological interpretation (Michelot et al., 2023; Northrup et al., 2013; Thurfjell et al., 2014). In RSFs and SSFs, the number of available samples can be based on a constant (e.g., 10,000, Aarts et al., 2012; Lele et al., 2013; Lele & Keim, 2006), point density (e.g., 1 available location per km², Hebblewhite & Merrill, 2008), or by a ratio of observed:available locations (e.g., 1:20, Northrup et al., 2013). Additionally, one can apply the RSF or SSF using a range of available sample size and chose the sample size where the coefficient estimates stabilized (i.e., Aarts et al., 2012; Fieberg et al., 2021; Northrup et al., 2013; Warton & Shepherd, 2010). As approximation accuracy increases with the size of the available sample, it is generally better to include a larger availability sample, especially given quick computation time (Michelot et al., 2023; Muff et al., 2020; Northrup et al., 2013).

2.3 Hidden Markov models (HMM)

In the context of movement ecology, hidden Markov models (HMMs) are principally used to classify movement into a finite number of discrete behavioural states (e.g., searching, resting, traveling), and differ conceptually from RSFs or SSFs as they generally do not make inference on habitat selection. However, HMMs can be used to investigate relationships between environmental

covariates and animal movement and behaviour (Glennie et al., 2023; Zucchini & MacDonald, 2009).

HMMs assume there is a hidden state process unfolding over time from which we obtain observation data (Fig. 3). That is, it is assumed that at any time, animals exhibit one of N discrete and temporally auto-correlated states (Z_1, \dots, Z_T) that represent hidden (unobserved) behaviours, where T is the total number of time steps. The hidden state process, Z_1, \dots, Z_T , is typically defined using a first-order Markov chain, where the probability of each state depends only on the state at the previous time (Fig. 3). These states are related to an observation time series (Y_1, \dots, Y_T). In contrast to the binary (observed/available) values used in the observation time series of RSFs and SSFs, the observation used in animal movement HMMs, generally comprise time series of step lengths and turning angles (Langrock et al., 2012; Morales et al., 2004) and can include additional data streams such as dive depth or acceleration (Ngô et al., 2019).




Figure 3. The structure of a hidden Markov model, where the hidden states (Z_t) depend only on the previous state, and the observed data (Y_t) depends on the hidden state. In the case of modeling animal movement, the observed process is composed of empirical data such as step lengths and

turning angles, and the hidden state may represent behaviours such as foraging, resting, and traveling.

For an N state HMM (e.g., $N = 3$ states, representing foraging, resting, and traveling), the state transition probability matrix (Γ) is a $N \times N$ matrix of $\gamma_{i,j}$, representing the probabilities of transitioning from each state to each other state. We define $\Gamma_t = (\gamma_{i,j})$, where $\gamma_{i,j}$ is the probability of switching from state i to state j :

Equation 6

$$\gamma_{i,j} = Pr(Z_{t+1} = j | Z_t = i),$$

and each row of Γ_t must sum to 1 (i.e., $\sum_{j=1}^N \gamma_{i,j} = 1$).

The observations are assumed to be independent of one another once we account for their dependence on the underlying hidden state, specifically the observation y_t at time t , is assumed to be dependent only on the state at time t :

Equation 7

$$Pr(y_t | Z_1, \dots, Z_t; y_1, \dots, y_{t-1}) = Pr(y_t | Z_t).$$

The conditional probability density, $P(y_t)$, is a $N \times N$ diagonal matrix:

Equation 8

$$P(y_t) = \text{diag}(p_1(y_t), \dots, p_N(y_t)),$$

where $p_i(y_t)$ represents the probability density of observation y given state i . When modeling animal movement, step length is often modeled with distributions such as gamma, exponential, or Weibull, while turning angle is modeled with circular distribution such as the von Mises or wrapped Cauchy (Langrock et al., 2012).

Taken together, the HMM likelihood function (L) is:

Equation 9

$$L = \delta P(y_1) \Gamma P(y_2) \Gamma P(y_3), \dots, \Gamma P(y_T) I'$$

where δ is the initial state distribution of the Markov chain, and I' is a row vector of ones. The parameters can be estimated using numerical likelihood maximization of this likelihood (eqn 9) or with the expectation-maximization (EM) algorithm (Zucchini & MacDonald, 2009).

Environmental covariates can be incorporated into two different parts of an HMM. First, to examine how covariates affect the probability to transition across states (e.g., how vegetation density may increase the probability to transition into foraging for an herbivore), they can be used to alter the transition probabilities, $\gamma_{i,j}$. Specifically, we can make transition probabilities, now denoted as $\gamma_{t,i,j}$, change at each time step as a function of a vector of environmental covariates (\mathbf{x}_t) and thus time t , e.g., via a multinomial logit link:

Equation 10

$$\gamma_{t,i,j} = \frac{\exp(\eta_{t,i,j})}{\sum_{l=1}^N \exp(\eta_{t,i,l})} ,$$

and

Equation 11

$$\eta_{t,i,j} = \beta_{0,i,j} + \beta_{1,i,j}x_{1,t} + \dots + \beta_{k,i,j}x_{k,t} ,$$

where $\beta_{0,i,j}$ is an intercept term and $\beta_{1,i,j}, \dots, \beta_{k,i,j}$ are the behaviour-specific regression coefficients for each of the k environmental covariates with values $\mathbf{x}_t = x_{1,t}, \dots, x_{k,t}$ at time t (McClintock et al., 2020; Michelot et al., 2016; Morales et al., 2004). To ensure model identifiability, $\beta_{0,i,j}$ and $\beta_{k,i,j}$ are fixed to 0 for one element in each row of the state transition probability matrix, typically, the diagonal when $i = j$.

Second, environmental covariates (typically abiotic) can be used to alter the observation probabilities, $p_i(x)$, which can be interpreted as examining how they affect the movement of the animal within a state (e.g., how snow depth affects the speed of a walking animal). To do so, the observation probability density functions will change through time, now denoted $p_{t,i}(x)$, and their parameters will depend on environmental covariates. For example, environmental covariates can affect step length mean $\mu_{t,i}^{(l)}$ and standard deviation $\sigma_{t,i}^{(l)}$, and turning angle concentration $\kappa_{t,i}^{(\theta)}$ for each state i at each time step t as

Equation 12

$$\ln(\mu_{t,i}^{(l)}) = \beta_{i,0}^{(\mu^{(l)})} + \beta_{l,i,j}^{(\mu^{(l)})} x_{l,t} + \dots + \beta_{k,i,j}^{(\mu^{(l)})} x_{k,t} ,$$

Equation 13

$$\ln(\sigma_{t,i}^{(l)}) = \beta_{i,0}^{(\sigma^{(l)})} + \beta_{l,i,j}^{(\sigma^{(l)})} x_{l,t} + \dots + \beta_{k,i,j}^{(\sigma^{(l)})} x_{k,t} , \text{ and}$$

Equation 14

$$\ln(\kappa_{t,i}^{(\theta)}) = \beta_{i,0}^{(\kappa^{(\theta)})} + \beta_{l,i,j}^{(\kappa^{(\theta)})} x_{l,t} + \dots + \beta_{k,i,j}^{(\kappa^{(\theta)})} x_{k,t} , \text{ respectively.}$$

Movement with orientation bias (i.e., biased random walks) can be modeled by defining parameters (b) such as the turning angle mean as a function of angular environmental covariates ψ_k (e.g., angle between animal orientation and ocean current):

Equation 15

$$\mu_{i,t}^{(\phi)} = \text{atan2}(\sum_{b=1}^k \beta_{i,k}^{\mu^{(\theta)}} \sin(\psi_{k,t}), \sum_{b=1}^k \beta_{i,k}^{\mu^{(\theta)}} \cos(\psi_{k,t})).$$

In addition to modeling bias directly toward or away from target, including both $\psi_{k,t}$ and $\psi_{k,t} + \frac{\pi}{2}$ as covariates can model movement with bias toward any angle relative to stimulus (e.g., olfactory search perpendicular to wind, Togunov et al., 2022).

Hidden Markov models are a specific type of state-space models (Auger-Méthé et al., 2021), where the states are discrete (i.e., there is a finite number of hidden states). Other state-

space models have been developed that describe animal movement behaviour as a continuum of move-persistence, γ_t , ranging from 0 to 1, where values closer to 0 are indicative of less move persistence (i.e., area-restricted search), and values closer to 1 are indicative of more persistent movement (i.e., traveling; Breed et al., 2012). Move-persistence can be modeled as a linear function of environmental predictors measured at each location,

Equation 16

$$\text{logit}(\gamma_t) = \beta_0 + \beta_1 x_{t,1} + \dots + \beta_k x_{t,k} + \varepsilon_t,$$

where $\varepsilon_t = N(0, \sigma_\gamma)$ (Jonsen et al., 2019). Such state-space models may be a more natural representation of animal movement as they provide an estimate of the “intensity” of the behaviour (Breed et al., 2012, e.g., onset of migration can be identified when move-persistence, γ_t , exceeds a threshold, Shuert et al., 2022). However, such state-space models assume that behaviour exists on a linear continuum of only two behaviours (e.g., area-restricted search to traveling; e.g., Gryba et al., 2019), and it is unclear where other behaviours (e.g., resting) would fall along that continuum (McClintock et al., 2014). Thus, models in discrete state-space may be easier to interpret, as each state has defined respective turning angle and step length distributions (McClintock et al., 2014).

3. Case study: Ringed seals in Hudson Bay, Canada

3.1 Background and methods

We used movement data of one ringed seal (*Pusa hispida*) to conduct a comparative analysis of different methods to characterize their relationship with one covariate. We chose prey diversity as the covariate of interest for exploration, given that ringed seals are known as opportunistic predators (Appendix 2; data from Florko et al., 2021a; Florko et al., 2021b). Providing an accurate description of how ringed seals interact with their environment is important because of their ecological and cultural significance. Ringed seals are facing a broad-scale demographic decline (Ferguson et al., 2017) underscoring the urgency to identify important areas of occupancy for this species.

We analyzed the estimated movement of a seal (Fig 4a) equipped with an ARGOS satellite telemetry transmitter from 29 Oct 2012 to 17 Mar 2013 in Hudson Bay, Canada (see Florko et al., 2023 for details). We first applied a state-space model to estimate more accurate locations and regularize the time interval to one location per day ($n = 140$, Patterson et al., 2010). Then, we matched each seal location with prey diversity in the corresponding grid cell (Fig. 4D, and used the prey diversity data as a covariate in an RSF, two SSFs (one with and one without movement-related covariates), and an HMM (in the transition probabilities, but see Appendix 4 for how to apply covariates to the observation probabilities). The selection functions included ten available locations per observed location. The HMM included three states which best captured variation in step length and turning angle (see Appendix 4 for two-state HMMs). We predicted the probability of use relative to prey diversity (log of the relative selection strength [\log -RSS] for selection functions) and in space (steady-state utilization distributions). Detailed methods are available in Appendix 2 and the data and a coding tutorial of the analysis are available in Appendix 3 and 4 (respectively).




Figure 4. The observed (used) and available locations for the A) resource selection function (RSF) and B) step-selection functions (SSF). Purple circles represent the observed locations and purple lines denote the path between observed points. Peach circles represent the available locations, although note that for the SSF, only a subset of these is available at a given time (see Fig 2). The RSF availability sample was generated randomly within the minimum convex polygon of the track. C) Decoded states from the hidden Markov model (HMM), including slow movement (green squares), moderate movement (blue circles), and fast movement (purple triangles). D) Ringed seal

estimated locations (yellow circles) and straight-line path between locations (yellow line) overlaid on estimated prey diversity.

3.2 Results and discussion

These models led to important differences in the magnitude, and sometimes direction, of the relationships with the same covariate, and thus resulted in different ecological inference. The estimated selection coefficient that represented the habitat selection relationship with prey diversity was positive for the RSF, but not significantly different from zero for the SSFs (Table S1, Fig. 5A-B). The different selection coefficient estimates between the RSF and the SSFs may be attributed to their distinct definition of available habitat. The RSF assumes the entire study area (within the minimum convex polygon (MCP) in this case study) is accessible to the animal, whereas the habitat available in an SSF changes at each time step and is limited to an area that the animal can reach within a time step (Fig. 4A-B). For the HMM, the probability of being in the slow behaviour, which is often of importance for conservation because it is thought to be associated with foraging (Pyke et al., 1977, but see Florke et al., 2023), had a positive relationship with prey diversity (Fig. 5C). The SSF with movement-related covariates showed no significant relationship between prey diversity and movement speed (Table S1), and as a result the relationship between selection and prey diversity was not significantly modified (Fig. 5B). Consequently, if these results were being used for conservation, the RSF and HMM would suggest that high prey diversity is important for ringed seal space use and important behaviours, while the SSFs would suggest that this covariate is not important. The HMM characterized the states mostly on movement speed (i.e., step length, Fig. S1), and the states had contrasting relationships with prey diversity (Fig. 5C). The probability of being in the moderate speed behaviour decreased with

prey diversity, while the probability of being in fast behaviour remained relatively constant with prey diversity, with a potential peak in low-medium prey diversity (Fig. 5C). The RSF and HMM slow movement state likely show similar results because slow movement is associated with greater residency (i.e., remaining in an area) and thus will result in more locations in an area. Note also that the RSF and SSFs model covariates linearly by default (although transformations can be applied) whereas the HMM models covariates to the transition probability matrix and are non-linear due to the multinomial link function (eqn. 10).




Figure 5. Log relative selection strength (log-RSS) from the A) RSF and SSF without movement-related covariates and B) SSF with movement-related covariates, where prey diversity affected step length; speeds were quantified as the 25th (slow), 50th (moderate), and 75th (fast) percentiles of step length, which we infer as “speed” since the locations are recorded at a fixed time interval. C) Estimated state probabilities from the HMM as a function of prey diversity (on the transition probability). Shaded areas represent the standard error. Note the bowtie shape of the standard error in A-B is due to log-RSS calculating selection strength *relative* to a starting step, in this case, where prey diversity is set to its mean.

Differences in coefficient estimates will affect maps of predicted probability of seal occurrence or behaviour (selection function or HMM, respectively; Fig. 6). Given the model's positive coefficient estimate with prey diversity, the RSF predicted that seals are most often found in a small portion of the study area where there is high prey diversity (Fig. 6A). The SSFs did not have significant relationships with prey diversity, thus, the maps showed that probability of use was more uniformly distributed in space (Fig. 6B-C). The HMM will produce one map per state (behaviour), and here with vastly contrasting spatial patterns. The slow movement state, which was positively related to prey diversity, showed a similar restricted spatial pattern to the RSF (Fig. 6D), whereas the moderate movement state, which had a negative relationship with prey diversity, showed the opposite pattern to the RSF prediction map (Fig. 6E). The fast movement state did not show much spatial variation (Fig. 6F), reflective of its non-significant relationship with prey diversity (Fig. 5F). The SSF-generated maps did not show any similarities with the other models' maps (or relative to prey diversity), likely due to prey diversity not being relevant at the step-scale. However, note that the SSF-generated maps are plotted at a different scale than the others, and that if the fast movement state map was on the same scale as the SSFs, it would likely present comparable results. Overall, we see that prey diversity is relevant at the home-range scale (RSF) and transition into states at the whole track-level scale (HMM). This demonstrates that choosing the wrong model for the question at hand can mislead ecological insight and could lead to damaging conservation and management decisions.




Figure 6. Estimated utilization distribution from A) resource selection function (RSF), B) step selection function (SSF) without movement-related covariates, C) SSF with movement-related covariates, and hidden Markov model (HMM) D) state 1: slow movement, E) state 2: moderate movement, and F) state 3: fast movement. Note that B and C were created using simulated paths with 10,000 locations.

In this case study, selection at the home range scale modeling approach (i.e., RSF) may be most relevant due to our goal of broadly identifying important areas for this ringed seal (see section 4: *Choosing a model*), and also given the coarse resolution of the prey data. However, to simplify comparisons across models, this case study was limited to one seal, one prey metric, and used the default settings for these models. Thus, to reflect realistic ecological dynamics, an


analysis of a more comprehensive ringed seal dataset (e.g., more individuals, longer tracking, and more prey metrics) should be performed in the future.

4. Choosing a model

By design, RSFs are well-suited to address large-scale questions on important areas for species (e.g., second-order selection; at the home range scale), whereas SSFs and HMMs are typically well-suited to address smaller-scale movement- or behaviour-specific questions (Box 1, e.g., third-order selection; at the habitat usage scale; Johnson, 1980; Nams, 2013; Owen, 1972; Sidrow et al., 2022). Since RSFs generate the availability sample within the home range (MCP) of the observed data, and SSFs generate the availability sample for each step within the range of possible steps (Avgar et al., 2016) the SSF prediction coefficients provide insight at a smaller scale than in the large-scale (home range) case of RSFs. Thus, predictions from estimated coefficients from SSFs (i.e., for prediction maps) can be misleading since the coefficients represent the probability of selection at the scale of individual steps by an animal, rather than habitat selection at the scale of the home range (Fieberg et al., 2021). In contrast, HMMs classify movement into discrete behavioural states, and thus do not consider habitat covariates as a driver of habitat selection, rather, as a mechanism related to estimated behavioural states (Glennie et al., 2023; Zucchini & MacDonald, 2009).

BOX 1. Which model to choose?

Decision tree for choosing an appropriate modeling approach for understanding animal relationships with environmental conditions using movement data, including if you are interested in how the environmental conditions promote occupancy, influence animal behaviour, or influence animal movement. For example, caribou movement in the winter could be related to snow depth. RSFs can be used for broad-scale inference relating caribou occupancy and snow depth, SSFs can be used to understand whether snow depth affect selection at finer scale, HMMs can identify the depths of snow associated with a behaviour (e.g., resting), and HMMs or SSFs can be used to assess whether snow depth slows down movement speed.



4.1. Examples of model application in ecology and conservation

Applying RSFs to animal movement has been particularly useful in conservation and to understand the spatial ecology of animals. For example, RSFs have been used to identify broad conservation corridors (Chetkiewicz & Boyce, 2009) and high wildlife density areas to improve rabies vaccination programs (McClure et al., 2022). Additionally, RSFs have been used to understand how habitat selection is affected by seasons (Martin et al., 2021) anthropogenic developments (Knopff et al., 2014), and the presence of predators (Florko et al., 2023), and can be used to understand the cumulative effects of such factors (Darlington et al., 2022).

SSFs have been used for understanding how environmental covariates affect animal movement characteristics (Squires et al., 2013). For example, SSFs have been used to show how roads increase step lengths (Roever et al., 2010). SSFs have also been used to understand how predators or social networks influence movement (Fortin et al., 2005; Webber et al., 2021). Further, SSFs have been used to identify that animals move slower and select shadier areas near mixed forests in high temperatures (Alston et al., 2020). Additionally, SSFs have been used to understand how slopes associated with thermal uplift affects soaring bird movement patterns, including potential stopover movements to access carrion (Eisaguirre et al., 2020).

Hidden Markov models provide another approach to identify habitat features relevant to animal movement (e.g., corridors) due to their ability to separate traveling behaviours (McClintock et al., 2020). HMMs have been used to quantify how foraging probability relates to canopy cover (Gardiner et al., 2019), how travel speed relates to snow depth (Chimienti et al., 2021), and how animal orientation relates to wind direction (Togunov et al., 2022). Further, HMMs have been used to show how wild dog movements were affected by anthropogenic disturbance within and outside of a National Park (Creel et al., 2020).

These models can be easily extended to incorporate inter- or intra-specific relationships. Most simply, interspecific relationships can be included by adding another species' probability of occupancy as a covariate (e.g., Florko et al., 2023; Muff et al., 2020; Warton et al., 2015). Additionally, more complex and dynamic intraspecific relationships can be included by

considering individual animal's movement relative to the group movement (see Langrock et al., 2014 for an HMM example).

Maps of probability of use are most easily generated using RSFs (Boyce & McDonald, 1999) and are particularly useful for conservation management and land-use planning (Chetkiewicz & Boyce, 2009; Gerber & Northrup, 2020; McLoughlin et al., 2010). Maps can be made using SSFs and HMMs (Signer et al., 2017, 2019), however the tools used to create these maps are less documented, and for SSFs, require time-intensive path simulation exercises with setting choices that are indistinct (e.g., choosing the number of tracks and steps in the simulation, the burn-in size, although see our step-by-step tutorial: Appendix 4).

4.3. Important considerations

Choosing an appropriate model can be difficult, and choosing an adequate model does not guarantee appropriate inference. For example, Buderman et al., 2021 used HMMs to identify white tailed deer (*Odocoileus virginianus*) male-female interaction events, but upon validation, found that the model incorrectly identified interactions and thus mislead inference about mating behaviours. Additionally, Florko et al., 2023 used move-persistence models similar to eqn. 16 to identify relationships between foraging behaviour and prey abundance, and found important considerations for matching foraging behaviour, prey biomass, and diving behaviour. These cautionary tales highlight the importance of understanding the chosen models' design, abilities, and limitations (Box 2). Where possible, validation exercises (e.g., leave-one-individual-out cross-validation, Boyce et al., 2002) and incorporation of additional data streams (e.g., dive data, Gryba et al., 2019) is recommended.

Selection functions are subject to several assumptions. Selection functions assume that animals make resource-driven choices where at each location they choose the best habitat to move to next, but this may not always be the case (e.g., as in migration, or when relying on memory). RSFs assume that locations are independent, and methods have been proposed to circumvent the violation of this assumption (e.g., data thinning, Northrup et al., 2022). Similarly, SSFs assume that the step length and turn angle are independent, which may not be true given the scale (e.g., animals may keep foraging in nearby locations; Hodel & Fieberg, 2022; Morales et al., 2004). Thus, sampling intervals need to be examined because small time intervals can lead to misleadingly narrow confidence intervals (Alston et al., 2023; Avgar et al., 2016). While many RSF and SSF models assume that individuals exhibit identical movement and selection, individual variation can be included in selection functions via random slopes in a mixed model (e.g., using the `glmmTMB` package in R, Muff et al., 2020) or by using a two-step approach of fitting individual models and averaging parameter estimates for inference at the population level (e.g., using the `TwoStepClogit` package in R, Fieberg et al., 2010; Squires et al., 2013).

Movement HMMs relax many assumptions of selection functions (RSF, SSF). HMMs acknowledge that telemetry data encompass distinct, unobserved behaviours and distinguish the state process from the observation process. HMMs also acknowledge that telemetry data are not independent, and explicitly model autocorrelation through the state process. Movement patterns and habitat associations are behaviour-specific, which can be modeled by the state-dependent distribution of the observation process and state transition probabilities.

BOX 2. Model comparison chart. A summary comparison of statistical models that use movement data to characterize species-habitat associations. Note that this table highlights the main intended use, assumptions, strengths, and weaknesses associated with each model and their implementation and is not an exhaustive list.

Intended use(s)	Main assumption(s)	Strengths	Considerations in implementation
Resource selection function (RSF)			
Relate environmental covariates to animal occurrence at home-range (large) scale	Habitat selection depends on the encountered environmental conditions; The probabilities of selection is constant during the period of investigation; Locations are independent ¹	Simple to use and interpret; Popular (can compare results to other studies)	Challenges in choosing the quantity of points for the availability sample ² and the extents of the available sample; Ignores potential movement barriers (i.e., the complete home range is assumed to be available to the animal) ³
Step selection function (SSF)			

<p>Relate environmental covariates to animal occurrence at a movement (small) scale;</p> <p>Can assess how movement characteristics (i.e., step length) are affected by environmental covariates[†]</p>	<p>Habitat selection is conditional on occurrence within a restricted range (i.e., available step-length is limited by observed movement);</p> <p>Habitat selection can be affected by environmental conditions[†]</p>	<p>Relatively easy to use;</p> <p>Accounts for autocorrelation;</p> <p>Creates availability samples based on observed movement patterns;</p> <p>Allows for simultaneous inference on both movement and habitat selection[†]</p>	<p>Challenges in choosing the quantity of points for the availability sample²;</p> <p>Selection of interval between movement locations must meet assumptions of independence;</p> <p>Scale of steps dictates habitat availability and the types of behaviours that can be modeled;</p> <p>Interpretation (e.g., prediction maps) is more nuanced because selection is relative to observed points (even more challenging when modeling how habitat affects movement[†]);</p>
--	---	--	--

Hidden Markov model (HMM)

<p>Assess how movement characteristics</p>	<p>Behaviour at time t depends only on behaviour at $t-1$</p>	<p>Ability to identify behavioural states and how their</p>	<p>Sampling interval dictates the types of behaviours that can be modeled;</p>
--	---	---	--

are affected by environmental covariates; Identify environmental covariates that promote certain behavioural states	(Markov property) ⁴ ; The distribution of observations (e.g. step length and turning angle) depend on the latent behaviour (state) but not the previous observations ⁴	occurrence is associated with environmental covariates; Ability to quantify how movement characteristics (e.g. step length) are affected by environmental covariates; State sequence can identify when and where behaviours occur	Limited to relationship with environmental condition experienced by the animals tagged, is not designed for habitat selection
--	---	---	---

¹In typical usage. Solutions exist in including an autocorrelation term in the model (e.g., when using integrated nested Laplace approximation [INLA], D. S. Johnson et al., 2013).

²In typical usage, but see Muff et al. (2020; Northrup et al., 2013).

³In typical usage, but see Brost et al. (2015).

⁴In typical usage, but all these assumptions have been relaxed in some form in the literature (e.g.,

semi-Markov model, (Langrock et al., 2012), conditionally autoregressive HMM, (Lawler et al., 2019), hierarchical HMM, (Sidrow et al., 2022)).

[†]Relevant if the SSF models the influence of habitat covariates on the movement kernel.

5. Future directions and conclusions

Selection functions and HMMs are widely applied models that provide essential insight from animal movement data, and several new and extended versions of these models provide deeper insight, although mainstream packages to implement these models are not yet available. For example, recent models have been developed that incorporate habitat availability into HMMs (termed HMM-SSF) and can give insight on behaviour-specific habitat selection (e.g., Klappstein et al., 2023; Nicosia et al., 2017; Prima et al., 2022). Additionally, there are a variety of machine learning approaches (e.g., MaxEnt, random forests) that we have not covered because our review is focused on statistical methods. However, it is important to note that MaxEnt is considered equivalent to an exponential RSF, and generally may be better suited for occupancy data (i.e., a location can only be used once, as in species distribution data, Elith et al., 2011), but can also be applied to movement data (Lele et al., 2013). Additionally, random forests may be used for modeling animal movement and generally relax the assumptions associated with parametric methods, but the results may be more challenging to interpret (Wijeyakulasuriya et al., 2020). If prediction is the goal, machine learning approaches including MaxEnt and random forests, or a parametric approach using LASSO for model selection, are often recommended (Elith & Leathwick, 2009; Gerber & Northrup, 2020).

The models reviewed in this paper for linking animals' movement data to the environment are widely applicable and can be easily implemented but provide different ecological insights. Our review highlights the conceptual differences that will aid ecologists in choosing a model based on their goals. For example, RSFs may be well suited for identifying broad corridors or protected areas, SSFs for understanding movement patterns, and HMM for understanding behaviours. Particularly, we highlight that RSFs are the most broad-scale model and are useful for home-range studies, whereas SSFs and ultimately HMMs get increasingly more movement-specific. Also, we show how RSFs show similar results to HMM states characterized by slow movement (step length). Further, the sampling – or exclusion as in the case of HMMs – of availability locations can influence the coefficient estimates, as can the inclusion of movement information in the model. All these models are invaluable for ecological research, yet careful consideration of the mathematical and conceptual underpinnings is necessary to address the ecological questions and applied conservation management that these results may inform.

Acknowledgements

We gratefully acknowledge the Sanikiluaq hunters, the Sanikiluaq Hunters and Trappers Association, and especially Lucassie and Johnnassie Ippak and their families for assistance in the field for the seal data used in the case study. We acknowledge the financial support from Fisheries and Oceans Canada, Nunavut Wildlife Management Board, ArcticNet, Natural Sciences and Engineering Research Council of Canada (NSERC), Ocean Leaders Graduate Fellowship, Canada Research Chairs program, Canada Foundation for Innovation, B.C. Knowledge Development Fund, Weston Family Foundation, Northern Scientific Training Program, Polar Knowledge

Canada, Earth Rangers, Social Sciences and Humanities Research Council. Animal handling was approved by the Freshwater Institute Animal Care Committee (FWI-ACC-2012) and Fisheries and Oceans Canada's Licenses to Fish for Scientific Purposes (S-05/06-09/13-1006-NU). Note that MAM is an Associate Editor for *Methods in Ecology and Evolution*.

Author Contributions

KRNF, RRT, and MAM designed the study. KRNF did the analysis and prepared figures and tables. KRNF wrote the manuscript with contributions from RRT, RG, ES, and MAM. KRNF wrote the code and Github tutorial with contributions from RRT and MAM. DJY and SHF collected the seal data. All authors contributed ideas and revised the manuscript.

Data Availability Statement

The data and code used in the case study are included as Appendix 3 and 4, respectively, and also available on Github (https://github.com/kflorko/movementstats_review)

References

- Aarts, G., Fieberg, J., & Matthiopoulos, J. (2012). Comparative interpretation of count, presence–absence and point methods for species distribution models. *Methods in Ecology and Evolution*, 3(1), 177–187.
- Alston, J. M., Fleming, C. H., Kays, R., Streicher, J. P., Downs, C. T., Ramesh, T., Reineking, B., & Calabrese, J. M. (2023). Mitigating pseudoreplication and bias in resource selection

- functions with autocorrelation-informed weighting. *Methods in Ecology and Evolution*, *14*(2), 643–654.
- Alston, J. M., Joyce, M. J., Merkle, J. A., & Moen, R. A. (2020). Temperature shapes movement and habitat selection by a heat-sensitive ungulate. *Landscape Ecology*, *35*, 1961–1973.
- Andrewartha, H. G., & Browning, T. O. (1961). An analysis of the idea of “resources” in animal ecology. *Journal of Theoretical Biology*, *1*(1), 83–97.
- Auger-Méthé, M., Newman, K., Cole, D., Empacher, F., Gryba, R., King, A. A., Leos-Barajas, V., Mills Flemming, J., Nielsen, A., & Petris, G. (2021). A guide to state–space modeling of ecological time series. *Ecological Monographs*, *91*(4), e01470.
- Avgar, T., Potts, J. R., Lewis, M. A., & Boyce, M. S. (2016). Integrated step selection analysis: bridging the gap between resource selection and animal movement. *Methods in Ecology and Evolution*, *7*(5), 619–630.
- Bachl, F. E., Lindgren, F., Borchers, D. L., & Illian, J. B. (2019). inlabru: an R package for Bayesian spatial modelling from ecological survey data. *Methods in Ecology and Evolution*, *10*(6), 760–766.
- Boyce, M. S., & McDonald, L. L. (1999). Relating populations to habitats using resource selection functions. *Trends in Ecology & Evolution*, *14*(7), 268–272.
- Boyce, M. S., Vernier, P. R., Nielsen, S. E., & Schmiegelow, F. K. A. (2002). Evaluating resource selection functions. *Ecological Modelling*, *157*(2–3), 281–300.
- Breed, G. A., Costa, D. P., Jonsen, I. D., Robinson, P. W., & Mills-Flemming, J. (2012). State-space methods for more completely capturing behavioral dynamics from animal tracks. *Ecological Modelling*, *235*, 49–58.

- Brost, B. M., Hooten, M. B., Hanks, E. M., & Small, R. J. (2015). Animal movement constraints improve resource selection inference in the presence of telemetry error. *Ecology*.
- Buderman, F. E., Gingery, T. M., Diefenbach, D. R., Gigliotti, L. C., Begley-Miller, D., McDill, M. M., Wallingford, B. D., Rosenberry, C. S., & Drohan, P. J. (2021). Caution is warranted when using animal space-use and movement to infer behavioral states. *Movement Ecology*, 9(1), 1–12.
- Chetkiewicz, C.-L. B., & Boyce, M. S. (2009). Use of resource selection functions to identify conservation corridors. *Journal of Applied Ecology*, 1036–1047.
- Chimienti, M., van Beest, F. M., Beumer, L. T., Desforges, J. P., Hansen, L. H., Stelvig, M., & Schmidt, N. M. (2021). Quantifying behavior and life-history events of an Arctic ungulate from year-long continuous accelerometer data. *Ecosphere*, 12(6).
<https://doi.org/10.1002/ecs2.3565>
- Creel, S., Merkle, J., Mweetwa, T., Becker, M. S., Mwape, H., Simpamba, T., & Simukonda, C. (2020). Hidden Markov Models reveal a clear human footprint on the movements of highly mobile African wild dogs. *Scientific Reports*, 10(1), 17908.
- Darlington, S., Ladle, A., Burton, A. C., Volpe, J. P., & Fisher, J. T. (2022). Cumulative effects of human footprint, natural features and predation risk best predict seasonal resource selection by white-tailed deer. *Scientific Reports*, 12(1), 1072.
- Eisaguirre, J. M., Booms, T. L., Barger, C. P., Lewis, S. B., & Breed, G. A. (2020). Novel step selection analyses on energy landscapes reveal how linear features alter migrations of soaring birds. *Journal of Animal Ecology*, 89(11), 2567–2583.

- Elith, J., & Leathwick, J. R. (2009). Species distribution models: ecological explanation and prediction across space and time. *Annual Review of Ecology, Evolution, and Systematics*, *40*, 677–697.
- Elith, J., Phillips, S. J., Hastie, T., Dudík, M., Chee, Y. E., & Yates, C. J. (2011). A statistical explanation of MaxEnt for ecologists. *Diversity and Distributions*, *17*(1), 43–57.
- Ferguson, S. H., Young, B. G., Yurkowski, D. J., Anderson, R., Willing, C., & Nielsen, O. (2017). Demographic, ecological and physiological responses of ringed seals to an abrupt decline in sea ice availability. *PeerJ, February*. <https://doi.org/10.7717/peerj.2957>
- Fieberg, J., Matthiopoulos, J., Hebblewhite, M., Boyce, M. S., & Frair, J. L. (2010). Correlation and studies of habitat selection: problem, red herring or opportunity? *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1550), 2233–2244.
- Fieberg, J., Signer, J., Smith, B., & Avgar, T. (2021). A ‘How to’ guide for interpreting parameters in habitat-selection analyses. *Journal of Animal Ecology*, *90*(5), 1027–1043.
- Fithian, W., & Hastie, T. (2012). Statistical models for presence-only data: finite-sample equivalence and addressing observer bias. *Ann Appl Stat.*
- Florko, K. R. N., Ross, T. R., Ferguson, S. H., Northrup, J. M., Obbard, M. E., Thiemann, G. W., Yurkowski, D. J., & Auger-Methe, M. (2023). The dynamic interaction between predator and prey drives mesopredator movement and foraging ecology. *BioRxiv*, 2023.04.27.538582. <https://doi.org/10.1101/2023.04.27.538582>
- Florko, K. R. N., Shuert, C. R., Cheung, W. W. L., Ferguson, S. H., Jonsen, I. D., Rosen, D. A. S., Sumaila, U. R., Tai, T. C., Yurkowski, D. J., & Auger-Méthé, M. (2023). Linking movement and dive data to prey distribution models: new insights in foraging behaviour and potential pitfalls of movement analyses. *Movement Ecology*, *11*(1), 17.

- Florko, K. R. N., Tai, T. C., Cheung, W. W. L., Ferguson, S. H., Sumaila, U. R., Yurkowski, D. J., & Auger-Méthé, M. (2021). Predicting how climate change threatens the prey base of Arctic marine predators. *Ecology Letters*, *24*, 2563–2575. <https://doi.org/10.1111/ele.13866>
- Florko, K. R. N., Tai, T. C., Cheung, W. W. L., Sumaila, U. R., Ferguson, S. H., Yurkowski, D. J., & Auger-Méthé, M. (2021). Predicting how climate change threatens the prey base of Arctic marine predators. *Dryad, dataset*. <https://doi.org/https://doi.org/10.5061/dryad.x69p8czjs>
- Forester, J. D., Im, H. K., & Rathouz, P. J. (2009). Accounting for animal movement in estimation of resource selection functions: sampling and data analysis. *Ecology*, *90*(12), 3554–3565.
- Fortin, D., Beyer, H. L., Boyce, M. S., Smith, D. W., Duchesne, T., & Mao, J. S. (2005). Wolves influence elk movements: Behavior shapes a trophic cascade in Yellowstone National Park. *Ecology*, *86*(5), 1320–1330. <https://doi.org/10.1890/04-0953>
- Gardiner, R., Hamer, R., Leos-Barajas, V., Peñaherrera-Palma, C., Jones, M. E., & Johnson, C. (2019). State-space modeling reveals habitat perception of a small terrestrial mammal in a fragmented landscape. *Ecology and Evolution*, *9*(17), 9804–9814.
- Gerber, B. D., & Northrup, J. M. (2020). Improving spatial predictions of animal resource selection to guide conservation decision making. *Ecology*, *101*(3), e02953.
- Glennie, R., Adam, T., Leos-Barajas, V., Michelot, T., Photopoulou, T., & McClintock, B. T. (2023). Hidden Markov models: Pitfalls and opportunities in ecology. *Methods in Ecology and Evolution*, *14*(1), 43–56.
- Godvik, I. M. R., Loe, L. E., Vik, J. O., Veiberg, V., Langvatn, R., & Mysterud, A. (2009). Temporal scales, trade-offs, and functional responses in red deer habitat selection. *Ecology*, *90*(3), 699–710.

- Gryba, R. D., Wiese, F. K., Kelly, B. P., Von Duyke, A. L., Pickart, R. S., & Stockwell, D. A. (2019). Inferring foraging locations and water masses preferred by spotted seals *Phoca largha* and bearded seals *Erignathus barbatus*. *Marine Ecology Progress Series*, *631*, 209–224.
- Hebblewhite, M., & Merrill, E. (2008). Modelling wildlife–human relationships for social species with mixed-effects resource selection models. *Journal of Applied Ecology*, *45*(3), 834–844.
- Hodel, F. H., & Fieberg, J. R. (2022). Circular–linear copulae for animal movement data. *Methods in Ecology and Evolution*, *13*(5), 1001–1013.
- Hooten, M. B., Hanks, E. M., Johnson, D. S., & Alldredge, M. W. (2013). Reconciling resource utilization and resource selection functions. *Journal of Animal Ecology*, *82*(6), 1146–1154.
- Hooten, M. B., Lu, X., Garlick, M. J., & Powell, J. A. (2020). Animal movement models with mechanistic selection functions. *Spatial Statistics*, *37*, 100406.
- Huey, R. B. (1991). Physiological consequences of habitat selection. *The American Naturalist*, *137*, S91–S115.
- Hussey, N. E., Kessel, S. T., Aarestrup, K., Cooke, S. J., Cowley, P. D., Fisk, A. T., Harcourt, R. G., Holland, K. N., Iverson, S. J., & Kocik, J. F. (2015). Aquatic animal telemetry: a panoramic window into the underwater world. *Science*, *348*, 1255642.
- Johnson, C. J., Nielsen, S. E., Merrill, E. H., McDONALD, T. L., & Boyce, M. S. (2006). Resource selection functions based on use-availability data: theoretical motivation and evaluation methods. *The Journal of Wildlife Management*, *70*(2), 347–357.
- Johnson, D. H. (1980). The comparison of usage and availability measurements for evaluating resource preference. *Ecology*, *61*(1), 65–71.
- Johnson, D. S., Hooten, M. B., & Kuhn, C. E. (2013). Estimating animal resource selection from telemetry data using point process models. *Journal of Animal Ecology*, *82*(6), 1155–1164.

- Jonsen, I. D., McMahon, C. R., Patterson, T. A., Auger-Méthé, M., Harcourt, R., Hindell, M. A., & Bestley, S. (2019). Movement responses to environment: fast inference of variation among southern elephant seals with a mixed effects model. *Ecology*, *100*(1), 1–8.
<https://doi.org/10.1002/ecy.2566>
- Joo, R., Boone, M. E., Clay, T. A., Patrick, S. C., Clusella-Trullas, S., & Basille, M. (2020). Navigating through the R packages for movement. *Journal of Animal Ecology*, *89*(1), 248–267.
- Klappstein, N. J., Thomas, L., & Michelot, T. (2023). Flexible hidden Markov models for behaviour-dependent habitat selection. *Movement Ecology*, *11*(1), 30.
- Knopff, A. A., Knopff, K. H., Boyce, M. S., & Clair, C. C. S. (2014). Flexible habitat selection by cougars in response to anthropogenic development. *Biological Conservation*, *178*, 136–145.
- Koper, N., & Manseau, M. (2012). A guide to developing resource selection functions from telemetry data using generalized estimating equations and generalized linear mixed models. *Rangifer*, *32*(2), 195–204.
- Langrock, R., Hopcraft, J. G. C., Blackwell, P. G., Goodall, V., King, R., Niu, M., Patterson, T. A., Pedersen, M. W., Skarin, A., & Schick, R. S. (2014). Modelling group dynamic animal movement. *Methods in Ecology and Evolution*, *5*(2), 190–199.
- Langrock, R., King, R., Matthiopoulos, J., Thomas, L., Fortin, D., & Morales, J. M. (2012). Flexible and practical modeling of animal telemetry data: hidden Markov models and extensions. *Ecology*, *93*(11), 2336–2342.
- Lawler, E., Whoriskey, K., Aeberhard, W. H., Field, C., & Mills Flemming, J. (2019). The conditionally autoregressive hidden Markov model (carhmm): Inferring behavioural states

from animal tracking data exhibiting conditional autocorrelation. *Journal of Agricultural, Biological and Environmental Statistics*, 24, 651–668.

Lele, S. R., & Keim, J. L. (2006). Weighted distributions and estimation of resource selection probability functions. *Ecology*, 87(12), 3021–3028.

Lele, S. R., Merrill, E. H., Keim, J., & Boyce, M. S. (2013). Selection, use, choice and occupancy: clarifying concepts in resource selection studies. *Journal of Animal Ecology*, 82(6), 1183–1191.

Manly, B. F. L., McDonald, L., & Thomas, D. L. (1993). *Resource Selection by Animals: Statistical Design and Analysis for Field Studies*. Chapman & Hall.

Martin, M. E., Moriarty, K. M., & Pauli, J. N. (2021). Landscape seasonality influences the resource selection of a snow-adapted forest carnivore, the Pacific marten. *Landscape Ecology*, 36, 1055–1069.

Matthiopoulos, J., Fieberg, J., Aarts, G., Beyer, H. L., Morales, J. M., & Haydon, D. T. (2015). Establishing the link between habitat selection and animal population dynamics. *Ecological Monographs*, 85(3), 413–436.

McClintock, B. T., Johnson, D. S., Hooten, M. B., Ver Hoef, J. M., & Morales, J. M. (2014). When to be discrete: the importance of time formulation in understanding animal movement. *Movement Ecology*, 2, 1–14.

McClintock, B. T., Langrock, R., Gimenez, O., Cam, E., Borchers, D. L., Glennie, R., & Patterson, T. A. (2020). Uncovering ecological state dynamics with hidden Markov models. *Ecology Letters*, 23(12), 1878–1903.

- McClintock, B. T., & Michelot, T. (2018). momentuHMM: R package for generalized hidden Markov models of animal movement. *Methods in Ecology and Evolution*, 9(6), 1518–1530. <https://doi.org/10.1111/2041-210X.12995>
- McClure, K. M., Bastille-Rousseau, G., Davis, A. J., Stengel, C. A., Nelson, K. M., Chipman, R. B., Wittemyer, G., Abdo, Z., Gilbert, A. T., & Pepin, K. M. (2022). Accounting for animal movement improves vaccination strategies against wildlife disease in heterogeneous landscapes. *Ecological Applications*, 32(4), e2568.
- McLoughlin, P. D., Morris, D. W., Fortin, D., Vander Wal, E., & Contasti, A. L. (2010). Considering ecological dynamics in resource selection functions. *Journal of Animal Ecology*, 79(1), 4–12.
- Mercker, M., Schwemmer, P., Peschko, V., Enners, L., & Garthe, S. (2021). Analysis of local habitat selection and large-scale attraction/avoidance based on animal tracking data: is there a single best method? *Movement Ecology*, 9(1), 20.
- Michelot, T., Klappstein, N. J., Potts, J. R., & Fieberg, J. (2023). Understanding step selection analysis through numerical integration. *ArXiv Preprint ArXiv:2308.15678*.
- Michelot, T., Langrock, R., & Patterson, T. A. (2016). moveHMM: an R package for the statistical modelling of animal movement data using hidden Markov models. *Methods in Ecology and Evolution*, 7(11), 1308–1315. <https://doi.org/10.1111/2041-210X.12578>
- Morales, J. M., Haydon, D. T., Frair, J., Holsinger, K. E., & Fryxell, J. M. (2004). Extracting more out of relocation data: building movement models as mixtures of random walks. *Ecology*, 85(9), 2436–2445.
- Morales, J. M., Moorcroft, P. R., Matthiopoulos, J., Frair, J. L., Kie, J. G., Powell, R. A., Merrill, E. H., & Haydon, D. T. (2010). Building the bridge between animal movement and

- population dynamics. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1550), 2289–2301.
- Morris, D. W. (2003). Toward an ecological synthesis: a case for habitat selection. *Oecologia*, 136, 1–13.
- Muff, S., Signer, J., & Fieberg, J. (2020). Accounting for individual-specific variation in habitat-selection studies: Efficient estimation of mixed-effects models using Bayesian or frequentist computation. *Journal of Animal Ecology*, 89(1), 80–92.
- Nams, V. O. (2013). Sampling animal movement paths causes turn autocorrelation. *Acta Biotheoretica*, 61, 269–284.
- Nathan, R., Monk, C. T., Arlinghaus, R., Adam, T., Alós, J., Assaf, M., Baktoft, H., Beardsworth, C. E., Bertram, M. G., & Bijleveld, A. I. (2022). Big-data approaches lead to an increased understanding of the ecology of animal movement. *Science*, 375(6582), eabg1780.
- Ngô, M. C., Heide-Jørgensen, M. P., & Ditlevsen, S. (2019). Understanding narwhal diving behaviour using Hidden Markov Models with dependent state distributions and long range dependence. *PLOS Computational Biology*, 15(3), e1006425.
<https://doi.org/10.1371/journal.pcbi.1006425>
- Nicosia, A., Duchesne, T., Rivest, L.-P., & Fortin, D. (2017). *A multi-state conditional logistic regression model for the analysis of animal movement*.
- Nielsen, S. E., Boyce, M. S., Stenhouse, G. B., & Munro, R. H. M. (2002). Modeling grizzly bear habitats in the Yellowhead ecosystem of Alberta: taking autocorrelation seriously. *Ursus*, 45–56.

- Northrup, J. M., Hooten, M. B., Anderson Jr, C. R., & Wittemyer, G. (2013). Practical guidance on characterizing availability in resource selection functions under a use–availability design. *Ecology*, *94*(7), 1456–1463.
- Northrup, J. M., Vander Wal, E., Bonar, M., Fieberg, J., Laforge, M. P., Leclerc, M., Prokopenko, C. M., & Gerber, B. D. (2022). Conceptual and methodological advances in habitat-selection modeling: guidelines for ecology and evolution. *Ecological Applications*, *32*(1), e02470.
- Owen, M. (1972). Some factors affecting food intake and selection in white-fronted geese. *The Journal of Animal Ecology*, 79–92.
- Patterson, T. A., McConnell, B. J., Fedak, M. A., Bravington, M. V., & Hindell, M. A. (2010). Using GPS data to evaluate the accuracy of state–space methods for correction of Argos satellite telemetry error. *Ecology*, *91*(1), 273–285.
- Prima, M.-C., Duchesne, T., Merkle, J. A., Chamaillé-Jammes, S., & Fortin, D. (2022). Multi-mode movement decisions across widely ranging behavioral processes. *Plos One*, *17*(8), e0272538.
- Pyke, G. H., Pulliam, H. R., & Charnov, E. L. (1977). Optimal foraging: a selective review of theory and tests. *The Quarterly Review of Biology*, *52*(2), 137–154.
- Rhodes, J. R., McAlpine, C. A., Lunney, D., & Possingham, H. P. (2005). A spatially explicit habitat selection model incorporating home range behavior. *Ecology*, *86*(5), 1199–1205.
- Roeber, C. L., Boyce, M. S., & Stenhouse, G. B. (2010). Grizzly bear movements relative to roads: application of step selection functions. *Ecography*, *33*(6), 1113–1122.
- Sells, S. N., Costello, C. M., Lukacs, P. M., Roberts, L. L., & Vinks, M. A. (2022). Grizzly bear habitat selection across the Northern Continental Divide Ecosystem. *Biological Conservation*, *276*, 109813.

- Shuert, C. R., Marcoux, M., Hussey, N. E., Heide-Jørgensen, M. P., Dietz, R., & Auger-Méthé, M. (2022). Decadal migration phenology of a long-lived Arctic icon keeps pace with climate change. *Proceedings of the National Academy of Sciences*, *119*(45), e2121092119.
- Sidrow, E., Heckman, N., Fortune, S. M. E., Trites, A. W., Murphy, I., & Auger-Méthé, M. (2022). Modelling multi-scale, state-switching functional data with hidden Markov models. *Canadian Journal of Statistics*, *50*(1), 327–356.
- Signer, J., Fieberg, J., & Avgar, T. (2017). Estimating utilization distributions from fitted step-selection functions. *Ecosphere*, *8*(4), e01771.
- Signer, J., Fieberg, J., & Avgar, T. (2019). Animal movement tools (amt): R package for managing tracking data and conducting habitat selection analyses. *Ecology and Evolution*, *9*(2), 880–890.
- Squires, J. R., DeCesare, N. J., Olson, L. E., Kolbe, J. A., Hebblewhite, M., & Parks, S. A. (2013). Combining resource selection and movement behavior to predict corridors for Canada lynx at their southern range periphery. *Biological Conservation*, *157*, 187–195.
- Thurfjell, H., Ciuti, S., & Boyce, M. S. (2014). Applications of step-selection functions in ecology and conservation. *Movement Ecology*, *2*, 1–12.
- Togunov, R. R., Derocher, A. E., Lunn, N. J., & Auger-Méthé, M. (2022). Drivers of polar bear behavior and the possible effects of prey availability on foraging strategy. *Movement Ecology*, *10*(1), 50.
- Valeix, M., Fritz, H., Sabatier, R., Murindagomo, F., Cumming, D., & Duncan, P. (2011). Elephant-induced structural changes in the vegetation and habitat selection by large herbivores in an African savanna. *Biological Conservation*, *144*(2), 902–912.

- Warton, D. I., Blanchet, F. G., O'Hara, R. B., Ovaskainen, O., Taskinen, S., Walker, S. C., & Hui, F. K. C. (2015). So many variables: joint modeling in community ecology. *Trends in Ecology & Evolution*, *30*(12), 766–779.
- Warton, D. I., & Shepherd, L. C. (2010). Poisson point process models solve the "pseudo-absence problem" for presence-only data in ecology. *The Annals of Applied Statistics*, 1383–1402.
- Webber, Q. M. R., Prokopenko, C. M., Kingdon, K. A., Turner, J. W., & Vander Wal, E. (2021). Effects of the social environment on movement-integrated habitat selection. *BioRxiv*, 2002–2021.
- Wiens, J. A. (1984). *Resource systems, populations and communities*. In: *A new Ecology: Novel Approaches to Interactive Systems* (P. Price, C. Slobodchikoff, W. Gaud, & Eds, Eds.). John Wiley and Sons, New York.
- Wijeyakulasuriya, D. A., Eisenhauer, E. W., Shaby, B. A., & Hanks, E. M. (2020). Machine learning for modeling animal movement. *Plos One*, *15*(7), e0235750.
- Zucchini, W., & MacDonald, I. L. (2009). *Hidden Markov models for time series: an introduction using R*. Chapman and Hall/CRC.

Appendix 1: Supplemental Figures and Tables




Figure S1. A) Step length and B) turning angle distributions of predicted states from the case study's main HMM (i.e., the HMM featured in the main paper, where the covariate, prey diversity, only affects the transition probability matrix and not the observation probabilities, called `hmm_trans_3` in the tutorial).




Figure S2. Simulated tracks ($n = 10,000$ locations) from the redistribution kernel that was generated from the step-selection functions where prey diversity A) without and B) with movement-related covariates. These simulated paths were used to generate a steady-state utilization distribution (SSUD) for each model. Note that we present here $n = 10,000$ locations (the sample size used in the tutorial) for visualization purposes. The SSUDs generated in Fig. 6 used $n = 100,000$ locations, which contained so many locations that they became indistinguishable from one another.

Table S1. Results from the case study RSF and SSFs regression coefficients.

Model	Parameter	β	SE	z-value	p-value
RSF					
	Intercept	-6.118	1.400	-4.369	<0.001
	Prey diversity	6.353	2.748	2.748	0.006
SSF without covariate affecting movement kernel (SSF1)					
	Prey diversity	0.948	6.837	0.139	0.890
	Step length	<0.001	<0.001	-0.383	0.701
	Log (step length)	-0.042	0.157	0.269	0.788
	Cos (turning angle)	0.031	0.131	0.235	0.814
SSF with covariate affecting movement kernel (SSF2)					
	Prey diversity	31.500	23.090	1.364	0.173
	Step length	<0.001	<0.001	-0.645	0.519
	Log (step length)	1.981	7.250	1.395	0.163
	Cos (turning angle)	0.021	1.022	0.163	0.871
	Prey diversity: log (step length)	-3.123	2.253	-1.386	0.166

β = parameter estimate, SE = standard error.

Table S2. Parameter estimates associated with the transition probability matrix of the case study’s main HMM (i.e., the HMM featured in the main paper, where the covariate, prey diversity, only affects the transition probability matrix, called `hmm_trans_3` in the tutorial) regression coefficients for the transition probabilities, presented as “estimate (standard error)”, on the beta (i.e., working) scale of the parameters. Note that the standard error for the transition from state 1 to 3 is NA since this transition does not occur in our short example track; using a longer track would likely allow the model to estimate a standard error.

	Estimate for transitioning from state_t -> state_{t+1}					
Parameter	1 -> 2	1 -> 3	2 -> 1	2 -> 3	3 -> 1	3 -> 2
Intercept	11.958 (6.491)	-28.986 (<0.001)	0.889 (6.070)	-6.008 (7.694)	52.188 (0.892)	-12.384 (9.787)
Prey diversity	-22.386 (10.919)	0.000 NA	-3.842 (9.974)	6.299 (12.461)	-97.113 (0.562)	19.088 (15.704)

State 1 = slow movement, state 2 = moderate movement, state 3 = fast movement.

Table S3. Parameter estimates associated with the observation probability matrix of the case study’s main HMM (i.e., the HMM featured in the main paper, where the covariate, prey diversity, only affects the transition probability matrix, called `hmm_trans_3` in the tutorial) including the regression coefficients for step length (mean and standard deviation [SD]) and turning angle (concentration), presented as “estimate (standard error)”, on the beta (i.e., working) scale of the parameters.

State:Parameter	Step length		Turning angle
	Mean	SD	Concentration
Slow movement: Intercept	1.544 (0.141)	1.144 (0.203)	-1.133 (0.579)
Moderate movement: Intercept	2.669 (0.056)	1.581 (0.132)	-0.351 (0.321)
Fast movement: Intercept	3.684 (0.030)	1.424 (0.203)	0.382 (0.379)

Table S4. Parameter estimates associated with the transition probability matrix from the tutorial’s example of a HMM where the covariate, prey diversity, only affects the observation probability (i.e., not the transition probability matrix, called `hmm_obs_3` in the tutorial), including regression coefficients for the transition probabilities, presented as “estimate (standard error)”, on the beta (i.e., working) scale of the parameters.

	Estimate for transitioning from state_t -> state_{t+1}					
Parameter	1 -> 2	1 -> 3	2 -> 1	2 -> 3	3 -> 1	3 -> 2
Intercept	-1.709 (0.474)	-18.576 (0.119)	-1.692 (0.449)	-2.287 (0.506)	-16.157 (0.059)	-0.441 (0.586)

State 1 = slow movement, state 2 = moderate movement, state 3 = fast movement.

Table S5. Parameter estimates associated with the observation probability matrix from the tutorial’s example of a HMM where the covariate, prey diversity, only affects the observation probability (i.e., not the transition probability matrix, called `hmm_obs_3` in the tutorial), including the regression coefficients for step length (mean and standard deviation [SD]) and turning angle (concentration), presented as “estimate (standard error)”, on the beta (i.e., working) scale of the parameters.

State:Parameter	Step length		Turning angle
	Mean	SD	Concentration
Slow movement: Intercept	2.690 (1.318)	1.082 (0.202)	8.038 (7.387)
Slow movement: Prey diversity	-1.957 (2.125)	Not modeled	-15.016 (12.647)
Moderate movement: Intercept	1.116 (0.674)	1.611 (0.119)	-3.286 (4.846)
Moderate movement: Prey diversity	2.495 (1.092)	Not modeled	4.794 (7.899)
Fast movement: Intercept	3.816 (0.411)	1.455 (0.241)	3.344 (4.104)
Fast movement: Prey diversity	-0.229 (0.678)	Not modeled	-4.922 (6.975)

Appendix 2: Full methods for case study on ringed seals in Hudson Bay

The data and code tutorial of analyses used for this case study are available as Appendix 3 and 4, respectively.

1. Data collection and preparation

The seal, a two-year-old male (tag ID: 116484), was captured on 29 Oct 2012 using a monofilament mesh net set perpendicular to the shore along the coast of the Belcher Islands, Nunavut. We equipped the seal with a 9000x data logger from the Sea Mammal Research Unit (SMRU, University of St. Andrews, UK). See Florko et al., 2023 for details. We used one segment of the tracking data for the purpose of this case study, which had consistent transmissions over the course of 4.5 months. We used a correlated random walk state-space model (`animotum` R package, Jonsen et al., 2023) to filter and regularize the location data at a 24-hr time step since ARGOS location data is observed irregularly in time and is prone to error (Costa et al., 2010). This procedure resulted in one location per day from 29 Oct 2012 to 17 Mar 2013 ($n = 140$).

We used estimated prey diversity using the estimated abundance of eight important ringed seal prey species: Arctic cod (*Boreogadus saida*), capelin (*Mallotus villosus*), northern sand lance (*Ammodytes dubius*), Pacific sand lance (*Ammodytes hexaoterus*), Arctic staghorn sculpin (*Gymnocanthus tricuspis*), shorthorn sculpin (*Myoxocephalus Scorpius*), mustache sculpin (*Triglops murrayi*), and rainbow smelt (*Osmerus mordax*) using Simpson's Diversity Index (Florko et al., 2021a; 2021b, Fig 4). The fish data were modeled at a yearly time step on a 0.5° longitude by 0.5° degree latitude grid, under both a low emission (representative concentration pathway, RCP 2.6) and a high emission (RCP 8.5) emission scenario. We extracted the fish

abundance data for each location in the corresponding grid cell. We used the high emission scenario as it aligned more closely to the emissions in 2012.

2. Statistical analysis

2.1 Resource selection function

We used the `amt` package in R (Signer et al., 2019) to prepare the data and fit the resource selection function (RSF) and step selection functions (SSFs). We used the `random_points()` function to generate random locations (i.e., availability sample) within the minimum convex polygon (MCP, clipped to exclude land, note a buffer around the MCP can also be added), which by default generated 10 random locations per observed location. We used the `extract_covariates()` function to extract the corresponding prey diversity estimates for each observed and available location. We fit the RSF using the `fit_rsf()` function (a wrapper for `fit_logit()`), including the covariate for prey diversity (*preydiv*). `fit_rsf()` estimates the RSF coefficients by fitting a logistic regression to the use-available observations $Y = (y_1, \dots, y_T)$, where $y_t \in \{0,1\}$. This is modeled using a Bernoulli distribution as:

Equation S 1

$$y_t \sim \text{Bernoulli}(P(Y = 1 | x_{preydiv,t})),$$

where we use a logit link function to keep the probability $P(Y = 1 | x_{preydiv,t})$ between 0 and 1:

Equation S 2

$$P(Y = 1 | x_{preydiv,t}) = \frac{\exp(\beta_0 + \beta_{preydiv} x_{preydiv,t})}{1 + \exp(\beta_0 + \beta_{preydiv} x_{preydiv,t})}.$$

By default, R fits logistic models with an intercept, however, in the context of an RSF based on use-available data, this intercept is not interpretable. Therefore, the RSF derived from this model excludes the intercept:

Equation S 3

$$w(x_{preydiv}) = \beta_{preydiv} x_{preydiv} .$$

We calculated the log of the relative selection strength (log-RSS) for prey diversity from our fitted model (Avgar et al., 2017; Fieberg et al., 2021), which is calculated as the relative frequency distribution based on location 1 (s_1) relative to location 2 (s_2):

Equation S 4

$$\frac{F^U(x_{preydiv}(s_1))}{F^U(x_{preydiv}(s_2))} = \frac{\exp(\beta_{preydiv} x_{preydiv}(s_1))}{\exp(\beta_{preydiv} x_{preydiv}(s_2))} .$$

Further, we predicted the relative probability of use in space given our fitted model by applying equation S2 (without the intercept term, β_0) to each cell in our study area, and normalizing the prediction following:

Equation S 5

$$\frac{w(x_{preydiv}) - \min(w(x_{preydiv}))}{\max(w(x_{preydiv})) - \min(w(x_{preydiv}))} .$$

2.2 Step selection function

We used the `steps()` function to turn the dataset into “steps” (i.e., with a start and end location) rather than just observations. Next, we used the `random_steps()` function to generate 10 random steps per each observed step. The `random_steps()` function also generated a “step ID” column which represents each observed step and their available (random) steps ($n = 140$ observed steps). Similar to the RSF, we used the `extract_covariates()` function to extract the prey diversity estimate for each observed and available step. We fit the SSF using the `fit_clogit()` function, including the covariate for prey diversity, the log of the step length, the cosine of the turning angle, as well as a `strata()` for step ID.

Using `fit_clogit()` to estimate SSF parameters requires that distributions of the movement kernel (i.e., step length and turn angle distributions) are of the exponential family, such that the product of movement kernel and habitat selection kernel can be defined in linear form. We used the gamma distribution for step length and von Mises distribution for turn angle, which defines the linear combination of SSF, $\eta_{t+\Delta t}$, predictors as follows (see Avgar et al., 2016 for full derivation):

Equation S 6

$$\begin{aligned}\eta_{t+\Delta t} &= w(x_{preydiv}(s_{t+\Delta t}))\phi(s_{t+\Delta t}, s_t) \\ &= \beta_{0,stepID} + \beta_{preydiv}x_{preydiv}(s_{t+\Delta t}) + \beta_l l(s_{t+\Delta t}, s_t) + \beta_{ln} \ln(l(s_{t+\Delta t}, s_t)) + \\ &\beta_{\theta} \cos(\theta(s_{t+\Delta t}, s_t)).\end{aligned}$$

As a computationally efficient method to approximate the integral in Equation 5 in the main text, we used a conditional logistic regression, where each observed step $S_{t+\Delta t, observed}$ was coupled with $n = 10$ available (random) sample steps, together, defining 11 steps $S_{t+\Delta t} \in \{S_{t+\Delta t,0}, \dots, S_{t+\Delta t,n}\}$ at each time step. In the case of fitting the SSFs using `amt`, we consider each step, t , as a stratum (coded as `strata(step_ID)`) the conditional probability of the observed step given available steps at each stratum (Avgar et al., 2016; Michelot et al., 2023), which is defined as:

Equation S 7

$$P(S_{t+\Delta t, observed} = 1 | S_{t+\Delta t}) = \frac{\exp(\eta_{t+\Delta t, observed})}{\sum_{i=0}^n \exp(\eta_{t+\Delta t, i})}.$$

We also fit a SSF where we allowed prey diversity to affect the movement (specifically, the shape of the gamma distribution); thus, this model was fit similarly but included an interaction between prey diversity and the natural log of the step length:

Equation S 8

$$\eta_{t+\Delta t} = \beta_{0, stepID} + \beta_{preydiv} x_{preydiv}(s_{t+\Delta t}) + \beta_l l(s_{t+\Delta t}, s_t) + \beta_{ln} \ln(l(s_{t+\Delta t}, s_t)) + \beta_{\theta} \cos(\theta(s_{t+\Delta t}, s_t)) + \beta_{preydiv:ln} x_{preydiv}(s_{t+\Delta t}) \ln(l(s_{t+\Delta t}, s_t)).$$

Note that the likelihoods between a conditional logistic regression and SSF are equivalent when the movement kernel and habitat selection function are modeled using a distribution from the exponential family. As such, when using a conditional logistic regression to fit a SSF, we are therefore limited to using distributions such as the exponential, gamma distributions, or half normal for step length, and von Mises distribution for turning angles (Avgar et al., 2016; Michelot

et al., 2023). However, using a general model (not conditional logistic regression) to consider other distributions (e.g., Weibull for step length and/or wrapped Cauchy for turning angle) may improve model fit (Michelot et al., 2023).

Similar to the RSF, we calculated the log-RSS for prey diversity from our fitted models (Avgar et al., 2017; Fieberg et al., 2021). For spatial inference, we estimated the utilization distribution (UD) from each model. The UD is defined as the two-dimensional relative frequency distribution of space use of an animal (Van Winkle, 1975). We calculated the steady-state UD, which is defined as the long-term expectation of the space-use distribution across the landscape (Signer et al., 2017). To do this, we simulated an animal movement track with 10,000 locations that started at the first observed location (Fig. S3), and we removed the first 200 locations (2%) as a burn-in to reduce bias associated with setting the starting point. We obtained the steady-state UD using the `hr_ud()` function in `amt` which counts the number of times each cell was visited by the simulated animal, and normalized these frequencies to sum to 1 over the study area (Signer et al., 2017).

2.3 Hidden Markov model

We used the `extract_covariates()` function from the `amt` package to extract the fish biomass estimates for the observed data. We used the `momentuHMM` package in R (McClintock & Michelot, 2018) for the rest of the data preparation and to fit the hidden Markov model (HMM). We used the `prepData()` function to convert the data into “steps”, and to compute step lengths and turning angles for each step. We assumed the step lengths followed a gamma distribution and

the turning angles followed a von Mises distribution. We initially fit a two-state HMM with prey diversity as a covariate in the transition probability (called `hmm_trans_2` in the tutorial), using the `fitHMM()` function, and the `getPar0()` function to compute starting parameter values for each state, including the mean and standard deviation of step length and the turning angle concentration. However, inspection of the histograms of step length and turning angle and pseudo-residual plots suggested an additional state was not being captured by the model (see Appendix 4). Therefore, we fit a three-state HMM, which improved the coverage in the histograms. We refit the three-state HMM using these new starting parameters (from `getPar0()`). The final model which was used in the main paper, a three-state HMM with prey diversity as a covariate on the transition probability (called `hmm_trans_3` in the tutorial), was:

Equation S 9

$$\gamma_{t,i,j} = \frac{\exp(\beta_{0,i,j} + \beta_{i,j} x_{preydiv,t})}{\sum_{i=1}^N \exp(\beta_{0,i,l} + \beta_{i,l} x_{preydiv,t})},$$

where $x_{preydiv,t}$ is the prey diversity at time t , N is the number of behavioural states (set to $N = 3$ in the final model), and $\beta_{0,i,j}$ and $\beta_{i,j}$ are the intercept and slope coefficient effect of prey diversity on transition probability from state i to j , respectively, which are fixed to $\beta_{0,i,j} = \beta_{k,i,j} = 0$ when $i = j$ (i.e., the diagonal of the transition probability matrix Γ).

We also show in the tutorial how covariates can alternatively be included to modify the observation probabilities, that is, a three-state model including the covariate effects on the step length mean and turning angle concentration (Table S3 and S4, called `hmm_obs_3` in the tutorial). This is typically used for factors that are unlikely to affect the probability of being in a

behaviour (e.g., abiotic factors that would affect movement capacity), however, to demonstrate how this is done in our tutorial, we allowed prey diversity to affect the step length mean $\mu^{(l)}$ and turning angle concentration $\kappa^{(\theta)}$ for each state i :

Equation S 10

$$\ln(\mu_{t,i}^{(l)}) = \beta_{i,0}^{(\mu^{(l)})} + \beta_{i,preydiv}^{(\mu^{(l)})} x_{preydiv,t} \ ,$$

Equation S 11

$$\ln(\kappa_{t,i}^{(\theta)}) = \beta_{i,0}^{(\kappa^{(\theta)})} + \beta_{i,preydiv}^{(\kappa^{(\theta)})} x_{preydiv,t} \ .$$

All behaviours were assumed to be correlated random walks with mean turn angle $\mu_i^{(\theta)}$ fixed to 0. Finally, we show how they can also be included in both transition and observation probabilities simultaneously (called `hmm_obstrans_2` and `hmm_obstrans_3` in the tutorial), although this model is complex and perhaps not well fitted in cases where the state-dependent distributions have a lot of overlap.

We extracted the stationary state probabilities using the `plotStationary()` function, and we used the `viterbi()` function to decode the hidden states.

References

Avgar, T., Lele, S. R., Keim, J. L., & Boyce, M. S. (2017). Relative selection strength: Quantifying effect size in habitat-and step-selection inference. *Ecology and Evolution*, 7(14), 5322–5330.

- Avgar, T., Potts, J. R., Lewis, M. A., & Boyce, M. S. (2016). Integrated step selection analysis: bridging the gap between resource selection and animal movement. *Methods in Ecology and Evolution*, 7(5), 619–630.
- Costa, D. P., Robinson, P. W., Arnould, J. P. Y., Harrison, A., Samantha, E., Hassrick, J. L., Hoskins, A. J., Kirkman, S. P., Oosthuizen, H., & Crocker, D. E. (2010). Accuracy of ARGOS locations of pinnipeds at-sea estimated using Fastloc GPS. *PLoS ONE*, 5(1).
<https://doi.org/10.1371/journal.pone.0008677>
- Fieberg, J., Signer, J., Smith, B., & Avgar, T. (2021). A ‘How to’ guide for interpreting parameters in habitat-selection analyses. *Journal of Animal Ecology*, 90(5), 1027–1043.
- Florko, K. R. N., Shuert, C. R., Cheung, W. W. L., Ferguson, S. H., Jonsen, I. D., Rosen, D. A. S., Sumaila, U. R., Tai, T. C., Yurkowski, D. J., & Auger-Méthé, M. (2023). Linking movement and dive data to prey distribution models: new insights in foraging behaviour and potential pitfalls of movement analyses. *Movement Ecology*, 11(1), 17.
- Florko, K. R. N., Tai, T. C., Cheung, W. W. L., Ferguson, S. H., Sumaila, U. R., Yurkowski, D. J., & Auger-Méthé, M. (2021a). Predicting how climate change threatens the prey base of Arctic marine predators. *Ecology Letters*, 24, 2563–2575. <https://doi.org/10.1111/ele.13866>
- Florko, K. R. N., Tai, T. C., Cheung, W. W. L., Sumaila, U. R., Ferguson, S. H., Yurkowski, D. J., & Auger-Méthé, M. (2021b). Predicting how climate change threatens the prey base of Arctic marine predators. *Dryad, dataset*. <https://doi.org/https://doi.org/10.5061/dryad.x69p8czjs>
- Jonsen, I. D., Grecian, W. J., Phillips, L., Carroll, G., McMahon, C., Harcourt, R. G., Hindell, M. A., & Patterson, T. A. (2023). aniMotum, an R package for animal movement data: Rapid quality control, behavioural estimation and simulation. *Methods in Ecology and Evolution*, 14(3), 806–816.

- McClintock, B. T., & Michelot, T. (2018). momentuHMM: R package for generalized hidden Markov models of animal movement. *Methods in Ecology and Evolution*, 9(6), 1518–1530. <https://doi.org/10.1111/2041-210X.12995>
- Michelot, T., Klappstein, N. J., Potts, J. R., & Fieberg, J. (2023). Understanding step selection analysis through numerical integration. *ArXiv Preprint ArXiv:2308.15678*.
- Signer, J., Fieberg, J., & Avgar, T. (2017). Estimating utilization distributions from fitted step-selection functions. *Ecosphere*, 8(4), e01771.
- Signer, J., Fieberg, J., & Avgar, T. (2019). Animal movement tools (amt): R package for managing tracking data and conducting habitat selection analyses. *Ecology and Evolution*, 9(2), 880–890.
- Van Winkle, W. (1975). Comparison of several probabilistic home-range models. *The Journal of Wildlife Management*, 118–123.

Appendix 3: Data for case study on ringed seals in Hudson Bay

[\[github link\]](#)

Appendix 4: Code tutorial for case study on ringed seals in Hudson Bay

[\[github link\]](#)