

On generalized covering and avoidance properties of finite groups and saturated fusion systems*

Shengmin Zhang, Zhencai Shen

College of Science, China Agricultural University, Beijing 100083, China

Abstract

A subgroup A of a finite group G is said to be a CAP -subgroup of G , if for any chief factor H/K of G , either $AH = AK$ or $A \cap H = A \cap K$. Let p be a prime, S be a p -group and \mathcal{F} be a saturated fusion system over S . Then \mathcal{F} is said to be supersolvable, if there exists a series of S , namely $1 = S_0 \leq S_1 \leq \cdots \leq S_n = S$, such that S_{i+1}/S_i is cyclic, and S_i is strongly \mathcal{F} -closed for any $i = 0, 1, \dots, n$. In this paper, we first introduce the concept of strong p - CAP -subgroups, and investigate the properties of finite groups under the assumption that some subgroups of G are partial CAP -subgroups or strong (p) - CAP -subgroups, and obtain some criteria for a group G to be p -supersolvable. Then, we investigate the characterizations for supersolvability of $\mathcal{F}_S(G)$ under the assumption that some subgroups of G are partial CAP -subgroups or strong (p) - CAP -subgroups of G , and obtain some criteria for a fusion system $\mathcal{F}_S(G)$ to be p -supersolvable.

Keywords: Strong p - CAP -subgroups, Partial CAP -subgroups, Saturated formation, p -supersolvable subgroups, Saturated fusion systems, Supersolvable fusion systems.

MSC:20D10,20D15, 20D20

1 Introduction

All group considered in this paper will be finite. We say a subgroup A of a group G satisfies the covering and avoidance property of G , if for any chief factor H/K of G , either $HA = KA$, or $H \cap A = K \cap A$. Such group A is called a CAP -subgroup of G . The covering and avoidance property was first introduced by W. Gaschütz in [10], and there have been a lot of generalizations of covering and avoidance property. These properties and characterizations of CAP -subgroups have been widely researched since it plays an important part in the structural study of soluble or supersolvable groups. In recent years, some authors give the structure of CAP -subgroups ([13]), some give the equivalent conditions ([15]), while others give some generalizations for the original definition as we have said before. In [12], partial CAP -subgroup was introduced as a weaker generalization, by changing any chief factor in to a chief series compared to CAP -subgroup. In [15], a generalized definition called CAP^* -subgroup is introduced to loose the condition into the

*E-mail addresses: zsmcau@cau.edu.cn (S. Zhang); zhencai688@sina.com (Z. Shen).

non-Frattini chief factors. Some equivalent criteria related to Hall groups, Sylow subgroups and maximal subgroups were given in this paper. In [4], stronger generalization named strong *CAP*-subgroup emerges. A subgroup A of a group G is called a strong *CAP*-subgroup of G , if A is a *CAP*-subgroup for any $H \geq A$. The reason to generalize the *CAP*-property in this way is that *CAP*-property does not satisfy the intermediate property ([7, A, Definition 10.8]). One can find that partial *CAP*-subgroup satisfies the intermediate property as well [5]. However, four subgroups mentioned above enjoy the same properties about the inheritance of quotient group as mentioned in [12, Lemma 2.6], [15, Lemma 2.3(1)], [4, Lemma 3(2)].

It is easy to find that only partial *CAP*-subgroup satisfies both intermediate property and inheritance property. Our aim is to find a new category of generalization of *CAP*-subgroup which still satisfies the two properties. As introduced in [9], we have the following definition:

Definition 1.1. A subgroup A of a group G is called a *p-CAP*-subgroup of G if A covers or avoids every *pd*-chief factor of G , where *pd*-chief factor denotes the chief factor of G with order divided by p .

It follows from [9, Lemma 2.1] that *p-CAP*-subgroup satisfies the inheritance of quotient group. However, no evidence shows that *p-CAP*-subgroup satisfies the intermediate property. Therefore, we can generalize the definition by strengthening our assumption into the following one.

Definition 1.2. A subgroup A of a group G is called a strong *p-CAP*-subgroup of G , if A is a *p-CAP*-subgroup for any $H \geq A$.

Let G be a finite group. Then for any prime divisor p of G , it is obvious that G and 1 are two trivial strong *p-CAP*-subgroup of G . Actually, non-trivial strong *p-CAP*-subgroups have frequent occurrence in the structure of finite groups.

Example 1.3. The alternating group on 4 letters A_4 has a maximal subgroup C_3 . Then clearly C_3 is a 3-*CAP*-subgroup of C_3 itself. On the other hand, there is only one 3*d*-chief factor A_4/C_2^2 , and we conclude immediately that $A_4 \cdot C_3 = A_4 = C_2^2 \cdot C_3$. Hence C_3 is a 3-*CAP*-subgroup of A_4 , and so C_3 is a strong *p-CAP*-subgroup of A_4 .

Now we would like to give an example to show that *p-CAP*-subgroups are not strong *p-CAP*-subgroups in general, which differs the concept of strong *p-CAP*-subgroups from *p-CAP*-subgroups.

Example 1.4. Consider the group $G := \text{SL}_2(\mathbb{F}_5)$. Then there exists a non-normal subgroup $\text{SL}_2(\mathbb{F}_3)$ of $\text{SL}_2(\mathbb{F}_5)$, and a subgroup C_4 of $\text{SL}_2(\mathbb{F}_3)$. Since there is only one non-trivial normal subgroup C_2 of G , and so C_2/C_1 is the only 2*d*-chief factor of G . As C_4 contains C_2 , it follows that $C_4 \cdot C_1 = C_4 = C_2 \cdot C_4$, and so C_4 covers C_2/C_1 , which implies that C_4 is a 2-*CAP*-subgroup of G . On the other hand, consider the subgroup C_4 again in the group $\text{SL}_2(\mathbb{F}_3)$. Then one can find that Q_8/C_2 is a 2*d*-chief factor of $\text{SL}_2(\mathbb{F}_3)$. However, since $C_2 \leq C_4 \leq Q_8$, it yields that $Q_8 \cdot C_4 = Q_8 \neq C_4 = C_4 \cdot C_2$ and $C_4 \cap Q_8 = C_4 \neq C_2 = C_4 \cap C_2$, and so C_4 is not a 2-*CAP*-subgroup of $\text{SL}_2(\mathbb{F}_3)$. Therefore C_4 is a 2-*CAP*-subgroup of G , but not a strong 2-*CAP*-subgroup of G .

In the first part of our paper (Section 3), we will give three characterizations for a group G to be p -supersolvable or in a saturated formation containing \mathfrak{U} under the assumption that some subgroups of G are strong p -CAP-subgroups or partial CAP-subgroups. In particular, Theorem 3.3 gives a new criteria for p -supersolvability under the assumption that all maximal subgroups of $S \in \text{Syl}_p(G)$ are partial CAP-subgroups, which is first introduced in our paper. This criteria also plays an important role in the research of characterizations for $\mathcal{F}_S(G)$, which will be mentioned in the next part of the paper.

Now, we are going to give a brief introduction to the basic theory of fusion systems, and begin to introduce our results on the characterization of the structure of $\mathcal{F}_S(G)$. Let S be a Sylow p -subgroup of G , where p is a prime divisor of $|G|$. Then the fusion system of G over S is defined in [6, Definition 1.34]. One can easily find that $\mathcal{F}_S(G)$ is exactly a saturated fusion system over S by [1, Theorem 2.3]. As is known to all, the structure of $\mathcal{F}_S(G)$ has a strong relationship with the structure of G . Hence some structures of finite groups can be generalized into the fusion system $\mathcal{F}_S(G)$. Recall that G is said to be supersolvable, if there exists a chief series, namely

$$1 = N_0 \leq N_1 \leq \cdots \leq N_t = G,$$

such that N_{i+1}/N_i is cyclic, $i = 0, 1, \dots, t-1$. As a natural way of generalization, we want to define a similar structure in \mathcal{F} , where \mathcal{F} is a fusion system over a p -group S . Since the objects of \mathcal{F} are exactly the subgroups of S , we may restrict the chief series of G into a series of S . Note that the normality of subgroups H of S in G represents the invariance of H under the morphisms induced by conjugation of G , we may change the normality of H in G into invariance of H under the morphisms in \mathcal{F} . Then one can easily find that the invariance of H under the morphisms in \mathcal{F} exactly suits the concept of weakly \mathcal{F} -closed property, hence we give the following definition which was introduced by N. Su in [17].

Definition 1.5. Let \mathcal{F} be a saturated fusion system over a p -group S . Then \mathcal{F} is called supersolvable, if there exists a series of subgroups of S , namely:

$$1 = S_0 \leq S_1 \leq \cdots \leq S_n = S,$$

such that S_i is strongly \mathcal{F} -closed, $i = 0, 1, \dots, n$, and S_{i+1}/S_i is cyclic for any $i = 0, 1, \dots, n-1$.

Now we would like to introduce some concepts which are useful for us to discover the structure of $\mathcal{F}_S(G)$. Let S be a p -group and P be a subgroup of S . Suppose that \mathcal{F} is a fusion system over S . Then P is called \mathcal{F} -centric, if $C_S(Q) = Z(Q)$ for all $Q \in P^{\mathcal{F}}$, where $P^{\mathcal{F}}$ denotes the set of all subgroups of S which are \mathcal{F} -conjugate to P . P is said to be fully normalized in \mathcal{F} , if $|N_S(P)| \geq |N_S(Q)|$ for all $Q \in P^{\mathcal{F}}$. P is said to be \mathcal{F} -essential, if P is \mathcal{F} -centric and fully normalized in \mathcal{F} , and $\text{Out}_{\mathcal{F}}(P)$ contains a strongly p -embedded subgroup (see [1, Definition A.6]). Now we are ready to introduce an important set, which was firstly introduced in [2, Notation 2.5].

Let p be a prime, \mathcal{F} be a saturated fusion system on a finite group S . Then define

$$\mathcal{E}_{\mathcal{F}}^* := \{Q \leq S \mid Q \text{ is } \mathcal{F}\text{-essential, or } Q = S\}.$$

These concepts will play an important role in the discovery of criteria for a fusion system $\mathcal{F}_S(G)$ to be supersolvable.

Let G be a finite group and S be a Sylow p -subgroup of G . In many cases, if certain subgroups of S satisfy certain properties, then G is p -nilpotent or p -supersolvable. On the other hand, on the assumption that a smaller collection of subgroups of S satisfy some properties, for example, they are strong p -CAP-subgroups of G , then G is not necessarily a p -supersolvable subgroup. But actually, this does not mean that there is nothing to do with the structure of the finite group G . Notice that the fusion system $\mathcal{F}_S(G)$ is a description of the structure of G , we wonder if we can characterize the structure of $\mathcal{F}_S(G)$. Fortunately, in this paper, we obtain several theorems and corollaries to characterize the structure of $\mathcal{F}_S(G)$ under the assumption that certain subgroups of S are strong p -CAP-subgroups, strong CAP-subgroups or partial CAP-subgroups of G , which consist of the second part (Section 4). In particular, Theorem 4.3 gives a new criteria for $\mathcal{F}_S(G)$ to be supersolvable under the assumption that every subgroup of $S \in \text{Syl}_p(G)$ with order $|D|$ ($1 \leq D < S$) is a strong CAP-subgroup of G , which is first introduced in our paper. These results highlight the paper considerably.

2 Preliminaries

In this section, several important lemmas are introduced to support our characterizations. Lemma 2.1 showcases the intermediate and inheritance property which have been frequently mentioned above. Theorem 2.2 plays an vital role in our characterizations. Lemma 2.4 and 2.5 are basic properties about supersolvability of fusion systems.

Lemma 2.1. *Let H be a strong p -CAP-subgroup of G and $N \trianglelefteq G$. Then the following statements are true.*

- (1) N is a strong p -CAP-subgroup of G .
- (2) H is a strong p -CAP-subgroup for any $K \geq H$.
- (3) If $H \geq N$, then H/N is a strong p -CAP-subgroup of G/N .
- (4) If $(|H|, |N|) = 1$, then HN/N is a strong p -CAP-subgroup of G/N .

Proof. (1) It follows from $N \trianglelefteq K$ for any $N \leq K \leq G$ and [9, Lemma 2.1(a)] that N is a p -CAP-subgroup for any $N \leq K \leq G$. Hence N is a strong p -CAP-subgroup of G .

(2) Since H is a strong p -CAP-subgroup of G , H is a p -CAP-subgroup for any $H \leq R \leq G$. Therefore H is a p -CAP-subgroup for any $H \leq R \leq K$. Hence we conclude that H is a strong p -CAP-subgroup of K .

(3) Since H is a p -CAP-subgroup for any $H \leq K \leq G$, it follows from [9, Lemma 2.1(b)] and $N \trianglelefteq K$ that H/N is a p -CAP-subgroup for any $H/N \leq K/N \leq G/N$. Thus it indicates that H/N is a strong p -CAP-subgroup of G/N .

(4) Since H is a p -CAP-subgroup for any $HN \leq K \leq G$, and $(|H|, |N|) = 1$, it follows from [9, Lemma 2.1(c)] and $N \trianglelefteq K$ that HN/N is a p -CAP-subgroup for any $HN/N \leq K/N \leq G/N$. Hence HN/N is a strong p -CAP-subgroup of G/N . \square

According to [16, Section 9.5], a class of finite groups \mathfrak{F} is said to be a formation if every image of an \mathfrak{F} -group is an \mathfrak{F} -group and if $G/N_1 \cap N_2$ belongs to \mathfrak{F} whenever G/N_1 and G/N_2 belong to \mathfrak{F} . A formation \mathfrak{F} is said to be saturated if a finite group $G \in \mathfrak{F}$ whenever $G/\Phi(G)$.

Let \mathfrak{U} denotes the class of all supersolvable groups and \mathfrak{U}_p denotes the class of all p -supersolvable groups, where p is a prime. It is well known that both \mathfrak{U} and \mathfrak{U}_p are saturated formations, and we have $\mathfrak{U}_p \supseteq \mathfrak{U}$.

Let \mathfrak{H} be a non-empty class of groups. According to [3, Definition 1.2.9, Definition 2.3.18], a chief factor H/K of a group G is called \mathfrak{H} -central in G if $[H/K] * G$ belongs to \mathfrak{H} , where $[H/K] * G$ is the semi-direct product $[H/K](G/C_G(H/K))$ if H/K is abelian and $G/C_G(H/K)$ if H/K is non-abelian. A normal subgroup N of a group G is said to be \mathfrak{H} -hypercentral in G if every chief factor of G below N is \mathfrak{H} -central in G . By generalised Jordan-Holder Theorem, we conclude that the product of \mathfrak{H} -hypercentral normal subgroups of a group G is \mathfrak{H} -hypercentral in G as well. Thus every group G possesses a unique maximal normal \mathfrak{H} -hypercentral subgroup namely the \mathfrak{H} -hypercentre of G and denoted by $Z_{\mathfrak{H}}(G)$. Apply the generalised Jordan-Holder Theorem again, it yields that every chief factor of G below $Z_{\mathfrak{H}}(G)$ is \mathfrak{H} -central in G .

Theorem 2.2. *Let p be a prime number and P be a normal p -subgroup of G . If every cyclic subgroup of P of prime order or 4 is a strong p -CAP-subgroup of G , then $P \leq Z_{\mathfrak{U}}(G)$, where $Z_{\mathfrak{U}}(G)$ denotes the \mathfrak{U} -hypercentre of G .*

Proof. Assume that the theorem is not true and choose G and P for which it fails. Then there exists a chief factor of G below P which is not of prime order. Between all of the non-cyclic chief factors of G below P , we choose L/K such that $|L|$ is of minimal size.

Assume there exists an element x of L of order p or 4 which is not in K . Then $\langle x \rangle$ is a strong p -CAP-subgroup of G . Consider a chief series of G passing through L , namely Γ :

$$1 = L_0 < L_1 < \cdots < L_{s-1} < L_s = L < L_{s+1} < \cdots < L_{m-1} < L_m = G.$$

It follows from L/K below P that $\langle x \rangle$ covers or avoids every L_{i+1}/L_i , $i = 0, 1, \dots, s-1$. Since $L_i \geq \langle x \rangle$, $i \geq s$, we conclude that $\langle x \rangle$ avoids L_{i+1}/L_i , $i \geq s$. Hence $\langle x \rangle$ covers or avoids each chief factor of Γ . By the choice of L/K , L_{i+1}/L_i , $i = 0, 1, \dots, s-1$ is of prime order. If $\langle x \rangle$ covers L_s/L_{s-1} , it follows from $L_s = L_{s-1}\langle x \rangle$ that L_s/L_{s-1} is of prime order. Hence we conclude that $L \leq Z_{\mathfrak{U}}(G)$. This implies that L/K is of prime order, a contradiction. Hence we have that $\langle x \rangle$ avoids L_s/L_{s-1} . Then $L_s = L_s \cap \langle x \rangle = L_{s-1} \cap \langle x \rangle$, i.e. $x \in L_{s-1}$. Thus we have that $L_{s-1}K > K$, which implies that $L_s = L_{s-1}K$. Since $L_{s-1}K/L_{s-1}$ is G -isomorphic to $L_{s-1}/(K \cap L_{s-1})$, we

conclude that $L_{s-1}/(K \cap L_{s-1})$ is a chief factor of G as well. By the minimality of $|L|$, we get that $L_{s-1}/(K \cap L_{s-1})$ is of prime order, i.e. L/K is of prime order, a contradiction. Hence it yields that every element of L of order p or 4 is contained in K .

Let X denote the intersection of the centralisers of the chief factors of G below K . Since all chief factor below K is of prime order, it follows that all of them are cyclic. Let

$$1 < K_1 < K_2 < \cdots < K_q = K.$$

denotes part of a chief series of G passing through K . Then $[K_i, K] \leq K_{i-1}$, $i = 2, 3, \dots, q$. Hence X stabilises a chain of subgroups of K . By applying [7, Chapter A, Corollary 12.4], $O^p(X)$ centralises K . In particular, $O^p(X)$ centralises every element of prime order or order 4 of L . By [14, Chapter IV, Satz 5.12], $O^p(X)$ centralises L . Thus $X/C_X(L/K)$ is a normal p -subgroup of $G/C_X(L/K)$. By [14, Chapter B, Proposition 3.12], X centralizes L/K . This implies that L/K can be regarded as an irreducible G/X -module over the finite field of p -elements. Note that every chief factor U/V of G below K is of order p , hence we conclude that $G/C_G(U/V)$ is cyclic of order dividing $p - 1$. Consequently, G/X is abelian of exponent dividing $p - 1$. By applying [7, Chapter B, Theorem 9.8], L/K has order p . This final contradiction shows that no such counterexample G exists. \square

It follows from every cyclic chief factor of order 2 in a chief series of a group is central that we have the following corollary:

Corollary 2.3. *Let P be a normal 2-subgroup of G . If every cyclic subgroup of P of order 2 or 4 is a strong p -CAP-subgroup of G , then P lies in the nilpotent hypercentre of G , namely $Z_\infty(G)$.*

Lemma 2.4 ([2, Lemma 2.6]). *Let p be a prime and \mathcal{F} be a saturated fusion system on a finite p -group S . Assume that the fusion system $N_{\mathcal{F}}(Q)$ is supersolvable for any $Q \in \mathcal{E}_{\mathcal{F}}^*$, then \mathcal{F} is supersolvable.*

Lemma 2.5 ([2, Lemma 2.9]). *Let G be a finite group, $p \in \pi(G)$, and S be a Sylow p -subgroup of G . Suppose that for any proper subgroup H of G with $O_p(G) < S \cap H$ and $S \cap H \in \text{Syl}_p(H)$, the fusion system $\mathcal{F}_{S \cap H}(H)$ is supersolvable. Assume additionally that $O_p(G) \leq Z_{\mathfrak{U}}(G)$. Then $\mathcal{F}_S(G)$ is supersolvable.*

Lemma 2.6 ([11, Proposition 3.1]). *Let P be a non-identity normal subgroup of G with $|P| = p^n$, where p is a prime divisor of $|G|$. Suppose that $n > 1$, and there exists an integer $n > k \geq 1$ that any subgroup of P of order p^k and any cyclic subgroup of P of order 4 (if P is a non-abelian 2-group and $n - 1 > k = 1$) are \mathfrak{U} -embedded in G . Then $P \leq Z_{\mathfrak{U}}(G)$.*

3 Characterizations for p -supersolvability of finite groups

Since we have several useful lemmas and theorems above, we are now ready to prove our main results.

Theorem 3.1. *Let p be a prime and G be a group of finite order. Suppose that every cyclic subgroup of G with order p or 4 (if $p = 2$) is a strong p -CAP-subgroup of G , then G is p -supersolvable.*

Proof. Suppose that the theorem is false and let G be a counterexample of minimal size. For the ease of reading we break the argument into separately stated steps.

Step 1. G is a minimal non- p -supersolvable group.

Let M be a maximal subgroup of G , and L be a subgroup of M of order p or 4 ($p = 2$). By Lemma 2.1(2), we conclude that L is a strong p -subgroup of M , hence M satisfies the assumption of the theorem. By the minimality of G , M is p -supersolvable. It follows from the inheritance of p -supersolvability that every proper subgroup of G is p -supersolvable, i.e. G is a minimal non- p -supersolvable group.

Step 2. $O_{p'}(G) = 1$, and $F(G) = O_p(G)$.

Set $\bar{G} = G/O_{p'}(G)$. Suppose that \bar{L} is a cyclic subgroup of \bar{G} of order p or 4 ($p = 2$). Then we can write $\bar{L} = LO_{p'}(G)/O_{p'}(G)$, where L is a cyclic subgroup of G of order p or 4. It follows from our assumption that L is a strong p -CAP-subgroup of G . Since $(|L|, |O_{p'}(G)|) = 1$, we conclude from Lemma 2.1(4) that $LO_{p'}(G)/O_{p'}(G) = \bar{L}$ is a strong p -CAP-subgroup of \bar{G} . Therefore \bar{G} satisfies the hypothesis of the theorem. Suppose that $O_{p'}(G) > 1$, then \bar{G} is p -supersolvable by our choice of G . Then by inheritance of p -supersolvability we get that G is p -supersolvable, which contradicts the fact that G is not p -supersolvable and step 2 has finished.

Step 3. $F(G) \leq Z_{\mathfrak{U}}(G)$.

Since $F(G) = O_p(G)$, a normal p -subgroup of G , by Theorem 2.2, it indicates that $O_p(G) \leq Z_{\mathfrak{U}}(G)$.

Step 4. $G = F^*(G)$.

Suppose that $F^*(G) < G$. Then by the p -supersolvability of $F^*(G)$ we get that $F^*(G) = F(G) = O_p(G)$. By step 3 we conclude that $F(G)$ is \mathfrak{U} -hypercentral in G . It follows from [7, Chapter IV, Theorem 6.10] that $[G^{\mathfrak{U}}, F(G)] = 1$, i.e. $G/C_G(F(G)) \in \mathfrak{U}$. By $C_G(F^*(G)) \leq F^*(G)$ and inheritance of supersolvability, we have that $G/F(G) \in \mathfrak{U}$. Since every G -chief factor below $F(G)$ is of order p , it follows that G is p -supersolvable, a contradiction. Hence it yields that $G = F^*(G)$.

Step 5. $G/Z(G)$ is non-abelian simple and G is perfect.

By step 4 we conclude that $G = F^*(G) = F(G)E(G)$. Since G is not nilpotent, it follows that $E(G)$ is not contained in $F(G)$. Hence we can find a component H of G . Then H is normal in $F^*(G) = G$, and $H/Z(H)$ is non-abelian simple. By step 2, it indicates that $p \mid |H|$. In particular, H is not p -supersolvable, hence we conclude that $H = G$, i.e. $G/Z(G)$ is non-abelian simple and G is perfect.

Step 6. Final contradiction.

Let M be a maximal subgroup of G which does not contain $Z(G)$. Then $MZ(G) = G$ and it yields that $M \triangleleft G$. Since $G/M = Z(G)M/M \cong Z(G)/M \cap Z(G)$, which is abelian, it follows that $G' \leq M$, contradicting the fact that G is perfect. Thus no such M exists and $Z(G) \leq \Phi(G)$. Now let G/A be a chief factor of G . Since $A \triangleleft G$, it follows that $A \leq Z(G)$, therefore we have that $A = Z(G)$. Hence $Z(G)$ belongs to all chief series of G . If $p \mid |G/Z(G)|$, let x be an element of order p or 4 ($p = 2$). It follows from our assumption that $\langle x \rangle$ is a strong p -CAP-subgroup of G . Then $\langle x \rangle$ covers or avoids $G/Z(G)$. If $G = G\langle x \rangle = Z(G)\langle x \rangle$, it follows that $G/Z(G)$ is abelian, a contradiction to the fact that $G = H$ is semi-simple. Hence we have that $G \cap \langle x \rangle = \langle x \rangle = Z(G) \cap \langle x \rangle$, i.e. $x \in Z(G)$. By the randomness of x , we conclude from [14, Chapter IV, Satz 5.5] that G is p -nilpotent, contradicting the fact that G is not p -supersolvable. Thus it yields that $p \nmid |G/Z(G)|$. If $Z(G)$ has non-trivial p' -part, it follows that $O_{p'}(G) > 1$, a contradiction to step 2. Thus $Z(G)$ is a p -group and then $Z(G) \leq O_p(G) = F(G)$. However, since G is semi-simple, it indicates from $F(G) \triangleleft G$ that $F(G) \leq Z(G)$. Thus we get that $F(G) = Z(G)$. By step 2, we conclude that every G -chief factor below $Z(G)$ is of order p . It follows that G is p -supersolvable, a contradiction. Finally we have that no such counterexample of G exists and we are done. \square

Theorem 3.2. *Let \mathfrak{F} be a saturated formation containing \mathfrak{U} , and let G be a group with a normal subgroup H such that $G/H \in \mathfrak{F}$. If for any $p \in \pi(G)$, every cyclic subgroup of $F^*(H)$ of order p or 4 (if $p = 2$) is a strong p -CAP-subgroup of G , then $G \in \mathfrak{F}$.*

Proof. We assume that $H > 1$, and p be a prime such that $p \mid |F^*(H)|$. By our assumption, Lemma 2.1(2) and Theorem 3.1 that $F^*(H)$ is p -supersolvable for all primes $p \mid |F^*(H)|$. Therefore $F^*(H)$ is supersolvable and so $F^*(H) = F(H)$. By definition of $F(H)$ and Theorem 2.2, we conclude that $F^*(H) = F(H) \leq Z_{\mathfrak{U}}(G)$. By [7, Chapter IV, Theorem 6.10], it follows that $G/C_G(F(H)) \in \mathfrak{U}$. Since $G/H \in \mathfrak{F}$ and $\mathfrak{F} \supseteq \mathfrak{U}$, it yields that $G/C_H(F(H)) \in \mathfrak{F}$. Since $C_H(F^*(H)) \leq F^*(H)$, it implies from the inheritance of \mathfrak{F} that $G/F^*(H) = G/F(H) \in \mathfrak{F}$. Since any G -chief factor below $F(H) \leq Z_{\mathfrak{U}}(G)$ is of prime order, and \mathfrak{F} contains \mathfrak{U} , we get that $G \in \mathfrak{F}$ and we are done. \square

Theorem 3.3. *Let G be a finite group and S a Sylow p -subgroup of G , where p is a prime divisor of $|G|$. If the order of S is larger than p , every maximal subgroup of S is a partial CAP-subgroup of G , every cyclic subgroup of order 4 of G is a partial CAP-subgroup of G ($p = 2$ and S is non-abelian), then G is p -supersolvable.*

Proof. Denote by $|S| \cdot p^{-1}$ by d . Suppose that $d = 1$, then we obtain that $|S| = p$, a contradiction to the fact that $|S| > p$. Hence we assert that $d > 1$. Then we obtain that every subgroup of S of order d is a partial CAP-subgroup of G , every cyclic subgroup of order 4 of G is a partial CAP-subgroup of G ($p = 2$ and S is non-abelian). Applying [5, Theorem C], we conclude that G is a p -soluble group. Now, we claim that G is p -supersolvable.

Suppose that our claim is false, in other words, there exists a group G satisfying such conditions that G is not p -supersolvable. Without loss of generality, let G be a counterexample with $|G|$ the smallest. At this time we would like M to be a minimal normal subgroup of G . Since G is already p -soluble, it follows that M is either a p -group or a p' -group. If M is a p' -group, let Q/M be a

maximal subgroup of SM/M . Then $Q \leq SM$, and so $M = O_{p'}(Q)$. Using the Schur-Zassenhaus Theorem, we get that there exists a p -group Q_1 such that $Q_1 \times M = Q$. Denote by S_1 the Sylow p -subgroup of SM containing Q_1 . Then $S_1^g = S$, where g is an element of SM by Sylow Theorem. Therefore $(Q_1M)^g = Q_1^gM = Q \leq SM$, and so $Q/M = Q_1^gM/M$ can be considered as the image of a maximal subgroup Q_1^g of S under the quotient morphism induced by M . Applying [8, Lemma 2], we obtain that every maximal subgroup Q/M of MS/M is a partial CAP -subgroup of G/M . Now suppose that M is a p -group, and so $M \leq O_p(G) \leq S$. Consider the maximal subgroup Q/M of S/M . Then we see that Q is a maximal subgroup of S containing M , and we conclude again from [8, Lemma 2] that Q/M is a partial CAP -subgroup of G/M . Therefore, every maximal subgroup of G/M is a partial CAP -subgroup, and so we obtain from the minimal choice of G that G/M is p -supersolvable.

Notice that the class of p -supersolvable groups is a saturated formation \mathfrak{U}_p , we may assume that M is the unique minimal normal subgroup of G . Assume that M is a p' -group. Then it follows from the fact G/M is p -supersolvable that G is p -supersolvable, a contradiction. Now, suppose on the contrary that M is a p -group. Then we claim that M is not contained in $\Phi(S)$. If $M \leq \Phi(S)$, then we conclude from the fact \mathfrak{U}_p is a saturated formation that G is p -supersolvable, another contradiction. Hence $M \not\leq \Phi(S)$, and so there exists a maximal subgroup S_1 of S such that $M \not\leq S_1$. Hence S_1 does not cover the chief factor $M/1$. Since S_1 is a partial CAP -subgroup of G , we obtain from the uniqueness of M that S_1 must avoid the chief factor $M/1$. Therefore $M \cap S_1 = 1 \cap S_1 = 1$, and so M is a cyclic subgroup of order p since $M \cdot S_1 = S$. Thus, as G/M is p -supersolvable, it follows directly that G is p -supersolvable, which leads to the final contradiction. Thus we assert that our claim is true, and so G is p -supersolvable. \square

As a direct application, we obtain the following two corollaries for characterizations of p -supersolvability of G under the assumption that certain subgroups of $S \in \text{Syl}_p(G)$ are CAP -subgroups or strong p - CAP -subgroups of G immediately.

Corollary 3.4. *Let G be a finite group and S a Sylow p -subgroup of G , where p is a prime divisor of $|G|$. If the order of S is larger than p , every maximal subgroup of S is a CAP -subgroup of G , every cyclic subgroup of order 4 of G is a CAP -subgroup of G ($p = 2$ and S is non-abelian), then G is p -supersolvable.*

Corollary 3.5. *Let G be a finite group and S a Sylow p -subgroup of G , where p is a prime divisor of $|G|$. Suppose that the order of S is larger than p , every maximal subgroup of S and every cyclic subgroup of order 4 of G ($p = 2$ and S is non-abelian) is a strong q - CAP -subgroup of G for any prime divisor q of $|G|$, then G is p -supersolvable.*

Proof. Let H be a subgroup of G such that H is a strong q - CAP -subgroup of G for any prime divisor q of $|G|$. Then for any chief factor U/V of G , it follows that there exists a prime divisor r of $|G|$ such that $r \mid |U/V|$. Hence we conclude that H covers or avoids U/V , and we obtain from the choice of U/V that H is a CAP -subgroup of G , and so a partial CAP -subgroup of G . Hence every maximal subgroup of S and every cyclic subgroup of order 4 of G ($p = 2$ and S is non-abelian) is

a partial CAP -subgroup of G , and so we conclude from Corollary 3.4 that G is p -supersolvable, as needed. \square

4 Characterizations for supersolvability of $\mathcal{F}_S(G)$

In this section, we investigate the structure of $\mathcal{F}_S(G)$ under the assumption that certain subgroups of G are strong p - CAP -subgroups, strong CAP -subgroups or partial CAP -subgroups of G , and obtain several criteria for fusion system $\mathcal{F}_S(G)$ to be supersolvable and finite group G to be p -nilpotent. The proofs of the theorems strongly rely on the proceeding results we have obtained in Section 3.

Theorem 4.1. *Let G be a finite group, p be a prime divisor of $|G|$, and S be a Sylow p -subgroup of G . Suppose that any cyclic subgroup of S with order p or 4 (if $p=2$) is a strong p - CAP -subgroup of G , then $\mathcal{F}_S(G)$ is supersolvable.*

Proof. Assume that the theorem is false, and let G be a counterexample of minimal order. Now denote $\mathcal{F}_S(G)$ by \mathcal{F} .

Step 1. Let H be a proper subgroup of G such that $S \cap H \in \text{Syl}_p(H)$ and $|S \cap H| \geq p^2$. Then $\mathcal{F}_{S \cap H}(H)$ is supersolvable.

By our hypothesis, every cyclic subgroup T of $S \cap H$ with order p or 4 (If $p = 2$) is a strong p - CAP -subgroup of G . Then every cyclic subgroup T of $S \cap H$ with order p or 4 (If $p = 2$) is a strong p - CAP -subgroup of H by Lemma 2.4 (2). Hence H satisfies the hypothesis of the theorem and it follows from the minimal choice of G that $\mathcal{F}_{S \cap H}(H)$ is supersolvable.

Step 2. Let $Q \in \mathcal{E}_{\mathcal{F}}^*$, then $|Q| \geq p^2$. If moreover that $Q \not\trianglelefteq G$, then $N_{\mathcal{F}}(Q)$ is supersolvable.

Suppose that there exists a subgroup $Q \in \mathcal{E}_{\mathcal{F}}^*$ such that $|Q| < p^2$. Then there is a subgroup R of S such that $|R| = p$, and $Q < R$. It follows directly that $R \leq C_S(Q)$. Since $Q < R \leq S$, we conclude from Q is a member of $\mathcal{E}_{\mathcal{F}}^*$ that Q is \mathcal{F} -essential. By the definition, Q is \mathcal{F} -centric. Hence $R \leq C_S(Q) = Z(Q) \leq Q$, a contradiction. Thus $|Q| \geq p^2$.

Assume that Q is not normal in G . Therefore $N_G(Q)$ is a proper subgroup of G . Since $Q \in \mathcal{E}_{\mathcal{F}}^*$, Q is fully \mathcal{F} -normalized or $Q = S$. Clearly S is fully \mathcal{F} -normalized, hence Q is always fully \mathcal{F} -normalized. By the argument below [1, Definition 2.4], $S \cap N_G(Q) = N_S(Q) \in \text{Syl}_p(N_G(Q))$. Since $|N_S(Q)| \geq |Q| \geq p^2$, it yields that $N_G(Q)$ satisfies the hypothesis of Step 1, and so $\mathcal{F}_{N_S(Q)}(N_G(Q)) = N_{\mathcal{F}}(Q)$ is supersolvable.

Step 3. $|O_p(G)| \geq p^2$.

Assume that there does not exist a subgroup $Q \in \mathcal{E}_{\mathcal{F}}^*$ such that $Q \trianglelefteq G$. Then for each $Q \in \mathcal{E}_{\mathcal{F}}^*$, the fusion system $N_{\mathcal{F}}(Q)$ is supersolvable by Step 2. By Lemma 2.4, \mathcal{F} is supersolvable, a contradiction. Thus there exists a subgroup $Q \in \mathcal{E}_{\mathcal{F}}^*$ such that $Q \trianglelefteq G$. Hence we conclude from Step 2 that $|O_p(G)| \geq |Q| \geq p^2$.

Step 4. $O_p(G) \leq Z_{\mathfrak{U}}(G)$.

It follows from $|O_p(G)| \geq p^2$ that any subgroup T of $O_p(G)$ of order p or any cyclic subgroup T of $O_p(G)$ of order 4 (If $p = 2$) is a strong p -CAP-subgroup of G . Then it yields from Theorem 2.5 that $O_p(G) \leq Z_{\mathfrak{U}}(G)$ and this part is complete.

Step 5. Final contradiction.

Suppose that H is a proper subgroup of G such that $O_p(G) < S \cap H$ and $S \cap H \in \text{Syl}_p(H)$. By Step 1 and Step 3, $|S \cap H| > |O_p(G)| \geq p^2$ and so $\mathcal{F}_{S \cap H}(H)$ is supersolvable. Since $O_p(G) \leq Z_{\mathfrak{U}}(G)$ by Step 4, it follows directly from Lemma 2.5 that $\mathcal{F}_S(G)$ is supersolvable, a contradiction. Hence our proof is complete. \square

As a direct application of the theorem above, we obtain the following characterization for the structure of finite groups under the assumption that any cyclic subgroup of S with order p or 4 (if $p=2$) is a strong p -CAP-subgroup of G .

Corollary 4.2. *Let G be a finite group and S a Sylow p -subgroup of G , where p is a prime divisor of $|G|$ such that $(p-1, |G|) = 1$. Suppose that any cyclic subgroup of S with order p or 4 (if $p=2$) is a strong p -CAP-subgroup of G , then G is p -nilpotent.*

Proof. It follows from Theorem 4.1 that $\mathcal{F}_S(G)$ is supersolvable. Thus we conclude from [17, Theorem 1.9] that G is p -nilpotent, as desired. \square

Theorem 4.3. *Let G be a finite group, p be a prime divisor of $|G|$, and S be a Sylow p -subgroup of G . Suppose that there exists a subgroup D of S with order $1 \leq |D| < |S|$ such that every subgroup of S with order $|D|$ and every cyclic subgroup of S with order 4 (if S is non-abelian and $|D| = p = 2$) is a strong CAP-subgroup of G , then $\mathcal{F}_S(G)$ is supersolvable.*

Proof. Assume that the theorem is false, and let G be a counterexample of minimal order. Now denote $\mathcal{F}_S(G)$ by \mathcal{F} .

Step 1. Suppose that H be a proper subgroup of G such that $S \cap H \in \text{Syl}_p(H)$ and $|S \cap H| \geq p|D|$.

Then we conclude that the fusion system $\mathcal{F}_{S \cap H}(H)$ is supersolvable.

By our hypothesis, every subgroup T of $S \cap H$ with order $|D|$ or 4 (If S is non-abelian and $|D| = p = 2$) is a strong CAP-subgroup of G . Then every cyclic subgroup T of $S \cap H$ with order $|D|$ or 4 (If $S \cap H$ is non-abelian and $|D| = p = 2$) is a strong CAP-subgroup of H by the definition of strong p -CAP-subgroup [4]. Hence H satisfies the hypothesis of the theorem and it follows from the minimal choice of G that $\mathcal{F}_{S \cap H}(H)$ is supersolvable.

Step 2. Let $Q \in \mathcal{E}_{\mathcal{F}}^*$, then $|Q| \geq p|D|$. If moreover that $Q \not\trianglelefteq G$, then $N_{\mathcal{F}}(Q)$ is supersolvable.

Suppose that there exists a subgroup $Q \in \mathcal{E}_{\mathcal{F}}^*$ such that $|Q| < p|D|$. Then there is a subgroup R of S such that $|R| = |D|$, and $Q < R$. It follows directly that $R \leq C_S(Q)$. Since $Q < R \leq S$, we conclude from Q is a member of $\mathcal{E}_{\mathcal{F}}^*$ that Q is \mathcal{F} -essential. By the definition, Q is \mathcal{F} -centric. Hence

$R \leq C_S(Q) = Z(Q) \leq Q$, a contradiction. Thus $|Q| \geq p|D|$. Now assume that Q is not normal in G . Therefore $N_G(Q)$ is a proper subgroup of G . Since $Q \in \mathcal{E}_{\mathcal{F}}^*$, Q is fully \mathcal{F} -normalized or $Q = S$. Clearly S is fully \mathcal{F} -normalized, hence Q is always fully \mathcal{F} -normalized. By the argument below [1, Definition 2.4], $S \cap N_G(Q) = N_S(Q) \in \text{Syl}_p(N_G(Q))$. Since $|N_S(Q)| \geq |Q| \geq p|D|$, it yields that $N_G(Q)$ satisfies the hypothesis of Step 1, and so $\mathcal{F}_{N_S(Q)}(N_G(Q)) = N_{\mathcal{F}}(Q)$ is supersolvable.

Step 3. $|O_p(G)| \geq p|D|$.

Assume that there does not exist a subgroup $Q \in \mathcal{E}_{\mathcal{F}}^*$ such that $Q \trianglelefteq G$. Then for each $Q \in \mathcal{E}_{\mathcal{F}}^*$, the fusion system $N_{\mathcal{F}}(Q)$ is supersolvable by Step 2. By Lemma 2.4, \mathcal{F} is supersolvable, a contradiction. Thus there exists a subgroup $Q \in \mathcal{E}_{\mathcal{F}}^*$ such that $Q \trianglelefteq G$. Hence we conclude from Step 2 that $|O_p(G)| \geq |Q| \geq p|D|$. Now we claim that $O_p(G) \leq Z_{\mathfrak{U}}(G)$. Actually, it follows from $|O_p(G)| \geq p|D|$ that any subgroup T of $O_p(G)$ of order $|D|$ or any cyclic subgroup T of $O_p(G)$ of order 4 (If S is non-abelian, $|D| = p = 2$) is a strong CAP -subgroup of G . Now let W be an arbitrary strong CAP -subgroup of G . Then for any chief factor U/V of G , W avoids or covers U/V . Suppose that W covers U/V , i.e. $WV \geq U$, then we conclude from Dedekind modular law that $V(U \cap W) = WV \cap U = U$, and so $|U : V(U \cap W)| = |G : N_G(V(U \cap W))| = 1$, which indicates that W is a generalized CAP -subgroup by [11, Definition 1.1]. Notice that $HG = G$, $G \cap H = H$ is a generalized CAP -subgroup of G , where G is obviously a K - \mathfrak{U} -subgroup of G by [3, Page 236], we obtain from [11, Definition 1.2] that H is \mathfrak{U} -embedded in G . By the choice of W , any subgroup T of $O_p(G)$ of order $|D|$ or any cyclic subgroup T of $O_p(G)$ of order 4 (If S is non-abelian, $|D| = p = 2$) is \mathfrak{U} -embedded in G . Therefore, applying Lemma 2.6, it yields directly that $O_p(G) \leq Z_{\mathfrak{U}}(G)$ and this part is complete.

Step 4. Final contradiction.

Suppose that H is a proper subgroup of G such that $O_p(G) < S \cap H$ and $S \cap H \in \text{Syl}_p(H)$. By Step 1 and Step 3, $|S \cap H| > |O_p(G)| \geq p|D|$ and so $\mathcal{F}_{S \cap H}(H)$ is supersolvable. Since $O_p(G) \leq Z_{\mathfrak{U}}(G)$ by Step 3, it follows directly from Lemma 2.5 that $\mathcal{F}_S(G)$ is supersolvable, a contradiction. Hence our proof is complete. \square

Corollary 4.4. *Let G be a finite group, p be a prime divisor of $|G|$, and S be a Sylow p -subgroup of G . Suppose that there exists a subgroup D of S with order $1 \leq |D| < |S|$ such that every subgroup of S with order $|D|$ and every cyclic subgroup of S with order 4 (if S is non-abelian and $|D| = p = 2$) is a strong q - CAP -subgroup for any prime divisor q of $|G|$, then $\mathcal{F}_S(G)$ is supersolvable.*

Proof. Let H be an arbitrary subgroup of G such that H is a strong q - CAP -subgroup for any prime divisor q of $|G|$. Then for any chief factor U/V of a subgroup T of G containing H , it follows that there exists a prime divisor r of $|T|$ such that $r \mid |U/V|$. Then it follows from the fact H is a strong r - CAP -subgroup of G that H covers or avoids U/V . By the choice of U/V , we conclude that H is a CAP -subgroup of T , and so we obtain from the choice of T that H is a strong CAP -subgroup of G . Hence we obtain that there exists a subgroup D of S with order $1 \leq |D| < |S|$ such that every subgroup of S with order $|D|$ and every cyclic subgroup of S with order 4 (if S is

non-abelian and $|D| = p = 2$) is a strong *CAP*-subgroup of G . Applying Theorem 4.3, we get that the fusion system $\mathcal{F}_S(G)$ is supersolvable. \square

As a direct application of the theorem above, we obtain the following characterization for the structure of finite groups under the assumption that any subgroup of S with order $1 \leq |D| < |S|$ for some subgroup D of S or 4 (if S is non-abelian and $|D| = 2 = p$) is a *CAP*-subgroup of G or a strong *q-CAP*-subgroup of G for any prime divisor q of $|G|$.

Corollary 4.5. *Let G be a finite group and S a Sylow p -subgroup of G , where p is a prime divisor of $|G|$ such that $(p - 1, |G|) = 1$. Suppose that there exists a subgroup D of S with order $1 \leq |D| < |S|$ such that every subgroup of S with order $|D|$ and every cyclic subgroup of S with order 4 (if S is non-abelian and $|D| = p = 2$) is a strong *CAP*-subgroup of G , then G is p -nilpotent.*

Proof. It follows from Theorem 4.3 that $\mathcal{F}_S(G)$ is supersolvable. Thus we conclude from [17, Theorem 1.9] that G is p -nilpotent, as desired. \square

Corollary 4.6. *Let G be a finite group and S a Sylow p -subgroup of G , where p is a prime divisor of $|G|$ such that $(p - 1, |G|) = 1$. Suppose that there exists a subgroup D of S with order $1 \leq |D| < |S|$ such that every subgroup of S with order $|D|$ and every cyclic subgroup of S with order 4 (if S is non-abelian and $|D| = p = 2$) is a strong *q-CAP*-subgroup of G for any prime divisor q of $|G|$, then G is p -nilpotent.*

Proof. It follows from Corollary 4.4 that $\mathcal{F}_S(G)$ is supersolvable. Thus we conclude from [17, Theorem 1.9] that G is p -nilpotent, as desired. \square

References

- [1] M. Aschbacher, R. Kessar, B. Oliver, *Fusion Systems in Algebra and Topology*, Cambridge University Press, New York, 2011.
- [2] F. Aseeri, J. Kaspczyk, Criteria for supersolvability of saturated fusion systems. <https://doi.org/10.48550/arXiv.2305.09008>.
- [3] A. Ballester-Bolinches, L. M. Ezquerro, *Classes of Finite Groups*. In: *Mathematics and its Applications*, vol. 584. New York: Springer, 2006.
- [4] A. Ballester-Bolinches, L. M. Ezquerro, A. N. Skiba, Subgroups of finite groups with a strong cover-avoidance property. *Bull. Aust. Math. Soc.* **79** (2009) 499–506.
- [5] A. Ballester-Bolinches, L. M. Ezquerro, Y. Li, N. Su, On partial *CAP*-subgroups of finite groups, *J. Algebra* **431** (2015) 196–208.
- [6] D. A. Craven, *The theory of fusion systems*, Cambridge University Press, New York.
- [7] K. Doerk, T. Hawkes, *Finite Soluble Groups*. In: *De Gruyter Expositions in Mathematics*, vol. 4. Berlin-New York: Walter de Gruyter, 1992.

- [8] Y. Fan, X.Y. Guo, K.P. Shum, Remarks on two generalizations of normality of subgroups, *Chinese Ann. Math. A* **2** (2006) 169–176.
- [9] Z. Gao, S. Qiao, H. Shi, L. Miao, On p -CAP-subgroups of finite groups, *Front. Math. China* **15** (2020) 915–921.
- [10] W. Gaschütz, Praefrattini gruppen, *Arch. Math.* **13** (1962) 418–426.
- [11] W. Guo, A. N. Skiba, N. Yang, A generalized CAP-subgroup of a finite group. *Sci. China Math.* **58** (2015) 2133–2144.
- [12] X. Guo, P. Guo, K.P. Shum, On semi cover-avoiding subgroups of finite groups, *Journal of Pure and Applied Algebra* **209** (2007) 151–158.
- [13] X. Guo, K.P. Shum, Cover-avoidance properties and the structure of finite groups. *Journal of Pure and Applied Algebra* **181** (2003) 297 – 308.
- [14] B. Huppert, Endliche Gruppen I. In: Grund Math Wiss, vol. 134. Berlin-Heidelberg-New York: Springer-Verlag, 1967.
- [15] J. Liu, X. Guo, S. Li, The Influence of CAP* -Subgroups on the Solvability of Finite Groups. *Bull. Malays. Math. Sci. Soc.* **35** (2012) 227–237.
- [16] D. J. S. Robinson, A Course in the Theory of Groups. Springer-Verlag New York, 1996.
- [17] N. Su, On supersolvable saturated fusion systems, *Monatsh Math* **187** (2018) 171–179.