

# Isolated Diophantine numbers

F. Argentieri, L. Chierchia

February 2, 2024

## Abstract

In this short note, we discuss the topology of Diophantine numbers, giving simple explicit examples of Diophantine isolated numbers (among those with same Diophantine constants), showing that, *Diophantine sets are not always Cantor sets*.

General properties of isolated Diophantine numbers are also briefly discussed.

**Keywords:** Diophantine sets, Diophantine conditions, Cantor sets, KAM Theory, small divisor problems.

**MSC2010:** 37J40, 70H08, 11D75

## 1 Introduction

Diophantine numbers are irrational numbers badly approximated by rationals, namely, real numbers  $\xi$  satisfying, for some  $\gamma, \tau > 0$ ,

$$|\xi q - p| \geq \frac{\gamma}{q^\tau}, \quad \forall p \in \mathbb{Z}, q \in \mathbb{N} = \{1, 2, \dots\}. \quad (1)$$

Such numbers, that form a set of full Lebesgue measure in<sup>1</sup>  $\mathbb{R}$ , arise naturally in number theory and in small divisor problems in dynamics. Indeed, since the seminal works of C.L. Siegel, in the context of linearization of holomorphic diffeomorphisms around a fixed point [10], and of A.N. Kolmogorov, in the context of Hamiltonian systems [8], Diophantine conditions as in (1) (or higher dimensional analogs) are ubiquitous in perturbative Hamiltonian dynamics both in finite and infinite dimensions.

Let us denote by  $D_{\gamma, \tau}$  the Diophantine set of all real numbers satisfying condition (1) with fixed<sup>2</sup>  $0 < \gamma < 1/2$  and  $\tau \geq 1$ . Clearly,  $D_{\gamma, \tau}$  is a closed and nowhere dense set. It is, therefore, natural to ask whether  $D_{\gamma, \tau}$  is actually a Cantor set, i.e., if it is also a perfect set (no isolated points).

A (authoritative) place where the term Cantor set appears in association with Diophantine sets is Chapter III of the fundamental and beautiful book, *Lectures on Celestial Mechanics*, by C.L. Siegel and J.K. Moser [11]. In § 32–36 of [11], Moser, extending the previous text of Siegel<sup>3</sup>, includes a proof, in the analytic case, of his theorem on the persistence of invariant curves for area-preserving twist diffeomorphisms of the annulus [9]. As well known, one of the main hypothesis is that  $\omega/2\pi$  belongs to  $D_{\gamma, \tau}$  for given  $\gamma, \tau$ , where  $\omega$  denotes the rotation number of the unperturbed invariant curve. Moser call *admissible* such numbers and, at page 245, writes<sup>4</sup>: “*the set of admissible values for  $\omega$  form a Cantor set of positive measure*”. Although, it does not appear a formal statement about the sets  $D_{\gamma, \tau}$ , reading p. 245 of [11], one might be lead to the belief that Diophantine sets are Cantor sets.

<sup>1</sup>See, e.g., [11], end of § 25.

<sup>2</sup>Notice that  $D_{\gamma, \tau} = \emptyset$  whenever  $\gamma \geq 1/2$  (trivially, taking  $q = 1$  and the minimum over  $p$ ) or, (by Dirichlet’s Theorem), when  $\tau < 1$ .

<sup>3</sup>Compare with the 1971 Preface to the English Edition of [11].

<sup>4</sup>Obviously, Moser considers Diophantine sets with exponent  $\tau > 1$  (see p. 242), as it is well known that for  $\tau = 1$ ,  $D_{\gamma, \tau}$  is not a Cantor set (see, e.g., [3]).

However, it turns out that, in general, this is *not* the case: In § 1, we show that the quadratic numbers  $\alpha := (n + \sqrt{n^2 + 4})/2 = [n, n, n, \dots] = [\bar{n}]$  (in continued fraction expansion) are, for any  $n \geq 2$ , isolated in  $D_{\gamma, \tau}$  with  $\gamma := 1/\alpha$  and  $\tau := \log \alpha / \log n$ .

In § 2, we briefly review some general properties of isolated Diophantine points, proven in [1, 2], which show, in particular, that isolated Diophantine points are not that rare; § 3 contains concluding remarks.

## 2 Elementary examples of isolated Diophantine numbers

**Theorem 1** *Let  $n \in \mathbb{N}$ ,  $n \geq 2$  and define*

$$\alpha := \frac{n + \sqrt{n^2 + 4}}{2}, \quad \gamma := \frac{1}{\alpha}, \quad \tau := \frac{\log \alpha}{\log n}. \quad (2)$$

*Then,  $\alpha$  is an isolated point of  $D_{\gamma, \tau}$ .*

**Remark** (i) By definition of  $D_{\gamma, \tau}$ , it follows immediately that

$$I_{\gamma, \tau}(p, q) := \left\{ \xi \in \mathbb{R} : \left| \xi - \frac{p}{q} \right| < \frac{\gamma}{q^{\tau+1}} \right\} \subseteq \mathbb{R} \setminus D_{\gamma, \tau}, \quad \forall q \in \mathbb{N}, \forall p \in \mathbb{Z}. \quad (3)$$

(ii)  $D_{\gamma, \tau}$  is invariant by translations by integers, as, for  $k \in \mathbb{Z}$ ,  $\xi \in D_{\gamma, \tau} \iff \xi + k \in D_{\gamma, \tau}$ . Therefore,  $\alpha - n = \frac{\sqrt{n^2 + 4} - n}{2} = [0, \bar{n}] \in (0, 1)$  is an isolated point of  $D_{\gamma, \tau} \cap [0, 1]$ .

As one may expect, proofs make use of the theory of continued fractions; see [6] for general information. Let  $\gamma > 0$ ,  $\tau \geq 1$ ; let  $\xi$  be an irrational number and let:

$$\xi = [a_0, a_1, a_2, \dots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots}}}$$

be its continued fraction expansion,  $p_k/q_k = [a_0, a_1, \dots, a_k]$  its  $k^{\text{th}}$  convergent, and  $a'_k := [a_k, a_{k+1}, \dots]$  its  $k^{\text{th}}$  complete quotient.

**Lemma** *A number  $\xi$  belongs to  $D_{\gamma, \tau}$  if and only if*

$$\frac{q_{k+1}}{q_k^\tau} + \frac{1}{a'_{k+2} q_k^{\tau-1}} \leq \frac{1}{\gamma}, \quad \forall k \geq 0. \quad (4)$$

**Proof** From continued fraction theory, one knows that<sup>5</sup>

$$\xi \in D_{\gamma, \tau} \iff \left| \xi - \frac{p_k}{q_k} \right| \geq \frac{\gamma}{q_k^{\tau+1}}, \quad \forall k \geq 0. \quad (5)$$

Then<sup>6</sup>,

$$\begin{aligned} \left| \xi - \frac{p_k}{q_k} \right| &= \frac{1}{q_k (a'_{k+1} q_k + q_{k-1})} = \frac{1}{q_k^{\tau+1}} \frac{q_k^\tau}{a'_{k+1} q_k + q_{k-1}} \\ &= \frac{1}{q_k^{\tau+1}} \frac{q_k^\tau}{a_{k+1} q_k + q_{k-1} + \frac{q_k}{a'_{k+2}}} \\ &= \frac{1}{q_k^{\tau+1}} \frac{q_k^\tau}{q_{k+1} + \frac{q_k}{a'_{k+2}}} \\ &= \frac{1}{q_k^{\tau+1}} \left( \frac{q_{k+1}}{q_k^\tau} + \frac{1}{a'_{k+2} q_k^{\tau-1}} \right)^{-1}, \end{aligned}$$

<sup>5</sup>For a proof, see, e.g., Lemma 1, Appendix 8, p. 122 of [5].

<sup>6</sup> $p_{-1} := 1$ ,  $q_{-1} := 0$ ; for the first equality, see [6, §10.7].

and the claim follows from (5).  $\blacksquare$

**Proof** (of the Theorem 1) One immediately verifies that

$$\begin{cases} \alpha = n + \frac{1}{\alpha}, & n^\tau = \alpha, \\ \alpha = [n, n, n, n, \dots], \\ p_0 = n, q_0 = 1, p_1 = n^2 + 1, q_1 = n, a'_k = \alpha, q_{k+1} = p_k \ (\forall k \geq 0). \end{cases} \quad (6)$$

Thus, for  $k = 0$  we have

$$\left| \alpha - \frac{p_0}{q_0} \right| \stackrel{(6)}{=} \alpha - n \stackrel{(6)}{=} \frac{1}{\alpha} \stackrel{(2)}{=} \gamma. \quad (7)$$

For  $k \geq 1$ , using (6) and the facts that  $p_k/q_k \leq p_1/q_1$  and  $q_k \geq q_1$ , one finds

$$\begin{aligned} \frac{q_{k+1}}{q_k^\tau} + \frac{1}{a'_{k+2} q_k^{\tau-1}} &= \frac{p_k}{q_k} \frac{1}{q_k^{\tau-1}} + \frac{1}{\alpha q_k^{\tau-1}} \\ &\leq \frac{p_1}{q_1} \frac{1}{q_1^{\tau-1}} + \frac{1}{\alpha q_1^{\tau-1}} = \frac{n^2 + 1}{n^\tau} + \frac{1}{n^{\tau-1} \alpha} \\ &= \frac{n^2 + 1}{\alpha} + \frac{n}{\alpha^2} = \frac{1}{\alpha} \left( n^2 + 1 + \frac{n}{\alpha} \right) \\ &= \frac{1}{\alpha} (\alpha n + 1) = n + \frac{1}{\alpha} = \alpha \\ &= \frac{1}{\gamma}, \end{aligned}$$

which, together with (7) and the Lemma, shows that  $\alpha \in D_{\gamma, \tau}$ .

Next, because of (6),

$$\begin{aligned} \left| \alpha - \frac{p_1}{q_1} \right| &= \frac{p_1}{q_1} - \alpha = \frac{n^2 + 1}{n} - \alpha = \frac{1}{n} + n - \alpha \\ &= \frac{1}{n} - \frac{1}{\alpha} = \frac{1}{n\alpha^2} = \frac{1}{\alpha} \frac{1}{q_1 n^\tau} = \frac{1}{\alpha q_1^{\tau+1}} \\ &= \frac{\gamma}{q_1^{\tau+1}}. \end{aligned}$$

Such relation, together with (7), shows that  $\alpha$  separates the two intervals<sup>7</sup>  $I_{\gamma, \tau}(p_0, q_0)$  and  $I_{\gamma, \tau}(p_1, q_1)$ , and, therefore,  $\alpha$  is an *isolated point* of  $D_{\gamma, \tau}$ .  $\blacksquare$

### 3 General properties of isolated Diophantine numbers

General properties of isolated Diophantine numbers have been investigated in [1, 2], where proofs may be found. Let us briefly report here the main results in [1, 2].

The first result in [1] shows that isolated points are not that rare: Indeed, *any* Diophantine number has at least one equivalent representative<sup>8</sup>, which is isolated in some Diophantine set:

<sup>7</sup>Recall the definition of the open intervals  $I_{\gamma, \tau}(p, q)$  in (3).

<sup>8</sup>Recall that two irrational numbers  $\xi$  and  $\xi'$  are *equivalent* if and only if  $\xi' = \frac{a\xi + b}{c\xi + d}$  with integers  $a, b, c, d$  satisfying  $ad - bc = \pm 1$ , and that happens if and only if the continued fractions of  $\xi$  and  $\xi'$  differ only by a finite number of terms; compare [6, §10.11].

**Theorem 2** [1, Theorem B] Fix  $\gamma \in (0, \frac{1}{2})$ ,  $\tau \geq 1$ ,  $\alpha \in D_{\gamma, \tau}$ , and let  $m := \lceil \frac{3 \cdot 2^\tau}{\gamma} \rceil$ . Then, the equivalent Diophantine number  $\alpha' := \frac{m\alpha + 1}{(2m + 1)\alpha + 2}$  is an isolated point of  $D_{\gamma_\alpha, \tau_\alpha}$  for suitable  $\tau_\alpha > \tau$  and  $\gamma_\alpha > 0$ .

Actually, it can happen that a Diophantine number is *simultaneously isolated for infinitely many Diophantine sets*. More precisely<sup>9</sup>:

**Theorem 3** [1, Theorem A] For all  $\tau \geq 1$ , there exist  $\gamma > 0$  and  $\alpha \in D_{\gamma, \tau}$  such that  $\alpha$  is an isolated point for  $D_{\gamma_n, \tau_n}$ , for suitable sequences  $\tau_n \searrow \tau$ ,  $\gamma_n \searrow \gamma$ .

Even though these two theorems show the existence of many isolated Diophantine numbers, from the metric point of view, the typical situation seems to be that Diophantine sets are Cantor sets:

**Theorem 4** [2] Let  $\tau > \tau_0 := \frac{3 + \sqrt{17}}{2} = \overline{[3, 1, 1]}$ . Then, for almost all  $\gamma \in (0, 1/2)$ ,  $D_{\gamma, \tau}$  is a Cantor set.

## 4 Remarks

(i) The Diophantine exponent  $\tau_0 := \frac{3 + \sqrt{17}}{2}$  in Theorem 4 is certainly not optimal, and it would not be difficult to improve it. On the other hand, it is not so obvious what is the *optimal*  $\tau_0$ , for which the statement of Theorem 4 holds.

(ii) Diophantine sets, as pointed out in the Introduction, play a fundamental rôle in Dynamics, e.g., in the theory of exact symplectic twist diffeomorphisms. Arithmetic properties of the rotation number of an invariant curve of a twist diffeomorphism are, in particular, relevant for the renormalization point of view; compare [7]. Now, even though Theorem 2 above indicates that the property of being isolated for Diophantine numbers may not be a stable property under renormalization, it would be quite interesting to see if such a property does have a counter part in dynamics. For example,

*Does there exist a  $C^r$  exact symplectic twist diffeomorphism  $f$ ,  $r \geq 2$ , having an isolated invariant curve of rotation number  $\alpha$  that is not of bounded type, with  $\alpha$  isolated point of a suitable Diophantine set?*

(iii) A final comment on higher dimensional Diophantine sets.

Let  $n \geq 1$ ,  $\gamma, \tau > 0$ , and define

$$D_{\gamma, \tau}^n := \{\alpha \in \mathbb{R}^n : |q \cdot \alpha - p| \geq \frac{\gamma}{|q|^\tau}, \forall q \in \mathbb{Z}^n \setminus \{0\}, p \in \mathbb{Z}\}.$$

The analogous problem discussed in this note is<sup>10</sup>:

*For  $n \geq 2$ , do there exist  $\gamma, \tau > 0$  such that  $D_{\gamma, \tau}^n$  is not a Cantor set?*

Clearly, such a question may be more difficult to analyse due to the lack of the beautiful and powerful theory of continued fractions.

**Acknowledgements** We are indebted with Michel Waldschmidt for useful discussions and for his encouragement. We also thank, Yann Bugeaud for interesting comments.

<sup>9</sup>In [1] something stronger is proven, in the sense that the sequence  $\tau_n$  in Theorem 3 can be assigned arbitrarily up to small errors.

<sup>10</sup>Homogeneous Diophantine sets  $\mathbb{R}_{\gamma, \tau}^n := \{\omega \in \mathbb{R}^n : |\omega \cdot k| \geq \gamma/|k|^\tau, \forall k \in \mathbb{Z}^n, k \neq 0\}$  are trivially Cantor sets, since if  $\omega \in \mathbb{R}_{\gamma, \tau}^n$ , then also  $t\omega \in \mathbb{R}_{\gamma, \tau}^n, \forall t \geq 1$ ; see [4] for related questions on  $\mathbb{R}_{\gamma, \tau}^n$ .

## References

- [1] F. Argentieri, *Isolated points of Diophantine sets*. arXiv:2011.10267
- [2] F. Argentieri, *Diophantine sets in general are Cantor sets*. arXiv:2012.13998
- [3] E. Bombieri, *Continued fractions and the Markoff tree*. Expo. Math. 25 (2007), n.3, 187-213.
- [4] H. Broer, *Do Diophantine vectors form a Cantor bouquet?* J. Difference Equ. Appl.16 (2010), no.5-6, 433-434.
- [5] A. Celletti, and L. Chierchia, *KAM stability and Celestial Mechanics*, Memoirs of the AMS, Vol. 187, Number 878, (2007)
- [6] G.H. Hardy, and E.M. Wright: *Introduction to the Theory of Numbers*. 4th Edition, Clarendon Press, Oxford (1962)
- [7] MacKay, R. S., Stark, J. *Locally most robust circles and boundary circles for area-preserving maps*. Nonlinearity (1992), no.4, 867–888.
- [8] A.N. Kolmogorov, *On the Conservation of Conditionally Periodic Motions under Small Perturbation of the Hamiltonian*, Dokl. akad. nauk SSSR,1954, vol.98, pp.527–530. Engl. transl.: Stochastic Behavior in Classical and Quantum Hamiltonian Systems, Volta Memorial conference, Como,1977, Lecture Notes in Physics, vol. 93, Springer, 1979, pp.51–56.
- [9] J.K. Moser, *On invariant curves of area-preserving mappings of an annulus*, Nachr. Akad. Wiss. Göttingen Math.–Phys. Kl. II (1962), 1–20.
- [10] C.L. Siegel, *Iteration of analytic functions*, Ann. Math., 1 **28**, 607 (1942)
- [11] C.L. Siegel, and J.K. Moser *Lectures on Celestial Mechanics*, Springer 1971