Valuative invariants for linearized line bundles on a spherical variety

Chenxi Yin

Abstract

We give formulas for various valuative invariants of linearized ample line bundles on a projective spherical variety. Then we show how to use these formulas to study g-solitons on a Fano spherical variety. As a corollary, we show that for a Fano horospherical manifold X, the corresponding cone $(K_X)^\times$ always admits a Calabi-Yau cone metric.

1 Introduction

Let X be a Fano manifold. In the quest of Kähler–Einstein metrics, the so called α −invariant was introduced in [\[Tia87,](#page-27-0) [TY87\]](#page-28-0). It is explained in [\[CS08\]](#page-26-0) that the α −invariant can be defined as:

 $\alpha(X, -K_X) := \inf \{ \text{lct}(X, D) | D \text{ is an effective } \mathbb{Q}-\text{divisor on } X \text{ and } D \sim_{\mathbb{Q}} -K_X \}$

where:

$$
let(X, D) := \inf_{E} \frac{A_X(E)}{\text{ord}_E(D)} = \inf_{v} \frac{A_X(v)}{\text{ord}_v(D)}
$$

Here A_X is the log discrepancy function, E runs through all the prime divisors over X. meaning that there is a proper birational morphism $\mu: Y \to X$ such that E is a prime divisor on Y , whereas v runs through all the non-trivial valuations with finite log discrepancy. A famous theorem of Tian says that:

Theorem 1.1. [\[Tia87\]](#page-27-0) Let X be a Fano manifold. If $\alpha(X, -K_X) > \frac{\dim(X)}{\dim(X)+1}$, then we have a Kähler–Einstein metric on X.

We see that the α −invariant is valuative and closely related to singularities. This suggests there should exist a theorem relating the existence of a Kähler–Einstein metric on a Fano manifold to the analysis of singularity via the valuative theory. This program was achieved after great efforts (see for example [\[XZ20,](#page-28-1) [Li22,](#page-27-1) [LXZ22\]](#page-27-2)). Several valuative invariants have been introduced in the process. In particular, the δ −invariant is defined in [\[FO18\]](#page-26-1) and the β -invariant is defined in [\[Li17,](#page-27-3) [Fuj19\]](#page-26-2). It is worth mentioning that the idea of linking the existence of a Kähler–Einstein metric and singularities is also in the pioneering paper of Nadel [\[Nad90\]](#page-27-4).

Another approach that is used a lot is the theory of K-stability or Ding-stability. To define these notions of stability, degenerations called test configurations of the manifold are considered. Then one can define numerical invariants such as Futaki invariants and Ding invariants for test configurations as obstructions to the existence of Kähler–Einstein metric. To consider all the test configurations together paves the way to the definitions of K-stability and Ding-stability. The direction from the existence of a Kähler–Einstein metric to the Kpolystability of a Q−Fano variety is proved in [\[Ber16\]](#page-25-0). The other direction is carried out by [\[CDS15a,](#page-26-3) [CDS15b,](#page-26-4) [CDS15c,](#page-26-5) [Tia15\]](#page-27-5). The equivalence between K-polystability and Dingpolystability for Q−Fano varieties is proved in [\[Fuj19\]](#page-26-2).

The two points of views mentioned above use seemingly different algebraic theories, but they are actually closely related and can be unified by the language of filtrations. To exchange information between the two aspects, one uses the so called special test configurations (Definition [7.1\)](#page-17-0).

Theories mentioned above are abstract in nature. It is usually difficult to say if a Fano manifold admits a Kähler–Einstein metric, or equivalently, is Ding (K) -polystable or not. It is seldom an easy task to compute out the valuative invariants. But the task becomes easier when a group acts on the manifold. Works in this direction include [\[WZ04,](#page-28-2) [Del17,](#page-26-6) [Del20,](#page-26-7) [BJ20,](#page-25-1) [Gol20\]](#page-26-8), to name just a few.

The purpose of this paper is to study valuative invariants for linearized ample line bundles on spherical varieties. Then we focus on Q−Fano case and use valuative information to study Ding-stability as well as the existence of different canonical metrics.

Spherical varieties are varieties with large symmetry. To be more precise, let G be a connected reductive complex algebraic group and B a Borel subgroup. Let X be a normal G−variety. We say it is G−spherical if it contains an open B−orbit. Toric varieties form a subclass of spherical varieties. There are a lot of combinatorial information associated to a spherical variety, the most important one being the valuation cone \mathcal{V} , which is the set of G−invariant Q−valued valuations. Usually the set of valuations for a variety is hard to describe, but ν is rather explicit and has a nice structure. It is actually a polyhedral convex cone inside a $\mathbb{Q}-\text{vector space }N\otimes\mathbb{Q}$. This nice property allows us to do computations and to have nice conclusions. Sometimes we also use the closure of it inside $N \otimes \mathbb{R}$, which we denote as $\mathcal{V}_{\mathbb{R}}$. When we have a G-linearized line bundle on a G-variety, we can define the moment polytope Δ^+ (Definition [4.2\)](#page-7-0), which encodes the most important combinatorial information about a G−linearized line bundle. We can also define a polytope Δ for every B−eigenvector $s \in H^0(X, L)$ with eigenvalue χ . More detailed information about spherical varieties and G-linearized line bundles are given in Section [4.](#page-6-0) We have $\Delta^+ = \Delta + \chi$. In this paper, we mainly deal with spherical X with \mathbb{Q} −Cartier K_X .

Our first result is a generalization of the results in [\[Del17,](#page-26-6) [BJ20,](#page-25-1) [Gol20\]](#page-26-8). After introducing different valuative invariants in Sections [2](#page-3-0) and [3,](#page-5-0) we can explicitly compute:

Theorem 1.2. Let G be a connected reductive algebraic group and X a G-spherical variety with Q-Cartier K_X . Let L be a G-linearized ample line bundle on X. Then we can compute the $\delta^{(p)}$ -invariant and α -invariant with G–action as

$$
\delta_G^{(p)}(X, L) = \min_{v \in E} \frac{h_{\mathcal{C}}(v)}{\left(\frac{\int_{\Delta} P(x)(\langle x, v \rangle + l_s(v))^p dx}{\int_{\Delta} P(x) dx}\right)^{1/p}},
$$

$$
\alpha_G(X, L) = \min_{v \in E} \min_{m \in \Delta} \frac{h_{\mathcal{C}}(v)}{\langle m, v \rangle + l_s(v)},
$$

where E is a finite set inside V. The set E and the functions $h_{\mathcal{C}}, l_s, P(x)$ are explicit and will be introduced in Section [4.](#page-6-0) The polytope Δ is the polytope associated to a section s of L. When $p = 1$, we get $\delta_G^{(1)}(X, L) = \delta_G(X, L)$, the δ -invariant with G -action.

The key to prove the theorem is to compute the valuative invariant $S(L, v)$ for $v \in V$ (more generally $S^{(p)}(L, v)$), the *expected vanishing order of L along v*. These invariants are introduced in Section [2.](#page-3-0)

Then we study g−solitons on a Q−Fano variety in the sense of [\[BN14\]](#page-25-2). In order to do this, we need to have a natural torus action at hand. For a G−spherical Q−Fano variety X which is the completion of the spherical homogeneous space G/H , $T = (N_G(H)/H)^0$ is a torus, and it is actually the identity component of $\text{Aut}_G(X)$. Like above, there is a moment polytope Δ_T^+ associated to the T-action. Fix now a smooth positive function g on Δ_T^+ .

First we can compute the valuative invariant $S^{g}(v)$, the g-weighted expected vanishing order. Then we can get δ –invariant specially designed for g–solitons, namely $\delta_G^{\rm g}(X, -K_X)$ in Section 6.

Theorem 1.3. Let $(X, -K_X)$ be a Q-Fano spherical G-variety which is a completion of G/H . Let Δ be the polytope associated to the G-action. Let $T = (N_G(H)/H)^0$ and we have the corresponding polytope Δ_T^+ $_T^+$. Assume ${\rm g\ }$ is a strictly positive smooth function on Δ_T^+ T . Then

$$
\delta_G^{\mathsf{g}}(X, -K_X) = \min_{v \in E} \frac{h_{\mathcal{C}}(v)}{h_{\mathcal{C}}(v) + \langle \operatorname{bar}_{\mathrm{DH}}^{\mathsf{g}}(\Delta), v \rangle},
$$

where

$$
\text{bar}_{\text{DH}}^{g}(\Delta) = \frac{\int_{\Delta} g(\bar{x}) P(x) x dx}{\int_{\Delta} g(\bar{x}) P(x) dx}.
$$

Here E is a finite set inside V. The set E and the functions h_c , $P(x)$ are explicit and will be introduced in Section [4.](#page-6-0) The polytope Δ is the polytope associated to a preferable section u of $-K_X$ introduced in Section [4.](#page-6-0) The \bar{x} denotes the projection of $x \in \Delta$ on Δ_T^+ $^+_T.$

Then we can directly get a result about the G-equivariantly g-weighted Ding-semistability.

Corollary 1.1. Let $(X, -K_X)$ be a Q-Fano spherical G-variety which is a completion of G/H . Let Δ be the polytope associated to the G-action. Let $T = (N_G(H)/H)^0$ and we have the corresponding polytope Δ_T^+ $_T^+$. Assume g is a smooth positive function on Δ_T^+ \overline{T} . Then the following statements are equivalent:

- 1. The g-weighted barycenter $bar_{DH}^g(\Delta)$ is in the dual cone of $-\mathcal{V}_{\mathbb{R}}$.
- 2. The pair $(X, -K_X)$ is G-equivariantly g-weighted Ding-semistable.

With extra work and using the invariant β^g , which is the generalization of the β −invariant defined in [\[Fuj19,](#page-26-2) [Li17\]](#page-27-3), we have the following theorem on the existence of g−solitons on a Q−Fano spherical G−variety.

Theorem 1.4. Let $(X, -K_X)$ be a Q-Fano spherical G-variety which is a completion of G/H . Let Δ be the polytope associated to the G-action. Let $T = (N_G(H)/H)^0$ and we have the corresponding polytope Δ_T^+ $_T^+$. Assume g is a smooth positive function on Δ_T^+ \overline{T} . Then the following three statements are equivalent:

- 1. The g-weighted barycenter $bar_{BH}^{g}(\Delta)$ is in the relative interior of the dual cone of $-\mathcal{V}_\mathbb{R}$.
- 2. The pair $(X, -K_X)$ is G-equivariantly g-weighted Ding-polystable for special test configurations.
- 3. There exists a g−soliton on X.

After completing our article, we got informed of the recent work of Li-Li-Wang [\[LLW22\]](#page-27-6), where they also prove our Theorem [1.4.](#page-3-1) Our approach is based on the valuative theory, thus the β −invariant seems to be the most natural tool to use. Li-Li-Wang [\[LLW22\]](#page-27-6) tackles the problem using test configurations. Of course, the two perspectives coincide on special test configurations. Notice that a theory of K−stability for g−solitons is also developed in [\[LLW22\]](#page-27-6), whereas we only discuss Ding-stability based on [\[Li21,](#page-27-7) [HL23\]](#page-26-9).

With the above theorem and a special choice of the function g, we obtain the following application.

Theorem 1.5. Let X be a smooth Fano horospherical variety. Then the corresponding cone $(K_X)^\times$ admits a Calabi-Yau cone metric, given by a Sasaki-Einstein structure on a unit circle inside K_X with respect to some hermitian metric on K_X .

Acknowledgement. The author would like to express his gratitude to his thesis supervisors Julien Keller and Vestislav Apostolov for valuable suggestions. The author also benefits a lot from discussions with Thibaut Delcroix.

2 Invariants for filtrations

Let X be complex normal projective variety with \mathbb{Q} −Cartier K_X . We say that a prime divisor E is over X if there is a birational proper morphism $\mu: Y \to X$ with normal Y such that E is a prime divisor on Y. For such a E , the log discrepancy function is given by $A_X(E) = \text{ord}_E(K_{Y/X}) + 1$, where $K_{Y/X}$ is the relative canonical divisor. We say X is klt or has klt singularities if $A_X(E) > 0$ for all prime divisors E over X.

Let X be a complex normal projective variety with klt singularities. Let L be an ample line bundle on X. Let $d_m = \dim H^0(X, mL)$. Without loss of generality, up to a multiple of L, we can assume that $d_m \neq 0$ for any $m \in \mathbb{N}$. Let $R = \bigoplus_{m \geq 0} R_m$, where $R_m = H^0(X, mL)$.

By saying that F is a filtration of the ring R, we mean that for any $\lambda \in \mathbb{R}_{\geq 0}$ and $m \in \mathbb{N}$, there is a vector subspace $\mathcal{F}^{\lambda}R_m \subset R_m$ satisfying the properties:

- 1. $\mathcal{F}^{\lambda}R_m \subset \mathcal{F}^{\lambda'}R_m$ for $\lambda \geq \lambda'$; 2. $\mathcal{F}^{\lambda}R_m = \bigcap_{\lambda' < \lambda} \mathcal{F}^{\lambda'}R_m;$ 3. $\mathcal{F}^{\lambda_1}R_{m_1}\cdot \mathcal{F}^{\lambda_2}R_{m_2}\subset \mathcal{F}^{\lambda_1+\lambda_2}R_{m_1+m_2};$
- 4. $\mathcal{F}^0 R_m = R_m$ and $\mathcal{F}^\lambda R_m = 0$ for $\lambda >> 0$.

The jumping numbers of the filtration $\mathcal F$ are:

$$
0 \le a_{m,1}(\mathcal{F}) \le a_{m,2}(\mathcal{F}) \le \cdots \le a_{m,d_m}(\mathcal{F}),
$$

where $a_{m,j}(\mathcal{F}) := \inf \{ \lambda \in \mathbb{R}_{\geq 0} | \operatorname{codim} \mathcal{F}^{\lambda} R_m \geq j \}.$

By the second condition in the definition of filtrations, we see that $\mathcal{F}^{\lambda_1}R_m = \mathcal{F}^{\lambda_2}R_m$ for $\lambda_1, \lambda_2 \in (a_{m,i}(\mathcal{F}), a_{m,i+1}(\mathcal{F})]$ when $a_{m,i}(\mathcal{F}) \neq a_{m,i+1}(\mathcal{F})$. For $p \in [1, +\infty)$, the following invariant is defined in [\[Zha22\]](#page-28-3):

$$
S_m^{(p)}(L,\mathcal{F}):=\frac{1}{d_m}\sum_{j=1}^{d_m}\left(\frac{a_{m,j}}{m}\right)^p.
$$

With a R-valued valuation v on X, we can define a filtration \mathcal{F}_v by setting $\mathcal{F}_v^{\lambda}R_m :=$ $\{s \in R_m | v(s) \geq \lambda\}$ for $\lambda \geq 0$ and $m \in \mathbb{N}$. In this case, we write $S_m^{(p)}(L, v)$ instead of $S_m^{(p)}(L, \mathcal{F}_v)$. Especially, a prime divisor F over X gives a valuation ord_F which then leads to a filtration. In this case, as in [\[Zha22\]](#page-28-3), we write $S_m^{(p)}(L, F)$ instead of $S_m^{(p)}(L, \mathcal{F}_{\text{ord}_F})$. In this case, $S_m^{(p)}(L, F)$ is called the p-th moment of the expected vanishing order of L along F at level m. A little bit more generally, valuations in the form of $c \cdot \text{ord}_F$ with $c > 0$ are called divisorial. They similarly provide a filtration.

Similarly we write (see for example [\[BJ20\]](#page-25-1))

$$
T_m(L, \mathcal{F}) := \frac{a_{m,d_m}}{m}.
$$

The limit $T(L, \mathcal{F}) := \lim_{m \to +\infty} T_m(L, \mathcal{F})$ always exists in $[0, +\infty]$ (see [\[BJ20,](#page-25-1) section 2.4]). We say that F is linearly bounded if $T(L, \mathcal{F}) < +\infty$. If \mathcal{F}_v is linearly bounded for some valuation v , we usually say that v is of linear growth.

We recall the following definition of [\[Zha22\]](#page-28-3):

Definition 2.1. For a linearly bounded filtration \mathcal{F} , we define

$$
S^{(p)}(L, \mathcal{F}) := \lim_{m \to +\infty} S_m^{(p)}(L, \mathcal{F}).
$$

Especially, the above limit exists and is a finite number([\[BJ20,](#page-25-1) [Zha22\]](#page-28-3)).

Since X is a complex normal projective variety with klt singularities, as explained in [\[BdFFU15\]](#page-25-3), we can define the log discrepancy function for general valuations

$$
A_X: \text{Val}_X \to [0, +\infty]
$$

where Val_X is the set of all R-valued valuations on X. This function is lower semi-continuous and satisfies $A_X(tv) = tA_X(v)$ for $\lambda \in \mathbb{R}_{\geq 0}$. What is also important is that $A_X(v) = 0$ if and only if v is the trivial valuation. The next lemma is shown in $[BJ20]$.

Lemma 2.1. Any $v \in Val_X$ satisfying $A_X(v) < +\infty$ has linear growth. Especially this is the case for a divisorial valuation.

We define as in [\[Zha22\]](#page-28-3) the $\delta^{(p)}$ -invariant:

Definition 2.2. For $p \in [1, +\infty)$, we set

$$
\delta^{(p)}(L) := \inf_{v \in \text{DivVal}_X^*} \frac{A_X(v)}{S^{(p)}(L, v)^{1/p}},
$$

where the first v runs through the set of non-trivial divisorial valuations over X , denoted by Div Val_X^* .

The classical α -invariant looks very similar:

Definition 2.3. We define

$$
\alpha(L) := \inf_{v \in \text{DivVal}_X^*} \frac{A_X(v)}{T(L, v)},
$$

where the first v runs through all non-trivial divisorial valuations over X .

Remark 2.1. It is shown that $[BJ20, Zha22]$ $[BJ20, Zha22]$:

$$
\delta^{(p)}(L) = \inf_{v \in \text{Val}_X^*} \frac{A_X(v)}{S^{(p)}(L, v)^{1/p}},
$$

$$
\alpha(L) = \inf_{v \in \text{Val}_X^*} \frac{A_X(v)}{T(L, v)},
$$

where v runs through non-trivial valuations with finite log discrepancy.

3 Group action

As in [\[Gol20\]](#page-26-8), we can take group actions into consideration. Let G be a complex connected algebraic group. Let us assume that there is a G -action on X and the ample line bundle L is G-linearized. In [\[Gol20\]](#page-26-8), the author only considers $G \subset Aut(X, L)$, but there is really no much difference in considering more general group actions. We can define:

Definition 3.1. Let X be a normal projective G -variety with klt singularities and L be a G-linearized ample line bundle on X. For $p \in [1, +\infty)$, we define the $\delta^{(p)}$ -invariant with G−action

$$
\delta_G^{(p)}(L) := \inf_{v \in \text{DivVal}_X^{G*}} \frac{A_X(v)}{S^{(p)}(L, v)^{1/p}},
$$

where v runs through all non-trivial G -invariant divisorial valuations over X .

The classical α_G -invariant has been introduced in the literature, see [\[Gol20\]](#page-26-8):

Definition 3.2. Let X be a normal projective G-variety with klt singularities and L be a G-linearized ample line bundle on X, the α -invariant with G-action is

$$
\alpha_G(L) := \inf_{v \in \text{DivVal}_X^{G*}} \frac{A_X(v)}{T(L, v)},
$$

where v runs through all non-trivial G -invariant divisorial valuations over X .

4 Valuative invariants for linearized line bundles on a spherical variety

In this section, we focus on a linearized line bundle on a spherical variety. The references we use on spherical varieties are mainly [\[Kno91,](#page-27-8) [Bri97b,](#page-26-10) [Tim11,](#page-27-9) [Per14\]](#page-27-10).

Let G be a connected reductive complex algebraic group, B be a Borel subgroup of G , T_{max} be a maximal torus of B.

Definition 4.1. A normal G−variety is spherical if it contains an open B−orbit.

Let X be a spherical G-variety, it must contain an open G-orbit. If we fix a point x in this orbit, we can write $G \cdot x \cong G/H$, where H is the closed subgroup of G which fixes x. We call G/H a spherical homogeneous space and X is a spherical embedding of G/H . We assume further that (X, x) is a complete spherical embedding of G/H . We have the spherical lattice:

 $M = \{ \chi \text{ a character of } B | b \cdot f_{\chi} = \chi(b) f_{\chi} \text{ for some } f_{\chi} \in \mathbb{C}(X)^* = \mathbb{C}(G/H)^* \text{ and all } b \in B \}$

This is a sub-lattice of the character group of B. Notice that f_{χ} is unique up to multiplication by \mathbb{C}^* . For convenience, we identify χ with f_{χ} in this paper when there is no possibility of confusion. So we can view χ as a rational function on G/H . Let $N = \text{Hom}_{\mathbb{Z}}(M, \mathbb{Z})$ be the dual lattice.

For a valuation $v \in Val_X$, we can define $\langle v, \chi \rangle = v(f_\chi)$. In this way we see that there is a map $\rho: \text{Val}_X \to N \otimes \mathbb{R}$. Let QVal_X^G be the set of G-invariant Q-valued valuations on X or equivalently on G/H . The map ρ identifies the set $Q_{\text{N}}^{\text{U}} \Omega_X^G$ with a polyhedral convex cone $\mathcal{V} \subset N \otimes \mathbb{Q}$ ([\[Bri97b,](#page-26-10) Section 4]). We always treat QVal_X^G as a subset of $N \otimes \mathbb{Q}$. We denote the closure of V inside $N \otimes \mathbb{R}$ as $\mathcal{V}_{\mathbb{R}}$.

Let $D(X)$ be the set of B-stable prime divisors in X. For a G-orbit Y in X, we let $D_Y(X) \subset D(X)$ be the set of B-stable prime divisors which contain Y. Let $G_Y(X) = \{D \in$ $D_Y(X)|D$ is G – invariant} and $\mathcal{F}_Y(X) = \{D \in D_Y(X)|D$ is not G – invariant}. Then we let $\mathcal{C}_Y(X)$ be the cone inside $N \otimes \mathbb{Q}$ generated by $\rho(\mathcal{F}_Y(X))$ and $\rho(G_Y(X))$. The pair $(\mathcal{C}_Y(X), \mathcal{F}_Y(X))$ is a so called colored cone. Let $(\mathcal{C}, \mathcal{F})$ be the set of all the colored cones $(\mathcal{C}_Y(X), \mathcal{F}_Y(X))$, where Y goes through all the G-orbits inside X, then $(\mathcal{C}, \mathcal{F})$ is called a colored fan. Since X is complete, we know that $\mathcal{V} \subset \bigcup_{(\mathcal{C}_Y(X), \mathcal{F}_Y(X)) \in (\mathcal{C}, \mathcal{F})} \mathcal{C}_Y(X) =: \mathcal{C}_X$.

Remark 4.1. We can see that every element in $QVal_X^G$ is divisorial. Actually for $v \in V$, after sufficiently subdividing the cones of C_X and taking out colors, we can always find a toroidal G-equivariant resolution Z of which v is on an extremal ray of some colored cone in the corresponding colored fan. Then v must be divisorial. On the other hand, every Ginvariant divisorial valuation is clearly proportional to some element in $\mathcal V$. As changing valuations by a positive constant doesn't modify ratios like $\frac{A_X(v)}{S(p)(I,v)}$ $\frac{A_X(v)}{S^{(p)}(L,v)^{1/p}}$. We can just think of $\mathcal V$ as DivVal_X. See more details in [\[Pas17,](#page-27-11) Remark 3.5] which is based on [\[Cox00\]](#page-26-11).

Now we can start the procedure of computing the valuative invariants. The method here is highly influenced by [\[BJ20,](#page-25-1) Section 7] on toric line bundles.

Now let L be an ample G-linearized line bundle on X. We assume that $s \in H^0(X, L)$ is a B-eigenvector with weight χ . We can write:

$$
div(s) = \sum_{D \in D(X)} n_D D.
$$

This divisor gives a piecewise linear function l_s on \mathcal{C}_X which is linear on each cone $\mathcal{C}_Y(X)$. We can explicitly describe the function as follows. For each G-orbit Y, We let $\chi_{s,Y}$ be an element in M such that $\langle \rho(D), \chi_{s,Y} \rangle = n_D$ for any $D \in D_Y(X)$. Then we let l_s be the function given by $\chi_{s,Y}$ on $\mathcal{C}_Y(X)$.

For $v \in \text{QVal}_X^G$, it is important to know what is $v(s)$.

Lemma 4.1. For $v \in \text{QVal}_X^G$, we have $v(s) = l_s(v)$.

Proof. We know that v is in the relative interior of a unique colored cone $(\mathcal{C}_Y(X), \mathcal{F}_Y(X))$. Then the center of v is Y ([\[Kno91,](#page-27-8) Theorem 2.5]). We consider the set $X_{Y,B} = \{x \in X | Y \subset Y\}$ Bx . This is an open affine subset of X. We know that Y is a spherical variety, so it must contain an open dense B-orbit. Especially, $Y \cap X_{Y,B} \neq \emptyset$. This shows that the valuation v has a center on $X_{Y,B}$. From this it is also clear that $D_Y(X) = \{D \in D(X) | D \cap X_{Y,B} \neq \emptyset\}.$ Now, we can write

$$
div(s)|_{X_{Y,B}} = \sum_{D \in D_Y(X)} n_D D,
$$

=
$$
\sum_{D \in D_Y(X)} \langle \chi_{s,Y}, \rho(v_D)D,
$$

=
$$
div(\chi_{s,Y})|_{X_{Y,B}}.
$$

Thus on $X_{Y,B}$, we have $s = \chi_{s,Y}e$, where e is a local trivialization of L. Thus $v(s)$ = $\langle \chi_{s,Y} , v \rangle$. This is the same as saying $v(s) = l_s(v)$. \Box

Below we need some classical results on linearized line bundles on a spherical varieties. A lot of them are mentioned in [\[Del23\]](#page-26-12) and we shall use the same notations.

There is a canonical polytope related to an ample G-linearized line bundle which is called the moment polytope:

Definition 4.2. Let $\mathcal{X}(B)$ be the character group of B. Let Δ_k^+ be the subset of $\mathcal{X}(B)$ such that

$$
H^0(X, kL) \cong \bigoplus_{\lambda \in \Delta_k^+} V_{\lambda},
$$

where V_{λ} corresponds to the simple G-module with highest weight λ . Notice that $H^{0}(X, kL)$ is multiplicity-free, meaning that each simple G-module can at most appear once in the decomposition. Then $\bigcup_{r\in\mathbb{N}^*}\frac{\Delta_k^+}{k}$ is a rational polytope inside $\mathcal{X}(B)\otimes\mathbb{Q}$ ([\[Bri97b\]](#page-26-10)). We define the moment polytope

$$
\Delta^+ = \overline{\bigcup_{r \in \mathbb{N}^*} \frac{\Delta_k^+}{k}},
$$

where the closure is taken in $\mathcal{X}(B) \otimes \mathbb{R}$.

The moment polytope is always contained in the closed positive Weyl chamber. There is another very useful polytope Δ in $M \otimes \mathbb{R}$ related to the section s:

$$
\Delta = \{ m \in M \otimes \mathbb{R} | \rho(D)(m) + n_D \ge 0 \text{ for all } D \in D(X) \}
$$

We have the relation $\Delta^+ = \Delta + \chi$ when we consider $M \otimes \mathbb{R}$ as a subspace of $\mathcal{X}(B) \otimes \mathbb{R}$.

We have an isomorphism of G -modules:

$$
H^0(X, kL) \cong \bigoplus_{m \in M \cap k\Delta} V_{k\chi+m}.
$$

Let R^+ be the positive root system of (G, B, T_{max}) and ρ be the half of the sum of positive roots. Then we have the Weyl formula

$$
dim(V_{\lambda}) = \prod_{\alpha \in R^{+}} \frac{\langle \lambda + \rho, \alpha \rangle}{\langle \rho, \alpha \rangle},
$$

where $\langle \beta, \alpha \rangle$ represents the pairing of β with the coroot corresponding to α .

Now we can talk about filtrations of $H^0(X, kL)$ induced by a valuation $v \in \text{Val}_X^G$. We denote again $H^0(X, kL)$ as R_k . Clearly because v is G-invariant, the set $\mathcal{F}^\lambda_v R_k = \{u \in \mathbb{R}^n : v \in \mathbb{R}^n\}$ $H^0(X, kL)|v(u) \geq \lambda$ is a G-module. Now $\mathcal{F}^{\lambda}_v R_k$ must be the direct sum of some simple G-module $V_{k\chi+m}$.

Look at each $V_{k\chi+m}$ individually. It contains $s^k\chi^m$, where χ^m is a B-eigenfunction associated to the lattice point m. Since $V_{k\chi+m}$ is a simple G-module, we can conclude that all the non-zero elements in $V_{k\chi+m}$ have the same value when paired with v, and that is $kl_s(v) + \langle m, v \rangle$.

Thus we must have

$$
\mathcal{F}_{v}^{\lambda} R_{k} \cong \bigoplus_{\substack{m \in M \cap k\Delta \\ \langle m, v \rangle + kl_{s}(v) \geq \lambda}} V_{k\chi+m}.
$$

We deduce the following

Lemma 4.2. For $v \in \text{QVal}_X^G$, we have

$$
S_k^{(p)}(L, v) = \frac{1}{\dim H^0(X, kL)} \sum_{m \in M \cap k\Delta} \dim V_{k\chi+m} \left(\frac{\langle m, v \rangle + k l_s(v)}{k}\right)^p,
$$

$$
= \frac{\sum_{m \in M \cap k\Delta} \left(\frac{\langle m, v \rangle + k l_s(v)}{k}\right)^p \prod_{\alpha \in R^+} \frac{\langle k \chi + m + \rho, \alpha \rangle}{\langle \rho, \alpha \rangle}}{\sum_{m \in M \cap k\Delta} \prod_{\alpha \in R^+} \frac{\langle k \chi + m + \rho, \alpha \rangle}{\langle \rho, \alpha \rangle}};
$$

$$
T_k(L, v) = \frac{\max_{m \in M \cap k\Delta} (\langle m, v \rangle + k l_s(v))}{k}.
$$

Taking limit, we get the following result.

Lemma 4.3. For $v \in \text{QVal}_X^G$, we have:

$$
S^{(p)}(L, v) = \frac{\int_{\Delta} P(x) (\langle x, v \rangle + l_s(v))^p dx}{\int_{\Delta} P(x) dx},
$$

$$
T(L, v) = \max_{m \in \Delta} (\langle m, v \rangle + l_s(v)),
$$

where $P(x) = \prod_{\alpha \in R_X^+}$ $\langle x+\chi,\alpha\rangle$ $\frac{\partial \phi + \chi,\alpha}{\partial \langle \rho,\alpha \rangle}$ and R_X^+ are the set of positive roots not orthogonal to $\Delta^+ = \emptyset$ $\chi + \Delta$.

Especially when p is 1, this provides

Lemma 4.4. Let we denote $\frac{\int_{\Delta} P(x) dx}{\int_{\Delta} P(x) dx}$ as $bar_{DH}(\Delta)$. For $v \in QVal_X^G$, we have

$$
S(L, v) = \frac{\int_{\Delta} P(x)(\langle x, v \rangle + l_s(v))dx}{\int_{\Delta} P(x)dx},
$$

$$
= \langle \operatorname{bar}_{DH}(\Delta), v \rangle + l_s(v),
$$

where $P(x) = \prod_{\alpha \in R_X^+}$ $(x+\chi,\alpha)$ $\frac{f(x+\chi,\alpha)}{(\rho,\alpha)}$ and R_X^+ are the set of positive roots not orthogonal to $\Delta^+=\chi$ $\chi + \Delta$.

From now on, we assume that K_X is a Q-Cartier divisor. This actually implies that X has klt singularities when X is spherical (see [\[Pas17,](#page-27-11) Proposition 5.6]).

From [\[GH15,](#page-26-13) Section 4] which is based upon [\[Bri97a\]](#page-25-4) (see also [\[Del20,](#page-26-7) Section 3.2]), there is a section u of $-K_X$ which is a B-eigenvector with weight $2\rho_P = \sum_{\alpha \in \Phi_{Ru(P)}} \alpha$ (see [\[Del20\]](#page-26-7) and [\[Bri89\]](#page-25-5)). Here P is the stabilizer of the open B-orbit, $R_u(P)$ is the unipotent radical of P, and then $\Phi_{R_u(P)}$ is the set of roots of $R_u(P)$. We also have

$$
div(u) = \sum_{D \text{ is } G-\text{stable}} D + \sum_{D \in D(G/H)} a_D D,
$$

where the notation $D(G/H)$ refers to the set of B-stable prime divisors of G/H , similar to the notation $D(X)$.

Remark 4.2. More precisely, up to some multiple, u is a section of some line bundle.

Similar to the construction of the piecewise-linear function l_s related to a section s, the assumption that K_X is a Q-Cartier divisor is equivalent to the assumption that there is a piecewise linear function $h_{\mathcal{C}} := l_u$ on $\mathcal{C}_X = \bigcup_{(\mathcal{C}_Y(X), \mathcal{F}_Y(X)) \in (\mathcal{C}, \mathcal{F})} \mathcal{C}_Y(X)$, linear on each cone $\mathcal{C}_Y(X)$ given by some $m \in M \otimes \mathbb{Q}$. It is a very important fact that this function $h_{\mathcal{C}}$ is exactly the log discrepancy function A_X when restricted on $\mathcal V$ (see [\[Pas17,](#page-27-11) Section 5]). Notice again that V is a subset of \mathcal{C}_X .

The polyhedral cone V is carved into finitely many smaller polyhedral cones $\mathcal{C}_Y(X) \cap \mathcal{V}$. Let's gather the primitive generators of all the extremal rays of all the cones $\mathcal{C}_Y(X) \cap \mathcal{V}$ in a finite set E . Then we have:

Theorem 4.1. Let G be a connected reductive algebraic group and X a spherical variety with \mathbb{Q} -Cartier K_X . Let L be a G-linearized ample line bundle on X. We have

$$
\delta_G^{(p)}(X, L) = \min_{v \in E} \frac{h_{\mathcal{C}}(v)}{\left(\frac{\int_{\Delta} P(x)(\langle x, v \rangle + l_s(v))^p dx}{\int_{\Delta} P(x) dx}\right)^{1/p}},
$$

$$
\alpha_G(X, L) = \min_{v \in E} \min_{m \in \Delta} \frac{h_{\mathcal{C}}(v)}{\langle m, v \rangle + l_s(v)}.
$$

Proof. By definition and considerations in remark [4.1,](#page-6-1) we have:

$$
\delta_G^{(p)}(L) = \inf_{v \in \mathcal{V} \setminus \{0\}} \frac{A_X(v)}{S^{(p)}(L, v)^{1/p}}
$$

The polyhedral cone V is carved into finitely many smaller polyhedral cones $\mathcal{C}_Y(X) \cap \mathcal{V}$. Let's say $\mathcal{C}_Y(X) \cap \mathcal{V}$ is generated by $\{v_1, \dots, v_n\}$. Then every element v in the cone can be written as $v = a_1 v_{i_1} + \cdots + a_k v_{i_k}$, where every $a_i > 0$. Then we see

$$
S^{(p)}(L, v)^{1/p} = \left(\frac{\int_{\Delta} P(x)(\langle x, v \rangle + l_s(v))^p dx}{\int_{\Delta} P(x) dx}\right)^{1/p},
$$

\n
$$
= \left(\frac{\int_{\Delta} P(x)(\sum_{j=1}^k a_j(\langle x, v_{i_j} \rangle + l_s(v_{i_j})))^p dx}{\int_{\Delta} P(x) dx}\right)^{1/p},
$$

\n
$$
\leq \sum_{j=1}^k a_j \left(\frac{\int_{\Delta} P(x)(\langle x, v_{i_j} \rangle + l_s(v_{i_j}))^p dx}{\int_{\Delta} P(x) dx}\right)^{1/p},
$$

\n
$$
= \sum_{j=1}^k a_j S^{(p)}(L, v_{i_j})^{1/p}.
$$

Thus we see that

$$
\frac{A_X(v)}{S^{(p)}(L,v)^{1/p}} \ge \min_{v_{ij}} \frac{h_{\mathcal{C}}(v_{i_j})}{S^{(p)}(L,v_{i_j})^{1/p}},
$$

since $h_{\rm C}$ is linear on the $\mathcal{C}_Y \cap \mathcal{V}$. Similarly:

$$
\frac{A_X(v)}{T(L,v)} \ge \min_{v_{i_j}} \frac{h_{\mathcal{C}}(v_{i_j})}{T(L,v_{i_j})}.
$$

 \Box

Especially, when p is 1, we get

Corollary 4.1. With the same setup as above, we have

$$
\delta_G(X, L) = \min_{v \in E} \frac{h_{\mathcal{C}}(v)}{\langle \operatorname{bar}_{DH}(\Delta), v \rangle + l_s(v)}.
$$

5 Examples

Several results in the literature can be seen as consequences of what we proved previously.

Example 5.1. For the Fano case and $L = -K_X$, as we mentioned above, we can take $s = u$. Then l_s coincides with h_c and $\chi = 2\rho_P$. As for $bar_{DH}(\Delta)$, we have

$$
\begin{split}\n\text{bar}_{DH}(\Delta) &= \frac{\int_{\Delta} \prod_{\alpha \in R_X^+} \frac{(x + \chi, \alpha)}{(\rho, \alpha)} x dx}{\int_{\Delta} \prod_{\alpha \in R_X^+} \frac{(x + \chi, \alpha)}{(\rho, \alpha)} dx}, \\
&= \frac{\int_{\Delta + \chi} \prod_{\alpha \in R_X^+} \frac{(x, \alpha)}{(\rho, \alpha)} (x - \chi) dx}{\int_{\Delta + \chi} \prod_{\alpha \in R_X^+} \frac{(x, \alpha)}{(\rho, \alpha)} dx}, \\
&= \frac{\int_{\Delta +} \prod_{\alpha \in R_X^+} \frac{(x, \alpha)}{(\rho, \alpha)} (x - 2\rho_P) dx}{\int_{\Delta +} \prod_{\alpha \in R_X^+} \frac{(x, \alpha)}{(\rho, \alpha)} dx} \\
&= \text{bar}_{DH}(\Delta^+) - 2\rho_P.\n\end{split}
$$

,

Thus for Fano case, we recover the result [\[Gol20,](#page-26-8) Propostion1.4]:

$$
\delta_G(X, -K_X) = \min_{v \in E} \frac{h_{\mathcal{C}}(v)}{h_{\mathcal{C}}(v) + \langle \operatorname{bar}_{DH}(\Delta^+) - 2\rho_P, v \rangle}.
$$

Remark 5.1. In [\[Gol20\]](#page-26-8), there is a constant V in front of the term $\langle \text{bar}_{DH}(\Delta^+) - 2\rho_P, v \rangle$. From our considerations, this constant is actually 1.

Example 5.2. In the papers [\[Del17\]](#page-26-6) and [\[Del20\]](#page-26-7), Delcroix studies group compactifications and compute α –invariant with group actions for ample line bundles. His formula for the anticanonical line bundle(which is positive) of the unique wonderful compactification of a semisimple adjoint group \tilde{G} (under changes of notations) is beautifully simple and illustrative

$$
\alpha_{\hat{G}\times\hat{G}}(X, -K_X) = \max\{c|2\rho \in (1-c)\Delta^+\}.
$$

Let's reconsider the formula and link it to our perspective. Let \hat{G} be a semisimple adjoint group. We assume the Borel subgroup \hat{B} and the maximal torus $\hat{T}_{max} \subset \hat{B}$ of \hat{G} are fixed. Thus we have also fixed positive roots, simple roots and the positive Weyl chamber C^+ of \hat{G} . The variety \hat{G} is considered as a $G = \hat{G} \times \hat{G}$ -variety by the action $(g, h) \cdot \underline{x} = g x h^{-1}$. The corresponding H is the diagonal of $\hat{G} \times \hat{G}$, so $\hat{G} \times \hat{G}/H \cong \hat{G}$ by identifying $\overline{(g,h)}$ and gh^{-1} . The corresponding Borel subgroup of G is $\hat{B} \times \hat{B}^-$. The corresponding maximal torus of G is $\hat{T}_{max} \times \hat{T}_{max}$. The spherical lattice of G/H can be identified with the character group of B denoted as $\mathcal{X}(\hat{B})$, by sending a character of $\hat{B} \times \hat{B}^-$ to the character of B^{oro}. With this identification, several objects, for example moment polytopes, can be identified with objects defined with B. From now on, with the identification of character groups as above, we work on \tilde{G} .

Let $\{\alpha_i | 1 \le i \le r\}$ be the set of simple roots, where r is the dimension of the torus \hat{T}_{max} . Let Q be the polytope constructed as the convex hull of the images of $2\rho + \sum_{i=1}^{r} \alpha_i$ under the action of the Weyl group. Then we have $\Delta^+ = Q \cap C^+$. We refer to [\[BK07\]](#page-25-6) and [\[AK05\]](#page-25-7).

Let's look at Δ^+ in this case more carefully. To each positive root β , there is a corresponding reflection s_β . This s_β moves any $x \in C^+$ in the $-\beta$ direction. Since simple roots generate positive roots in a nonnegative way, we see that the extremal rays coming from $2\rho + \sum_{i=1}^r \alpha_i$ are give by $-\alpha_i$. Thus the codimensional 1 faces containing $2\rho + \sum_{i=1}^r \alpha_i$ are generated by $\{\alpha_{i_1}, \cdots, \alpha_{i_{r-1}}\}$. In other words, each of the face is perpendicular to some v_i , where $\{v_i | 1 \le i \le r\}$ is the dual basis of α_i in $N \otimes \mathbb{R}$. Because of all the considerations,

$$
\Delta^{+} = \{x \in C^{+}|\langle 2\rho + \sum_{i=1}^{r} \alpha_i, v_i \rangle \ge \langle x, v_i \rangle \text{ for all } v_i\},
$$

=
$$
\{x \in C^{+}|\langle 2\rho, v_i \rangle + 1 \ge \langle x, v_i \rangle \text{ for all } v_i\}.
$$

From this we get:

$$
\alpha_G(X, -K_X) = \max\{c|2\rho \in (1-c)\Delta^+\},
$$

= 1 - \min\{\lambda|\lambda(\langle 2\rho, v_i \rangle + 1) \ge \langle 2\rho, v_i \rangle \text{ for all } v_i\},
= 1 - \max_i \left\{\frac{\langle 2\rho, v_i \rangle}{\langle 2\rho, v_i \rangle + 1}\right\},
= \min_i \left\{\frac{1}{1 + \langle 2\rho, v_i \rangle}\right\}.

Let's link this to our formula in Theorem [4.1.](#page-10-0) Wonderful compactifications enjoys two properties: they are both symmetric and toroidal. The former implies that the valuation cone $\mathcal V$ is exactly the dual of the negative Weyl chamber, that is to say, the cone generated by $\{-v_i|1 \leq i \leq r\}$. The latter implies that there is no color, that is to say every B-stable divisor on it which contains a G-orbit is actually G−invariant, see [\[Tim11,](#page-27-9) Section 26] or [\[Del20,](#page-26-7) Section 5.4.2].

In this case, the section u mentioned above has weight $2\rho_P = 2\rho$. And h_C is 1 on each $-v_i$ and extends linearly. Now we use Theorem [4.1:](#page-10-0)

$$
\alpha_G(X, -K_X) = \min_{i} \min_{m \in \Delta} \left\{ \frac{1}{\langle m, -v_i \rangle + 1} \right\},
$$

=
$$
\min_{i} \min_{m \in \Delta^+} \left\{ \frac{1}{\langle m - 2\rho, -v_i \rangle + 1} \right\}
$$

=
$$
\min_{i} \left\{ \frac{1}{\langle 2\rho, v_i \rangle + 1} \right\},
$$

,

since clearly $\langle m, -v_i \rangle \leq 0$ for $m \in \Delta^+ \subset C^+$.

Remark 5.2. Let's discuss $P(x)$ in the computation of $\delta_G^{(p)}(-K_X)$ in the previous example. Let's denote the set of positive roots of \hat{G} with respect to \hat{B} as Φ^+ . After identifying the spherical lattice of G/H with $\mathcal{X}(\hat{B})$, we should write $P(x) = \prod_{\alpha \in \Phi^+} \left(\frac{\langle x+2\rho, \alpha \rangle}{\langle \rho, \alpha \rangle} \right)$ $\frac{+2\rho,\alpha\rangle}{\langle\rho,\alpha\rangle}$ ² instead of $\prod_{\alpha\in \Phi^+}$ $\langle x+2\rho,\alpha\rangle$ $\frac{+2\rho,\alpha}{\langle\rho,\alpha\rangle}$, where ρ is the half sum of positive roots of \hat{G} with respect to \hat{B} . The square here really corresponds to the fact that we need to consider representations of G instead of \tilde{G} in the process of the last section.

Now let's look at a last and concrete example.

Example 5.3. Let G be $\text{PGL}_2(\mathbb{C}) \times \text{PGL}_2(\mathbb{C})$. Let H be the diagonal of $\text{PGL}_2(\mathbb{C}) \times \text{PGL}_2(\mathbb{C})$. Then $G/H \cong \text{PGL}_2(\mathbb{C})$ identifying $\overline{(a,b)}$ with ab^{-1} . Then the induced group action on $PGL_2(\mathbb{C})$ is $(g, h) \cdot x = gxh^{-1}$ for $g, h, x \in \text{PGL}_2(\mathbb{C})$. A usual choice of a Borel subgroup is $B = \{(b_1, b_2) \in G | b_1 \text{ upper-triangular}, b_2 \text{ lower-triangular}\}\.$ Let $T_{max} = \{(b_1, b_2) \in G | b_1 \text{ upper-triangular}\}\.$ $G|b_1$ and b_2 are diagonal}. Define:

$$
f\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}\right) = \frac{d^2}{ad - bc}.
$$

It is easy to check that f is a B -eigenvector, actually:

$$
\left(\left(\begin{bmatrix} \lambda_1 & \lambda_2 \\ 0 & \lambda_3 \end{bmatrix}, \begin{bmatrix} \beta_1 & 0 \\ \beta_2 & \beta_3 \end{bmatrix} \right) \cdot f \right) \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \right) = \frac{\lambda_1}{\lambda_3} \frac{\beta_3}{\beta_1} \frac{d^2}{ad - bc}.
$$

In this case, the spherical lattice of G/H is isomorphic to $\mathbb Z$ and is generated by the element ω given by $\omega\left(\begin{bmatrix} \lambda_1 & \lambda_2 \\ 0 & \lambda \end{bmatrix}\right)$ $0 \lambda_3$ $\begin{bmatrix} \beta_1 & 0 \\ 0 & 0 \end{bmatrix}$ $\left[\begin{matrix} \beta_1 & 0 \ \beta_2 & \beta_3 \end{matrix} \right] \bigg) = \frac{\lambda_1}{\lambda_3}$ λ_3 β_3 $\frac{\beta_3}{\beta_1}$. We can identify ω with $\omega_1\left(\begin{bmatrix} \lambda_1 & \lambda_2 \\ 0 & \lambda_2 \end{bmatrix}\right)$ $\begin{pmatrix} \lambda_1 & \lambda_2 \ 0 & \lambda_3 \end{pmatrix} \bigg) =$ λ_1 $\frac{\lambda_1}{\lambda_3}$. The character ω_1 generates the character group of the Borel subgroup of the first component of G. We identify ω or ω_1 with 1.

The wonderful compactification of $PGL_2(\mathbb{C})$ is $X = \mathbb{P}(M_{2\times 2}(\mathbb{C}))$. The only closed G-orbit of X is the prime divisor $D = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} | ad - bc = 0 \right\}$. The embedding X is now simple, meaning that there is only one closed \overline{G} -orbit, and complete. It is colorless simply because the only closed G-orbit is of codimension 1. Clearly $v_D(f) = -1$. The valuation cone is generated by $\rho(v_D)$ in a nonnegative way. This cone can be easily identified with the dual of the negative Weyl chamber of the first component of G.

Now let's look at the section u of the anticanonical bundle of X. Clearly $-K_X \cong \mathcal{O}(4)$. We claim that the section u is identified with the section $(ad - bc)d^2$ of $\mathcal{O}(4)$. We check now this fact.

Actually, the section $(ad-bc)d^2$, when written locally in the chart $\{d \neq 0\}$ and considered as a section of $-K_X$, is $(a - bc)\frac{\partial}{\partial a} \wedge \frac{\partial}{\partial b} \wedge \frac{\partial}{\partial c}$. We have:

$$
\begin{bmatrix} \lambda_1 & \lambda_2 \\ 0 & \lambda_3 \end{bmatrix} \begin{bmatrix} a & b \\ c & 1 \end{bmatrix} = \begin{bmatrix} \lambda_1 a + \lambda_2 c & \lambda_1 b + \lambda_2 \\ \lambda_3 c & \lambda_3 \end{bmatrix} = \begin{bmatrix} \frac{\lambda_1 a + \lambda_2 c}{\lambda_3} & \frac{\lambda_1 b + \lambda_2}{\lambda_3} \\ c & 1 \end{bmatrix}.
$$

Let $a' = \frac{\lambda_1 a + \lambda_2 c}{\lambda_3}, b' = \frac{\lambda_1 b + \lambda_2}{\lambda_3}, c' = c$. Then $a = \frac{\lambda_3 a' - \lambda_2 c'}{\lambda_1}, b = \frac{\lambda_3 b' - \lambda_2}{\lambda_1}, c = c'$. Then:

$$
\begin{bmatrix} \lambda_1 & \lambda_2 \\ 0 & \lambda_3 \end{bmatrix}_* \left((a - bc) \frac{\partial}{\partial a} \wedge \frac{\partial}{\partial b} \wedge \frac{\partial}{\partial c} \right) = \frac{\lambda_1}{\lambda_3} \left((a' - b'c') \frac{\partial}{\partial a'} \wedge \frac{\partial}{\partial b'} \wedge \frac{\partial}{\partial c'} \right).
$$

Similar computation holds if we let $\begin{bmatrix} \beta_1 & 0 \\ 0 & \beta \end{bmatrix}$ β_2 β_3 1 act on the right side of the section. This shows that $(ad - bc)d^2$, when considered as a section of $-K_X$, is a B-eigenvector with weight ω . So it must be the preferable section u .

It is not hard to see that the moment polytope Δ^+ is [0, 2]. Actually there are only two B-stable divisors ,namely D and the divisor given by $\{d = 0\}$. Then we can compute the polytope associated to the section u and then get the moment polytope. Otherwise we can also conclude that by considering dimensions. We know that $H^0(X, \mathcal{O}(4))$ is of dimension 35. On the other hand, $\dim(V_i) = (2i+1)^2$ for $i = 0, 1, 2$. At the same time, we see that the polytope Δ associated to u is [-1, 1].

Now using the formula from Theorem [4.1:](#page-10-0)

$$
\alpha_G(X, -K_X) = \min_{m \in [-1,1]} \left\{ \frac{1}{\langle m, -1 \rangle + 1} \right\} = \frac{1}{2}.
$$

This was also shown in [\[Del17\]](#page-26-6). For the $\delta_G^{(p)}$ -invariant, We obtain

$$
\delta_G^{(p)}(X, -K_X) = \frac{1}{\left(\frac{\int_{-1}^1 (x+1,1)^2((x,-1)+1)^p dx}{\int_{-1}^1 (x+1,1)^2 dx}\right)^{1/p}} = \frac{1}{2} \left(\frac{(p+1)(p+2)(p+3)}{6}\right)^{1/p}
$$

.

6 Valuative invariants for g-solitons

A Q−Fano variety is a normal projective Q-Gorenstein Fano variety with klt singularities. Assume that X is a \mathbb{Q} –Fano variety of dimension m. Assume also that there is a torus action on X. We denote the algebraic torus as T and T^c be the compact real torus inside T. We introduce g−solitons as in [\[HL23\]](#page-26-9).

Let h be a smooth positive hermitian metric on $-K_X$ with positive curvature ω representing $c_1(X)$. We can define a global measure on X as (see [\[BBE](#page-25-8)+19])

$$
d\mu_0 = |s|_{h}^{\frac{2}{r}} ((\sqrt{-1})^{rm^2} s^* \wedge \overline{s^*})^{\frac{1}{r}},
$$

where r is sufficiently divisible so that $-rK_X$ is a line bundle, s is a nowhere-zero local section of $-rK_X$, s^* is the dual of s .

For any Kähler form $\omega_{\varphi} = \omega + \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \varphi$ inside $c_1(X)$, there is an associated moment map:

$$
m_{\varphi}: X \to \text{Lie}(T^c) \cong M_T \otimes \mathbb{R},
$$

where Lie(T^c) is the lie algebra of T^c and M_T is the character lattice of T.

The image of m_{φ} is the moment polytope Δ_T^+ , which is independent of the choice of ω_{φ} . Let g be a strictly positive smooth function on Δ_T^+ . The g-soliton equation is the following non-linear PDE in φ (we refer to [\[BN14,](#page-25-2) [HL23,](#page-26-9) [Li21\]](#page-27-7)):

$$
g \circ m_{\varphi} \frac{\omega_{\varphi}^n}{n!} = e^{-\varphi} d\mu_0.
$$

With a careful choice of g, we can get the equation of Mabuchi soliton

In [\[RTZ21\]](#page-27-12), the authors introduce δ invariants for g-solitons. We recall some of their definitions. The setup is as above. From the torus action, we have a decomposition $H^0(X, -rK_X) = \bigoplus_{\alpha} R_{r,\alpha}$, here α is a character of the torus and $R_{r,\alpha}$ represents the subspace

of eigenvalue α . Denote $\Delta_{T,r}^+$ as the set of α such that $R_{r,\alpha}$ is non-trivial. We denote the character lattice of T as M_T . Then let

$$
\Delta_T^+ = \overline{\cup_{r \in \mathbb{N}^*} \frac{\Delta_{T,r}}{r}}
$$

which is a polytope in $M_T \otimes \mathbb{R}$. By [\[Bri87\]](#page-25-9), this polytope is just the moment polytope in the symplectic sense introduced above, justifying the same notation for the two. Again let g be a strictly positive smooth function on Δ_T^+ .

Let d_r be the dimension of $H^0(X, -rK_X)$ and $d_{r,\alpha}$ be the dimension of $R_{r,\alpha}$. Let $\bar{g}_r = \frac{1}{d_s}$ $\frac{1}{d_r} \sum_{\alpha \in \Delta_{T,r}^+} g(\frac{\alpha}{r}) d_{r,\alpha}$. Let F be a linearly bounded T-invariant filtration of $R =$ $\bigoplus_{r\geq 0} H^0(X, -rK_X)$ on X, meaning that every $\mathcal{F}^{\lambda}R_m$ is T-invariant. Let $a_{r,\alpha,j}$ be the jth jumping number of the filtration on $R_{r,\alpha}$. Then

$$
S_r^{\mathsf{g}}(\mathcal{F}) := \frac{1}{r d_r \bar{\mathsf{g}}_r} \sum_{\alpha \in \Delta_{T,r}^+} \sum_{j \ge 1} \mathsf{g} \left(\frac{\alpha}{r}\right) a_{r,\alpha,j},
$$

$$
S^{\mathsf{g}}(\mathcal{F}) := \lim_{r \to \infty} S_m^{\mathsf{g}}(\mathcal{F}).
$$

Now we define

.

$$
\delta^{\mathsf{g}}(X,-K_X):=\inf_{v\in{\rm DivVal}^{T*}_X}\frac{A_X(v)}{S^{\mathsf{g}}(v)}
$$

At the moment, X is a T-variety. Now we assume that it is also a G -variety for some connected reductive G , and the G-action commutes with the T-action. In this way, X is a $T \times G$ -variety.

Definition 6.1. Let X be a $\mathbb{Q}-$ Fano variety. Assume that there is a $T \times G$ -action on X, where G is a connected complex reductive algebraic group, and T is a algebraic torus. Let g be a strictly positive smooth function on Δ_T^+ . Then we introduce

$$
\delta_G^{\mathsf{g}}(X, -K_X) := \inf_{v \in \text{DivVal}_X^{T \times G*}} \frac{A_X(v)}{S^{\mathsf{g}}(v)},
$$

where $DivVal_X^{T \times G*}$ denotes the set of non-trivial $T \times G$ -invariant divisorial valuations over X .

We should point out that, in the Fano case, δ -invariant is closely related to β -invariant. We recall the definition of the β -invariant here.

Definition 6.2. Let X be a \mathbb{Q} –Fano variety. Assume that there is a T-action on X, where T is a algebraic torus. Let g be a strictly positive smooth function on Δ_T^+ . Then

$$
\beta^{\mathsf{g}}(v) := A_X(v) - S^{\mathsf{g}}(v),
$$

where v is a T -invariant divisorial valuation.

When we study g−solitons on a spherical variety, we need to have a natural torus action. We know that the G-equivariant automorphism group of a spherical homogeneous space G/H is $N_G(H)/H$, where $N_G(H)$ is the normalizer of H inside G. The group action of $N_G(H)/H$ on G/H is $pH \cdot gH = gp^{-1}H$. The group $N_G(H)/H$ is diagonalizable ([\[Bri97b,](#page-26-10) Theorem 4.3]). We elaborate on this fact a little bit more. For a B-eigenfunction $f_{\chi} \in \mathbb{C}(G/H)$ with eigenvalue χ , because the $N_G(H)/H$ -action commutes with the G-action, we see that $\gamma \cdot f_\chi$ is another B–eigenvector with eigenvalue χ . Since $\mathbb{C}(G/H)$ is multiplicity-free, thus there is a non-zero complex number $\theta_{\chi}(\gamma)$ such that $\gamma \cdot f_{\chi} = \theta_{\chi}(\gamma) f_{\chi}$. In this way, there is a group homomorphism:

$$
\theta: N_G(H)/H \to \text{Hom}(M, \mathbb{C}^*),
$$

where M is the spherical lattice. It turns out this homomorphism is injective and the image is Hom $(M/\langle\Sigma\rangle, \mathbb{C}^*)$. See [\[Bri97b\]](#page-26-10) for the explication of Σ .

The neutral component of $N_G(H)/H$ is an algebraic torus, and it is also the neutral component of the group of $G-$ equivariant automorphisms of X ([\[Del20,](#page-26-7) Section 3.1.3]). The neutral component $(N_G(H)/H)^0$ is the algebraic torus that we are going to use. Notice that $(N_G(H)/H)^0$ can be identified with $\text{Hom}(M/\langle \Sigma \rangle, \mathbb{C}^*)$, where $M/\langle \Sigma \rangle$ is $M/\langle \Sigma \rangle$ quotient by its torsion part.

The situation here is simpler than general cases thanks to the following lemma.

Lemma 6.1. Assume $v \in \text{QVal}_{X}^{G}$. Then v is also $N_G(H)/H$ -invariant.

Proof. The two groups G and $N_G(H)/H$ commute with each other when they act on G/H , then they commute when they act on the space of rational functions on G/H , then it follows that they commute when they act on the space of valuations on G/H . Then we see easily that, if v is G-invariant, $\gamma \cdot v$ is also G-invariant for $\gamma \in N_G(H)/H$. We just need to show that $\gamma \cdot v = v$.

We just need to show that for any B-eigenvector $f_{\chi} \in \mathbb{C}(G/H)$ with eigenvalue χ , we have $v(f_\chi) = (\gamma \cdot v)(f)$. This follows from $(\gamma \cdot v)(f_\chi) = v(\gamma^{-1} \cdot f_\chi) = v(\theta_\chi(\gamma^{-1})f_\chi) = v(f_\chi)$.

From now on, we use T to denote $(N_G(H)/H)^0$. Because of the above lemma, DivVal $^{T \times G}_{X}$ is our valuation cone V , and $DivVal_X^{T \times G*} = V \setminus \{0\}.$

Let X be a Q-Fano spherical variety. As before, we have

$$
H^{0}(X, -rK_{X}) \cong \bigoplus_{m \in M \cap r\Delta} V_{2r\rho_{P}+m}.
$$

Here again $2\rho_P$ is the weight of the preferable B−eigenvector u, and Δ is the polytope associated to u.

Because G and T commute with each other when they act on X , from multiplicityfreeness we see easily that u is also a T−eigenvector. Each $V_{2r\rho_P+m}$ is generated by the B-eigenvector $u^r \chi^m$ with eigenvalue $2r\rho_P + m$. Then each element in $V_{2r\rho_P+m}$ has the same T-eigenvalue as $u^r \chi^m$ because $u^r \chi^m$ generates $V_{2r\rho_P+m}$ as a G -module and G commutes with T .

The section u actually has the trivial T −eigenvalue. This can be seen by the construction before the proposition 4.1 in $\left[\text{GH15} \right]$ where s stands for our eigenvectoru. It is given at each point as the wedge of some holomorphic vector fields induced by one-parameter subgroups of G. Because T and the one-parameter subgroups of G commute, the holomorphic vector fields are all preserved by the action of T. Then u is preserved by the action of T. Thus $u^r \chi^m$ has the T-eigenvalue \bar{m} , where \bar{m} represents the projection of m on $M/\langle\Sigma'\rangle$.

We have two polytopes with respect to the action of G , namely the moment polytope Δ^+ , and the polytope Δ related to u. They are related by $\Delta^+ = \Delta + 2\rho_P$. Because u has trivial T-eigenvalue, we just have one polytope with respect to the action of T , namely the moment polytope Δ_T^+ . The projection from M to $M/\langle\Sigma'\rangle$ provides us with a projection from Δ to Δ_T^+ . We assume that we have a strictly positive smooth function g on Δ_T^+ .

Now for $v \in V \setminus \{0\}$, we know that v is also T-invariant. So we can consider S_k^g $_{k}^{\mathsf{g}}(v)$. Using the same argument in Section 4, we see,

$$
S_k^{\mathbf{g}}(v) = \frac{\sum_{m \in M \cap k\Delta} \mathbf{g}(\frac{\bar{m}}{k}) \prod_{\alpha \in R^+} \frac{\langle \alpha, 2k\rho_P + m + \rho \rangle}{\langle \alpha, \rho \rangle} \frac{\langle m, v \rangle + k h_{\mathcal{C}}(v)}{k}}{\sum_{m \in M \cap k\Delta} \mathbf{g}(\frac{\bar{m}}{k}) \prod_{\alpha \in R^+} \frac{\langle \alpha, 2k\rho_P + m + \rho \rangle}{\langle \alpha, \rho \rangle}}
$$

.

Taking limit, we get:

$$
S^{g}(v) = \frac{\int_{\Delta} g(\bar{x}) P(x) (\langle x, v \rangle + h_{\mathcal{C}}(v)) dx}{\int_{\Delta} g(\bar{x}) P(x) dx},
$$

$$
= h_{\mathcal{C}}(v) + \langle \text{bar}_{DH}^{g}(\Delta), v \rangle,
$$

where

$$
\text{bar}_{\text{DH}}^{g}(\Delta) = \frac{\int_{\Delta} g(\bar{x}) P(x) x dx}{\int_{\Delta} g(\bar{x}) P(x) dx}.
$$

Then eventually:

Theorem 6.1. Let $(X, -K_X)$ be a Q–Fano spherical G-variety which is a completion of G/H . Let Δ be the polytope associated to the G-action. Let $T = (N_G(H)/H)^0$ and we have the corresponding polytope Δ_T^+ $_T^+$. Assume ${\rm g\,}$ is a strictly positive smooth function on Δ_T^+ T . Then

$$
\delta_G^{\mathrm{g}}(X, -K_X) = \min_{v \in E} \frac{h_{\mathcal{C}}(v)}{h_{\mathcal{C}}(v) + \langle \mathrm{bar}_{\mathrm{DH}}^{\mathrm{g}}(\Delta), v \rangle}.
$$

7 Ding-stability For g-solitons

In this section we are going to study Ding-stability for g−solitons on a Q-Fano spherical variety.

First we recall some definitions from [\[HL23,](#page-26-9) [Li21\]](#page-27-7).

Definition 7.1. Let $(X, -K_X)$ be a Q-Fano variety. Assume that there is a $T \times G$ -action on X, where T is a complex algebraic torus and G is a connected reductive algebraic group. A T × G-equivariant test configuration $(\mathcal{X}, \mathcal{L})$ of $(X, -K_X)$ is the following data:

• X is a normal variety with a $\mathbb{C}^* \times T \times G$ –action on it. We have a \mathbb{C}^* -equivariant flat proper morphism $\pi : \mathcal{X} \to \mathbb{C}$. The $T \times G$ -action only acts on fibers of π .

- $\mathcal L$ is π -ample Q-line bundle on $\mathcal X$. There is a $\mathbb C^* \times T \times G$ -action on $\mathcal L$ which is a lift of the $\mathbb{C}^* \times T \times G$ -action on X.
- Over \mathbb{C}^* , we have a $\mathbb{C}^* \times T \times G$ -equivariant isomorphism $(\mathcal{X}, \mathcal{L}) \times_{\mathbb{C}^*} \mathbb{C} \cong \mathbb{C}^* \times (X, -K_X)$, where on the right side \mathbb{C}^* only acts on \mathbb{C}^* and $T \times G$ only acts on $(X, -K_X)$.

We can twist a $T \times G$ –equivariant test configuration:

- Let $\mathbb T$ be the connected part of the center of G. Let $N_{\mathbb T}$ be the group of one-parameter subgroups of \mathbb{T} . For any $\xi \in N_{\mathbb{T}}$, we can define the twist $(\mathcal{X}_{\xi}, \mathcal{L}_{\xi})$ of a $T \times G$ -equivariant test configuration $(\mathcal{X}, \mathcal{L})$ by changing the original \mathbb{C}^* -action on $(\mathcal{X}, \mathcal{L})$ to a new one which is the composition of the original one with the \mathbb{C}^* -action on $(\mathcal{X}, \mathcal{L})$ induced by ξ. The pair $(\mathcal{X}_{\xi}, \mathcal{L}_{\xi})$ is a new $T \times G$ -equivariant test configuration.
- For $\xi \in N_{\mathbb{T}} \otimes \mathbb{Q}$, assume that $k\xi \in N_{\mathbb{T}}$. We first use the base change $z \mapsto z^k$ to define a $T \times G$ -equivariant test configuration $(\mathcal{X}_k, \mathcal{L}_k)$. There is an induced \mathbb{C}^* -action on $(\mathcal{X}_k, \mathcal{L}_k)$. Notice that for $\lambda \in \mathbb{C}^*$, the λ -action on $(\mathcal{X}_k, \mathcal{L}_k)$ actually commutes with λ^k –action on $(\mathcal{X}, \mathcal{L})$ through the pullback morphism between them. Now the oneparameter subgroup corresponding to $k\xi$ gives a \mathbb{C}^* -action on $(\mathcal{X}_k, \mathcal{L}_k)$ through the T–action on it. Then we twist the \mathbb{C}^* –action on $(\mathcal{X}_k, \mathcal{L}_k)$ with this action. The test configuration after twist is thought of as the k times of $(\mathcal{X}_{\xi}, \mathcal{L}_{\xi})$.

Taking G to be trivial, we get the definition of a T-equivariant test configuration. Taking T and G to be trivial, we get the definition of a test configuration. Some important classes of test configurations are:

- A test configuration is a product if there is a \mathbb{C}^* -action on $(X, -K_X)$ and that $(\mathcal{X}, \mathcal{L}) \cong$ $\mathbb{C} \times (X, -K_X)$ in a \mathbb{C}^* – equivariant way, where the action on the right hand side is given by the multiplication of \mathbb{C}^* on \mathbb{C} and the \mathbb{C}^* -action on $(X, -K_X)$.
- A test configuration $(\mathcal{X}, \mathcal{L})$ is special if $\mathcal{L} \cong -K_{\mathcal{X}/\mathbb{C}}$ and \mathcal{X}_0 is \mathbb{Q} -Fano.

When we have a test configuration $(\mathcal{X}, \mathcal{L})$, we have a corresponding filtration $\mathcal{F}(\mathcal{X}, \mathcal{L})$ of $R = \bigoplus_{r \in \mathbb{N}} H^0(X, -rK_X)$. This is for example described in [\[BHJ17\]](#page-25-10).

For a \mathbb{Q} -Fano variety with a T-action, we assume as before that there is a strictly positive smooth function g on Δ_T^+ .

Definition 7.2. Given a T–equivariant test configuration $(\mathcal{X}, \mathcal{L})$, we have the following non-Archimedean functionals:

- $\mathbf{L}^{\text{NA}}(\mathcal{X}, \mathcal{L}) = \text{lct}(\mathcal{X}, -(K_{\mathcal{X}} + \mathcal{L}); \mathcal{X}_0) 1,$
- $\mathbf{E}_{\text{g}}^{\text{NA}}(\mathcal{X}, \mathcal{L}) = S^{\text{g}}(\mathcal{F}(\mathcal{X}, \mathcal{L})),$
- \bullet $\mathbf{D}_{\mathrm{g}}^{\mathrm{NA}} = \mathbf{L}^{NA} \mathbf{E}_{\mathrm{g}}^{\mathrm{NA}}$ g ,
- \bullet $\, \Lambda_{\scriptscriptstyle \sigma}^{\rm NA}$ $g_{\text{g}}^{\text{NA}}(\mathcal{X}, \mathcal{L}) = \mathbf{\Lambda}^{\text{NA}}(\mathcal{X}, \mathcal{L}) = T(\mathcal{F}(\mathcal{X}, \mathcal{L})),$
- $J_g^{NA} = \Lambda_g^{NA} E_g^{NA},$

• $J_{g,T}^{NA}(\mathcal{X}, \mathcal{L}) = \inf_{\xi \in N_T \otimes \mathbb{Q}} J_g^{NA}(\mathcal{X}_{\xi}, \mathcal{L}_{\xi}).$

The functional \mathbf{E}_{g}^{NA} ^{NA}(\mathcal{X}, \mathcal{L}) is called the g-weighted non-Archimedean Monge-Ampère energy functional. The functional D_g^{NA} is called the g-weighted Ding functional.

When $g = 1$, we omit g so it is compatible with notations in literature.

Remark 7.1. What is going to be important to us is that if $(\mathcal{X}, -\mathcal{K}_{\mathcal{X}})$ is a special T−equivariant test configuration, we have $\mathbf{D}_{g}^{NA}(\mathcal{X}, -\mathcal{K}_{\mathcal{X}}) = \beta^{g}(v) = A_{X}(v) - S^{g}(v)$, where v is the restriction of $v_{\mathcal{X}_0}$ to $\mathbb{C}(X)$. This is for example shown in the proof of [\[Li21,](#page-27-7) Theorem 1.16].

Now we can proceed to definitions of stability:

Definition 7.3. Let $(X, -K_X)$ be a Q-Fano variety. Assume that there is a $T \times G$ -action on X , where T is a complex algebraic torus and G is a connected reductive algebraic group. We assume that there is a strictly positive smooth function g on Δ_T^+ .

- We say that X is G-equivariantly g-weighted Ding-semistable if $\mathbf{D}_{g}^{NA}(\mathcal{X}, \mathcal{L}) \geq 0$ for every $T \times G$ –equivariant test configuration.
- We say that X is G -equivariantly g-weighted Ding-polystable if it is G -equivariantly g–weighted Ding-semistable, and we have equality if and only if $(\mathcal{X}, \mathcal{L})$ is a product test configuration.
- We say that X is G-equivariantly g-weighted uniformly Ding-stable if there exists γ > 0 such that $\mathbf{D}_{g}^{NA}(X, L) \geq \gamma \cdot \mathbf{J}_{g}^{NA}$ $g_{\rm g}^{\rm NA}(\mathcal{X}, \mathcal{L})$ for every $T \times G$ -equivariant test configuration.
- We say that X is G-uniformly g-weighted Ding-stable if there exists a constant $\gamma > 0$ such that $\mathbf{D}_{g}^{NA}(X, L) \geq \gamma \cdot \mathbf{J}_{g, \mathbb{T}}^{NA}(\mathcal{X}, \mathcal{L})$ for every $T \times G$ -equivariant test configuration.

Remark 7.2. Notice that J_{ϱ}^{NA} $_{\rm g}^{\rm NA}$ and $\bm J^{\rm NA}$ are bounded by each other. See for example [\[Li21,](#page-27-7) Lemma 1.4]. So we can also define G−equivariantly g−weighted uniformly Ding-stable with J^{NA} and G -uniformly g-weighted Ding-stable with $J^{\text{NA}}_{\mathbb{T}}$.

Although not written down explicitly, the following proposition is actually in [\[Li21,](#page-27-7) [HL23\]](#page-26-9).

Proposition 7.1. Let $(X, -K_X)$ be a Q-Fano variety. Assume that there is a $T \times G$ action on X , where T is a complex algebraic torus and G is a connected reductive algebraic group. We assume that there is a strictly positive smooth function g on Δ_T^+ $\stackrel{+}{T}$. Then X is G−equivariantly g−weighted Ding-semistable if and only if $\delta_G^{\rm g}(X, -K_X) \geq 1$, X is G –equivariantly g—weighted uniformly Ding-stable if and only if $\delta_G^{\mathbf{g}}(X, -K_X) > 1$.

The 'only if' part of the proof of the above proposition is in [\[HL23,](#page-26-9) Theorem 7.8]. Notice that in the proof of [\[HL23,](#page-26-9) Theorem 7.8], the twist can be dropped because of the considerations in [\[Li22,](#page-27-1) 4.1]. The 'if' part is a consequence of [\[Li21,](#page-27-7) Theorem C.7].

Because of the above proposition and Theorem [6.1,](#page-17-1) we immediately get the following result.

Corollary 7.1. Let $(X, -K_X)$ be a Q-Fano spherical G-variety which is a completion of G/H . Let Δ be the polytope associated to the G-action. Let $T = (N_G(H)/H)^0$ and we have the corresponding polytope Δ_T^+ $_T^+$. Assume g is a smooth positive function on Δ_T^+ $\dot{\overline{T}}$. Then the following statements are equivalent:

- 1. The g-weighted barycenter $bar_{DH}^g(\Delta)$ is in the dual cone of $-\mathcal{V}_{\mathbb{R}}$.
- 2. The pair $(X, -K_X)$ is G-equivariantly g-weighted Ding-semistable.

Now we study when a Q−Fano spherical G-variety admits a g−soliton. This follows Golota's idea in [\[Gol20,](#page-26-8) Proposition 5.10].

Let X be a Q–Fano spherical G-variety which is a completion of some G/H . From the last section, we have the torus $T = (N_G(H)/H)^0$. We have the corresponding polytope Δ_T^+ , and g is a strictly positive smooth function on Δ_T^+ .

In [\[Del20\]](#page-26-7) and [\[Del23\]](#page-26-12), Delcroix describes G-equivariant test configurations of $(X, -K_X)$. We follow more closely the Section 4 of [\[Del23\]](#page-26-12). First, we should remark that any $G-$ equivariant test configuration is $T \times G$ –equivariant ([\[Del20,](#page-26-7) section 3.4.1]).

Like in the toric case, a G-equivariant test configuration $(\mathcal{X}, \mathcal{L})$ of the pair $(X, -K_X)$ correspond to a positive rational piecewise linear functions on the moment polytope Δ associated to the preferable B−eigenvector as before. After communications with Delcroix, we find that there is a minor typo in \lfloor Del23. This piecewise linear function is g/r instead of g in Delcroix's terminology. So let's write in this way:

$$
\frac{g(x)}{r} = \inf_{(v,s)\in\mathcal{A}} \left(\frac{v(x)}{-s} + \frac{n_{v,s}}{-rs} \right).
$$

Here $v \in V$, s is a negative integer and A is the set of divisorial valuations on X given by the irreducible component of the central fiber \mathcal{X}_0 . Notice that every element in $\mathcal A$ is $G \times \mathbb{C}^*$ –invariant. Each (v, s) corresponds to an irreducible component of the central fibre \mathcal{X}_0 where v(respectively s) represents the the restriction of the valuation on $\mathbb{C}(X)$ (respectively $\mathbb{C}(t)$ through the inclusion $\mathbb{C}(X) \hookrightarrow \mathbb{C}(\mathcal{X})$ (respectively $\mathbb{C}(t) \hookrightarrow \mathbb{C}(\mathcal{X})$). The pair (v, s) is a primitive element in $N \times \mathbb{Z} \cap \mathcal{V} \times \mathbb{Q}$, where $\mathcal{V} \times \mathbb{Q}$ is actually the cone of $G \times \mathbb{C}^*$ - invariant valuations over $X \times \mathbb{C}^*$. We also recall that $N = \text{Hom}_{\mathbb{Z}}(M, \mathbb{Z})$, where M is the spherical lattice of X. By description above we see that $(\mathcal{X}, \mathcal{L})$ is an integral test configuration, meaning that the scheme theoretic central fibre \mathcal{X}_0 is integral, if and only if there is only one element in \mathcal{A} and $s = -1$. When $(\mathcal{X}, \mathcal{L})$ is integral here, the central fiber \mathcal{X}_0 is a $G \times \mathbb{C}^*$ -stable subvariety of X, then it is also a $G \times \mathbb{C}^*$ -spherical variety. So \mathcal{X}_0 is normal. Then by [\[Ber16,](#page-25-0) Lemma 2.2], with a suitable constant term of the piecewise linear function, we have a special test configuration. Notice also that the special test configuration is furthermore a product test configuration if and only if $v \in \text{Lin}(\mathcal{V}) \cap N$, where $\text{Lin}(\mathcal{V})$ is the maximal Q–linear subspace contained in $\mathcal V$. As mentioned before, the non-Archimedean Ding functional related to the g−soliton takes a very simple form for T-equivariant special test configurations:

$$
D_g(\mathcal{X}, -K_{\mathcal{X}}) = \beta^g(v),
$$

= $A_X(v) - S^g(v),$
= $h_c(v) - \frac{\int_{\Delta} g(\bar{x}) P(x) (\langle x, v \rangle + h_c(v)) dx}{\int_{\Delta} g(\bar{x}) P(x) dx},$
= $-\left\langle \frac{\int_{\Delta} g(\bar{x}) P(x) x dx}{\int_{\Delta} g(\bar{x}) P(x) dx}, v \right\rangle,$
= $-\langle bar_{DH}^g(\Delta), v \rangle.$

Then we get:

Proposition 7.2. Let $(X, -K_X)$ be a Q–Fano spherical G-variety which is a completion of G/H . Let Δ be the polytope associated to the G-action. Let $T = (N_G(H)/H)^0$ and we have the corresponding polytope Δ_T^+ $_T^+$. Assume g is a strictly positive smooth function on Δ_T^+ $^+_T.$ Then $(X, -K_X)$ is G-equivariantly g-weighted Ding-polystable for special test configurations if and only if $bar_{DH}^g(\Delta)$ is in the relative interior of the dual cone of $-\mathcal{V}_{\mathbb{R}}$.

Now we get $T \times G$ -uniformly g–weighted Ding-stability for special test configurations from G-equivariantly g-weighted Ding-polystable for special test configurations. We use 'T × G-uniformly' here since Delcroix twists by $N_T \otimes \mathbb{Q}$ in his paper [\[Del23\]](#page-26-12).

Proposition 7.3. Assume the setup from the last proposition and also that $(X, -K_X)$ is G-equivariantly g-weighted Ding-polystable for special test configurations, then $(X, -K_X)$ is $T \times G$ -uniformly g–weighted Ding-stable for special test configurations.

Proof. The non-Archimedean functionals for spherical test configurations are described in [\[Del23,](#page-26-12) Section 5]. Following the description of J^{NA} there, we just need to prove that there is a $\epsilon > 0$ such that the following inequality holds for any $v \in \mathcal{V}$:

$$
-\langle bar_{DH}^{\mathbf{g}}(\Delta), v \rangle \ge \epsilon \inf_{l \in \text{Lin}(\mathcal{V})} \int_{\Delta} (\max_{\Delta} (v+l) - (v+l)) P dx.
$$

We denote the weighted barycenter $bar_{BH}^g(\Delta)$ by b. By assumption, b is in the relative interior of $-\mathcal{V}_{\mathbb{R}}^{\vee}$.

We pick a complement W of Lin(V) in $N \otimes \mathbb{Q}$. Now any $v \in V$ is written uniquely as $v = v_1 + v_2$, where $v_1 \in \text{Lin}(\mathcal{V})$ and $v_2 \in \mathcal{W} \cap \mathcal{V}$. The set $\mathcal{W} \cap \mathcal{V}$ is a strictly convex polyhedral cone inside W. We know that $-b$ as a function vanishes on Lin(V) and is strictly positive on $W \cap V \setminus \{0\}.$

Now we endow $M \otimes \mathbb{R}$ and $N \otimes \mathbb{R}$ with inner products so they are dual Euclidean spaces. Then:

$$
\inf_{l \in \text{Lin}(\mathcal{V})} \int_{\Delta} (\max_{\Delta} (v+l) - (v+l)) P dx \le \int_{\Delta} (\max_{\Delta} v_2 - v_2) P dx, \n\le C \max_{x,y \in \Delta} \langle v_2, x - y \rangle, \n\le C |v_2| \text{diam}(\Delta),
$$

where diam(Δ) is the diameter of Δ .

We have $-\langle b, v \rangle = -\langle b, v_2 \rangle$. The set $\{v \in \overline{\mathcal{W} \cap \mathcal{V}} \subset \mathcal{W} \otimes \mathbb{R} \text{ such that } |v| = 1\}$ is compact. The function $-b$ is strictly positive on it because $\overline{\mathcal{W} \cap \mathcal{V}}$ is a polyhedral cone generated by some elements in $W \cap V$. So $-b$ is bounded from below by a positive constant on $\{v \in \overline{W} \cap V \subset W \otimes \mathbb{R} \mid |v| = 1\}$. Then we have $-\langle b, v_2 \rangle > C' |v_2|, C' > 0$. on $\{v \in \overline{\mathcal{W} \cap \mathcal{V}} \subset \mathcal{W} \otimes \mathbb{R} \mid |v| = 1\}$. Then we have $-\langle b, v_2 \rangle \ge C' |v_2|, C' > 0$.

Then we get the following theorem:

Theorem 7.1. Let $(X, -K_X)$ be a Q-Fano spherical G-variety which is a completion of G/H . Let Δ be the polytope associated to the G-action. Let $T = (N_G(H)/H)^0$ and we have the corresponding polytope Δ_T^+ $_T^+$. Assume g is a smooth positive function on Δ_T^+ \overline{T} . Then the following three statements are equivalent:

- 1. The g-weighted barycenter $bar_{BH}^g(\Delta)$ is in the relative interior of the dual cone of $-\mathcal{V}_{\mathbb{R}}$.
- 2. The pair $(X, -K_X)$ is G-equivariantly g-weighted Ding-polystable for special test configurations.
- 3. There exists a g−soliton on X.

Proof. The first two are equivalent because of the proposition [7.2.](#page-21-0) We can get (3) from (1) by using the proposition [7.3](#page-21-1) and [\[HL23,](#page-26-9) Theorem 1.7]. If we have (3), we get (2) by using [\[Li21,](#page-27-7) Theorem 1.17] and [\[Li21,](#page-27-7) Theorem 1.21]. \Box

Theorem [7.1](#page-22-0) a generalization of Theorem A in [\[Del20\]](#page-26-7).

As written in the introduction, after we finish writing, we are informed that the theorem above is already given in [\[LLW22\]](#page-27-6). Lemma 6.1 in their paper basically serves the purpose of proposition [7.2.](#page-21-0) Our points of views are slightly different. We use valuative theory, so β -invariant is the most natural tool to us. They approach the problem with test configurations. But of course, the two points of views coincide on special test configurations.

Remark 7.3. There is another way to get (3) from (2) in the above theorem. By Minimal Model Program developed in $[LX14]$, $[Fuj19]$, we actually know that G-equivariantly integral g-weighted Ding-polystability is equivalent to G-equivariantly g-weighted Ding-polystability. The latter is equivalent to reduced uniformly g–weighted stability by [\[Li21,](#page-27-7) Theorem 1.17] and [\[BLXZ23,](#page-25-11) Theorem 1.3]. Then reduced uniformly g−weighted stability is equivalent to the existence of the g−soliton by [\[Li21,](#page-27-7) Theorem 1.21].

8 Applications to Sasaki-Einstein metrics

In [\[AJL23,](#page-25-12) proposition 2], the authors describe an interesting relationship between a special type of g−soliton and the existence of Sasaki-Einstein structure.

Proposition 8.1. let X be a smooth Fano variety of complex dimension m. Let T be a complex torus acting on X with the canonical moment polytope Δ_T^+ $_{T}^{+}$. Let l be a positive affine-linear function on Δ_T^+ written as $l(x) = \langle \xi, x \rangle + a$. Let T^c be the real compact torus

inside T. A T^c - invariant Kähler metric $\omega \in 2\pi c_1(X)$ is an $l^{-(m+2)}$ -soliton if and only if the lift $\hat{\xi}$ of ξ to K_X via l is the Reeb vector field of a Sasaki-Einstein structure defined on the unit circle bundle inside K_X with respect to the hermitian metric on K_X with curvature $-\omega$.

The authors of [\[AJL23\]](#page-25-12) use this proposition to prove the existence of a Sasaki-Einstein structure on a unit circle bundle associated to K_X for a Fano semi-simple principle bundle X with a smooth toric Fano fiber.

We can use their strategy to get a similar result for smooth Fano horospherical manifolds. First we need to introduce some backgrounds here.

Let G, B be as above. Let U be the unipotent radical of B. Let R be the set of roots of (G, T) and R^+ be the set of positive roots associated to B. Thus we have the root decomposition:

$$
\mathfrak{g}=\mathfrak{t}\oplus\bigoplus_{\alpha\in R}\mathfrak{g}_{\alpha}
$$

Let S be the set of simple roots. Let H be a closed subgroup of G which contains U . Then we call H a horospherical subgroup of G (associated to B) and G/H a horospherical homogeneous space. Let $P = N_G(H)$, this is a parabolic subgroup of G containing B. The parabolic subgroup has the unipotent radical $R_u(P)$. Let's denote $I \subset S$ as the set of $\alpha \in S$ such that \mathfrak{g}_{α} is not in the Lie algebra of $R_u(P)$. If we denote R_I as the sub root system of R generated by I, then the unique Levi subgroup L of P which contains T has the Lie algebra

$$
\mathfrak{l}=\mathfrak{t}\oplus\bigoplus_{\alpha\in R_I}\mathfrak{g}_{\alpha}
$$

In this situation, P/H is actually a torus and the morphism θ introduced in section 6 is an isomorphism. What makes the combinatorial criterion in Section 7 simpler is the fact that the valuation cone V is the whole $N \otimes \mathbb{Q}$ in this situation, so the dual cone of $-\mathcal{V}$ is just $\{0\}$.

Now we let X be a smooth projective Fano completion of G/H . In this case, we have $2\rho_P = \sum_{\alpha \in R^+ \setminus R_I} \alpha \text{ and } P(x) = \prod_{\alpha \in R^+ \setminus R_I}$ $\langle x+2\rho_P , \alpha \rangle$ $\frac{\langle \rho \rangle \langle \rho \rangle}{\langle \rho \rangle \langle \rho \rangle}.$

We shall prove the following theorem:

Theorem 8.1. Let X be a smooth Fano horospherical variety. Then the corresponding cone $(K_X)^\times$ admits a Calabi-Yau cone metric, given by a Sasaki-Einstein structure on a unit circle inside K_X with respect to some hermitian metric on K_X .

Proof. We follow the argument of [\[AJL23\]](#page-25-12). Assume that our X has complex dimension m complex dimension of P/H is r. So r is also the real dimension of Δ . We have:

$$
m = \dim(X) = \dim(P/H) + \dim(G/P)
$$

$$
= r + \#R^+\backslash R_I.
$$

Let's consider the set $\Delta^* = \{\xi \in N \otimes \mathbb{R} | \langle \xi, x \rangle + 1 \geq 0 \text{ for } x \in \Delta \}.$ This is the so called dual polytope of Δ . Polytopes like Δ^* play a very important role in classifying Fano horospherical varieties, see for example $|Pas08|$.

Because of Theorem [7.1](#page-22-0) and Proposition [8.1,](#page-22-1) we just need to find $\xi \in \text{Int}(\Delta^*)$ such that:

$$
\int_{\Delta} (\langle \xi, x \rangle + 1)^{-m-2} x P(x) dx = 0 \tag{1}
$$

In order to do this, we consider the functional:

$$
\xi \in \mathrm{Int}(\Delta^*) \mapsto \int_{\Delta} (\langle \xi, x \rangle + 1)^{-m-1} P(x) dx.
$$

This functional is clearly strictly convex since we can take derivatives twice and see that the Hessian is strictly positive. Now we have to show that this functional is also proper, meaning the functional goes to infinity when ξ goes to some boundary point of Δ^* . If this happens, then the functional has a unique minimum point $\xi \in \text{Int}(\Delta^*)$, then take the derivative we see that ξ satisfies the equation [1.](#page-24-0) Now let's say that some sequence $\{\xi_i\} \subset \text{Int}(\Delta^*)$ converges to some boundary point ξ . For any small positive c, we consider the set $\Delta_c = \{x \in \Delta | \langle \xi, x \rangle + 1 \geq c\}.$ Clearly

$$
\int_{\Delta_c} (\langle \xi_i, x \rangle + 1)^{-m-1} P(x) dx \to \int_{\Delta_c} (\langle \xi, x \rangle + 1)^{-m-1} P(x) dx.
$$

Making c smaller and smaller we only need to show:

$$
\int_{\Delta} (\langle \xi, x \rangle + 1)^{-m-1} P(x) dx = +\infty
$$

for $\xi \in \partial \Delta^*$.

This is rather clear if $P(x)$ is strictly positive on Δ , which is the case when X is a toroidal horospherical Fano manifold (see [\[Pas08,](#page-27-14) section 3]). But $P(x)$ can have zeroes on $\partial \Delta$. Let's explain how to deal with this case by giving first an intuitive argument. Since $\xi \in \partial \Delta^*,$ we must have $\langle \xi, y \rangle + 1 = 0$ for some vertex y of Δ . Around y, the term $(\langle \xi, x \rangle + 1)^{-m-1}$ provides a pole of $(m + 1)$ order. The term $P(x)$ provides zero of at most $\#R^+\backslash R_I$ order. The quotient will then have a pole of at least $(r + 1)$ order. Let's provide now more details.

Since $\xi \in \partial \Delta^*$, we must have $\langle \xi, y \rangle + 1 = 0$ for some vertex y of Δ . Let's denote $J = {\alpha \in S | y + 2\rho_P, \alpha \rangle = 0}.$ Notice that simple roots in I are all orthogonal to $2\rho_P + \Delta$, so $I \subset J$. We have a sub-root system R_J generated by J, let's denote the positive part as R_J^+ $_J^+$. Clearly if a positive root α satisfies $\langle y+2\rho_P, \alpha \rangle = 0$, we must have $\alpha \in R_J^+$ j^+ . We can pick a very large positive number c and consider the set $\Delta' = \Delta \cap \bigcap_{\alpha \in R_J^+ \setminus R_I} \{c\langle x + 2\rho_P, \alpha \rangle \geq \langle x, \xi \rangle + 1\}.$ Clearly Δ' is just $\Delta \cap \bigcap_{\alpha \in J\setminus I} \{c\langle x + 2\rho_P, \alpha \rangle \geq \langle x, \xi \rangle + 1\}$. Now in some region Δ'' inside Δ' and close enough to z we have that

$$
(\langle \xi, x \rangle + 1)^{-m-1} P(x) \ge C(\langle \xi, x \rangle + 1)^{-m-1+\#R_J^+ \setminus R_I} \ge C(\langle \xi, x \rangle + 1)^{-r-1},
$$

for some positive constant C. Hence we obtain as expected:

$$
\int_{\Delta} (\langle \xi, x \rangle + 1)^{-m-1} P(x) dx \ge C \int_{\Delta''} (\langle \xi, x \rangle + 1)^{-r-1} dx = +\infty.
$$

 \Box

Remark 8.1. In [\[Ngh23\]](#page-27-15), Nghiem establishes establish the equivalence between a weighted volume minimization principle and the existence of a conical Calabi-Yau structure on horospherical cones with klt singularities. One can get Theorem [8.1](#page-23-0) by using his method.

References

- [AJL23] Vestislav Apostolov, Simon Jubert, and Abdellah Lahdili. Weighted K–stability and coercivity with applications to extremal Kähler and Sasaki metrics. Geom. Topol., 27(8):3229–3302, 2023.
- [AK05] Valery Alexeev and Ludmil Katzarkov. On K-stability of reductive varieties. Geometric & Functional Analysis GAFA, $15(2):297-310$, 2005.
- [BBE⁺19] Robert J Berman, Sebastien Boucksom, Philippe Eyssidieux, Vincent Guedj, and Ahmed Zeriahi. Kähler–einstein metrics and the kähler–ricci flow on log fano varieties. Journal für die reine und angewandte Mathematik (Crelles Journal), 2019(751):27–89, 2019.
- [BdFFU15] S. Boucksom, T. de Fernex, C. Favre, and S. Urbinati. Valuation spaces and multiplier ideals on singular varieties. In Recent advances in algebraic geometry, volume 417 of London Math. Soc. Lecture Note Ser., pages 29–51. Cambridge Univ. Press, Cambridge, 2015.
- [Ber16] Robert J Berman. K-polystability of Q-Fano varieties admitting Kähler-Einstein metrics. Inventiones mathematicae, 203(3):973–1025, 2016.
- [BHJ17] S´ebastien Boucksom, Tomoyuki Hisamoto, and Mattias Jonsson. Uniform Kstability, Duistermaat-Heckman measures and singularities of pairs. Ann. Inst. Fourier (Grenoble), 67(2):743–841, 2017.
- [BJ20] Harold Blum and Mattias Jonsson. Thresholds, valuations, and K-stability. Advances in Mathematics, 365:107062, 2020.
- [BK07] Michel Brion and Shrawan Kumar. Frobenius splitting methods in geometry and representation theory, volume 231. Springer Science & Business Media, 2007.
- [BLXZ23] Harold Blum, Yuchen Liu, Chenyang Xu, and Ziquan Zhuang. The existence of the Kähler-Ricci soliton degeneration. Forum Math. Pi , 11: Paper No. e9, 28, 2023.
- [BN14] Robert J Berman and David Witt Nystrom. Complex optimal transport and the pluripotential theory of Kähler-Ricci solitons. $arXiv$ preprint $arXiv:1401.8264$, 2014.
- [Bri87] Michel Brion. Sur l'image de l'application moment. In Séminaire d'Algèbre Paul Dubreil et Marie-Paule Malliavin: Proceedings, Paris 1986, pages 177–192. Springer, 1987.
- [Bri89] Michel Brion. Groupe de Picard et nombres caractéristiques des variétés sphériques. *Duke Math. J.*, 58(2):397–424, 1989.
- [Bri97a] Michel Brion. Curves and divisors in spherical varieties. Algebraic groups and Lie groups, 9:21–34, 1997.
- [Bri97b] Michel Brion. Variétés sphériques. Notes de cours, 1997.
- [CDS15a] Xiuxiong Chen, Simon Donaldson, and Song Sun. K¨ahler-Einstein metrics on Fano manifolds. I: Approximation of metrics with cone singularities. Journal of the American Mathematical Society, 28(1):183–197, 2015.
- [CDS15b] Xiuxiong Chen, Simon Donaldson, and Song Sun. K¨ahler-Einstein metrics on Fano manifolds. II: Limits with cone angle less than 2π . J. Amer. Math. Soc., 28(1):199–234, 2015.
- [CDS15c] Xiuxiong Chen, Simon Donaldson, and Song Sun. K¨ahler-Einstein metrics on Fano manifolds. III: Limits as cone angle approaches 2π and completion of the main proof. J. Amer. Math. Soc., 28(1):235–278, 2015.
- [Cox00] David A Cox. Toric varieties and toric resolutions. Resolution of Singularities: A research textbook in tribute to Oscar Zariski Based on the courses given at the Working Week in Obergurgl, Austria, September 7–14, 1997, pages 259–284, 2000.
- [CS08] Ivan A Cheltsov and Konstantin A Shramov. Log canonical thresholds of smooth Fano threefolds. Russian Mathematical Surveys, 63(5):859, 2008.
- [Del17] Thibaut Delcroix. Log canonical thresholds on group compactifications. Algebr. Geom., 4(2):203–220, 2017.
- [Del20] Thibaut Delcroix. K-stability of Fano spherical varieties. Ann. Sci. Ec. Norm. $Supér. (4), 53(3):615–662, 2020.$
- [Del23] Thibaut Delcroix. Uniform K-stability of polarized spherical varieties. $Epijournal Géom. Algébrique, 7:Art. 9, 27, 2023. With an appendix by Yuji$ Odaka.
- [FO18] Kento Fujita and Yuji Odaka. On the K-stability of Fano varieties and anticanonical divisors. Tohoku Mathematical Journal, 70(4):511–521, 2018.
- [Fuj19] Kento Fujita. A valuative criterion for uniform K-stability of Q-Fano varieties. J. Reine Angew. Math., 751:309–338, 2019.
- [GH15] Giuliano Gagliardi and Johannes Hofscheier. Gorenstein spherical Fano varieties. Geometriae Dedicata, 178(1):111–133, 2015.
- [Gol20] Aleksei Golota. Delta-invariants for Fano varieties with large automorphism groups. International Journal of Mathematics, 31(10):2050077, 2020.
- [HL23] Jiyuan Han and Chi Li. On the Yau-Tian-Donaldson Conjecture for Generalized Kähler-Ricci Soliton Equations. Communications on Pure and Applied Mathematics, 76(9):1793–1867, 2023.
- [Kno91] Friedrich Knop. The Luna-Vust theory of spherical embeddings. In *Proceedings* of the Hyderabad Conference on Algebraic Groups (Hyderabad, 1989), volume 225, page 249. Citeseer, 1991.
- [Li17] Chi Li. K-semistability is equivariant volume minimization. Duke Math. J., 166(16):3147–3218, 2017.
- [Li21] Chi Li. Notes on weighted Kähler-Ricci solitons and application to Ricci-flat Kähler cone metrics. arXiv preprint arXiv:2107.02088, 2021.
- [Li22] Chi Li. G-uniform stability and Kähler–Einstein metrics on Fano varieties. Inventiones mathematicae, 227(2):661–744, 2022.
- [LLW22] Yan Li, ZhenYe Li, and Feng Wang. Weighted K-stability of Q-Fano spherical varieties. arXiv preprint arXiv:2208.02708, 2022.
- [LX14] Chi Li and Chenyang Xu. Special test configuration and K-stability of Fano varieties. Annals of mathematics, pages 197–232, 2014.
- [LXZ22] Yuchen Liu, Chenyang Xu, and Ziquan Zhuang. Finite generation for valuations computing stability thresholds and applications to K-stability. Annals of Mathematics, 196(2):507–566, 2022.
- [Nad90] Alan Michael Nadel. Multiplier ideal sheaves and Kahler-Einstein metrics of positive scalar curvature. Annals of Mathematics, pages 549–596, 1990.
- [Ngh23] Tran-Trung Nghiem. Spherical cones: classification and a volume minimization principle. The Journal of Geometric Analysis, 33(7):221, 2023.
- $[Pas08]$ Boris Pasquier. Variétés horosphériques de Fano. Bulletin de la société mathématique de France, $136(2):195-225$, 2008.
- [Pas17] Boris Pasquier. A survey on the singularities of spherical varieties. EMS Surveys in Mathematical Sciences, 4(1):1–19, 2017.
- [Per14] Nicolas Perrin. Introduction to spherical varieties. Lecture notes, 2014.
- [RTZ21] Yanir A Rubinstein, Gang Tian, and Kewei Zhang. Basis divisors and balanced metrics. Journal für die reine und angewandte Mathematik (Crelles Journal), 2021(778):171–218, 2021.
- [Tia87] Gang Tian. On K¨ahler-Einstein metrics on certain K¨ahler manifolds with $C_1(M)$ >0. Inventiones mathematicae, 89:225–246, 1987.
- [Tia15] Gang Tian. K-stability and K¨ahler-Einstein metrics. Communications on Pure and Applied Mathematics, 68(7):1085–1156, 2015.
- [Tim11] Dmitry A Timashev. Homogeneous spaces and equivariant embeddings, volume 138. Springer Science & Business Media, 2011.
- [TY87] Gang Tian and Shing-Tung Yau. Kähler-Einstein metrics on complex surfaces with $C_1 > 0$. Comm. Math. Phys., 112(1):175-203, 1987. [WZ04] Xu-Jia Wang and Xiaohua Zhu. Kähler–Ricci solitons on toric manifolds with positive first Chern class. Advances in Mathematics, 188(1):87–103, 2004. [XZ20] Chenyang Xu and Ziquan Zhuang. On positivity of the CM line bundle on K-moduli spaces. Annals of mathematics, 192(3):1005–1068, 2020.
- [Zha22] Kewei Zhang. Valuative invariants with higher moments. The Journal of Geometric Analysis, 32(1):10, 2022.

Chenxi Yin, Département de mathématiques, UQAM, Montréal, Canada Email address: yin.chenxi@courrier.uqam.ca