A PARALLEL DOMAIN DECOMPOSITION METHOD FOR SOLVING ELLIPTIC EQUATIONS ON MANIFOLDS

LIZHEN QIN *, FENG WANG [†], AND YUN WANG [‡]

Abstract. We propose a new numerical domain decomposition method for solving elliptic equations on compact Riemannian manifolds. One advantage of this method is its ability to bypass the need for global triangulations or grids on the manifolds. Additionally, it features a highly parallel iterative scheme. To verify its efficacy, we conduct numerical experiments on some 4-dimensional manifolds without and with boundary.

AMS subject classifications. Primary 65N30; Secondary 58J05, 65N55.

Key words. Riemannian manifolds, elliptic problems, domain decomposition methods, finite element methods

1. Introduction. Elliptic equations on Riemannian manifolds are important both in analysis and in geometry (see e.g. [46] and [34]). These equations appear in many areas, such as multifluid dynamics, micromagnetics, and image processing (see e.g. [3, 38, 45, 17, 30, 29, 6, 2, 32, 33, 4]). A simple and important example of such equations is

$$(1.1) \qquad \qquad -\Delta u + bu = f.$$

Here Δ is the Laplace-Beltrami operator, or Laplacian for brevity, defined on a *d*dimensional Riemannian manifold *M*. Many manifolds are naturally submanifolds of Euclidean spaces, where the "dimension" of a submanifold is referred to as the topological dimension of the manifold, not the dimension of its ambient Euclidean space. For example, the *n*-dimensional unit sphere S^n is embedded in \mathbb{R}^{n+1} .

When the manifold M is a two dimensional Riemannian submanifold in \mathbb{R}^3 , i.e. a surface, the numerical methods to solve differential equations, particularly (1.1), on M have been extensively studied for a long history (see e.g. [40], [39], [5], [19] and [20]). A conventional and popular approach is to solve the equations by finite element methods based on a global grid or triangulation on M. Such a grid can be obtained by polyhedron approximation or level set method. This approach has been far-reaching developed and widely applied (see e.g. [15, 21, 14, 1, 44, 41], see also e.g. [13, 22, 7] for surveys and bibliographies).

However, practically, it's of essential difficulty to build a global triangulation on a higher dimensional manifold for the purpose of numerical computation. The difficulty comes from at least three aspects. Firstly, high dimensional spaces challenge significantly people's geometric intuition. The experience of tessellation in \mathbb{R}^2 and \mathbb{R}^3 cannot be simply ported to higher dimensions. Secondly, many important and interesting examples of high dimensional manifolds are not submanifolds of Euclidean spaces by definition. A case in point is the complex projective spaces \mathbb{CP}^n which is foundational to algebraic geometry. The usual polytopal approximation or level set

^{*}Mathematics Department, Nanjing University, Nanjing, Jiangsu, China (qinlz@nju.edu.cn).

[†]Key Laboratory of Numerical Simulation of Large Scale Complex Systems (Nanjing Normal University), Ministry of Education; Jiangsu International Joint Laboratory of Big Data Modeling, Computing, and Application; School of Mathematical Sciences, Nanjing Normal University, Nanjing Jiangsu, China (fwang@njnu.edu.cn).

[‡]School of Mathematical Sciences, Center for Dynamical Systems and Differential Equations, Soochow University, Suzhou, China (ywang3@suda.edu.cn).

method is not applicable to such manifolds. Thirdly, even if M is a submanifold of some \mathbb{R}^k by definition, if the codimension of M in \mathbb{R}^k is greater than 1, which is often the case, the triangulation of M is horribly difficult in general due to topological and geometrical complexity. We do acknowledge that J. H. C. Whitehead proved ([50]) that every smooth manifold can be globally triangulated in an abstract way. However, to the best of our knowledge, in practice, there has been no efficient algorithm to construct concrete triangulations over a high dimensional manifold in general.

To circumvent the above difficulty, in [42], Qin–Zhang–Zhang proposed a new idea to numerically solve elliptic PDEs on manifolds by avoiding global triangulations. Since a *d*-dimensional manifold M has local coordinate charts by definition, M can be decomposed into finitely many subdomains that carry local coordinates. Consequently, an elliptic equation on each subdomain can be transformed to one on a domain in \mathbb{R}^d . Thus an elliptic problem on M can be solved by Domain Decomposition Methods (DDMs) with subproblems on Euclidean domains.

The idea of [42] was developed by Cao–Qin in [10]. They extended a DDM proposed by P. L. Lions ([36, Section I. 4]) for solving problems on Euclidean domains to a DDM on manifolds. In this paper, we combine the ideas of [42] and [37], which yields an adaption and generalization of Lions' another DDM ([37, p. 66]). This DDM was also proposed by Lions for solving continuous problems on Euclidean domains. We found that it can also be well-adapted to a DDM on manifolds. One notable advantage of this method is its high level of parallelism, making it particularly valuable when dealing with a large number of subdomains.

Here is an overview of the method in this paper. First of all, global triangulation/grid is completely avoided. There is also no requirement of any compatibility among local grids, i.e. grids over subdomains. (As explained in the introduction of [10], this incompatibility actually reflects the high flexibility of the numerical methods.) The numerical difficulty caused by the nonmatching grids is easily resolved by the technique of interpolation. Furthermore, the subproblems on subdomains can be localized and then be reduced to problems on Euclidean domains. These features bear resemblance to that method described in [10]. However, this paper is remarkably different from [10] in that the current method is highly parallel. In fact, this work follows [37], while [10] follows [36]. Lions described the work [36, Section I. 4] as "sequential" (see the sentence right before (26) in [36]), which means the subproblems on subdomains have to be solved one by one sequentially. On the contrary, the work [37] was claimed as "parallel" (see a few sentences before (37) in [37]), which means the subproblems on subdomains can be solved independently and parallelly at each iterative step.

It is worth noting that, even in the special case where the target manifold is a Euclidean domain, our algorithm (see Algorithms 2.1 and 3.1) is different from [37, p. 66] though we follow it largely. Typically, we employ a technique of partition of unity which was not utilized in [37, p. 66]. Due to this technique, we obtain a better convergence theory (see Theorem 2.4 and Remark 2.5).

Overall, our method is an overlapping DDM on manifolds which avoids global grids. There has been a long history of DDM. It was originated from the work of H. A. Schwarz [47] who invented his famous alternating method. The seminal works [36, 37] of P. L. Lions are natural and wonderful extensions of Schwarz alternating method. From then on, the area of DDM has been developed explosively and greatly (see e.g. [43, 48, 49, 16]). The later development often take DDM schemes, including both overlapping and nonoverlapping ones, as preconditioners for globally discretized problems (see e.g. [8, 51, 9, 52, 18]). However, a globally discretized problem on a

manifold M should be based on a global grid on M, which is not accessible in our case. Therefore, we follow more closely Lions' original approach rather than the later development.

Our DDM in this paper is numerically verified on examples of 4-dimensional manifolds. They are \mathbb{CP}^2 , $S^2 \times S^2$, B^4 and $B^2 \times S^2$, where \mathbb{CP}^2 is the complex projective plane, S^2 is the 2-dimensional unit sphere, and B^d is the *d*-dimensional unit ball. The former two manifolds are without boundary, while the latter two are with boundary. The numerical results show that our method solves effectively the equation (1.1) on these manifolds.

The outline of this paper is as follows. In Section 2, we shall adapt and extend P. L. Lions' method in [37, p. 66] to a DDM for continuous problems on manifolds and prove its convergence. In Section 3, we shall propose our numerical DDM which is a discrete imitation of Lions' method. In Section 4, we shall introduce a technique to decompose manifolds with boundary. Finally, some numerical results will be presented in Section 5.

2. Theory on Continuous Problems. In this section, we shall firstly formulate an elliptic model problem of second order on manifolds. Then we introduce a domain decomposition method (DDM) to solve the model problem (see Algorithm 2.1 below). This DDM iteration procedure was originally proposed by P. L. Lions in [37, p. 66] for solving differential equations in a Euclidean domain. Unlike those DDMs in [36] and [10], this iterative procedure is highly parallel. More precisely, the subproblem in each subdomain can be solved independently at each iterative step. This procedure motivates us to propose a highly parallel numerical procedure (Algorithm 3.1) in the next section which provides numerical approximations to the solution of the model problem.

2.1. Model Problem and Continuous Algorithm. We shall formulate a model problem, the continuous Algorithm 2.1, and Theorem 2.4 on convergence. The proof of convergence is much more complicated than those theorems in [36] and [10], and will occupy the next two subsections.

Let M be a d-dimensional compact smooth manifold without or with boundary ∂M . Equip M with a Riemannian metric g. Then the Laplace operator Δ , also named the Laplace-Beltrami operator, is defined on M. Note that neither g nor Δ can be expressed by coordinates globally in general because M does not necessarily have a global coordinate chart. In a local chart with coordinates (x_1, \ldots, x_d) , the Riemannian metric tensor g is expressed as

(2.1)
$$g = \sum_{\alpha,\beta=1}^{d} g_{\alpha\beta} \mathrm{d} x_{\alpha} \otimes \mathrm{d} x_{\beta},$$

where the matrix $(g_{\alpha\beta})_{d\times d}$ is symmetric and positive definite. The Laplace operator Δ is expressed as

$$\Delta u = \frac{1}{\sqrt{G}} \sum_{\alpha=1}^{d} \frac{\partial}{\partial x_{\alpha}} \left(\sum_{\beta=1}^{d} g^{\alpha\beta} \sqrt{G} \frac{\partial u}{\partial x_{\beta}} \right),$$

where $G = \det((g_{\alpha\beta})_{d\times d})$ is the determinant of the matrix $(g_{\alpha\beta})_{d\times d}$ and $(g^{\alpha\beta})_{d\times d}$ is the inverse of $(g_{\alpha\beta})_{d\times d}$. It's well-known that Δ is an elliptic differential operator of second order. We consider the following model problem on M

(2.2)
$$\begin{cases} Lu = -\Delta u + bu = f, \\ u|_{\partial M} = \varphi, \end{cases}$$

where $b \ge 0$ is a constant and $f \in L^2(M)$. It suffices to solve (2.2) on each component of M. Therefore, without loss of generality, M is assumed to be connected. Furthermore, it is possible for ∂M to be either empty or nonempty. In the case where $\partial M \ne \emptyset$, we assume $\varphi \in C^0(\partial M) \cap H^{\frac{1}{2}}(\partial M)$. On the other hand, if $\partial M = \emptyset$, the boundary condition $u|_{\partial M} = \varphi$ is vacuously satisfied. However, to guarantee (2.2) is well-posed, we need to further assume b > 0 if $\partial M = \emptyset$. With these assumptions, the problem (2.2) always has a unique solution $u \in C^0(M) \cap H^1(M)$.

Now we describe a domain decomposition iteration procedure to solve (2.2). This method was originally proposed by P. L. Lions in [37, p. 66] for solving differential equations in a Euclidean domain. We found that the method is well-adapted to solve problems on manifolds. Suppose M is decomposed into m subdomains, i.e. $M = \bigcup_{i=1}^{m} M_i$. Here M_i is a closed subdomain (submanifold with codimension 0) of M with Lipschitz boundary. Let $\gamma_i = \partial M_i \setminus \partial M$. We make the following assumption on decomposition.

ASSUMPTION 2.1. Suppose $M = \bigcup_{i=1}^{m} (M_i \setminus \overline{\gamma_i})$, and $S := \bigcup_{i \neq j} (\gamma_i \cap \gamma_j)$ has a finite (d-2)-dimensional Hausdorff measure.

The requirement of the Hausdorff measure of S can be easily satisfied. For example, we may choose M_i with piecewise C^1 boundary and arrange that all γ_i meet transversely with each other. Transversality is a generic property which, therefore, can be satisfied almost trivially. (For a detailed theory on transversality, see e.g. [28, Chapter 3].)

Suppose $\{\rho_i \mid 1 \leq i \leq m\}$ is a partition of unity of M, i.e. these ρ_i are a family of nonnegative functions on M and $\sum_{i=1}^{m} \rho_i = 1$.

ASSUMPTION 2.2. The partition of unity $\{\rho_i \mid 1 \leq i \leq m\}$ is subordinate to the decomposition in Assumption 2.1, i.e. $\operatorname{supp}\rho_i \subset M_i \setminus \overline{\gamma_i}$, where $\operatorname{supp}\rho_i$ is the support of ρ_i . Furthermore, each ρ_i is Lipschitz continuous.

Now we are ready to formulate the desired algorithm of solving continuous problem (2.2).

Algorithm 2.1 A DDM to solve (2.2).

- 1: Choose an arbitrary initial guess $u^0 \in C^0(M) \cap H^1(M)$ with $u^0|_{\partial M} = \varphi$ for (2.2).
- 2: For each n > 0, assuming u^{n-1} has been obtained, for $1 \le i \le m$, find a function u_i^n on M_i such that

(2.3)
$$\begin{cases} Lu_i^n(x) = f(x), & x \in M_i \setminus \partial M_i; \\ u_i^n(x) = u^{n-1}(x), & x \in \partial M_i. \end{cases}$$

3: Let
$$u^n = \sum_{i=1}^m \rho_i u_i^n$$
.

Unlike those methods in [36, Section I. 4] and [10], Algorithm 2.1 is highly parallel. Obviously, the subproblems in (2.3) are solved independently on M_i for all *i*. It's necessary to point out that u_i^n in (2.3) is merely defined on M_i . However, u^n is well-defined on the whole M since $\operatorname{supp} \rho_i \subset M_i$. (More precisely, the function $\rho_i u_i^n$ on M_i is extended as 0 outside of M_i .) We shall prove that $\{u_i^n\}$ and $\{u^n\}$ converge to the solution u in certain sense.

The boundary value condition of u_i^n in Algorithm 2.1 is slightly different from Lions' original approach in [37]. More precisely, a partition of unity was not employed in [37], instead, the boundary value $u_i^n|_{\gamma_i}$ was chosen as an arbitrary function v such that (see [37, (42) & (43)])

$$\min\{u_{j}^{n-1}(x) \mid x \in M_{j}, j \neq i\} \le v(x) \le \max\{u_{j}^{n-1}(x) \mid x \in M_{j}, j \neq i\}.$$

Our $u_i^n|_{\gamma_i}$ in Algorithm 2.1 certainly satisfies the above inequalities. Moreover, by a partition of unity satisfying Assumption 2.2, the iteration in Algorithm 2.1 even has a better theory of convergence.

First of all, the u_i^n in [37] was usually not continuous on M_i since $u_i^n|_{\gamma_i}$ was not necessarily continuous. However, the following lemma shows our u_i^n in (2.3) has better regularity. Let $u_i^0 = u^0|_{M_i}$. In the following, all statements on $u_i^n|_{\partial M}$ are vacuously true if $M_i \cap \partial M = \emptyset$.

LEMMA 2.3. Under the Assumptions 2.1 and 2.2, in (2.3), we have $\forall n, \forall i, u^n \in C^0(M) \cap H^1(M), u_i^n \in C^0(M_i) \cap H^1(M_i), u^n|_{\partial M} = \varphi$, and $u_i^n|_{\partial M} = \varphi$.

Proof. We already know the conclusion holds for n = 0. Suppose the conclusion is true for n - 1. Then $u^{n-1} - u \in C^0(M) \cap H^1(M)$. Furthermore, u_i^n satisfies

$$\begin{cases} L(u_i^n - u)(x) = 0, & x \in M_i \setminus \partial M_i; \\ (u_i^n - u)(x) = (u^{n-1} - u)(x), & x \in \partial M_i. \end{cases}$$

By [25, Corollary 8.28], we infer $u_i^n - u \in C^0(M_i) \cap H^1(M_i)$. As $u \in C^0(M) \cap H^1(M)$, we see $u_i^n \in C^0(M_i) \cap H^1(M_i)$. By Rademacher's theorem (see Theorem 5 in [24, §4.2.3]), the Lipschitz continuity of ρ_i means $\rho_i \in W^{1,\infty}(M)$, which implies $u^n \in C^0(M) \cap H^1(M)$. Since $u^{n-1}|_{\partial M} = \varphi$, we have $u_i^n|_{\partial M} = \varphi$ and hence $u^n|_{\partial M} = \varphi$. \Box

Secondly, we have the following theorem on convergence.

THEOREM 2.4. Under the Assumptions 2.1 and 2.2, suppose the solution $u \in C^0(M) \cap H^1(M)$. Assume further u and the initial guess u^0 are Lipschitz continuous if $\partial M \neq \emptyset$. Let

$$\theta^n = \max\{\|u_i^n - u\|_{C^0(M_i)} \mid 1 \le i \le m\}.$$

Then the following holds:

(1) $\lim_{n \to \infty} \theta^n = 0;$ (2) $\forall n > 0, \|u^n - u\|_{C^0(M)} \le \theta^n \text{ and } \|u^n - u\|_{H^1(M)} \le C_1 \theta^n;$ (3) $\forall n > 1, \max\{\|u_i^n - u\|_{H^1(M_i)} \mid 1 \le i \le m\} \le C_2 \theta^{n-1}.$ Here C_1 and C_2 are constants independent of n and i.

Theorem 2.4 was essentially proved by Lions in [37, p. 66-67] in the case that M is a Euclidean domain. Even in that special case, the proof was complicated. Actually, the proof in [37] was presented in a dense style. We shall follow and adapt Lions' argument and then extend it to the case of general manifolds. Our proof will be given in the next two subsections. To ensure the readability for the reader, we shall provide plenty of details of argument.

REMARK 2.5. Even in the case that M is a Euclidean domain, the statement on convergence in [37] is slightly different from Theorem 2.4. In [37], the function u_i^n was not necessarily continuous on M_i , the uniform convergence and H^1 -convergence of u_i^n

on M_i were not claimed either. However, it was proved that u_i^n converges uniformly to u on each compact set $K_i \subset M_i \setminus \overline{\gamma_i}$ there.

REMARK 2.6. One may obtain a convergence theory better than Theorem 2.4 if one imposes an additional assumption as follows: every three distinct subdomains have an empty intersection (cf. [37, (36)]). As pointed in [37, Proposition 2 and p. 63-65], under this assumption, one can prove that the convergence is geometric, which is stronger than the conclusion of Theorem 2.4, by an alternative and much easier approach. However, this additional assumption is stringent in practice. In fact, none of our numerical examples satisfies this assumption.

2.2. Subsolutions and Supersolutions. The key ingredient of the proof of Theorem 2.4 is the technique of subsolutions and supersolutions. By considering $u_i^n - u$, it suffices to study the following homogeneous problem on M

(2.4)
$$\begin{cases} L\hat{u} = 0, \\ \hat{u}|_{\partial M} = 0, \end{cases}$$

which obviously has a unique solution $\hat{u} = 0$.

As mentioned before, L can be expressed as

(2.5)
$$Lu = -\frac{1}{\sqrt{G}} \sum_{\alpha=1}^{d} \frac{\partial}{\partial x_{\alpha}} \left(\sum_{\beta=1}^{d} g^{\alpha\beta} \sqrt{G} \frac{\partial u}{\partial x_{\beta}} \right) + bu$$

in a local chart with coordinates (x_1, \ldots, x_d) . Let's recall the definition of subsolutions and supsolutions in the sense of viscosity (cf. [12, Definition 2.2]).

DEFINITION 2.7. A function \underline{u} (resp. \overline{u}) on $M \setminus \partial M$ is a subsolution (resp. supersolution) of the equation $L\hat{u} = 0$ if it satisfies:

(1) \underline{u} (resp. \overline{u}) is upper (resp. lower) semicontinuous;

(2) for each $\hat{x} \in M \setminus \partial M$ and each C^2 function ψ on $M \setminus \partial M$ such that \hat{x} is a local maximum (resp. minimum) of $\underline{u} - \psi$ (resp. $\overline{u} - \psi$), we have

$$\begin{split} & -\frac{1}{\sqrt{G}}\sum_{\alpha=1}^{d}\frac{\partial}{\partial x_{\alpha}}\left(\sum_{\beta=1}^{d}g^{\alpha\beta}\sqrt{G}\frac{\partial\psi}{\partial x_{\beta}}\right)(\hat{x}) + b\underline{u}(\hat{x}) \leq 0\\ (\textit{resp.} \quad & -\frac{1}{\sqrt{G}}\sum_{\alpha=1}^{d}\frac{\partial}{\partial x_{\alpha}}\left(\sum_{\beta=1}^{d}g^{\alpha\beta}\sqrt{G}\frac{\partial\psi}{\partial x_{\beta}}\right)(\hat{x}) + b\bar{u}(\hat{x}) \geq 0), \end{split}$$

where L is expressed as (2.5) in a local chart containing \hat{x} .

REMARK 2.8. There is also a notion of subsolutions and supersolutions in the sense of distribution. It is actually equivalent to that in the sense of viscosity (see e.g. [35, Appendix], [31], and [26]).

Consider two domain decomposition iterative schemes as follows: Choose arbitrarily an initial guess $\underline{u}^0 \in C^0(M)$ (resp. $\overline{u}^0 \in C^0(M)$) with $\underline{u}^0|_{\partial M} = \overline{u}^0|_{\partial M} = 0$ such that \underline{u}^0 (resp. \overline{u}^0) is a subsolution (resp. supersolution) of (2.4). Let $\underline{u}^0_i = \underline{u}^0|_{M_i}$. For each $n \geq 1$, define

(2.6)
$$\begin{cases} L\underline{u}_{i}^{n}(x) = 0, & x \in M_{i} \setminus \partial M_{i}; \\ \underline{u}_{i}^{n}(x) = \min\{\underline{u}_{j}^{n-1}(x) \mid x \in M_{j}, j \neq i\}, & x \in \gamma_{i}; \\ \underline{u}_{i}^{n}(x) = 0, & x \in \partial M_{i} \cap \partial M. \end{cases}$$

Similarly, let $\bar{u}_i^0 = \bar{u}^0|_{M_i}$. For each $n \ge 1$, define

(2.7)
$$\begin{cases} L\bar{u}_i^n(x) = 0, & x \in M_i \setminus \partial M_i; \\ \bar{u}_i^n(x) = \max\{\bar{u}_j^{n-1}(x) \mid x \in M_j, j \neq i\}, & x \in \gamma_i; \\ \bar{u}_i^n(x) = 0, & x \in \partial M_i \cap \partial M. \end{cases}$$

Here $\underline{u}_i^n|_{\partial M_i}$ and $\overline{u}_i^n|_{\partial M_i}$ are not necessarily continuous. We take \underline{u}_i^n and \overline{u}_i^n as the Perron solutions with Dirichlet boundary value on M_i (see [25, Theorem 6.11]). The goal of this subsection is to prove Proposition 2.11 below.

In the following, all arguments cover both the cases $\partial M \neq \emptyset$ and $\partial M = \emptyset$. In the case of $\partial M = \emptyset$, the proof is even simpler. All statements related to ∂M and $M_i \cap \partial M$ are vacuously true if $\partial M = \emptyset$ and $M_i \cap \partial M = \emptyset$ respectively.

LEMMA 2.9. $\{\underline{u}_i^n\}$ (resp. $\{\overline{u}_i^n\}$) is an increasing (resp. decreasing) sequence of solutions to $L\hat{u} = 0$ with an upper (resp. lower) bound 0. Furthermore, $\{\underline{u}_i^n\}$ and $\{\bar{u}_i^n\}$ are continuous on $M_i \setminus (\gamma_i \cap S)$. Here S is the set defined in Assumption 2.1.

Proof. We only prove the case of $\{\underline{u}_i^n\}$ since the proof of the other case is similar. The argument is an induction on n. The conclusion is true for n = 0 by assumption.

Firstly, since $L\underline{u}_i^1 = 0$, $\underline{u}_i^1|_{\partial M_i} = \underline{u}^0$ is continuous, and ∂M_i is Lipschitz, we have $\underline{u}_i^1 \in C^0(M_i)$. By the assumption that \underline{u}^0 is a subsolution and $\underline{u}^0|_{\partial M} = 0$, we infer $\underline{u}^0 \leq 0$, which further implies $\underline{u}_i^0 = \underline{u}^0|_{M_i} \leq 0$. Since \underline{u}_i^1 is a solution and $\underline{u}_i^0|_{\partial M_i} \leq 0$, we have $\underline{u}_i^0 \leq \underline{u}_i^1 \leq 0$. The conclusion is verified for n = 1. Suppose the conclusion is true for $n \leq k-1$. By inductive hypothesis, $\forall i, \underline{u}_i^{k-2} \leq \underline{u}_i^{k-1} \leq 0$ and \underline{u}_i^{k-1} is continuous in $M_i \setminus (\gamma_i \cap S)$.



FIG. 1. An illustration of domain decomposition.

We claim $\underline{u}_i^k|_{\partial M_i}$ is continuous on $\partial M_i \setminus (\gamma_i \cap S)$. Note that

(2.8)
$$\partial M_i \setminus S = \partial M_i \setminus (\gamma_i \cap S) = (\partial M_i \cap \partial M) \cup (\gamma_i \setminus S).$$

Assuming $x \in \partial M_i \setminus S$ and $y \in \partial M_i$, we need to show $\underline{u}_i^k(y)$ tends to $\underline{u}_i^k(x)$ when y tends to x. Suppose firstly $x \in \partial M_i \cap \partial M$. (See Fig. 1 for an illustration, where M is decomposed into three subdomains and the shadowed part is M_i .) By the boundary value condition, $\underline{u}_i^k(x) = \underline{u}_i^k|_{\partial M} = 0$. Thus $\underline{u}_i^k(y)$ tends to 0 when $y \in \partial M_i \cap \partial M$ (see the left hand side of Fig. 1). On the other hand, $\forall y \in \gamma_i$ (see the right hand side of Fig. 1),

(2.9)
$$\underline{u}_i^k(y) = \min\{\underline{u}_j^{k-1}(y) \mid y \in M_j, j \neq i\}.$$

and $\underline{u}_{j}^{k-1}|_{\partial M} = 0$ for all j. Since these \underline{u}_{j}^{k-1} are continuous at x, we also infer $\underline{u}_{i}^{k}(y)$ tends to 0 when $y \in \gamma_{i}$. In summary, $\underline{u}_{i}^{k}|_{\partial M_{i}}$ is continuous at $x \in \partial M_{i} \cap \partial M$. Let's

assume $x \in \gamma_i \setminus S$ further. (See Fig. 2 for an illustration.) There exists a neighborhood U_x of x such that, $\forall y \in U_x \cap \partial M_i$, we have $y \notin \partial M \cup S$ and y belongs to the same subdomains M_j in (2.9) as x does. By (2.9) and the continuity of \underline{u}_j^{k-1} at x again, $\underline{u}_i^k|_{\partial M_i}$ is continuous at x. Overall, our claim is proved by (2.8).



FIG. 2. An illustration of domain decomposition.

Therefore, \underline{u}_i^k is continuous in $M_i \setminus (\gamma_i \cap S)$ since it is a solution and ∂M_i is Lipschitz. We also infer $\underline{u}_i^{k-1}|_{\partial M_i} \leq \underline{u}_i^k|_{\partial M_i}$ by the fact $\underline{u}_j^{k-2} \leq \underline{u}_j^{k-1}$ for all j. This further yields $\underline{u}_i^{k-1} \leq \underline{u}_i^k$. Since $\underline{u}_j^{k-1} \leq 0$ for all j, we also have $\underline{u}_i^k|_{\partial M_i} \leq 0$ which implies $\underline{u}_i^k \leq 0$. The conclusion holds for n = k.

LEMMA 2.10. $\{\underline{u}_i^n\}$ (resp. $\{\overline{u}_i^n\}$) converges to a function $\underline{u}_i \leq 0$ (resp. $\overline{u}^i \geq 0$). The convergence is uniform on each $K \subset M_i \setminus (\gamma_i \cap S)$. (Here " $\subset \subset$ " means K is a compact subset.) Furthermore, \underline{u}_i and \overline{u}^i vanish on ∂M and are continuous on $M_i \setminus (\gamma_i \cap S)$.

Proof. We only prove the case of $\{\underline{u}_i^n\}$. By Lemma 2.9, $\{\underline{u}_i^n\}$ converges increasingly to a function $\underline{u}_i \leq 0$. Clearly, $\underline{u}_i|_{\partial M} = 0$ since so do all \underline{u}_i^n . Lemma 2.9 indicates the continuity of \underline{u}_i^n on $M_i \setminus (\gamma_i \cap S)$. By Dini's Theorem, the uniform convergence on K would follow from the continuity of \underline{u}_i . So we only need to prove \underline{u}_i is continuous on $M_i \setminus (\gamma_i \cap S)$.

As these \underline{u}_i^n are solutions, by Harnack's inequality, the monotone convergence of $\{\underline{u}_i^n\}$ is uniform on each $K' \subset \subset M_i \setminus \partial M_i$ which implies the continuity of \underline{u}_i on $M_i \setminus \partial M_i$. It remains to show \underline{u}_i is continuous at each $x \in \partial M_i \setminus (\gamma_i \cap S)$. By (2.8), we may check the continuity of \underline{u}_i at $x \in \gamma_i \setminus S$ and at $x \in \partial M_i \cap \partial M$ separately.

We firstly claim $\underline{u}_i|_{\partial M_i}$ is continuous at each $x \in \gamma_i \setminus S$. (See Fig. 2 for an illustration.) Suppose $y \in \partial M_i$. Since $x \notin S$, we see $y \notin \partial M \cup S$ when y is close enough to x. In other words, $y \notin \partial M$ and $y \notin \partial M_j$ for all $j \neq i$. Thus, when $y \in \partial M_i$ is near x, we have y belongs to the same subdomains M_j as x does and

$$\underline{u}_i^n(y) = \min\{\underline{u}_j^{n-1}(y) \mid y \in M_j, j \neq i\} = \min\{\underline{u}_j^{n-1}(y) \mid y \in M_j \setminus \partial M_j, j \neq i\}.$$

Since $\{\underline{u}_j^n\}$ converges uniformly on $K' \subset M_j \setminus \partial M_j$, we also have $\underline{u}_i^n|_{\partial M_i}$ converges uniformly near x. By Lemma 2.9, \underline{u}_i^n is continuous near x. Now the continuity of $\underline{u}_i|_{\partial M_i}$ at x follows from that of $\underline{u}_i^n|_{\partial M_i}$ at x.

Let \tilde{u}_i be the Perron solution on M_i with $\tilde{u}_i|_{\partial M_i} = \underline{u}_i|_{\partial M_i}$. By the fact, $\forall n$, \underline{u}_i^n is also a solution and $\underline{u}_i^n|_{\partial M_i} \leq \underline{u}_i|_{\partial M_i} = \tilde{u}_i|_{\partial M_i}$, we have $\underline{u}_i^n \leq \tilde{u}_i$ and hence $\underline{u}_i^n \leq \underline{u}_i \leq \tilde{u}_i$. Let $x \in \gamma_i \setminus S$. Then \tilde{u}_i is continuous at x since so is $\tilde{u}_i|_{\partial M_i}$. For each $\epsilon > 0$, we have $0 \leq \tilde{u}_i(x) - \underline{u}_i^N(x) < \epsilon$ for some N since $\{\underline{u}_i^n(x)\}$ converges to $\tilde{u}_i(x)$. Furthermore, there exists a neighborhood U_x of x such that $\forall y \in U_x \cap M_i$,

$$|\underline{u}_i^N(x) - \underline{u}_i^N(y)| < \epsilon$$
 and $|\tilde{u}_i(x) - \tilde{u}_i(y)| < \epsilon$

because \underline{u}_i^N and \tilde{u}_i are continuous at x. Thus

$$\underline{u}_i(x) - 2\epsilon = \tilde{u}_i(x) - 2\epsilon < \underline{u}_i^N(x) - \epsilon < \underline{u}_i^N(y)$$

$$\leq \underline{u}_i(y) \leq \tilde{u}_i(y) < \tilde{u}_i(x) + \epsilon = \underline{u}_i(x) + \epsilon$$

i.e. $\underline{u}_i(x) - 2\epsilon < \underline{u}_i(y) < \underline{u}_i(x) + \epsilon$. So \underline{u}_i is continuous at each $x \in \gamma_i \setminus S$.

Finally, suppose $x \in \partial M_i \cap \partial M$. Since \underline{u}_i^0 is continuous, $\forall \epsilon > 0$, there exists a neighborhood U_x of x such that $\forall y \in U_x \cap M_i$, $\underline{u}_i^0(x) - \epsilon < \underline{u}_i^0(y)$. We already know $\underline{u}_i^0 \leq \underline{u}_i \leq 0$ and $\underline{u}_i^0|_{\partial M} = \underline{u}_i|_{\partial M} = 0$. Therefore, $\underline{u}_i(x) = \underline{u}_i^0(x) = 0$ and

$$-\epsilon = \underline{u}_i^0(x) - \epsilon < \underline{u}_i^0(y) \le \underline{u}_i(y) \le 0$$

We see $|\underline{u}_i(y) - \underline{u}_i(x)| < \epsilon$ and hence \underline{u}_i is continuous at $x \in \partial M_i \cap \partial M$. We finish the proof by (2.8).

PROPOSITION 2.11. $\{\underline{u}_i^n\}$ and $\{\overline{u}_i^n\}$ converge to 0 on $M_i \setminus (\gamma_i \cap S)$. The convergence is uniform on each $K \subset M_i \setminus (\gamma_i \cap S)$.

Proof. By Lemma 2.10, it suffices to show $\underline{u}_i = \overline{u}_i = 0$ on $M_i \setminus (\gamma_i \cap S)$. We only prove the case of \overline{u}_i since the proof of the other case is similar.

We already know these \bar{u}_i^n are solutions. By the uniform convergence on $K \subset \subset M_i \setminus (\gamma_i \cap S)$, we know that \bar{u}_i is a solution in $M_i \setminus \partial M_i$. Furthermore, $\forall x \in \gamma_i$,

$$\bar{u}_i^n(x) = \max\{\bar{u}_j^{n-1}(x) \mid x \in M_j, j \neq i\}.$$

We infer, $\forall x \in \gamma_i$,

(2.10)
$$\bar{u}_i(x) = \max\{\bar{u}_j(x) \mid x \in M_j, j \neq i\} = \max\{\bar{u}_j(x) \mid x \in M_j\}.$$

Define a function w on $M \setminus S$ as

(2.11)
$$w(y) = \max\{\bar{u}_j(y) \mid y \in M_j\}$$

We shall prove that w = 0. Clearly, $w|_{\partial M} = 0$ and $0 \le w \le \bar{u}^0$ since $\bar{u}_j|_{\partial M} = 0$ and $0 \le \bar{u}_j \le \bar{u}^0$ for all j.

We firstly claim that w is continuous on $M \setminus S$. Note that

$$M \setminus S = \partial M \cup \left[M \setminus \left(\partial M \cup \bigcup_{i=1}^m \gamma_i \right) \right] \cup \bigcup_{i=1}^m (\gamma_i \setminus S).$$

The continuity at each $x \in \partial M$ is obvious since $\bar{u}_j(x) = 0$ and \bar{u}_j are continuous at x for all j. If $x \notin \partial M \cup \bigcup_{i=1}^m \gamma_i$, there exists a neighborhood U_x of x such that, $\forall y \in U_x$, y belongs to the same subdomains as x does. The continuity of w at x follows from that of all \bar{u}_j . Now suppose $x \in \gamma_i \setminus S$. There exists a neighborhood U_x of x such that γ_i divides U_x into two sides. Furthermore, The points y on one side belong to exactly one more subdomain, i.e. M_i , than the points on the other side do. (See Fig. 3 for an illustration, where y_1 and y_2 are on different sides of γ_i . Both x and y_1 belong to M_i , while y_2 does not.) The continuity of w at $x \in \gamma_i \setminus S$ follows from the continuity of all \bar{u}_j together with (2.10).

Secondly, we show that w is a subsolution of (2.4) in $M \setminus S$. It suffices to check w is a subsolution locally in $M \setminus (\partial M \cup S)$ since w is continuous on $M \setminus S$ and $M \setminus (\partial M \cup S)$ is the interior of $M \setminus S$. This fact is obvious in $M \setminus (\partial M \cup \bigcup_{i=1}^{m} \gamma_i)$. By (2.11), w is the maximum of a fixed family of \bar{u}_j near each $x \in M \setminus (\partial M \cup \bigcup_{i=1}^{m} \gamma_i)$.



FIG. 3. An illustration of domain decomposition.

The fact follows because all \bar{u}_j are solutions. By an argument similar to the proof of the (iii) in [25, p. 103], we also see w is a subsolution near each $x \in \gamma_i \setminus S$. More precisely, near x, the family of \bar{u}_j in (2.11) on one side of γ_i has one more member, i.e. \bar{u}_i , than the family on the other side does. (See also Fig. 3 for an illustration.) So locally, w can be viewed as a lifting of a subsolution on the first side which is again a subsolution.

By Assumption 2.1, the (d-2)-dimensional Hausdorff measure of S is finite. We also know w is bounded. Thus by [27, Theorem 12.1 (c)], there is a canonical way (see [27, (3.1)]) to extend w to be a subsolution of (2.4) on M, i.e. $w|_{M\setminus\partial M}$ satisfies Definition 2.7 and w is continuous at each $x \in \partial M$. Now we claim the extended $w \leq 0$. If $\partial M \neq \emptyset$, by the fact that $w|_{\partial M} = 0$ and w is a subsolution, we see $w \leq 0$ on M. On the other hand, supposing $\partial M = \emptyset$, we have b > 0 in (2.5) by assumption. Since w is upper semicontinuous and M is compact, w has a global maximum point \hat{x} . This \hat{x} is in $M \setminus \partial M$ since $\partial M = \emptyset$. Taking $\underline{u} = w$ and $\psi = 0$ in Definition 2.7, we infer $w(\hat{x}) \leq 0$. Therefore, $w \leq 0$ on M.

However, $w|_{M\setminus S} \ge 0$, thus $w|_{M\setminus S} = 0$. Since $0 \le \bar{u}^i|_{M_i\setminus S} \le w|_{M_i\setminus S}$, we obtain $\bar{u}^i|_{M_i\setminus S} = 0$. Thus $\bar{u}^i|_{M_i\setminus(\gamma_i\cap S)} = 0$ by the continuity of $\bar{u}^i|_{M_i\setminus(\gamma_i\cap S)}$.

2.3. Proof of Convergence. Again, in the following, all arguments cover both the case $\partial M \neq \emptyset$ and $\partial M = \emptyset$. All statements related to ∂M are vacuously true if $\partial M = \emptyset$. In addition, $C_0^0(M) = C^0(M)$ if $\partial M = \emptyset$.

LEMMA 2.12. Let u^0 be the initial guess in Algorithm 2.1. Suppose further u and u^0 are Lipschitz continuous if $\partial M \neq \emptyset$. Then there exist a subsolution $\underline{u}^0 \in C_0^0(M)$ and a supersolution $\overline{u}^0 \in C_0^0(M)$ of (2.4) such that $\underline{u}^0 \leq u^0 - u \leq \overline{u}^0$.

Proof. Firstly, let's assume $\partial M \neq \emptyset$. Clearly, the boundary value problem

$$\begin{cases} Lv = 1, & \text{in } M, \\ v = 0, & \text{on } \partial M, \end{cases}$$

has a unique solution $v \in C^{\infty}(M)$. By Hopf's Lemma and strong maximum principle ([25, § 3.2]), we have $\frac{\partial v}{\partial \nu} < 0$ and $v|_{M \setminus \partial M} > 0$, where $\frac{\partial v}{\partial \nu}$ is the outer normal derivative of v on ∂M . Since $u^0 - u$ is Lipschitz and $(u^0 - u)|_{\partial M} = 0$, there exists an open neighborhood V of ∂M such that $|u^0 - u| \leq kv$ in V for sufficiently large constant k. Furthermore, $v|_{M \setminus V}$ has a positive minimum. We see $|u^0 - u| \leq kv$ on the whole M for sufficiently large k. We can define $\underline{u}^0 = -kv$ and $\overline{u}^0 = kv$ for some large k.

On the other hand, suppose $\partial M = \emptyset$, then we have b > 0 in (2.2) as assumed. Since $u^0 - u$ is continuous and hence bounded on M. We can easily define $\underline{u}^0 = -k$ and $\bar{u}^0 = k$ for some large constant k.

LEMMA 2.13. Let $\{\underline{u}_i^n\}$ (resp. $\{\overline{u}_i^n\}$) be the sequence generated in (2.6) (resp. (2.7)) with \underline{u}^0 (resp. \overline{u}^0) satisfying the conclusion of Lemma 2.12. Then, $\forall n, \forall i, \underline{u}_i^n \leq u_i^n - u \leq \overline{u}_i^n$.

Proof. We only prove $\underline{u}_i^n \leq u_i^n - u$ because the proof of $u_i^n - u \leq \overline{u}_i^n$ is similar. Since $\underline{u}_i^0 = \underline{u}^0|_{M_i}$ and $u_i^0 = u^0|_{M_i}$, the conclusion is true for n = 0.

Suppose the conclusion is true for n-1. Since $L\underline{u}_i^n = L(u_i^n - u) = 0$, it suffices to prove that, $\forall x \in \partial M_i, \underline{u}_i^n(x) \leq u_i^n(x) - u(x)$. This is certainly true for $x \in \partial M$ since $\underline{u}_i^n(x) = u_i^n(x) - u(x) = 0$. For $x \in \partial M_i \setminus \partial M = \gamma_i$, by inductive hypothesis,

$$\underline{u}_{i}^{n}(x) = \min\{\underline{u}_{j}^{n-1}(x) \mid x \in M_{j}, j \neq i\} \leq \min\{u_{j}^{n-1}(x) - u(x) \mid x \in M_{j}, j \neq i\}$$
$$\leq \sum_{j=1}^{m} \rho_{j}(x)[u_{j}^{n-1}(x) - u(x)] = u_{i}^{n}(x) - u(x).$$

Note that we have used the fact $\rho_i(x) = 0$ when $x \in \gamma_i$ and $\rho_j(x) = 0$ when $x \notin M_j$ in the last inequality. The proof is finished.

Now we are ready to prove Theorem 2.4.

Proof of Theorem 2.4. (1) It suffices to prove $\lim_{n\to\infty} ||u_i^n - u||_{C^0(M_i)} = 0$ for each *i*. Since $L(u_i^n - u) = 0$, by the maximum principle, we only need to show that $\{(u_i^n - u)|_{\partial M_i}\}$ converges uniformly to 0. Let $\{\underline{u}_j^n\}$ and $\{\overline{u}_j^n\}$ be the sequences stated in Lemma 2.13. By Assumption 2.2, we have $\operatorname{supp} \rho_j \subset M_j \setminus \overline{\gamma_j}$. By Proposition 2.11, $\forall j, \{\underline{u}_j^n\}$ and $\{\overline{u}_j^n\}$ converge uniformly to 0 on $\operatorname{supp} \rho_j$. By Lemma 2.13, so does $\{u_j^n - u\}$. Since $(u_i^n - u)|_{\partial M_i} = \sum_{j=1}^m \rho_j(u_j^{n-1} - u)$, we infer $\{(u_i^n - u)|_{\partial M_i}\}$ converges uniformly to 0.

(2) By the fact $u^n - u = \sum_{i=1}^m \rho_i(u_i^n - u)$, it's obvious to see $||u^n - u||_{C^0(M)} \le \theta^n$ and $||u^n - u||_{L^2(M)} \le C_3 \theta^n$ for some constant C_3 independent of n and i.

Secondly, we claim that $\forall K_i \subset \subset M_i \setminus \overline{\gamma_i}$, there exists a constant $C_4(K_i)$ independent of n such that, $\forall n$,

(2.12)
$$\|\nabla u_i^n - \nabla u\|_{L^2(K_i)} \le C_4(K_i) \|u_i^n - u\|_{C^0(M_i)}.$$

Since $L(u_i^n - u) = 0$ in $M_i \setminus \partial M_i$ and $(u_i^n - u)|_{\partial M} = 0$, the claim (2.12) follows from the regularity theory for elliptic equations (cf. [23, p. 331] and [23, p. 340]).

Furthermore, since ρ_i is Lipschitz continuous, we have $\rho_i \in W^{1,\infty}(M)$ (see Theorem 5 in [24, §4.2.3]) and

$$\nabla(u^n - u) = \sum_{i=1}^m \nabla \rho_i \cdot (u_i^n - u) + \sum_{i=1}^m \rho_i \cdot \nabla(u_i^n - u),$$

which implies

$$\|\nabla(u^n - u)\|_{L^2(M)} \le \sum_{i=1}^m \|\nabla\rho_i\|_{L^\infty(M_i)} \cdot \|u_i^n - u\|_{L^2(M_i)} + \sum_{i=1}^m \|\nabla u_i^n - \nabla u\|_{L^2(\mathrm{supp}\rho_i)}.$$

Clearly, $||u_i^n - u||_{L^2(M_i)} \leq C_5 \theta^n$ for some constant C_5 independent of n and i. By (2.12) and the fact $\operatorname{supp} \rho_i \subset \subset M_i \setminus \overline{\gamma_i}$, we also have

$$\|\nabla u_i^n - \nabla u\|_{L^2(\mathrm{supp}\rho_i)} \le C_4(\mathrm{supp}\rho_i) \cdot \theta^n.$$

Now the conclusion $||u^n - u||_{H^1(M)} \leq C_1 \theta^n$ follows.

(3) As
$$L(u_i^n - u) = 0$$
 and $u_i^n|_{\partial M_i} = u^{n-1}|_{\partial M_i}$, we have $u_i^n - u^{n-1} \in H_0^1(M_i)$ and
 $L(u_i^n - u^{n-1}) = L(u_i^n - u) + L(u - u^{n-1}) = L(u - u^{n-1}).$

By the Poincaré inequality and the fact $b \ge 0$ in (2.2), there exist positive constants C_6 and C_7 independent of n and i such that

$$C_{6} \|u_{i}^{n} - u^{n-1}\|_{H^{1}(M_{i})}^{2} \leq (L(u_{i}^{n} - u^{n-1}), u_{i}^{n} - u^{n-1})_{M_{i}}$$

= $(L(u - u^{n-1}), u_{i}^{n} - u^{n-1})_{M_{i}} \leq \|L(u^{n-1} - u)\|_{H^{-1}(M_{i})} \cdot \|u_{i}^{n} - u^{n-1}\|_{H^{1}(M_{i})}$
 $\leq C_{7} \|u^{n-1} - u\|_{H^{1}(M_{i})} \cdot \|u_{i}^{n} - u^{n-1}\|_{H^{1}(M_{i})},$

where $(\cdot, \cdot)_{M_i}$ is the dual pairing between $H^{-1}(M_i)$ and $H^1_0(M_i)$. Thus

$$||u_i^n - u^{n-1}||_{H^1(M_i)} \le C_6^{-1}C_7||u^{n-1} - u||_{H^1(M_i)}$$

and

$$||u_i^n - u||_{H^1(M_i)} \le (1 + C_6^{-1}C_7)||u^{n-1} - u||_{H^1(M_i)}.$$

Now $||u_i^n - u||_{H^1(M_i)} \le C_2 \theta^{n-1}$ follows from the fact $||u^{n-1} - u||_{H^1(M)} \le C_1 \theta^{n-1}$.

3. Numerical Scheme. In this section, we propose a numerical DDM iteration procedure (Algorithm 3.1 below) to obtain approximations to the solution to (2.2). The idea is as follows. We decompose M into overlapping subdomains M_i satisfying Assumption 2.1. Furthermore, each M_i is required in a coordinate chart. Then apply a DDM iteration procedure similar to Algorithm 2.1. Since M_i is in a coordinate chart, an elliptic problem on M_i is converted to one on a domain in a Euclidean space and then can be solved approximately by usual finite element methods. The transition of information among subdomains is by interpolation.

Unlike the numerical DDM procedure in [10, Algorithm 3.1], the iteration of this algorithm is highly parallel. As a result, this algorithm is of particular interest when the number of subdomains is large.

3.1. Finite Element Spaces over a d-Rectangle. Our numerical algorithm is based on finite element methods. Suppose a manifold M has dimension d. As indicated above, each subdomain M_i of M will be converted to a domain $D_i \subset \mathbb{R}^d$. To minimize the difficulty of coding, as in [10], we shall choose D_i as a d-rectangle and use the multilinear finite element space of d-rectangles, i.e. the space of d-rectangle of type (1) in [11, p. 56-64].

Recall that a d-rectangle D is

$$D = \prod_{i=1}^{d} [a_i, b_i] = \{ (x_1, \cdots, x_d) \mid \forall i, x_i \in [a_i, b_i] \}.$$

We refine each coordinate factor interval $[a_i, b_i]$ by adding points of partition:

$$a_i = c_{i,0} < c_{i,1} < \dots < c_{i,N_i} = b_i.$$

Then $[a_i, b_i]$ is divided into N_i subintervals. The refinement of all such $[a_i, b_i]$ provides a grid on D. This divides D into $\prod_{i=1}^d N_i$ many small d-rectangles

(3.1)
$$\prod_{i=1}^{d} [c_{i,t_i-1}, c_{i,t_i}],$$

12

where $1 \le t_i \le N_i$ for all *i*. Each small *d*-rectangle (3.1) is an element of the grid. We refer to [10, § 3.1] (see also [11, p. 56-64]) for a detailed description of the finite element space of *d*-rectangles of type (1) based on such a grid.

3.2. Numerical Algorithm. Let M be a d-dimensional compact Riemannian manifold with or without boundary. We shall propose a numerical DDM based on finite element methods to find numerical solutions to (2.2)

Suppose $M = \bigcup_{i=1}^{m} (M_i \setminus \overline{\gamma_i})$ as Assumption 2.1. Assume further there is a smooth diffeomorphism $\phi_i : D_i \to M_i$ for each *i*, where D_i is a *d*-rectangle in \mathbb{R}^d . Theoretically, we can always get such triples (M_i, D_i, ϕ_i) . The reason has been explained in [10, § 3.2] provided that $\partial M = \emptyset$. On the contrary, if $\partial M \neq \emptyset$, special technique should be employed to deal with the boundary. We shall explain this technique in Section 4.

We also need a partition of unity $\{\rho_i \mid 1 \leq i \leq m\}$ satisfying Assumption 2.2. Such ρ_i can be constructed as follows. Choose nonnegative Lipschitz continuous functions σ_i on M such that $\operatorname{supp} \sigma_i \subset M_i \setminus \overline{\gamma_i}$ and $\sum_{i=1}^m \sigma_i > 0$ on M. We define desired ρ_i as

(3.2)
$$\rho_i = \frac{\sigma_i}{\sum_{j=1}^m \sigma_j}.$$

To obtain σ_i , it suffices to define $\sigma_i \circ \phi_i$ which is a function on D_i and hence can be defined simply and quite arbitrarily in terms of elementary functions.

Our numerical algorithm is a discrete imitation of Algorithm 2.1. Since the numerical procedure is based on finite element methods, it's necessary to convert the first line of (2.3) to its weak form: $\forall v \in H_0^1(M_i)$,

$$\int_{M_i} (\langle \nabla u_i^n, \nabla v \rangle + b u_i^n v) \operatorname{dvol} = \int_{M_i} f v \operatorname{dvol}.$$

Here ∇u_i^n and ∇v are the gradients of u_i^n and v with respect to the Riemannian metric $g, \langle \nabla u_i^n, \nabla v \rangle$ is the inner product of ∇u_i^n and ∇v , and dvol is the volume form. In terms of the local coordinates (x_1, \ldots, x_d) provided by $\phi_i : D_i \to M_i$,

$$\langle \nabla u_i^n, \nabla v \rangle = \sum_{\alpha, \beta=1}^d g^{\alpha\beta} \frac{\partial u_i^n}{\partial x_\alpha} \frac{\partial v}{\partial x_\beta} \quad \text{and} \quad \text{dvol} = \sqrt{G} \mathrm{d}x_1 \cdots \mathrm{d}x_d.$$

Thus we define an energy bilinear form on $H^1(D_i)$ as

$$a_i(w,v) = \int_{D_i} \left(\sum_{\alpha,\beta=1}^d g^{\alpha\beta} \frac{\partial w}{\partial x_\alpha} \frac{\partial v}{\partial x_\beta} + bwv \right) \sqrt{G} \mathrm{d}x_1 \cdots \mathrm{d}x_d.$$

Define a bilinear form $(\cdot, \cdot)_i$ on $L^2(D_i)$ as

$$(w,v)_i = \int_{D_i} wv \sqrt{G} \mathrm{d}x_1 \cdots \mathrm{d}x_d.$$

The first line of (2.3) is converted to an equation on D_i in weak form: $\forall v \in H_0^1(D_i)$,

(3.3)
$$a_i(u_i^n \circ \phi_i, v) = (f \circ \phi_i, v)_i.$$

Create a grid of *d*-rectangles over D_i . Let $V_{h,i}$ be the finite element space of *d*-rectangles of type (1) over D_i . A discrete imitation of the first line of (2.3) would be: find a $u_{h,i}^n \in V_{h,i}$ such that, $\forall v_h \in V_{h,i} \cap H_0^1(D_i)$,

$$a_i(u_{h,i}^n, v_h) = (f \circ \phi_i, v_h)_i.$$

However, this discrete problem is not well-posed because the degrees of freedom of $u_{h,i}^n$ on ∂D_i are undetermined. As an imitation of the second line of (2.3), we should evaluate these degrees of freedom by the data in *d*-rectangles D_j for $j \neq i$. So we have to investigate the transitions of coordinates.

For $i \neq j$, let $D_{ij} = \phi_i^{-1}(M_i \cap M_j) \subseteq D_i$ and $D_{ji} = \phi_j^{-1}(M_i \cap M_j) \subseteq D_j$. Then

$$\phi_i^{-1} \circ \phi_i : D_{ij} \to D_{ji}$$

is a diffeomorphism which is the transition of coordinates on the overlapping between M_i and M_j . We refer to [10, § 3.2] for a detailed explanation of the transition maps. (Particularly, Fig. 1 therein provides an illustration.) We still wish to point out that $\phi_j^{-1} \circ \phi_i$ preserves neither nodes nor grid necessarily. In other words, $\phi_j^{-1} \circ \phi_i$ may neither map a node in D_{ij} to a node in D_{ji} , nor map the grid over D_{ij} to the one over D_{ji} . As emphasized in [10, § 3.2], this incompatibility among the grids over different D_i actually shows the high flexibility of our approach.

Now we are in a position to propose our numerical Algorithm 3.1. Define

$$V_h = \bigoplus_{i=1}^m V_{h,i}.$$

Algorithm 3.1 A numerical DDM to solve (2.2).

- 1: Choose an arbitrary initial guess $u_h^0 = (u_{h,1}^0, \cdots, u_{h,m}^0) \in V_h$ such that, $\forall i, u_{h,i}^0(\xi) = \varphi(\phi_i(\xi))$ if $\xi \in \partial D_i$ is a node and $\phi_i(\xi) \in \partial M$.
- 2: For each n > 0, assuming u_h^{n-1} has been obtained, for $1 \le i \le m$, find a $u_{h,i}^n \in V_{h,i}$ as follows. Suppose $\xi \in \partial D_i$ is a node. If $\phi_i(\xi) \in \partial M$, let $u_{h,i}^n(\xi) = \varphi(\phi_i(\xi))$. Otherwise, define

(3.4)
$$u_{h,i}^{n}(\xi) = \sum_{j=1}^{m} \rho_j(\phi_i(\xi)) \cdot u_{h,j}^{n-1}(\phi_j^{-1} \circ \phi_i(\xi)).$$

The interior degrees of freedom of $u_{h,i}^n$ are determined by $\forall v_h \in V_{h,i} \cap H_0^1(D_i)$,

$$a_i(u_{h,i}^n, v_h) = (f \circ \phi_i, v_h)_i$$

3: Define $u_h^n = (u_{h,1}^n, \cdots, u_{h,m}^n) \in V_h$.

The computation in (3.4) needs more explanation. If $\phi_i(\xi) \notin M_j$, then $\rho_j(\phi_i(\xi)) = 0$ and the summand $\rho_j(\phi_i(\xi)) \cdot u_{h,j}^{n-1}(\phi_j^{-1} \circ \phi_i(\xi))$ is actually not needed. On the contrary, $\phi_i(\xi) \in M_j$,

$$\rho_j(\phi_i(\xi)) = (\rho_j \circ \phi_j)(\phi_j^{-1} \circ \phi_i(\xi)).$$

Here $\rho_j \circ \phi_j$ is a function on D_j with an explicit formula. Thus $\rho_j(\phi_i(\xi))$ can be obtained once we get the coordinates of $\phi_j^{-1} \circ \phi_i(\xi)$. Similarly, $u_{h,j}^{n-1}(\phi_j^{-1} \circ \phi_i(\xi))$ can also be evaluated. Since $\phi_j^{-1} \circ \phi_i(\xi)$ is not necessarily a node, the evaluation $u_{h,j}^{n-1}(\phi_j^{-1} \circ \phi_i(\xi))$ is essentially an interpolation of the degrees of freedom of $u_{h,j}^{n-1}$.

Now the u_h^n in Algorithm 3.1 is the *n*th iterated numerical approximation to the solution to (2.2). Unlike the u^n in Algorithm 2.1 which is globally defined on M, the

 u_h^n consists of a group of numerical solutions $u_{h,i}^n$ on D_i for $1 \le i \le m$. In fact, the $u_{h,i}^n$ in Algorithm 3.1 is a numerical imitation of the u_i^n in Algorithm 2.1. As suggested by Theorem 2.4, $u_{h,i}^n$ should approximate $u \circ \phi_i$ well on D_i when the grid scale is small and n is large. Our numerical experiments show that it is indeed the case.

It's worth emphasizing that, for a fixed n, these $u_{h,i}^n$ $(1 \le i \le m)$ in Algorithm 3.1 can be solved independently. Thus Algorithm 3.1 is a highly parallel iterative procedure, which is remarkably different from that numerical method in [10]. The current method would have more advantages when the number of subdomains increases.

4. Manifolds with Boundary. We explain a technique to decompose M as $M = \bigcup_{i=1}^{m} (M_i \setminus \overline{\gamma_i})$ such that M_i can be parameterized by *d*-rectangles. For the case of $\partial M = \emptyset$, this has been explained in [10, § 3.2].

Now suppose $\partial M \neq \emptyset$. The difficulty occurs only in the part near the boundary. But this difficulty can be indeed overcome theoretically. By the Collar Neighborhood Theorem (see Theorem 6.1 in [28, Chapter 4]), there exists a smooth diffeomorphism

$$\psi: \ [\delta,\eta] \times \partial M \to V \subset M$$

such that V is a neighborhood of ∂M in M. Such a V is called a *collar neighborhood* of ∂M . One may even arrange $\psi(\{\delta\} \times \partial M) = \partial M$ or $\psi(\{\eta\} \times \partial M) = \partial M$ as one wishes. Let's assume $\psi(\{\eta\} \times \partial M) = \partial M$. (See Fig. 4 for an illustration, the disk therein is a manifold with boundary, the part with segments is a collar neighborhood $\psi((\delta, \eta] \times \partial M)$, each segment is parameterized by $(\delta, \eta]$.)



FIG. 4. An illustration of collar neighborhood.

For each $\zeta \in M \setminus \partial M$, there is an open chart neighborhood $U_{\zeta} \subseteq M \setminus \partial M$ of ζ , i.e. there is a diffeomorphism $\phi_{\zeta} : \Omega_{\zeta} \to U_{\zeta}$, where Ω_{ζ} is an open subset of \mathbb{R}^d . Since $\phi_{\zeta}^{-1}(\zeta)$ is an interior point of Ω_{ζ} , we can choose a rectangular neighborhood D_{ζ} of $\phi_{\zeta}^{-1}(\zeta)$ such that $D_{\zeta} \subseteq \Omega_{\zeta}$. This yields a diffeomorphism $\phi_{\zeta} : D_{\zeta} \to M_{\zeta} \subset U_{\zeta}$, where M_{ζ} is a neighborhood of ζ . Now

$$\{M_{\zeta} \setminus \partial M_{\zeta} \mid \zeta \in M \setminus \partial M\} \cup \{\psi((\delta, \eta] \times \partial M)\}$$

is an open covering of M. By the compactness of M, we obtain a finite subcovering

$$\{M_{\zeta_i} \setminus \partial M_{\zeta_i} \mid 1 \le i \le m_1\} \cup \{\psi((\delta, \eta] \times \partial M)\}.$$

Obviously, $\psi((\delta, \eta] \times \partial M)$ appears in the subcovering necessarily since these $M_{\zeta_i} \setminus \partial M_{\zeta_i}$ are contained in $M \setminus \partial M$. For brevity, let (M_i, D_i, ϕ_i) denote $(M_{\zeta_i}, D_{\zeta_i}, \phi_{\zeta_i})$. Since $M_i \subset M \setminus \partial M$, we have $\partial M_i = \partial M_i \setminus \partial M = \gamma_i$.

It remains to decompose $V = \psi([\delta, \eta] \times \partial M)$ into subdomains which can be parameterized by rectangles. Let's decompose $[\delta, \eta] \times \partial M$ firstly. Since ∂M is a manifold without boundary, we can find a desired decomposition of ∂M . In other

words, there are triples $\{(M'_j, D'_j, \phi'_j) \mid 1 \leq j \leq m_2\}$ such that D'_j are (d-1)rectangles and $\partial M = \bigcup_{j=1}^{m_2} (M'_j \setminus \partial M'_j)$. Note that $[\delta, \eta] \times \partial M$ is a product manifold. As pointed out in [10, § 4], decompositions of factor manifolds canonically result in a decomposition of their product. The situation of $[\delta, \eta] \times \partial M$ is even simpler because $[\delta, \eta]$ is already a 1-rectangle. Let $D_{m_1+j} = [\delta, \eta] \times D'_j$. Then all

$$\mathrm{id} \times \phi'_j: \ D_{m_1+j} = [\delta, \eta] \times D'_j \to [\delta, \eta] \times M'_j \subset [\delta, \eta] \times \partial M$$

provide a desired decomposition of $[\delta, \eta] \times \partial M$, where id is the identity on $[\delta, \eta]$. Finally, define

$$\phi_{m_1+j} = \psi \circ (\mathrm{id} \times \phi'_j) : \ D_{m_1+j} \to M.$$

Let $M_{m_1+j} = \phi_{m_1+j}(D_{m_1+j})$ and $\gamma_{m_1+j} = \partial M_{m_1+j} \setminus \partial M$. Then

$$M_{m_1+j} \setminus \overline{\gamma_{m_1+j}} = \phi_{m_1+j}((\delta,\eta] \times (D_{m_1+j} \setminus \partial D_{m_1+j})).$$

It's easy to see that $\{(M_i, D_i, \phi_i) \mid 1 \leq i \leq m_1 + m_2\}$ is a decomposition of M such that $M = \bigcup_{i=1}^{m_1+m_2} (M_i \setminus \overline{\gamma_i})$. We would like to mention that the transition map between D_{m_1+j} and D_{m_1+k}

has a simple expression. In fact,

$$\begin{split} \phi_{m_1+k}^{-1} \circ \phi_{m_1+j} &= (\psi \circ (\operatorname{id} \times \phi'_k))^{-1} \circ (\psi \circ (\operatorname{id} \times \phi'_j)) \\ &= (\operatorname{id} \times \phi'_k)^{-1} \circ \psi^{-1} \circ \psi \circ (\operatorname{id} \times \phi'_j) = (\operatorname{id} \times \phi'_k)^{-1} \circ (\operatorname{id} \times \phi'_j) = \operatorname{id} \times (\phi'_k^{-1} \circ \phi'_j). \end{split}$$

Here $\phi'_k^{-1} \circ \phi'_i$ is exactly a transition map for the decomposition of ∂M .

5. Numerical Experiments. We perform several numerical tests of Algorithm 3.1. The proposed method certainly applies to problems in all dimensions. However, the sizes of the linear systems derived from subdomains will increase exponentially with respect to the dimension. This difficulty is so called "the curse of dimensionality". Due to the constraint of computing resources, just as in [10], we only deal with problems with dimension no more than 4. Nevertheless, the numerical examples in [10] are all on manifolds without boundary. We now deal with manifolds with boundary too.

Our examples of manifolds are \mathbb{CP}^2 , $S^2 \times S^2$, B^4 and $B^2 \times S^2$. All of them are of dimension 4, the former two are without boundary, and the latter two are with boundary. It's also worth noting that, if a manifold can be decomposed into two subdomains, say the *d*-dimensional unit sphere S^d , the proposed method coincides essentially with the Schwarz Alternating Method and the method in [10]. As a result, we do not perform test on S^4 now since such an experiment has been done in [10].

5.1. A Problem on \mathbb{CP}^2 . Let $M = \mathbb{CP}^2$ be the complex projective plane. It is a compact complex manifold with complex dimension 2. Certainly, it can be considered as a real manifold with dimension 4.

In this example, the definition of \mathbb{CP}^2 , the Riemannian metric, the concrete problem of (2.2), the decomposition of \mathbb{CP}^2 , and the transition maps are the same as those in $[10, \S 5.2]$. We refer to [10] for a thorough and detailed description. Here we only briefly overview the decomposition.

Now \mathbb{CP}^2 is decomposed into 3 subdomains M_j for $0 \leq j \leq 2$. (Note that, following the convention of algebraic geometry, the index j is chosen to start from 0.) In the following, $z_j = x_j + \sqrt{-1}y_j \in \mathbb{C}, x_j \in \mathbb{R}$, and $y_j \in \mathbb{R}$. We shall identify the complex number z_j with the 2-dimensional real vector (x_j, y_j) . Let $D_j = [-r, r]^4 \subset \mathbb{R}^4 \simeq \mathbb{C}^2$. For $0 \leq j \leq 2$, we have the following diffeomorphisms

$$\phi_0: D_0 \to M_0 \subset \mathbb{CP}^2$$
$$(z_1, z_2) \mapsto [1, z_1, z_2],$$
$$\phi_1: D_1 \to M_1 \subset \mathbb{CP}^2$$
$$(z_0, z_2) \mapsto [z_0, 1, z_2],$$

and

$$\phi_2: D_2 \to M_2 \subset \mathbb{CP}^2$$
$$(z_0, z_1) \mapsto [z_0, z_1, 1],$$

where $[w_0, w_1, w_2]$ are called the *homogeneous coordinates* of \mathbb{CP}^2 . (For the definition of homogeneous coordinates, see [10].) To guarantee $\mathbb{CP}^2 = \bigcup_{j=0}^2 (M_j \setminus \partial M_j)$, we have to let r > 1. The larger r is, the more overlapping there will be.

Unlike the method in [10], we now need a partition of unity $\{\rho_j \mid 0 \leq j \leq 2\}$ satisfying Assumption 2.2. As mentioned before, it suffices to define the σ_j in (3.2), which is in turn reduced to the definition of $\sigma_j \circ \phi_j$ on D_j . Let r' = 0.9r + 0.1. Then 1 < r' < r. Suppose $x = (x_1, x_2, x_3, x_4) \in D_j = [-r, r]^4$. Define

(5.1)
$$\sigma_j \circ \phi_j(x) = \begin{cases} 0, & \max\{|x_j| \mid 1 \le j \le 4\} > r'; \\ \prod_{j=1}^4 (1 - (\frac{x_j}{r'})^2), & \text{otherwise.} \end{cases}$$

It's easy to check that $\operatorname{supp} \sigma_j = \phi_j([-r', r']^4) \subset M_j \setminus \partial M_j$, σ_j is Lipschitz continuous, and $\sum_{j=0}^2 \sigma_j > 0$. Hence the resulted family of ρ_j satisfy Assumption 2.2.

For the discretization of $D_j = [-r, r]^4$, we divide each coordinate interval [-r, r]into N equal parts. The scale of the grid is thus h = 2r/N. There are $(N + 1)^4$ nodes on D_j , most rows of the stiffness matrix have $3^4 = 81$ nonzero entries. We keep $N \leq 80$ due to the memory limitation of the hardware.

To get the *n*-th numerical approximation $u_h^n = (u_{h,0}^n, u_{h,1}^n, u_{h,2}^n)$, we need to solve a linear system $A_j X_j^n = b_{n,j}$ for $0 \le j \le 2$, where X_j^n provides the interior degrees of freedom of $u_{h,j}^n$. We use the Conjugate Gradient Method (CG) to find X_j^n . As a result, the process to generate the sequence $\{u_h^n\}$ is a nested iteration. The outer iteration is the DDM procedure Algorithm 3.1. The initial guess is chosen as $u_h^0 = 0$. For each *n*, the inner iteration is the CG iteration to solve $A_j X_j^n = b_{n,j}$ for $0 \le j \le 2$. Note that A_j remains the same when *n* changes, whereas $b_{n,j}$ varies because of the evaluation of $u_{h,j}^n|_{\partial D_j}$. If $\{u_h^n\}$ does converge, X_j^{n-1} will be close to X_j^n when *n* is large enough. Thus, we choose the initial guess of X_j^n as X_j^{n-1} . The tolerance for CG is set as

$$||A_j X_j^n - b_{n,j}||_2 / ||b_{n,j}||_2 \le 10^{-8}.$$

Our numerical results show that u_h^n becomes stable when $n = n_0$ for some n_0 , i.e. $u_h^n = u_h^{n_0}$ up to machine precision for all $n \ge n_0$. Actually, if

$$||A_j X_j^n - b_{n+1,j}||_2 / ||b_{n+1,j}||_2 \le 10^{-8}$$

for all j, then the inner iteration terminates for n+1 and $X_j^{n+1} = X_j^n$. We found that inner iteration terminates for all $n > n_0$. In other words, practically, the sequence $\{u_h^n\}$ reaches its limit

$$u_h^\infty = u_h^{n_0}$$

at step n_0 .

In the following tables,

$$I_h u = (I_h u_0, I_h u_1, I_h u_2) \in V_h,$$

where $I_h u_j \in V_{h,j}$ is the interpolation of $u \circ \phi_j$ on D_j . We define the energy norm of the error as

$$\|I_h u - u_h^{\infty}\|_a = \max\{a_j(I_h u_j - u_{h,j}^{\infty}, I_h u_j - u_{h,j}^{\infty})^{\frac{1}{2}} \mid 0 \le j \le 2\}.$$

The L^2 -norm $||I_h u - u_h^{\infty}||_{L^2}$, L^{∞} -norm $||I_h u - u_h^{\infty}||_{L^{\infty}}$ and H^1 -seminorm $|I_h u - u_h^{\infty}||_{H^1}$ are defined in similar ways. The numerical results are as follows in Tables 1 and 2, where, for each norm, the data on the left side of each cell are errors and orders of convergence are appended to the right.

h	$ I_h u - u_h^{\infty} _{L^{\infty}}$		$ I_h u - u_h^{\infty} $	$ _{L^2}$	$ I_h u - u_h^{\infty} _{H^1}$		$\ I_h u - u_h^\infty\ _a$		n_0
0.24	0.0376		0.0451		0.1548		0.0715		32
0.12	0.0103	1.9	0.0116	2.0	0.0438	1.8	0.0203	1.8	31
0.06	0.0024	2.1	0.0029	2.0	0.0126	1.8	0.0057	1.8	31
0.03	6.1841e - 4	2.0	7.3551e - 4	2.0	0.0039	1.7	0.0017	1.7	30
			TA	ble 1					

Numerical result on \mathbb{CP}^2 for r = 1.2.

h	$ I_hu-u $	$u_h^\infty \ _{L^\infty}$	$ I_hu-u $	$\iota_h^\infty \ _{L^2}$	$ I_hu-u $	$u_h^{\infty} _{H^1}$	$ I_hu-u $	$u_h^\infty \ _a$	n_0
0.4	0.1004		0.3600		0.7681		0.2134		10
0.2	0.0307	1.7	0.0826	2.1	0.2358	1.7	0.0647	1.7	11
0.1	0.0093	1.7	0.0219	1.9	0.0745	1.7	0.0192	1.8	11
0.05	0.0020	2.2	0.0052	2.1	0.0240	1.6	0.0059	1.7	11

TABLE 2 Numerical result on \mathbb{CP}^2 for r = 2.

We see that the error $I_h u - u_h^{\infty}$ decays in the optimal order when h decreases. Furthermore, n_0 decreases when r becomes larger, i.e. u_h^n reaches its limit u_h^∞ fast provides that the overlapping between subdomains is large.

5.2. A Problem on $S^2 \times S^2$. Let $M = S^2 \times S^2$, where S^2 is the unit sphere in \mathbb{R}^3 , i.e.

$$S^{2} = \left\{ (y_{1}, y_{2}, y_{3}) \in \mathbb{R}^{3} \middle| \sum_{i=1}^{3} y_{i}^{2} = 1 \right\}.$$

In this example, the Riemannian metric, the concrete problem of (2.2), the decomposition of M, and the transition maps are the same as those in [10, § 5.3]. We refer to [10] for a thorough and detailed description. Here we only briefly overview the decomposition.

We decompose $S^2 \times S^2$ into 4 subdomains. In the following, let $x = (x_1, x_2)$ and $x' = (x'_1, x'_2)$ denote the coordinates of \mathbb{R}^2 . Let $||x|| = \sqrt{\sum_{i=1}^2 x_i^2}$ and ||x'|| = $\sqrt{\sum_{i=1}^{2} x'_{i}^{2}}$. For $1 \le i \le 4$, let $D_i = [-r, r]^4 = \{(x, x') \mid x \in [-r, r]^2, x' \in [-r, r]^2\}.$

18

For $1 \leq i \leq 4$, we have the following diffeomorphisms

$$\phi_1: D_1 \to M_1 \subset S^2 \times S^2 \subset \mathbb{R}^3 \times \mathbb{R}^3$$
$$(x, x') \mapsto \left(\frac{2x}{1+\|x\|^2}, \frac{1-\|x\|^2}{1+\|x\|^2}, \frac{2x'}{1+\|x'\|^2}, \frac{1-\|x'\|^2}{1+\|x'\|^2}\right),$$

$$\phi_2: D_2 \to M_2 \subset S^2 \times S^2 \subset \mathbb{R}^3 \times \mathbb{R}^3$$
$$(x, x') \mapsto \left(\frac{2x}{1+\|x\|^2}, \frac{1-\|x\|^2}{1+\|x\|^2}, \frac{2x'}{1+\|x'\|^2}, \frac{-1+\|x'\|^2}{1+\|x'\|^2}\right),$$

$$\phi_3: \ D_3 \to M_3 \subset S^2 \times S^2 \subset \mathbb{R}^3 \times \mathbb{R}^3 (x, x') \mapsto \left(\frac{2x}{1+\|x\|^2}, \frac{-1+\|x\|^2}{1+\|x\|^2}, \frac{2x'}{1+\|x'\|^2}, \frac{1-\|x'\|^2}{1+\|x'\|^2}\right),$$

and

$$\phi_4: \ D_4 \to M_4 \subset S^2 \times S^2 \subset \mathbb{R}^3 \times \mathbb{R}^3$$
$$(x, x') \mapsto \left(\frac{2x}{1+\|x\|^2}, \frac{-1+\|x\|^2}{1+\|x\|^2}, \frac{2x'}{1+\|x'\|^2}, \frac{-1+\|x'\|^2}{1+\|x'\|^2}\right).$$

To guarantee $S^2 \times S^2 = \bigcup_{i=1}^4 (M_i \setminus \partial M_i)$, we have to let r > 1. The larger r is, the more overlapping there will be.

To obtain a partition of unity $\{\rho_i \mid 1 \leq i \leq 4\}$, it suffices to define the σ_j in (3.2). We define σ_j as (5.1) again. The grids on D_i are the same as those for \mathbb{CP}^2 . We choose the initial guess as 0. The numerical results are as follows in Tables 3 and 4.

h	$\ I_h u - u_h^\infty\ _{L^\infty}$		$ I_h u - u_h^{\infty} $	L^2	$ I_h u - u $	$ h_h^{\infty} _{H^1}$	$\ I_h u - u_h^\infty\ _a$		n_0
0.24	0.0192		0.0503		0.1444		0.2093		21
0.12	0.0043	2.2	0.0126	2.0	0.0410	1.8	0.0577	1.9	21
0.06	0.0011	2.0	0.0031	2.0	0.0115	1.8	0.0154	1.9	21
0.03	2.8668e - 4	1.9	7.6697e - 4	2.0	0.0041	1.5	0.0050	1.6	21
			TA	ble 3					

Numerical	result	on S^2	$ imes S^2$	for r	= 1.	2.
-----------	--------	----------	-------------	---------	------	----

h	$ I_hu-u $	${}_{h}^{\infty}\ _{L^{\infty}}$	$ I_hu-u $	$\iota_h^\infty \ _{L^2}$	$ I_hu-u $	$ h_h^{\infty} _{H^1}$	$ I_hu-u $	$u_h^\infty \ _a$	n_0
0.4	0.1381		0.6327		0.9428		0.9869		9
0.2	0.0228	2.6	0.1331	2.2	0.3062	1.6	0.2908	1.8	8
0.1	0.0085	1.4	0.0363	1.9	0.0996	1.6	0.0857	1.8	8
0.05	0.0015	2.5	0.0084	2.1	0.0329	1.6	0.0260	1.7	9

TABLE 4 Numerical result on $S^2 \times S^2$ for r = 2.

5.3. A Problem on B^4 . Let $M = B^4$ be the unit ball in \mathbb{R}^4 , i.e.

$$B^{4} = \left\{ (y_{1}, y_{2}, y_{3}, y_{4}) \in \mathbb{R}^{4} \left| \sum_{i=1}^{4} y_{i}^{2} \leq 1 \right\}.$$

This is a manifold with boundary S^3 . Following the suggestion in Section 4, to decompose B^4 , we firstly choose a collar neighborhood of $\partial B^4 = S^3$. Let

$$\psi : \ [\delta, 1] \times S^3 \to B^4$$
$$(t, y_1, y_2, y_3, y_4) \mapsto t(y_1, y_2, y_3, y_4)$$

be a collar mapping, where $0 < \delta < 1$. (Again, see Fig. 4 for an illustration.) Let $M_1 = [-s, s]^4$, we have

$$B^4 = \psi((\delta, 1] \times S^3) \cup (M_1 \setminus \partial M_1)$$

provided that $0 < \delta < s < 0.5$. To obtained a desired decomposition of the collar neighborhood, as pointed out in Section 4, it suffices to decompose S^3 . We employ the stereographic projection to do this job as in [10, § 5.1].

In summary, a decomposition with 3 subdomains is provided by the following diffeomorphisms:

$$\phi_1: D_1 = [-s, s]^4 \to M_1$$
$$x \mapsto x,$$

$$\phi_2: D_2 = [\delta, 1] \times [-r, r]^3 \to M_2$$
$$(t, \check{x}) \mapsto t\left(\frac{2\check{x}}{1 + \|\check{x}\|^2}, \frac{1 - \|\check{x}\|^2}{1 + \|\check{x}\|^2}\right),$$

and

$$\phi_3: D_3 = [\delta, 1] \times [-r, r]^3 \to M_3$$
$$(t, \check{x}) \mapsto t\left(\frac{2\check{x}}{1 + \|\check{x}\|^2}, \frac{-1 + \|\check{x}\|^2}{1 + \|\check{x}\|^2}\right),$$

where $\check{x} = (x_1, x_2, x_3) \in [-r, r]^3$ and $||\check{x}|| = \sqrt{\sum_{i=1}^3 x_i^2}$. The transitions of coordinates are as follows:

$$\phi_2^{-1} \circ \phi_1(x) = \left(\|x\|, \frac{\check{x}}{\|x\| + x_4} \right) \text{ and } \phi_3^{-1} \circ \phi_1(x) = \left(\|x\|, \frac{\check{x}}{\|x\| - x_4} \right),$$

where $x = (x_1, x_2, x_3, x_4)$ and $\check{x} = (x_1, x_2, x_3)$ consists of the first three components of x;

$$\phi_1^{-1} \circ \phi_2(t, \check{x}) = t \left(\frac{2\check{x}}{1 + \|\check{x}\|^2}, \frac{1 - \|\check{x}\|^2}{1 + \|\check{x}\|^2} \right),$$

$$\phi_1^{-1} \circ \phi_3(t, \check{x}) = t \left(\frac{2\check{x}}{1 + \|\check{x}\|^2}, \frac{-1 + \|\check{x}\|^2}{1 + \|\check{x}\|^2} \right),$$

 $\phi_3^{-1} \circ \phi_2 = \phi_2^{-1} \circ \phi_3$, and

$$\phi_3^{-1} \circ \phi_2(t, \check{x}) = \left(t, \frac{\check{x}}{\|\check{x}\|^2}\right).$$

In the above, we have $0 < \delta < s < 0.5$ and r > 1. The larger $s - \delta$ and r are, the more overlapping there will be.

To obtain the partition of unity $\{\rho_i \mid 1 \leq i \leq 3\}$, it suffices to define σ_j in (5.1). Let $\delta' = 0.9\delta + 0.1s$, $s' = 0.1\delta + 0.9s$, and r' = 0.9r + 0.1. Define

$$\sigma_1 \circ \phi_1(x) = \begin{cases} 0, & \max\{|x_j| \mid 1 \le j \le 4\} > s'; \\ \prod_{j=1}^4 (1 - (\frac{x_j}{s'})^2), & \text{otherwise;} \end{cases}$$

 $\sigma_2 \circ \phi_2 = \sigma_3 \circ \phi_3$; and

$$\sigma_2 \circ \phi_2(t, \check{x}) = \begin{cases} 0, & t < \delta' \text{ or } \max\{|x_j| \mid 1 \le j \le 3\} > r'; \\ \frac{t - \delta'}{1 - \delta'} \prod_{j=1}^3 (1 - (\frac{x_j}{r'})^2), & \text{otherwise.} \end{cases}$$

Equip B^4 with the Riemannian metric g inherited from the standard one on \mathbb{R}^4 . Then Δ on B^4 coincides with the usual $\Delta = \sum_{\alpha=1}^4 \frac{\partial^2}{\partial y_{\alpha}^2}$ on \mathbb{R}^4 . Clearly, on D_1 , the metric is expressed as

$$g = \sum_{\alpha=1}^{4} \mathrm{d}x_{\alpha} \otimes \mathrm{d}x_{\alpha}.$$

The equation (3.3) on D_1 is: $\forall v \in H_0^1(D_1)$,

$$\int_{D_1} \left(\sum_{\alpha=1}^4 \frac{\partial u \circ \phi_1}{\partial x_\alpha} \frac{\partial v}{\partial x_\alpha} + b \cdot u \circ \phi_1 \cdot v \right) \mathrm{d}x_1 \mathrm{d}x_2 \mathrm{d}x_3 \mathrm{d}x_4 = \int_{D_1} f \circ \phi_1 \cdot v \mathrm{d}x_1 \mathrm{d}x_2 \mathrm{d}x_3 \mathrm{d}x_4.$$

On D_2 and D_3 , the Riemannian metric is expressed as

$$g = \mathrm{d}t \otimes \mathrm{d}t + 4t^2(1 + \|\check{x}\|^2)^{-2} \sum_{\alpha=1}^3 \mathrm{d}x_\alpha \otimes \mathrm{d}x_\alpha.$$

The equation (3.3) on D_i (i = 2, 3) is: $\forall v \in H_0^1(D_i)$,

$$\int_{D_i} \left[8t^3 (1+\|\check{x}\|^2)^{-3} \frac{\partial u \circ \phi_i}{\partial t} \frac{\partial v}{\partial t} + 2t(1+\|\check{x}\|^2)^{-1} \sum_{\alpha=1}^3 \frac{\partial u \circ \phi_i}{\partial x_\alpha} \frac{\partial v}{\partial x_\alpha} + 8t^3 (1+\|\check{x}\|^2)^{-3} b \cdot u \circ \phi_i \cdot v \right] dt dx_1 dx_2 dx_3$$
$$= \int_{D_i} 8t^3 (1+\|\check{x}\|^2)^{-3} f \circ \phi_i \cdot v dt dx_1 dx_2 dx_3.$$

We choose the true solution u in B^4 as

$$u(y_1, y_2, y_3, y_4) = \sin(\pi y_4).$$

Then f in (2.2) is

$$f = (b + \pi^2)u$$

In D_i , $u \circ \phi_i$ has the expression

$$u \circ \phi_1(x) = \sin(\pi x_4),$$

$$u \circ \phi_2(t, x) = \sin\left(\pi t \frac{1 - \|\check{x}\|^2}{1 + \|\check{x}\|^2}\right), \quad \text{and} \quad u \circ \phi_3(t, x) = \sin\left(\pi t \frac{-1 + \|\check{x}\|^2}{1 + \|\check{x}\|^2}\right).$$

Now we choose b = 0 in (2.2). For the discretization on D_i , we divide each coordinate interval [-s, s] and $[\delta, 1]$ into N_1 equal parts, divide each coordinate interval [-r, r] into N_2 equal parts. Set $N_1 = 0.4N_2$. Thus the scale of the grid is

$$h = \max\{\frac{2s}{N_1}, \frac{1-\delta}{N_1}, \frac{2r}{N_2}\}.$$

We choose N_2 as 10, 20, 40, and 80. For the initial guess, we set the degrees of freedom off ∂M as 0. The numerical results are as follows in Tables 5 and 6.

h	$ I_hu-u $	$\iota_h^\infty \ _{L^\infty}$	$ I_hu-u $	$\iota_h^\infty \ _{L^2}$	$ I_h u - u $	$\iota^\infty_h _{H^1}$	$ I_hu-c $	$u_h^\infty \ _a$	n_0			
0.24	0.1049		0.0604		0.3642		0.2278		13			
0.12	0.0267	2.0	0.0177	1.8	0.1305	1.5	0.0799	1.5	13			
0.06	0.0073	1.9	0.0047	1.9	0.0413	1.7	0.0251	1.7	13			
0.03	0.0020	1.9	0.0012	2.0	0.0127	1.7	0.0078	1.7	13			
	TABLE 5											

Numerical result on B^4 for $(s, \delta, r) = (0.4, 0.2, 1.2)$.

h	$ I_hu-u $	$\iota_h^\infty \ _{L^\infty}$	$ I_hu-u $	$\iota_h^\infty \ _{L^2}$	$ I_hu-u $	$u_h^{\infty} _{H^1}$	$ I_hu-a $	$u_h^\infty \ _a$	n_0
0.4	0.2251		0.1389		0.6561		0.3443		8
0.2	0.0582	2.0	0.0418	1.7	0.2854	1.2	0.1182	1.5	9
0.1	0.0216	1.4	0.0120	1.8	0.1080	1.4	0.0361	1.7	9
0.05	0.0054	2.0	0.0031	2.0	0.0351	1.6	0.0111	1.7	9
				TABLE	6				

Numerical result on B^4 for $(s, \delta, r) = (0.4, 0.1, 2)$.

5.4. A Problem on $B^2 \times S^2$. Let $M = B^2 \times S^2$, where B^2 is the unit ball (disk) in \mathbb{R}^2 and S^2 is the unit sphere in \mathbb{R}^3 . This is a manifold with boundary $S^1 \times S^2$.

As pointed out in [10, § 4], decompositions of B^2 and S^2 yield a canonical product decomposition of $B^2 \times S^2$. Thus it suffices to decompose both factor manifolds.

We use a decomposition of B^2 similar to that of B^4 above. In other words, it is provided by

$$\phi_1: D_1 = [-s, s]^2 \to M_1 \subset B^2$$
$$x \mapsto x.$$

$$\phi_2: D_2 = [\delta, 1] \times [-r, r] \to M_2 \subset B^2$$
$$(t, \check{x}) \mapsto t\left(\frac{2\check{x}}{1 + \check{x}^2}, \frac{1 - \check{x}^2}{1 + \check{x}^2}\right).$$

and

$$\phi_3: D_3 = [\delta, 1] \times [-r, r] \to M_3 \subset B^2$$
$$(t, \check{x}) \mapsto t\left(\frac{2\check{x}}{1 + \check{x}^2}, \frac{-1 + \check{x}^2}{1 + \check{x}^2}\right)$$

Here $0 < \delta < s < \frac{1}{\sqrt{2}}$ and r > 1. The transitions of coordinates are as follows:

$$\phi_2^{-1} \circ \phi_1(x) = \left(\|x\|, \frac{x_1}{\|x\| + x_2} \right) \text{ and } \phi_3^{-1} \circ \phi_1(x) = \left(\|x\|, \frac{x_1}{\|x\| - x_2} \right),$$

where $x = (x_1, x_2);$

$$\phi_1^{-1} \circ \phi_2(t, \check{x}) = t \left(\frac{2\check{x}}{1 + \check{x}^2}, \frac{1 - \check{x}^2}{1 + \check{x}^2} \right), \qquad \phi_1^{-1} \circ \phi_3(t, \check{x}) = t \left(\frac{2\check{x}}{1 + \check{x}^2}, \frac{-1 + \check{x}^2}{1 + \check{x}^2} \right),$$

 $\phi_3^{-1} \circ \phi_2 = \phi_2^{-1} \circ \phi_3$, and

$$\phi_3^{-1} \circ \phi_2(t, \check{x}) = \left(t, \frac{1}{\check{x}}\right)$$

We use stereographic projection to decompose S^2 . In other words, the decomposition is provided by

$$\phi_1': D_1' = [-r, r]^2 \to M_1' \subset S^2 \subset \mathbb{R}^3$$
$$x' \mapsto \left(\frac{2x'}{1 + \|x'\|^2}, \frac{1 - \|x'\|^2}{1 + \|x'\|^2}\right)$$

and

$$\begin{aligned} \phi_2': \ D_2' &= [-r,r]^2 \to M_2' \subset S^2 \subset \mathbb{R}^3 \\ x' &\mapsto \left(\frac{2x'}{1+\|x'\|^2}, \frac{-1+\|x'\|^2}{1+\|x'\|^2}\right). \end{aligned}$$

Here r > 1. The transitions of coordinates are given by

$$\phi'_{2}^{-1} \circ \phi'_{1}(x') = \phi'_{1}^{-1} \circ \phi'_{2}(x') = \frac{x'}{\|x'\|^{2}}.$$

The above decompositions result in a decomposition of $B^2 \times S^2$ with $3 \times 2 = 6$ subdomains. Define $D_{i,i'} = D_i \times D'_{i'}$, $M_{i,i'} = M_i \times M'_{i'}$ and $\phi_{i,i'} = \phi_i \times \phi'_{i'}$. The decomposition of $B^2 \times S^2$ are provided by the diffeomorphisms

$$\phi_{i,i'}: D_{i,i'} \to M_{i,i'} \subset B^2 \times S^2$$

for $1 \leq i \leq 3$ and $1 \leq i' \leq 2$. The transition maps are

$$\phi_{j,j'}^{-1} \circ \phi_{i,i'} = (\phi_j \times \phi'_{j'})^{-1} \circ (\phi_i \times \phi'_{i'}) = (\phi_j^{-1} \circ \phi_i) \times (\phi'_{j'}^{-1} \circ \phi'_{i'}).$$

Note that $D_{1,1} = D_{1,2} = [-s,s]^2 \times [-r,r]^2$ and other $D_{i,i'}$ are $[\delta,1] \times [-r,r]^3$. Here $0 < \delta < s < \frac{1}{\sqrt{2}}$ and r > 1. The larger $s - \delta$ and r are, the more overlapping there will be.

We need to define a partition of unity $\{\rho_{i,i'} \mid 1 \leq i \leq 3, 1 \leq i' \leq 2\}$ such that $\operatorname{supp}\rho_{i,i'} \subset M_{i,i'} \setminus \overline{\gamma_{i,i'}}$, where $\gamma_{i,i'} = \partial M_{i,i'} \setminus \partial M$. Let $\delta' = 0.9\delta + 0.1s$, $s' = 0.1\delta + 0.9s$, and r' = 0.9r + 0.1. On $D_{1,1} = D_{1,2} = [-s,s]^2 \times [-r,r]^2$, define

$$\sigma_{1,i'} \circ \phi_{1,i'}(x_1, x_2, x_3, x_4) = \begin{cases} 0, & \max\{|x_1|, |x_2|\} > s' \text{ or } \max\{|x_3|, |x_4|\} > r'; \\ \prod_{j=1}^2 (1 - (\frac{x_j}{s'})^2) \prod_{j=3}^4 (1 - (\frac{x_j}{r'})^2), & \text{otherwise.} \end{cases}$$

On other $D_{i,i'} = [\delta, 1] \times [-r, r]^3$, define

$$\begin{split} &\sigma_{i,i'} \circ \phi_{i,i'}(t,x_1,x_2,x_3) \\ &= \begin{cases} 0, & t < \delta' \text{ or } \max\{|x_1|,|x_2|,|x_3|\} > r'; \\ \frac{t-\delta'}{1-\delta'} \prod_{j=1}^3 (1-(\frac{x_j}{r'})^2), & \text{otherwise.} \end{cases} \end{split}$$

Applying a formula similar to (3.2), we obtain a desired $\{\rho_{i,i'} \mid 1 \le i \le 3, 1 \le i' \le 2\}$.

Equip B^2 with the Riemannian metric inherited from the standard one on \mathbb{R}^2 . Equip S^2 with the Riemannian metric inherited from the standard one on \mathbb{R}^3 . Consider the product metric g on $B^2 \times S^2$. On $D_{1,i'}$, the coordinates are (x, x'), where $x = (x_1, x_2)$ and $x' = (x'_1, x'_2)$, the metric g is expressed as

$$g = \sum_{\alpha=1}^{2} \mathrm{d}x_{\alpha} \otimes \mathrm{d}x_{\alpha} + 4(1 + \|x'\|^2)^{-2} \sum_{\alpha'=1}^{2} \mathrm{d}x'_{\alpha'} \otimes \mathrm{d}x'_{\alpha'}.$$

The equation (3.3) is

$$\begin{split} \int_{D_{1,i'}} \left[4(1+\|x'\|^2)^{-2} \sum_{\alpha=1}^2 \frac{\partial u \circ \phi_{1,i'}}{\partial x_{\alpha}} \frac{\partial v}{\partial x_{\alpha}} + \sum_{\alpha'=1}^2 \frac{\partial u \circ \phi_{1,i'}}{\partial x'_{\alpha'}} \frac{\partial v}{\partial x'_{\alpha'}} \right. \\ \left. + 4(1+\|x'\|^2)^{-2} b \cdot u \circ \phi_{1,i'} \cdot v \right] \mathrm{d}x_1 \mathrm{d}x_2 \mathrm{d}x'_1 \mathrm{d}x'_2 \\ = \int_{D_{1,i'}} 4(1+\|x'\|^2)^{-2} \cdot f \circ \phi_{1,i'} \cdot v \mathrm{d}x_1 \mathrm{d}x_2 \mathrm{d}x'_1 \mathrm{d}x'_2. \end{split}$$

On $D_{i,i'}$ for $2 \le i \le 3$, the coordinates are (t, \check{x}, x') , where $x' = (x'_1, x'_2)$,

$$g = \mathrm{d}t \otimes \mathrm{d}t + 4t^2 (1 + \check{x}^2)^{-2} \mathrm{d}\check{x} \otimes \mathrm{d}\check{x} + 4(1 + ||x'||^2)^{-2} \sum_{\alpha'=1}^2 \mathrm{d}x'_{\alpha'} \otimes \mathrm{d}x'_{\alpha'}.$$

The equation (3.3) is

$$\begin{split} &\int_{D_{i,i'}} \left[8t(1+\check{x}^2)^{-1}(1+\|x'\|^2)^{-2} \frac{\partial u \circ \phi_{i,i'}}{\partial t} \frac{\partial v}{\partial t} \right. \\ &+ 2t^{-1}(1+\check{x}^2)(1+\|x'\|^2)^{-2} \frac{\partial u \circ \phi_{i,i'}}{\partial \check{x}} \frac{\partial v}{\partial \check{x}} + 2t(1+\check{x}^2)^{-1} \sum_{\alpha'=1}^2 \frac{\partial u \circ \phi_{i,i'}}{\partial x'_{\alpha'}} \frac{\partial v}{\partial x'_{\alpha'}} \\ &+ 8t(1+\check{x}^2)^{-1}(1+\|x'\|^2)^{-2} b \cdot u \circ \phi_{i,i'} \cdot v \right] dt d\check{x} dx'_1 dx'_2 \\ &= \int_{D_{i,i'}} 8t(1+\check{x}^2)^{-1}(1+\|x'\|^2)^{-2} \cdot f \circ \phi_{i,i'} \cdot v dt d\check{x} dx'_1 dx'_2. \end{split}$$

We set the true solution u on $B^2 \times S^2$ as

$$u(y_1, y_2, y_1', y_2', y_3') = \sin(\pi y_2) + y_3'.$$

Then f in (2.2) is

$$f(y_1, y_2, y_1', y_2', y_3') = (b + \pi^2)\sin(\pi y_2) + (b + 2)y_3'$$

On $D_{i,i'}$, we have

$$u \circ \phi_{i,i'}(x_1, x_2, x_3, x_4) = u_i(x_1, x_2) + u'_{i'}(x_3, x_4)$$

and

$$f \circ \phi_{i,i'}(x_1, x_2, x_3, x_4) = f_i(x_1, x_2) + f'_{i'}(x_3, x_4),$$

where

$$u_1(x_1, x_2) = \sin(\pi x_2), \quad u_2(t, \check{x}) = \sin\left(\pi t \frac{1 - \check{x}^2}{1 + \check{x}^2}\right), \quad u_3(t, \check{x}) = \sin\left(\pi t \frac{-1 + \check{x}^2}{1 + \check{x}^2}\right),$$
$$u_1'(x') = \frac{1 - \|x'\|^2}{1 + \|x'\|^2}, \qquad u_2'(x') = \frac{-1 + \|x'\|^2}{1 + \|x'\|^2},$$

 $f_i = (b + \pi^2)u_i$, and $f'_{i'} = (b + 2)u'_{i'}$. Now we choose b = 1 in (2.2). For the discretization on $D_{i,i'}$, we divide each coordinate interval [-s, s] and $[\delta, 1]$ into N_1 equal parts, divide each coordinate interval [-r, r] into N_2 equal parts. Set $N_1 = 0.4N_2$. We choose N_2 as 10, 20, 40, and 80. For the initial guess, we set the degrees of freedom off ∂M as 0. The numerical results are as follows in Tables 7 and 8.

h	$ I_h u - u_h^{\infty} $	$\ L^{\infty}\ $	$ I_hu-v $	$\iota_h^\infty \ _{L^2}$	$ I_h u - u $	$u_h^\infty _{H^1}$	$ I_hu-u $	$\ u_h^\infty\ _a$	n_0
3/10	0.1046		0.0916		0.3938		0.4863		35
3/20	0.0343	1.6	0.0261	1.8	0.1180	1.7	0.1373	1.8	36
3/40	0.0083	2.0	0.0072	1.9	0.0332	1.8	0.0373	1.9	33
3/80	0.0021	2.0	0.0018	2.0	0.0103	1.7	0.0113	1.7	31
				TABLE	7				

h	$ I_hu-u $	$\iota_h^\infty \ _{L^\infty}$	$ I_hu-v $	$\iota_h^\infty \ _{L^2}$	$ I_hu-u $	$\iota_h^\infty _{H^1}$	$ I_hu-c $	$u_h^\infty \ _a$	n_0
0.4	0.2555		0.4455		1.7415		1.3294		18
0.2	0.0831	1.6	0.1344	1.7	0.6474	1.4	0.4392	1.6	18
0.1	0.0229	1.9	0.0247	2.4	0.1639	2.0	0.1320	1.7	19
0.05	0.0059	2.0	0.0061	2.0	0.0519	1.7	0.0415	1.7	19
				TADIE	8				

Numerical result on $B^2 \times S^2$ for $(s, \delta, r) = (0.7, 0.1, 2)$.

Acknowledgements. We thank Jun Li, Yuanming Xiao and Yiyan Xu for various discussions. The first author was partially supported by NSFC 11871272. The second author was partially supported by NSFC 12071227 and 12371369, and the National Key Research and Development Program of China 2020YFA0713803. The third author was partially supported by NSFC 12171349 and 12271389.

REFERENCES

- P. ANTONIETTI, A. DEDNER, P. MADHAVAN, S. STANGALINO, B. STINNER, AND M. VERANI, High order discontinuous Galerkin methods for elliptic problems on surfaces, SIAM Journal on Numerical Analysis, 53 (2015), pp. 1145–1171.
- [2] E. BACHINI, M. FARTHING, AND M. PUTTI, Intrinsic finite element method for advectiondiffusion-reaction equations on surfaces, Journal of Computational Physics, 424 (2021), p. 109827.
- [3] E. BANSCH, P. MORIN AND R. NOCHETTO, A finite element method for surface diffusion: the parametric case, J. Comp. Physics, 203 (2005), 321–343. 1
- [4] J. BARRERA, T. KOLEV, K. MITTAL AND V. TOMOV, High-order mesh morphing for boundary and interface fitting to implicit geometries, Comput.-Aided Des. 158 (2023), Paper No. 103499, 11 pp. 1
- J. BAUMGARDNER AND P. FREDERICKSON, Icosahedral discretization of the two-sphere, SIAM Journal on Numerical Analysis, 22 (1985), pp. 1107–1115.
- [6] A. BONITO, A. DEMLOW, AND M. LICHT, A divergence-conforming finite element method for the surface Stokes equation, SIAM Journal on Numerical Analysis, 58 (2020), pp. 2764–2798.
 1
- [7] A. BONITO, A. DEMLOW, AND R. NOCHETTO, Finite element methods for the Laplace-Beltrami operator, in Handbook of Numerical Analysis, vol. 21, Elsevier, 2020, pp. 1–103. 1
- [8] J. BRAMBLE, J. PASCIAK, J. WANG, AND J. XU, Convergence estimates for product iterative methods with applications to domain decomposition, Mathematics of Computation, 57 (1991), pp. 1–21. 2
- X.-C. CAI AND Y. SAAD, Overlapping domain decomposition algorithms for general sparse matrices, Numerical linear algebra with applications, 3 (1996), pp. 221–237.
- [10] S. CAO AND L. QIN, A numerical domain decomposition method for solving elliptic equations on manifolds, To appear in SIAM Journal on Scientific Computing, arXiv:2212.04079. 2, 3, 4, 12, 13, 14, 15, 16, 17, 18, 20, 22
- P. CIARLET, The finite element method for elliptic problems, Society for Industrial and Applied Mathematics, Philadelphia, PA, 2002. 12, 13
- [12] M. CRANDALL, H. ISHII, AND P. L. LIONS, User's guide to viscosity solutions of second order partial differential equations, Bull. Amer. Math. Soc. (N.S.) 27 (1992), no.1, 1–67. 6

- [13] K. DECKELNICK, G. DZIUK, AND C. ELLIOTT, Computation of geometric partial differential equations and mean curvature flow, Acta numerica, 14 (2005), pp. 139–232. 1
- [14] A. DEMLOW, Higher-order finite element methods and pointwise error estimates for elliptic problems on surfaces, SIAM Journal on Numerical Analysis, 47 (2009), pp. 805–827. 1
- [15] A. DEMLOW AND G. DZIUK, An adaptive finite element method for the Laplace-Beltrami operator on implicitly defined surfaces, SIAM Journal on Numerical Analysis, 45 (2007), pp. 421–442. 1
- [16] V. DOLEAN, P. JOLIVET, AND N. NATAF, An introduction to domain decomposition methods, Algorithms, theory, and parallel implementation, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2015. 2
- [17] V. DOBREV, J.-L. GUERMOND, AND B. POPOV, Surface reconstruction and image enhancement via L¹-minimization, SIAM Journal on Scientific Computing, 32 (2010), pp. 1591–1616. 1
- [18] M. DRYJA, J. GALVIS, AND M. SARKIS, A FETI-DP preconditioner for a composite finite element and discontinuous Galerkin method, SIAM J. Numer. Anal. 51 (2013), no.1, 400– 422. 2
- [19] G. DZIUK, Finite elements for the Beltrami operator on arbitrary surfaces, Partial differential equations and calculus of variations, (1988), pp. 142–155. 1
- [20] G. DZIUK, An algorithm for evolutionary surfaces, Numerische Mathematik, 58 (1990), pp. 603– 611. 1
- [21] G. DZIUK AND C. ELLIOTT, Finite elements on evolving surfaces, IMA journal of numerical analysis, 27 (2007), pp. 262–292. 1
- [22] G. DZIUK AND C. ELLIOTT, Finite element methods for surface PDEs, Acta Numerica, 22 (2013), pp. 289–396. 1
- [23] L. EVANS, Partial differential equations, Second edition, Grad. Stud. Math. 19, American Mathematical Society, Providence, RI, 2010. 11
- [24] L. EVANS AND R. GARIEPY, Measure theory and fine properties of functions, Stud. Adv. Math., CRC Press, Boca Raton, FL, 1992. 5, 11
- [25] D. GILBARG AND N. TRUDINGER, Elliptic partial differential equations of second order, Springer-Verlag, Berlin, 2001. 5, 7, 10
- [26] F. HARVEY AND H. LAWSON, The equivalence of viscosity and distributional subsolutions for convex subequations-a strong Bellman principle, Bull. Braz. Math. Soc. (N.S.) 44 (2013), no.4, 621–652. 6
- [27] F. HARVEY AND H. LAWSON, Removable singularities for nonlinear subequations, Indiana Univ. Math. J. 63 (2014), no.5, 1525–1552. 10
- [28] M. HIRSCH, Differential topology, Graduate Texts in Mathematics, No. 33. Springer-Verlag, New York-Heidelberg, 1976. 4, 15
- [29] M. HOLST, O. SARBACH, M. TIGLIO AND M. VALLISNERI, The emergence of gravitational wave science: 100 years of development of mathematical theory, detectors, numerical algorithms, and data analysis tools, Bull. Amer. Math. Soc. (N.S.) 53 (2016), no.4, 513–554.
- [30] M. HOLST AND A. STERN, Geometric variational crimes: Hilbert complexes, finite element exterior calculus, and problems on hypersurfaces, Found. Comput. Math. 12 (2012), no.3, 263–293. 1
- [31] H. ISHII, On the equivalence of two notions of weak solutions, viscosity solutions and distribution solutions, Funkcial. Ekvac. 38 (1995), no. 1, 101–120. 6
- [32] T. JANKUHN, M. OLSHANSKII, A. REUSKEN, AND A. ZHILIAKOV, Error analysis of higher order trace finite element methods for the surface Stokes equation, Journal of Numerical Mathematics, 29 (2021), pp. 245–267. 1
- [33] M. JIN, X. FENG, AND K. WANG, Gradient recovery-based adaptive stabilized mixed fem for the convection-diffusion-reaction equation on surfaces, Computer Methods in Applied Mechanics and Engineering, 380 (2021), p. 113798. 1
- [34] J. JOST, Riemannian geometry and geometric analysis, Seventh edition, Universitext, Springer, Cham, 2017. 1
- [35] P. L. LIONS, Optimal control of diffusion processes and Hamilton-Jacobi-Bellman equations II, viscosity solutions and uniqueness, Comm. Partial Differential Equations 8 (1983), no. 11, 1229–1276. 6
- [36] P. L. LIONS, On the Schwarz alternating method I, in the First International Symposium on Domain Decomposition Methods for Partial Differential Equations, 1988. 2, 3, 4
- [37] P. L. LIONS, On the Schwarz alternating method II, in the Second International Symposium on Domain Decomposition Methods for Partial Differential Equations, 1989. 2, 3, 4, 5, 6
- [38] A. MOHAMED AND C. DAVATZIKOS, Finite element modeling of brain tumor mass-effect from 3d medical images, in International conference on medical image computing and computer-

assisted intervention, Springer, 2005, pp. 400–408. 1

- [39] J. NEDELEC, Curved finite element methods for the solution of singular integral equations on surfaces in r3, Computer Methods in Applied Mechanics and Engineering, 8 (1976), pp. 61–80. 1
- [40] J. NEDELEC AND J. PLANCHARD, Une méthode variationnelle d'éléments finis pour la résolution numérique d'un problème extérieur dans R³, Revue française d'automatique, informatique, recherche opérationnelle. Analyse numérique, 7 (1973), pp. 105–129. 1
- [41] M. OLSHANSKII AND A. REUSKEN, Trace finite element methods for PDEs on surfaces, in Geometrically unfitted finite element methods and applications, Springer, 2017, pp. 211– 258. 1
- [42] L. QIN, S. ZHANG AND Z. ZHANG, Finite element formulation in flat coordinate spaces to solve elliptic problems on general closed Riemannian manifolds, SIAM J. Sci. Comput. 36 (2014), no. 5, A2149–A2165. 2
- [43] A. QUARTERONI AND A. VALLI, Domain decomposition methods for partial differential equations, The Clarendon Press, Oxford University Press, New York, 1999. 2
- [44] A. REUSKEN, Analysis of trace finite element methods for surface partial differential equations, IMA Journal of Numerical Analysis, 35 (2015), pp. 1568–1590.
- [45] M. REUTER, S. BIASOTTI, D. GIORGI, G. PATANÈ, AND M. SPAGNUOLO, Discrete Laplace-Beltrami operators for shape analysis and segmentation, Computers & Graphics, 33 (2009), pp. 381–390. 1
- [46] R. SCHOEN AND S. T. YAU, Lectures on Differential Geometry, International Press, Cambridge, MA, 1994. 1
- [47] H. A. SCHWARZ, Ueber einige Abbildungsaufgaben, Journal f
 ür die Reine und Angewandte Mathematik, 70 (1869), 105–120. 2
- [48] B. SMITH, P. BJØRSTAD AND W. GROPP, Domain decomposition, Parallel multilevel methods for elliptic partial differential equations, Cambridge University Press, Cambridge, 1996. 2
- [49] A. TOSELLI AND O. WIDLUND, Domain Decomposition Methods-Algorithms and Theory, vol. 34, Springer Science & Business Media, 2004. 2
- [50] J. H. C. WHITEHEAD, On C¹-complexes, Ann. of Math. (2) 41 (1940), 809–824. 2
- [51] J. Xu, Iterative methods by space decomposition and subspace correction, SIAM Rev. 34 (1992), no.4, 581–613. 2
- [52] J. XU AND J. ZOU, Some nonoverlapping domain decomposition methods, SIAM Rev. 40 (1998), no.4, 857–914. 2