

A characterization of extremal non-transmission-regular graphs by the distance (signless Laplacian) spectral radius

Jingfen Lan^{*a} and Lele Liu^b

^aSchool of Mathematics and Statistics, Xidian University, Xi'an 710126, China.

(jflan@xidian.edu.cn)

^bSchool of Mathematical Sciences, Anhui University, Hefei 230601, China.

(liu@ahu.edu.cn)

Abstract

Let G be a simple connected graph of order n and $\partial(G)$ is the spectral radius of the distance matrix $D(G)$ of G . The transmission D_i of vertex i is the i -th row sum of $D(G)$. Denote by $D_{\max}(G)$ the maximum of transmissions over all vertices of G , and $\partial^Q(G)$ is the spectral radius of the distance signless Laplacian matrix $D(G) + \text{diag}(D_1, D_2, \dots, D_n)$. In this paper, we present a sharp lower bound of $2D_{\max}(G) - \partial^Q(G)$ among all n -vertex connected graphs, and characterize the extremal graphs. Furthermore, we give the minimum values of respective $D_{\max}(G) - \partial(G)$ and $2D_{\max}(G) - \partial^Q(G)$ on trees and characterize the extremal trees.

AMS classification: 05C50

Keywords: Distance (signless Laplacian) matrix; Distance (signless Laplacian) spectral radius; Non-transmission-regular graph.

1 Introduction

All graphs considered in this paper are undirected and simple. Let G be a connected graph of order n with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$. The distance between two vertices v_i and v_j , denoted by d_{ij} , is the length of the shortest path between v_i and v_j . The distance matrix $D(G)$ of graph G is the $n \times n$ matrix with d_{ij} as the (i, j) -entry, whose largest eigenvalue, denoted by $\partial(G)$, is called the *distance spectral radius* of G . The *transmission* $D_i(G)$ of vertex v_i is the sum of distance from v_i to all over vertices of G , i.e., the i -th row sum of $D(G)$. Denote by $D_{\max}(G)$ and $D_{\min}(G)$ respectively the maximum and minimum of the transmissions over all vertices of G . If there is no risk of confusion, we denote $D_i(G)$, $D_{\max}(G)$ and $D_{\min}(G)$ simply by D_i , D_{\max} and D_{\min} respectively. Let $Q(G) = D(G) + \text{diag}(D_1, D_2, \dots, D_n)$ be the *distance signless Laplacian matrix* of graph G , whose largest eigenvalue, denoted by $\partial^Q(G)$,

*Corresponding author

is called the *distance signless Laplacian spectral radius* of G . The *distinguished vertex deleted regular graph* (DVDR), introduced by Atic and Panigrahi [1], is a connected graph G with a vertex v such that the degree $d(v)$ of v equals to $n - 1$ and $G - v$ is regular. The vertex v is called a *distinguished vertex* of the DVDR graph G . If $G - v$ is r -regular, we say G is an r -DVDR graph. For example, wheel graphs are 2-DVDR graphs and complete graphs are $(n - 1)$ -DVDR graphs. The *Wiener index* $W(G)$ of a graph G is the sum of distances between all unordered pairs of vertices of G , that is, the half sum of all the entries of $D(G)$,

$$W(G) = \frac{1}{2} \sum_{i=1}^n D_i.$$

It is well-known that the adjacency spectral radius, denoted by $\rho(G)$, is no more than the maximum degree $\Delta(G)$ of a graph G , and $\rho(G) = \Delta(G)$ if and only if G is regular for a connected graph G (see [5]). Hence, $\Delta(G) - \rho(G)$ can be regarded as a measure of irregularity for a nonregular graph G ([6], p. 242), and many estimates on it have been obtained ([4, 7, 14, 13, 3]). Similarly, $2\Delta(G) - q(G)$ is also used to qualify how close a graph to being regular ([11, 12]), where $q(G)$ is the signless Laplacian spectral radius of G . As a natural generalization, graph regularity associated with the distance matrix is defined. A connected graph G is *transmission-regular* if $D_{\max} = D_{\min}$; otherwise G is *non-transmission-regular*. It is known that the distance (signless Laplacian) spectral radius of a graph G satisfies $D_{\min}(G) \leq \partial(G) \leq D_{\max}(G)$ (and $2D_{\min}(G) \leq \partial^Q(G) \leq 2D_{\max}(G)$), with the equalities if and only if G is a transmission-regular graph [10]. Naturally, Liu, Shu and Xue [9] asked the question: how small can $D_{\max}(G) - \partial(G)$ and $2D_{\max}(G) - \partial^Q(G)$ be for non-transmission-regular graphs? They gave some lower bounds for the two indicators. In [8], Liu, Shan and He confirmed a conjecture with respect to $D_{\max}(G) - \partial(G)$ posed by Liu, Shu and Xue [9] by giving a tight lower bound for graphs G of order n , and determining the extremal graphs to be DVDR graphs. For a connected graph G of order n , write

$$\begin{aligned} \sigma(G) &:= D_{\max}(G) - \partial(G), \\ \tau(G) &:= 2D_{\max}(G) - \partial^Q(G). \end{aligned}$$

Using the method in [8], we give a tight lower bound for $\tau(G)$ and characterize the extremal graphs which are the same DVDR graphs as in the case of $\sigma(G)$. Furthermore, we give lower bounds respectively for $\sigma(G)$ and $\tau(G)$ on trees and characterize the extremal graphs which are stars.

Theorem 1.1. *Let G be a connected non-transmission-regular graph on n vertices.*

(1) *If n is odd, then*

$$\tau(G) \geq \frac{n + 2 - \sqrt{n^2 + 4n - 4}}{2}.$$

Equality holds if and only if $G \cong K_{1,2,\dots,2}$.

(2) *If n is even, then*

$$\tau(G) \geq \frac{n + 4 - \sqrt{n^2 + 8n}}{2}.$$

Equality holds if and only if G is isomorphic to some $(n - 4)$ -DVDR graph.

Consider $\sigma(G)$ and $\tau(G)$ only on trees, which are also non-transmission-regular graphs. We obtain the lower bounds and the extremal trees.

Theorem 1.2. *Let T be a tree on n vertices. We have*

$$\sigma(T) \geq \frac{2n - 2 - \sqrt{4n^2 - 12n + 12}}{2}$$

and

$$\tau(T) \geq \frac{3n - 4 - \sqrt{9n^2 - 32n + 32}}{2},$$

with the equalities if and only if $T \cong K_{1,n-1}$.

2 Preliminaries

Let M be a real symmetric matrix associated with a graph G . An equitable partition $\pi : V(G) = X_1 \cup X_2 \cup \dots \cup X_k$ of the vertex set $V(G)$ can divide M into the block form $M = (M_{ij})$ such that any block M_{ij} has a constant row sum b_{ij} , and then the eigenvalues of the $k \times k$ matrix $B = (b_{ij})$ are also eigenvalues of M (see [2, p. 24]). Furthermore, if M is nonnegative and irreducible then B and M have the same spectral radius. Based on this understanding, let us calculate $\tau(K_{1,2,\dots,2})$ and $\tau(G)$ of an $(n - 4)$ -DVDR graph G .

Lemma 2.1. *Let $K_{1,2,\dots,2}$ be the r -partite graph on n vertices with $n = 2r - 1$ and G is an $(n - 4)$ -DVDR graph. Then*

$$\tau(K_{1,2,\dots,2}) = \frac{n + 2 - \sqrt{n^2 + 4n - 4}}{2} \quad \text{and} \quad \tau(G) = \frac{n + 4 - \sqrt{n^2 + 8n}}{2}.$$

Proof. Note that $K_{1,2,\dots,2}$ is an $(n - 3)$ -DVDR graph. Denote by v the distinguish vertex of $K_{1,2,\dots,2}$ (or G). The quotient matrix of $Q(K_{1,2,\dots,2})$ with respect to the equitable partition $\{v\} \cup (V(K_{1,2,\dots,2}) \setminus \{v\})$ is

$$\begin{bmatrix} n - 1 & n - 1 \\ 1 & 2n - 1 \end{bmatrix},$$

which has the same spectral radius as $Q(K_{1,2,\dots,2})$, i.e.,

$$\partial^Q(K_{1,2,\dots,2}) = \frac{3n - 2 + \sqrt{n^2 + 4n - 4}}{2}.$$

Since $D_{\max}(K_{1,2,\dots,2}) = n$, we have

$$\tau(K_{1,2,\dots,2}) = 2n - \partial^Q(K_{1,2,\dots,2}) = \frac{n + 2 - \sqrt{n^2 + 4n - 4}}{2}.$$

The quotient matrix of $Q(G)$ with respect to the equitable partition $\{v\} \cup (V(G) \setminus \{v\})$ is

$$\begin{bmatrix} n - 1 & n - 1 \\ 1 & 2n + 1 \end{bmatrix},$$

which has the same spectral radius as $Q(G)$, i.e.,

$$\partial^Q(G) = \frac{3n + \sqrt{n^2 + 8n}}{2}.$$

Since $D_{\max}(G) = n + 1$, we have

$$\tau(K_{1,2,\dots,2}) = 2(n + 1) - \partial^Q(G) = \frac{n + 4 - \sqrt{n^2 + 8n}}{2}.$$

The proof is completed. \square

In the following, we denote

$$\tau_n := \frac{n + 2\gamma_n - \sqrt{(n + 2\gamma_n)^2 - 8\gamma_n}}{2},$$

where $\gamma_n = 1$ if n is odd; $\gamma_n = 2$ if n is even. We can see that γ_n satisfies the equality

$$\tau_n^2 - (n + 2\gamma_n)\tau_n + 2\gamma_n = 0. \quad (2.1)$$

By a similar calculation as the proof of Lemma 2.1, we obtain $\sigma(K_{1,n-1})$ and $\tau(K_{1,n-1})$.

Lemma 2.2. *Let $K_{1,n-1}$ be the star graph on n vertices. We have*

$$\sigma(K_{1,n-1}) = \frac{2n - 2 - \sqrt{4n^2 - 12n + 12}}{2} \quad \text{and} \quad \tau(K_{1,n-1}) = \frac{3n - 4 - \sqrt{9n^2 - 32n + 32}}{2}.$$

We denote

$$\sigma'_n := \frac{n + \eta_n - \sqrt{(n + \eta_n)^2 - 4\eta_n}}{2},$$

$$\tau'_n := \frac{n + 2\eta_n - \sqrt{(n + 2\eta_n)^2 - 8\eta_n}}{2}$$

with $\eta_n = n - 2$. We see that σ'_n and τ'_n satisfy the following equalities

$$\sigma_n'^2 - (n + \eta_n)\sigma'_n + \eta_n = 0, \quad (2.2)$$

$$\tau_n'^2 - (n + 2\eta_n)\tau'_n + 2\eta_n = 0. \quad (2.3)$$

3 Proof of Theorem 1.1

In this section, we assume that G^* is a graph attaining the minimum of $\tau(G)$ among connected non-transmission-regular graphs G of order n . Let \mathbf{x} be the unit positive eigenvector of $Q(G^*)$ corresponding to $\partial^Q(G^*)$. For convenience, we denote $x_{\max} := \max\{x_v \mid v \in V(G^*)\}$ and $x_{\min} := \min\{x_v \mid v \in V(G^*)\}$. Let u and v be respectively the vertices such that $x_u = x_{\max}$ and $x_v = x_{\min}$. By the equation $Q(G^*)\mathbf{x} = \partial^Q(G^*)\mathbf{x}$ with respect to the vertex u , we see

$$\partial^Q(G^*)x_u = D_u x_u + \sum_{w \in V(G^*)} d(u, w)x_w$$

$$\begin{aligned}
&= D_u x_u + \sum_{w \in V(G^*) \setminus \{v\}} d(u, w) x_w + d(u, v) x_v \\
&\leq (2D_{\max} - d(u, v)) x_u + d(u, v) x_v,
\end{aligned}$$

which yields that

$$2D_{\max} - \partial^Q(G^*) \geq d(u, v) \left(1 - \frac{x_v}{x_u}\right) \geq 1 - \frac{x_v}{x_u}.$$

On the other hand, $\tau(G^*) = 2D_{\max} - \partial^Q(G^*) \leq \tau_n$, we obtain

$$\frac{x_{\max}}{x_{\min}} = \frac{x_u}{x_v} \leq \frac{1}{1 - \tau_n}. \quad (3.1)$$

Theorem 1.1 follows from Lemmas 3.1 and 3.2 below.

Lemma 3.1. $\tau(G^*) = \tau_n$.

Proof. By the minimality of $\tau(G^*)$, we have $\tau(G^*) \leq \tau_n$. Noting that

$$(2D_{\max} - \partial^Q(G^*)) \sum_{i=1}^n x_i = 2 \sum_{i=1}^n (D_{\max} - D_i) x_i \geq 2(nD_{\max} - 2W) x_{\min},$$

together with Eqs. (2.1) and (3.1), we find

$$\begin{aligned}
nD_{\max} - 2W &\leq \frac{\tau_n(x_{\min} + (n-1)x_{\max})}{2x_{\min}} \\
&= \frac{\tau_n}{2} \left(1 + (n-1) \cdot \frac{x_{\max}}{x_{\min}}\right) \\
&\leq \frac{\tau_n(n - \tau_n)}{2(1 - \tau_n)} = \gamma_n.
\end{aligned} \quad (3.2)$$

Hence, $nD_{\max} - 2W = 1$ if n is odd; If n is even, then $nD_{\max} - 2W$ is an even positive integer and equals to 2. Anyway, we have

$$nD_{\max} - 2W = \gamma_n. \quad (3.3)$$

Thus, the first equality of Eq. (3.2) holds and implies $\tau(G^*) = \tau_n$. \square

Lemma 3.2. *If n is odd, then $G^* \cong K_{1,2,\dots,2}$; If n is even, then G^* is isomorphic to some $(n-4)$ -DVDR graph.*

Proof. By the first equality of Eq. (3.2), there are $n-1$ vertices attaining x_{\max} and the remaining vertex attains x_{\min} . Our proof relies on the the following two claims.

Claim 1. There are $n-1$ vertices attaining D_{\max} .

Let $S = \{w \in V(G^*) : D_w = D_{\max}\}$. From Eq. (3.3), we infer that $|S| = n-1$ if n is odd, and $|S| \in \{n-1, n-2\}$ if n is even. Suppose that $|S| = n-2$, in which case the remaining two vertices both attain D_{\min} . There must be a vertex u such that $x_u = x_{\max}$ and $D_u = D_{\min}$. However, the eigenvalue equation

$$2D_{\min} x_u < \partial^Q(G^*) x_u = \sum_{w \in V(G^*)} d(u, w) x_w + D_u x_u \leq 2D_u x_u$$

implies $D_{\min} < D_u$. A contradiction! Thus, $|S| = n - 1$ for either odd or even n .

Claim 2. $D_{\max} = n + \gamma_n - 1$ and $D_{\min} = n - 1$.

By the proof of Claim 1, the $n - 1$ vertices attaining x_{\max} also attain D_{\max} . Recalling that $x_v = x_{\min}$, we have $D_v = D_{\min} = D_{\max} - \gamma_n$ by Eq. (3.3). It can be derived from the eigenvalue equation that

$$\begin{aligned} (2D_{\max} - \tau_n)x_{\min} &= \partial^Q(G^*)x_v = \sum_{u \in V(G^*)} d(v, u)x_u + D_v x_v \\ &= D_v x_u + D_v x_v = (D_{\max} - \gamma_n)(x_{\max} + x_{\min}). \end{aligned}$$

Besides, the equalities in Eq. (3.2) imply $\frac{x_{\max}}{x_{\min}} = \frac{1}{1-\tau_n}$. Together with Eq. (2.1), it follows that

$$\begin{aligned} D_{\max} &= \frac{\gamma_n(1 + \frac{x_{\max}}{x_{\min}}) - \tau_n}{\frac{x_{\max}}{x_{\min}} - 1} = \frac{\frac{\gamma_n(2-\tau_n)}{1-\tau_n} - \tau_n}{\frac{\tau_n}{1-\tau_n}} \\ &= \tau_n - \gamma_n + \frac{2\gamma_n}{\tau_n} - 1 = n - 1 + \gamma_n. \end{aligned}$$

Hence, $D_{\min} = D_{\max} - \gamma_n = n - 1$.

Now we continue the proof of the lemma. Since $D_v = D_{\min} = n - 1$, we have $d(v) = n - 1$, $d(u, v) = 1$ and $d(u, w) \leq 2$ for all $u, w \in V(G^*) \setminus \{v\}$, which yields that

$$D_u = \sum_{w \in V(G^*)} d(u, w) = d(u) + 2(n - 1 - d(u)) = 2(n - 1) - d(u).$$

Therefore, for each vertex $u \in V(G^*) \setminus \{v\}$,

$$d(u) = 2(n - 1) - D_u = n - 1 - \gamma_n = \begin{cases} n - 2, & n \text{ is odd;} \\ n - 3, & n \text{ is even.} \end{cases}$$

Hence, $G^* \cong K_{1,2,\dots,2}$ if n is odd, and G^* is isomorphic to some $(n - 4)$ -DVDR graph if n is even. \square

4 Proof of Theorem 1.2

Suppose that D_{\max} and W are respectively the maximum transmission and Wiener index of a tree T . It is easy to see that

$$nD_{\max} - 2W = \sum_{i=1}^n (D_{\max} - D_i).$$

Let $e = v_1 v_2$ be a leaf of T with $d(v_1) = 1$. Since $d(v_1, v) = d(v_2, v) + 1$ for any vertex $v \in V(T) \setminus \{v_1, v_2\}$, we have $D_{v_1} - D_{v_2} = n - 2$. Thus,

$$nD_{\max} - 2W \geq D_{\max} - D_{v_2} \geq D_{v_1} - D_{v_2} = n - 2 = \eta_n. \quad (4.1)$$

We prove Theorem 1.2 for cases of $\tau(T)$ and $\sigma(T)$ separately by Lemmas 4.1 and 4.2 below.

Assume that T^* is a tree attaining the minimum of $\tau(T)$ among all trees T of order n . Let \mathbf{x} be the unit positive eigenvector of $Q(T^*)$ corresponding to $\partial^Q(T^*)$. For convenience, we still denote $x_{\max} := \max\{x_v \mid v \in V(T^*)\}$ and $x_{\min} := \min\{x_v \mid v \in V(T^*)\}$. Let u and v be respectively the vertices such that $x_u = x_{\max}$ and $x_v = x_{\min}$. By a similar derivation of Eq. (3.1), we get

$$\frac{x_{\max}}{x_{\min}} = \frac{x_u}{x_v} \leq \frac{1}{1 - \tau'_n}. \quad (4.2)$$

Lemma 4.1. $\tau(T^*) = \tau'_n$ and $T^* \cong K_{1, n-1}$.

Proof. By the minimality of $\tau(T^*)$, we have $\tau(T^*) \leq \tau'_n$. Noting that

$$(2D_{\max} - \partial^Q(T^*)) \sum_{i=1}^n x_i = 2 \sum_{i=1}^n (D_{\max} - D_i)x_i \geq 2(nD_{\max} - 2W)x_{\min},$$

together with Eqs. (2.3) and (4.2), we see

$$\begin{aligned} nD_{\max} - 2W &\leq \frac{\tau'_n(x_{\min} + (n-1)x_{\max})}{2x_{\min}} \\ &= \frac{\tau'_n}{2} \left(1 + (n-1) \cdot \frac{x_{\max}}{x_{\min}} \right) \\ &\leq \frac{\tau'_n(n - \tau'_n)}{2(1 - \tau'_n)} = \eta_n. \end{aligned} \quad (4.3)$$

Hence, by Eq. (4.1), we have

$$nD_{\max} - 2W = \eta_n. \quad (4.4)$$

Thus, all equalities of Eq. (4.3) hold. We get $\tau(T^*) = \tau'_n$ and there are $n-1$ vertices attaining x_{\max} and the remaining vertex attains x_{\min} in T^* . For the same reason of Claim 1 of Lemma 3.2, the $n-1$ vertices attaining x_{\max} do not attain D_{\min} . By the argument of Eq. (4.1), the equality (4.4) implies that v is the unique vertex attaining D_{\min} and the other $n-1$ vertices attain D_{\max} , i.e., $D_v = D_{\min} = D_{\max} - \eta_n$. Noting the similarity of Eq. (2.3) and Eq. (2.1), a similar deduction as Claim 2 of Lemma 3.2 can conclude $D_{\max} = n-1 + \eta_n$ and $D_{\min} = D_{\max} - \eta_n = n-1$.

For the tree T^* , $D_v = D_{\min} = n-1$ forces $d(u, v) = 1$, which means v is adjacent to any other vertex of T^* . Hence, $T^* \cong K_{1, n-1}$. \square

In the following, we still denote by T^* an extremal tree attaining the minimum of $\sigma(T)$ among all trees of order n and \mathbf{x} is the unit positive eigenvector of $D(T^*)$ corresponding to $\partial(T^*)$. Let the vertices u, v such that $x_u = x_{\max}$ and $x_v = x_{\min}$ have the same meaning as before. By the equation $D(G^*)\mathbf{x} = \partial\mathbf{x}$ with respect to the vertex u , we see

$$\partial(T^*)x_u = \sum_{w \in V(G^*)} d(u, w)x_w$$

$$\begin{aligned}
&= \sum_{w \in V(G^*) \setminus \{v\}} d(u, w)x_w + d(u, v)x_v \\
&\leq (D_{\max} - d(u, v))x_u + d(u, v)x_v,
\end{aligned}$$

which yields that

$$D_{\max} - \partial(T^*) \geq d(u, v) \left(1 - \frac{x_v}{x_u}\right) \geq 1 - \frac{x_v}{x_u}.$$

On the other hand, $\sigma(T^*) = D_{\max} - \partial(T^*) \leq \sigma'_n$, we obtain

$$\frac{x_{\max}}{x_{\min}} = \frac{x_u}{x_v} \leq \frac{1}{1 - \sigma'_n}. \quad (4.5)$$

Lemma 4.2. $\sigma(T^*) = \sigma'_n$ and $T^* \cong K_{1, n-1}$.

Proof. By the eigenvalue equation

$$(D_{\max} - \partial(T^*)) \sum_{i=1}^n x_i = \sum_{i=1}^n (D_{\max} - D_i)x_i \geq (nD_{\max} - 2W)x_{\min},$$

together with Eqs. (2.2) and (4.4), we find

$$\begin{aligned}
nD_{\max} - 2W &\leq \frac{\sigma'_n(x_{\min} + (n-1)x_{\max})}{x_{\min}} \\
&= \sigma'_n \left(1 + (n-1) \cdot \frac{x_{\max}}{x_{\min}}\right) \\
&\leq \frac{\sigma'_n(n - \sigma'_n)}{1 - \sigma'_n} = \eta_n.
\end{aligned} \quad (4.6)$$

Hence, by Eq. (4.1), we have

$$nD_{\max} - 2W = \eta_n. \quad (4.7)$$

Thus, all equalities of Eq. (4.6) hold. We get $\sigma(T^*) = \sigma'_n$ and there are $n-1$ vertices attaining x_{\max} and the remaining vertex attains x_{\min} in T^* . By the argument of Eq. (4.1), the equality (4.7) implies that v is the unique vertex attaining D_{\min} and the other $n-1$ vertices attain D_{\max} , i.e., $D_v = D_{\min} = D_{\max} - \eta_n$. Using the eigenvalue equation, we have

$$(D_{\max} - \sigma'_n)x_{\min} = \partial(T^*)x_v = \sum_{u \in V(G^*)} d(v, u)x_u = D_v x_u = (D_{\max} - \eta_n)x_{\max}.$$

The equalities in Eq. (4.6) imply $\frac{x_{\max}}{x_{\min}} = \frac{1}{1 - \sigma'_n}$. Together with Eq. (2.2), it follows that

$$D_{\max} = \frac{\eta_n \cdot \frac{x_{\max}}{x_{\min}} - \sigma'_n}{\frac{x_{\max}}{x_{\min}} - 1} = \frac{\frac{\eta_n}{1 - \sigma'_n} - \sigma'_n}{\frac{\sigma'_n}{1 - \sigma'_n}} = \sigma'_n + \frac{\eta_n}{\sigma'_n} - 1 = n - 1 + \eta_n.$$

Hence, $D_v = D_{\min} = D_{\max} - \eta_n = n - 1$, which means $T^* \cong K_{1, n-1}$. \square

5 Concluding Remark

Although it is easy to see that $\sigma(G) < \tau(G)$ for non-transmission-regular graphs, the results of this paper show that the extremal graphs attaining minimum $\sigma(G)$ and those attaining minimum $\tau(G)$ are the same, and the extremal trees attaining minimal $\sigma(T)$ and those attaining minimal $\tau(T)$ are also the same. This may mean that the two indicators for measuring non-transmission-regular graphs are similar to some extent. In addition, one can find that the the minimum values σ_n [8] and τ_n decrease and converge to zero as n increases to infinity, while σ'_n and τ'_n increase as n increases. This means that there is a big gap between trees and cyclic graphs in terms of σ and τ . However, we know that the stars take the maximal value of $\Delta(G) - \rho(G)$. Our results reveal that σ and τ are quite different from $\Delta(G) - \rho(G)$ in measuring the irregularity of graphs.

References

- [1] Atik F., Panigrahi P.: On the distance and distance signless Laplacian eigenvalues of graphs and the smallest Gersgorin disc, *Electron. J. Linear Algebra* 34, 191–204 (2018)
- [2] Brouwer A.E., Haemers W.H.: *Spectra of Graphs*, Springer, Berlin (2011)
- [3] Chen X., Hou Y.: The extreme eigenvalues and maximum degree of k -connected irregular graphs, *Linear Algebra Appl.* 463, 33–44 (2014)
- [4] Cioabă S.M., Gregory D.A., Nikiforov V.: Extreme eigenvalues of nonregular graphs, *J. Combin. Theory Ser. B* 97, 483–486 (2007)
- [5] Collatz L., Sinogowitz U.: *Spektren endlicher Grafen*, *Abh. Math. Sem. Univ. Hamburg* 21, 63–77 (1957)
- [6] Cvetkovic D., Rowlinson P., Simic S.: *An Introduction to the Theory of Graph Spectra*, Cambridge University Press, (2010)
- [7] Liu B., Huang Y., You Z.: On λ_1 -extremal non-regular graphs, *Electron. J. Linear Algebra* 18, 735–744 (2009)
- [8] Liu L.L., Shan H.Y., He C.X.: A proof of a conjecture on the distance spectral radius and maximum transmission of graph, *Graphs Combin.* 38, 49 (2022)
- [9] Liu S., Shu J., Xue J.: On the largest distance (signless Laplacian) eigenvalue of non-transmission-regular graphs, *Electron. J. Linear Algebra* 34, 459–471 (2018)
- [10] Minc H.: *Nonnegative Matrices*, John Wiley, Sons Inc, New York (1988)
- [11] Ning W., Li H., Lu M.: On the signless Laplacian spectral radius of irregular graphs, *Linear Algebra Appl.* 438, 2280–2288 (2013)

- [12] Shiu W., Huang P., Sun P.: The signless Laplacian spectral radius of k -connected irregular graphs, *Linear Multilinear Algebra* 65, 830–839 (2017)
- [13] Shi L.S.: The spectral radius of irregular graphs, *Linear Algebra Appl.* 431, 189–196 (2009)
- [14] Zhang X.: Eigenvectors and eigenvalues of nonregular graphs, *Linear Algebra Appl.* 409, 79–86 (2005)