

A note on the number of plane partitions and r -component multipartitions of n

Mircea Cimpoeaş¹ and Alexandra Teodor²

Abstract

Using elementary methods, we prove new formulas for $\text{pp}(n)$, the number of plane partitions of n , $\text{pp}_r(n)$, the number of plane partitions of n with at most r rows, $\text{pp}^s(n)$, the number of strict plane partitions of n and $\text{pp}^{so}(n)$, the number of symmetric plane partitions of n . Also, we give new formulas for $P_r(n)$, the number of r -component multipartitions of n .

Keywords: Integer partition, Restricted partition function, Plane partition, Multipartition.

MSC2010: 11P81, 11P83.

1 Introduction

Let n be a positive integer. We denote $[n] = \{1, 2, \dots, n\}$. A partition of n is a non-increasing sequence $\lambda = (\lambda_1, \dots, \lambda_m)$ of positive integers such that $|\lambda| = \lambda_1 + \dots + \lambda_m = n$. We define $p(n)$ as the number of partitions of n and for convenience, we define $p(0) = 1$. This notion has the following generalization: A *plane partition* of n is an array $(n_{ij})_{i,j \in [n]}$ of nonnegative integers such that

$$\sum_{i,j \in [n]} n_{ij} = n \text{ and } n_{ij} \geq n_{i'j'} \text{ for all } i, j, i', j' \in [n] \text{ such that } i \leq i' \text{ and } j \leq j'.$$

If $n_{ij} > n_{i(j+1)}$ whenever $n_{ij} \neq 0$, then we shall call such a partition *strict*. If $n_{ij} = n_{ji}$ for all i and j , then the partition is called *symmetric*.

For example, there are 6 plane partitions of $n = 3$, namely:

$$\begin{array}{cccccc} 3 & 0 & 0 & 2 & 1 & 0 & 2 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0, & 0 & 0 & 0, & 1 & 0 & 0, & 0 & 0 & 0, & 1 & 0 & 0, & 1 & 0 & 0. \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{array}$$

¹Mircea Cimpoeaş, University Politehnica of Bucharest, Faculty of Applied Sciences, Bucharest, 060042, Romania and Simion Stoilow Institute of Mathematics, Research unit 5, P.O.Box 1-764, Bucharest 014700, Romania, E-mail: mircea.cimpoeas@upb.ro, mircea.cimpoeas@imar.ro

²Alexandra Teodor, University Politehnica of Bucharest, Faculty of Applied Sciences, Bucharest, 060042, E-mail: alexandra.teodor@upb.ro

Note that four of them are strict partitions and two of them are symmetric. Also, three of them have nonzero entries only on the first row and five of them have nonzero entries on the first two rows.

We denote $\text{pp}(n)$ the total number of plane partitions of n and, we define $\text{pp}(0) = 1$. The properties of $\text{pp}(n)$ have been extensively studied in literature. As a curiosity, we mention that the function $\text{pp}(n)$ appears in physics in connection with the enumeration of small black holes in string theory, see [5, Appendix E].

Let $k \geq 1$ be an integer. We denote $\text{pp}_r(n)$, the number of plane partitions with at most r rows, and $\text{pp}_k(0) = 1$. Note that $\text{pp}_1(n) = \text{p}(n)$ for all $n \geq 0$. Also, if $k \geq n$ then $\text{pp}_k(n) = \text{pp}(n)$. In the example above, we have $\text{pp}_1(3) = \text{p}(3) = 3$, $\text{pp}_2(3) = 5$ and $\text{pp}_3(3) = \text{pp}(3) = 6$.

We denote $\text{pp}^s(n)$, the number of strict plane partitions of n , and $\text{pp}^{so}(n)$, the number of strict plane partitions of n with odd parts. We set $\text{pp}^s(0) = \text{pp}^{so}(0) = 1$. It is well known that $\text{pp}^{so}(n)$ counts also the number of symmetric plane partitions of n .

Let $\mathbf{a} := (a_1, a_2, \dots, a_r)$ be a sequence of positive integers, $r \geq 1$. The *restricted partition function* associated to \mathbf{a} is $p_{\mathbf{a}} : \mathbb{N} \rightarrow \mathbb{N}$, $p_{\mathbf{a}}(n) :=$ the number of integer solutions (x_1, \dots, x_r) of $\sum_{i=1}^r a_i x_i = n$ with $x_i \geq 0$. Note that the generating function of $p_{\mathbf{a}}(n)$ is

$$\sum_{n=0}^{\infty} p_{\mathbf{a}}(n) z^n = \frac{1}{(1 - z^{a_1}) \cdots (1 - z^{a_r})}, \quad |z| < 1. \quad (1.1)$$

See [1, Chapter 5] for further details.

Our aim is to provide new formulas for $\text{pp}(n)$, $\text{pp}_r(n)$, $\text{pp}^s(n)$ and $\text{pp}^{so}(n)$ using their generating functions and the relation with the restricted partition function $p_{\mathbf{a}}(n)$; see Proposition 2.1. In Theorem 2.2, we prove a new formula for $\text{pp}(n)$ in terms of coefficients of the polynomials

$$f_s(z) = (1 + z + \cdots + z^{\alpha_s})^s, \quad \text{where } \alpha_s = \frac{D_n}{s} - 1,$$

where $1 \leq s \leq n$ and D_n is the least common multiple of $1, 2, \dots, n$. Similarly, we provide formulas for $\text{pp}_r(n)$, $\text{pp}^s(n)$ and $\text{pp}^{so}(n)$ in Theorem 2.5, Theorem 2.7 and Theorem 2.9, respectively. Also, using a result from [3] regarding the restricted partition function, we deduce other formulas for $\text{pp}(n)$, $\text{pp}_r(n)$, $\text{pp}^s(n)$ and $\text{pp}^{so}(n)$; see Theorem 2.12.

A r -component multipartition of n is a r -tuple $\lambda = (\lambda^1, \dots, \lambda^r)$ of partitions of n such that $|\lambda| = |\lambda^1| + \cdots + |\lambda^r| = n$; see [2]. We denote $P_r(n)$, the number of r -component multipartitions of n and $P_r(0) = 1$. In Proposition 3.4 we show that

$$\text{pp}_r(n) = \sum_{\substack{0 \leq t_1 \leq r-1 \\ \vdots \\ 0 \leq t_{r-1} \leq 1}} (-1)^{t_1 + \cdots + t_{r-1}} P_r(n - t_1 - 2t_2 - \cdots - (r-1)t_{r-1}),$$

where $P_r(j) = 0$ for $j < 0$.

In Theorem 3.2 we prove a new formula for $P_r(n)$ and we deduce from it a new expression for $\text{pp}_r(n)$; see Corollary 3.5. Also, in Theorem 3.6 we obtain other formula for $P_r(n)$.

2 New formulas for the number of plane partitions

- Let n, k be two positive integers.
- Let $\text{pp}(n)$ be the number of plane partitions of n . We define $\text{pp}(0) = 1$.
- Let $\text{pp}_r(n)$ be the number of plane partitions of n with at most r rows. We also define $\text{pp}_r(0) = 1$. Note that, if $n \leq r$ then $\text{pp}(n) = \text{pp}_r(n)$.
- Let $\text{pp}^s(n)$ be the number of strict plane partitions of n . We set $\text{pp}^s(0) = 1$.
- Let $\text{pp}^{so}(n)$ be the number of strict plane partitions of n with odd parts. We set $\text{pp}^{so}(n) = 1$. As it was shown in [6], $\text{pp}^{so}(n)$ is equal to the number of symmetric plane partitions of n .

MacMahon [8] proved that

$$\sum_{n=0}^{\infty} \text{pp}(n)z^n = \prod_{n=1}^{\infty} \frac{1}{(1-z^n)^n} \text{ for } |z| < 1. \quad (2.1)$$

A refinement of this result is the following

$$\sum_{n=0}^{\infty} \text{pp}_r(n)z^n = \prod_{n=1}^{\infty} \frac{1}{(1-z^n)^{\min\{n,r\}}} \text{ for } |z| < 1, \quad (2.2)$$

see [1, Equation (10.1)].

Gordon and Houten [7] proved that

$$\sum_{n=0}^{\infty} \text{pp}^s(n)z^n = \prod_{n=1}^{\infty} \frac{1}{(1-z^n)^{\lfloor (n+1)/2 \rfloor}} \text{ for } |z| < 1. \quad (2.3)$$

Also, Gordon [6] proved that

$$\sum_{n=0}^{\infty} \text{pp}^{so}(n)z^n = \prod_{n=1}^{\infty} \frac{1}{(1-z^{2n+1})} \prod_{n=1}^{\infty} \frac{1}{(1-z^{2n})^n} \text{ for } |z| < 1. \quad (2.4)$$

Give n, k two positive integers, we denote $n^{[k]}$ the sequence n, n, \dots, n of length k , e. g. $2^{[3]} = 2, 2, 2$. We consider the following sequences of integers:

$$\begin{aligned} \mathbf{n} &:= (1, 2^{[2]}, 3^{[3]}, \dots, n^{[n]}), \\ \mathbf{n}_r &:= (1, 2^{[\min\{2,r\}]}, 3^{[\min\{3,r\}]}, \dots, n^{[\min\{n,r\}]}), \\ \mathbf{n}^s &:= (1, 2^{[1]}, 3^{[2]}, 4^{[2]}, \dots, n^{[\lfloor (n+1)/2 \rfloor]}), \text{ and} \\ \mathbf{n}^{so} &:= (1, 2^{[1]}, 3^{[1]}, 4^{[2]}, \dots, n^{[\lambda(n)]}), \end{aligned}$$

$$\text{where } \lambda(n) = \begin{cases} 1, & n \text{ is odd} \\ \frac{n}{2}, & n \text{ is even} \end{cases}.$$

Proposition 2.1. *Let n, k be two positive integers. We have that:*

- (1) $\text{pp}(n) = p_{\mathbf{n}}(n)$ for all $n \geq 0$.
- (2) $\text{pp}_r(n) = p_{\mathbf{n}_r}(n)$ for all $n \geq 0$.
- (3) $\text{pp}^s(n) = p_{\mathbf{n}^s}(n)$ for all $n \geq 0$.
- (4) $\text{pp}^{so}(n) = p_{\mathbf{n}^{so}}(n)$ for all $n \geq 0$.

Proof. (1) From (1.1), it follows that

$$\sum_{m=0}^{\infty} p_{\mathbf{n}}(m)z^m = \prod_{j=1}^n \frac{1}{(1-z^j)^j} = (1+z+z^2+\dots)(1+z^2+z^4+\dots)\cdots(1+z^n+z^{2n}+\dots). \quad (2.5)$$

On the other hand, from (2.1), we have

$$\sum_{m=0}^{\infty} \text{pp}(m)z^m = (1+z+z^2+\dots)(1+z^2+z^4+\dots)\cdots(1+z^n+z^{2n}+\dots)(1+z^{n+1}+\dots)\cdots. \quad (2.6)$$

Comparing (2.5) and (2.6), it follows that

$$\text{pp}(m) = p_{\mathbf{n}}(m) \text{ for all } 0 \leq m \leq n,$$

thus, in particular, $\text{pp}(n) = p_{\mathbf{n}}(n)$, as required.

(2,3,4) The proof is similar to the proof of (1), so we omit it. \square

Let D_n be the least common multiple of $1, 2, \dots, n$. For all $1 \leq s \leq n$, we define

$$f_s(z) := (1+z+\dots+z^{\alpha_s})^s, \text{ where } \alpha_s = \frac{D_n}{s} - 1.$$

We write

$$f_s(z) = f_{s,0} + f_{s,1}z + \dots + f_{s,s\alpha_s}z^{s\alpha_s}. \quad (2.7)$$

From the definition of $f_s(z)$, we note that

$$f_{s,\ell} = \#\{(j_1, \dots, j_s) : j_1 + \dots + j_s = \ell, \text{ where } 0 \leq j_t \leq \alpha_s, 1 \leq t \leq s\}. \quad (2.8)$$

Note that the polynomial $f_s(z)$ is reciprocal, that is,

$$x^{s\alpha_s} f_s\left(\frac{1}{x}\right) = f_s(x),$$

so that $f_{s,i} = f_{s,s\alpha_s-i}$ for all i . On the other hand, we have

$$f_s(z) = \left(\frac{1-z^{\frac{D_n}{s}}}{1-z}\right)^s = (1-z^{\frac{D_n}{s}})^s (1-z)^{-s}.$$

Using the binomial expansion of $(1 - z)^{-s}$, it follows that

$$f_s(z) = \sum_{i=0}^s (-1)^i \binom{s}{i} z^{i \frac{D_n}{s}} \times \sum_{j=0}^{\infty} \binom{j+s-1}{j} z^j. \quad (2.9)$$

Comparing (2.7) with (2.9) we deduce that:

$$f_{s,\ell} = \sum_{i,j \geq 0 \text{ with } \frac{iD_n}{s} + j = \ell} (-1)^i \binom{s}{i} \binom{j+s-1}{j}, \quad (2.10)$$

for all $0 \leq \ell \leq s\alpha_s$.

Theorem 2.2. *Let $n \geq 3$ be an integer. We have that*

$$\text{pp}(n) = \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_n} \prod_{s=2}^n \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ \frac{i_s D_n}{s} + j_s = \ell_s}} (-1)^{i_s} \binom{s}{i_s} \binom{j_s + s - 1}{j_s},$$

where $\mathbf{A}_n := \{(\ell_1, \dots, \ell_n) \in \mathbb{N}^n : \ell_1 + 2\ell_2 + \dots + n\ell_n = n\}$.

Proof. From Proposition 2.1(1) it follows that

$$\text{pp}(n) = \#\{(j_1, \dots, j_{\binom{n+1}{2}}) \in \mathbb{N}^{\binom{n+1}{2}} : j_1 + 2j_2 + 2j_3 + \dots + nj_{\binom{n}{2}+1} + \dots + nj_{\binom{n+1}{2}} = n\}. \quad (2.11)$$

We denote $\ell_1 = j_1$, $\ell_2 = j_2 + j_3, \dots, \ell_n = j_{\binom{n}{2}+1} + \dots + j_{\binom{n+1}{2}}$.

Since $n \geq 3$, it follows that

$$\ell_s \leq \frac{n}{s} \leq \frac{D_n}{s} - 1 \text{ for all } 2 \leq s \leq n.$$

Hence, from (2.8) and (2.11) it follows that

$$\text{pp}(n) = \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_n} \prod_{s=2}^n f_{s,\ell_s}. \quad (2.12)$$

Therefore, from (2.12) and (2.10) we get the required formula. \square

Remark 2.3. Note that $p(n) = \#\mathbf{A}_n$ for all $n \geq 3$.

Example 2.4. If we take $n = 3$ in Theorem 2.2, we have $\mathbf{A}_3 = \{(3, 0, 0), (1, 1, 0), (0, 0, 1)\}$ and thus

$$\text{pp}(3) = \sum_{(\ell_1, \ell_2, \ell_3) \in \mathbf{A}_3} \prod_{s=2}^3 \sum_{\frac{6i_s}{s} + j_s = \ell_s} (-1)^{i_s} \binom{j_s + s - 1}{j_s} = 1 + \binom{1}{1} \binom{2}{1} + \binom{3}{1} = 6.$$

Theorem 2.5. *Let $n > r \geq 2$ be two positive integers. We have that*

$$\text{pp}_r(n) = \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_n} \prod_{s=2}^n \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ \frac{i_s D_n}{\min\{s, r\}} + j_s = \ell_s}} (-1)^{i_s} \binom{\min\{s, r\}}{i_s} \binom{j_s + \min\{s, r\} - 1}{j_s}.$$

Proof. From Proposition 2.1(2) it follows that

$$\begin{aligned} \text{pp}_r(n) = \#\{(j_1, \dots, j_{rn-\binom{r}{2}}) \in \mathbb{N}^{rn-\binom{r}{2}} : j_1 + 2j_2 + 2j_3 + \dots + rj_{\binom{r}{2}+1} + \dots + rj_{\binom{r+1}{2}} + \\ + \dots + nj_{rn-\binom{r+1}{2}+1} + \dots + nj_{rn-\binom{r}{2}} = n\}. \end{aligned} \quad (2.13)$$

We denote $\ell_1 = j_1$, $\ell_2 = j_2 + j_3, \dots, \ell_n = j_{rn-\binom{r+1}{2}+1} + \dots + j_{rn-\binom{r}{2}}$.

Since $n \geq 3$ and $r \geq 2$ it follows that

$$\ell_s \leq \frac{n}{s} \leq \frac{D_n}{s} - 1 \leq \frac{D_n}{\min\{r, s\}} - 1 \text{ for all } 2 \leq s \leq n.$$

Hence, from (2.13) and (2.8) it follows that:

$$\text{pp}_r(n) = \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_n} \prod_{s=2}^n f_{\min\{r, s\}, \ell_s} \quad (2.14)$$

Therefore, from (2.14) and (2.10) we get the required formula. \square

Example 2.6. Let $n = 3$ and $r = 2$. Since $\mathbf{A}_3 = \{(3, 0, 0), (1, 1, 0), (0, 0, 1)\}$, according to Theorem 2.5, the number of plane partitions of 3 with at most 2 rows is:

$$\text{pp}_2(3) = \sum_{(\ell_1, \ell_2, \ell_3) \in \mathbf{A}_3} \prod_{s=2}^3 \sum_{i_s, j_s \geq 0, 3i_s + j_s = \ell_s} \binom{2}{i_s} \binom{j_s + 1}{j_s} = 1 + \binom{2}{0} \binom{2}{1} + \binom{2}{0} \binom{2}{1} = 5.$$

Theorem 2.7. *Let $n \geq 3$ be an integer. We have that*

$$\text{pp}^s(n) = \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_n} \prod_{s=3}^n \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ \frac{i_s D_n}{\lfloor \frac{s+1}{2} \rfloor} + j_s = \ell_s}} (-1)^{i_s} \binom{\lfloor \frac{s+1}{2} \rfloor}{i_s} \binom{j_s + \lfloor \frac{s+1}{2} \rfloor - 1}{j_s}.$$

Proof. Assume $n = 2p$. From Proposition 2.1(3) it follows that

$$\text{pp}^s(n) = \#\{(j_1, \dots, j_{p^2+p}) : j_1 + 2j_2 + 3j_3 + 3j_4 + \dots + nj_{p^2+1} + \dots + nj_{p^2+p} = n\} \quad (2.15)$$

We denote $\ell_1 = j_1$, $\ell_2 = j_2$, $\ell_3 = j_3 + j_4, \dots, \ell_n = j_{p^2+1} + \dots + j_{p^2+p}$.

Since $n \geq 3$ and $k \geq 2$ it follows that

$$\ell_s \leq \frac{n}{s} \leq \frac{D_n}{s} - 1 \leq \frac{D_n}{\lfloor \frac{s+1}{2} \rfloor} - 1 \text{ for all } 1 \leq s \leq n.$$

Hence, from (2.15) and (2.8) it follows that:

$$\text{pp}^s(n) = \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_n} \prod_{s=3}^n f_{\lfloor \frac{s+1}{2} \rfloor, \ell_s} \quad (2.16)$$

Therefore, from (2.16) and (2.10) we get the required formula. The case $n = 2p + 1$ is similar. \square

Example 2.8. Let $n = 3$. Since $\mathbf{A}_3 = \{(3, 0, 0), (1, 1, 0), (0, 0, 1)\}$, according to Theorem 2.7, the number of strict plane partitions of 3 is:

$$\text{pp}^s(3) = \sum_{(\ell_1, \ell_2, \ell_3) \in \mathbf{A}_3} \sum_{i_3, j_3 > 0, 3i_3 + j_3 = \ell_3} (-1)^{i_3} \binom{2}{i_3} \binom{j_3 + 1}{j_3} = 1 + 1 + \binom{2}{0} \binom{2}{1} = 4.$$

Theorem 2.9. Let $n \geq 3$ be a positive integer. We have that

$$\text{pp}^{so}(n) = \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_n} \prod_{s=2}^{\lfloor \frac{n}{2} \rfloor} \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ i_s \frac{D_n}{s} + j_s = \ell_{2s}}} (-1)^{i_s} \binom{s}{i_s} \binom{j_s + s - 1}{j_s}.$$

Proof. Assume $n = 2p$. From Proposition 2.1(4) it follows that

$$\begin{aligned} \text{pp}^s(n) = \#\{(j_1, \dots, j_{p^2+p}) : j_1 + 2j_2 + 3j_3 + 4j_4 + 4j_5 + 5j_6 + \dots + (n-1)j_{\binom{p}{2}} + \\ + nj_{\binom{p}{2}+1} + \dots + nj_{\binom{p+1}{2}} = n\}. \end{aligned} \quad (2.17)$$

We denote $\ell_1 = j_1$, $\ell_2 = j_2$, $\ell_3 = j_3, \dots$, $\ell_{n-1} = j_{\binom{p}{2}}$, $\ell_n = j_{\binom{p}{2}+1} + \dots + j_{\binom{p+1}{2}}$.

Since $n \geq 3$ and $k \geq 2$ it follows that

$$\ell_{2s} \leq \frac{n}{2s} \leq \frac{D_n}{2s} - 1 \leq \frac{D_n}{s} - 1 \text{ for all } 2 \leq s \leq p.$$

Hence, from (2.15) and (2.8) it follows that:

$$\text{pp}^s(n) = \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_n} \prod_{s=2}^{\lfloor \frac{n}{2} \rfloor} f_{s, \ell_{2s}} \quad (2.18)$$

Therefore, from (2.18) and (2.10) we get the required formula. The case $n = 2p + 1$ is similar. \square

Example 2.10. Let $n = 3$. Since $\mathbf{A}_3 = \{(3, 0, 0), (1, 1, 0), (0, 0, 1)\}$ and $\lfloor \frac{3}{2} \rfloor = 1$, according to Theorem 2.9, the number of symmetric plane partition of 3 is:

$$\text{pp}^{so}(3) = p(3) = \#\mathbf{A}_3 = 3.$$

The unsigned Stirling numbers $\begin{bmatrix} r \\ k \end{bmatrix}$'s are defined by

$$\binom{n+r-1}{r-1} = \frac{1}{n(r-1)!} n^{(r)} = \frac{1}{(r-1)!} \left(\begin{bmatrix} r \\ r \end{bmatrix} n^{r-1} + \cdots + \begin{bmatrix} r \\ 2 \end{bmatrix} n + \begin{bmatrix} r \\ 1 \end{bmatrix} \right). \quad (2.19)$$

We recall the following result:

Theorem 2.11. (see [3, Theorem 2.8(2)] and [4]) Let $\mathbf{a} = (a_1, \dots, a_r)$ be sequence of positive integers and let D be the least common multiple of a_1, \dots, a_r . We have that

$$p_{\mathbf{a}}(n) = \frac{1}{(r-1)!} \sum_{m=0}^{r-1} \sum_{\substack{0 \leq j_1 \leq \frac{D}{a_1} - 1, \dots, 0 \leq j_r \leq \frac{D}{a_r} - 1 \\ a_1 j_1 + \cdots + a_r j_r \equiv n \pmod{D}}} \sum_{k=m}^{r-1} \begin{bmatrix} r \\ k+1 \end{bmatrix} (-1)^{k-m} \binom{k}{m} D^{-k} (a_1 j_1 + \cdots + a_r j_r)^{k-m} n^m.$$

Using Theorem 2.11 we are able to obtain new formulas for $pp(n)$, $pp_k(n)$, $pp^s(n)$ and $pp^{so}(n)$, where $n \geq 3$ is an integer:

Theorem 2.12. (1) For $n \geq 3$ we have

$$\begin{aligned} pp(n) &= \frac{1}{\left(\binom{n+1}{2} - 1\right)!} \sum_{m=0}^{\binom{n+1}{2} - 1} \sum_{\substack{0 \leq \ell_1 \leq D_n - 1, \dots, 0 \leq \ell_n \leq D_n - n \\ \ell_1 + 2\ell_2 + \cdots + n\ell_n \equiv n \pmod{D_n}}} \prod_{s=2}^n \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ \frac{i_s D_n}{s} + j_s = \ell_s}} (-1)^{i_s} \times \\ &\times \binom{s}{i_s} \binom{j_s + s - 1}{j_s} \sum_{k=m}^{\binom{n+1}{2} - 1} \begin{bmatrix} \binom{n+1}{2} \\ k+1 \end{bmatrix} (-1)^{k-m} \binom{k}{m} D_n^{-k} (\ell_1 + 2\ell_2 + \cdots + n\ell_n)^{k-m} n^m. \end{aligned}$$

(2) For $n \geq 3$ and $2 \leq r \leq n-1$ we have

$$\begin{aligned} pp_r(n) &= \frac{1}{\left(nr - \binom{r}{2} - 1\right)!} \sum_{m=0}^{nr - \binom{r}{2} - 1} \sum_{\substack{0 \leq \ell_1 \leq D_n - \min\{1, r\}, \dots, 0 \leq \ell_n \leq D_n - \min\{n, r\} \\ \ell_1 + 2\ell_2 + \cdots + n\ell_n \equiv n \pmod{D_n}}} \times \\ &\times \prod_{s=2}^n \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ \frac{i_s D_n}{\min\{s, r\}} + j_s = \ell_s}} (-1)^{i_s} \binom{\min\{s, r\}}{i_s} \binom{j_s + \min\{s, r\} - 1}{j_s} \times \\ &\times \sum_{k=m}^{nr - \binom{r}{2} - 1} \begin{bmatrix} nr - \binom{r}{2} \\ k+1 \end{bmatrix} (-1)^{k-m} \binom{k}{m} D_n^{-k} (\ell_1 + 2\ell_2 + \cdots + n\ell_n)^{k-m} n^m. \end{aligned}$$

(3) For $n \geq 3$ we have

$$\begin{aligned} \text{pp}^s(n) &= \frac{1}{\left(\left\lfloor \frac{n+1}{2} \right\rfloor \cdot \left\lfloor \frac{n+2}{2} \right\rfloor - 1\right)!} \sum_{m=0}^{\left\lfloor \frac{n+1}{2} \right\rfloor \cdot \left\lfloor \frac{n+2}{2} \right\rfloor - 1} \sum_{\substack{0 \leq \ell_1 \leq D_n - 1, \dots, 0 \leq \ell_n \leq D_n - \left\lfloor \frac{n+1}{2} \right\rfloor \\ \ell_1 + 2\ell_2 + \dots + n\ell_n \equiv n \pmod{D_n}}} \times \\ &\times \prod_{s=3}^n \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ \frac{i_s D_n}{\left\lfloor \frac{s+1}{2} \right\rfloor} + j_s = \ell_s}} (-1)^{i_s} \binom{\left\lfloor \frac{s+1}{2} \right\rfloor}{i_s} \binom{j_s + \left\lfloor \frac{s+1}{2} \right\rfloor - 1}{j_s} \times \\ &\times \sum_{k=m}^{\left\lfloor \frac{n+1}{2} \right\rfloor \cdot \left\lfloor \frac{n+2}{2} \right\rfloor - 1} \left[\begin{matrix} \left\lfloor \frac{n+1}{2} \right\rfloor \cdot \left\lfloor \frac{n+2}{2} \right\rfloor \\ k+1 \end{matrix} \right] (-1)^{k-m} \binom{k}{m} D_n^{-k} (\ell_1 + 2\ell_2 + \dots + n\ell_n)^{k-m} n^m. \end{aligned}$$

(4) For $n \geq 3$ we have

$$\begin{aligned} \text{pp}^{so}(n) &= \frac{1}{\left(\left(\left\lfloor \frac{n}{2} \right\rfloor\right) + \varepsilon(n) - 1\right)!} \sum_{m=0}^{\left(\left\lfloor \frac{n}{2} \right\rfloor\right) + \varepsilon(n) - 1} \sum_{\substack{0 \leq \ell_1 \leq D_n - \lambda(1), \dots, 0 \leq \ell_n \leq D_n - \lambda(n) \\ \ell_1 + 2\ell_2 + \dots + n\ell_n \equiv n \pmod{D_n}}} \times \\ &\times \prod_{s=2}^{\left\lfloor \frac{n}{2} \right\rfloor} \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ \frac{i_s D_n}{s} + j_s = \ell_{2s}}} (-1)^{i_s} \binom{s}{i_s} \binom{j_s + s - 1}{j_s} \times \\ &\times \sum_{k=m}^{\left(\left\lfloor \frac{n}{2} \right\rfloor\right) + \varepsilon(n) - 1} \left[\begin{matrix} \left(\left\lfloor \frac{n}{2} \right\rfloor\right) + \varepsilon(n) \\ k+1 \end{matrix} \right] (-1)^{k-m} \binom{k}{m} D_n^{-k} (\ell_1 + 2\ell_2 + \dots + n\ell_n)^{k-m} n^m, \end{aligned}$$

$$\text{where } \lambda(n) = \begin{cases} 1, & n \text{ is odd} \\ \frac{n}{2}, & n \text{ is even} \end{cases} \text{ and } \varepsilon(n) = \begin{cases} 1, & n \text{ is odd} \\ 0, & n \text{ is even} \end{cases}.$$

Proof. (1) Note that the length of the sequence $\mathbf{n} = (1, 2^{[2]}, \dots, n^{[n]})$ is $\binom{n+1}{2}$. From Proposition 2.1(1) and Theorem 2.11 it follows that

$$\begin{aligned} \text{pp}(n) = \text{p}_{\mathbf{n}}(n) &= \frac{1}{\left(\binom{n+1}{2} - 1\right)!} \sum_{m=0}^{\binom{n+1}{2} - 1} \sum_{(j_1, \dots, j_{\binom{n+1}{2}}) \in \mathbf{C}_n} \sum_{k=m}^{\binom{n+1}{2} - 1} \left[\begin{matrix} \binom{n+1}{2} \\ k+1 \end{matrix} \right] \times \\ &\times (-1)^{k-m} \binom{k}{m} D_n^{-k} (j_1 + 2j_2 + 2j_3 + \dots + nj_{\binom{n}{2}+1} + \dots + nj_{\binom{n+1}{2}})^{k-m} n^m, \end{aligned} \quad (2.20)$$

where $\mathbf{C}_n = \{(j_1, \dots, j_{\binom{n+1}{2}}) : 0 \leq j_1 \leq D_n - 1, 0 \leq j_2 \leq \frac{D_n}{2} - 1, 0 \leq j_3 \leq \frac{D_n}{2} - 1, \dots, 0 \leq j_{\binom{n}{2}+1} \leq \frac{D_n}{n} - 1, \dots, 0 \leq j_{\binom{n+1}{2}} \leq \frac{D_n}{n} - 1 \text{ such that } j_1 + 2j_2 + 2j_3 + \dots + nj_{\binom{n}{2}+1} + \dots + nj_{\binom{n+1}{2}} \equiv n \pmod{D_n}\}$.

We let $\ell_1 = j_1$, $\ell_2 = j_2 + j_3$, \dots , $\ell_n = j_{\binom{n}{2}+1} + \dots + j_{\binom{n+1}{2}}$.

Note that if $(j_1, j_2, \dots, j_{\binom{n+1}{2}}) \in \mathbf{C}_n$ then $0 \leq \ell_t \leq D_n - t$ for all $1 \leq t \leq n$ and, moreover, $\ell_1 + 2\ell_2 + \dots + n\ell_n \equiv n \pmod{D_n}$. From (2.20), using a similar argument as in the proof of Theorem 2.2, we get the required result.

(2) Note that the length of the sequence $\mathbf{n}_r := (1, 2^{[\min\{2,r\}], 3^{[\min\{3,r\}]}, \dots, n^{[\min\{n,r\}]}])$ is $\binom{r+1}{2} + r(n-r) = nr - \binom{r}{2}$. The rest of the proof is similar to the proof of (1), using Proposition 2.1(2), Theorem 2.5 and Theorem 2.11.

(3) Note that the length of the sequence $\mathbf{n}^s := (1, 2^{[1]}, 3^{[2]}, 4^{[2]}, \dots, n^{[\lfloor (n+1)/2 \rfloor]})$ is $p^2 + p$ if $n = 2p$ and $(p+1)^2$ if $n = 2p+1$. Hence, in both cases, the length is $\lfloor \frac{n+1}{2} \rfloor \cdot \lfloor \frac{n+2}{2} \rfloor$. The rest of the proof is similar to the proof of (1), using Proposition 2.1(3), Theorem 2.7 and Theorem 2.11.

(4) Note that the length of the sequence $\mathbf{n}^{so} := (1, 2^{[1]}, 3^{[1]}, 4^{[2]}, \dots, n^{[\lambda(n)]})$ is $\binom{p}{2}$ if $n = 2p$ and $\binom{p}{2} + 1$ if $n = 2p+1$. Hence, in both cases, the length is $\binom{\lfloor \frac{n}{2} \rfloor}{2} + \varepsilon(n)$. The rest of the proof is similar to the proof of (1), using Proposition 2.1(4), Theorem 2.9 and Theorem 2.11. \square

3 New formulas for the number of multipartitions with r components

If we denote $P_r(n)$ the number of r -component multipartitions of n , we have that

$$\sum_{n=0}^{\infty} P_r(n) z^n = \prod_{n=1}^{\infty} \frac{1}{(1-z^n)^r}, \quad (3.1)$$

where $P_r(0) = 0$ by convention; see [2]. We consider the sequence

$$\mathbf{n}^r = (1^{[r]}, 2^{[r]}, \dots, n^{[r]}).$$

Proposition 3.1. *With the above notations, we have $P_r(n) = p_{\mathbf{n}^r}(n)$ for all $n \geq 0$.*

Proof. It follows from (1.1) and (3.1), using a similar argument as in the proof of Proposition 2.1(1). \square

Theorem 3.2. *Let $n \geq 4$ and $r \geq 2$ be two positive integers such that $n > r$. We have that*

$$P_r(n) = \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_n} \prod_{s=1}^n \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ \frac{i_s D_n}{r} + j_s = \ell_s}} (-1)^{i_s} \binom{r}{i_s} \binom{j_s + r - 1}{j_s}.$$

Proof. From Proposition 3.1 it follows that

$$P_r(n) = \#\{(j_1, \dots, j_{nr}) : j_1 + \dots + j_r + 2j_{r+1} + \dots + 2j_{2r} + \dots + nj_{nr-r+1} + \dots + nj_{nr} = n\} \quad (3.2)$$

We denote $\ell_1 = j_1 + \cdots + j_r$, $\ell_2 = j_{r+1} + \cdots + j_{2r}$, \dots , $\ell_n = j_{nr-r+1} + \cdots + j_{nr}$.

Since $n \geq 4$ and $n > r \geq 2$ it follows that

$$\ell_s \leq \frac{n}{s} \leq \frac{D_n}{r} - 1 \text{ for all } 1 \leq s \leq n.$$

Hence, from (3.2) and (2.8) it follows that:

$$P_r(n) = \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_n} \prod_{s=1}^n f_{r, \ell_s} \quad (3.3)$$

Therefore, from (3.3) and (2.10) we get the required formula. \square

Example 3.3. Let $n = 4$ and $r = 2$. Since $\mathbf{A}_4 = \{(4, 0, 0, 0), (2, 1, 0, 0), (1, 0, 1, 0), (0, 2, 0, 0), (0, 0, 0, 1)\}$ and $f_{2, \ell} = \ell + 1$ for $0 \leq \ell \leq 4$, from Theorem 3.2 we have that

$$P_2(4) = 5 + 3 \cdot 2 + 2 \cdot 2 + 3 + 2 = 20.$$

Let $\text{pp}_r(n)$ be the number of plane partitions of n with at most r rows. From (2.2) and (3.1) it follows that

$$\begin{aligned} \sum_{n=0}^{\infty} \text{pp}_r(n) z^n &= \prod_{j=1}^{r-1} (1 - z^j)^{r-j} \sum_{n=0}^{\infty} P_r(n) z^n = \\ &= \sum_{\substack{0 \leq t_1 \leq r-1 \\ 0 \leq t_2 \leq r-2 \\ \vdots \\ 0 \leq t_{r-1} \leq 1}} (-1)^{t_1 + \cdots + t_{r-1}} \sum_{n=0}^{\infty} P_r(n - t_1 - 2t_2 - \cdots - (r-1)t_{r-1}) z^n, \end{aligned} \quad (3.4)$$

where $P_r(j) = 0$ for $j < 0$. As a direct consequence of (3.4), we get the following result:

Proposition 3.4. *For all $n \geq 0$ we have that*

$$\text{pp}_r(n) = \sum_{\substack{0 \leq t_1 \leq r-1 \\ \vdots \\ 0 \leq t_{r-1} \leq 1}} (-1)^{t_1 + \cdots + t_{r-1}} P_r(n - t_1 - 2t_2 - \cdots - (r-1)t_{r-1}).$$

As a consequence of Proposition 3.4 and Theorem 3.2, we get:

Corollary 3.5. *For all $n \geq 4$ and $n > r \geq 2$, we have that:*

$$\begin{aligned} \text{pp}_r(n) &= \sum_{\substack{0 \leq t_1 \leq r-1 \\ \vdots \\ 0 \leq t_{r-1} \leq 1 \\ n < 4 + t_1 + 2t_2 + \cdots + (r-1)t_{r-1}}} (-1)^{t_1 + \cdots + t_{r-1}} P_r(n - t_1 - 2t_2 - \cdots - (r-1)t_{r-1}) + \\ &\sum_{\substack{0 \leq t_1 \leq r-1 \\ \vdots \\ 0 \leq t_{r-1} \leq 1 \\ n \geq 4 + t_1 + 2t_2 + \cdots + (r-1)t_{r-1}}} (-1)^{t_1 + \cdots + t_{r-1}} \sum_{(\ell_1, \dots, \ell_n) \in \mathbf{A}_{n-t_1-2t_2-\cdots-(r-1)t_{r-1}}} \prod_{s=1}^n \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ \frac{i_s D_n}{r} + j_s = \ell_s}} (-1)^{i_s} \binom{r}{i_s} \binom{j_s + r - 1}{j_s}. \end{aligned}$$

Theorem 3.6. *Let $n \geq 4$ and $r \geq 2$ be two positive integers such that $n \geq k$. We have that*

$$P_r(n) = \frac{1}{(nr-1)!} \sum_{m=0}^{nr-1} \sum_{\substack{0 \leq \ell_1 \leq D_n-1, \dots, 0 \leq \ell_n \leq D_n-n \\ \ell_1 + 2\ell_2 + \dots + n\ell_n \equiv n \pmod{D_n}}} \prod_{s=1}^n \sum_{\substack{i_s, j_s \geq 0 \text{ with} \\ \frac{i_s D_n}{r} + j_s = \ell_s}} (-1)^{i_s} \binom{r}{i_s} \binom{j_s + k - 1}{j_s} \times \\ \times \sum_{k=m}^{nr-1} \left[\begin{matrix} nr \\ k+1 \end{matrix} \right] (-1)^{k-m} \binom{k}{m} D_n^{-k} (\ell_1 + 2\ell_2 + \dots + n\ell_n)^{k-m} n^m.$$

Proof. From Proposition 3.1 and Theorem 2.11 it follows that

$$P_r(n) = \frac{1}{(nr-1)!} \sum_{m=0}^{nr-1} \sum_{\substack{0 \leq j_1 \leq D_n-1, \dots, 0 \leq j_r \leq D-1 \\ \vdots \\ 0 \leq j_{nr-r+1} \leq \frac{D_n}{n}-1, \dots, 0 \leq j_{nr} \leq \frac{D_n}{n}-1 \\ j_1 + \dots + j_r + \dots + nj_{nr-r+1} + \dots + nj_{nr} \equiv n \pmod{D_n}}} \sum_{k=m}^{nr-1} \left[\begin{matrix} nr \\ k+1 \end{matrix} \right] (-1)^{k-m} \times \\ \times \binom{k}{m} D_n^{-k} (j_1 + \dots + j_r + \dots + nj_{nr-r+1} + \dots + nj_{nr})^{k-m} n^m. \quad (3.5)$$

We let $\ell_1 = j_1 + \dots + j_r, \dots, \ell_n = j_{nr-r+1} + \dots + j_{nr}$. It is easy to see that $0 \leq \ell_t \leq D_n - t$ for $1 \leq t \leq n$. Therefore, from (3.5), using a similar argument as in the proof of Theorem 3.2, it follows the required formula. \square

4 Conclusions

We proved new formulas for $\text{pp}(n)$, the number of plane partitions of n , $\text{pp}_k(n)$, the number of plane partitions with at most k rows of n , $\text{pp}^s(n)$, the number of strict plane partitions of n , $\text{pp}^{so}(n)$, the number of symmetric plane partitions, and $P_k(n)$, the number of multi-partitions with k components.

Further investigations include the study of restricted plane partitions with at most r rows and c columns or, more generally, with a given shape.

Acknowledgments

The first author (Mircea Cimpoeaş) was supported by a grant of the Ministry of Research, Innovation and Digitization, CNCS - UEFISCDI, project number PN-III-P1-1.1-TE-2021-1633, within PNCDI III.

References

- [1] G. E. Andrews, K. Eriksson, *Integer Partitions*, Cambridge University Press, Cambridge, 2004.
- [2] G. E. Andrews G.E. *A Survey of Multipartitions Congruences and Identities*, In: Surveys in Number Theory. Developments in Mathematics (Diophantine Approximation: Festschrift for Wolfgang Schmidt), vol 17. Springer, New York, NY, 2008.
- [3] M. Cimpoeaş, F. Nicolae, *On the restricted partition function*, Ramanujan J. 47, **no. 3**, (2018), 565-588.
- [4] M. Cimpoeaş, F. Nicolae, *Corrigendum to "On the restricted partition function"*, Ramanujan J. 49 **no. 3**, (2019), 699–700.
- [5] A. Dabholkar, F. Denef, G. W. Moore, B. Pioline, *Precision counting of small black holes*, Journal of High Energy Physics **10** (2005), 89pp.
- [6] B. Gordon, *Notes on plane partitions V*, J. Combinatorial Theory Series A, no. **11** (1971), 157-168.
- [7] B. Gordon, L. Houten, *Notes on plane partitions*, J. Combinatorial Theory Series A, no. **4** (1968), 72-99.
- [8] P. A. MacMahon, *Collected papers* Vol.1 (G. E. Andrews Ed.) M.I.T. Press Cambridge, M A, 1978.