

On the power of linear programming for K-means clustering ^{*}

Antonio De Rosa [†] Aida Khajavirad [‡] Yakun Wang [§]

Abstract

In [8], the authors introduced a new linear programming (LP) relaxation for K-means clustering. In this paper, we further investigate both theoretical and computational properties of this relaxation. As evident from our numerical experiments with both synthetic real-world data sets, the proposed LP relaxation is almost always tight; i.e., its optimal solution is feasible for the original nonconvex problem. To better understand this unexpected behaviour, on the theoretical side, we focus on K-means clustering with two clusters, and we obtain sufficient conditions under which the LP relaxation is tight. We further analyze the sufficient conditions when the input is generated according to a popular stochastic model and obtain recovery guarantees for the LP relaxation. We conclude our theoretical study by constructing a family of inputs for which the LP relaxation is never tight. Denoting by n the number of data points to be clustered, the LP relaxation contains $\Omega(n^3)$ inequalities making it impractical for large data sets. To address the scalability issue, by building upon a cutting-plane algorithm together with the GPU implementation of PDLP, a first-order method LP solver, we develop an efficient algorithm that solves the proposed LP and hence the K-means clustering problem, for up to $n \leq 4000$ data points.

Key words. *K-means clustering; Linear programming relaxation; Tightness; Recovery guarantee; Cutting-plane algorithm; First-order methods.*

1 Introduction

Clustering data points into a small number of groups according to some similarity measure is a common task in unsupervised machine learning. K-means clustering, one of the oldest and the most popular clustering techniques, partitions the data points into clusters by minimizing the total squared distance between each data point and the corresponding cluster center. Let $\{x^i\}_{i=1}^n$ denote a set of n data points in \mathbb{R}^m , and denote by K the number of desired clusters. Define a partition of $[n] := \{1, \dots, n\}$ as a family $\{\Gamma_k\}_{k=1}^K$ of non-empty subsets of $[n]$ such that $\Gamma_a \cap \Gamma_b = \emptyset$ for all $a \neq b \in [K]$ and $\cup_{k \in [K]} \Gamma_k = [n]$. Then K-means clustering can be formulated as a combinatorial optimization problem:

$$\min \sum_{k=1}^K \sum_{i \in \Gamma_k} \left\| x^i - \frac{1}{|\Gamma_k|} \sum_{j \in \Gamma_k} x^j \right\|_2^2 \quad (1)$$

^{*}The authors were partially funded by AFOSR grant FA9550-23-1-0123.

[†]Department of Decision Sciences and BIDS, Bocconi University, Via Röntgen 1, 20136, Milano, MI, Italy. E-mail: antonio.derosa@unibocconi.it.

[‡]Department of Industrial and Systems Engineering, Lehigh University, Bethlehem, PA 18015, USA. E-mail: aida@lehigh.edu.

[§]Department of Industrial and Systems Engineering, Lehigh University, Bethlehem, PA 18015, USA. E-mail: yaw220@lehigh.edu.

s.t. $\{\Gamma_k\}_{k \in [K]}$ is a partition of $[n]$.

It is well-known that Problem (1) is NP-hard even when there are only two clusters [3] or when the data points are in \mathbb{R}^2 [19]. The most popular techniques for solving Problem (1) are heuristics such as Lloyd’s algorithm [16], approximation algorithms [13, 10], and convex relaxations [26, 25, 5, 22, 12, 15, 8]. The two prominent types of convex relaxations for K-means clustering are semi-definite programming (SDP) relaxations [25] and linear programming (LP) relaxations [8]. The theoretical properties of SDP relaxations for K-means clustering have been thoroughly studied in the literature [5, 22, 12, 28, 15]. In this paper we investigate the power of LP relaxations for K-means clustering. In the following, we present an alternative formulation for Problem (1), which we use to construct our relaxations.

Consider a partition $\{\Gamma_k\}_{k=1}^K$ of $[n]$; let $\mathbf{1}_{\Gamma_k}$, $k \in [K]$ be the indicator vector of the k th cluster; i.e., the i th component of $\mathbf{1}_{\Gamma_k}$ is defined as: $(\mathbf{1}_{\Gamma_k})_i = 1$ if $i \in \Gamma_k$ and $(\mathbf{1}_{\Gamma_k})_i = 0$ otherwise. Define the associated *partition matrix* as:

$$X = \sum_{k=1}^K \frac{1}{|\Gamma_k|} \mathbf{1}_{\Gamma_k} \mathbf{1}_{\Gamma_k}^T. \quad (2)$$

Define $d_{ij} := \|x^i - x^j\|_2^2$ for all $i, j \in [n]$. Then Problem (1) can be equivalently written as (see [15] for detailed derivation):

$$\begin{aligned} \min \quad & \sum_{i,j \in [n]} d_{ij} X_{ij} \\ \text{s.t.} \quad & X \text{ is a partition matrix defined by (2).} \end{aligned} \quad (3)$$

It can be checked that any partition matrix X is positive semidefinite. Using this observation, one can obtain the following SDP relaxation of Problem (3):

$$\begin{aligned} \min \quad & \sum_{i,j \in [n]} d_{ij} X_{ij} \\ \text{s.t.} \quad & \text{Tr}(X) = K, \quad \sum_{j=1}^n X_{ij} = 1, \quad \forall i \in [n], \\ & X \succeq 0, \quad X_{ij} \geq 0, \quad X_{ij} = X_{ji}, \quad \forall 1 \leq i < j \leq n, \end{aligned} \quad (\text{PW})$$

where $\text{Tr}(X)$ is the trace of the matrix X , and $X \succeq 0$ means that X is positive semidefinite. The above relaxation was first proposed in [25] and is often referred to as the ‘‘Peng-Wei SDP relaxation’’. Both theoretical and numerical properties of this relaxation have been thoroughly investigated in the literature [5, 22, 12, 15, 27].

1.1 LP relaxations for K-means clustering

In [8], the authors introduced the *ratio-cut polytope*, defined as the convex hull of ratio-cut vectors corresponding to all partitions of n points in \mathbb{R}^m into at most K clusters. They showed this polytope is closely related to the convex hull of the feasible region of Problem (3). The authors then studied the facial structure of the ratio-cut polytope, which in turn enabled them to obtain a new family

of LP relaxations for K-means clustering. Fix a parameter $t \in \{2, \dots, K\}$; then an LP relaxation for K-means clustering is given by:

$$\begin{aligned}
\min \quad & \sum_{i,j \in [n]} d_{ij} X_{ij} && (\text{LPK}_t) \\
\text{s.t.} \quad & \text{Tr}(X) = K, \quad \sum_{j=1}^n X_{ij} = 1, \quad \forall i \in [n], \\
& \sum_{j \in S} X_{ij} \leq X_{ii} + \sum_{j,k \in S: j < k} X_{jk}, \quad \forall i \in [n], \forall S \subseteq [n] \setminus \{i\} : 2 \leq |S| \leq t, \\
& X_{ij} \geq 0, \quad X_{ij} = X_{ji}, \quad \forall 1 \leq i < j \leq n.
\end{aligned} \tag{4}$$

By Proposition 3 of [8], inequalities (4) define facets of the convex hull of the feasible region of Problem (3). It then follows that for any $t < t'$, the feasible region of Problem $(\text{LPK}_{t'})$ is strictly contained in the feasible region of Problem (LPK_t) . However, it is important to note that Problem (LPK_t) contains $\Theta(n^{t+1})$ inequalities; moreover, an inequality of the form (4) with $|S| = t'$ contains $\frac{t'(t'+1)}{2} + 1$ variables with nonzero coefficients. That is, while increasing t improves the quality of the relaxation, it significantly increases the computational cost of solving the LP as well. Hence, a careful selection of t is of crucial importance; we address this question in Section 4. In [5], the authors propose and study a different LP relaxation for K-means clustering. As it is detailed in [8], for any $t \geq 2$, the feasible region of Problem (LPK_t) is strictly contained in that of the LP relaxation considered in [5] (see Remark 1 in [8]). In fact, the numerical experiments on synthetic data in [8] indicate that Problem (LPK_t) with $t = 2$ is almost always *tight*; i.e., the optimal solution of the LP is a partition matrix. To better understand this unexpected behaviour, in this paper, we perform a theoretical study of the tightness of the (LPK_t) relaxation. As a first step, we consider the case of two clusters. Performing a similar type of analysis for $K > 2$ is a topic of future research.

Our main contributions are summarized as follows:

- (i) Consider any partition of $[n]$ and the associated partition matrix X defined by (2). By constructing a dual certificate, we obtain a sufficient condition, often referred to as a “proximity condition”, under which X is the unique optimal solution of Problem (LPK_t) (see Proposition 1 and Proposition 3). This result can be considered as a generalization of Theorem 1 in [8] where the optimality of *equal-size* clusters is studied. Our proximity condition is overly conservative since, to obtain an explicit condition, we fix a subset of the dual variables to zero. To address this, we propose a simple algorithm to carefully assign values to dual variables previously set to zero, leading to a significantly better dual certificate for X (see Proposition 5).
- (ii) Consider a generative model, referred to as the *stochastic sphere model* (SSM), in which there are two clusters of possibly different size in \mathbb{R}^m , and the data points in each cluster are sampled from a uniform distribution on a sphere of unit radius. Using the result of part (i) we obtain a sufficient condition in terms of the distance between sphere centers under which the (LPK_t) relaxation recovers the planted clusters with high probability. By high probability we mean the probability tending to one as the number of data points tends to infinity (see Proposition 4).
- (iii) Since in all our numerical experiments Problem (LPK_t) is tight, it is natural to ask whether the (LPK_t) relaxation is tight with high probability under reasonable generative models. We

present a family of inputs for which the (LPK_t) relaxation is never tight (see Proposition 6 and Corollary 1).

- (iv) Motivated by the surprising performance of the LP relaxation, we propose a scalable cutting plane algorithm, which relies on an efficient separation algorithm for inequalities (4), lower bounding and upper bounding techniques, and a GPU implementation of PDLP, a first-order primal-dual LP solver. Thanks to this algorithm, we are able to solve Problem (LPK_t) for real-world instances with up to 4000 data points in less than two and half hours. Interestingly for *all* instances, by solving the LP relaxation, we solve the original K-means clustering problem to global optimality.

1.2 Organization

The rest of the paper is structured as follows. In Section 2, we perform a theoretical study of the tightness of the LP relaxation for two clusters. In Section 3, we present a family of inputs for which the optimal value of Problem (LPK_t) is strictly smaller than the optimal value of the K-means clustering problem. Finally, in Section 4 we develop a customized algorithm for solving the LP relaxation and present extensive numerical experiments.

2 The tightness of the LP relaxation for two clusters

In this section, we consider the K-means clustering problem with $K = 2$ and examine the strength of the LP relaxation theoretically. Recall that for $K = 2$, we must have $t = 2$; therefore, Problem (LPK_t) simplifies to:

$$\min \sum_{i,j \in [n]} d_{ij} X_{ij} \tag{LP2}$$

$$\text{s.t. } \text{Tr}(X) = 2, \tag{5}$$

$$\sum_{j=1}^n X_{ij} = 1, \quad \forall 1 \leq i \leq n, \tag{6}$$

$$X_{ij} + X_{ik} \leq X_{ii} + X_{jk}, \quad \forall i \neq j \neq k \in [n], j < k, \tag{7}$$

$$X_{ij} \geq 0, \quad \forall 1 \leq i < j \leq n, \tag{8}$$

where as before we set $X_{ji} = X_{ij}$ for all $1 \leq i < j \leq n$. Throughout this section, whenever we say “the LP relaxation,” we mean the LP defined by Problem (LP2) and whenever we say “the SDP relaxation,” we mean the SDP defined by Problem (PW). To motivate our theoretical study, in the following we present a simple numerical experiment to convey the power of the LP relaxation for clustering real-world data sets. Solving both LP and SDP becomes computationally expensive as we increase the number of points n ; that is, to solve an instance with $n \gtrsim 200$ efficiently, one needs to design a specialized algorithm for both LP and SDP. Indeed, in [27] the authors design a specialized algorithm for the SDP relaxation and solve problems with $n \leq 4000$. In Section 4, we present a customized algorithm to solve the LP relaxation. In this section, however, we are interested in performing a theoretical analysis of the tightness of the LP and we are using our numerical experiments to motivate this study. Hence, we limit ourselves to solving problem instances with $n \leq 200$ variables.

Table 1: Comparing the strength of the LP versus the SDP for clustering real-world data with $K = 2$: t_{LP} denotes the tightness rate of the LP, t_{SDP} denotes the tightness rate of the SDP, g_{rel} denotes the (average) percentage of relative gap between the optimal values of SDP and LP.

Data set	n	m	t_{LP}	t_{SDP}	g_{rel} (%)
Voice	126	310	1.0	0.0	5.40
Iris	150	4	1.0	0.0	1.09
Wine	178	13	1.0	0.0	3.44
Seeds	210	7	1.0	0.0	3.80
Accent	329	12	1.0	0.3	0.02
ECG5000	500	140	1.0	0.1	0.07
Hungarian	522	20	1.0	0.0	1.01
Wdbc	569	30	1.0	0.0	5.51
Strawberry	613	235	1.0	0.0	5.40
Energy	768	16	1.0	1.0	0.0
SalesWeekly	810	106	1.0	0.0	1.74
Vehicle	846	18	1.0	0.0	1.11
Wafer	1000	152	1.0	0.3	0.06
Ethanol	2000	27	1.0	1.0	0.0
Rice	3810	7	1.0	0.0	1.70

We say that a convex relaxation is tight if its optimal solution is a partition matrix. In [26], the authors prove that a symmetric matrix X with unit row-sums and trace K is a partition matrix if and only if it is a *projection* matrix; i.e., $X^T X = X$. Hence to check whether the solution of the LP relaxation or the SDP relaxation is tight, we check whether it is a projection matrix. We collected a set of 15 data sets from the UCI machine learning repository [9]. The data sets and their characteristics, i.e., the number of points n and the input dimension m are listed in columns 1-3 of Table 1. For data sets with $n \leq 200$ (i.e., Voice, Iris, and Wine), we solve the clustering problem over the entire data set. For each of the remaining data sets, to control the computational cost of both LP and SDP, we sample 10 times $n' = 200$ data points and solve LP and SDP using the 10 random instances. All experiments are performed on the NEOS server [6]; LPs are solved with GAMS/Gurobi [11] and SDPs are solved with GAMS/MOSEK [1]. The tightness rate of the LP (t_{LP}), the tightness rate of the SDP (t_{SDP}), and the (average) relative gap between the optimal values of SDP and LP are listed in columns 4-6 of Table 1. In *all* instances, the LP relaxation is tight, while the SDP is only tight in a few instances. The remarkable performance of the LP relaxation is indeed unexpected and hence in this paper, it is our goal to better understand this behaviour through a theoretical study.

2.1 A sufficient condition for the tightness of the LP relaxation

We first obtain a sufficient “proximity” condition under which Problem (LP2) is tight; i.e., the optimal solution of the LP relaxation is a partition matrix. A proximity condition for the SDP relaxation defined by Problem (PW) is presented in [15]. As our proximity condition is overly conservative, we then present a simple algorithm that certifies the optimality of a given partition matrix. To establish our proximity condition, in Proposition 1, we first obtain a sufficient condition

under which a given partition matrix is an optimal solution of Problem (LP2). Subsequently in Proposition 3, we address the question of uniqueness of the optimal solution. The next proposition can be considered as a generalization of Theorem 1 in [8] where the authors study the optimality of *equal-size* clusters.

In the following, for every $A \subseteq [n]$ and $f : A \rightarrow \mathbb{R}$, we define

$$\sum_{i \in A} f(i) := \frac{1}{|A|} \sum_{i \in A} f(i).$$

Moreover, given a partition $\{\Gamma_1, \Gamma_2\}$ of $[n]$, for each $i \in \Gamma_l$, $l \in \{1, 2\}$, we define

$$d_i^{\text{in}} := \sum_{j \in \Gamma_l} d_{ij}, \quad d_i^{\text{out}} := \sum_{j \in [n] \setminus \Gamma_l} d_{ij}.$$

We are now ready to state our sufficient condition.

Proposition 1. *Let $\{\Gamma_1, \Gamma_2\}$ denote a partition of $[n]$ and assume without loss of generality that $|\Gamma_1| \leq |\Gamma_2|$. Define*

$$r_1 := \frac{2|\Gamma_1|}{|\Gamma_1| + |\Gamma_2|}, \quad r_2 := \frac{2|\Gamma_2|}{|\Gamma_1| + |\Gamma_2|}, \quad (9)$$

and

$$\eta := \frac{r_2}{2} \left(\left(1 - \frac{r_1}{r_2}\right) \max_{k \in \Gamma_1} d_k^{\text{in}} + \left(1 - \frac{r_2}{r_1}\right) \min_{k \in \Gamma_2} d_k^{\text{in}} + \frac{r_1}{r_2} \sum_{k \in \Gamma_1} d_k^{\text{in}} + \frac{r_2}{r_1} \sum_{k \in \Gamma_2} d_k^{\text{in}} \right). \quad (10)$$

Suppose that

$$\sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} - d_{ij} \geq \eta, \quad \forall i < j \in \Gamma_1, \quad (11)$$

and

$$\sum_{k \in \Gamma_1} \min\{r_1 d_{ik} + d_j^{\text{in}}, r_1 d_{jk} + d_i^{\text{in}}\} - d_{ij} \geq \frac{r_1}{r_2} \eta, \quad \forall i < j \in \Gamma_2. \quad (12)$$

Then, an optimal solution of Problem (LP2) is given by the partition matrix

$$\bar{X} = \frac{1}{|\Gamma_1|} \mathbf{1}_{\Gamma_1} \mathbf{1}_{\Gamma_1}^T + \frac{1}{|\Gamma_2|} \mathbf{1}_{\Gamma_2} \mathbf{1}_{\Gamma_2}^T. \quad (13)$$

Proof. We start by constructing the dual of Problem (LP2). Define dual variables ω associated with (5), μ_i for all $i \in [n]$ associated with (6), λ_{ijk} for all $(i, j, k) \in \Omega := \{(i, j, k) : i \neq j \neq k \in [n], j < k\}$ associated with (7), and σ_{ij} for all $1 \leq i < j \leq n$ associated with (8). The dual of Problem (LP2) is then given by:

$$\begin{aligned} \max \quad & -(2\omega + \sum_{i \in [n]} \mu_i) \\ \text{s.t.} \quad & \mu_i + \mu_j + \sum_{k \in [n] \setminus \{i, j\}} (\lambda_{ijk} + \lambda_{jik} - \lambda_{kij}) + 2d_{ij} - \sigma_{ij} = 0, \quad \forall 1 \leq i < j \leq n, \end{aligned} \quad (14)$$

$$\omega + \mu_i - \sum_{j, k \in [n] \setminus \{i\} : j < k} \lambda_{ijk} = 0, \quad \forall i \in [n], \quad (15)$$

$$\lambda_{ijk} \geq 0, \quad \forall (i, j, k) \in \Omega, \quad \sigma_{ij} \geq 0, \quad \forall 1 \leq i < j \leq n,$$

where we let $\lambda_{ikj} = \lambda_{ijk}$ for all $(i, j, k) \in \Omega$. To establish the optimality of \bar{X} defined by (13), it suffices to construct a dual feasible point $(\bar{\lambda}, \bar{\mu}, \bar{\omega}, \bar{\sigma})$ that together with \bar{X} satisfies the complementary slackness. Without loss of generality, assume that $i < j$ for all $i \in \Gamma_1$ and $j \in \Gamma_2$. Then \bar{X} and $(\bar{\lambda}, \bar{\mu}, \bar{\omega}, \bar{\sigma})$ satisfy the complementary slackness if and only if:

- (i) $\bar{\lambda}_{ijk} = 0$ if $i \in \Gamma_1$ and $j, k \in \Gamma_2$ or if $i \in \Gamma_2$ and $j, k \in \Gamma_1$,
- (ii) $\bar{\sigma}_{ij} = 0$ if $i, j \in \Gamma_1$ or $i, j \in \Gamma_2$.

After projecting out $\bar{\sigma}_{ij}$ for $i \in \Gamma_1$ and $j \in \Gamma_2$, we deduce that it suffices to find $(\bar{\lambda}, \bar{\mu}, \bar{\omega})$ satisfying:

$$\bar{\mu}_i + \bar{\mu}_j + \sum_{k \notin \Gamma_l} (\bar{\lambda}_{ijk} + \bar{\lambda}_{jik}) + \sum_{k \in \Gamma_l \setminus \{i, j\}} (\bar{\lambda}_{ijk} + \bar{\lambda}_{jik} - \bar{\lambda}_{kij}) + 2d_{ij} = 0, \quad \forall i < j \in \Gamma_l, l \in \{1, 2\} \quad (16)$$

$$\bar{\mu}_i + \bar{\mu}_j + \sum_{k \in \Gamma_1 \setminus \{i\}} (\bar{\lambda}_{ijk} - \bar{\lambda}_{kij}) + \sum_{k \in \Gamma_2 \setminus \{j\}} (\bar{\lambda}_{jik} - \bar{\lambda}_{kij}) + 2d_{ij} \geq 0, \quad \forall i \in \Gamma_1, j \in \Gamma_2 \quad (17)$$

$$\bar{\omega} + \bar{\mu}_i - \sum_{\substack{j \in \Gamma_l \setminus \{i\}, \\ k \notin \Gamma_l}} \bar{\lambda}_{ijk} - \sum_{j < k \in \Gamma_l \setminus \{i\}} \bar{\lambda}_{ijk} = 0, \quad \forall i \in \Gamma_l, l \in \{1, 2\}. \quad (18)$$

To this end, let

$$\bar{\lambda}_{ijk} - \bar{\lambda}_{jik} = \frac{d_{jk} - d_{ik}}{n/2} + \frac{d_i^{\text{in}} - d_j^{\text{in}}}{|\Gamma_2|}, \quad \forall i, j \in \Gamma_1, k \in \Gamma_2 \quad (19)$$

$$\bar{\lambda}_{ijk} - \bar{\lambda}_{jik} = \frac{d_{jk} - d_{ik}}{n/2} + \frac{d_i^{\text{in}} - d_j^{\text{in}}}{|\Gamma_1|}, \quad \forall i, j \in \Gamma_2, k \in \Gamma_1 \quad (20)$$

$$\bar{\mu}_i = -d_i^{\text{in}} - r_2 d_i^{\text{out}} + \eta, \quad \forall i \in \Gamma_1 \quad (21)$$

$$\bar{\mu}_i = -d_i^{\text{in}} - r_1 d_i^{\text{out}} + \frac{r_1}{r_2} \eta, \quad \forall i \in \Gamma_2 \quad (22)$$

$$\bar{\omega} = -\frac{1}{2} \sum_{i \in [n]} (d_i^{\text{in}} + \bar{\mu}_i), \quad (23)$$

where r_1, r_2 and η are as defined by (9) and (10), respectively.

First let us examine the validity of inequalities (17). Substituting (19)-(22) in (17) yields:

$$\begin{aligned} & -d_i^{\text{in}} - r_2 d_i^{\text{out}} + \eta - d_j^{\text{in}} - r_1 d_j^{\text{out}} + \frac{r_1}{r_2} \eta + \sum_{k \in \Gamma_1 \setminus \{i\}} \left(\frac{d_{jk} - d_{ij}}{n/2} + \frac{d_i^{\text{in}} - d_k^{\text{in}}}{|\Gamma_2|} \right) \\ & + \sum_{k \in \Gamma_2 \setminus \{j\}} \left(\frac{d_{ik} - d_{ij}}{n/2} + \frac{d_j^{\text{in}} - d_k^{\text{in}}}{|\Gamma_1|} \right) + 2d_{ij} = \\ & -d_i^{\text{in}} - r_2 d_i^{\text{out}} + \eta - d_j^{\text{in}} - r_1 d_j^{\text{out}} + \frac{r_1}{r_2} \eta + r_1 d_j^{\text{out}} - r_1 d_{ij} + \frac{r_1}{r_2} d_i^{\text{in}} - \frac{r_1}{r_2} \sum_{k \in \Gamma_1} d_k^{\text{in}} + r_2 d_i^{\text{out}} - r_2 d_{ij} + \\ & \frac{r_2}{r_1} d_j^{\text{in}} - \frac{r_2}{r_1} \sum_{k \in \Gamma_2} d_k^{\text{in}} + 2d_{ij} = \end{aligned}$$

$$\left(1 + \frac{r_1}{r_2}\right)\eta - \left(1 - \frac{r_1}{r_2}\right)d_i^{\text{in}} - \left(1 - \frac{r_2}{r_1}\right)d_j^{\text{in}} - \frac{r_1}{r_2}\sum_{k \in \Gamma_1} d_k^{\text{in}} - \frac{r_2}{r_1}\sum_{k \in \Gamma_2} d_k^{\text{in}} \geq 0,$$

where the last inequality follows from the definition of η given by (10) and the identity $r_1 + r_2 = 2$.

We now show that equalities (18) are implied by equalities (16). In the following we prove that all equalities of the form (18) with $l = 1$ are implied by equalities of the form (16) with $l = 1$; the proof of the case with $l = 2$ follows from a similar line of arguments. Using (19) to eliminate $\bar{\lambda}_{jik}$ for $i, j \in \Gamma_1$ and $k \in \Gamma_2$, and using (21) to eliminate $\bar{\mu}_i$, $i \in \Gamma_1$, it follows that equalities (16) with $l = 1$ can be equivalently written as:

$$\sum_{k \in \Gamma_2} \bar{\lambda}_{ijk} + \frac{1}{2} \sum_{k \in \Gamma_1 \setminus \{i, j\}} (\bar{\lambda}_{ijk} + \bar{\lambda}_{jik} - \bar{\lambda}_{kij}) = d_i^{\text{in}} + r_2 d_j^{\text{out}} - d_{ij} - \eta, \quad \forall i < j \in \Gamma_1. \quad (24)$$

Using (21)-(23), inequalities (18) can be written as:

$$\sum_{\substack{j \in \Gamma_1 \setminus \{i\}, \\ k \in \Gamma_2}} \bar{\lambda}_{ijk} + \sum_{j < k \in \Gamma_1 \setminus \{i\}} \bar{\lambda}_{ijk} = r_2 \sum_{j \in \Gamma_1} d_j^{\text{out}} - d_i^{\text{in}} - r_2 d_i^{\text{out}} - (|\Gamma_1| - 1)\eta, \quad \forall i \in \Gamma_1, \quad (25)$$

where we used the identity $r_2 \sum_{j \in \Gamma_1} d_i^{\text{out}} = r_1 \sum_{j \in \Gamma_2} d_i^{\text{out}}$. Moreover, it can be checked that

$$\sum_{j \neq k \in \Gamma_1 \setminus \{i\}} (\bar{\lambda}_{ijk} + \bar{\lambda}_{jik} - \bar{\lambda}_{kij}) = 2 \sum_{j < k \in \Gamma_1 \setminus \{i\}} \bar{\lambda}_{ijk}.$$

Therefore, to show that equalities (25) are implied by equalities (24), it suffices to have:

$$\sum_{j \in \Gamma_1 \setminus \{i\}} (d_i^{\text{in}} + r_2 d_j^{\text{out}} - d_{ij} - \eta) = r_2 \sum_{j \in \Gamma_1} d_j^{\text{out}} - d_i^{\text{in}} - r_2 d_i^{\text{out}} - (|\Gamma_1| - 1)\eta,$$

whose validity follows since $\sum_{j \in \Gamma_1 \setminus \{i\}} (d_i^{\text{in}} - d_{ij}) = (|\Gamma_1| - 1)d_i^{\text{in}} - |\Gamma_1|d_i^{\text{in}} = d_i^{\text{in}}$.

Therefore, it remains to prove the validity of equalities (16). First consider the case with $l = 1$. By (19) and nonnegativity of $\bar{\lambda}_{ijk}$ for all $(i, j, k) \in \Omega$, we deduce that

$$\bar{\lambda}_{ijk} + \bar{\lambda}_{jik} \geq \text{abs} \left(\frac{d_{jk} - d_{ik}}{n/2} + \frac{d_i^{\text{in}} - d_j^{\text{in}}}{|\Gamma_2|} \right) = \frac{1}{|\Gamma_2|} \text{abs} \left((d_i^{\text{in}} + r_2 d_{jk}) - (d_j^{\text{in}} + r_2 d_{ik}) \right), \quad \forall i, j \in \Gamma_1, k \in \Gamma_2,$$

where $\text{abs}(\cdot)$ denote the absolute value function. Hence, using (21) to eliminate $\bar{\mu}_i$, we conclude that equalities (16) with $l = 1$ can be satisfied if

$$\frac{1}{2} \sum_{k \in \Gamma_1 \setminus \{i, j\}} (\bar{\lambda}_{ijk} + \bar{\lambda}_{jik} - \bar{\lambda}_{kij}) \leq \sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} - d_{ij} - \eta, \quad \forall i < j \in \Gamma_1, \quad (26)$$

where we used the identity $\text{abs}(a - b) = a + b - 2 \min\{a, b\}$. Letting $\bar{\lambda}_{ijk} = 0$ for all $(i, j, k) \in \Gamma_1$ and using (11), we conclude that inequalities (26) are valid. Similarly, it can be checked that inequalities (17) with $l = 2$ can be satisfied if

$$\frac{1}{2} \sum_{k \in \Gamma_2 \setminus \{i, j\}} (\bar{\lambda}_{ijk} + \bar{\lambda}_{jik} - \bar{\lambda}_{kij}) \leq \sum_{k \in \Gamma_1} \min\{r_1 d_{ik} + d_j^{\text{in}}, r_1 d_{jk} + d_i^{\text{in}}\} - d_{ij} - \frac{r_1}{r_2} \eta, \quad \forall i < j \in \Gamma_2. \quad (27)$$

Letting $\bar{\lambda}_{ijk} = 0$ for all $(i, j, k) \in \Gamma_2$ and using (12), we conclude that inequalities (27) are valid and this completes the proof. \square

We now consider the question of uniqueness of the optimal solution. To this end, we make use of the following result:

Proposition 2 (Part (iv) of Theorem 2 in [20]). *Consider an LP whose feasible region is defined by $Ax = b$ and $Cx \leq d$, where $x \in \mathbb{R}^n$ denotes the vector of optimization variables, and b, d, A, C are vectors and matrices of appropriate dimensions. Let \bar{x} be an optimal solution of this LP and denote by \bar{u} the dual optimal solution corresponding to the inequality constraints. Let C_i denote the i -th row of C . Define $Q = \{i : C_i \bar{x} = d_i, \bar{u}_i > 0\}$, $L = \{i : C_i \bar{x} = d_i, \bar{u}_i = 0\}$. Let \mathcal{C}_Q and \mathcal{C}_L be the matrices whose rows are C_i , $i \in Q$ and C_i , $i \in L$, respectively. Then \bar{x} is the unique optimal solution of the LP, if there exists no x different from the zero vector satisfying*

$$Ax = 0, \quad \mathcal{C}_Q x = 0, \quad \mathcal{C}_L x \leq 0. \quad (28)$$

We are now ready to establish our uniqueness result:

Proposition 3. *Suppose that inequalities (11) and (12) are strictly satisfied. Then \bar{X} defined by (13) is the unique optimal solution of Problem (LP2).*

Proof. Consider the dual certificate $(\bar{\lambda}, \bar{\mu}, \bar{\omega}, \bar{\sigma})$ constructed in the proof of Proposition 1. We consider a slightly modified version of this certificate by redefining η defined in (10) as follows:

$$\eta = \frac{r_2}{2} \left(\left(1 - \frac{r_1}{r_2}\right) \max_{k \in \Gamma_1} d_k^{\text{in}} + \left(1 - \frac{r_2}{r_1}\right) \min_{k \in \Gamma_2} d_k^{\text{in}} + \frac{r_1}{r_2} \sum_{k \in \Gamma_1} d_k^{\text{in}} + \frac{r_2}{r_1} \sum_{k \in \Gamma_2} d_k^{\text{in}} \right) + \epsilon,$$

for some $\epsilon > 0$. This in turn will imply that $\bar{\sigma}_{ij} > 0$ for all $i \in \Gamma_1, j \in \Gamma_2$. Notice that this is an admissible modification as we are assuming that inequalities (11) and (12) are strictly satisfied. In addition, we can choose

$$\begin{aligned} \bar{\lambda}_{ijk} &> \max \left\{ 0, \frac{d_{jk} - d_{ik}}{n/2} + \frac{d_i^{\text{in}} - d_j^{\text{in}}}{|\Gamma_2|} \right\}, \quad \bar{\lambda}_{jik} > \max \left\{ 0, \frac{d_{ik} - d_{jk}}{n/2} + \frac{d_j^{\text{in}} - d_i^{\text{in}}}{|\Gamma_2|} \right\}, \quad \forall i, j \in \Gamma_1, k \in \Gamma_2 \\ \bar{\lambda}_{ijk} &> \max \left\{ 0, \frac{d_{jk} - d_{ik}}{n/2} + \frac{d_i^{\text{in}} - d_j^{\text{in}}}{|\Gamma_1|} \right\}, \quad \bar{\lambda}_{jik} > \max \left\{ 0, \frac{d_{ik} - d_{jk}}{n/2} + \frac{d_j^{\text{in}} - d_i^{\text{in}}}{|\Gamma_1|} \right\}, \quad \forall i, j \in \Gamma_2, k \in \Gamma_1. \end{aligned}$$

Therefore, by Proposition 2, the matrix \bar{X} defined by (13) is the unique optimal solution of Problem (LP2), if there exists no $X \neq 0$ satisfying:

$$\sum_{j=1}^n X_{ij} = 0, \quad \forall 1 \leq i \leq n \quad (29)$$

$$X_{ij} = 0, \quad \forall i \in \Gamma_1, j \in \Gamma_2 \quad (30)$$

$$X_{ij} + X_{ik} = X_{ii} + X_{jk}, \quad \forall i, j \in \Gamma_1, k \in \Gamma_2, \text{ or } i, j \in \Gamma_2, k \in \Gamma_1. \quad (31)$$

From (29) and (30) it follows that

$$X_{ii} + \sum_{j \in \Gamma_1} X_{ij} = 0, \quad \forall i \in \Gamma_1, \quad X_{ii} + \sum_{j \in \Gamma_2} X_{ij} = 0, \quad \forall i \in \Gamma_2. \quad (32)$$

Moreover, from equations (31) we deduce that

$$X_{ij} = \frac{X_{ii} + X_{jj}}{2}, \quad \forall i, j \in \Gamma_1 \text{ or } i, j \in \Gamma_2. \quad (33)$$

Substituting (33) in (32) we obtain: $X_{ii} = X_{ij} = 0$ for all $i, j \in \Gamma_1$ and for all $i, j \in \Gamma_2$, which together with (30) completes the proof. \square

To have an intuitive understanding of our proximity condition, consider the special case with $d_i^{\text{in}} = \alpha$ for all $i \in [n]$ and $|\Gamma_1| = |\Gamma_2|$. This assumption is for instance satisfied, if $|\Gamma_1| = |\Gamma_2|$ and Γ_1 and Γ_2 are uniformly distributed on two spheres S_1 and S_2 of equal radii. Denote by c_1 the center of S_1 . By Γ_1 being uniformly distributed on S_1 , we imply that for any rotation matrix $R \in \mathbb{R}^{d \times d}$ for which there exist two points $x^i, x^j \in \Gamma_1$ such that $R(x^i - c_1) = (x^j - c_1)$, we have $R(\Gamma_1 - c_1) = (\Gamma_1 - c_1)$. In this setting we have $\eta = 1$ and assumptions (11) and (12) simplify to:

$$\sum_{k \in \Gamma_2} \min\{d_{ik}, d_{jk}\} \geq d_{ij}, \quad \forall i < j \in \Gamma_1, \quad \sum_{k \in \Gamma_1} \min\{d_{ik}, d_{jk}\} \geq d_{ij}, \quad \forall i < j \in \Gamma_2.$$

These inequalities requires that the distance between any two points x_i, x_j in the same cluster should be smaller than the average distance between the sets $\{x_i, x_j\}$ and $\{x_k\}$, where x_k denotes a point in the other cluster.

Using our proximity condition given by inequalities (11) and (12), we next obtain a recovery guarantee for the LP relaxation under a popular stochastic model for the input data.

2.2 Recovery guarantee for the stochastic sphere model

It is widely understood that worst-case guarantees for optimization algorithms are often too pessimistic. A recent line of research in data clustering is concerned with obtaining sufficient conditions under which a planted clustering corresponds to the unique optimal solution of a convex relaxation under suitable stochastic models [5, 22, 12, 15, 8]. Such conditions are often referred to as (exact) *recovery* conditions and are used to compare the strength of various convex relaxations for NP-hard problems. Henceforth, we say that an optimization problem *recovers* the planted clusters if its unique optimal solution corresponds to the planted clusters.

Perhaps the most popular generative model for K-means clustering is the *stochastic ball model*, where the points are sampled from uniform distributions on K unit balls in \mathbb{R}^m . As before, we let $K = 2$ and we denote by Δ the distance between the ball centers. Notice that the question of recovery only makes sense when $\Delta > 2$. We denote by Γ_1 and Γ_2 the set of points sampled from the first and second balls, respectively and without loss of generality we assume $|\Gamma_1| \leq |\Gamma_2|$. In the following, whenever we say *with high probability*, we mean the probability tending to one as $n \rightarrow \infty$. In [15], the authors prove that the Peng-Wei SDP relaxation defined by Problem (PW) recovers the planted clusters with high probability if

$$\Delta > 2 \left(1 + \sqrt{\frac{1}{r_1(m+2)}} \right), \quad (34)$$

where r_1 is defined by (9). In the special case of equal-size clusters, i.e., $|\Gamma_1| = |\Gamma_2|$, the authors of [5] show that the Peng-Wei SDP relaxation recovers the planted clusters with high probability, if $\Delta > 2\sqrt{2}(1 + \frac{1}{\sqrt{m}})$, while the authors of [12] show that the same SDP recovers the planted clusters

with high probability if $\Delta > 2(1 + \frac{2}{m})$. Again, for equal-size clusters, in [8], the authors show that Problem (LP2) recovers the planted clusters with high probability, if $\Delta > 1 + \sqrt{3}$.

In this section, we obtain a recovery guarantee for the LP relaxation for two clusters of arbitrary size. For simplicity, we consider a slightly different stochastic model, which we refer to as the stochastic sphere model (SSM), where instead of a ball, points in each cluster are sampled from a sphere (i.e., the boundary of a ball). We prove that our deterministic condition given by inequalities (11)-(12) implies that Problem (LP2) recovers the planted clusters with high probability, if

$$\Delta > 1 + \sqrt{1 + \frac{2}{r_1}}. \quad (35)$$

We should mention that, at the expense of a significantly longer proof, one can obtain the same recovery guarantee (35) for the LP relaxation under the SBM. We do not include the latter result in this paper because, while the proof is more technical and longer, it does not contain any new ideas and closely follows the path of our proof for the SSM. Indeed, in case of equal-size clusters; i.e., $r_1 = 1$, inequality (35) simplifies to the recovery guarantee of [8] for SBM: $\Delta > 1 + \sqrt{3}$. Also note that while the recovery guarantee for the LP (35) is better than the the recovery guarantee for the SDP (34) for $1 \leq m \leq 5$, it becomes weaker for larger dimensions. As we detail in the next section, condition (35) can be significantly improved via a more careful selection of the dual certificate.

Throughout this section, for an event A , we denote by $\mathbb{P}(A)$ the probability of A . For a random variable Y , we denote by $\mathbb{E}[Y]$ its expected value. In case of a multivariate random variable X_{ij} , the conditional expected value in j , with i fixed, will be denoted either with $\mathbb{E}_i[X]$ or with $\mathbb{E}^j[X]$. We denote by $\partial\mathcal{B}_1$ and $\partial\mathcal{B}_2$ the spheres corresponding to the first and second clusters, respectively. Up to a rotation we can assume that the centers of $\partial\mathcal{B}_1$ and $\partial\mathcal{B}_2$ are 0 and Δe_1 , respectively, where e_1 is the first vector of the standard basis of \mathbb{R}^m . For a continuous function $f : \partial\mathcal{B}_1 \rightarrow \mathbb{R}$ (and analogously for $\partial\mathcal{B}_2$), we define

$$\int_{\partial\mathcal{B}_1} f(x) d\mathcal{H}^{m-1}(x) := \frac{1}{\mathcal{H}^{m-1}(\partial\mathcal{B}_1)} \int_{\partial\mathcal{B}_1} f(x) d\mathcal{H}^{m-1}(x),$$

where $\mathcal{H}^{m-1}(x)$ denotes the $(m-1)$ -dimensional Hausdorff measure.

We are now ready to state our recovery result.

Proposition 4. *Suppose that the points are generated according to the SSM. Then Problem (LP2) recovers the planted clusters with high probability if $\Delta > \Delta_0 := 1 + \sqrt{1 + \frac{2}{r_1}}$, where r_1 is defined by (9).*

Proof. By Proposition 3, it suffices to show that for $\Delta > \Delta_0$, inequalities (11)-(12) are strictly satisfied with high probability. Namely, we show that there exists a universal constant $C > 0$ such that, for $\Delta > \Delta_0$ we have

$$\mathbb{P}\left(\bigcap_{i,j \in \Gamma_1} \left\{d_{ij} + \eta - \sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} < 0\right\}\right) \geq 1 - Cn^2 e^{-\frac{n\epsilon_2^2}{C}} \quad (36)$$

and

$$\mathbb{P}\left(\bigcap_{i,j \in \Gamma_2} \left\{d_{ij} + \frac{r_1}{r_2} \eta - \sum_{k \in \Gamma_1} \min\{r_1 d_{ik} + d_j^{\text{in}}, r_1 d_{jk} + d_i^{\text{in}}\} < 0\right\}\right) \geq 1 - Cn^2 e^{-\frac{n\epsilon_1^2}{C}},$$

where $\epsilon_1, \epsilon_2 > 0$ are as defined in the statement of Lemma 1. Since the two inequalities are symmetric, their proof is similar and we will only prove inequality (36). To this aim, for notational simplicity, define

$$t_{ij} := \mathbb{E}^k \left[\sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + \mathbb{E}_j[d_j^{\text{in}}], r_2 d_{jk} + \mathbb{E}_i[d_i^{\text{in}}]\} \right].$$

Then we can compute

$$\begin{aligned} & \mathbb{P} \left(\bigcap_{i,j \in \Gamma_1} \left\{ d_{ij} + \eta - \sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} < 0 \right\} \right) \\ & \geq \mathbb{P} \left(\bigcap_{i,j \in \Gamma_1} \left\{ d_{ij} + \eta - d_{ij} - 2r_2 + t_{ij} - \sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} < 4\epsilon_2 \right\} \right) \\ & \geq \mathbb{P} \left(\left\{ |\eta - 2r_2| < 2\epsilon_2 \right\} \cap \bigcap_{i,j \in \Gamma_1} \left\{ \left| t_{ij} - \mathbb{E}^k \left[\sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} \right] \right| < \epsilon_2 \right\} \right) \\ & \cap \bigcap_{i,j \in \Gamma_1} \left\{ \left| \mathbb{E}^k \left[\sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} \right] - \sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} \right| < \epsilon_2 \right\} \\ & \geq 1 - \mathbb{P}(\{|\eta - 2r_2| \geq 2\epsilon_2\}) - \mathbb{P} \left(\bigcup_{i,j \in \Gamma_1} \left\{ \left| t_{ij} - \mathbb{E}^k \left[\sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} \right] \right| \geq \epsilon_2 \right\} \right) \\ & - \mathbb{P} \left(\bigcup_{i,j \in \Gamma_1} \left\{ \left| \mathbb{E}^k \left[\sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} \right] - \sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} \right| \geq \epsilon_2 \right\} \right) \end{aligned} \quad (37)$$

The first inequality follows from (47) in Lemma 1, since $\Delta > \Delta_0$; the second inequality holds by set inclusion, and the third inequality is obtained by taking the union bound. To complete the proof, we next estimate each of the terms in the last two lines of (37). In the following, C will always denote a universal positive constant, which may increase from one line to the next line and we will not relabel it for the sake of exposition. First, to estimate $\mathbb{P}(\{|\eta - 2r_2| \geq 2\epsilon_2\})$, we define

$$\eta_1 := \frac{r_2}{2} \left(\left(1 - \frac{r_1}{r_2}\right) \max_{k \in \Gamma_1} d_k^{\text{in}} + \left(1 - \frac{r_2}{r_1}\right) \min_{k \in \Gamma_2} d_k^{\text{in}} \right), \quad \eta_2 := \frac{r_2}{2} \left(\frac{r_1}{r_2} \sum_{k \in \Gamma_1} d_k^{\text{in}} + \frac{r_2}{r_1} \sum_{k \in \Gamma_2} d_k^{\text{in}} \right), \quad (38)$$

so that $\eta = \eta_1 + \eta_2$. Recall that $|\Gamma_1| = r_1 \frac{n}{2}$, $|\Gamma_2| = r_2 \frac{n}{2}$, $r_1 \in (0, 1]$ and $r_2 \in [1, 2)$. Since for $\Delta > 4$, the recovery follows from a simple thresholding argument, we can restrict our attention to $\Delta \leq 4$, i.e., we assume that $r_1 \geq \frac{1}{4}$. It then follows that

$$\begin{aligned} & \mathbb{P}(\{|\eta_2 - \mathbb{E}[\eta_2]| \geq \epsilon_2\}) \\ & = \mathbb{P} \left(\left| r_1 \left(\sum_{i,j \in \Gamma_1} d_{ij} - \mathbb{E} \left[\sum_{i,j \in \Gamma_1} d_{ij} \right] \right) + \frac{r_2}{r_1} \left(\mathbb{E} \left[\sum_{i,j \in \Gamma_2} d_{ij} \right] - \sum_{i,j \in \Gamma_2} d_{ij} \right) \right| \geq 2\epsilon_2 \right) \\ & \leq \mathbb{P} \left(\left\{ \left| \sum_{i,j \in \Gamma_1} d_{ij} - \mathbb{E} \left[\sum_{i,j \in \Gamma_1} d_{ij} \right] \right| \geq \frac{\epsilon_2}{C} \right\} \cup \left\{ \left| \mathbb{E} \left[\sum_{i,j \in \Gamma_2} d_{ij} \right] - \sum_{i,j \in \Gamma_2} d_{ij} \right| \geq \frac{\epsilon_2}{C} \right\} \right) \\ & \leq \mathbb{P} \left(\left| \sum_{i,j \in \Gamma_1} d_{ij} - \mathbb{E} \left[\sum_{i,j \in \Gamma_1} d_{ij} \right] \right| \geq \frac{\epsilon_2}{C} \right) + \mathbb{P} \left(\left| \mathbb{E} \left[\sum_{i,j \in \Gamma_2} d_{ij} \right] - \sum_{i,j \in \Gamma_2} d_{ij} \right| \geq \frac{\epsilon_2}{C} \right) \leq C e^{-\frac{n^2 \epsilon_2^2}{\sigma}}. \end{aligned} \quad (39)$$

The first inequality holds by set inclusion and the third inequality follows from Hoeffding's inequality (see for example Theorem 2.2.6 in [29]), since d_{ij} , $i, j \in \Gamma_l$ are i.i.d. random variables for every $l \in \{1, 2\}$ and $d_{ij} \in [0, 4]$. Next we show that

$$\mathbb{P}\left(\left|\eta_1 - \left(2r_2 - r_1 - \frac{r_2^2}{r_1}\right)\right| \geq \epsilon_2\right) \leq Cne^{-\frac{n\epsilon_2^2}{\sigma}}. \quad (40)$$

For notational simplicity, we denote the i th point in Γ_1 by x . We notice that for any $i \in \Gamma_1$ we have

$$\mathbb{E}_i[d_i^{\text{in}}] = \int_{\partial\mathcal{B}_1} \|x - z\|^2 d\mathcal{H}^{m-1}(z) = \|x\|^2 + \int_{\partial\mathcal{B}_1} \|z\|^2 d\mathcal{H}^{m-1}(z) - 2x^T \int_{\partial\mathcal{B}_1} z d\mathcal{H}^{m-1}(z) = 2. \quad (41)$$

By symmetry, the same calculation holds for $\mathbb{E}_i[d_i^{\text{in}}]$ with $i \in \Gamma_2$. By (41), we have

$$\mathbb{E}\left[\sum_{k \in \Gamma_1} d_k^{\text{in}}\right] = \mathbb{E}\left[\sum_{k \in \Gamma_2} d_k^{\text{in}}\right] = 2. \quad (42)$$

Using Hoeffding's inequality together with (41), we get

$$\mathbb{P}(\{d_k^{\text{in}} > 2 + \epsilon_2\}) \leq Ce^{-\frac{n\epsilon_2^2}{\sigma}}, \quad \mathbb{P}(\{d_k^{\text{in}} < 2 - \epsilon_2\}) \leq Ce^{-\frac{n\epsilon_2^2}{\sigma}}, \quad \forall k \in \Gamma_1 \cup \Gamma_2.$$

Hence, by the union bound, we obtain

$$\mathbb{P}(\{|\max_{k \in \Gamma_1} d_k^{\text{in}} - 2| \geq \epsilon_2\}) \leq Cne^{-\frac{n\epsilon_2^2}{\sigma}}, \quad \mathbb{P}(\{|\min_{k \in \Gamma_2} d_k^{\text{in}} - 2| \geq \epsilon_2\}) \leq Cne^{-\frac{n\epsilon_2^2}{\sigma}},$$

from which we conclude the validity of (40). Since by (38) and (42) we have

$$\mathbb{E}[\eta_2] = r_2\left(\frac{r_1}{r_2} + \frac{r_2}{r_1}\right) = \left(r_1 + \frac{r_2^2}{r_1}\right),$$

we can combine (39) with (40) to conclude that

$$\mathbb{P}(\{|\eta - 2r_2| \geq 2\epsilon_2\}) \leq Cne^{-\frac{n\epsilon_2^2}{\sigma}}. \quad (43)$$

We now observe that

$$\begin{aligned} & \mathbb{P}\left(\bigcup_{i,j \in \Gamma_1} \left\{\left|t_{ij} - \mathbb{E}^k \left[\sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\}\right]\right| \geq \epsilon_2\right\}\right) \\ & \leq \mathbb{P}\left(\bigcup_{i \in \Gamma_1} \left\{\left|d_i^{\text{in}} - \mathbb{E}_i[d_i^{\text{in}}]\right| \geq \epsilon_2/2\right\}\right) \leq Cne^{-\frac{n\epsilon_2^2}{\sigma}}, \end{aligned} \quad (44)$$

where the first inequality follows from the linearity of expectation and the second inequality follows from the application of Hoeffding's inequality and taking the union bound. Finally, by Hoeffding's inequality we have

$$\begin{aligned} & \mathbb{P}\left(\bigcup_{i,j \in \Gamma_1} \left\{\left|\mathbb{E}^k \left[\sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\}\right] - \sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\}\right| \geq \epsilon_2\right\}\right) \\ & \leq Cn^2 e^{-\frac{n\epsilon_2^2}{\sigma}}. \end{aligned} \quad (45)$$

Plugging inequalities (43)-(44)-(45) in (37), we conclude the claimed inequality (36). \square

In order to prove our recovery result in Proposition 4, we made use of the following lemma.

Lemma 1. *Suppose that the random points are generated according to the SSM. Then the following inequalities hold provided that $\Delta > \Delta_0$:*

$$4\epsilon_1 := \inf_{i,j \in \Gamma_2} \mathbb{E}^k \left[\sum_{k \in \Gamma_1} \min\{r_1 d_{ik} + \mathbb{E}_j[d_j^{in}], r_1 d_{jk} + \mathbb{E}_i[d_i^{in}]\} \right] - d_{ij} - 2r_1 > 0, \quad (46)$$

$$4\epsilon_2 := \inf_{i,j \in \Gamma_1} \mathbb{E}^k \left[\sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + \mathbb{E}_j[d_j^{in}], r_2 d_{jk} + \mathbb{E}_i[d_i^{in}]\} \right] - d_{ij} - 2r_2 > 0. \quad (47)$$

Proof. We first prove the following:

Claim 1 *Inequalities (46)-(47) can be equivalently written as:*

$$\max_{x,y \in \partial \mathcal{B}_1} r \int_{\partial \mathcal{B}_2} \max\{x^T z, y^T z\} d\mathcal{H}^{m-1}(z) - x^T y < \frac{r}{2} \Delta^2, \quad \text{for } r = r_1, r_2. \quad (48)$$

Proof of claim As before, for notational simplicity, we denote the i th (resp. j th, k th) point by x (resp. y, z). By (41) and (42), inequalities (46)-(47) read respectively

$$\begin{aligned} \min_{x,y \in \partial \mathcal{B}_2} r_1 \int_{\partial \mathcal{B}_1} \min\{\|x - z\|^2, \|y - z\|^2\} d\mathcal{H}^{m-1}(z) - \|x - y\|^2 &> 2(r_1 - 1), \\ \min_{x,y \in \partial \mathcal{B}_1} r_2 \int_{\partial \mathcal{B}_2} \min\{\|x - z\|^2, \|y - z\|^2\} d\mathcal{H}^{m-1}(z) - \|x - y\|^2 &> 2(r_2 - 1). \end{aligned}$$

The first inequality, up to a change of variable, reads:

$$\min_{x,y \in \partial \mathcal{B}_1} r_1 \int_{\partial \mathcal{B}_2} \min\{\|x - z\|^2, \|y - z\|^2\} d\mathcal{H}^{m-1}(z) - \|x - y\|^2 > 2(r_1 - 1),$$

hence inequalities (46)-(47) read

$$\min_{x,y \in \partial \mathcal{B}_1} r \int_{\partial \mathcal{B}_2} \min\{\|x - z\|^2, \|y - z\|^2\} d\mathcal{H}^{m-1}(z) - \|x - y\|^2 > 2(r - 1), \quad \text{for } r = r_1, r_2$$

which, expanding the squares, gives

$$\min_{x,y \in \partial \mathcal{B}_1} r \int_{\partial \mathcal{B}_2} \|z\|^2 + 1 + \min\{-2x^T z, -2y^T z\} d\mathcal{H}^{m-1}(z) + 2x^T y - 2 > 2(r - 1), \quad \text{for } r = r_1, r_2. \quad (49)$$

Via a change of variables, we have

$$\begin{aligned} \int_{\partial \mathcal{B}_2} \|z\|^2 d\mathcal{H}^{m-1}(z) &= \int_{\partial \mathcal{B}_1} \|\Delta e_1 + z\|^2 d\mathcal{H}^{m-1}(z) \\ &= \int_{\partial \mathcal{B}_1} \|\Delta e_1\|^2 + \|z\|^2 + 2\Delta e_1^T z d\mathcal{H}^{m-1}(z) = \Delta^2 + 1. \end{aligned}$$

Hence (49) reads

$$\min_{x,y \in \partial \mathcal{B}_1} r \int_{\partial \mathcal{B}_2} \min\{-2x^T z, -2y^T z\} d\mathcal{H}^{m-1}(z) + 2x^T y > -r\Delta^2, \quad \text{for } r = r_1, r_2,$$

which is equivalent to (48). \diamond

Thanks to Claim 1, to conclude the proof of the lemma it suffices to show the following:

Claim 2 *Inequalities (48) hold if and only if $\Delta > \Delta_0$.*

Proof of claim We will prove that the maximum of the left-hand side of inequalities (48) over all $x, y \in \partial\mathcal{B}_1$ is attained at $(e_1, -e_1)$, for all $r \in (0, 2)$. Since $r_1 \in (0, 1]$ and $r_2 \in [1, 2)$, this in turn implies that inequalities (48) are satisfied if and only if

$$r\Delta + 1 = r \int_{\partial\mathcal{B}_2} z^T e_1 d\mathcal{H}^{m-1}(z) + 1 < \frac{r}{2}\Delta^2, \quad \text{for } r = r_1, r_2,$$

which, by $r_1 \leq r_2$, is true if and only if $\Delta > \Delta_0$; i.e., the desired condition. Define

$$F_r(x, y) := r \int_{\partial\mathcal{B}_2} \max\{x^T z, y^T z\} d\mathcal{H}^{m-1}(z) - x^T y. \quad (50)$$

Our goal is to show that for every $r \in (0, 2)$

$$\max_{x, y \in \partial\mathcal{B}_1} F_r(x, y) = F_r(e_1, -e_1) = r\Delta + 1. \quad (51)$$

In Lemma 2, we prove that

$$\max_{x, y \in \partial\mathcal{B}_1} F_2(x, y) = F_2(e_1, -e_1) = 2\Delta + 1. \quad (52)$$

Hence, recalling that $r \in (0, 2)$ and that $1 - \frac{r}{2} \geq -(1 - \frac{r}{2})x^T y$ for every $x, y \in \partial\mathcal{B}_1$, we conclude with the following chain of inequalities:

$$\begin{aligned} r\Delta + 1 &= \frac{r}{2}(2\Delta + 1) + (1 - \frac{r}{2}) = \frac{r}{2} \max_{x, y \in \partial\mathcal{B}_1} F_2(x, y) + (1 - \frac{r}{2}) \\ &= \max_{x, y \in \partial\mathcal{B}_1} \frac{r}{2} F_2(x, y) + (1 - \frac{r}{2}) \geq \max_{x, y \in \partial\mathcal{B}_1} F_r(x, y). \end{aligned} \quad (53)$$

◇

□

In order to prove Lemma 1, we made use of the next lemma, for which we provide a proof that is closely related to the proof of Lemma 1 in [8], in which the authors prove

$$\max_{x, y \in \mathcal{B}_1} F_1(x, y) = F_1(e_1, -e_1) = \Delta + 1,$$

where F_r is defined by (50) and \mathcal{B}_1 denotes a ball of radius one as defined in the SBM. For brevity, in the following, we only include the parts of the proof that are different, and when possible, we refer to the relevant parts of the proof of Lemma 1 in [8].

Lemma 2. (52) *holds.*

Proof. For any $x \in \mathbb{R}^m$, we denote by x_i the i th component of x . We divide the proof in several steps:

Step 1. Slicing:

Let z, w be any pair of points in $\partial\mathcal{B}_2$ satisfying $z_1 = w_1$, $z_2 = -w_2 \geq 0$, $z_j = w_j = 0$ for all $j \in \{3, \dots, m\}$. In the special case $m = 1$, we consider $z = w \in \partial\mathcal{B}_2$. Define

$$H(x, y) := \max\{x^T z, y^T z\} + \max\{x^T w, y^T w\} - x^T y.$$

Then (52) holds if the following holds

$$\max_{x, y \in \partial\mathcal{B}_1} H(x, y) = H(e_1, -e_1) \quad (54)$$

for any pair of points $z, w \in \partial\mathcal{B}_2$ satisfying $z_1 = w_1$, $z_2 = -w_2 \geq 0$, $z_j = w_j = 0$ for all $j \in \{3, \dots, m\}$ in case $m \geq 2$, or, in case $m = 1$, for any point $z = w \in \partial\mathcal{B}_2$.

Proof of Step 1. Since

$$F_2(x, y) = \frac{1}{\mathcal{H}^{m-1}(\partial B_2)} \int_{\Delta-1}^{\Delta+1} \int_{\{z_1=s\} \cap \partial B_2} 2 \max\{x^T z, y^T z\} - x^T y d\mathcal{H}^{m-2}(z) ds,$$

to show (52) it is enough to show that the function

$$G(x, y) := \int_{\{z_1=s\} \cap \partial B_2} 2 \max\{x^T z, y^T z\} - x^T y d\mathcal{H}^{m-2}(z)$$

is maximized in $x = e_1$, $y = -e_1$, for every $s \in [\Delta - 1, \Delta + 1]$. Denoting $A := \{z_1 = s, z_2 \geq 0\} \cap \partial B_2$, then

$$\frac{1}{2} G(x, y) = \int_A \max\{x^T z, y^T z\} + \max\{x^T(2se_1 - z), y^T(2se_1 - z)\} - x^T y d\mathcal{H}^{m-1}(z).$$

Hence, it is enough to prove that for every $s \in [\Delta - 1, \Delta + 1]$ and for every $z \in A$,

$$\max_{x, y \in \partial B_1} \{ \max\{x^T z, y^T z\} + \max\{x^T(2se_1 - z), y^T(2se_1 - z)\} - x^T y \}, \quad (55)$$

is achieved at $(e_1, -e_1)$. Since Problem (55) is invariant under a rotation of the space around the axis generated by e_1 , we conclude that solving Problem (55) is equivalent to solving Problem (54).

Step 2. Symmetric distribution of the maxima:

Let z, w be any pair of points as defined in Step 1. Define

$$I(x, y) := x^T z + y^T w - x^T y.$$

In order to show that (54) holds, it suffices to prove that for $m \geq 2$

$$\max_{\substack{x, y \in \partial B_1 \\ x^T z \geq y^T z, y^T w \geq x^T w}} \{I(x, y)\} \leq H(e_1, -e_1) = 2z_1 + 1. \quad (56)$$

Proof of Step 2. The proof is identical (up to trivial changes) to the proof of Step 2 of Lemma 1 in [8].

Step 3. Reduction from spheres to circles:

To show the validity of (56), we can restrict to dimension $m = 2$.

Proof of Step 3. The proof repeats verbatim as in the proof of Step 3 of Lemma 1 in [8].

Step 4. *Symmetric local maxima:*

For any pair $x, y \in \partial\mathcal{B}_1$ of the form $x_1 = y_1$ and $x_2 = -y_2$, we have

$$I(x, y) \leq H(e_1, -e_1).$$

Proof of Step 4. Given such symmetric pair (x, y) , the objective function evaluates to $I(x, y) = 2z_1x_1 + 2z_2x_2 - x_1^2 + x_2^2$. Using $x_1^2 + x_2^2 = 1$ and $z_2\sqrt{1-x_1^2} \leq z_2$, it suffices to show that

$$2z_2 \leq 2x_1^2 - 2z_1x_1 + 2z_1, \quad \forall x_1 \in [-1, 1]. \quad (57)$$

Since the function $\hat{f}(x_1) := 2x_1^2 - 2z_1x_1 + 2z_1$ on the right hand side of (57) is a convex parabola in x_1 , its minimum is either attained at one of the end points or at $\tilde{x}_1 = \frac{z_1}{2}$, provided that $-1 \leq \frac{z_1}{2} \leq 1$. Since $\Delta - 1 \leq z_1 \leq \Delta + 1$, the point \tilde{x}_1 lies in the domain only if $\Delta - 1 \leq z_1 \leq \min\{2, \Delta + 1\} = 2$. The value of \hat{f} at $x_1 = -1$ and $x_1 = 1$ evaluates to $2 + 4z_1$ and 2 , respectively, both of which are bigger than $2z_2$. Hence it remains to show that $\hat{f}(\tilde{x}_1) \geq 2z_2$ if $\Delta - 1 \leq z_1 \leq 2$, that is we have to show that

$$2\sqrt{1-u^2} \leq 2(u+\Delta) - \frac{(u+\Delta)^2}{2}, \quad \text{if } -1 \leq u \leq 2-\Delta,$$

where we set $u := z_1 - \Delta$ and we use that $z_2 = \sqrt{1-u^2}$. Since $u + \Delta \leq 2$, the right hand side of the above inequality is increasing in Δ ; hence it suffices to show its validity at $\Delta = 2$; i.e.,

$$2\sqrt{1-u^2} \leq 2(u+2) - \frac{(u+2)^2}{2}, \quad \text{if } -1 \leq u \leq 0,$$

which can be easily proved.

Step 5. *Decomposition of the circle:*

To solve Problem (56), it suffices to solve

$$\max_{\substack{x, y \in \partial\mathcal{B}_1 \cap \{x_1 \leq 0, y_1 \geq 0\} \\ x^T z \geq y^T z, y^T w \geq x^T w}} I(x, y) \leq H(e_1, -e_1) = 2z_1 + 1, \quad (58)$$

for every $z \in \partial\mathcal{B}_2$, $z_1 = w_1$, $z_2 = -w_2 \geq 0$.

Proof of Step 5. The proof is identical (up to trivial changes) to the proof of Step 6 of Lemma 1 in [8].

Step 6. *We solve Problem (58).*

Proof of Step 6. The proof is identical (up to trivial changes) to the proof of Step 7 of Lemma 1 in [8]. □

While the recovery guarantee of Proposition 4 is the first of its kind for an LP relaxation of K-means clustering, it is too conservative. We demonstrate this fact via numerical simulations. We let $n = 100$, $m = 2$, $r_1 \in \{0.6, 0.8, 1.0\}$, and $\Delta \in [2.0 : 0.01 : 3.2]$. For each fixed configuration, we generate 20 random trials according to the SSM. We count the number of times the optimization algorithm returns the planted clusters as the optimal solution; dividing this number by the total number of trials, we obtain the empirical recovery rate. We use the same set up as before to solve all LPs and SDPs. Our results are shown in Figure 1, where as before we compare the LP relaxation

defined by Problem (LP2) with the SDP relaxation defined by Problem (PW). Clearly, in all cases the LP outperforms the SDP. Moreover, our results indicate that the recovery threshold of the LP relaxation for SSM is significantly better than the one given by Proposition 4. For example, for $r_1 = 0.6$, condition (35) gives the recovery threshold $\Delta_0 \approx 3.08$, while Figure 1(c) suggests $\Delta_0 \approx 2.6$. We should also remark that in *all* these experiments, the LP relaxation is tight; i.e., whenever the LP fails in recovering the planted clusters, its optimal solution is still a partition matrix. In contrast, the SDP relaxation is not tight in almost all cases for which it does not recover the planted clusters.




Figure 1: The recovery rate of the LP versus the SDP when the input is generated according to the SSM with $n = 100$ and $m = 2$.

2.3 A stronger dual certificate

By Proposition 1, if inequalities (11) and (12) are satisfied then the partition matrix \bar{X} defined by (13) is an optimal solution of Problem (LP2) and as a result \bar{X} is an optimal solution of the K-means clustering problem. However, as evident from Proposition 4 and Figure 1, our proximity condition given by inequalities (11) and (12) are overly conservative. In the following we present a simple algorithm that leads to significantly better recovery guarantees.

Recall that in the last step of the proof of Proposition 1, our task is to identify conditions under which inequalities (26) and (27) can be satisfied. In the proof we let $\bar{\lambda}_{ijk} = 0$ for all $(i, j, k) \in \Gamma_1$ and $\bar{\lambda}_{ijk} = 0$ for all $(i, j, k) \in \Gamma_2$, which in turn gives us inequalities (26) and (27). As we show

next, a careful selection of these multipliers will lead to significantly better recovery results. Define

$$\gamma_{ij} := 2 \sum_{k \in \Gamma_2} \min\{r_2 d_{ik} + d_j^{\text{in}}, r_2 d_{jk} + d_i^{\text{in}}\} - 2d_{ij} - 2\eta, \quad \forall i < j \in \Gamma_1,$$

and

$$\gamma_{ij} := 2 \sum_{k \in \Gamma_1} \min\{r_1 d_{ik} + d_j^{\text{in}}, r_1 d_{jk} + d_i^{\text{in}}\} - 2d_{ij} - 2\frac{r_1}{r_2}\eta, \quad \forall i < j \in \Gamma_2.$$

We define $\gamma_{ji} := \gamma_{ij}$ for all $i < j$. Then, by the proof of Proposition 1, the partition matrix \bar{X} defined by (13) is an optimal solution of Problem (LP2) if the following system of inequalities is feasible:

$$\begin{aligned} \sum_{k \in \Gamma_{l'} \setminus \{i, j\}} (\lambda_{ijk} + \lambda_{jik} - \lambda_{kij}) &\leq \gamma_{ij}, \quad \forall i < j \in \Gamma_l, l \neq l' \in \{1, 2\} \\ \lambda_{ijk} &\geq 0, \quad \forall (i, j, k) \in \Gamma_l, l \in \{1, 2\}, \end{aligned} \quad (59)$$

where as before we let $\lambda_{ikj} = \lambda_{ijk}$ for $j < k$. We next present a simple algorithm whose successful termination serves as a sufficient condition for feasibility of system (59):

Certify : The algorithm for constructing a dual certificate

Input: Given γ_{ij} for all $i < j \in \Gamma_l, l \in \{1, 2\}$

Output: A Boolean success

Initialize $\bar{\lambda}_{ijk} = \bar{\lambda}_{jik} = \bar{\lambda}_{kij} = 0$ for all $(i, j, k) \in \Gamma_l, l \in \{1, 2\}$, $\bar{r}_{ij} = \gamma_{ij}$ for all $(i, j) \in \Gamma_l, l \in \{1, 2\}$ and $\mathcal{N} = \{(i, j) \in \Gamma_l, l \in \{1, 2\} : i < j, \bar{r}_{ij} < 0\}$.

while $\mathcal{N} \neq \emptyset$, **do**

set success = **.false.**

 select some $(i, j) \in \mathcal{N}$

for each $k \in \Gamma_l \setminus \{i, j\}$, **do**

 let $\omega = \min\{-\bar{r}_{ij}, \bar{r}_{ik}, \bar{r}_{jk}\}$.

 if $\omega \leq 0$, then cycle

 update $\bar{r}_{ik} \leftarrow \bar{r}_{ik} - \omega, \bar{r}_{jk} \leftarrow \bar{r}_{jk} - \omega$, and $\bar{r}_{ij} \leftarrow \bar{r}_{ij} + \omega$

 update $\bar{\lambda}_{kij} \leftarrow \bar{\lambda}_{kij} + \omega, \bar{\lambda}_{kji} \leftarrow \bar{\lambda}_{kji} + \omega$

if $\bar{r}_{ij} \geq 0$, **then**

 update success = **.true.**

 update $\mathcal{N} \leftarrow \mathcal{N} \setminus \{(i, j)\}$

exit the k-loop

if success = **.false.**, then **return**

Proposition 5. *If Algorithm Certify terminates with success = .true., then the partition matrix \bar{X} defined by (13) is an optimal solution of Problem (LP2). Moreover, Algorithm Certify runs in $\Theta(n^3)$ operations.*

Proof. We prove that the system defined by all inequalities in system (59) with $l = 1$ is feasible; the proof for $l = 2$ then follows. Define

$$\bar{r}_{ij} := \gamma_{ij} + \sum_{k \in \Gamma_2 \setminus \{i, j\}} (\bar{\lambda}_{kij} - \bar{\lambda}_{ijk} - \bar{\lambda}_{jik}), \quad \forall i < j \in \Gamma_1. \quad (60)$$

Then the system defined by all inequalities in system (59) with $l = 1$ can be equivalently written as:

$$\bar{r}_{ij} \geq 0, \quad \forall i < j \in \Gamma_1, \quad \bar{\lambda}_{ijk} \geq 0, \quad \forall (i, j, k) \in \Gamma_1. \quad (61)$$

If $\mathcal{N} = \emptyset$, then by letting $\bar{\lambda}_{ijk} = \bar{\lambda}_{jik} = \bar{\lambda}_{kij} = 0$ for all $(i, j, k) \in \Gamma_1$, we obtain a feasible solution and the algorithm terminates with `success = .true..` Hence, suppose that $\mathcal{N} \neq \emptyset$; that is, the initialization step violates inequalities $\bar{r}_{ij} \geq 0$ for all $(i, j) \in \mathcal{N}$. Consider an iteration of the algorithm for some $(\bar{i}, \bar{j}) \in \mathcal{N}$; we claim that if this iteration is completed with `success = .true..`, all nonnegative \bar{r}_{ij} , $i < j \in \Gamma_1$ remain nonnegative (even though their values may decrease), and we will have $\bar{r}_{\bar{i}\bar{j}} \geq 0$. Since the value of $\bar{\lambda}$ does not decrease over the course of the algorithm, this in turn implies that if the algorithm terminates with `success = .true..`, $(\bar{\lambda}, \bar{r})$ is feasible for system (61). To see this, consider some $\bar{k} \in \Gamma_1 \setminus \{\bar{i}, \bar{j}\}$ for which we have $\omega = \min\{-\bar{r}_{\bar{i}\bar{j}}, \bar{r}_{\bar{i}\bar{k}}, \bar{r}_{\bar{j}\bar{k}}\} > 0$. Notice that variable $\bar{\lambda}_{\bar{k}\bar{i}\bar{j}}$ (which is equal to $\bar{\lambda}_{\bar{k}\bar{j}\bar{i}}$) appears only in three equations of (60) defining $\bar{r}_{\bar{i}\bar{j}}$ (with positive coefficient), $\bar{r}_{\bar{i}\bar{k}}$ (with negative coefficient), and $\bar{r}_{\bar{j}\bar{k}}$ (with negative coefficient). Since $\bar{r}_{\bar{i}\bar{k}} \geq \omega$, $\bar{r}_{\bar{j}\bar{k}} \geq \omega$, and $\omega > 0$, we can increase the value of $\bar{\lambda}_{\bar{k}\bar{i}\bar{j}}$ by ω and keep the system feasible, this in turn implies that the value of $\bar{r}_{\bar{i}\bar{j}}$ will increase by ω , while the values of $\bar{r}_{\bar{i}\bar{k}}$ and $\bar{r}_{\bar{j}\bar{k}}$ decrease by ω . Therefore, if the algorithm terminates with `success=.true..`, the assignment $(\bar{\lambda}, \bar{r})$ is a feasible solution for system (61). It is simple to verify that this algorithm runs in $\Theta(n^3)$ operation. \square

Let us now comment on the power of Algorithm `Certify` for recovering the planted clusters under the SBM. Since we do not have an explicit proximity condition, we are unable to perform a rigorous probabilistic analysis similar to that of Section 2.2. However, in dimension $m = 2$, we can perform a high precision simulation as the computational cost of Algorithm `Certify` is very low. First, we consider the case of equal-size clusters; i.e., $r_1 = r_2 = 1$. We set $n = 20000$ and we observe that for $\Delta > 2.14$, Algorithm `Certify` terminates with `success = .true..` We then conjecture the following:

Conjecture 1. *Let $m \geq 2$, and suppose that the points are generated according to the SBM with equal-size clusters. If $\Delta > 2.14$, then Problem (LP2) recovers the planted clusters with high probability.*

If true, the recovery guarantee of Conjecture 1 is better than the recovery guarantee of the SDP relaxation given by (34) for $m \leq 202$. Clearly, any convex relaxation succeeds in recovering the underlying clusters only if the original problem succeeds in doing so. To this date, the recovery threshold for K-means clustering under the SBM for $m > 1$ remains an open question. Let us briefly discuss special cases $m = 1$ and $m = 2$. In [12], the authors prove that a necessary condition for recovery of K-means clustering in dimension one is $\Delta > 1 + \sqrt{3}$. In the same paper, the recovery threshold of the SDP relaxation in dimension one is conjectured to be $\Delta_0 = 4$ (see Section 2.3 of [12] for a detailed discussion). In [8], the authors prove that Problem (LP2) recovers the planted clusters with high probability (for every $m \geq 1$), if $\Delta > 1 + \sqrt{3}$. It then follows that for $m = 1$, the K-means clustering problem recovers the planted clusters with high probability if and only if $\Delta > 1 + \sqrt{3}$. For $m = 2$, in [12], via numerical simulations, the authors show that K-means clustering recovers the planted clusters with high probability only if $\Delta > 2.08$. If true, Conjecture 1 implies that in dimension two, the recovery thresholds for K-means clustering and the LP relaxation are fairly

close. In [15], the authors prove that, if

$$\Delta < 1 + \sqrt{1 + \frac{2}{m+2}}, \quad (62)$$

then the SDP relaxation fails in recovering the planted clusters with high probability. They also state that “it remains unclear whether this necessary condition (i.e., inequality (62)) is only necessary for the SDP relaxation or is necessary for the K-means itself.” If true, Conjecture 1 implies that inequality (62) is *not* a necessary condition for the K-means clustering problem.

We further use Algorithm `Certify` to estimate the recovery threshold for the case with different cluster sizes, i.e. $r_1 \neq 1$, in dimension $m = 2$. Our results are depicted in Figure 2. As can be seen from the figure, the recovery guarantee given by Algorithm `Certify` is significantly better than that of Proposition 4. For comparison, we have also plotted the recovery threshold for the SDP relaxation given by condition (34) for different input dimensions m . While Figure 2 suggests that the recovery threshold of the LP as given by Algorithm `Certify` quickly degrades as r_1 decreases, our numerical experiments with the LP relaxation, depicted in Figure 1 for $0.6 \leq r_1 \leq 1.0$, indicates otherwise. Notice that Algorithm `Certify` provides a *sufficient condition* for feasibility of system (59). Obtaining a better sufficient condition or, if possible, a tractable necessary and sufficient condition for the feasibility of system (59) remains an open question.




Figure 2: Comparing the recovery threshold Δ_0 of the LP relaxation versus the SDP relaxation for K-means clustering when the input is generated according to the SBM.

3 A Counter example for the tightness of the LP relaxation

As we discussed in the previous section, in all our experiments with both synthetic and real-world data sets, Problem (LP2) is tight; i.e., its optimal solution is a partition matrix. In particular, a large number of our experiments are done under the SBM and SSM. Hence, it is natural to ask whether the LP relaxation is tight with high probability under reasonable generative models. In the following we present a family of inputs for which the LP relaxation is never tight. This family of inputs subsumes as a special case the points generated by a variation of the SBM. To prove our result, we make use of the following lemma, which essentially states that if several sets of points have identical optimal cluster centers, then those cluster centers are optimal for the union of all points as well:

Lemma 3. Let $\{x^{i,l} : i \in [n_l], l \in [L]\}$ denote a set of $N := \sum_{l \in [L]} n_l$ points in \mathbb{R}^m for some $n_1, \dots, n_L, L \in \mathbb{N}$, with $n_1, \dots, n_L, L \geq 1$. For each $l \in [L]$, consider an optimal solution of Problem (1) for clustering the n_l points $\{x^{i,l}\}_{i=1}^{n_l}$ and denote by $\bar{c}^l \in \mathbb{R}^K$ the vector with entries equal to the K centers of the optimal clusters. If $\bar{c}^l = \bar{c}^{l'}$ for all $l, l' \in [L]$, then there exists an optimal solution of Problem (1) for clustering the entire set of N points with cluster centers vector $\bar{c} \in \mathbb{R}^K$ satisfying $\bar{c}^l = \bar{c}$ for all $l \in [L]$.

Proof. Let $\{y^i\}_{i=1}^n$ denote a set of n points in \mathbb{R}^m that we would like to put into K clusters and denote by $c \in \mathbb{R}^K$ the vector of cluster centers. We start by reformulating Problem (1) as follows:

$$\min_{c \in \mathbb{R}^K} \sum_{i=1}^n \min \left\{ \|y^i - c_1\|_2^2, \dots, \|y^i - c_K\|_2^2 \right\}. \quad (63)$$

Notice that by solving Problem (63) one directly obtains a vector of optimal cluster centers and subsequently can assign each point y^i to a cluster at which $\min \left\{ \|y^i - c_1\|_2^2, \dots, \|y^i - c_K\|_2^2 \right\}$ is attained. Now for each $l \in [L]$, define

$$f_l(c) := \sum_{i=1}^{n_l} \min \left\{ \|x^{i,l} - c_1\|_2^2, \dots, \|x^{i,l} - c_K\|_2^2 \right\}.$$

It then follows that the K-means clustering problem for clustering the n_l points $\{x^{i,l}\}_{i=1}^{n_l}$ for some $l \in [L]$ can be written as:

$$\min_{c \in \mathbb{R}^K} f_l(c), \quad (64)$$

while the K-means clustering problem for the entire set of N points can be written as

$$\min_{c \in \mathbb{R}^K} \sum_{l \in [L]} f_l(c). \quad (65)$$

Clearly, $\sum_{l \in [L]} \min_{c \in \mathbb{R}^K} f_l(c) \leq \min_{c \in \mathbb{R}^K} \sum_{l \in [L]} f_l(c)$. Hence, if for each $l \in [L]$, there exists a minimizer \bar{c}^l of Problem (64) such that $\bar{c}^l = \bar{c}^{l'}$ for all $l, l' \in [L]$, we conclude that \bar{c}^l is a minimizer of Problem (65) as well. \square

Now let us investigate the tightness of the LP relaxation defined by Problem (LP2). First note that by Proposition 5 in [8], if $n \leq 4$, then the feasible region of Problem (LP2) coincides with the convex hull of the feasible region of Problem (3). Hence, to find an instance for which the LP relaxation is not tight we must have $n \geq 5$. Consider the following $n = 5$ points in dimension $m = 3$, which we refer to as the *five-point input*:

$$x^1 = \left(0, \frac{\sqrt{3}}{3}, 0\right), x^2 = \left(\frac{1}{2}, -\frac{\sqrt{3}}{6}, 0\right), x^3 = \left(-\frac{1}{2}, -\frac{\sqrt{3}}{6}, 0\right), x^4 = \left(0, 0, \frac{1}{2}\right), x^5 = \left(0, 0, -\frac{1}{2}\right). \quad (66)$$

In this case the vector of squared pair-wise distances $d = (d_{ij})_{1 \leq i < j \leq 5}$ is given by

$$d = \left(1, 1, \frac{7}{12}, \frac{7}{12}, 1, \frac{7}{12}, \frac{7}{12}, \frac{7}{12}, \frac{7}{12}, 1\right). \quad (67)$$

By direct calculation it can be checked that the partition $\Gamma_1 = \{1, 4\}$ and $\Gamma_2 = \{2, 3, 5\}$ is a minimizer of the K-means clustering problem (1) with the optimal objective value given by

$$f_{\text{Kmeans}}^* = \frac{146}{72} > 2.$$

Now consider the following matrix

$$\tilde{X} = \frac{1}{14} \begin{bmatrix} 6 & 1 & 1 & 3 & 3 \\ 1 & 6 & 1 & 3 & 3 \\ 1 & 1 & 6 & 3 & 3 \\ 3 & 3 & 3 & 5 & 0 \\ 3 & 3 & 3 & 0 & 5 \end{bmatrix}. \quad (68)$$

Notice that \tilde{X} is not a partition matrix. It is simple to verify the feasibility of \tilde{X} for the LP relaxation. Moreover, the objective value of Problem (LP2) at \tilde{X} evaluates to $\hat{f}_{\text{LP}} = \frac{54}{28} < 2$. Denote by f_{LP}^* the optimal value of Problem (LP2). Since $f_{\text{LP}}^* \leq \hat{f}_{\text{LP}} < f_{\text{Kmeans}}^*$, we conclude that for the five-point input, the LP relaxation is not tight.

Now consider the following set of points, which we will refer to as the *five-ball input*. Instead of $n = 5$ points located at each x^p , $p \in \{1, \dots, 5\}$ defined by (66), suppose that we have $n = 5n'$ points for some $n' \in \mathbb{N}$, $n' \geq 1$, such that for each $p \in \{1, \dots, 5\}$, we have n' points located inside a closed ball of radius $r \geq 0$ centered at x^p . In the following we show that for $r \leq 3 \times 10^{-3}$, the LP relaxation always fails in finding the optimal clusters.

Proposition 6. *Consider an instance of the five-ball input for some $r \leq 3 \times 10^{-3}$. Then the LP relaxation is not tight; i.e., the optimal value of Problem (LP2) is strictly smaller than that of Problem (1).*

Proof. We denote by \mathcal{B}_p the index set of points located in the p th ball; without loss of generality, let $\mathcal{B}_p = \{(p-1)n' + 1, \dots, pn'\}$ for all $p \in \{1, \dots, 5\}$. Let us denote these points by y^j , $j \in \mathcal{B}_p$, $p \in \{1, \dots, 5\}$. We start by computing an upper bound on the optimal value of the LP relaxation. Consider the matrix \hat{X} defined as follows:

$$\begin{aligned} \hat{X}_{ij} &= \frac{6}{14n'}, & \forall i, j \in \mathcal{B}_p, p \in \{1, 2, 3\} \\ \hat{X}_{ij} &= \frac{5}{14n'}, & \forall i, j \in \mathcal{B}_p, p \in \{4, 5\} \\ \hat{X}_{ij} &= \frac{1}{14n'}, & \forall i \in \mathcal{B}_p, j \in \mathcal{B}_{p'}, p \neq p' \in \{1, 2, 3\} \\ \hat{X}_{ij} &= \frac{3}{14n'}, & \forall i \in \mathcal{B}_p, j \in \mathcal{B}_{p'}, p \in \{1, 2, 3\}, p' \in \{4, 5\} \\ \hat{X}_{ij} &= 0, & \forall i \in \mathcal{B}_p, j \in \mathcal{B}_{p'}, p \neq p' \in \{4, 5\}. \end{aligned}$$

Clearly, \hat{X} is not a partition matrix. First we show that \hat{X} is feasible for Problem (LP2). The equality constraint (5) is satisfied as we have $\text{Tr}(\hat{X}) = \sum_{i \in [n]} \hat{X}_{ii} = 3n'(\frac{6}{14n'}) + 2n'(\frac{5}{14n'}) = 2$. Equalities (6) are also satisfied since we have $\sum_{j=1}^n \hat{X}_{ij} = n'(\frac{6}{14n'}) + 2n'(\frac{1}{14n'}) + 2n'(\frac{3}{14n'}) = 1$ for all $i \in \mathcal{B}_p, p \in \{1, 2, 3\}$, and $\sum_{j=1}^n \hat{X}_{ij} = 3n'(\frac{3}{14n'}) + n'(\frac{5}{14n'}) + n'(0) = 1$ for all $i \in \mathcal{B}_p, p \in \{4, 5\}$. Hence it remains to check the validity of inequalities (7): first note that if $i \in \mathcal{B}_p, j \in \mathcal{B}_{p'}, k \in \mathcal{B}_{p''}$ for $p \neq p' \neq p''$, then the validity of $\hat{X}_{ij} + \hat{X}_{ik} \leq \hat{X}_{ii} + \hat{X}_{jk}$ follows from the validity of inequalities (7) at \tilde{X} defined by (68). If $i, j, k \in \mathcal{B}_p, p \in \{1, 2, 3\}$, then $\hat{X}_{ij} + \hat{X}_{ik} = \frac{6}{14n'} + \frac{6}{14n'} \leq \frac{6}{14n'} + \frac{6}{14n'} = \hat{X}_{ii} + \hat{X}_{jk}$ for all $i \neq j \neq k, j < k$. If $i, j, k \in \mathcal{B}_p, p \in \{4, 5\}$, then $\hat{X}_{ij} + \hat{X}_{ik} = \frac{5}{14n'} + \frac{5}{14n'} \leq \frac{5}{14n'} + \frac{5}{14n'} = \hat{X}_{ii} + \hat{X}_{jk}$ for all $i \neq j \neq k, j < k$. The remaining cases, i.e., when two of the points are in the same ball, while the third point is in a different ball, can be checked in a similar fashion.

Denote by d_{ij}^r , $1 \leq i < j \leq n$, the vector of squared pairwise distances. We have:

$$\begin{aligned} d_{ij}^r &\leq 4r^2, \quad \forall i, j \in \mathcal{B}_p, p \in \{1, \dots, 5\}, \\ d_{ij}^r &\leq (1 + 2r)^2, \quad \forall i \in \mathcal{B}_p, j \in \mathcal{B}_{p'}, \forall p \neq p' \in \{1, 2, 3\}, \text{ or } p \neq p' \in \{4, 5\}, \\ d_{ij}^r &\leq \left(\sqrt{\frac{7}{12}} + 2r \right)^2, \quad \forall i \in \mathcal{B}_p, j \in \mathcal{B}_{p'}, \forall p \in \{1, 2, 3\}, p' \in \{4, 5\}. \end{aligned}$$

Hence, an upper bound $\hat{g}_{\text{LP}}(r)$ on the optimal value of Problem (LP2) is given by

$$\begin{aligned} g_{\text{LP}}^*(r) &\leq 6 \binom{n'}{2} \frac{6}{14n'} (4r^2) + 4 \binom{n'}{2} \frac{5}{14n'} (4r^2) + 6n'^2 \frac{1}{14n'} (1 + 2r)^2 + 12n'^2 \frac{3}{14n'} \left(\sqrt{\frac{7}{12}} + 2r \right)^2 \\ &= n' \left(\frac{54}{28} + 20r^2 + 12 \left(\frac{\sqrt{21} + 1}{7} \right) r - 8 \frac{r^2}{n'} \right) \leq \left(\frac{54}{28} + 10r \right) n' := \hat{g}_{\text{LP}}(r), \end{aligned} \quad (69)$$

where the last inequality follows since by assumption $r \leq 3 \times 10^{-3}$, and where $g_{\text{LP}}^*(r)$ denotes the optimal value of Problem (LP2) for the five-ball input.

Denote by $g_{\text{Kmeans}}^*(r)$ the optimal value of the K-means clustering problem (1) for the five-ball input. Next, we obtain a lower bound on $g_{\text{Kmeans}}^*(r)$, and show that this value is strictly larger than $\hat{g}_{\text{LP}}(r)$ defined by (69) for any $0 \leq r \leq 3 \times 10^{-3}$, implying that Problem (LP2) is not tight. To this end, first consider the case with $r = 0$. Define the set of points $Y^l := \{y^l, y^{n'+l}, y^{2n'+l}, y^{3n'+l}, y^{4n'+l}\}$ for all $l \in [n']$. Notice that since $r = 0$, we have $Y^l = Y^{l'}$ for all $l, l' \in [n']$. For each $l \in [n']$, consider the K-means clustering problem with $K = 2$ for five input points Y^l , and denote the optimal cluster centers by (c_1^l, c_2^l) . We then have $(c_1^l, c_2^l) = (c_1, c_2)$ for all $l \in [n']$, where (c_1, c_2) denotes the optimal cluster centers for the five-point input (66). Therefore, by Lemma 3, we conclude that an optimal cluster centers for the five-ball input with $r = 0$, coincides with an optimal cluster centers for the five-point input. This in turn implies that

$$g_{\text{Kmeans}}^*(0) = n' f_{\text{Kmeans}}^* = \frac{146}{72} n'. \quad (70)$$

Now let us consider the five-ball input for some $r > 0$. We observe that

$$\sqrt{d_{ij}^r} \geq \sqrt{d_{ij}^0} - 2r, \quad \forall 1 \leq i < j \leq n.$$

Since $d_{ij}^0 \leq 1$ for all $1 \leq i < j \leq n$, we have $d_{ij}^r \geq d_{ij}^0 - 4r$ for all $1 \leq i < j \leq n$. Therefore, for every partition matrix X we compute

$$\sum_{i,j \in [n]} d_{ij}^r X_{ij} \geq \sum_{i,j \in [n]} d_{ij}^0 X_{ij} - 4r \sum_{i,j \in [n]} X_{ij} = \sum_{i,j \in [n]} d_{ij}^0 X_{ij} - 20rn',$$

where the last equality follows from constraints (6). Hence, taking the minimum over all partition matrices on both sides and using (70), we deduce that

$$g_{\text{Kmeans}}^*(r) \geq g_{\text{Kmeans}}^*(0) - 20rn' = \left(\frac{146}{72} - 20r \right) n'. \quad (71)$$

From (69) and (71) it follows that by choosing

$$r < \frac{\frac{146}{72} - \frac{54}{28}}{30},$$

we get $g_{\text{LP}}^*(r) < g_{\text{Kmeans}}^*(r)$ and this completes the proof. \square

As a corollary of Proposition 6, we find that the LP relaxation is not tight for a variant of the SBM:

Corollary 1. *Let $r \leq 3 \times 10^{-3}$. Suppose that the points are generated in \mathbb{R}^m , for some $m \geq 3$ according to any generative model consisting of five measures μ_p , $p = 1, \dots, 5$, each supported in the ball $B((x^p, 0, \dots, 0), r)$, where x^p is defined by (66). Then Problem (LP2) is not tight.*

Recall that in all our previous numerical experiments with synthetic and real-world data sets, the LP relaxation outperforms the SDP relaxation. That is, the optimal value of the LP is always at least as large as that of the SDP. Hence, one wonders whether such a property can be proved in a general setting. Interestingly, the stochastic model defined in Corollary 1 provides the first counter example, which we illustrate via a numerical experiment. We consider the stochastic model defined in Corollary 1, where we assume the points supported by each ball are sampled from a uniform distribution. We set $m = 3$ and generate $n' = 20$ points in each of the five balls to get a total of $n = 100$ points. Moreover we set ball radii $r \in [0.0 : 0.01 : 0.5]$ and for each fixed r we generate 20 random instances. Our results are depicted in Figure 3, where as before we compare the LP, i.e., Problem (LP2), with the SDP, i.e., Problem (PW), with respect to their tightness rate and average relative gap. Recall that the relative gap is defined as $g_{\text{rel}} = \frac{f_{\text{LP}} - f_{\text{SDP}}}{f_{\text{LP}}} \times 100$, where f_{LP} and f_{SDP} denote the optimal values of the LP and the SDP, respectively. That is, a negative relative gap means that the SDP relaxation is stronger than the LP relaxation. As can be seen from the figure, while the SDP is never tight, for $r \lesssim 0.15$, we often have $f_{\text{SDP}} > f_{\text{LP}}$.




Figure 3: Comparing the strength of the LP relaxation versus the SDP relaxation for K-means clustering when the input is generated according to the stochastic five-ball input with $n = 100$ and $m = 3$.

We conclude this section by noting that the family of examples considered above are very special and our numerical experiments with real-world data sets suggest that such special configurations do not appear in practice. Hence, it is highly plausible that one can establish the tightness of the LP relaxation under a fairly general family of inputs. We leave this as an open question.

4 A customized algorithm for solving the LP relaxation

As we detailed in the previous section, Problem (LPK_t) contains $\Theta(n^{t+1})$ inequalities of the form (4). This in turn implies that even in the cheapest case; i.e., $t = 2$, the computational cost of solving the LP relaxation quickly increases making it impractical for many real-world applications. To further illustrate this fact, we consider three state-of-the-art LP solvers:

- **Gurobi 11.02** [11], we use the interior-point algorithm and turn off the **crossover** operator to expedite the solver,
- **PDLP 9.10** [4], a primal-dual first-order method for solving large-scale LPs that is implemented in Google’s OR-Tools,
- **cuPDLP-C** [18], an improved implementation of **cuPDLP** in the **C** language, where **cuPDLP** [17] is a GPU implementation of **PDLP** and is written in **Julia**.

We first show that even for relatively small data sets i.e., $n < 500$, none of the solvers listed above are able to solve Problem (LPK_t) within 4 hours. Again, we consider a number of data sets from the UCI machine learning repository [9]. The data sets and their characteristics, i.e., the number of points n and the number of clusters K are listed in columns 1-3 of Table 2. We solve Problem (LPK_t) with $t = 2$ using the three LP solvers listed above. The size of each LP, i.e., the number of variables n_v , the number of constraints n_e , and the number of nonzeros in the constraint matrix n_z are listed in columns 4-6 of Table 2. Henceforth, all experiments are conducted on **Google Colab** using an Intel(R) Xeon(R) CPU @ 2.20GHz with 8 cores and 50 GB of RAM. The GPU used is an NVIDIA L4 with 22.5 GB of RAM. The total time in seconds of solving the LP using each solver is listed in the last three columns of Table 2. As can be seen from this table, for all problems that fit in the GPU memory, **cuPDLP-C** significantly outperforms other solvers. Nonetheless, for problems with $n \geq 400$, the LP becomes excessively expensive for all solvers. It is interesting to note that with the exception of **glass** and **WholeSale** (with $K = 6$) data sets, for all instances, the LP relaxation is tight; that is, it returns a partition matrix upon termination. As we detail shortly, for **glass** and **WholeSale** data sets, Problem (LPK_t) is tight if $t = 3$ and $t = 6$, respectively.

Motivated by this numerical study, we next present a cutting-plane based algorithm that enables us to solve Problem (LPK_t) for significantly larger data sets. The key to the success of this algorithm is to solve a sequence of LPs each of which contains a small but carefully selected subset of inequalities (4). Next, we describe the main ingredients of our algorithm¹.

4.1 Initialization

Lloyd’s algorithm and its enhanced implementations such as **k-means++** are extremely fast and are often successful in finding high quality solutions for the K-means clustering problem. In our framework, we make use of **k-means++**’s solution at the initialization step. First, the cost associated with this solution serves as an (often very good) upper bound on the optimal value of Problem (LPK_t). Second, to construct the first LP, we select inequalities (4) with $t = 2$ that are satisfied tightly at the **k-means++**’s solution. This in turn implies that if the **k-means++**’s solution is globally optimal, and if the optimal value of Problem (LPK_t) with $t = 2$ coincides with that of the K-means clustering problem, then our algorithm terminates after one iteration. For large scale data sets, however, the number of inequalities (4) with $t = 2$ that are satisfied tightly at the **k-means++**’s solution might be too large; in those cases, we randomly select a subset of at most p_{init} of such inequalities.

¹The source code as well as the data sets are available at <https://github.com/Yakun1125/cutLPK/tree/main>

Table 2: Execution time (secs) for solving Problem (LPK_t) with $t = 2$ using different LP solvers for small data sets from UCI.

Data set	n	K	n_v	n_e	n_z	cuPDLP-C	PDLP	Gurobi
ruspini	75	4	2850	2.03×10^5	8.16×10^5	1.0	4.0	3.9
iris	150	2	11325	1.65×10^6	6.64×10^6	5.6	31.0	33.4
		3				7.9	46.6	45.1
		4				7.2	41.1	59.8
wine	178	2	15931	2.77×10^6	1.11×10^7	14.4	73.7	79.4
		7				20.0	98.9	195.8
gr202	202	6	20503	4.06×10^6	1.63×10^7	131.9	403.1	868.7
seeds	210	2	22155	4.56×10^6	1.83×10^7	24.3	207.2	163.1
		3				19.3	115.6	200.6
glass	214	3 ¹	23005	4.83×10^6	1.94×10^7	116.2	845.1	651.0
		6 ¹				168.3	864.3	1394.6
CatsDogs	328	2	53956	1.75×10^7	7.00×10^7	1245.8	2894.2	1104.2
accent	329	2	54285	1.76×10^7	7.07×10^7	148.9	713.8	1395.9
		6				271.0	982.7	8401.9
ecoli	336	3	56616	1.88×10^7	7.53×10^7	129.7	547.3	5729.5
RealEstate	414	3	85905	3.52×10^7	1.41×10^8	N/A ²	3974.6	> 14400
		5				N/A	5708.4	> 14400
wholesale	440	5	97020	4.23×10^7	1.69×10^8	N/A	2944.4	N/A
		6 ¹				N/A	> 14400	N/A

¹ Problem (LPK_t) with $t = 2$ is not tight.

² Exceeds the maximum memory of the machine.

4.2 Safe lower bounds

To further expedite the cutting plane algorithm for large scale problems, in the first few iterations, we solve LPs with a larger feasibility tolerance and terminate early, if needed. In such cases the solution returned by the LP solver is often infeasible. To generate valid lower bounds on the optimal value of Problem (LPK_t), we make use of a technique that is has been used in mixed-integer programming solvers [24]. Consider the following pair of primal-dual LPs:

$$\begin{aligned}
 \min \quad & c^T x & \max \quad & b^T y \\
 \text{s.t.} \quad & Ax = b, \quad Qx \leq 0 & \text{s.t.} \quad & A^T y + Q^T z \leq c \\
 & x \geq 0, & & z \leq 0.
 \end{aligned}$$

Denote by $(\hat{x}, \hat{y}, \hat{z})$ an infeasible primal-dual pair returned by the LP solver satisfying $\hat{z} \leq 0$. Define the dual residual r as $r := \max\{A^T \hat{y} + Q^T \hat{z} - c, 0\}$. We then have $c \geq A^T \hat{y} + Q^T \hat{z} - r$. Let x^* denote an optimal solution for the primal LP. Using $Ax^* = b$, we get $c^T x^* \geq \hat{y}^T b + \hat{z}^T Qx^* - r^T x^*$. From $Qx^* \leq 0$ and $\hat{z} \leq 0$, we deduce that $\hat{z}^T Qx^* \geq 0$ implying that $c^T x^* \geq \hat{y}^T b - r^T x^*$. Denote by \bar{x} an upper bound on x^* . We then obtain the following “safe lower bound” on the optimal value of the primal LP:

$$c^T x^* \geq \hat{y}^T b - r^T \bar{x}.$$

4.3 Rounding

Denote by X_{lb} an optimal solution of the LP relaxation at an iteration of the cutting plane algorithm and suppose that X_{lb} is not a partition matrix. It is then desirable to “round” this solution to obtain a partition matrix X_{ub} whose cost may serve as good upper bound on the optimal clustering cost. We make use of the rounding scheme proposed in [25], which is given below.

round : The rounding scheme to obtain a partition matrix

Input: Data points: $\{x^i\}_{i=1}^n$, number of clusters: K , LP solution X_{lb}

Output: Partition matrix X_{ub}

Apply Singular Value Decomposition (SVD) to compute the K largest eigenvalues of X_{lb} with the corresponding eigen vectors v_i , $i \in [K]$. Compute initial cluster centers:

$$c_i = v_i^T W, \quad \forall i \in [K],$$

where W denotes a matrix whose i th column equals x^i .

Apply Lloyd’s algorithm using $\{c_i\}_{i=1}^K$ as the initial cluster centers to obtain a partition matrix X_{ub} .

4.4 The separation algorithm

Recall that the key to the efficiency of a cutting plane algorithm to solve Problem (LPK_{*t*}) is the efficient separation of inequalities (4). That is, each intermediate LP should contain a small number of inequalities (4) that are as sparse as possible. Let \tilde{X} be the solution to the current LP. Fix $t \in \{2, \dots, K\}$. For each $i \in [n]$, the separation problem is to find a nonempty subset $S \subseteq [n] \setminus \{i\}$ with $|S| \leq t$ such that

$$w_i(S) := \sum_{j \in S} \tilde{X}_{ij} - \tilde{X}_{ii} - \sum_{j, k \in S: j < k} \tilde{X}_{jk} > 0,$$

or to prove that no such subset exists. This separation problem is similar to the separation of clique inequalities (see for example [21]). In this setting, we can avoid repeated calculations using the following relation:

$$w_i(S \cup \{k\}) = w_i(S) + \tilde{X}_{ik} - \sum_{v \in S} \tilde{X}_{vk},$$

where we use the identity $\tilde{X}_{ij} = \tilde{X}_{ji}$ for all $i < j$. However, this enumeration scheme is too expensive for large scale problems. To this end, we utilize the greedy strategy proposed in [21] for separating clique inequalities. Our heuristic is as follows: for each $i \in [n]$, instead of enumerating all subsets S , we start by setting $S = \{j\}$ for some $j \in [n] \setminus \{i\}$. The algorithm then proceeds by expanding S iteratively, where in each iteration, an index $k \in \{j + 1, \dots, n\}$ that maximizes $w_i(S \cup \{k\})$ is appended to S .

An outline of the separation scheme is provided in Algorithm **separate**. Clearly, this algorithm is not exact, as the greedy expansion of S may lead to exclusion of some violated inequalities. Hence we apply the following two step approach. We first run Algorithm **separate**; if no violated inequalities are found and $t_{\max} = K$ while $r_g > \epsilon_{\text{opt}}$, then we switch to enumeration. However, in

separate : The heuristic algorithm for separating inequalities (4)

Input: LP solution \tilde{X} , violation tolerance ϵ_{vio} , and t_{max}

Output: A number of inequalities of the form (4) violated at \tilde{X} .

for $i \in [n]$ *in parallel* **do**

for $j \in [n] \setminus \{i\}$ **do**

 Initialize $S = \{j\}$, $w_i(S) = X_{ij} - X_{ii}$, and $c = j$

while $|S| < t_{\text{max}}$ **do**

 select $k \in \{c + 1, \dots, n\} \setminus (S \cup \{i\})$ that maximizes $\gamma_i(k) = X_{ik} - \sum_{l \in S, l < k} X_{lk}$.

 Update $\bar{S} = S \cup \{k\}$, and $c = k$

 Compute $w_i(\bar{S}) = w_i(S) + \gamma_i(k)$.

if $w_i(\bar{S}) > \epsilon_{\text{vio}}$ **then**

 Add $\sum_{k \in \bar{S}} X_{ik} \leq X_{ii} + \sum_{l, k \in \bar{S}: l < k} X_{lk}$ to the system of violated inequalities.

 Update $S = \bar{S}$

our numerical experiments, for all problems instances, Algorithm **separate** was enough to close the optimality gap r_g . We should also remark that, to separation inequalities (4), we run n separation problems, each for a fixed $i \in [n]$, in parallel. We make use of **OpenMP** [7] to do the parallel computations.

We are now ready to present our iterative cutting-plane based algorithm. An overview of this algorithm is given in Algorithm **Iterative LP solver**. Notice that in **Iterative LP solver**, the sparsity of the inequalities added to the LP is controlled by parameter t_{max} , which is initially set to $t_{\text{max}} = 2$, and is increased by one, only if the number of violated inequalities (4) with $t \leq t_{\text{max}}$ that are found by Algorithm **separate** is below a certain threshold.

4.5 Numerical Experiments

We start by testing **Iterative LP solver** using our three LP solvers as before for small data sets from UCI. Results are shown in Table 3. Notice that **Iterative LP solver** is not a deterministic algorithm; that is, at initialization, if the total number of eligible inequalities exceeds p_{init} , a random subset is selected and added to the LP. Hence, for a fair comparison, we run each instance 5 times and report average time and average number of iterations along with standard deviations. We set the optimality tolerance $\epsilon_{\text{opt}} = 10^{-4}$. Results are summarized in Table 3. As can be seen from the table, even with the cut generation scheme, **Gurobi** is unable to solve instances with more than 400 data points within 4 hours of CPU time. On the other hand, both first-order method LP solvers solve all instances within the time limit. *Surprisingly, in all cases, the LP relaxation is tight; i.e., the optimality gap r_g upon termination is smaller than the optimality tolerance ϵ_{opt} .* Moreover, **cuPDLP-C** is on average 5.7 times faster than **PDLP**. As a byproduct, we can verify that out of 19 instances, the SDP relaxation is tight for 2 instances only. In addition, **k-means++**'s solution is optimal for 12 instances.

Motivated by the impressive performance of **Iterative LP solver** for small data sets, next we consider larger data sets. We consider the first-order LP solvers only, as **Gurobi** is unable to solve any of the larger instances. Results are shown in Table 4. While **cuPDLP-C** solves all 32 instances, **PDLP** solves only 24 instances within the time limit. Moreover, for the latter 24 instances, **cuPDLP-C**

Iterative LP solver : The cutting plane algorithm for solving Problem (LPK_t)

Input: Data points $\{x^i\}_{i=1}^n$, number of clusters K , optimality tolerance ϵ_{opt} , initial number of inequalities p_{init} , number of inequalities added at each round of cut generation p_{max} , and the solver time limit for each intermediate LP T .

Output: Partition matrix X_{ub} and optimality gap r_g .

Initialize: Set the lower bound $f_{lb} = -\infty$ and the optimality gap $r_g = +\infty$. Run **k-means++** to get a partition matrix X_{ub} . Set the upper bound f_{ub} as the cost of X_{ub} . Randomly select at most p_{init} of inequalities (4) with $t = 2$ that are active at X_{ub} in the LP. Set $t_{\text{max}} = 2$.

while *there exists a violated inequality of form (4)*, **do**

```

    Solve the LP with time limit  $T$  to obtain a safe lower bound  $\bar{f}_{lb}$  and an optimal
    solution  $X_{lb}$ .
    if  $\bar{f}_{lb} > f_{lb}$ , then
        ⊥ Update  $f_{lb} = \bar{f}_{lb}$ 
    Run Algorithm round to round  $X_{lb}$  and get a partition matrix  $\bar{X}_{ub}$  with cost  $\bar{f}_{ub}$ .
    if  $\bar{f}_{ub} < f_{ub}$ , then
        ⊥ Update  $X_{ub} = \bar{X}_{ub}$  and  $f_{ub} = \bar{f}_{ub}$ 
    Update  $r_g = (f_{ub} - f_{lb})/f_{ub}$ 
    if  $r_g \leq \epsilon_{\text{opt}}$ , then
        ⊥ Terminate
    Remove from the LP, inequalities (4) that are not satisfied tightly at  $X_{lb}$ .
    Run Algorithm separate to obtain a number of inequalities of the form (4) with
     $t \leq t_{\text{max}}$  that are violated at  $X_{lb}$ . Add at most  $p_{\text{max}}$  of the most violated inequalities
    to the LP.
    if the number of violated inequalities returned by Algorithm separate is small, then
        ⊥ Update  $t_{\text{max}} = \min\{K, t_{\text{max}} + 1\}$ 

```

is on average 11.6 times faster than PDLP. *Again, for all 32 instances, the LP relaxation is tight; i.e., the optimality gap r_g upon termination is smaller than the optimality tolerance ϵ_{opt} .* On the contrary, the SDP relaxation is tight only for 5 instances. Moreover, the **k-means++**'s solution is optimal for 22 instances.

In Tables 3 and 4, all instances for which Problem (LPK_t) with $t = 2$ is not tight or is not solvable within 4 hours are marked. As we mentioned before, by letting $t > 2$, all such instances are solved to global optimality within the time limit, indicating the importance of selecting t in a dynamic fashion. Recall that by construction, if $K = 2$ or if an instance is solved in one iteration, then we have $t = 2$. For the remaining problems, in Table 5, we show the distribution of t associated with all inequalities added to the LP in the course of **Iterative LP solver**. As can be seen from this table, while for the majority of the inequalities we have $t \in \{2, 3\}$, for some instances adding inequalities with $t \in \{4, 5, 6\}$ is beneficial as well.

The termination criterion for **Iterative LP solver** is based on the optimality tolerance r_g . An alternative strategy is to terminate when the LP solution is a partition matrix. In this case, no upper bounding is needed. However, we need to solve each intermediate LP with high accuracy implying that we cannot rely on early stopping and safe lower bounds. The impact of this alternative strategy (labeled as “without upper bounding”) on the performance of **Iterative LP solver** is detailed

Table 3: Average run time and average number of iterations of the cutting plane algorithm with different LP solvers for small data sets. The numbers in parentheses denote the standard deviations across five random trials. No parentheses implies identical results for all trials.

Data set	n	K	cuPDLP-C		PDLP		Gurobi	
			time(s)	iteration	time(s)	iteration	time(s)	iteration
ruspini	75	4*	0.5	1.0	3.0	1.0	1.1	1.0
iris	150	2	5.1	1.0	18.8	1.0	20.7	1.0
		3	4.4	1.0	23.3	1.0	23.8	1.0
		4	3.2	1.0	22.9	1.0	15.8	1.0
wine	178	2	11.5	1.0	42.8	1.0	47.2	1.0
		7 ¹	4.3	1.0	23.3 (1.4)	1.0	23.1 (1.9)	1.0
gr202	202	6 ¹	34.1 (1.5)	1.0	82.5 (12.7)	1.0	232.2 (31.5)	1.0
seeds	210	2 ¹	17.6 (0.2)	1.0	110.0 (3.6)	1.0	140.6 (11.3)	1.0
		3	10.2	1.0	61.2	1.0	143.4	1.0
glass	214	3 [†]	119.3	3.0	426.7	2.0	421.9	1.0
		6 [†]	97.1 (21.2)	3.2 (0.5)	541.4 (78.7)	4.0 (1.4)	952.7 (47.0)	3.0
CatsDogs	328	2 ¹	72.3 (31.0)	1.0	780.7 (328.7)	1.2 (0.5)	1844.3 (153.6)	1.00
accent	329	2*	88.2 (6.4)	1.0	707.4 (52.0)	1.0	1617.1 (183.7)	1.0
		6 ¹	78.7 (19.5)	1.0	512.7 (36.5)	1.0	8443.7 (407.7)	9.2 (0.5)
ecoli	336	3	339.8 (267.8)	1.8 (0.8)	217.8 (19.2)	1.0	3202.2 (232.0)	1.0
RealEstate	414	3	73.2 (2.8)	1.0	641.7 (65.6)	1.0	> 14400	N/A
		5	130.3 (21.5)	1.0	1308.3 (348.6)	1.8 (0.5)	> 14400	N/A
Wholesale	440	5 ¹	141.1 (9.4)	1.0	717.7 (23.8)	1.0	> 14400	N/A
		6 [†]	1106.6 (276.3)	5.6 (2.1)	4854.6 (1627.0)	6.8 (0.8)	> 14400	N/A

¹ **k-means++**'s solution is not globally optimal for all five random trials.

[†] Problem (LPK_t) with $t = 2$ is not tight.

* The SDP relaxation, i.e., Problem (PW) is tight.

in Table 6. In this table, we include the data sets for which the LP solution in the first iteration of **Iterative LP solver** is not a partition matrix; as otherwise the two strategies coincide. As can be seen from this table, without upper bounding, **cuPDLP-C** solves only 9 out of 19 instances within the time limit. Moreover, for these 9 instances, the algorithm with upper bounding is on average 1.3 times faster than the one without upper bounding.

Finally, in Table 7 we list the average time spent for different components of **Iterative LP solver** with **cuPDLP-C** as the LP solver. As can be seen from this table, the vast majority of the time is spent for solving the LPs. The initialization time is particularly big for large scale problems in which case we need to select a random subset of inequalities that are binding at the **k-means++** solution. Other major components such as separation time and rounding time are rather insignificant.

We conclude this paper by acknowledging that further work is needed to make **Iterative LP solver** relevant for larger data sets. Of course, as first-order LP solvers become faster and more powerful GPUs become available, one can solve larger instances. Exploring sketching techniques [23, 2] for tackling larger data sets is an interesting topic of future research as well. An alternative strategy is to incorporate the proposed LP relaxation within mixed-integer nonlinear programming solvers [14] to expedite their convergence.

Table 4: Average run time and average number of iterations of the cutting plane algorithm with PDLP and cuPDLP-C for large data sets. The numbers in parentheses denote the standard deviations across five random trials. No parentheses implies identical results for all trials.

Data set	n	K	cuPDLP-C		PDLP	
			time(s)	iteration	time(s)	iteration
ECG5000	500	2	120.7 (15.7)	1.0	484.3 (43.6)	1.0
		5 [†]	581.7 (93.8)	3.2 (0.5)	3727.6 (1115.2)	5.4 (1.1)
Hungarian	522	2	96.9 (11.3)	1.0	599.8 (109.4)	1.0
		Wdbc	569	88.9 (12.2)	1.0	218.8 (457.7)
Control	600	5 ¹	23.4 (20.1)	1.0	1448.6 (437.7)	1.8 (0.5)
		3	93.9 (7.0)	1.0	532.3 (41.6)	1.0
Heartbeat	606	2 [*]	158.1 (39.6)	1.0	712.8 (31.4)	1.0
Strawberry	613	2	94.7 (5.5)	1.0	1039.7 (525.4)	1.4 (0.6)
Energy	768	2 [*]	50.3 (0.4)	1.0	169.5 (13.2)	1.0
		12 [†]	158.5 (91.2)	1.8 (1.1)	869.5 (514.7)	2.8 (2.1)
Gene	801	5 ¹	211.2 (104.3)	1.2 (0.5)	1104.7 (414.6)	1.4 (0.6)
		6 ¹	707.4 (111.2)	3.0	974.1 (340.4)	1.2 (0.5)
SalesWeekly	811	2 ¹	70.1 (3.9)	1.0	1620.7 (118.1)	2.0
		3	65.3 (2.1)	1.0	565.1 (50.10)	1.0
		5	89.9 (2.7)	1.0	1417.4 (433.1)	1.8 (0.5)
Vehicle	846	2	61.3 (1.5)	1.0	866.8 (326.1)	1.2 (0.5)
Arcene	900	2	115.5 (5.7)	1.0	273.6 (15.0)	1.0
		3 [*]	74.6 (3.4)	1.0	292.1 (8.1)	1.0
		5 [†]	759.7 (78.7)	3.8 (0.5)	> 14400	N/A
Wafer	1000	2	67.4 (1.5)	1.0	316.9 (32.1)	1.0
		4 [†]	643.1 (76.3)	3.6 (0.6)	4602.1 (1101.9)	5.2 (1.1)
Power	1096	2	95.9 (19.9)	1.0	1543.2 (508.9)	2.2 (0.8)
Phising	1353	9 [†]	840.5 (47.2)	20.4 (1.8)	> 14400	N/A
Aspirin	1500	3	230.8 (119.1)	1.4 (0.6)	1651.6 (894.8)	2.4 (1.5)
Car	1727	4	1504.6 (198.9)	17.4 (1.1)	10587.8 (1872.9)	18.8 (1.6)
Ethanol	2000	2 [*]	210.2 (4.1)	1.0	10330.9 (2217.6)	7.0 (0.7)
Wifi	2000	5 [†]	3309.2 (702.6)	8.0 (1.6)	> 14400	N/A
Mallat	2400	3 [‡]	5934.9 (1110.8)	10.6 (1.5)	> 14400	N/A
		4 [‡]	7820.3 (1941.1)	15.2 (2.2)	> 14400	N/A
Advertising	3279	2 [‡]	4318.1 (622.6)	8.4 (2.1)	> 14400	N/A
		8 [‡]	7410.7 (2367.0)	12.0 (2.9)	> 14400	N/A
Rice	3810	2 [‡]	8314.7 (669.2)	11.8 (0.5)	> 14400	N/A

¹ **k-means++**'s solution is not globally optimal for all five random trials.

[†] Problem LPK_t with $t = 2$ is not tight.

[‡] Unable to verify whether Problem (LPK_t) with $t = 2$ is tight or not due to exceeding the time limit.

^{*} The SDP relaxation, i.e., Problem (PW) is tight.

Table 5: Distribution of different types of inequalities (4) (in percentage) added to the LP relaxation during cut generation with cuPDLP-C as the LP solver. Inequalities (4) with $t = 2$ added during the initialization step are not included.

Data set	n	K	$t = 2$	$t = 3$	$t = 4$	$t = 5$	$t = 6$
glass	214	3	13.2	86.8	0.0	0.0	0.0
		6	67.5	32.5	0.0	0.0	0.0
ecoli	336	3	80.6	19.4	0.0	0.0	0.0
Wholesale	440	6	51.0	41.5	6.2	1.2	0.1
ECG5000	500	5	53.7	46.3	0.0	0.0	0.0
Energy	768	12	100.0	0.0	0.0	0.0	0.0
		5	100.0	0.0	0.0	0.0	0.0
Gene	801	6	43.5	56.5	0.0	0.0	0.0
		5	37.4	62.6	0.0	0.0	0.0
Arcene	900	5	37.4	62.6	0.0	0.0	0.0
Wafer	1000	4	46.5	53.5	0.0	0.0	0.0
Phising	1353	9	90.5	9.5	0.0	0.0	0.0
Aspirin	1500	3	100.0	0.0	0.0	0.0	0.0
Car	1727	4	77.9	17.7	4.4	0.0	0.0
Wifi	2000	5	37.5	49.2	10.8	2.5	0.0
Mallat	2400	3	32.2	67.8	0.0	0.0	0.0
		4	37.2	40.3	22.5	0.0	0.0
Advertising	3279	8	84.9	12.1	2.9	0.1	0.0

Table 6: The impact of upper rounding on the performance of the cutting plane algorithm, in terms of average time and average number of iterations, with cuPDLP-C as the LP solver. The numbers in parentheses denote the standard deviations across five random trials. No parentheses implies identical results for all trials.

Data set	n	K	with upper bounding		without upper bounding	
			time(s)	iteration	time(s)	iteration
glass	214	3	119.3	3.0	119.3	3.0
		6	97.1 (21.2)	3.2 (0.5)	129.8 (39.4)	3.6 (0.9)
ecoli	336	3	339.8 (267.8)	1.8 (0.8)	205.5 (167.2)	1.0
Wholesale	440	6	1106.6 (276.3)	5.6 (2.1)	2722.8 (1535.3)	3.80 (0.5)
ECG5000	500	5	581.7 (93.8)	3.2(0.5)	> 14400	N/A
Energy	768	12	158.5 (91.2)	1.8 (1.1)	250.1 (206.3)	1.6 (0.6)
		5	211.2 (104.3)	1.2 (0.5)	239.8 (169.8)	1.0
Gene	801	6	707.4 (111.2)	3.0	676.6 (194.2)	1.2 (0.5)
		5	759.7 (78.7)	3.8 (0.5)	1423.6 (165.9)	3.2 (0.5)
Wafer	1000	4	643.1 (76.3)	3.6 (0.6)	679.0 (119.9)	3.0
Phising	1353	9	840.5 (47.2)	20.4 (1.8)	> 14400	N/A
Aspirin	1500	3	230.8 (119.1)	1.4 (0.6)	> 14400	N/A
Car	1727	4	1504.6 (198.9)	17.4 (1.1)	> 14400	N/A
Wifi	2000	5	3309.2 (702.6)	8.0 (1.6)	> 14400	N/A
Mallat	2400	3	5934.9 (1110.8)	10.6 (1.5)	> 14400	N/A
		4	7820.3 (1941.1)	15.2 (2.2)	> 14400	N/A
Advertising	3279	2	4318.1 (622.6)	8.4 (2.1)	> 14400	N/A
		8	7410.7 (2367.0)	12.0 (2.9)	> 14400	N/A
Rice	3810	2	8314.7 (669.2)	11.8 (0.5)	> 14400	N/A

Table 7: Average percentage of time consumed by different components of the cutting plane algorithm with cuPDLP-C as the LP solver. These components are: LP solver time (solver), initialization time (init), separation time (sep), rounding time (round), and other tasks such as model building (others). The numbers in parentheses denote the standard deviations across five random trials. No parentheses implies identical results for all trials.

Data set	n	K	solver	init	sep	round	others
ruspini	75	4	89.1	8.2	0.0	0.0	2.7
iris	150	2	84.5	12.7	0.0	0.1	2.7
		3	86.8	10.7	0.0	0.1	2.4
		4	85.4	11.8	0.0	0.2	2.6
wine	178	2	88.3	9.5	0.0	0.1	2.1
		7	86.9	10.6	0.0	0.2	2.3
gr202	202	6	96.9 (0.1)	2.1 (0.5)	0.0	0.0	0.9 (0.6)
seeds	210	2	87.9 (0.2)	9.9 (0.2)	0.0	0.1	2.1
		3	84.6	12.6	0.0	0.1	2.7
glass	214	3	93.0	1.6	0.0	0.0	5.4
		6	93.3 (0.9)	1.2 (0.2)	0.0	0.0	5.5 (0.7)
CatsDogs	328	2	81.7 (5.0)	15.8 (4.3)	0.0	0.2 (0.1)	2.3 (0.6)
accent	329	2	90.7 (0.7)	8.0 (0.6)	0.0	0.1	1.2 (0.1)
		6	94.8 (1.3)	4.2 (1.1)	0.0	0.1	0.9 (0.2)
ecoli	336	3	93.5 (3.4)	3.9 (4.0)	0.0	0.1	2.5 (0.9)
RealEstate	414	3	87.0 (0.5)	10.8 (0.4)	0.0	0.2	2.0 (0.1)
		5	92.9 (1.0)	5.9 (0.8)	0.0	0.1	1.1 (0.1)
Wholesale	440	5	93.9 (0.4)	5.1 (0.3)	0.0	0.1	1.0 (0.1)
		6	94.4 (0.3)	0.7 (0.1)	0.0	0.1	4.9 (0.4)
ECG5000	500	2	91.0 (1.1)	7.5 (0.9)	0.0	0.2	1.3 (0.2)
		5	94.1 (0.4)	1.2 (0.2)	0.0	0.1	4.6 (0.3)
Hungarian	522	2	88.5 (1.3)	9.6 (1.1)	0.0	0.3	1.6 (0.2)
Wdbc	569	2	86.2 (1.7)	11.5 (1.3)	0.0	0.4 (0.1)	1.9 (0.2)
		5	91.1 (1.4)	7.4 (1.1)	0.0	0.3 (0.1)	1.2 (0.2)
Control	600	3	88.3 (0.8)	9.7 (0.7)	0.0	0.3	1.7 (0.1)
Heartbeat	606	2	90.7 (2.4)	7.9 (2.0)	0.0	0.3 (0.1)	1.1 (0.3)
Strawberry	613	2	86.4 (0.8)	11.4 (0.7)	0.0	0.4	1.8 (0.1)
Energy	768	2	67.8 (0.3)	27.0 (0.3)	0.0	1.8	3.4 (0.1)
		12	89.6 (3.6)	6.1 (4.6)	0.1 (0.1)	1.0 (0.3)	3.1 (1.4)
Gene	801	5	67.8 (8.0)	29.3 (9.4)	0.0	1.2 (0.1)	1.7 (1.6)
		6	84.1 (2.3)	10.0 (1.4)	0.1	0.9 (0.1)	4.9 (0.7)
SalesWeekly	811	2	74.9 (1.4)	21.1 (1.2)	0.0	1.4 (0.1)	2.5 (0.1)
		3	74.5 (0.8)	21.3 (0.7)	0.0	1.5 (0.1)	2.7 (0.1)
		5	83.3 (0.5)	13.7 (0.4)	0.0	1.1 (0.1)	1.9 (0.1)
Vehicle	846	2	69.7 (1.0)	25.6 (0.8)	0.0	1.8 (0.2)	2.9
Arcene	900	2	75.0 (1.4)	21.9 (1.3)	0.0	1.5 (0.1)	1.6 (0.1)
		3	60.2 (1.2)	35.1 (1.0)	0.0	2.3 (0.2)	2.4 (0.1)
		5	86.8 (1.1)	5.4 (0.7)	0.2	0.9 (0.1)	6.6 (0.7)
Wafer	1000	2	62.6 (0.9)	31.8 (0.7)	0.0	2.7 (0.2)	2.9 (0.1)
		4	88.8 (1.4)	3.0 (0.4)	0.4 (0.1)	1.0 (0.2)	6.8 (1.0)
Power	1096	2	67.0 (6.5)	28.1 (5.4)	0.0	2.7 (0.6)	2.2 (0.5)
Phising	1353	9	74.5 (1.8)	3.2 (0.2)	6.2 (0.6)	10.7 (0.8)	5.3 (0.4)
Aspirin	1500	3	59.7 (11.8)	32.9 (14.4)	0.4 (0.5)	3.4 (0.4)	3.6 (2.6)
Car	1727	4	70.1 (2.6)	4.4 (0.7)	7.4 (0.3)	11.1 (1.2)	7.0 (0.4)
Ethanol	2000	2	21.4 (0.5)	70.8 (0.8)	0.0	6.2 (0.3)	1.6
Wifi	2000	5	84.6 (1.0)	3.1 (0.8)	3.4 (0.4)	3.0 (0.2)	5.9 (0.4)
Mallat	2400	3	83.2 (1.4)	3.9 (0.6)	4.0 (0.3)	4.5 (0.4)	4.4 (0.4)
		4	82.0 (1.7)	2.9 (0.6)	4.9 (0.3)	4.7 (0.4)	5.5 (0.6)
Advertising	3279	2	64.4 (1.9)	18.5 (3.1)	8.5 (1.0)	2.4 (0.6)	6.1 (1.1)
		8	68.6 (3.0)	11.3 (4.1)	9.0 (0.9)	4.3 (1.1)	6.8 (1.4)
Rice	3810	2	68.2 (1.6)	12.6 (0.8)	10.2 (0.5)	3.5 (0.5)	5.4 (0.2)

References

- [1] *MOSEK 9.2*, 2019. <http://docs.mosek.com/9.0/faq.pdf>.
- [2] P. Abdalla and A. S. Bandeira. Community detection with a subsampled semidefinite program. *Sampling Theory, Signal Processing, and Data Analysis*, 20(1):6, 2022.
- [3] D. Aloise, A. Deshpande, P. Hansen, and P. Popat. NP-hardness of Euclidean sum-of-squares clustering. *Machine Learning*, 75:245–248, 2009.
- [4] D. Applegate, M. Díaz, O. Hinder, H. Lu, M. Lubin, B. O’Donoghue, and W. Schudy. Practical large-scale linear programming using primal-dual hybrid gradient. *Advances in Neural Information Processing Systems*, 34:20243–20257, 2021.
- [5] P. Awasthi, A. S. Bandeira, M. Charikar, R. Krishnaswamy, S. Villar, and R. Ward. Relax, no need to round: Integrality of clustering formulations. *Proceedings of the 2015 Conference on Innovations in Theoretical Computer Science*, 165:191–200, 2015.
- [6] J. Czyzyk, M. P. Mesnier, and J. J. More. The NEOS server. *IEEE Journal on Computational Science and Engineering*, 5(3):68–75, 1998.
- [7] L. Dagum and R. Menon. OpenMP: an industry standard api for shared-memory programming. *IEEE computational science and engineering*, 5(1):46–55, 1998.
- [8] A. De Rosa and A. Khajavirad. The ratio-cut polytope and K-means clustering. *SIAM Journal on Optimization*, 32(1):173–203, 2022.
- [9] D. Dua, C. Graff, et al. UCI machine learning repository. URL <http://archive.ics.uci.edu/ml>, 2017.
- [10] Z. Friggstad, M. Rezapour, and Salavatipour M. R. Local search yields a PTAS for K-means in doubling metrics. *SIAM Journal on Computing*, 48(2):452–480, 2019.
- [11] Gurobi Optimization, LLC. Gurobi Optimizer Reference Manual, 2021. URL: <https://www.gurobi.com>.
- [12] T. Iguchi, D. G. Mixon, J. Peterson, and S. Villar. Probably certifiably correct K-means clustering. *Mathematical Programming*, 165:605–642, 2017.
- [13] T. Kanungo, D.M. Mount, N.S. Netanyahu, C.D. Piatko, R. Silverman, and A.Y. Wu. A local search approximation algorithm for K-means clustering. *Proceedings of the 18th Annual ACM Symposium on Computational Geometry*, pages 10–18, 2002.
- [14] A. Khajavirad and N. V. Sahinidis. A hybrid LP/NLP paradigm for global optimization relaxations. *Mathematical Programming Computation*, 10:383–421, 2018. doi:10.1007/s12532-018-0138-5.
- [15] X. Li, Y. Li, S. Ling, T. Strohmer, and K. Wei. When do birds of a feather flock together? K-means, proximity, and conic programming. *Mathematical Programming*, 179:295–341, 2020.
- [16] S. Lloyd. Least squares quantization in PCM. *IEEE Transactions on Information Theory*, 28(2):129–137, 1982.
- [17] H. Lu and J. Yang. cuPDLP.jl: A GPU implementation of restarted primal-dual hybrid gradient for linear programming in julia. *arXiv preprint arXiv:2311.12180*, 2023.
- [18] H. Lu, J. Yang, H. Hu, Q. Huangfu, J. Liu, T. Liu, Y. Ye, C. Zhang, and D. Ge. cuPDLP-C: A strengthened implementation of cupdlp for linear programming by c language. *arXiv preprint arXiv:2312.14832*, 2023.
- [19] M. Mahajan, P. Nimbhorkar, and K. Varadarajan. The planar K-means problem is NP-hard. In *WALCOM: Algorithms and Computation*, pages 274–285. Springer Berlin Heidelberg, 2009.
- [20] O.L. Mangasarian. Uniqueness of solution in linear programming. *Linear Algebra and its Applications*, 25:151–162, 1979.
- [21] F. Marzi, F. Rossi, and S. Smriglio. Computational study of separation algorithms for clique inequalities. *Soft Computing*, 23(9):3013–3027, 2019.
- [22] D. G. Mixon, S. Villar, and R. Ward. Clustering subgaussian mixtures by semidefinite programming. *Information and Inference: A Journal of the IMA*, 6(4):389–415, 2017.
- [23] D. G. Mixon and K. Xie. Sketching semidefinite programs for faster clustering. *IEEE Transactions on Information Theory*, 67(10):6832–6840, 2021.
- [24] A. Neumaier and O. Shcherbina. Safe bounds in linear and mixed-integer linear programming. *Mathematical Programming*, 99:283–296, 2004.

- [25] J Peng and Y Wei. Approximating K-means-type clustering via semidefinite programming. *SIAM Journal on Optimization*, 18(1):186–205, 2007.
- [26] J. Peng and Y. Xia. *A New Theoretical Framework for K-Means-Type Clustering*, pages 79–96. Springer Berlin Heidelberg, 2005.
- [27] V. Piccialli, A. M. Sudoso, and A. Wiegele. SOS-SDP: An exact solver for minimum sum-of-squares clustering. *INFORMS Journal on Computing*, 34(4):2144–2162, 2022.
- [28] M. N. Prasad and G. A. Hanasusanto. Improved conic reformulations for K-means clustering. *SIAM Journal on Optimization*, 28(4):3105–3126, 2018.
- [29] R. Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge University Press, 2018.