Enumeration of linear codes with different hulls

Iliya Bouyukliev[∗] and Stefka Bouyuklieva†

Abstract

The hull of a linear code C is the intersection of C with its dual code. We present and analyze the number of linear q -ary codes of the same length and dimension but with different dimensions for their hulls. We prove that for given dimension k and length $n \geq 2k$ the number of all $[n, k]_q$ linear codes with hull dimension l decreases as l increases. We also present classification results for binary and ternary linear codes with trivial hulls (LCD and self-orthogonal) for some values of the length n and dimension k , comparing the obtained numbers with the number of all linear codes for the given n and k .

1 Introduction

Let \mathbb{F}_q be a finite field with q elements and \mathbb{F}_q^n be the *n*-dimensional vector space over \mathbb{F}_q . Suppose that \mathbb{F}_q^n is equipped with an inner product. Any linear subspace C of \mathbb{F}_q^n is called a linear q-ary code of length n, and its orthogonal complement C^{\perp} (with respect to the defined inner product) is its dual (or orthogonal) code.

The hull of a linear code C was introduced by Assmus, Jr. and Key $[4]$, and it is the intersection of C with its dual code C^{\perp} . The study of hulls also has applications ranging from the construction of quantum error correcting codes to post-quantum cryptography [\[16\]](#page-14-0). If the hull contains only the zero vector, C is called a linear complementary dual (LCD) code. LCD codes were introduced by Massey [\[17\]](#page-14-1) and gave an optimum linear coding solution for the two user binary adder channel. The determination of the largest minimum weight among all LCD $[n, k]$ codes and the classification of all or only optimal LCD $[n, k]$ codes are fundamental problems in coding theory. Recently, much work has been done concerning this fundamental problem (see [\[7,](#page-11-1) [10,](#page-14-2) [11,](#page-14-3) [14\]](#page-14-4)). It has been shown in [\[7\]](#page-11-1) that if the considered inner product is Euclidean and $q > 3$, any linear code over \mathbb{F}_q is equivalent to an (Euclidean) LCD code, and if the inner product is Hermitian and $q > 4$, any q-ary linear code is equivalent to a Hermitian LCD code. Note that a Hermitian inner product is considered only if q is an even power of the characteristic of the field.

The other extremal case is when the hull of a linear code C coincides with the code itself. Such a code is called self-orthogonal (SO) and it is a subset of its dual code. If C coincides with C^{\perp} , the code is self-dual. Self-dual codes have been a topic of great interest since the 1960s, with connections and applications to various branches of mathematics and coding theory.

[∗] I. Bouyukliev is with the Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, P.O. Box 323, Veliko Tarnovo, BULGARIA. Email: iliyab@math.bas.bg

[†]S. Bouyuklieva is with the Faculty of Mathematics and Informatics, St. Cyril and St. Methodius University of Veliko Tarnovo, BULGARIA. Email: stefka@ts.uni-vt.bg. The research of this author was supported, in part, by a Bulgarian NSF contract KP-06-N32/2-2019.

Theoretically, self-dual codes are closely connected to other mathematical structures such as designs, lattices, graphs and modular forms [\[9\]](#page-14-5). Practically, many of the best-known codes are actually self-dual codes.

Self-dual and LCD codes are widely studied in the literature not only for their mathematical attraction, but also for their applications. In this study, we are also interested in linear codes that lie between these two extremal cases. If $h(C)$ is the dimension of the hull of the linear code C , then

$$
0 \le h(C) \le \min\{\dim C, \dim C^{\perp}\}.
$$

Sendrier in [\[20\]](#page-14-6) gave a formula for the number of linear q -ary codes of a given length n, dimension k and hull dimension l. He proved that, asymptotically, the proportion of q -ary linear codes whose hull has dimension l to all q -ary linear codes of the same length and dimension, only depends on l and q . It turns out that for a fixed length and dimension and for LCD codes $(l = 0)$, this proportion is approximately $1 - 1/q$. Hence most linear codes have hull dimension 0 (i.e. LCD) when q is large.

By $A_{n,k,l}$ we denote the number of the linear $[n, k]_q$ codes whose hull has dimension l (for fixed q). The main result of the present study is contained in the following theorem.

Theorem 1. For fixed positive integers k and $n \geq 2k$, the following inequalities hold:

$$
A_{n,k,0} > A_{n,k,1} > \cdots > A_{n,k,k}.\tag{1}
$$

Note that $A_{n,k,0}$ and $A_{n,k,k}$ are the numbers of all LCD and SO q-ary codes of length n and dimension k , respectively. To prove the main theorem, we use some known formulae which we present in Section 2. Section 3 is devoted to Theorem [1](#page-1-0) and its proof. Note that $A_{n,k,l}$ is the number of all different codes with the corresponding parameters, but the number of the inequivalent among them is much smaller. Since we do not have a formula for this number, we did some experiments and classified the binary and ternary linear $[n, k]$ codes for some values of n and k. We present and analyze the obtained results in Section 4.

2 Preliminaries

In this section, we present the basic definitions, theorems and formulas that we need for the proof of the main theorem, as well as for the computational results in Section [4.](#page-9-0)

This study only applies to linear codes over a finite field. A linear $[n, k, d]$ q-ary code C is a k-dimensional subspace of the vector space \mathbb{F}_q^n , and d is the smallest weight among all non-zero codewords of C , called the *minimum weight* (or minimum distance) of the code. A code is called even if all its codewords have even weights. Let $(u, v) : \mathbb{F}_q^n \times \mathbb{F}_q^n \to \mathbb{F}_q$ be an inner product in \mathbb{F}_q^n . The orthogonal complement of C according to the defined inner product is called the *dual* code of C and denoted by C^{\perp} , i.e. $C^{\perp} = \{u \in \mathbb{F}_q^n : (u, v) = 0 \text{ for all } v \in C\}$. Obviously, C^{\perp} is a linear $[n, n-k]$ code. The intersection $C \cap C^{\perp}$ is called the hull of the code and denoted by $\mathcal{H}(C)$. As we mentioned in Introduction, the dimension $h(C)$ of the hull can be at least 0 and at most k, as $h(C) = k$ if and only if the code is self-orthogonal, and $h(C) = 0$ if and only if C is an LCD code.

Two linear q -ary codes C_1 and C_2 are equivalent if the codewords of C_2 can be obtained from the codewords of C_1 via a sequence of transformations of the following types:

(1) permutation on the set of coordinate positions;

- (2) multiplication of the elements in a given position by a non-zero element of \mathbb{F}_q ;
- (3) application of a field automorphism to the elements in all coordinate positions.

An automorphism of a linear code C is a sequence of the transformations $(1) - (3)$ which maps each codeword of C onto a codeword of the same code. The set of all automorphisms of C forms a group called the automorphism group $Aut(C)$ of the code. The presented formulas do not count the equivalence but in Section [4](#page-9-0) we enumerate only inequivalent codes.

If the considered inner product is Euclidean, i.e. $(x, y) = \sum_{i=1}^{n} x_i y_i$ for any two vectors $x=(x_1,\ldots,x_n), y=(y_1,\ldots,y_n)$ from \mathbb{F}_q^n , the dimension of the hull is an invariant of equivalent codes in the cases $q = 2$ and $q = 3$, but if $q \ge 4$, any linear code over \mathbb{F}_q is equivalent to an Euclidean LCD code [\[7\]](#page-11-1). If $q = p^{2s}$, where p is the characteristic of the field, we can consider the Hermitian inner product over \mathbb{F}_q defined by

$$
(x,y) = \sum_{i=1}^{n} x_i y_i^{\sqrt{q}}, \ \forall x = (x_1,\ldots,x_n), y = (y_1,\ldots,y_n) \in \mathbb{F}_q^n.
$$

For this inner product, the dimension of the hull is an invariant in the case of the quaternary codes $(q = 4)$, and if $q > 4$, any linear code over \mathbb{F}_q , $q = p^{2s}$, is equivalent to an Hermitian LCD code [\[7\]](#page-11-1).

Formulae for counting the number $\sigma(n, k)$ of all different q-ary self-orthogonal codes of length n and dimension k were proven in [\[19\]](#page-14-7).

Theorem 2. Let $m = \lfloor n/2 \rfloor$ and $\pi_{n,k} = \prod_{i=1}^{k}$ $\frac{q^{2m-2i+2}-1}{q^i-1}$. For all $1 \leq k \leq m$, we have

$$
\sigma_{n,k} = \begin{cases}\n\pi_{n,k}, & \text{if } n \text{ is odd,} \\
\frac{q^{n-k}-1}{q^{n-1}} \pi_{n,k}, & \text{if } n \text{ and } q \text{ are even,} \\
\frac{q^{m-k}-1}{q^{m-k}-1} \pi_{n,k}, & \text{if } n \equiv 2 \pmod{4} \text{ and } q \equiv 3 \pmod{4}, \\
\frac{q^{m-k}+1}{q^{m+1}} \pi_{n,k}, & \text{if } (n \equiv 0 \pmod{4}) \\
& \text{or } (n \equiv 2 \pmod{4}) \text{ and } q \equiv 1 \pmod{4}).\n\end{cases}
$$

Note that $\sigma_{n,0} = 1$. We need also the following corollary.

Corollary 3. Let $m = |n/2|$ and $0 \le k \le m - 1$. Then

$$
\frac{\sigma_{n,k}}{\sigma_{n,k+1}} = \tau_{n,k} \frac{q^{k+1} - 1}{q^{2m-2k} - 1},\tag{2}
$$

where

$$
\tau_{n,k} = \begin{cases}\n1, & \text{if } n \text{ is odd,} \\
\frac{q^{n-k}-1}{q^{n-k-1}-1}, & \text{if } n \text{ and } q \text{ are even,} \\
\frac{q^{m-k}+\epsilon}{q^{m-k-1}+\epsilon}, & \text{if } n \text{ is even and } q \text{ is odd,}\n\end{cases}
$$
\n(3)

with $\epsilon = -1$ if $n \equiv 2 \pmod{4}$ and $q \equiv 3 \pmod{4}$, and $\epsilon = 1$ otherwise. *Proof.* The formula [\(2\)](#page-2-0) follows directly from Theorem [2.](#page-2-1)

The following theorem is the main tool in our investigations.

Theorem 4. [\[20\]](#page-14-6) Let k and $n \geq 2k$ be positive integers. If $\sigma_{n,i}$ is the number of all selforthogonal $[n, i]_q$ codes, $i \leq k$, then the number of all $[n, k]_q$ codes whose hull has dimension l, $l < k$, is equal to

$$
A_{n,k,l} = \sum_{i=l}^{k} \begin{bmatrix} n-2i \\ k-i \end{bmatrix} \begin{bmatrix} i \\ l \end{bmatrix} (-1)^{i-l} q^{\binom{i-l}{2}} \sigma_{n,i}.
$$
 (4)

In the above formula, $\begin{bmatrix} n \\ k \end{bmatrix}$ $\binom{n}{k}$ is the Gaussian binomial coefficient, defined by

$$
\begin{bmatrix} n \\ 0 \end{bmatrix} = 1, \quad \begin{bmatrix} n \\ k \end{bmatrix} = \frac{(q^n - 1)(q^{n-1} - 1) \dots (q^{n-k+1} - 1)}{(q^k - 1)(q^{k-1} - 1) \dots (q - 1)}, \text{if } 1 \le k \le n. \tag{5}
$$

We use the following properties of the Gaussian coefficients for $0 \le k \le n - 1$:

$$
\begin{bmatrix} n \\ n-k \end{bmatrix} = \begin{bmatrix} n \\ k \end{bmatrix}, \tag{6}
$$

$$
\begin{bmatrix} n+1 \\ k \end{bmatrix} = \frac{q^{n+1}-1}{q^{n-k+1}-1} \begin{bmatrix} n \\ k \end{bmatrix},\tag{7}
$$

$$
\begin{bmatrix} n \\ k+1 \end{bmatrix} = \frac{q^{n-k}-1}{q^{k+1}-1} \begin{bmatrix} n \\ k \end{bmatrix}, \tag{8}
$$

$$
\begin{bmatrix} n-2i \\ k-i \end{bmatrix} = \frac{(q^{n-2i}-1)(q^{n-2i-1}-1)}{(q^{n-k-i}-1)(q^{k-i}-1)} \begin{bmatrix} n-2i-2 \\ k-i-1 \end{bmatrix}.
$$
 (9)

Remark 1. In [\[8\]](#page-14-8), the authors give more explicit formulas for the number of different LCD codes with a given length and dimension in the cases when $q = 2$ and $q = p^s$ for an odd prime p.

3 Number of linear codes whose hull has dimension l for different values of l

For the calculations, we need to evaluate a few constants.

Lemma 5. If $n = 2m$ and $k \leq m$ are positive integers, then

$$
\tau_{n,k} \ge \begin{cases} q, & \text{if } q \text{ is even, or } (n \equiv 2 \pmod{4} \text{ and } q \equiv 3 \pmod{4}) \\ (q+1)/2, & \text{otherwise.} \end{cases}
$$

Proof. We use the formula [\(3\)](#page-2-2):

$$
\frac{q^{n-k}-1}{q^{n-k-1}-1} = q + \frac{q-1}{q^{n-k-1}-1} > q, \quad \frac{q^{m-k}-1}{q^{m-k-1}-1} = q + \frac{q-1}{q^{m-k-1}-1} > q,
$$

$$
\frac{q^{m-i}+1}{q^{m-i-1}+1} = q - \frac{q-1}{q^{m-i-1}+1} \ge q - \frac{q-1}{2} = \frac{q+1}{2}.
$$

This proves the lemma in all cases. \Box

Lemma 6. Let $m = \lfloor n/2 \rfloor$, $0 \le i < k \le m$, and

$$
\lambda_{n,i} = \frac{(q^{n-2i} - 1)(q^{n-2i-1} - 1)}{(q^{2m-2i} - 1)(q^{n-k-i} - 1)(q^{k-i} - 1)} = \begin{cases} \frac{q^{n-2i} - 1}{(q^{n-k-i} - 1)(q^{k-i} - 1)} & \text{if } n = 2m + 1, \\ \frac{q^{n-2i-1} - 1}{(q^{n-k-i} - 1)(q^{k-i} - 1)} & \text{if } n = 2m, \end{cases}
$$

Then $q^{2m+1-n}\lambda_{n,i} > 1 + \frac{1}{q^{k-i}-1} > 1$.

Proof. We have that

$$
\frac{q^{n-2i}-1}{(q^{n-k-i}-1)(q^{k-i}-1)} = 1 + \frac{1}{q^{n-k-i}-1} + \frac{1}{q^{k-i}-1} > 1
$$

and

$$
\frac{q^{n-2i} - q}{(q^{n-k-i} - 1)(q^{k-i} - 1)} = 1 + \frac{1}{q^{n-k-i} - 1} + \frac{1}{q^{k-i} - 1} - \frac{q-1}{(q^{n-k-i} - 1)(q^{k-i} - 1)}
$$

\n
$$
\geq 1 + \frac{1}{q^{n-k-i} - 1} + \frac{1}{q^{k-i} - 1} - \frac{q^{k-i} - 1}{(q^{n-k-i} - 1)(q^{k-i} - 1)}
$$

\n
$$
= 1 + \frac{1}{q^{k-i} - 1} > 1,
$$

therefore $\lambda_{n,i} > 1$ if $n = 2m + 1$, and $q\lambda_{n,i} > 1$ if $n = 2m$.

We use the constant $\lambda_{n,i}$ in the following formula

$$
\begin{bmatrix} n-2i \ k-i \end{bmatrix} = (q^{2m-2i}-1)\lambda_{n,i} \begin{bmatrix} n-2i-2 \ k-i-1 \end{bmatrix}.
$$
 (10)

Consider the difference $A_{n,k,l}-A_{n,k,l+1}$ for $0\leq l\leq k-1.$

$$
A_{n,k,l} - A_{n,k,l+1} =
$$
\n
$$
= \sum_{i=l}^{k} \begin{bmatrix} n-2i \\ k-i \end{bmatrix} \begin{bmatrix} i \\ l \end{bmatrix} (-1)^{i-l} q^{\binom{i-l}{2}} \sigma_{n,i} - \sum_{i=l+1}^{k} \begin{bmatrix} n-2i \\ k-i \end{bmatrix} \begin{bmatrix} i \\ l+1 \end{bmatrix} (-1)^{i-l-1} q^{\binom{i-l-1}{2}} \sigma_{n,i}
$$
\n
$$
= \begin{bmatrix} n-2l \\ k-l \end{bmatrix} \sigma_{n,l} + \sum_{i=l+1}^{k} (-1)^{i-l} \begin{bmatrix} n-2i \\ k-i \end{bmatrix} \left(\begin{bmatrix} i \\ l \end{bmatrix} q^{\binom{i-l}{2}} + \begin{bmatrix} i \\ l+1 \end{bmatrix} q^{\binom{i-l-1}{2}} \right) \sigma_{n,i}
$$
\n
$$
= \begin{bmatrix} n-2l \\ k-l \end{bmatrix} \sigma_{n,l} + \sum_{i=l+1}^{k} (-1)^{i-l} a_{n,k,l,i},
$$

where

$$
a_{n,k,l,i} = \begin{bmatrix} n-2i \\ k-i \end{bmatrix} \left(\begin{bmatrix} i \\ l \end{bmatrix} q^{i-l} + \begin{bmatrix} i \\ l+1 \end{bmatrix} q^{i-l-1} \right) \sigma_{n,i}
$$

\n
$$
= \begin{bmatrix} n-2i \\ k-i \end{bmatrix} \begin{bmatrix} i \\ l \end{bmatrix} (q^{i-l-1} + \frac{q^{i-l}-1}{q^{l+1}-1}) q^{i-l-1} \sigma_{n,i}
$$

\n
$$
= \begin{bmatrix} n-2i \\ k-i \end{bmatrix} \begin{bmatrix} i \\ l \end{bmatrix} \frac{q^{i} - q^{i-l-1} + q^{i-l} - 1}{q^{l+1}-1} q^{i-l-1} \sigma_{n,i},
$$

\n
$$
= \begin{bmatrix} n-2i \\ k-i \end{bmatrix} \begin{bmatrix} i \\ l \end{bmatrix} \frac{\alpha_{l,i}}{q^{l+1}-1} q^{i-l-1} \sigma_{n,i}, \quad l+1 \leq i \leq k,
$$

$$
\alpha_{l,i} = q^{i-l-1}(q^{l+1} + q - 1) - 1 \ge q^{l+1} + q - 2 \ge 2q - 2.
$$

We take $a_{n,k,l,l} = \begin{bmatrix} n-2l \\ k-l \end{bmatrix}$ $\sum_{k-l}^{i-2l} \sigma_{n,l}$. Obviously, the coefficients $a_{n,k,l,i}$ are positive for all possible values of n, k, l and i . Firstly, we prove two lemmas about these coefficients.

Lemma 7. If $0 \le l \le k - 1 \le m - 1$, $m = \lfloor n/2 \rfloor$, and $l + 1 \le i \le k - 1$, then $a_{n,k,l,i} > a_{n,k,l,i+1}$. Proof. We evaluate

$$
\frac{a_{n,k,l,i}}{a_{n,k,l,i+1}} = \frac{\begin{bmatrix} n-2i \\ k-i \end{bmatrix} \begin{bmatrix} i \\ l \end{bmatrix} \alpha_{l,i} q^{\binom{i-l-1}{2}} \sigma_{n,i}}{\begin{bmatrix} n-2i-2 \\ k-i-1 \end{bmatrix} \begin{bmatrix} i+1 \\ l \end{bmatrix} \alpha_{l,i+1} q^{\binom{i-l}{2}} \sigma_{n,i+1}} \\
= \frac{(q^{2m-2i}-1) \lambda_{n,i} (q^{i-l+1}-1) \alpha_{l,i} (q^{i+1}-1) \tau_{n,i}}{(q^{i+1}-1) \alpha_{l,i+1} q^{i-l-1} (q^{2m-2i}-1)} \\
= \frac{\lambda_{n,i} (q^{i-l+1}-1) \alpha_{l,i} \tau_{n,i}}{\alpha_{l,i+1} q^{i-l-1}} = \lambda_{n,i} \tau_{n,i} (q^2 - \frac{1}{q^{i-l-1}}) \frac{\alpha_{l,i}}{\alpha_{l,i+1}}.
$$

Using that

$$
\alpha_{l,i+1} = q^{i-l}(q^{l+1} + q - 1) - 1 = q\alpha_{l,i} + q - 1,\tag{11}
$$

we obtain

$$
\frac{\alpha_{l,i}}{\alpha_{l,i+1}} = \frac{\alpha_{l,i}}{q \alpha_{l,i} + q - 1} = \frac{1}{q + \frac{q-1}{\alpha_{l,i}}} \ge \frac{1}{q + \frac{1}{2}} = \frac{2}{2q + 1},\tag{12}
$$

and therefore

$$
\frac{a_{n,k,l,i}}{a_{n,k,l,i+1}} \ge \frac{2\lambda_{n,i}\tau_{n,i}}{2q+1}(q^2 - \frac{1}{q^{i-l-1}}) \ge \frac{2\lambda_{n,i}\tau_{n,i}(q^2-1)}{2q+1}.
$$

According to Lemma [6](#page-4-0) and Lemma [5,](#page-3-0) we have that $\lambda_{n,i} \tau_{n,i} \geq 1$ in the cases when n is odd, or n and q are even, or $n \equiv 2 \pmod{4}$ and $q \equiv 3 \pmod{4}$. Hence in these cases

$$
\frac{a_{n,k,l,i}}{a_{n,k,l,i+1}} \ge \frac{2(q^2-1)}{2q+1} > 1.
$$

In the case when $n = 2m$ is even and q is odd $(q \ge 3)$, such that (1) $n \equiv 2 \pmod{4}$ and $q \equiv 1 \pmod{4}$, or (2) $n \equiv 0 \pmod{4}$, $\tau_{n,i} \geq (q+1)/2$, so

$$
\frac{a_{n,k,l,i}}{a_{n,k,l,i+1}} \ge \frac{2(q^2-1)(q+1)}{2q(2q+1)} > \frac{(q^2-1)}{2q} \ge \frac{8}{6} > 1.
$$

Hence $a_{n,k,l,i} > a_{n,k,l,i+1}$ for $l+1 \leq i \leq k-1$ in all cases.

Lemma 8. If $q \ge 3$ and $0 \le l \le k-1$, then $a_{n,k,l,l} > a_{n,k,l,l+1}$.

Proof. Now

$$
\frac{a_{n,k,l,l}}{a_{n,k,l,l+1}} = \frac{\binom{n-2l}{k-l}\left(q-1\right)\sigma_{n,l}}{\binom{n-2l-2}{k-l-1}\alpha_{l,l+1}\sigma_{n,l+1}}
$$
\n
$$
= \frac{\left(q^{2m-2l}-1\right)\lambda_{n,l}\left(q-1\right)\tau_{n,l}\left(q^{l+1}-1\right)}{\left(q^{l+1}+q-2\right)\left(q^{2m-2l}-1\right)}
$$
\n
$$
= \frac{\lambda_{n,l}\tau_{n,l}\left(q-1\right)\left(q^{l+1}-1\right)}{\left(q^{l+1}+q-2\right)}.
$$

As in the previous proof,

$$
\frac{a_{n,k,l,l}}{a_{n,k,l,l+1}} \ge \frac{(q-1)(q^{l+1}-1)}{q^{l+1}+q-2} = \frac{1}{\frac{1}{q-1}+\frac{1}{q^{l+1}-1}} \ge \frac{q-1}{2} \ge 1
$$

in the cases (1) $n = 2m + 1$, (2) $n = 2m$ and $q = 2^s$, and (3) $n = 2m \equiv 2 \pmod{4}$ and $q \equiv 3$ (mod 4).

Let now $n \equiv 0 \pmod{4}$ and $q \equiv 1 \pmod{2}$, or $n \equiv 2 \pmod{4}$ and $q \equiv 1 \pmod{4}$. In this case

$$
\frac{a_{n,k,l,l}}{a_{n,k,l,l+1}} \ge \frac{(q+1)(q-1)(q^{l+1}-1)}{2(q^{l+1}+q-2)} \ge \frac{(q+1)(q-1)}{4} \ge \frac{q^2-1}{4} \ge 2.
$$

This gives us that $a_{n,k,l,l} > a_{n,k,l,l+1}$ if $q \geq 3$.

Consider the binary case separately, i.e. $q = 2$. Now

$$
a_{n,k,l,i} = \begin{bmatrix} n-2i \\ k-i \end{bmatrix} \begin{bmatrix} i \\ l \end{bmatrix} \frac{\alpha_{l,i}}{2^{l+1}-1} 2^{\binom{i-l-1}{2}} \sigma_{n,i}, \quad l+1 \leq i \leq k, \quad a_{n,k,l,l} = \begin{bmatrix} n-2l \\ k-l \end{bmatrix} \sigma_{n,l},
$$

where $\alpha_{l,i} = 2^{i-l-1}(2^{l+1} + 1) - 1$ for $l+1 \leq i \leq k$. Using that

$$
\alpha_{l,l+1} = 2^{l+1}, \quad \alpha_{l,l+2} = 2^{l+2} + 1, \quad \alpha_{l,l+3} = 2^{l+3} + 3,
$$

we obtain

$$
a_{n,k,l,l+1} = \begin{bmatrix} n-2l-2 \\ k-l-1 \end{bmatrix} \begin{bmatrix} l+1 \\ l \end{bmatrix} \frac{\alpha_{l,l+1}}{2^{l+1}-1} \sigma_{n,l+1} = 2^{l+1} \begin{bmatrix} n-2l-2 \\ k-l-1 \end{bmatrix} \sigma_{n,l+1},
$$

$$
a_{n,k,l,l+2} = \begin{bmatrix} n-2l-4 \\ k-l-2 \end{bmatrix} \begin{bmatrix} l+2 \\ l \end{bmatrix} \frac{\alpha_{l,l+2}}{2^{l+1}-1} \sigma_{n,l+2} = \frac{(2^{l+2}-1)(2^{l+2}+1)}{3} \begin{bmatrix} n-2l-4 \\ k-l-2 \end{bmatrix} \sigma_{n,l+2},
$$

$$
a_{n,k,l,l+3} = \begin{bmatrix} n-2l-6 \ k-l-3 \end{bmatrix} \begin{bmatrix} l+3 \ l \end{bmatrix} \frac{\alpha_{l,l+3}}{2^{l+1}-1} 2^{\binom{2}{2}} \sigma_{n,l+3}
$$

\n
$$
= \begin{bmatrix} n-2l-6 \ k-l-3 \end{bmatrix} \frac{(2^{l+3}-1)(2^{l+2}-1)(2^{l+1}-1)(2^{l+3}+3)}{21(2^{l+1}-1)} 2\sigma_{n,l+3}
$$

\n
$$
= \frac{(2^{l+3}-1)(2^{l+2}-1)(2^{l+4}+6)}{21} \begin{bmatrix} n-2l-6 \ k-l-3 \end{bmatrix} \sigma_{n,l+3}.
$$

Next, we consider separately the cases $k \le 2l + 1$ and $k > 2l + 1$.

Lemma 9. If $q = 2$ and $k \leq 2l + 1$ then $a_{n,k,l,l} > a_{n,k,l,l+1}$.

Proof. In this case

$$
a_{n,k,l,l} - a_{n,k,l,l+1} = \binom{n-2l}{k-l} \sigma_{n,l} - 2^{l+1} \binom{n-2l-2}{k-l-1} \sigma_{n,l+1}
$$

$$
= [\lambda_{n,l} \tau_{n,l} (2^{l+1} - 1) - 2^{l+1}] \binom{n-2l-2}{k-l-1} \sigma_{n,l+1}.
$$

According to Lemma [6,](#page-4-0) $2^{2m+1-n}\lambda_{n,l} > 1 + \frac{1}{2^{k-l}-1} = \frac{2^{k-l}}{2^{k-l}-1}$ $\frac{2^{n}}{2^{k-l}-1}$. If *n* is odd then $\tau_{n,l}=1$ and therefore $\lambda_{n,l}\tau_{n,l} > \frac{2^{k-l}}{2^{k-l}}$ $\frac{2^{k-l}}{2^{k-l}-1}$. We have the same inequality in the opposite case, when *n* is even, because then

$$
\lambda_{n,l}\tau_{n,l} > \frac{2^{k-l}(2^{n-l}-1)}{2(2^{k-l}-1)(2^{n-l-1}-1)} = \frac{2^{k-l}(2^{n-l}-1)}{(2^{k-l}-1)(2^{n-l}-2)} > \frac{2^{k-l}}{2^{k-l}-1}.
$$

Hence

$$
a_{n,k,l,l} - a_{n,k,l,l+1} > \left[\frac{2^{k-l}(2^{l+1}-1)}{2^{k-l}-1} - 2^{l+1}\right] \begin{bmatrix} n-2l-2\\ k-l-1 \end{bmatrix} \sigma_{n,l+1}
$$

=
$$
\frac{2^{l+1}-2^{k-l}}{2^{k-l}-1} \begin{bmatrix} n-2l-2\\ k-l-1 \end{bmatrix} \sigma_{n,l+1}
$$

=
$$
\frac{2^{2l+1}-2^k}{2^k-2^l} \begin{bmatrix} n-2l-2\\ k-l-1 \end{bmatrix} \sigma_{n,l+1} \ge 0
$$

when $k \leq 2l + 1$.

We are now ready to prove the main theorem.

Proof of Theorem [1](#page-1-0):

We use that

$$
A_{n,k,l} - A_{n,k,l+1} = \sum_{i=l}^{k} (-1)^{i-l} a_{n,k,l,i} = \sum_{j=0}^{k-l} (-1)^j a_{n,k,l,j+l}.
$$
 (13)

We proved in Lemma [7,](#page-5-0) that $a_{n,k,l,i} > a_{n,k,l,i+1}$ if $l+1 \leq i \leq k-1$ for all possible values of l, k, n and q. From Lemma [8](#page-5-1) we have that $a_{n,k,l,l} > a_{n,k,l,l+1}$, if $q \geq 3$, and the same inequality holds in the binary case if $k \leq 2l + 1$. In these cases

• if $k - l = 2r + 1$ then

$$
\sum_{j=0}^{k-l} (-1)^j a_{n,k,l,j+l} = \sum_{s=0}^r (a_{n,k,l,2s} - a_{n,k,l,2s+1}) > 0,
$$

• and if $k - l = 2r$ is even, we have

$$
A_{n,k,l} - A_{n,k,l+1} = \sum_{s=0}^{r-1} (a_{n,k,l,2s} - a_{n,k,l,2s+1}) + a_{n,k,l,2r} > 0.
$$

Consider now the case $q = 2$ and $k > 2l + 1$. Then $k \geq l + 3$ except when $l = 0$ and $k = 2$, but we will leave this special case for last. Let $a = a_{n,k,l,l}-a_{n,k,l,l+1}+a_{n,k,l,l+2}-a_{n,k,l,l+3}$, $k \ge \max\{2l + 2, l + 3\}.$ We have

$$
a_{n,k,l,l+2} - a_{n,k,l,l+3} = \frac{(2^{l+2} - 1)(2^{l+2} + 1)}{3} \begin{bmatrix} n-2l-4 \ k-l-2 \end{bmatrix} \sigma_{n,l+2}
$$

$$
-\frac{(2^{l+3}-1)(2^{l+2}-1)(2^{l+4}+6)}{21}\begin{bmatrix}n-2l-6\\k-l-3\end{bmatrix}\sigma_{n,l+3}
$$

=
$$
[\frac{(2^{2m-2l-4}-1)\lambda_{n,l+2}(2^{l+2}-1)(2^{l+2}+1)\tau_{n,l+2}(2^{l+3}-1)}{3(2^{2m-2l-4}-1)}
$$

$$
-\frac{(2^{l+3}-1)(2^{l+2}-1)(2^{l+4}+6)}{21}\begin{bmatrix}n-2l-6\\k-l-3\end{bmatrix}\sigma_{n,l+3}
$$

$$
=[\frac{\lambda_{n,l+2}(2^{l+2}+1)\tau_{n,l+2}}{3}-\frac{(2^{l+4}+6)}{21}](2^{l+2}-1)(2^{l+3}-1)\begin{bmatrix}n-2l-6\\k-l-3\end{bmatrix}\sigma_{n,l+3}
$$

$$
>(7(2^{l+2}+1)-2^{l+4}-6)\begin{bmatrix}n-2l-6\\k-l-3\end{bmatrix}\sigma_{n,l+3}
$$

For the last inequality we use that $\lambda_{n,l+2}\tau_{n,l+2} > \frac{2^{k-l-2}}{2^{k-l-2}}$ $\frac{2^{k-l-2}}{2^{k-l-2}-1} > 1$, and $(2^{l+2}-1)(2^{l+3}-1) \ge$ 21. It follows that

$$
a_{n,k,l,l+2} - a_{n,k,l,l+3} > (3.2^{l+2} + 1) \binom{n-2l-6}{k-l-3} \sigma_{n,l+3}.
$$

Let
$$
\mu = \min\{\left[\frac{n-2l-6}{k-l-3}\right]\sigma_{n,l+3}, \left[\frac{n-2l-2}{k-l-1}\right]\sigma_{n,l+1}\} > 0
$$
. Then
\n
$$
a > \left(3 \cdot 2^{l+2} + 1 - \frac{2^k - 2^{2l+1}}{2^k - 2^l}\right) \mu = \left(3 \cdot 2^{l+2} + \frac{2^{2l+1} - 2^l}{2^k - 2^l}\right) \mu > 0.
$$

To prove that $A_{n,k,l} > An, k, l+1$, we consider the same two cases depending on the parity of $k - l$.

• If $k - l = 2r + 1$ then

$$
\sum_{j=0}^{k-l} (-1)^j a_{n,k,l,j+l} = a + \sum_{s=2}^r (a_{n,k,l,2s} - a_{n,k,l,2s+1}) > 0.
$$

• In the case when $k - l = 2r$ is even, we have

$$
A_{n,k,l} - A_{n,k,l+1} = a + \sum_{s=2}^{r-1} (a_{n,k,l,2s} - a_{n,k,l,2s+1}) + a_{n,k,l,2r} > 0.
$$

In the last remaining case we have $k = 2$ and $l = 0$. Then

$$
A_{n,2,0} - A_{n,2,1} = {n \choose 2} \sigma_{n,0} - 2 {n-2 \choose 1} \sigma_{n,1} + 5 {n-4 \choose 0} \sigma_{n,2}
$$

=
$$
\frac{(2^n - 1)(2^{n-1} - 1)}{3} - 2(2^{n-2} - 1)\sigma_{n,1} + 5\sigma_{n,2}.
$$

After the calculations, we obtain

$$
A_{n,2,0} - A_{n,2,1} = \begin{cases} 2^{n-3}(2^{n-1} - 1)/3 & \text{if } n = 2m + 1 \\ 2^{n-1}(2^{n-3} + 1)/3 & \text{if } n = 2m. \end{cases}
$$

Hence $A_{n,k,l} > A_{n,k,l+1}$ over \mathbb{F}_q in all cases.

4 Computational results

In this section, we present computational results with the number of inequivalent linear $[n, k, \geq 2]$ codes of different types for given length and dimension over \mathbb{F}_2 and \mathbb{F}_3 , whose dual distance is at least 2 (this means that their generator matrices have no zero columns). Denote by $\mathcal{B}_q(n, k, l)$ the set of all inequivalent q -ary linear codes whose hull has dimension l. In the binary case, the mass formula, that helps to verify classification results for the linear $[n, k]_2$ codes C with $h(C) = l$, is the following

$$
A_{n,k,l} = \sum_{C \in \mathcal{B}_2(n,k,l)} \frac{n!}{\text{Aut}(C)}\tag{14}
$$

In the ternary case we have

$$
A_{n,k,l} = \sum_{C \in \mathcal{B}_3(n,k,l)} \frac{2^n n!}{\text{Aut}(C)}.
$$
 (15)

More information on these formulas for self-dual codes can be found in [\[15\]](#page-14-9). Araya and Harada in [\[1\]](#page-11-2) used them in the classification of the binary LCD codes of length $n \leq 13$ and ternary LCD codes of length $n \leq 10$. They described clearly how to use these mass formulas for the verification of the computational results for classification of the binary LCD [6, 3] codes with all possible minimum distances $d \geq 1$ and dual distances $d^{\perp} \geq 1$.

We classify the binary and ternary linear, self-orthogonal and LCD codes of a given dimension $3 \leq k \leq 10$, length $k + 3 \leq n \leq 20$, minimum distance $d > 2$ and dual distance $d^{\perp} \geq 2$. In the binary case, we classify also the even codes with the corresponding length and dimension, i.e. all linear codes whose codewords have only even weights. We do not count codes with dual distance $d^{\perp} = 1$ because if C is an $[n, k, d]$ code with $d^{\perp} = 1$, then all its codewords share a common zero coordinate, so $C = (0|C_1)$ where C_1 is an $[n-1, k, d]$ code. In this case, $C^{\perp} = (0|C_1^{\perp}) \cup (1|C_1^{\perp}), \mathcal{H}(C) = (0|\mathcal{H}(C_1) \text{ and } h(C) = h(C_1).$ If C is an $[n, k, 1]$ code then $C \cong (0|C_1) \cup (1|C_1)$, where C_1 is an $[n-1, k-1]$ code, and then $C^{\perp} \cong (0|C_1^{\perp})$. This gives us that $\mathcal{H}(C) = (0|\mathcal{H}(C_1)$ and $h(C) = h(C_1)$. Therefore, if we have the number $LCD^*(n, k)$ of all $[n, k, \geq 2]$ LCD codes with dual distance ≥ 2 for all length $\leq n$ and dimensions $\leq k$, we can easily compute the number $LCD(n, k)$ of LCD $[n, k, \ge 1]$ codes with dual distance ≥ 1 , using the following formula

$$
LCD(n,k) = \sum_{m=k+1}^{n} LCD^{*}(m,k) + LCD(n,k-1).
$$

We give a simple example that can be followed by hand.

Example 1. Let $n = 4$, $k = 2$ and $q = 2$. In this case

 $A_{4,2,2} = \sigma_{4,2} = 3, A_{4,2,1} = 12, A_{4,2,0} = 20, B_2(4,2,2) = B_2(4,2,1) = 1, B_2(4,2,0) = 4.$

There are six inequivalent [4, 2] binary codes. The first one is obtained from \mathbb{F}_2^2 by adding two zero columns and it is an LCD code. There are two more binary [4, 2] inequivalent codes with zero columns, and these are the codes $C_2 = \{0000, 0110, 0001, 0111\}$, $h(C_2) = 1$, and $C_3 =$ $\{0000, 0110, 0101, 0011\}, h(C_3) = 0.$ The remaining three codes are $C_4 = \{0000, 1110, 0001, 1111\},\$ $h(C_4) = 0, C_5 = \{0000, 1110, 0101, 1011\}, h(C_5) = 0, \text{ and the self-dual } [4, 2, 2] \text{ code } C_6 =$ $\{0000, 1100, 0011, 1111\}, h(C_6) = 2.0$ nly two of these codes have minimum and dual distance $d = d^{\perp} = 2$ and these are C_5 and C_6 .

We use the following properties of the considered types of binary and ternary codes:

- If C is an LCD code, its dual code C^{\perp} is also LCD. It follows that $LCD^*(n, k) =$ $LCD^*(n, n - k)$ and $LCD(n, k) = LCD(n, n - k)$ both in the binary and the ternary case. The same holds for the number of all linear codes of length n and dimension k .
- Self-orthogonal codes exist only when $k \leq n/2$.
- All binary self-orthogonal codes are even.
- Ternary self-dual codes exist only for lengths a multiple of 4 and only have codewords of Hamming weight a multiple of 3.

We obtain the classification results by the program GENERATION of the software package $QExTNEWE\text{DITION}$ [\[5\]](#page-11-3). The computations were executed on a Windows 11 OS in a single core of an Intel Xeon Gold 5118 CPU with a 2.30 GHz clock frequency.

In the binary case, we classify all linear, even, self-orthogonal and LCD codes of length $n \leq 20$ and dimension $k \leq 10$. The results are presented in Table [1.](#page-12-0) For some values of n and k, when all inequivalent codes are too many (more than a milion), we classify only optimal $[n, k]$ codes. In this cases, we put a $*$ after the number of codes. Consider for example $n = 17$ and $k = 7$. The largest possible minimum distance for a binary [17, 7] code is $d = 6$ (see [\[12\]](#page-14-10)). There are exactly 377 binary linear [17, 7, 6] codes. Exactly 329 of these codes are even, 7 are LCD, but none of them is self-orthogonal. Furthermore, there are 497119 even, 58 self-orthogonal, and 14 734 654 LCD [17, 7, ≥ 2] binary codes with dual distance $d^{\perp} \geq 2$. Since we have the count of both all $[17, 7, 2]$ and optimal $[17, 7, 6]$ even codes without zero columns, we denote this in the table by 497119(329*). The optimal binary self-orthogonal codes with the parameters presented in Table [1](#page-12-0) have been also classified in [\[13\]](#page-14-11).

The results in the table confirm the fact that LCD codes are much more than selforthogonal and even more than even codes for a given length and size, even when considering only inequivalent codes. However, this is not the case if we consider only the optimal codes. In most cases, when the optimal code is unique, it is not LCD. The optimal $[19, 7, 8]$ is selforthogonal, as is the optimal $[20, 8, 8]$ code. The optimal $[17, 8, 6]$ and $[17, 9, 5]$ codes are LCD, but the optimal $[19, 8, 7]$, $[20, 9, 7]$ and $[18, 9, 6]$ are neither self-orthogonal, nor LCD. We see interesting examples in the optimal codes of dimension 7. Out of all 377 optimal [17, 7, 6] codes, none is self-orthogonal, but 7 are LCD codes. We have the opposite situation for length 20, namely out of all 26 optimal [20, 7, 8] codes, none is LCD, but four are self-orthogonal. It is also worth noting the optimal [20, 10, 6] codes, where out of all 1682 codes only one is odd-like (it contains codewords of odd weight), but none of the 1681 even codes is self-orthogonal.

In the ternary case, we consider linear, self-orthogonal and LCD codes. We classify all $[n, k, \geq 2]$ linear, LCD and SO codes without zero columns for $k = 3$ and $n \leq 20$, $k = 4$ and $n \le 15$, $k = 5$ and $n \le 14$, $6 \le k \le 9$ and $n \le 13$. For $k = 7$ and $n = 15, 16, 17$, $k = 8$ and $n = 16, 17, 18,$ and $k = 9, n = 19$ we classify only the self-orthogonal codes. The question mark (?) in the table means that there are too many corresponding codes (more than a million). Furthermore, we classify all optimal codes of these three types with 21 parameters. The results are presented in Table [2.](#page-13-0) Let's describe the codes of dimension 5 in more detail. If we look at the number of codes with length n, $10 \le n \le 14$, we see that the LCD codes are more than half of all inequivalent $[n, 5, \geq 2]$ linear codes. The situation with self-orthogonal codes is quite different - they occur much less often. For example, for length 13, the linear codes are more

than a million, but only 17 of them are self-orthogonal. For the optimal codes, we have: (1) of four [15, 5, 8] linear codes one is LCD (none is self orthogonal since 8 is not a multiple of 3), (2) the only $[16, 5, 9]$ code is self-orthogonal, (3) there are 1804 linear $[17, 5, 9]$ codes, 35 of which are self-orthogonal and 400 are LCD, (4) none of the seven [18, 5, 10] and both [19, 5, 11] codes is LCD, nor self-orthogonal, (5) there are two linear [20, 5, 12] codes and both are self-orthogonal.

We have interesting results with the optimal $[19, 7, 9]_3$ and $[20, 8, 9]_3$ codes. In these cases, all optimal linear codes are self-orthogonal.

5 Conclusion

By a result of Sendrier [\[20\]](#page-14-6), it is known that most linear codes are LCD when q is large. Moreover, the proportion of q-ary linear codes of length n, dimension k and specified hull dimension l to all q -ary linear codes of the same length and dimension is convergent when n and k goes to infinity. Using the limit, Sendrier proved that the average dimension of the hull of a q-ary linear code is asymptotically equal to $\sum_{i\geq 1} \frac{1}{q^i+1}$ $\frac{1}{q^i+1}$ [\[20\]](#page-14-6).

In this paper, we obtain general results on hulls of linear codes, proving that the number of all q -ary linear codes of a given length n and dimension k decreases when the hull dimension increases, for all values of n, k and q .

In addition, we classify all binary linear, even, self-orthogonal and LCD $[n \leq 20, k \leq 1]$ $10, d \geq 2$ codes and the ternary linear, self-orthogonal and LCD $[n \leq 19, k \leq 10, d \geq 2]$ codes (with a few exceptions). For some considered values of n and k , when the number of all inequivalent linear codes is huge, we classify only the optimal codes. The results are listed in Tables [1](#page-12-0) and [2.](#page-13-0)

References

- [1] Makoto Araya, Masaaki Harada, On the classification of linear complementary dual codes, Discrete Mathematics, Volume 342, Issue 1, 2019, 270–278.
- [2] M. Araya, M. Harada, On the minimum weights of binary linear complementary dual codes, Cryptogr. Commun. 12, 285–300 (2020).
- [3] M. Araya, M. Harada and K. Saito, Characterization and classification of optimal LCD codes, Des. Codes Cryptogr. 89, 617–640 (2021).
- [4] Assmus, Jr. E. F., Key, J. D.: Affine and projective planes. Discrete Math. 83, 161–187 (1990).
- [5] I. Bouyukliev, The Program Generation in the Software Package QextNewEdition. In: Bigatti A., Carette J., Davenport J., Joswig M., de Wolff T. (eds) Mathematical Software – ICMS 2020, Lecture Notes in Computer Science, vol 12097. Springer, Cham (2020).
- [6] C. Carlet and S. Guilley, Complementary Dual Codes for Counter-measures to Side-Channel Attacks, Advances in Mathematics of Communications 10, 131–150 (2016).
- [7] C. Carlet, S. Mesnager, C. Tang, Y. Qi and R. Pellikaan, Linear codes over \mathbb{F}_q are equivalent to LCD codes for $q > 3$, IEEE Trans. Inform. Theory 64, 3010-3017 (2018).

 $\frac{1}{1}$

Table 1: Classification of binary linear codes $(d^{\perp} \geq 2)$

	$k=3$																	
$n =$	6	$\overline{7}$	8	9	10	11	12		13	14	15	16		17	18		19	20
linear	14	31	68	137	263	484	878		1538	2649	4474	7421		12093	19420		30680	47793
SO	θ	$\mathbf{1}$	$\mathbf{1}$	3	$\overline{4}$	$\overline{5}$	10		15	17	31	44		$54\,$	91		126	160
$\rm LCD$	$\overline{7}$	15	33	67	132	253	471		839	1491	2560	4294		7142	11583		18423	29070
$k=4$																		
$n =$	7	8 9		10		11	12		13	14		15		[16, 4, 9]		[17, 4, 10]		$\sqrt{18}, 4, 11$
linear	31	129	460	1638		5701		19996	69536	239681		809694		$317*$		$18*$		$2*$
SO	÷.	$\mathbf{1}$	$\mathbf{1}$	3		$\bf 5$	16		26	$52\,$		121		$255(13*)$		523		1267
LCD	15	33	220	839		3077	11228		40668	144447		497679		$124*$		$8*$		0^*
$k=5$																		
$\boldsymbol{n} =$	8	9	10		11		12	13		$\overline{14}$		[15, 5, 8]		16, 5, 9		[17, 5, 9]		$[\overline{1}8,5,10]$
linear	68	460	3221		24342		202064	1767647		15604611		$4*$		$1*$	$1804*$			$7*$
SO			$\overline{0}$		3		$\overline{7}$		17	44		156		$523(1^*)$	$1981(35*)$			9460
LCD	33	220	1681		13537	118878		1080479		9737965		$1*$			0^*		$400*$	0^*
	$k=6$ $k=7$																	
$\boldsymbol{n} =$	$\boldsymbol{9}$	10		11		12		13	10	11		12		13	15	16	17	20
linear	137		1638 24342		474106		10956955		$\overline{263}$	5701		202064		10956955	$\overline{\mathcal{E}}$	$\overline{\mathcal{C}}$	$\overline{?}$	$\overline{\mathcal{P}}$
SO	\bar{a}				3		$\overline{4}$		÷,						12	50	249	2287775
$\rm LCD$	67	839		13537		283650		6807504	132	3077	118878			$\overline{\mathcal{L}}$ 6807504		$\overline{\mathcal{L}}$ $\ddot{?}$		$\overline{?}$
$k=8$ $k=9$																		
$n =$	11	12		13		16 17		18 12		13		14		19				
linear	484	19996		1767647		$\overline{\cdot}$ $\overline{?}$		$\overline{\cdot}$	878	69536		15604611		$\overline{?}$				
_{SO}						$\overline{7}$ 16		137	\overline{a}					56				
$\rm LCD$	253	γ $\overline{\mathcal{L}}$ $\overline{\mathcal{L}}$ 11228 1080479 471								40668		11835111		$\ddot{?}$				
									Optimal codes									
		[19, 4, 12]		[20, 4, 12]		[19, 5, 11]			[20, 5, 12]	[14, 6, 6]		[15, 6, 7]		[16, 6, 7]			$\overline{[17,6,8]}$	
linear		$1*$		$84*$		$2*$			$2*$	47674*		$22*$		$>10^8*$		2145181*		
SO		$2867(1^{*})$		$6893(32*)$		50618		$294990(2^*)$		$15(4^*)$		61		286		1504		
LCD	0^*			$14*$		0^*			0^*	27776*		$10*$		53236943*		807993*		
									Optimal codes									
$\boldsymbol{n} =$	[18, 6, 9]			[19, 6, 9]			[18, 7, 8]			[19,7,9]		[19, 8, 8]		[20, 8, 9]		[20, 9, 8]		
linear	$171*$			$\overline{\mathcal{P}}$			827459*		$61*$			1508*		$23*$		$\overline{32^*}$		
SO	$13831(105*)$			184980(18019*)			2486		$57551(61*)$			2281		$112899(23*)$		1122		
LCD	$4*$?				450403^*		0^*		$363*$		0^*		$2*$		

Table 2: Classification of ternary linear codes $(d^{\perp} \geq 2)$

- [8] C. Carlet, S. Mesnager, C. Tang, Y. Qi, New characterization and parametrization of LCD codes, IEEE Trans. Inf. Theory, 65(1), 39–49 (2019).
- [9] Conway, J.H., Sloane, N.J.A., Sphere Packings, Lattices and Groups, third ed., Springer, New York, 1999.
- [10] S.T. Dougherty, J.-L. Kim, B. Ozkaya, L. Sok and P. Solé, The combinatorics of LCD codes: linear programming bound and orthogonal matrices, Int. J. Inf. Coding Theory 4, 116–128 (2017).
- [11] L. Galvez, J.-L. Kim, N. Lee, Y.G. Roe and B.-S. Won, Some bounds on binary LCD codes, Cryptogr. Commun. 10, 719–728 (2018).
- [12] M. Grassl, Code Tables: Bounds on the parameters of various types of codes, <http://www.codetables.de/>.
- [13] I. Bouyukliev, S. Bouyuklieva, T. A. Gulliver, and P. R. Östergård, "Classification of optimal binary self-orthogonal codes," Journal of Combinatorial Mathematics and Combinatorial Computing, vol. 59, pp. 33-87, 2006.
- [14] M. Harada, K. Saito, Binary linear complementary dual codes, Cryptogr. Commun. 11, 677–696 (2019).
- [15] W. C. Huffman and V. Pless, Fundamentals of Error-Correcting Codes, Cambridge Univ. Press, 2003.
- [16] Luo, G., Cao, X., Chen, X., MDS codes with hulls of arbitrary dimensions and their quantum error correction. IEEE Transactions on Information Theory, 65: 2944-2952, 2019.
- [17] J.L. Massey, Linear codes with complementary duals, Discrete Math. 106/107, 337–342 (1992).
- [18] McKay, B.: Isomorph-free exhaustive generation. J.Algorithms 26, 306–324 (1998).
- [19] Vera Pless, The number of isotropic subspaces in a finite geometry, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Nat. (8)39(1965), 418–421.
- [20] Nicolas Sendrier: On the Dimension of the Hull. SIAM J. Discret. Math. 10(2): 282-293 (1997).