Arindam Khan 🖂 💿

Indian Institute of Science, Bengaluru, India

Aditya Subramanian

Indian Institute of Science, Bengaluru, India

Tobias Widmann Technical University, Munich, Germany

Andreas Wiese Technical University, Munich, Germany

— Abstract -

We study the problem of stabbing rectilinear polygons, where we are given n rectilinear polygons in the plane that we want to stab, i.e., we want to select horizontal line segments such that for each given rectilinear polygon there is a line segment that intersects two opposite (parallel) edges of it. Our goal is to find a set of line segments of minimum total length such that all polygons are stabbed. For the special case of rectangles, there is a O(1)-approximation algorithm and the problem is NP-hard [Chan, van Dijk, Fleszar, Spoerhase, and Wolff, 2018]. Also, the problem admits a QPTAS [Eisenbrand, Gallato, Svensson, and Venzin, 2021] and even a PTAS [Khan, Subramanian, and Wiese, 2022]. However, the approximability for the setting of more general polygons, e.g., L-shapes or T-shapes, is completely open.

In this paper, we characterize the conditions under which the problem admits a $(1 + \varepsilon)$ approximation algorithm. We assume that each input polygon is composed of rectangles that are placed on top of each other, such that for each pair of adjacent edges between rectangles, one edge contains the other. We show that if all input polygons satisfy the *hourglass condition*, then the problem admits a quasi-polynomial time approximation scheme. In particular, it is thus unlikely that this case is APX-hard. Furthermore, we show that there exists a PTAS if each input polygon is composed out of rectangles with a bounded range of widths. On the other hand, if the input polygons do *not* satisfy these conditions, we prove that the problem is APX-hard, already if all input polygons have only eight edges. We remark that all polygons with fewer edges automatically satisfy the hourglass condition. On the other hand, for arbitrary rectilinear polygons we even show a lower bound of $\Omega(\log n)$ for the possible approximation ratio, which implies that the best possible ratio is in $\Theta(\log n)$ since the problem is a special case of SET COVER.

2012 ACM Subject Classification Theory of computation \rightarrow Computational geometry

Keywords and phrases Approximation Algorithms, Stabbing, Rectangles, Rectilinear Polygons, QPTAS, APX-hardness

Funding Arindam Khan: Research partly supported by Pratiksha Trust Young Investigator Award, Google India Research Award, and SERB Core Research Grant (CRG/2022/001176) on "Optimization under Intractability and Uncertainty".

1 Introduction

The STABBING problem is a geometric setting of the well-studied SET COVER problem. We are given a set of geometric objects in the plane. The goal is to compute a set of horizontal line segments of minimum total length such that each given object R is *stabbed*, i.e., there is a line segment ℓ for which $R \setminus \ell$ consists of two connected components. The problem was introduced by Chan, van Dijk, Fleszar, Spoerhase, and Wolff [9] for the case that each given object is an axis-parallel rectangle. In particular, they argued that this case models a

resource allocation problem for frequencies. In this application, the x-axis models time and the y-axis represents a frequency spectrum. Each given rectangle represents a request for a time window $[t_1, t_2]$ and a frequency band $[f_1, f_2]$ that needs to be fulfilled. Each selected segment $[t'_1, t'_2] \times \{f'\}$ corresponds to opening a communication channel f' during a time interval $[t'_1, t'_2]$ which then serves each request whose time window is contained in $[t'_1, t'_2]$ and for which f is a frequency in its corresponding band $[f_1, f_2]$. Also, Chan et al. [9] showed a connection to the GENERALIZED MINIMUM MANHATTAN NETWORK problem.

The first result for the case of rectangles was a polynomial time O(1)-approximation due to Chan et al. [9]. Subsequently, Eisenbrand, Gallato, Svensson, and Venzin improved the approximation ratio to 8 and provided a QPTAS, i.e., a $(1 + \varepsilon)$ -approximation algorithm that runs in quasi-polynomial time [15]. In particular, this implies that the problem is unlikely to be APX-hard. After that, Khan, Subramanian, and Wiese presented a polynomial time approximation scheme (PTAS) for rectangles [28].

A natural question is the STABBING problem for geometric shapes that are more general than rectangles. We restrict ourselves to rectilinear polygons since, e.g., a triangle can be stabbed at one of its vertices at essentially zero cost. A rectilinear polygon can model more general types of requests in the resource allocation application above, for example, requests for which the allowed frequency bands depend on the time window. Also from a theoretical point of view, it is natural to ask which approximation ratios are possible for more general geometric objects.

As mentioned above, STABBING admits a $(1 + \varepsilon)$ -approximation algorithm when all given objects are rectangles [28]. However, is this also true for slightly more general polygons, e.g., that have the shape of an L or a T, or even for arbitrary rectilinear polygons? If not, under which conditions on the input objects is a $(1 + \varepsilon)$ -approximation still possible? Also, given that STABBING is a special case of SET COVER, another natural question is whether it is strictly easier than this problem.

In this paper, we investigate the question above. We focus on a type of rectilinear polygons that we call k-shapes. Intuitively, a k-shape is formed by k rectangles that are stacked on top of each other such that for any two consecutive rectangles, the top edge of the bottom rectangle is contained in the bottom edge of the top rectangle, or vice versa, see Figure 2.

1.1 Our contribution

In this paper, we characterize under which conditions of the k-shapes in the input the STABBING problem admits a $(1 + \varepsilon)$ -approximation algorithm in (quasi-)polynomial time, which makes it unlikely that it is APX-hard in these cases. We provide two separate conditions for this. Also, we prove that if the input objects (slightly) violate these conditions, then the problem becomes APX-hard. For arbitrary k-shapes, we prove even that the problem is as difficult as general SET COVER, which yields a lower bound of $\Omega(\log n)$ for the possible approximation ratio.

Our first condition on the input k-shapes is the hourglass condition. It requires intuitively that the rectangles of each k-shape in the input are stacked like an hourglass (see Figure 1 and Definitions 1 and 3). Formally, it states that if we consider the rectangle of each k-shape of the smallest width, then the rectangles on top of it are ordered non-decreasingly by width, and an analogous mirrored ordering holds for the rectangles below it. For example, L-shapes and T-shapes fulfill this condition. We prove that this setting admits a $(1 + \varepsilon)$ -approximation algorithm for any $\varepsilon > 0$ in quasi-polynomial running time, i.e., in time $n^{(\log n/\varepsilon)^{O(1)}}$. In particular, this makes it unlikely that this case is APX-hard. Our algorithm generalizes the

known QPTAS for the case of rectangles [15]. However, it is arguably simpler. For example, it does not need an O(1)-approximation algorithm for the problem as a subroutine. Instead, we show that the calls to this subroutine in [15] can be replaced by suitable guessing steps and by a $O(\log n)$ -approximation algorithm for general SET COVER.

Our algorithm is based on a hierarchical decomposition of the plane into smaller and smaller rectangular regions. Intuitively, given such a region R, we guess all line segments that are relatively long compared to the width of R. Then, we partition R into smaller rectangular regions inside which we will select only shorter line segments. It can happen that a k-shape K contained in R is composed of at least one wide rectangle (of similar width as the guessed long line segments) and of at least one narrow rectangle. If the guessed long line segments do not stab K, then it is clear that K needs to be stabbed by a short line segment (that we select in one of the subproblems that we recurse into). Such line segments can stab only the narrow rectangles of K. Therefore, in this case we remove the wide rectangle from K and hence make K smaller. The hourglass condition ensures that after this removal, the remainder of K still consists of only one connected component. We crucially need this property in order to ensure that the subproblems of R we recurse into form independent subproblems. This would not be the case if the remainder of K consisted of two connected components such that each of them lies in a different subproblem.

While the hourglass condition is crucial for our algorithm above, it could be that it is not needed in an alternative algorithmic approach that computes a $(1 + \varepsilon)$ -approximation for general k-shapes. However, we prove that this is not the case. We show that our problem is APX-hard, already if the input consists only of 3-shapes that do not satisfy the hourglass condition. On the other hand, note that each 2-shape automatically satisfies the hourglass condition by definition.

In our proof of this APX-hardness result, we construct 3-shapes that are composed out of three rectangles whose widths differ a lot. We prove that the latter is necessary in order to prove that our problem is APX-hard. To this end, we show that it admits a polynomial time $(1 + \varepsilon)$ -approximation algorithm for any $\varepsilon > 0$ if each k-shape is composed out of rectangles whose widths are in a *constant* range. This yields our second condition under which our problem admits a $(1 + \varepsilon)$ -approximation. Our algorithm is a generalization of the PTAS for rectangles in [28]. One crucial insight is that if the widths of the rectangles of each input k-shape differ by at most a constant factor of $1/\delta$, then we can reduce our problem to the setting of rectangles by losing only a factor of $O(1/\delta)$. To do this, we simply replace each k-shape K by the smallest rectangle that contains K. We use this insight in one step of our algorithm where we need a O(1)-approximation algorithm as a black-box. More precisely, we again partition the input plane hierarchically into smaller and smaller rectangular regions. In the process, we repeatedly need to compute constant factor approximations for certain sets of k-shapes that intuitively admit a solution whose cost is at most $O(\delta \in OPT)$; for those, we use the mentioned algorithm. We stab all other k-shapes with segments whose total cost is at most $(1 + \varepsilon)$ OPT, which yields a PTAS.

We round up our results by showing that for general k-shapes and, more generally, even arbitrary rectilinear polygons that are composed of k rectangles each, STABBING admits a polynomial time O(k)-approximation algorithm. A natural question is whether the dependence on k (and the input size) in the approximation ratio can be avoided and there is, e.g., also a O(1)-approximation. We show that this is not the case: for arbitrary k, we prove that STABBING for k-shapes is as difficult as arbitrary instance of SET COVER, which yields a lower bound of $\Omega(\log n)$ for our approximation ratio.

1.2 Other related work

As mentioned above, the STABBING problem is a special case of SET COVER which is NP-hard [20] and which does not admit a $(c \cdot \ln n)$ -approximation algorithm for SET COVER for any c < 1, assuming that $\mathsf{P} \neq \mathsf{NP}$ [14] (see also [16]). On the other hand, a simple polynomial time greedy algorithm [12] achieves an approximation ratio of $O(\log n)$.

Das, Fleszar, Kobourov, Spoerhase, Veeramoni, and Wolff [13] studied approximation algorithms for the GENERALIZED MINIMUM MANHATTAN NETWORK (GMMN) problem, where given a set of n pairs of terminal vertices, the goal is to find a minimum-length rectilinear tree such that each pair is connected by a Manhattan path. The currently best known approximation ratio for this problem is $(4 + \varepsilon) \log n$, due to Khan, Subramanian, and Wiese [28] by using their PTAS for STABBING as a subroutine in a variant of the algorithm of Das et al. [13].

Gaur, Ibaraki, and Krishnamurti [21] studied the problem of stabbing rectangles by a minimum number of axis-aligned lines and obtained an LP-based 2-approximation algorithm. Kovaleva and Spieksma [30] studied a weighted generalization of this problem and gave an O(1)-approximation algorithm.

Geometric set cover is a related geometric special case of general SET COVER, where the given sets are geometric objects. Brönnimann and Goodrich [5] first gave an $O(d \log(d \cdot \text{OPT}))$ -approximation algorithm for unweighted geometric set cover where d is the dual VC-dimension of the set system and and OPT is the value of the optimal solution. Aronov, Ezra, and Sharir [3] utilized ε -nets to design an $O(\log \log \text{OPT})$ -approximation algorithm for the hitting set problem involving axis-parallel rectangles. Varadarajan [37] providing an improved approximation algorithm for weighted geometric set cover for fat triangles or disks, and his techniques were extended by Chan, Grant, Könemann, and Sharpe [7] to any set system with low shallow cell complexity. Subsequently, Chan and Grant [6], and Mustafa, Raman, and Ray [33] have settled the APX-hardness statuses of (almost) all natural variants for this problem. Recently, these problem are studied under online and dynamic setting as well [2, 8, 25].

Maximum Independent Set of Rectangles is another related problem. The problems admits a QPTAS [1], and recently a breakthrough O(1)-approximation algorithm was given by Mitchell [32]. Subsequently, a $(2 + \varepsilon)$ -approximation guarantee [18] was achieved.

Rectangle packing and covering problems such as two-dimensional knapsack [23, 26, 17], two-dimensional bin packing [4, 27], strip packing [22, 24] etc. are well-studied in computational geometry and approximation algorithms. We refer the readers to [11] for a survey on the approximation/online algorithms related to rectangles.

Rectilinear polygons appear naturally in the context of circuit design [31], architectural design [35], geometric information systems [10], computer graphics [36], etc. In computational geometry, often problems (for general polygons) are studied in the rectilinear setting, e.g., the art gallery problem [38], rectilinear convex hull [34], and rectilinear steinter tree [19].

2 Preliminaries

We start with some basic definitions and notation. We represent a given axis-aligned rectangle R_i as the Cartesian product of two given closed and bounded intervals, i.e., $R_i = [x_i^\ell, x_i^r] \times [y_i^b, y_i^t]$ for given coordinates $x_i^\ell, x_i^r, y_i^b, y_i^t \in \mathbb{N}$, where $x_i^\ell \leq x_i^r$ and $y_i^b \leq y_i^t$. The following notation will be useful: we define

- $b(R_i) := [x_i^{\ell}, x_i^r] \times \{y_i^b\}$ as the bottom edge of R_i ,
- $t(R_i) := [x_i^{\ell}, x_i^r] \times \{y_i^t\}$ as the top edge of R_i , and

• $w(R_i) := (x_i^r - x_i^\ell)$ as the width of R_i .

A horizontal line segment $s \in \mathbb{R}^2$ is a Cartesian product $s = [x^{\ell}, x^r] \times \{y\}$ with coordinates $x^{\ell}, x^r, y \in \mathbb{N}$ and $x^{\ell} \leq x^r$. We say that s stabs the rectangle R_i if and only if $R_i \cap s = x_i^{\ell}, x_i^r \times \{y\}$. Also, we define $|s| := x^r - x^{\ell}$ is the *length* or the cost of s. We will study the STABBING problem in the setting where each given object is a k-shape.

▶ **Definition 1** (k-shape). Let $k \in \mathbb{N}$. A k-shape K is the union of at most k axis-aligned rectangles $R_1 \cup \cdots \cup R_k = K$ such that $t(R_i) \subseteq b(R_{i+1})$ or $t(R_i) \supseteq b(R_{i+1})$ for each $i \in \{1, \ldots, k-1\}$.



Figure 1 Examples of *k*-shapes satisfying the hourglass condition



Figure 2 A 3-shape not satisfying the hourglass condition (left), and a stack of rectangles that does not form a k-shape (right).

We say that a k-shape $K = R_1 \cup \cdots \cup R_k$ is stabled by a line segment s, if there exists an index $i \in \{1, \ldots, k\}$ such that the rectangle R_i is stabled by s. This leads to the following formal definition of the STABBING problem for k-shapes.

▶ **Definition 2.** Let $k \in \mathbb{N}$. An instance of the k-STABBING problem for k-shapes is a finite set of k-shapes \mathcal{K} , where the objective is to find a set \mathcal{S} of horizontal line segments of minimum total length, such that every k-shape in \mathcal{K} is stabled by a segment in \mathcal{S} .

In the following section, we shall use the term OPT interchangeably to refer to the optimal solution to the problem, and also to represent its cost, i.e., the total length of segments in the set. Similarly, SOL will be used to represent a solution set and also its cost.

3 Quasi-polynomial-time approximation scheme

In this section, we present our QPTAS for k-STABBING. The algorithm is an adaptation of the QPTAS for STABBING [15] to the more general case of k-shapes.

Let $\varepsilon > 0$ and suppose we are given a set of k-shapes \mathcal{K} . In this section, we assume that each given k-shape $K \in \mathcal{K}$ satisfies the hourglass condition (see Figure 1).

6

▶ **Definition 3.** A k-shape $K = R_1 \cup \cdots \cup R_k$ satisfies the hourglass condition if there is no value $i \in \{2, ..., k-1\}$ such that both $w(R_{i-1}) < w(R_i)$ and $w(R_{i+1}) < w(R_i)$.

For each given k-shape K, we define $w_{\max}(K) \coloneqq \max_{i \in \{1,\dots,k\}} w(R_i)$ and $w_{\min}(K) \coloneqq \min_{i \in \{1,\dots,k\}} w(R_i)$ which are the widths of the widest and most marrow parts of K, respectively. For sets of k-shapes $\mathcal{K}' \subseteq \mathcal{K}$, we define accordingly $w_{\max}(\mathcal{K}') \coloneqq \max_{K \in \mathcal{K}'} w_{\max}(K)$, $w_{\min}(\mathcal{K}') \coloneqq \min_{K \in \mathcal{K}'} w_{\min}(K)$. Moreover, we define $w_{\operatorname{range}}(\mathcal{K}') \coloneqq \min\{w | \exists x \forall K \in \mathcal{K}' : K \subseteq [x, x + w] \times \mathbb{R}\}$ as the width of the most narrow strip that contains all k-shapes in \mathcal{K}' . Further, we note here that there are n given k-shapes and each is described by at most 2k distinct points. Therefore, the solution to the instance has only $\binom{2kn}{2}$ combinatorially distinct candidate segments, which is a polynomial in n (we shall use the notation that the number of candidate segments is $\operatorname{poly}(n)$).

▶ Lemma 4. By losing a factor of $1 + \varepsilon$ in our approximation ratio, we assume that $\frac{\varepsilon}{n} < w_{\min}(\mathcal{K}) \leq w_{\max}(\mathcal{K}) \leq \log n$ and $w_{\text{range}}(\mathcal{K}) \leq n \log n$.

Let $\mu := \varepsilon/\log^2 n$. We partition the plane into relatively wide vertical strips of width $w_{\max}(\mathcal{K})/\mu$ each. We do this such that, intuitively, almost all input shapes are contained in one of our strips, and the remaining shapes, which are intersected by the vertical grid lines, can be stabbed very cheaply. To construct this partition, we define vertical grid lines with a spacing of $w_{\max}(\mathcal{K})/\mu$ and give them a random horizontal shift (see Figure 3). Then, each shape in \mathcal{K} intersects one of these grid lines only with very small probability. Therefore, we can show that there exists a specific way to perform the shift of our grid lines such that all input shapes intersecting our grid lines can be stabbed with line segments whose cost is at most $\mu \cdot \text{OPT}$.





Formally, we invoke the following lemma with our choice for μ defined above. It guesses a set of line segments that yield our desired partition into narrow strips, i.e., it produces a polynomial number of candidate sets such that one of the has the claimed property. Algorithmically, we recurse on each of these polynomially many options and at the end output the returned solution with the smallest total cost.

▶ Lemma 5 (Partitioning into narrow strips). Let $\mu > 0$ such that $\mu/n < w_{\min}(\mathcal{K})$. In polynomial time, we can guess a partition of \mathcal{K} into sets $\mathcal{K}_0, \ldots, \mathcal{K}_t$ and one special set \mathcal{K}_{rest} such that

- (i) OPT $\geq \sum_{\ell=1}^{t} \text{OPT}(\mathcal{K}_{\ell}),$
- (ii) $OPT(\mathcal{K}_{rest}) \leq 8\mu \cdot OPT$, and
- (iii) $w_{\text{range}}(\mathcal{K}_i) \leq w_{\max}(\mathcal{K})/\mu$ for each $i \in \{1, \ldots, t\}$.

We compute an $O(\log n)$ -approximate solution for stabbing \mathcal{K}_{rest} by reducing our problem to an instance of SET COVER (see Appendix A.1 for details). By our choice of μ , the resulting cost is at most $O(\log n \cdot \mu \cdot OPT) = O(OPT \cdot \varepsilon / \log n)$.

Now let \mathcal{K}_i be one of the sets of k-shapes due to Lemma 5. We define $S_i := [a, b] \times \mathbb{R}$ for some values $a, b \in \mathbb{R}$ with $b - a \leq w_{\max}(\mathcal{K})/\mu$ such that each k-shape in \mathcal{K}_i is contained in S_i . We want to partition S_i along horizontal lines into rectangular pieces such that each resulting piece contains line segments from $OPT(\mathcal{K}_i)$ of total cost at most $O(w_{\max}(\mathcal{K})/\mu^2)$. To this end, we guess whether the segments in $OPT(\mathcal{K}_i)$ have a total cost of at most $w_{\max}(\mathcal{K})/\mu^2$. If this is not the case, we guess a line segment $s = [a, b] \times \{h\}$ for some value $h \in \mathbb{N}$ according to the following lemma, which intuitively partitions S_i in a balanced way according to the segments in $OPT(\mathcal{K}_i)$. We call such a segment s a balanced horizontal cut.

▶ Lemma 6. If $OPT(\mathcal{K}_i) > w_{\max}(\mathcal{K})/\mu^2$ then in polynomial time we can guess a value $h \in \mathbb{N}$ and a corresponding line segment $s = [a, b] \times \{h\}$ such that each connected component C of $S_i \setminus s$ contains segments from $OPT(\mathcal{K}_i)$ whose total cost is at least $OPT(\mathcal{K}_i)/2 - w_{\max}(\mathcal{K})/\mu$.

We add s to our solution and recurse on each connected component C of $S_i \setminus s$ separately. The resulting subproblem is to stab all input shapes that are contained in C. Observe that s stabs all k-shapes contained in S_i that intersect both connected components of $S_i \setminus s$. Given C, we guess again whether OPT(C), i.e., the optimal solution for all k-shapes contained in C, has a total cost of at most $w_{\max}(\mathcal{K})/\mu^2$, and if not, we guess a corresponding horizontal line segment. Note that we stop after at most $O(\log n)$ recursion levels if all guesses are correct, since $OPT(S_i) \leq OPT \leq n \log n$ due to our preprocessing in Lemma 4. We enforce that in any case we stop after $O(\log n)$ recursion levels in order to guarantee a quasi-polynomial bound on the running time later.

▶ Lemma 7. If all guesses for the balanced horizontal cut are correct, then their total cost is bounded by $3\mu \cdot \text{OPT}(\mathcal{K}_i)$.

At the end, each resulting subproblem is characterized by a rectangle C of width at most $w_{\max}(\mathcal{K})/\mu$ and for which $OPT(C) \leq w_{\max}(\mathcal{K})/\mu^2$. We guess all line segments in OPT(C) whose width is larger than $\varepsilon w_{\max}(\mathcal{K})$. Since $OPT(C) \leq w_{\max}(\mathcal{K})/\mu^2$ there can be at most $1/\varepsilon\mu^2 = \varepsilon^{-3}\log^2 n$ of them, and for each of them there are only poly(n) options. Hence, we can guess them in time $n^{O(\varepsilon^{-3}\log^2 n)}$. Let \mathcal{S}_C denote the guessed segments.

Our next step crucially deviates from the known (Q)PTASs for stabbing rectangles [15, 28]. Inside C, there might be a k-shape K that is not stabbed by any segment in \mathcal{S}_C but for which one of its rectangles R_i satisfies that $w(R_i) > \varepsilon w_{\max}(\mathcal{K})$. Since we have guessed all segments in C of width larger than $\varepsilon w_{\max}(\mathcal{K})$ and did not yet stab K, we know that the optimal solution does not stab K by stabbing R_i (but by stabbing another rectangle that K is composed of). Therefore, we modify K by removing R_i from K. We do this for each rectangle R_i with $w(R_j) > \varepsilon w_{\max}(\mathcal{K})$ that is part of a k-shape K that is contained in C but not yet stabbed. Denote by $\mathcal{K}'(C)$ the resulting set of k-shapes. Importantly, the hourglass property implies that still each K shapes has only one single connected component. This is the reason why we imposed this property.

Observe that for each $K \in \mathcal{K}'(C)$ we have that $w_{\max}(K) \leq \varepsilon \cdot w_{\max}(K)$. Thus, we made progress in the sense that the maximum width of any k-shape reduces by a factor of ε . Also, if all our guesses are correct, then our total cost is small, i.e., $O(\mu \cdot \text{OPT})$. Also, the number of guesses is quasi-polynomially bounded since for each guess there are only $n^{O(\varepsilon^{-3} \log^2 n)}$ many options and our recursion depth is only $O(\log n)$.

▶ Lemma 8. If all our guesses are correct, then the total cost for the selected line segments due to Lemmas 5 and Lemmas 7 is bounded by $O(\mu \cdot \text{OPT})$. Also, the total number of

(combinations of) guesses is bounded by $n^{O(\varepsilon^{-3}\log^2 n)}$.

We continue recursively with each resulting subproblem. Since initially $\frac{\varepsilon}{n} < w_{\min}(\mathcal{K}) \leq w_{\max}(\mathcal{K}) \leq \log n$, we stop after applying the algorithm above for $O(\log(n/\varepsilon))$ levels. Each level incurs in total at most $n^{O(\varepsilon^{-3}\log^2 n)}$ guesses, which yields a total running time of $n^{O(\varepsilon^{-4}\log^3 n)}$. Also, our approximation ratio can easily be bounded by $(1 + \mu)^{O(\log n)} = 1 + O(\varepsilon)$.

Theorem 9. There is a QPTAS for the stabbing problem for k-shapes that satisfy the hourglass condition.

4 PTAS if pieces have bounded ratio of widths

In this section, we improve our QPTAS from Section 3 to a PTAS in the special case that for each given k-shape, for any two of its rectangles R_i, R_j , it holds that $\delta w(R_j) \leq w(R_i) \leq w(R_j)/\delta$ for a given constant $\delta > 0$.

Let α be a constant for which the problem admits an α -approximation algorithm (We show the existence of such an algorithm in Section B.1). Without loss of generality, we assume that $\alpha, (1/\varepsilon) \in \mathbb{N}$, and we say that an x-coordinate $x \in \mathbb{R}$ is discrete if x is an integral multiple of ε^d , where we define $d \in \mathbb{N}$ such that $\varepsilon^3/n < \varepsilon^d \le \varepsilon^2/n$; note that hence d is unique. Similarly a y-coordinate is called *discrete* if it is integral. A point is called *discrete* if its x and y coordinates are discrete, and similarly a segment or a rectangle is said to be *discrete* if both of its end points, or both of its diagonally opposite corners are discrete.

▶ Lemma 10. Let α be a constant for which k-STABBING admits an α -approximate algorithm and let $\varepsilon > 0$ with $\varepsilon < 1/3$. In polynomial time we can compute a new instance of k-STABBING, in which each $K \in \mathcal{K}$ satisfies,

(i) $\frac{\alpha\varepsilon}{n} < w_{\min}(K) \le w_{\max}(K) \le \alpha$,

(ii) all points defining K are discrete,

(iii) K lies within a bounding box of $[0, \alpha n] \times [0, (k+1)n]$,

and this new instance admits a solution of cost at most $(1 + O(\varepsilon)) \cdot \text{OPT}$ with each segment in the solution being discrete, and having length at most α/ε .

First, we apply Lemma 10 in order to preprocess our instance. In our algorithm, we intuitively embed the recursion of our QPTAS in Section 3 into a polynomial time dynamic program. The idea is to construct a DP-table that contains one cell for each possible subproblem of a recursive call. Formally, we introduce one DP-cell DP(R, S) for each combination of

a closed rectangle $R \subseteq [0, \alpha n] \times [0, (k+1)n]$ with discrete coordinates,

= a set S of at most ε^{-3} discrete horizontal line segments, that all intersect R.

This DP-cell encodes the subproblem of stabbing all input k-shapes that are contained in R and that are not already stabbed by the segments in S. Clearly, the DP-cell DP($[0, \alpha n] \times [0, (k+1)n], \emptyset$) corresponds to our given problem.

Given a DP-cell DP(R, S), we compute its solution as follows. The base case occurs when the line segments in S already stab all k-shapes that are contained in R. Then we define $DP(R, S) := \emptyset$. Another easy case occurs when there is a line segment $\ell \in S$ that stabs the interior of R, i.e., $R \setminus \ell$ has two connected components R_1 and R_2 . Assume that S_1 and S_2 are parts of the line segments from S that intersect R_1 and R_2 , respectively. Then we define $DP(R, S) := DP(R_1, S_1) \cup DP(R_2, S_2) \cup \{\ell\}$. We will refer to this later as the *trivial operation*.

Otherwise, we compute a polynomial number of candidate solutions as follows,

- 1. Add operation. For each set S' of discrete segments contained in R for which $|S| + |S'| \le 3\varepsilon^{-3}$ holds, we generate the candidate solution $S' \cup DP(R, S \cup S')$.
- 2. Line operation. Consider each vertical line ℓ that intersects the interior of R. Let \mathcal{K}_{ℓ} denote the set of k-shapes contained in R that intersect with ℓ . For each $K \in \mathcal{K}_{\ell}$ we construct the smallest axis-parallel rectangle that contains K, let \mathcal{R}_{ℓ} denote the resulting set of rectangles. We apply the PTAS for stabbing rectangles [28] to \mathcal{R}_{ℓ} , let \mathcal{S}_{ℓ} denote the computed set of segments. We will show later that the optimal solution for \mathcal{R}_{ℓ} is by at most a factor $O(1/\delta)$ more expensive that the optimal solution for \mathcal{K}_{ℓ} , and that this approximation ratio is good enough for our purposes in this step. Denote by R_1 and R_2 the connected components of $R \setminus \ell$ and by \mathcal{S}_1 and \mathcal{S}_2 the parts of segments from \mathcal{S} that intersect R_1 and R_2 , respectively. We define the candidate solution $\mathcal{S}_{\ell} \cup DP(R_1, \mathcal{S}_1) \cup DP(R_2, \mathcal{S}_2)$.

We store in DP(R, S) the candidate solution with smallest cost. Finally, we output the solution stored in the cell $DP([0, \alpha n] \times [0, 2kn], \emptyset)$.

4.1 Analysis

We first note that all DP subproblems and operations are defined on discrete coordinates, and since there are only a polynomial $\frac{\alpha n}{\varepsilon^d} \times 2kn \leq 2\alpha k\varepsilon^{-3}n^3$ number of discrete points, the running time of the dynamic program is also polynomial.

▶ Lemma 11. The running time of the above dynamic program is $(kn/\varepsilon)^{O(1/\varepsilon^3)}$.

Our consideration of the approximation factor is similar to the analysis of the PTAS by Khan, Subramanian, and Wiese [28] and our QPTAS in Section 3. We describe here its main structure and highlight the key differences. We refer to Appendix B for all details.

The solution computed by our DP corresponds to performing a sequence of trivial, add, and *line* operations, and recursing on the respective subproblems. It is sufficient to argue that there exists a sequence of these operations such that

- there exists a DP-cell for each arising subproblem; in particular, the number of line segments in each subproblem is bounded by $3\varepsilon^{-3}$ and,
- the total cost of the computed solution is bounded by $(1 + O(\varepsilon))$ OPT.

We now describe this sequence. It is based on a hierarchical grid of vertices lines, shifted by a random offset $r \in \{0, \varepsilon^d, 2\varepsilon^d, \ldots, \alpha\varepsilon^{-2}\}$ that we will fix later. For each level $j \in \mathbb{N}_0$, we define a grid line $\{r + t \cdot \alpha\varepsilon^{j-2}\} \times \mathbb{R}$ for each $t \in \mathbb{Z}$. Note that for all $j \leq d+2$, grid lines of level j have discrete x-coordinates. We say that a line segment $\ell \in \text{OPT}$ is of level j if the length of ℓ is in $(\alpha\varepsilon^j, \alpha\varepsilon^{j-1}]$. We say that a line segment of some level j is well-aligned if its left and right endpoint lies on a grid line of level j + 3, and if the y-coordinate of both endpoints is discrete. We can extend each line segment $\ell \in \text{OPT}$ so that it becomes well-aligned, by increasing its length by at most a factor of $1 + O(\varepsilon)$.

▶ Lemma 12. For any value of our offset, by losing a factor of $1+O(\varepsilon)$ in our approximation ratio, we can assume that each line segment $\ell \in \text{OPT}$ is well-aligned.

Note that each horizontal segment $\ell \in \text{OPT}$ satisfies that $\alpha \varepsilon/n < |\ell| \le \alpha \varepsilon^{-1}$. By our choice of d we have $\varepsilon^{d-1} \le \varepsilon/n < \varepsilon^{d-2}$ which implies $\alpha \varepsilon^{d-1} < |\ell| \le \alpha \varepsilon^{-1}$. Since a segment is of level j if its length is in the range $(\alpha \varepsilon^j, \alpha \varepsilon^{j-1}]$, we can conclude that all segments in OPT belong to levels in the range $\{0, \ldots, d-1\}$. From this we can infer that any well-aligned horizontal segment is aligned to a vertical grid line of level at most d+2, which as we noted earlier has discrete x-coordinates.

In our sequence of operations, we first perform one *line operation* for each (vertical) grid line of level j = 0. This is similar as partitioning the instance into narrow strips as we did it in Lemma 5. However, now each strip has a width of $\alpha \varepsilon^{-2}$ instead of $w_{\max}(\mathcal{K})/\mu$. In our following operations, we add horizontal line segments to partition each vertical strip, similar to Section 3. Formally, we sort the segments from OPT of level j = 0 in increasing order of their *y*-coordinates, and pick every (ε^{-3})-th segment, and do an add operation along the (strip wide) line along it. This leads to a trivial operation immediately after that. Finally, we perform add operations for all line segments of level j = 0 in OPT. We call the above operations to be *operations of level* 0.

With the above operations for level j = 0 done, in increasing order of level j = 1, 2, ...we do *operations of level j* similarly as follows:

- line operations on vertical grid lines of level j,
- any valid trivial operations (this step is not done for level 0),
- add, and trivial operations to divide the vertical strips into smaller subproblems,
- \blacksquare and finally the add operations on the segments from OPT of level j,

mimicking the recursive structure from the analysis of the QPTAS.

▶ Lemma 13. The above sequence of operations always leads to valid DP subproblems.

Proof. Consider a subproblem (R, S) obtained at any stage of application of the above operations. The rectangular cell R is always discrete and a subset of $[0, \alpha n] \times [0, (k+1)n]$ since the line and trivial operations are done only on discrete lines. So the only property we need to show is that $|S| \leq 3\varepsilon^{-3}$. Let the last segment added to S be of level j. An add operation of level j is preceded by *line operations* of level j, and hence any segment of level j - 3 already in S gets removed from S by trivial operations, by virtue of it being well-aligned (Lemma 10). Therefore, S only contains segments from the levels j, j - 1, and j - 2. By construction, when we perform add operations, we add at most ε^{-3} segments of any particular level j to S, and hence there are at most $3\varepsilon^{-3}$ segments in S.

We wish to bound the cost of the the above operations. Suppose that we perform a line operation with a vertical line ℓ and let \mathcal{K}_{ℓ} denote the k-shapes that ℓ intersects. Recall that for each *line operation*, we compute a solution that stabs all k-shapes in \mathcal{K}_{ℓ} (and in fact every rectangle in \mathcal{R}_{ℓ}). Note that any horizontal line segment $\ell' \in \text{OPT}$ of some level $j' \geq j$ stabs a k-shape in \mathcal{K}_{ℓ} only if the distance between ℓ and ℓ' is at most $\alpha \varepsilon^{j-1}$. Another key other insight is that since the ratio between the widest and the narrowest part of any $K \in \mathcal{K}_{\ell}$ is $1/\delta$, the solution we compute is also a $O(1/\delta + \varepsilon)$ -approximate solution. Using the above facts, we claim that if we choose our offset r uniformly at random from the range $\{0, \varepsilon^d, 2\varepsilon^2 d, \ldots, \alpha\varepsilon^{-2}\}$, then the overall cost of these *line operations* is only $O(\varepsilon) \cdot \text{OPT}$. Further to bound the cost of the add operations, we note that each add operation is either done on a segment in OPT, or is an operation that created a subproblem. We will show that we can charge the latter operations to segments from OPT inside the subproblem thus created, whose total cost is at least ε^{-1} times the width of the subproblem. We refer to Appendix B for a formal description of our analysis.

▶ Lemma 14. There is a discrete value for the offset $r \in \{0, \varepsilon^d, 2\varepsilon^d, \dots, \varepsilon^{-2}\}$ such that the described sequence of operations produces a solution of cost at most $(1 + O(\varepsilon))$ OPT.

▶ **Theorem 15.** For each constant $k \in \mathbb{N}$ there is a PTAS for the k-STABBING problem when each given k-shape consists of pieces of a constant range of widths that are placed strictly on top of each other.



Figure 4 Construction of *k*-STABBING instance in our reduction from vertex cover.

5 General case

In this section, we study the general case of stabbing rectilinear polygons. In contrast to the cases studied in Sections 3 and 4, we show that it does *not* admit a $(1 + \varepsilon)$ -approximation algorithm, even for only slightly more general types of instances.

5.1 APX-hardness

Formally, we prove that stabbing is APX-hard, already if each input polygon is a 3-shape.

▶ **Theorem 16.** *The stabbing problem for 3-shapes is* APX*-hard.*

On the other hand, any 2-shape satisfies the hourglass property; hence, stabbing is unlikely to be APX-hard for this class of objects since we have a QPTAS for this case.

▶ Proposition 17. Each 2-shape satisfies the hourglass property.

In the remainder of this subsection, we prove Theorem 16. We give an L-reduction (with $\alpha = \beta = 1$) from the vertex cover problem to stabbing for 3-shapes. Note that it is NP-hard to approximate vertex cover with a strictly better approximation factor than $\sqrt{2}$ [29]. We will obtain the same lower bound for stabbing.

Consider a given instance G = (V, E) of vertex cover. Remember that in vertex cover, we are required to select a subset $S \subseteq V$ of smallest size such that for each $e \in E$ one of its end points is in S. We construct an instance of k-STABBING corresponding to G as follows. Assume that $V = \{v_1, \ldots, v_n\}$. For each $v_i \in V$ construct a 1×1 square s_i , such that they are all arranged in a column separated by 1 unit distance each (see Figure 4). Formally, for each $v_i \in V$ the top-left corner of the square s_i has the coordinates (0, 2i - 1). Note that the squares s_1, \ldots, s_n do not belong to our input shapes, but they only help us to construct the latter. For each edge $\{v_i, v_j\} \in E$ we define a 3-shape $r_{i,j}$ as the union of the three rectangles $s_i, [0, n + 1] \times [2i - 1, 2j - 2]$ and s_j (see Figure 4).

Note that none of the resulting shapes satisfies the hourglass property, and also for neither of them the widths of its three rectangles are in a constant range. The width of the widest



Figure 5 The k-shape constructed for set $S = \{v_1, v_2, v_4, v_n\} \in \mathcal{F}$

rectangle of each constructed 3-shape is greater than n, but there is always a feasible solution with cost n that simply stabs the square s_i for each vertex $v_i \in V$. Thus, in any given solution to the stabbing instance, we can assume w.l.o.g. that no 3-shape is stabbed across its widest rectangle.

▶ Lemma 18. For each $\gamma \in \mathbb{N}$, the given instance of vertex cover instance has a solution of size γ if and only if the corresponding k-STABBING instance has a solution of cost γ .

This yields the proof of Theorem 16.

5.2 Set-Cover hardness

In this section, we further show that k-STABBING for arbitrary k-shapes cannot be approximated with a ratio of $o(\log n)$, unless $\mathsf{P} = \mathsf{NP}$. In fact, we show that the problem is as hard as general instances of SET COVER, for which it is known that it does not admit an $o(\log n)$ approximation algorithm, unless $\mathsf{P} \neq \mathsf{NP}$ [14].

▶ **Theorem 19.** The STABBING problem for k-shapes does not admit an $o(\log n)$ -approximation algorithm.

In the remainder of this subsection, we prove Theorem 19. We reduce from the hitting set problem, which is known to be equivalent to SET COVER. In hitting set, we are given a set of elements $P = \{p_1, p_2, \ldots, p_n\}$, and a family of their subsets \mathcal{F} . The aim is to compute a minimum size subset $H \subseteq P$ such that every set in \mathcal{F} , contains a point in H.

Our construction here is similar to the reduction from vertex cover above. For each element $p_i \in P$, we construct a unit square s_i with its top left corner being located at (0, 2i - 1). For each set $S_i = \{p_{i_1}, p_{i_2}, \ldots, p_{i_f}\} \in \mathcal{F}$ we construct a k-shape which is a stack of the rectangles $s_{i_1}, [0, n + 1] \times [2i_1 - 1, 2i_2 - 2], s_{i_2}, [0, n + 1] \times [2i_3 - 1, 2i_3 - 2], s_{i_3}, \ldots, s_{i_{f-1}}, [0, n + 1] \times [2i_{f-1} - 1, 2i_f - 2], s_{i_f}$ (see Figure 5).

Again, these constructed k-shapes neither satisfy the hourglass property nor the condition that the widths of its rectangles are in a bounded range. As before, for each given solution,

we can assume w.l.o.g. that no k-shape is stabled across one of its wide rectangles (i.e., of width n + 1).

▶ Lemma 20. For each $\gamma \in \mathbb{N}$, the given instance of hitting set has a solution of size γ if and only if the constructed k-STABBING instance has a solution of cost γ .

Therefore, we constructed an appropriate approximation preserving solution from SET COVER, which yields the proof of Theorem 19.

5.3 Approximation algorithm

We show that there is a polynomial time O(k)-approximation algorithm for stabbing for k-shapes. Our algorithm is *LP*-relative, meaning that it outputs a solution whose cost is at most by a factor of O(k) larger than the cost of the optimal solution to the canonical LP-formulation for the problem.

Suppose we are given an instance \mathcal{K} of k-STABBING with $n := |\mathcal{K}|$. In principle, there is an infinite set of possible line segments that we could use for our solution. However, it is sufficient to restrict ourselves to a polynomial number of line segments which we construct using the following lemma.

Lemma 21. In polynomial time, we can construct a set C of line segments with the following properties:

- \blacksquare C contains $O((kn)^3)$ segments,
- \square C contains no redundant segments, where a segment is redundant if it stabs exactly the same k-shapes as another segment, or no k-shapes at all, and
- K admits an optimal solution using only the segments from C.
 Using C, we define a linear program that corresponds to K.

$$\min \sum_{s \in \mathcal{C}} |s| \cdot z_s$$
s.t.
$$\sum_{s \in \mathcal{C}: s \text{ stabs } K} z_s \ge 1 \qquad \forall K \in \mathcal{K}$$

$$z_s \ge 0 \qquad \forall s \in \mathcal{C}.$$
(1)

If each k-shape $K \in \mathcal{K}$ is a rectangle, then it was shown by Chan et al. [9] that this LP has a constant integrality gap.

▶ Theorem 22 ([9]). If each k-shape $K \in \mathcal{K}$ is a rectangle, then there is a constant α such that for any solution z to LP (1), in polynomial time we can compute an integral solution to (1) whose cost is at most $\alpha \sum_{s \in \mathcal{F}} |s| \dot{z}_s$.

Using Theorem 22, we construct now an $(\alpha \cdot k)$ -approximation algorithm for arbitrary kshapes. Suppose we are given an optimal solution z^* to the LP (1). We define a new solution \tilde{z} by setting $\tilde{z}_s := k \cdot z_s^*$ for each segment $s \in \mathcal{F}$. Each k-shape $K \in \mathcal{K}$ is composed out of at most k rectangles. Thus, for each k-shape $K \in \mathcal{K}$ there is one of these rectangles R for which $\sum_{s \in \mathcal{F}: s \text{ stabs } R} z_s^* \geq 1/k$ and, therefore, $\sum_{s \in \mathcal{F}: s \text{ stabs } R} \tilde{z}_s \geq 1$. Let \mathcal{R} denote the set of all these rectangles for all k-shapes in \mathcal{K} . We apply Theorem 22 on \tilde{z}_s and \mathcal{R} which yields a set of segments $\tilde{\mathcal{S}}$ whose cost is at most $\alpha \cdot \sum_{s \in \mathcal{F}} |s| \cdot \tilde{z}_s = \alpha k \cdot \sum_{s \in \mathcal{F}} |s| \cdot z_s^* \leq \alpha k \cdot \text{OPT}$. Since $\tilde{\mathcal{S}}$ stabs \mathcal{R} , it also stabs \mathcal{K} . Hence, $\tilde{\mathcal{S}}$ yields an O(k)-approximation to our problem.

Theorem 23. There is a polynomial time O(k)-approximation algorithm for k-STABBING.

We remark that our algorithm extends also to the setting in which each given shape consists of at most k rectangles that are not necessarily connected, but such that still at least one of them needs to be stabled.

— References

- 1 Anna Adamaszek, Sariel Har-Peled, and Andreas Wiese. Approximation schemes for independent set and sparse subsets of polygons. J. ACM, 66(4):29:1–29:40, 2019.
- 2 Pankaj K. Agarwal, Hsien-Chih Chang, Subhash Suri, Allen Xiao, and Jie Xue. Dynamic geometric set cover and hitting set. *ACM Trans. Algorithms*, 18(4):40:1–40:37, 2022.
- 3 Boris Aronov, Esther Ezra, and Micha Sharir. Small-size ε-nets for axis-parallel rectangles and boxes. SIAM J. Comput., 39(7):3248–3282, 2010.
- 4 Nikhil Bansal and Arindam Khan. Improved approximation algorithm for two-dimensional bin packing. In *SODA*, pages 13–25, 2014.
- 5 Hervé Brönnimann and Michael T. Goodrich. Almost optimal set covers in finite vc-dimension. Discret. Comput. Geom., 14(4):463–479, 1995.
- **6** Timothy M. Chan and Elyot Grant. Exact algorithms and apx-hardness results for geometric packing and covering problems. *Comput. Geom.*, 47(2):112–124, 2014.
- 7 Timothy M. Chan, Elyot Grant, Jochen Könemann, and Malcolm Sharpe. Weighted capacitated, priority, and geometric set cover via improved quasi-uniform sampling. In SODA, pages 1576–1585, 2012.
- 8 Timothy M. Chan, Qizheng He, Subhash Suri, and Jie Xue. Dynamic geometric set cover, revisited. In SODA, pages 3496–3528, 2022.
- 9 Timothy M. Chan, Thomas C. van Dijk, Krzysztof Fleszar, Joachim Spoerhase, and Alexander Wolff. Stabbing rectangles by line segments how decomposition reduces the shallow-cell complexity. In *ISAAC*, pages 61:1–61:13, 2018.
- 10 Kang-Tsung Chang. Introduction to geographic information systems (4. ed.). McGraw-Hill, 2008.
- 11 Henrik I Christensen, Arindam Khan, Sebastian Pokutta, and Prasad Tetali. Approximation and online algorithms for multidimensional bin packing: A survey. *Computer Science Review*, 24:63–79, 2017.
- 12 Vasek Chvátal. A greedy heuristic for the set-covering problem. Math. Oper. Res., 4(3):233–235, 1979.
- 13 Aparna Das, Krzysztof Fleszar, Stephen G. Kobourov, Joachim Spoerhase, Sankar Veeramoni, and Alexander Wolff. Polylogarithmic approximation for generalized minimum manhattan networks. 2012. arXiv:1203.6481.
- 14 Irit Dinur and David Steurer. Analytical approach to parallel repetition. In *STOC*, pages 624–633, 2014.
- 15 Friedrich Eisenbrand, Martina Gallato, Ola Svensson, and Moritz Venzin. A QPTAS for stabbing rectangles. 2021. arXiv:2107.06571.
- 16 Uriel Feige. A threshold of $\ln n$ for approximating set cover. J. ACM, 45(4):634–652, 1998.
- 17 Waldo Gálvez, Fabrizio Grandoni, Sandy Heydrich, Salvatore Ingala, Arindam Khan, and Andreas Wiese. Approximating geometric knapsack via l-packings. In FOCS, pages 260–271, 2017.
- 18 Waldo Gálvez, Arindam Khan, Mathieu Mari, Tobias Mömke, Madhusudhan Reddy Pittu, and Andreas Wiese. A $(2+\epsilon)$ -approximation algorithm for maximum independent set of rectangles. 2021. arXiv:2106.00623.
- 19 M. R. Garey and David S. Johnson. The rectilinear steiner tree problem is NP complete. SIAM Journal of Applied Mathematics, 32:826–834, 1977.
- 20 M. R. Garey and David S. Johnson. Computers and Intractability: A Guide to the Theory of NP-Completeness. W. H. Freeman, 1979.
- 21 Daya Ram Gaur, Toshihide Ibaraki, and Ramesh Krishnamurti. Constant ratio approximation algorithms for the rectangle stabbing problem and the rectilinear partitioning problem. J. Algorithms, 43(1):138–152, 2002.
- 22 Rolf Harren, Klaus Jansen, Lars Prädel, and Rob van Stee. A (5/3 + ε)-approximation for strip packing. Comput. Geom., 47(2):248–267, 2014.

- 23 Klaus Jansen, Arindam Khan, Marvin Lira, and K. V. N. Sreenivas. A PTAS for packing hypercubes into a knapsack. In *ICALP*, pages 78:1–78:20, 2022.
- 24 Arindam Khan, Aditya Lonkar, Arnab Maiti, Amatya Sharma, and Andreas Wiese. Tight approximation algorithms for two-dimensional guillotine strip packing. In *ICALP*, pages 80:1–80:20, 2022.
- 25 Arindam Khan, Aditya Lonkar, Saladi Rahul, Aditya Subramanian, and Andreas Wiese. Online and dynamic algorithms for geometric set cover and hitting set. In SoCG, pages 46:1–46:17, 2023.
- 26 Arindam Khan, Arnab Maiti, Amatya Sharma, and Andreas Wiese. On guillotine separable packings for the two-dimensional geometric knapsack problem. In SoCG, pages 48:1–48:17, 2021.
- 27 Arindam Khan and Eklavya Sharma. Tight approximation algorithms for geometric bin packing with skewed items. In APPROX/RANDOM, pages 22:1–22:23, 2021.
- 28 Arindam Khan, Aditya Subramanian, and Andreas Wiese. A PTAS for the horizontal rectangle stabbing problem. In *IPCO*, pages 361–374, 2022.
- 29 Subhash Khot, Dor Minzer, and Muli Safra. Pseudorandom sets in grassmann graph have near-perfect expansion. In FOCS, pages 592–601, 2018.
- 30 Sofia Kovaleva and Frits C. R. Spieksma. Approximation algorithms for rectangle stabbing and interval stabbing problems. *SIAM J. Discret. Math.*, 20(3):748–768, 2006.
- 31 Jens Lienig and Juergen Scheible. *Fundamentals of layout design for electronic circuits*. Springer, 2020.
- 32 Joseph S. B. Mitchell. Approximating maximum independent set for rectangles in the plane. In FOCS, pages 339–350, 2021.
- 33 Nabil H. Mustafa, Rajiv Raman, and Saurabh Ray. Settling the apx-hardness status for geometric set cover. In FOCS, pages 541–550, 2014.
- 34 Thomas Ottmann, Eljas Soisalon-Soininen, and Derick Wood. On the definition and computation of rectilinear convex hulls. *Information Sciences*, 33(3):157–171, 1984.
- 35 Helmut Pottmann, Andreas Asperl, Michael Hofer, Axel Kilian, and Daril Bentley. Architectural geometry, volume 724. Bentley Institute Press Exton, 2007.
- **36** Peter Shirley, Michael Ashikhmin, and Steve Marschner. *Fundamentals of computer graphics*. AK Peters/CRC Press, 2009.
- 37 Kasturi R. Varadarajan. Weighted geometric set cover via quasi-uniform sampling. In STOC, pages 641–648, 2010.
- 38 Chris Worman and J Mark Keil. Polygon decomposition and the orthogonal art gallery problem. Int. J. Comput. Geom. Appl., 17(02):105–138, 2007.

A Missing proofs from Section 3

A.1 Reduction to Set Cover

In this subsection we show that the k-STABBING problem can be reduced to a general weighted set cover instance. We are given an instance of k-STABBING with a set of n k-shapes, \mathcal{K} . Since each k-shape can be described by at most 2k points (2 each describing each of its constituent rectangular sections), we have a total of 2nk points, and only $\binom{2nk}{2} = O(n^2k^2)$ possible combinatorially distinct segments in the solution.

We create an instance of set cover as follows. The universe of elements is given by the set of all k-shapes. Now for each of the $O(n^2k^2)$ possible segments ℓ in the solution, we create a set of weight $|\ell|$, which contains all the k-shapes that the segment stabs. This forms our family of subsets. Now a solution of weight w to the set cover instance, corresponds to a set of segments of weight w which stab all k-shapes in \mathcal{K} . Hence, we have an approximation preserving

reduction from k-STABBING to weighted set cover, and by extension, a $(\log n)$ -approximate algorithm for k-STABBING.

A.2 Proof of Lemma 4

▶ Lemma 4. By losing a factor of $1 + \varepsilon$ in our approximation ratio, we assume that $\frac{\varepsilon}{n} < w_{\min}(\mathcal{K}) \leq w_{\max}(\mathcal{K}) \leq \log n$ and $w_{\text{range}}(\mathcal{K}) \leq n \log n$.

Proof. Using the $(\log n)$ -approximation algorithm, obtain a solution to \mathcal{K} and determine its $\cot \gamma \in [\operatorname{OPT}(\mathcal{K}), \log n \cdot \operatorname{OPT}(\mathcal{K})]$. Consequently, $\operatorname{OPT}(\mathcal{K}) \in [\frac{1}{\log n}\gamma, \gamma]$.

Scaling each k-shape in \mathcal{K} along the x-axis by a factor of $\beta \coloneqq \log n \frac{1}{\gamma}$ yields an instance \mathcal{K}_{β} with $\operatorname{OPT}(\mathcal{K}_{\beta}) = \beta \cdot \operatorname{OPT}(\mathcal{K}) \in [1, \log n]$, which has the following implications for solving \mathcal{K}_{β} :

- Any parts R of k-shapes in \mathcal{K}_{β} with $w(R) > \log n$ can be discarded, since an optimal solution cannot stab them.
- Greedily stabbing all k-shapes $K \in \mathcal{K}_{\beta}$ with $w_{\min}(K) \leq \frac{\varepsilon}{n}$ requires segments of total length at most $\varepsilon \leq \varepsilon \cdot \text{OPT}(\mathcal{K}_{\beta})$.

An algorithm for solving \mathcal{K}_{β} can perform this preprocessing and continue to operate only on k-shapes K with $\frac{\varepsilon}{n} < w_{\min}(K) \leq w_{\max}(K) \leq \log n$. Furthermore, it can be assumed w.l.o.g. that $w_{\text{range}}(\mathcal{K}_{\beta}) \leq n \log n$; otherwise, \mathcal{K}_{β} could be partitioned into independent sub-instances to be solved separately.

A.3 Proof of Lemma 5

▶ Lemma 5 (Partitioning into narrow strips). Let $\mu > 0$ such that $\mu/n < w_{\min}(\mathcal{K})$. In polynomial time, we can guess a partition of \mathcal{K} into sets $\mathcal{K}_0, \ldots, \mathcal{K}_t$ and one special set \mathcal{K}_{rest} such that

- (i) OPT $\geq \sum_{\ell=1}^{t} \text{OPT}(\mathcal{K}_{\ell}),$
- (ii) $OPT(\mathcal{K}_{rest}) \leq 8\mu \cdot OPT$, and
- (iii) $w_{\text{range}}(\mathcal{K}_i) \leq w_{\max}(\mathcal{K})/\mu$ for each $i \in \{1, \ldots, t\}$.

Proof. For the sake of conciseness, define $w \coloneqq w_{\max}(\mathcal{K})$. Define $Z \coloneqq \{i \cdot \mu/n \in [0, w/\mu) \mid i \in \mathbb{Z}\}$ to be the set of offsets. For every $z \in Z$, $\mathcal{L}_z \coloneqq \{\{z + i \cdot w/\mu\} \times \mathbb{R} \mid i \in \mathbb{Z}\}$ is a set of uniformly spaced vertical grid lines, and let $\mathcal{K}_{\operatorname{rest},z} \subseteq \mathcal{K}$ be the set of k-shapes that are intersected by a line from \mathcal{L}_z . For simplicity we shall henceforth use $\mathcal{K}_{\operatorname{rest}}$ where z is clear from the context. Also let the set of k-shapes completely contained in $[z + i \cdot w/\mu, z + (i+1) \cdot w/\mu]$ belong to the set \mathcal{K}_{i+1} , and similarly let K_0 be the set of all k-shapes fully contained inside the strip [0, z]. Clearly each of the \mathcal{K}_i satisfies that $w_{\operatorname{range}}(\mathcal{K}_i) \leq w_{\max}(\mathcal{K})/\mu$ (proving property (*iii*)).

Clearly the \mathcal{K}_i sets are disjoint subsets of \mathcal{K} , so any solution (and in particular the optimal solution) to \mathcal{K} already stabs \mathcal{K}_i for all *i*. This gives us that $\text{OPT} \geq \sum_{\ell=0}^t \text{OPT}(\mathcal{K}_\ell)$ (proving property (*i*)).

We now need to show that there is a choice of $z \in Z$ such that $OPT(\mathcal{K}_{rest}) \leq 8\mu \cdot OPT(\mathcal{K})$. To this end, suppose \mathcal{S}_{OPT} is an optimal solution to the entire instance \mathcal{K} , and fix some choice of $z \in Z$. The idea is to collect all (parts of) segments from \mathcal{S}_{OPT} that are needed in order to stab \mathcal{K}_{rest} , and estimate their total cost.

Since no k-shape is wider than w, every k-shape in \mathcal{K}_{rest} must be entirely contained within the [-w, +w]-strip around some vertical line from \mathcal{L}_z . It is therefore sufficient to collect all intersections of segments in \mathcal{S}_{OPT} with such strips, obtaining a set $\mathcal{S}_{\mathcal{K}_{rest}}$ that

fully stabs $\mathcal{K}_{\text{rest}}$ (This differs from the case of stabbing rectangles, where only the segments directly intersected by \mathcal{L}_z are needed).

For estimating the cost of $\mathcal{S}_{\mathcal{K}_{rest}}$, denote by $\sigma_z(s)$ the total cost of the segments in $\mathcal{S}_{\mathcal{K}_{rest}}$ generated by $s \in \mathcal{S}_{OPT}$. Notice that $s = [x_1, x_2] \times \{y\}$ intersects the [-w, +w]-strip around a line $\ell = \{x_\ell\} \times \mathbb{R}$, if and only if $x_\ell \in [x_1 - w, x_2 + w]$. Counting the intersections of swith those strips is therefore equivalent to counting the intersections of a segment of length |s| + 2w with the lines themselves.

There are two mutually exclusive cases:

Case 1. $|s| + 2w \ge w/\mu$.

This implies that there is at least one intersection. Because the distance between the lines in \mathcal{L}_z is w/μ , there are at most $\frac{|s|+2w}{w/\mu} + 1$ intersections, each of which costs at most $\min\{|s|, 2w\}$. Therefore,

$$\begin{aligned} \sigma_z(s) &\leq \left(\frac{|s|+2w}{w/\mu}+1\right) \cdot \min\{|s|, 2w\} \\ &= \left(\frac{|s|+2w}{w/\mu}+\frac{w/\mu}{w/\mu}\right) \cdot \min\{|s|, 2w\} \\ &\leq \left(\frac{|s|+2w}{w/\mu}+\frac{|s|+2w}{w/\mu}\right) \cdot \min\{|s|, 2w\} \\ &= (2/w)\mu|s| \cdot \min\{|s|, 2w\} + 4\mu \cdot \min\{|s|, 2w\} \\ &\leq 4\mu|s|+4\mu|s| \\ &= 8\mu|s|. \end{aligned}$$

Case 2. $|s| + 2w < w/\mu$.

This implies that there is at most one intersection. To determine its probability, consider the set $\mathcal{L} := \bigcup_{z \in \mathbb{Z}} \mathcal{L}_z = \{k \cdot \mu/n \mid k \in \mathbb{Z}\}$. Since this is a disjoint union, each line $\ell \in \mathcal{L}$ can be associated with a unique value of z it was generated by, i.e., there exists a unique $z \in \mathbb{Z}$ such that $\ell \in \mathcal{L}_z$.

Now counting how many choices of z produce an intersection is equivalent to counting the intersections of \mathcal{L} with the (elongated) segment: Because $|s| + 2w < w/\mu$, it cannot intersect more than one line from the same \mathcal{L}_z , precluding double counting.

intersect more than one line from the same \mathcal{L}_z , precluding double counting. Hence there are at most $\frac{|s|+2w}{\mu/n} + 1$ choices of z producing an intersection. Division by the total number of choices $|Z| = wn/\mu^2$ yields

$$\begin{split} \mathbb{E}_{z}[\sigma_{z}(s)] &\leq \left(\frac{|s|+2w}{\mu/n}+1\right) \cdot \frac{1}{wn/\mu^{2}} \cdot \min\{|s|, 2w\} \\ &= \left(\frac{\mu}{w} \cdot |s|+2\mu + \frac{\mu^{2}}{wn}\right) \cdot \min\{|s|, 2w\} \\ &= \frac{\mu}{w} \cdot |s| \cdot \min\{|s|, 2w\} + 2\mu \cdot \min\{|s|, 2w\} + \frac{\mu^{2}}{wn} \cdot \min\{|s|, 2w\} \\ &\leq 2\mu|s| + 2\mu|s| + \frac{\mu^{2}}{wn} \cdot 2w \\ &\leq 2\mu|s| + 2\mu|s| + 2\mu \cdot \frac{\mu}{n} \\ &\leq 2\mu|s| + 2\mu|s| + 2\mu \cdot \frac{\mu}{n} \\ &\leq 2\mu|s| + 2\mu|s| + 2\mu|s| \qquad (\text{since } \frac{\mu}{n} \leq w_{\min}(\mathcal{K}) \leq |s|) \\ &= 6\mu|s|. \end{split}$$

The above argument shows that $\mathbb{E}[OPT(\mathcal{K}_{rest})] \leq O(\mu) \cdot OPT$, and hence there is at least one offset satisfying $OPT(\mathcal{K}_{rest}) \leq O(\mu) \cdot OPT$. For μ that is polynomial in n, there are only

a polynomial $\frac{w/\mu}{\mu/n} = \frac{wn}{\mu^2}$ number of possible offsets, and hence we can guess it in polynomial time.

A.4 Proof of Lemma 6

▶ Lemma 6. If $OPT(\mathcal{K}_i) > w_{max}(\mathcal{K})/\mu^2$ then in polynomial time we can guess a value $h \in \mathbb{N}$ and a corresponding line segment $s = [a, b] \times \{h\}$ such that each connected component C of $S_i \setminus s$ contains segments from $OPT(\mathcal{K}_i)$ whose total cost is at least $OPT(\mathcal{K}_i)/2 - w_{max}(\mathcal{K})/\mu$.

Proof. Since we use only horizontal segments to stab k-shapes w.l.o.g. (by scaling along the y direction) we can assume that the at most 2kn points describing the instance occupy consecutive integral y-coordinates, starting at y = 0.

Consider the segments from $OPT(\mathcal{K}_i)$. Starting from y = 0 and going up, we can start counting the cumulative cost of segments in OPT. Let h be the y-coordinate at which this cumulative cost crosses $OPT(\mathcal{K}_i)/2$, and $s = [a, b] \times \{h\}$ be the corresponding segment. Since the width of S_i is at most $w_{\max}(\mathcal{K})/\mu$, no segment in $OPT(\mathcal{K}_i)$ is wider than $w_{\max}(\mathcal{K})/\mu$. From this we can infer that cost of segments from $OPT(\mathcal{K}_i)$, below (and similarly, above) the segment s should have been at least $OPT(\mathcal{K}_i)/2 - w_{\max}(\mathcal{K})/\mu$.

Since there are only a polynomial 2kn number of possible y-coordinates, we can guess this value h in polynomial time by enumeration.

A.5 Proof of Lemma 7

▶ Lemma 7. If all guesses for the balanced horizontal cut are correct, then their total cost is bounded by $3\mu \cdot \text{OPT}(\mathcal{K}_i)$.

Proof. After our sequence of (correctly guessed) balanced horizontal cuts, let us assume that there are t connected components, with cost at least $w_{\max}(\mathcal{K}_i)/2\mu^2 - w_{\max}(\mathcal{K})/\mu$. This can happen only if there were t-1 such cuts applied. If we charge the cost of every cut s to the cost of segments of $\text{OPT}(\mathcal{K}_i)$ within a cell C, we get

$$\frac{|s|}{\operatorname{OPT}(C)} = \frac{w_{\max}(\mathcal{K})/\mu}{w_{\max}(\mathcal{K})/2\mu^2 - w_{\max}(\mathcal{K})/\mu} = \frac{2\mu}{1 - 2\mu} \le 3\mu.$$

Where the last inequality follows under the assumption of $\mu \leq \varepsilon < 1/3$.

Summing over all such horizontal cuts, we get the total cost to be at most $3\mu \cdot w_{\max}(\mathcal{K}_i)$.

A.6 Proof of Lemma 8

▶ Lemma 8. If all our guesses are correct, then the total cost for the selected line segments due to Lemmas 5 and Lemmas 7 is bounded by $O(\mu \cdot \text{OPT})$. Also, the total number of (combinations of) guesses is bounded by $n^{O(\varepsilon^{-3} \log^2 n)}$.

Proof. The bound on the cost of segments directly follows from Lemma 5 and Lemma 7. The above Lemmas also guarantee that the number of guesses for applying them is polynomial. The overall number of guesses is $n^{O(\varepsilon^{-3} \log^2 n)}$ because while guessing the segments within a cell (formed as a result of Lemmas 5 and Lemmas 7) there are only a polynomial number of (combinatorially distinct) possible segments, and there are at most $\varepsilon^{-3} \log^2 n$ such segments within each cell.

B Missing proofs and details from Section 4

B.1 Constant Factor Approximation

In this subsection, we give a simple $O(1/\delta)$ -approximation algorithm for k-STABBING when for each k-shape the widths of any two of its rectangles differ by at most a factor of $1/\delta$. Given such an instance of k-STABBING and let OPT denote its optimal solution. We create an instance of STABBING of rectangles as follows. For each given k-shape K, we take the rectangle of smallest width and height that contains K. We add all these rectangles to our constructed instance of STABBING for rectangles. Let OPT' denote the optimal solution to that instance. On this instance, we apply the known PTAS for STABBING for rectangles [28].

We observe that $\text{OPT}' \leq O(\text{OPT}/\delta)$ since we can simply take each segment in OPT and extend it by a factor of $1/\delta$ in each direction. Due to our assumption about the widths of the input rectangles, this yields a feasible solution to our initially given instance to k-STABBING. Therefore, the solution computed by our PTAS yields a solution with cost at most $O(\text{OPT}/\delta)$, and hence a $O(1/\delta)$ -approximation.

B.2 Proof of Lemma 10, Preprocessing Step

We note here that the discretization steps here are similar to the case of rectangles as done by Khan, Subramanian, and Wiese [28].

▶ Lemma 10. Let α be a constant for which k-STABBING admits an α -approximate algorithm and let $\varepsilon > 0$ with $\varepsilon < 1/3$. In polynomial time we can compute a new instance of k-STABBING, in which each $K \in \mathcal{K}$ satisfies,

(i) $\frac{\alpha\varepsilon}{n} < w_{\min}(K) \le w_{\max}(K) \le \alpha$,

(ii) all points defining K are discrete,

(iii) K lies within a bounding box of $[0, \alpha n] \times [0, (k+1)n]$,

and this new instance admits a solution of cost at most $(1 + O(\varepsilon)) \cdot \text{OPT}$ with each segment in the solution being discrete, and having length at most α/ε .

Proof. Using the α -approximation algorithm, obtain a solution to \mathcal{K} and determine its cost $C \in [OPT(\mathcal{K}), \alpha OPT(\mathcal{K})]$. Consequently, $OPT(\mathcal{K}) \in [\frac{1}{\alpha}C, C]$.

Scaling each k-shape in \mathcal{K} along the x-axis by a factor of $\beta \coloneqq (1-2\varepsilon)\frac{\alpha}{C}$ yields an instance \mathcal{K}_{β} with $\operatorname{OPT}(\mathcal{K}_{\beta}) = \beta \cdot \operatorname{OPT}(\mathcal{K}) \in [(1-2\varepsilon), (1-2\varepsilon)\alpha]$. Now, greedily stabbing all k-shapes $K \in \mathcal{K}_{\beta}$ with $w_{\min}(K) \leq \frac{\alpha\varepsilon}{n}$ requires segments of total length at most $\alpha\varepsilon$. This accounts for a factor

$$\frac{\text{SOL}}{\text{OPT}} \le \frac{\text{OPT} + \alpha\varepsilon}{\text{OPT}} \le 1 + \frac{\alpha\varepsilon}{1 - 2\varepsilon} \le 1 + \varepsilon \cdot \frac{\alpha}{1 - 2(1/3)} = 1 + O(\varepsilon),$$

increase in the cost of the solution.

Now we try to discretize the x-coordinates of all k-shapes. Each k-shape consists of up to k rectangular parts. Extend each such part on both sides to make their x-coordinates align with the next nearest multiple of ε^d . Since this involves extension by at most $2\varepsilon^d < 2\varepsilon/n$ to the width of every rectangular section, the total cost of the solution increases by at most a factor of $1 + 2\varepsilon$.

In other words we can also say that the cost of an optimal solution goes up, to $OPT(\mathcal{K}_{\beta}) \in [1, \alpha]$. Now we notice that any parts R of k-shapes in \mathcal{K}_{β} with $w(R) > \alpha$ can be discarded, since an optimal solution cannot stab them. Hence, the above steps ensure that $\frac{\alpha \varepsilon}{n} < w_{\min}(\mathcal{K}) \leq w_{\max}(\mathcal{K}) \leq \alpha$ (proving property (i)), and that each x-coordinate of any k-shape is discretized.

Since we stab the k-shapes using horizontal lines, we can stretch the a k-shape vertically without affecting the solution cost. Since each k-shape has at most k+1 distinct y-coordinates, there are at most (k+1)n distinct y-coordinates in the problem instance, and we can stretch the instance in such a manner that these y-coordinates are consecutive integers between 0 to (k+1)n, ensuring that all y-coordinates are also discretized (proving property (ii)).

It can be assumed w.l.o.g. that $w_{\text{range}}(\mathcal{K}_{\beta}) \leq \alpha n$; otherwise, \mathcal{K}_{β} could be partitioned into independent sub-instances to be solved separately. Similarly since the *y*-coordinates are also already shown to lie between 0 and 2kn, we have also shown property (*iii*).

We now consider the final property. Let OPT' be the optimal solution to the instance obtained after applying the three steps above. Then the cost of OPT' can exceed the cost of the original optimal solution OPT by at most a factor of $1 + O(\varepsilon)$. Consider any horizontal segment $\ell \in \text{OPT'}$ that is longer than α/ε . From the left endpoint, we divide the segment into consecutive smaller segments of length $\alpha/\varepsilon - 2\alpha$ each, with one potential last piece being smaller than $\alpha/\varepsilon - 2\alpha$. Now, for each smaller segment, we extend it on both sides in such a way that it completely stabs the k-shapesthat it intersects (i.e., that it intersects before the extension). Since the maximum width of a k-shape α , we extend each such segment by at most 2α units. We denote by $|\ell|$ the length of ℓ and conclude that we increase the length of ℓ by at most a factor of

$$\left(\left| \ell \right| + \left\lceil \frac{\left| \ell \right|}{\alpha/\varepsilon - 2\alpha} \right\rceil \cdot 2\alpha \right) \cdot \frac{1}{\left| \ell \right|} \le 1 + \frac{2\alpha}{\left| \ell \right|} \cdot \left\lceil \frac{\varepsilon \left| \ell \right|}{\alpha - 2\alpha\varepsilon} \right\rceil$$

$$\le 1 + \frac{2\alpha}{\left| \ell \right|} \cdot \left(1 + \frac{\varepsilon \left| \ell \right|}{\alpha - 2\alpha\varepsilon} \right)$$

$$\le 1 + 2\varepsilon + 2\varepsilon \cdot \frac{\alpha}{\alpha - 2\alpha\varepsilon}$$

$$\le 1 + 2\varepsilon + 2\varepsilon \cdot \frac{1}{1 - 2/3}$$

$$(since \ \varepsilon < 1/3)$$

$$\le 1 + 8\varepsilon.$$

This completes the proof of the lemma.

B.3 Proof of Lemma 11

▶ Lemma 11. The running time of the above dynamic program is $(kn/\varepsilon)^{O(1/\varepsilon^3)}$.

Proof. We know that the DP always picks the sequence of operations that gives a result of minimum cost. Since there is a sequence of operations that produces a solution of cost $(1 + \varepsilon)$ OPT, the DP returns a solution of cost at most that.

Now let us consider the running time of the algorithm. Since a DP problem is defined on a discrete rectangular cell, there are at most

$$\frac{\alpha n}{\varepsilon^d} \times (k+1)n \le \alpha (k+1)\varepsilon^{-3}n^3 \qquad (\text{since } \varepsilon^3/n < \varepsilon^d \le \varepsilon^2/n)$$

possibilities for a corner vertex of a rectangle, and hence $\binom{\alpha(k+1)\varepsilon^{-3}n^3}{2} = O(k^2n^6/\varepsilon^6)$ possible rectangles.

Similarly, the subproblem definition also includes a set of segments \mathcal{L} of size at most $3\varepsilon^{-3}$. Since the segments are discrete we can count the number of horizontal segments by picking two points (corresponding to the starting and ending points) from the available discrete

points on a horizontal line, and then fixing its y-coordinate.

$$\binom{\alpha n/\varepsilon^d}{2} \times (k+1)n \le \binom{\alpha n^2/\varepsilon^3}{2} \times (k+1)n \le \frac{\alpha^2 (k+1)n^5}{\varepsilon^6}$$

that is, $O(kn^5/\varepsilon^6)$ possible segments. So there are at most $(kn/\varepsilon)^{O(1/\varepsilon^3)}$ subsets of segments \mathcal{L} of size at most $3\varepsilon^{-3}$, and at most $(kn/\varepsilon)^{O(1/\varepsilon^3)}$ valid DP-cells.

For each DP-cell, we have to consider all possible candidate solutions and select the minimum. There are at most $\alpha n/\varepsilon^d = O(n^2/\varepsilon^3)$ possible *line operations* and $(kn/\varepsilon)^{O(1/\varepsilon^3)}$ possible add operations. Note that if the DP performs a trivial operation, then there is no choice to make here, but the trivial operation is selected automatically.

Hence, the total number of possible operations for a given DP-cell is $(kn/\varepsilon)^{O(1/\varepsilon^3)}$. For each *line operation* we call the O(1)-approximation algorithm which is also runs in polynomial time [9]. Since we have $(kn/\varepsilon)^{O(1/\varepsilon^3)}$ DP-cells, our overall running time is bounded by $(kn/\varepsilon)^{O(1/\varepsilon^3)}$.

B.4 Proof of Lemma 12

▶ Lemma 12. For any value of our offset, by losing a factor of $1+O(\varepsilon)$ in our approximation ratio, we can assume that each line segment $\ell \in \text{OPT}$ is well-aligned.

Proof. A segment ℓ in some level j will be of length in $(\alpha \varepsilon^j, \alpha \varepsilon^{j-1}]$. To align it to a grid line of level j + 3 we would need to extend it by at most $\alpha \varepsilon^{j+1}$ on each side. The new segment ℓ' thus obtained is of length

$$|\ell'| \le |\ell| + 2\alpha\varepsilon^{j+1} \le |\ell| \left(1 + \frac{2\alpha\varepsilon^{j+1}}{|\ell|}\right) < |\ell| \left(1 + \frac{2\alpha\varepsilon^{j+1}}{\alpha\varepsilon^j}\right) = |\ell| \cdot (1 + 2\varepsilon).$$

Therefore, the sum of weights over all the segments in OPT is

$$\sum_{\ell \in \text{OPT}} |\ell'| \le \sum_{\ell \in \text{OPT}} |\ell| \cdot (1 + 2\varepsilon) = (1 + 2\varepsilon) \cdot \text{OPT.}$$

B.5 Approximation Factor: Proof of Lemma 14

▶ Lemma 14. There is a discrete value for the offset $r \in \{0, \varepsilon^d, 2\varepsilon^d, \dots, \varepsilon^{-2}\}$ such that the described sequence of operations produces a solution of cost at most $(1 + O(\varepsilon))$ OPT.

Proof. In the described sequence of operations, some add operations are applied on segments from OPT (or their parts), and hence the cost across all such add operations is at most cost of OPT. Similarly, all trivial operations are applied on segments that were 'added' before, and hence their cost is also already accounted for. So we are left with analyzing the cost of stabbing the rectangles which are intersected by the lines along which we apply the line operations, and the add operations which are not which are not applied on a segment from OPT (that is, the ones that are used to partition a vertical strip). We claim that for a discretized random offset $r \in \{0, \varepsilon^d, 2\varepsilon^d, \dots, \alpha\varepsilon^{-2}\}$, the expected cost is $O(\varepsilon \cdot \text{OPT})$, which would give us the required result.

Let us first consider any add operation of level j that is applied to a horizontal line ℓ (that is not in OPT). We do such an operation only after accounting for ε^{-3} segments from OPT of level j, that is, segments of cost at least $\varepsilon^{-3} \cdot \alpha \varepsilon^j = \alpha \varepsilon^{j-3}$. Since a segment of width $\alpha \varepsilon^{j-2}$ (width of strip) is sufficient to stab all the k-shapes stabled by ℓ , we see that this horizontal segment only takes ε times the cost of the segments in OPT that we have

already accounted for. We charge the cost of adding this segment to the solution, to the ε^{-3} segments from OPT that were counted before adding it. Since any segment in OPT gets charged only once, we can infer that the cost of such add operations is at most $2\varepsilon \cdot \text{OPT}$.

Now, let us consider the *line operations* applied to vertical grid lines. Consider a grid line ℓ of level j. We wish to bound the cost of stabbing all the rectangles in \mathcal{R}_{ℓ} intersected by grid lines, over all levels j. Let $OPT_{\mathcal{R}_{\ell}}$ be the set of minimum cost that stabs all segments in \mathcal{R}_{ℓ} , and let $OPT_{\mathcal{K}_{\ell}}$ be the set of segments from OPT that stab the corresponding k-shapes in \mathcal{K}_{ℓ} . From Appendix B.1, we know that $OPT_{\mathcal{R}_{\ell}} \leq 1/\delta \cdot OPT_{\mathcal{K}_{\ell}}$. So instead of bounding the cost of stabbing \mathcal{R}_{ℓ} , we instead focus on bounding the cost of stabbing \mathcal{K}_{ℓ} .

A horizontal line segment $\ell' \in \text{OPT}$ of some level $j' \geq j$ stabs a k-shape in \mathcal{K}_{ℓ} only if the distance between ℓ and ℓ' is at most $\alpha \varepsilon^{j-1}$. So we need to bound the cost, over all levels j, of line segments of level j in OPT (call this set OPT_j) intersected or close to grid lines of level j. For a horizontal segment $\ell' \in \text{OPT}_j$, let $I_{\ell'}$ be the indicator variable representing the event that ℓ' is within distance $\alpha \varepsilon^{j-1}$ of ℓ . Now, since any two consecutive grid lines of level j are separated by $\alpha \varepsilon^{j-2}$, there are $(\alpha \varepsilon^{j-2})/\varepsilon^d$ possible shifts for these grid lines due to our offset, and each of these shifts has the same probability. Similarly, $(2\alpha \varepsilon^{j-1}/\varepsilon^d) + 1$ of these offsets would allow ℓ' to be within $\alpha \varepsilon^{j-1}$ distance of ℓ . So, if we take a random discrete offset $r \in \{0, \varepsilon^d, 2\varepsilon^d, \dots, \varepsilon^{-2}\}$, we have that

$$\mathbb{E}[I_{\ell'}] \le \frac{(2\alpha\varepsilon^{j-1}/\varepsilon^d) + 1}{\alpha\varepsilon^{j-2}/\varepsilon^d} = 2\varepsilon + \frac{\varepsilon^{2+(d-j)}}{\alpha} \le 3\varepsilon. \qquad (\text{since } j \in \{0, 1, \dots, d-1\})$$

With the expectation computed above, we can upper bound the expected cost of segments in $OPT_{\mathcal{K}_{\ell}}$ as:

$$\mathbb{E}\left[\sum_{j}\sum_{\ell\in \mathrm{OPT}_{j}}I_{\ell}\cdot|\ell|\right] = \sum_{j}\sum_{\ell\in \mathrm{OPT}_{j}}\mathbb{E}\left[I_{\ell}\cdot|\ell|\right]$$
$$= \sum_{j}\sum_{\ell\in \mathrm{OPT}_{j}}|\ell|\cdot\mathbb{E}[I_{\ell}]$$
$$\leq \sum_{j}\sum_{\ell\in \mathrm{OPT}_{j}}|\ell|\cdot(3\varepsilon)$$
$$= 3\varepsilon\cdot\mathrm{OPT}$$

Since $\operatorname{OPT}_{\mathcal{R}_{\ell}} \leq 1/\delta \cdot \operatorname{OPT}_{K_{\ell}}$, we get that $\mathbb{E}[\operatorname{OPT}_{\mathcal{R}_{\ell}}] \leq 3\varepsilon/\delta \cdot \operatorname{OPT}$. And since we use a PTAS (which, let's say has an approximation factor of $(1 + \varepsilon')$) for computing the cost of $\operatorname{OPT}_{\mathcal{R}_{\ell}}$, the solution returned by our algorithm takes an additional cost of $\frac{3\varepsilon(1+\varepsilon')}{\delta} \cdot \operatorname{OPT}$.

Now we prove our main theorem.

Proof. We gave a DP algorithm in Section 4 which was shown to have the required running time in Lemma 11. Further in Lemma 14 we showed the correctness and that the solution computed by the DP is actually a $(1 + O(\varepsilon))$ -approximation of the solution.

C Missing proofs and details from Section 5

C.1 Proof of Proposition 17

▶ Proposition 17. Each 2-shape satisfies the hourglass property.

Proof. If k = 2 then there is no value $i \in \{2, ..., k - 1\}$ and hence the hourglass property is trivially satisfied.

C.2 Proof of Lemma 18

▶ Lemma 18. For each $\gamma \in \mathbb{N}$, the given instance of vertex cover instance has a solution of size γ if and only if the corresponding k-STABBING instance has a solution of cost γ .

Proof. We first show that if there is a vertex cover of size γ then there is a solution of cost γ solution to our instance of k-STABBING. Given a solution $S = \{v_1, v_2, \ldots, v_{\gamma}\}$ to the vertex cover instance, construct a solution to the stabbing instance as follows: for each $v_i \in S$, stab the corresponding s_i along its top edge by a segment of length one. Clearly the cost of this set of segments is γ . Now we notice that every k-shape $r_{i,j}$ corresponds to an edge $e(v_i, v_j)$ in the graph. Since this edge has been covered by one of its adjacent vertices $v_i \in S$, $r_{i,j}$ is also stabbed by the segment that stabs s_i . We know that every edge of the graph is covered by some vertex in S, and hence every k-shape in the instance is also stabbed in the solution we constructed.

Next, we argue that a solution of $\cot \gamma$ to our instance of yields a solution to vertex cover of size at most γ . Consider any solution to the stabbing instance of $\cot \gamma$. We can assume that there are no segments of length greater than one in this solution, since any segment of length at least n + 1, can be broken down into at most n segments of length 1 stabbing the same set of k-shapes, but along their bordering squares; and segments of length in the range (1, n + 1) can stab only one k-shape, and hence be shortened to length one. Further segments in any solution can also not be of length less than one, since such a segment cannot stab any k-shape. Hence we conclude that all segments in the solution are of length one, and by extension that they stab any k-shape along one of its bordering squares.

Now we construct a vertex cover solution by picking the vertices v_i , that correspond to any square s_i that has been stabbed by the given (or modified as mentioned above) k-STABBING solution. Note that every k-shape is stabbed by the given solution, and hence a vertex adjacent to every edge in the vertex cover instance has been picked by us. This shows that the selected set is in fact a valid vertex set, and is of size at most γ .

C.3 Proof of Lemma 20

▶ Lemma 20. For each $\gamma \in \mathbb{N}$, the given instance of hitting set has a solution of size γ if and only if the constructed k-STABBING instance has a solution of cost γ .

Proof. Given a solution $H = \{v_1, v_2, \ldots, v_{\gamma}\}$ to the hitting set instance, construct a solution to the stabbing instance as follows: for each $v_i \in H$, stab the corresponding s_i along its top edge by a segment of length one. Clearly the cost of this set of segments is γ . Now we notice that every k-shape r_{S_i} corresponds to an set $S_i \in \mathcal{F}$. Since this set has been hit by a vertex, say $v_i \in H$, r_{S_i} is also stabbed by the segment that stabs s_i . We know that every set of the family \mathcal{F} is covered by some element in H, and hence every k-shape in the instance is also stabbed in the solution we constructed.

Similar to the argument in proof of Lemma 18, we can modify the solution without increasing its cost so that every segment is of length one. Now we construct a hitting set solution by picking the vertices v_i , that correspond to any square s_i that has been stabbed by the given (or modified as mentioned above) k-STABBING solution. Note that every k-shape is stabbed by the given solution, and hence an element from each set in \mathcal{F} has been picked by us. This shows that the selected set is in fact a valid hitting set, and is of size at most γ .

C.4 Proof of Lemma 21

Lemma 21. In polynomial time, we can construct a set C of line segments with the following properties:

- \square C contains $O((kn)^3)$ segments,
- C contains no redundant segments, where a segment is redundant if it stabs exactly the same k-shapes as another segment, or no k-shapes at all, and
- \blacksquare \mathcal{K} admits an optimal solution using only the segments from \mathcal{C} .

Proof. Let S_{OPT} be the set of line segments of an optimal solution to \mathcal{K} . Any segment $s = [x^{\ell}, x^{r}] \times \{y\} \in S_{\text{OPT}}$ must start and end on vertical boundaries of k-shapes. Furthermore, s can be translated along the y-direction to the nearest horizontal boundary of some part of a k-shape without changing the set of k-shapes it stabs. Therefore, it is sufficient to consider O(kn) choices each for x^{ℓ}, x^{r} , and y, obtaining $O((kn)^{3})$ combinations.

Candidate segments not stabbing any k-shapes can be discarded. Suppose that there are multiple candidate segments stabbing the same set of k-shapes. Any optimal solution uses at most one of them, specifically a shortest one. Discard the rest. This yields a set C with no redundant segments.