

A note on vertex Turán problems in the Kneser cube

Dániel Gerbner* Balázs Patkós†

Abstract

The Kneser cube Kn_n has vertex set $2^{[n]}$ and two vertices F, F' are joined by an edge if and only if $F \cap F' = \emptyset$. For a fixed graph G , we are interested in the most number $\text{vex}(n, G)$ of vertices of Kn_n that span a G -free subgraph in Kn_n . We show that the asymptotics of $\text{vex}(n, G)$ is $(1 + o(1))2^{n-1}$ for bipartite G and $(1 - o(1))2^n$ for graphs with chromatic number at least 3. We also obtain results on the order of magnitude of $2^{n-1} - \text{vex}(n, G)$ and $2^n - \text{vex}(n, G)$ in these two cases. In the case of bipartite G , we relate this problem to instances of the forbidden subposet problem.

1 Introduction

Graph Turán problems are among the most studied problems in extremal graph theory [12]. They ask for how large a subgraph F of a given host graph H can be if F does not contain G as a subgraph, i.e. the input of the problem is (G, H) . Largeness is usually measured by the number of edges, but there are other parameters like spectral radius. Recently there have been interest [2, 14, 19, 23] in *vertex Turán problems*, where one is interested in the most number of vertices that a vertex subset U of H can have such that the induced subgraph $H[U]$ is G -free. One particular host graph where this problem has been studied is the *Kneser graph* $Kn(n, k)$ with vertex set $\binom{[n]}{k} := \{F \subseteq [n] : |F| = k\}$, where $[n] = \{1, 2, \dots, n\}$, and F, F' are connected by an edge if and only if F and F' are disjoint. Vertex Turán problems in the Kneser graph are usually phrased as extremal finite set theory problems. The first such result is the Erdős-Ko-Rado theorem [7], which determines the largest family of pairwise intersecting k -sets. In other words, the Erdős-Ko-Rado theorem determines the largest independent set, i.e., the largest K_2 -free vertex subset of $Kn(n, k)$. The Erdős matching problem [6] deals with the largest family without $r + 1$ pairwise disjoint sets, i.e., the largest set of vertices without K_{r+1} . For results on this problem, see [10] and the citations in it. Gerbner, Lemons, Palmer, Patkós and Szécsi [13] studied the largest ℓ -almost intersecting families, where each set is disjoint with at most ℓ other sets, i.e., the largest $S_{\ell+1}$ -free sets of vertices. Katona and Nagy [21] studied (s, t) -union intersecting families, which have the property that the union of any s sets intersect the union of any t sets, i.e., $K_{s,t}$ -free sets. Alishahi and Taherkhani [1] united these notions and determined for every F the largest F -free set of vertices in the Kneser graph $Kn(n, k)$, provided n is sufficiently large. More results in this topic can be found in [15, 25].

*HUN-REN Alfréd Rényi Institute of Mathematics, Budapest, 1053, Reáltanoda utca 13-15, e-mail: gerbner@renyi.hu, Research supported by NKFIH under grants FK132060 and KKP-133819.

†HUN-REN Alfréd Rényi Institute of Mathematics, Budapest, 1053, Reáltanoda utca 13-15, e-mail: patkos@renyi.hu Research supported by NKFIH under grant FK132060.

In each of the above specific problems on set families, the non-uniform version has also been studied, but not for the general problem. Here we address this. Our host graph is the Kneser cube Kn_n with vertex set $2^{[n]} := \{F \subseteq [n]\}$, where F, F' are connected by an edge if and only if F and F' are disjoint. We denote by $\text{vex}(n, G)$ the most number of vertices of Kn_n that induce a G -free subgraph in Kn_n . Note that $\text{vex}(n, G) \geq 2^{n-1}$ if G contains an edge, as shown by the family of sets each containing a fixed element. Therefore, the order of magnitude of $\text{vex}(n, G)$ is known, and the more interesting question may be to determine or bound the order of magnitude of $2^n - \text{vex}(n, G)$ or $\text{vex}(n, G) - 2^{n-1}$.

If $G = K_{r+1}$, then the problem is equivalent to the non-uniform Erdős matching problem. Kleitman [22] obtained a general and very nice upper bound on $\text{vex}(n, K_{r+1})$. Sharpenings for special cases of r and the residue class of n modulo $r + 1$ have been obtained in (among others) [8, 9, 11]. A simple lower bound is obtained by taking each set of size larger than $n/(r + 1)$; all the above results show upper bounds close to this.

Gerbner, Lemons, Palmer, Patkós and Szécsi [13] determined $\text{vex}(n, S_2)$ and $\text{vex}(n, S_3)$ exactly. Katona and Nagy [21] also determined $\text{vex}(n, C_4)$ exactly and for $t \geq 4$ they showed that $\text{vex}(n, S_t) - 2^{n-1} = (\frac{1}{2} + o(1))\binom{n}{\lfloor n/2 \rfloor}$. For $t \geq s \geq 2$, Katona and Nagy showed that $\text{vex}(n, K_{s,t}) - 2^{n-1} = (1 + o(1))\binom{n}{\lfloor n/2 \rfloor}$.

We shall consider all graphs G and the corresponding vertex Turán problem in the Kneser cube. It turns out that there are three major graph classes with respect to the value of $\text{vex}(n, G)$: matchings, other bipartite graphs and graphs with chromatic number at least 3. The following proposition about matchings is rather a simple exercise. We include its proof for completeness in Section 2.1, the interested reader is welcome to figure out the proof on its own. M_k denotes the matching with k edges.

Proposition 1.1. $\text{vex}(n, M_{k+1}) = 2^{n-1} + k$.

The situation is somewhat different for bipartite graphs other than matchings. The asymptotics of $\text{vex}(n, G)$ is still 2^{n-1} , but the second order term is different. Our next result determines the order of magnitude of $2^{n-1} - \text{vex}(n, G)$.

Theorem 1.2. *For any bipartite graph G that is not a matching, we have*

$$2^{n-1} + \left(\frac{1}{2} - o(1)\right) \binom{n}{n/2} \leq \text{vex}(n, G) \leq 2^{n-1} + (1 + o(1)) \binom{n}{\lfloor n/2 \rfloor}.$$

Both bounds of Theorem 1.2 can be tight. In Section 2.1, we shall relate the problem of finding $\text{vex}(n, G)$ for bipartite graphs G to a forbidden subposet problem (see the definitions there). Whenever the corresponding problem is solved, we will be able to determine $\text{vex}(n, G)$ up to a $o(\binom{n}{n/2})$ term and whenever that forbidden subposet problem is settled, the coefficient of $\binom{n}{n/2}$ is always either 1/2 or 1.

Next, we shall turn our attention to non-bipartite graphs. We introduce the following notation: $\binom{n}{\leq m} := \sum_{i=0}^m \binom{n}{i}$ and $\binom{n}{\geq m} := \sum_{i=m}^n \binom{n}{i}$. The following simple proposition shows that if the chromatic number of G is at least 3, then almost all vertices can be selected to obtain a G -free induced subgraph of Kn_n .

Proposition 1.3. *For any graph G with chromatic number at least 3, let $\ell = \ell(G)$ be the length of the shortest odd cycle of G with $\ell = 2k + 1$. Then $2^n - \text{vex}(n, G) \leq \binom{n}{\leq \lfloor \frac{kn}{2k+1} \rfloor}$.*

Our final theorem establishes that for odd cycles, the correct order of magnitude of $2^n - \text{vex}(n, C_{2k+1})$ is given by the bound of Proposition 1.3. In the next theorem and later, \log denotes the natural logarithm.

Theorem 1.4. *For any positive integer k , we have $\text{vex}(n, C_{2k+1}) \leq \binom{n}{\geq \lceil \frac{kn}{2k+1} \rceil - \lfloor 2(k+1) \log(2k) \rfloor}$. Therefore, $2^n - \text{vex}(n, C_{2k+1}) = \Theta_k\left(\binom{n}{\leq \lfloor \frac{kn}{2k+1} \rfloor}\right)$.*

2 Proofs

2.1 Bipartite graphs

We start with the proof of Proposition 1.1.

Proof of Proposition 1.1. Let $\mathcal{F} \subseteq 2^{[n]}$ be such that $\text{Kn}_n[\mathcal{F}]$ is M_{k+1} -free. To avoid a copy of M_{k+1} , out of every pair of complement sets at most k can exist with both sets belonging to \mathcal{F} . This shows the upper bound. And for any family \mathcal{F} containing exactly one of $F, [n] \setminus F$ for $2^{n-1} - k$ pairs of complements and both of them for the remaining k pairs, the induced subgraph $\text{Kn}_n[\mathcal{F}]$ is M_k with isolated vertices. \square

To prove Theorem 1.2, we need some auxiliary definitions and results. We say that a family \mathcal{G} is a copy of a poset $P = (P, \prec)$ if there exists a bijection $b : P \rightarrow \mathcal{G}$ such that $p \prec q$ implies $b(p) \subseteq b(q)$ for any $p, q \in P$. A family \mathcal{F} is P -free if it does not contain any copy of P . The problem of finding the size of the largest P -free family $\mathcal{F} \subseteq 2^{[n]}$, denoted by $\text{La}(n, P)$, is called the forbidden subposet problem (for a survey see [17] and Chapter 7 of [16]). If set \mathcal{P} of posets is forbidden, then $\text{La}(n, \mathcal{P})$ denotes the maximum size of a family $\mathcal{F} \subseteq 2^{[n]}$ that is P -free for all $P \in \mathcal{P}$. It follows from an old result of Erdős [5] that for any poset P , we have $\text{La}(n, P) \leq (|P| - 1) \cdot \binom{n}{\lfloor n/2 \rfloor}$, but there exist stronger results (and used to be a stronger conjecture - see at the end of this subsection). We will use a special case of the following theorem of Bukh [3]. The *Hasse diagram* of a poset (P, \prec) is the oriented graph with vertex set P and \vec{pq} is an arc if $p \prec q$ and there is no $z \in P$ with $p \prec z \prec q$. A poset is called a *tree poset* if its Hasse diagram is a tree. The height $h(P)$ of a poset P is the size of its longest chain.

Theorem 2.1 (Bukh [3]). *For any tree poset T , we have $\text{La}(n, T) = (h(T) + o(1)) \binom{n}{\lfloor n/2 \rfloor}$.*

Let $G = (A, B, E)$ be a bipartite graph. We define $P_{G,A}$ to be the poset with Hasse diagram G and each edge oriented towards its end-vertex in A . The poset $P_{G,B}$ is the *dual poset* of $P_{G,A}$, that is, the poset with the same Hasse diagram but with all edges oriented in the opposite direction.

Lemma 2.2. *Let G be a connected bipartite graph, and let $\mathcal{F} \subseteq 2^{[n]}$ be such that $\text{Kn}_n[\mathcal{F}]$ is G -free. Then $\mathcal{F}_{\text{sym}} := \mathcal{F} \cap \mathcal{F}^c \cap \bigcup_{i=-n^{2/3}}^{n^{2/3}} \binom{[n]}{\lfloor n/2 \rfloor + i}$ is $P_{G,A}$ -free if n is large enough.*

Proof. Suppose towards a contradiction that \mathcal{F}_{sym} contains a copy \mathcal{H} of $P_{G,A}$. Let \mathcal{H}_A and \mathcal{H}_B be the parts of \mathcal{H} that correspond to vertices of A and B , respectively. First we show that we cannot have $F \in \mathcal{H}_A$ and $F^c \in \mathcal{H}_B$. This is because G is connected, and therefore \mathcal{H} is connected. As a result, there is a path $v_1 v_2 \dots v_{2k}$ in \mathcal{H} with $v_1 = F$, $v_{2k} = F^c$ and

$2k \leq |V(G)|$. Observe that the Hamming distance between v_i and the complement of v_{i+1} is at most $2n^{2/3}$ for any i . Therefore, the Hamming distance between v_i and v_{i+2} is at most $4n^{2/3}$. This implies that the Hamming distance between v_1 and v_{2k-1} is at most $2(2k-2)n^{2/3}$, thus v_{2k-1} is not disjoint from the complement of v_1 , a contradiction.

Consider the family $\mathcal{H}_B \cup \mathcal{H}_A^c$. If there is a relation, i.e., a directed edge between $H \in \mathcal{H}_B$ and H' in $G_{P,A}$, then the edge goes to H' , i.e., $H \subset H'$. This implies $H \cap H'^c = \emptyset$, and thus H and H'^c are connected by an edge. So by definition of $P_{G,A}$, the family $\mathcal{H}_B \cup \mathcal{H}_A^c$ induces a copy of G in Kn_n , a contradiction completing the proof. \square

Now we are ready to prove Theorem 1.2.

Proof of Theorem 1.2. Let G be any bipartite graph that has at least one vertex of degree at least 2, i.e. not a matching. First we show the lower bound $\text{vex}(n, G) \geq 2^{n-1} + \frac{1}{2} \binom{n}{n/2}$ if n is even, and $\text{vex}(n, G) \geq 2^{n-1} + \binom{n-1}{\lfloor n/2 \rfloor - 1}$ if n is odd. In the former case the family of all the sets of size at least $n/2$ induce a matching in Kn_n , and the same holds in the odd n case for the family consisting of all the sets of size larger than $n/2$ together with the sets of size $\lfloor n/2 \rfloor$ sharing a common element.

To see the upper bound, let $\mathcal{F} \subseteq 2^{[n]}$ be a family with $Kn_n[\mathcal{F}]$ being G -free. Observe first that by the well-known Chernoff bound, $|\binom{[n]}{\leq n/2 - n^{2/3}} \cup \binom{[n]}{\geq n/2 + n^{2/3}}| = o(\binom{n}{n/2})$. We claim that $\mathcal{F}_{\text{sym}} := \mathcal{F} \cap \mathcal{F}^c \cap \bigcup_{i=n/2 - n^{2/3}}^{n/2 + n^{2/3}} \binom{[n]}{i}$ have size at most $\text{La}(n, P_{G,A}) + 2|G|$. Let C^1, C^2, \dots, C^m be the components of G and $A^j \subset A$ is the part of C^j that belongs to A . If $|\mathcal{F}_{\text{sym}}| > \text{La}(n, P_{G,A}) + 2|G|$, then by Lemma 2.2 and by $\text{La}(n, P_{C^j, A^j}) \leq \text{La}(n, P_{G,A})$ for any $1 \leq j \leq m$, we obtain a copy of C^j . Removing this copy and the complement sets, the remaining family is still large enough to find the second component, and so on. As each time we throw away $2|C^j|$ sets, therefore even before finding the last component we still have more than $|\mathcal{F}| - 2|G| > \text{La}(n, P_{G,A})$ sets. So we found a copy of G and this contradiction yields $|\mathcal{F}| \leq 2^{n-1} + o(\binom{n}{n/2}) + \frac{1}{2} \text{La}(n, P_{G,A}) + 2|G|$.

Observe that if G is bipartite, then $P_{G,A}$ has height 2 (all chains have length at most 2) and all such posets are subsets of $K_{s,1,t}$ for appropriately chosen s and t , where $K_{s,1,t}$ is the so-called complete 3-level poset with elements $u_1, u_2, \dots, u_s, v, w_1, w_2, \dots, w_t$ and $u_i \prec v \prec w_j$ for any $1 \leq i \leq s, 1 \leq j \leq t$ being all its relations. As $K_{s,1,t}$ is a tree poset, Theorem 2.1 implies that for all bipartite G , we have $\text{La}(n, P_{G,A}) \leq (2 + o(1)) \binom{n}{n/2}$. Plugging this in the upper bound at the end of the previous paragraph, we obtain the statement of the theorem. \square

In obtaining the upper bound of Theorem 1.2, we used the inequality $\text{vex}(n, G) \leq 2^{n-1} + \frac{1}{2} \text{La}(n, P_{G,A}) + o(\binom{n}{n/2})$. In a similar fashion, $\text{vex}(n, G) \leq 2^{n-1} + \frac{1}{2} \text{La}(n, P_{G,B}) + o(\binom{n}{n/2})$ and

$$\text{vex}(n, G) \leq 2^{n-1} + \frac{1}{2} \text{La}(n, \{P_{G,A}, P_{G,B}\}) + o\left(\binom{n}{n/2}\right)$$

hold, where $\text{La}(n, \{P_1, P_2\})$ denotes the maximum size of a family $\mathcal{F} \subseteq 2^{[n]}$ that is both $P_{G,A}$ - and $P_{G,B}$ -free. Also, if $\text{La}_{\text{sym}}(n, P)$ denotes the maximum size of a P -free family $\mathcal{F} \subseteq 2^{[n]}$ that is complement closed, i.e. $\mathcal{F}^c = \mathcal{F}$, then the above reasoning yields

$$\text{vex}(n, G) \leq 2^{n-1} + \frac{1}{2} \text{La}_{\text{sym}}(n, P_{G,A}) + o\left(\binom{n}{n/2}\right).$$

We conjecture that this upper bound gives the correct asymptotics of $\text{vex}(n, G)$ for all bipartite graphs that are not matchings.

Conjecture 2.3. *For any bipartite graph G that is not a matching, we have $\text{vex}(n, G) = 2^{n-1} + \frac{1}{2} \text{La}_{\text{sym}}(n, P_{G,A}) + o\left(\binom{n}{n/2}\right)$.*

There is a very important poset parameter that is relevant in forbidden subposet problems. For a poset P , let $e(P)$ denotes the largest integer k such that for any n and $j < n$ the family $\bigcup_{i=1}^k \binom{[n]}{j+i}$ is P -free. In words, $e(P)$ denote the maximum number of consecutive levels of the Boolean cube that we can have without creating a copy of P . This means that if we consider the family containing the middle $e(P)$ levels of the Boolean cube, then we obtain the lower bound $(e(P) + o(1))\binom{n}{n/2} \leq \text{La}(n, P)$. There used to be a very important conjecture of Bukh [3] and Griggs, Lu [18] in the area of forbidden subposet problems. It stated that this construction is asymptotically optimal for any poset P , and so $\text{La}(n, P) = (e(P) + o(1))\binom{n}{n/2}$. This has been very recently disproved by Ellis, Ivan, and Leader [4], but their smallest counterexample is the Boolean poset B_4 of dimension 4 with $e(P) = 4$. Our posets of interest all have height 2.

What bearings does the parameter $e(P)$ have on $\text{vex}(n, G)$ for bipartite graphs G ? As we discussed in the proof of Theorem 1.2, $P_{G,A}$ has height 2 and thus it is a subposet of $K_{s,1,t}$ for s, t large enough, and thus $e(P_{G,A}) \leq 2$, and so $e(P_{G,A})$ is either 1 or 2. Note that $e(P_{G,A}) = e(P_{G,B})$ by the definition of e and $(e(P_{G,A}) + o(1))\binom{n}{n/2} \leq \text{La}(n, \{P_{G,A}, P_{G,B}\})$. Can we turn the middle level construction showing this lower bound into a construction that gives the analogous lower bound for $\text{vex}(n, G)$? The answer is positive, but let us start with the simpler case $e(P_{G,A}) = 1$. Putting together the lower bound of Theorem 1.2 and the upper bound $\text{vex}(n, G) \leq 2^{n-1} + o\left(\binom{n}{n/2}\right) + \frac{1}{2} \text{La}(n, P_{G,A})$, we obtain the following corollary for those bipartite graphs G for which the generally false conjecture by Bukh and Griggs-Lu holds for $P_{G,A}$ with $e(P_{G,A}) = 1$.

Corollary 2.4. *If for a bipartite graph $G = (A, B, E)$ that is not a matching, we have $e(P_{G,A}) = 1$ and $\text{La}(n, P_{G,A}) = (1 + o(1))\binom{n}{n/2}$, then $\text{vex}(n, G) = 2^{n-1} + (1/2 + o(1))\binom{n}{n/2}$ holds.*

Let us now consider bipartite graphs with $e(P_{G,A}) = 2$. This means that the union of the two middle levels does not contain $P_{G,A}$ nor $P_{G,B}$. For a family $\mathcal{F} \subseteq 2^{[n]}$ let us write $\mathcal{U}(\mathcal{F}) := \{G \subseteq [n] : \exists F \in \mathcal{F}, F \subseteq G\}$. Let us introduce the following families:

$$\mathcal{F}_0^{\text{even}} = \{F \subseteq [n] : 1 \in F, |F| = n/2 - 1, n/2\} \cup \{F \subseteq [n] : 1 \notin F, |F| = n/2, n/2 + 1\},$$

$$\mathcal{F}_0^{\text{odd}} = \binom{[n]}{\lfloor n/2 \rfloor} \cup \binom{[n]}{\lceil n/2 \rceil}, \quad \mathcal{F}^{\text{odd}} = \mathcal{U}(\mathcal{F}_0^{\text{odd}}), \quad \mathcal{F}^{\text{even}} = \mathcal{U}(\mathcal{F}_0^{\text{even}}).$$

Now $\mathcal{F}_0^{\text{odd}}$ cannot contain $P_{G,A}$ nor $P_{G,B}$ by definition of e , while $\mathcal{F}_0^{\text{even}}$ has this property as there is no containment between $\{F \subseteq [n] : 1 \in F, |F| = n/2 - 1, n/2\}$ and $\{F \subseteq [n] : 1 \notin F, |F| = n/2, n/2 + 1\}$, and separately these two subfamilies are two consecutive levels of Boolean cubes of dimension $n - 1$. Next, we claim that $Kn_n[\mathcal{F}^{\text{odd}}]$ and $Kn_n[\mathcal{F}^{\text{even}}]$ are G -free. Indeed, vertices corresponding to sets in $\mathcal{F}^{\text{even}} \setminus \mathcal{F}_0^{\text{even}}$ and $\mathcal{F}^{\text{odd}} \setminus \mathcal{F}_0^{\text{odd}}$ are isolated in $Kn_n[\mathcal{F}^{\text{even}}]$ and $Kn_n[\mathcal{F}^{\text{odd}}]$, respectively. So a copy of G should be in $Kn_n[\mathcal{F}_0^{\text{even}}]$ or in $Kn_n[\mathcal{F}_0^{\text{odd}}]$. Note that if G has components C^1, C^2, \dots, C^r , then $e(P_{G,A}) = \max\{e(P_{C^i, A_i}) : 1 \leq i \leq r\}$, so there exists a C^i with $e(P_{C^i, A_i}) = 2$. A copy of C^i in $Kn_n[\mathcal{F}^{\text{even}}]$ or $Kn_n[\mathcal{F}^{\text{odd}}]$

would yield a copy of P_{C^i, A_i} in $Kn_n[\mathcal{F}^{even}]$ or $Kn_n[\mathcal{F}^{odd}]$ as in the proof of Theorem 1.2. But that contradicts $e(P_{C^i, A_i}) = 2$. So we obtained a lower bound $\text{vex}(n, G) \geq 2^{n-1} + (1+o(1))\binom{n}{n/2}$ for arbitrary bipartite graphs G with $e(P_{G,A}) = 2$. Together with the upper bound of Theorem 1.2, it yields the following corollary.

Corollary 2.5. *If for a bipartite graph $G = (A, B, E)$, we have $e(P_{G,A}) = 2$, then $\text{vex}(n, G) = 2^{n-1} + (1 + o(1))\binom{n}{n/2}$ holds.*

Let us finish this subsection with two remarks on the smallest/simplest examples of bipartite graphs G for which $\text{vex}(n, G)$ is not known up the constant term of $\binom{n}{n/2}$. Among even cycles we have $e(P_{C_4, A}) = 2$, and for any $t \geq 3$, we have $e(P_{C_{2t}, A}) = 1$. The corresponding posets have special names: $P_{C_4, A}$ is the butterfly poset \bowtie and $P_{C_{2t}, A}$ for ≥ 3 is the crown poset also denoted by C_{2t} . (Note that $P_{C_{2t}, A} = P_{C_{2t}, B}$ for any $t \geq 2$.) Corollary 2.5 gives the asymptotics of $\text{vex}(n, C_4)$, but, as mentioned in the introduction, its *exact value* was determined by Katona and Nagy [21]. The equality $\text{La}(n, P) = (e(P) + o(1))\binom{n}{n/2}$ is known to be true for C_{2t} (for t even proved by Griggs and Lu [18], for t odd proved by Lu [24]), with the exception of C_6 and C_{10} . So these two graphs are natural targets of future research.

As we have mentioned several times, Ellis, Ivan, and Leader [4] disproved the conjecture $\text{La}(n, P) = (e(P) + o(1))\binom{n}{n/2}$. They constructed a family $\mathcal{F} \subseteq \binom{[n]}{n/2}$ of size $0.29\binom{n}{n/2}$ that does not contain all six sets of any 'middle 4-cube', i.e. for any $S \subset [n]$ with $|S| = n/2 - 2$, $|T| = n/2 + 2$, we have $\{G : S \subset G \subset T, |G| = n/2\} \not\subseteq \mathcal{F}$. Adding sets of size $n/2 \pm 1, 2$ shows $\text{La}(n, B_4) \geq (4.29 + o(1))\binom{n}{n/2}$, where B_4 is the Boolean lattice of dimension 4. Also, let G_4 be the bipartite incidence graph of the complete graph K_4 , that is $G_4 = (A, B, E)$ with $A = E(K_4)$, $B = V(K_4)$ and $(e, v) \in E$ if and only if $v \in e$. Then the above \mathcal{F} together with all the sets of size $n/2 - 1$ shows that $\text{La}(n, P_{G_4, A}) \geq (1.29 + o(1))\binom{n}{n/2}$. Similarly, $\mathcal{F} \cup \binom{[n]}{n/2+1}$ shows $\text{La}(n, P_{G_4, B}) \geq (1.29 + o(1))\binom{n}{n/2}$. On the other hand, the first family contains a copy of $P_{G_4, B}$ while the second contains a copy of $P_{G_4, A}$. If the original conjecture would have been true, then the parameter $\text{La}(n, P)$ would have been *principal*, i.e. $\text{La}(n, \mathcal{P}) = (1 + o(1)) \min\{\text{La}(n, P) : P \in \mathcal{P}\}$ for any family \mathcal{P} of posets. In view of the above, we conjecture that this is not true, in particular we make the following conjecture.

Conjecture 2.6. $\text{La}(n, P_{G_4, A}) - \text{La}(n, \{P_{G_4, A}, P_{G_4, B}\}) = \Omega\left(\binom{n}{n/2}\right)$.

Let us point out that by a result of Griggs and Lu (Theorem 1.4 in [18]) $\text{La}(n, P_{G_4, A}) \leq (1 + \sqrt{2/3} + o(1))\binom{n}{n/2}$ and so if the limit $\lim_n \frac{\text{La}(n, P_{G_4, A})}{\binom{n}{n/2}}$ exists, then its value is a non-integer.

2.2 Non-bipartite graphs

To address the problem of estimating $\text{vex}(n, G)$ for non-bipartite graphs G , we start with the proof of Proposition 1.3.

Proof of Proposition 1.3. Consider the family $\mathcal{F}_k = \{F \subseteq [n] : |F| > \frac{k}{2k+1}n\}$. We claim that $Kn_n[\mathcal{F}_k]$ is C_{2k+1} -free. Indeed, for any $F_1, F_2, \dots, F_{2k+1} \in \mathcal{F}_k$, we have $\frac{\sum_{i=1}^{2k+1} |F_i|}{n} > (2k+1) \cdot \frac{k}{2k+1} = k$, and thus there exists an element $x \in [n]$ that belongs to at least $k+1$

F_i s. Therefore these sets form an independent set in Kn_n , which does not exist in C_{2k+1} . The size of \mathcal{F}_k is $2^n - \binom{n}{\leq 2k+1}$. This completes the proof of the proposition. \square

Now we turn to the proof of Theorem 1.4. We will use Katona's cycle method [20]. Let π be a cyclic permutation of $[n]$. We call a subset S of $[n]$ an *interval with respect to π* if $S = \{\pi(i), \pi(i+1), \pi(i+2), \dots, \pi(i+j)\}$ for some $1 \leq i \leq n, 0 \leq j \leq n-1$, where addition is modulo n , so for example $\{\pi(n-1), \pi(n), \pi(1)\}$ is an interval. We will omit "with respect to π " whenever it is clear from context. For a family $\mathcal{F} \subseteq 2^{[n]}$ and a cyclic permutation π of $[n]$, we define $\mathcal{F}_\pi = \{F \in \mathcal{F} : F \text{ is an interval with respect to } \pi\}$. We shall also need the following weight function: $w_n : [n] \rightarrow \mathbb{N}$ with $w_n(i) = \binom{n}{i}$. The weight $w_n(F)$ of a set F is $w_n(|F|)$, and we let $w_n(\mathcal{F}) = \sum_{F \in \mathcal{F}} w_n(F)$, and we will sometimes omit the subscript n if it is clear from context. The following lemma reduces our problem "to the cycle".

Lemma 2.7. *Let \mathbf{G} be a set of pair (\mathcal{G}, π) with \mathcal{G} being a family of intervals with respect to π and π is a cyclic permutation of $[n]$. Assume that $w(\mathcal{G}) \leq n \cdot B$ for every \mathcal{G} for which there exists a π such that $(\mathcal{G}, \pi) \in \mathbf{G}$. Suppose further that for some $\mathcal{F} \subseteq 2^{[n]}$ it holds that for any cyclic permutation π , we have $(\mathcal{F}_\pi, \pi) \in \mathbf{G}$. Then $|\mathcal{F}| \leq B$.*

This lemma is well-known, we include a proof for the sake of completeness.

Proof. Let us count $\sum_\pi w(\mathcal{F}_\pi)$ in two ways, where the summation runs over all cyclic permutation π of $[n]$. By the assumptions of the lemma, it is at most $(n-1)! \cdot nB = n!B$, as the number of cyclic permutations is $(n-1)!$. On the other hand, every $F \subseteq [n]$ is an interval with respect to exactly $|F|!(n-|F|)!$ cyclic permutations, so the sum equals

$$\sum_{F \in \mathcal{F}} |F|!(n-|F|)! \cdot w(F) = \sum_{F \in \mathcal{F}} n!.$$

We obtained

$$|\mathcal{F}| \cdot n! = \sum_{F \in \mathcal{F}} n! \leq n!B,$$

and dividing by $n!$ yields the statement of the lemma. \square

Let us continue with the proof of Theorem 1.4.

Proof of Theorem 1.4. Based on Lemma 2.7, it is enough to prove that for any family \mathcal{G} of intervals such that $Kn_n[\mathcal{G}]$ is C_{2k+1} -free, we have

$$w_n(\mathcal{G}) \leq n \sum_{i=\lceil \frac{kn}{2k+1} \rceil - \lceil 2(k+1) \log(2k) \rceil}^n \binom{n}{i}.$$

For any $j \leq \frac{kn}{2k+1}$ let $m(j)$ denote the maximum number m such that $\lfloor \frac{kn}{2k+1} \rfloor - j + 2km \leq kn$. Observe that $m(j)$ is monotone increasing in j and $m(j+2k) = m(j) + 1$. Also, $m(j) \leq n/2$ with equality only if n is even and $j = \lfloor \frac{kn}{2k+1} \rfloor$. We also have $m(j) \geq \lceil \frac{kn}{2k+1} \rceil - \lceil (k+1) \log(2k) \rceil$.

Consider the following mapping: for any interval F of size $\lfloor \frac{kn}{2k+1} \rfloor - j$, let $f(F) = \{F_1, F_2, \dots, F_{2k}\}$, where all F_i s are intervals of size $m(j)$ such that F_1 is the interval with left endpoint immediately after F 's right endpoint, F_2 is the interval with left endpoint immediately after F_1 's right endpoint, and so on.

By definition and by the sizes of the intervals F is disjoint with F_1 , F_i is disjoint with F_{i+1} for all $1 \leq i \leq 2k - 1$. Also, by the definition of $m(j)$, F is disjoint with F_{2k} . Indeed, if $F = \{1, \dots, \lfloor \frac{kn}{2k+1} \rfloor - j\}$, then the last element of F_{2k} is $\lfloor \frac{kn}{2k+1} \rfloor - j + 2km \leq kn$, and the first element is more than $\lfloor \frac{kn}{2k+1} \rfloor - j$ if n is sufficiently large. Therefore, F and the sets in $f(F)$ form a C_{2k+1} in Kn_n . This means that for every $G \in \mathcal{G}$ at least one set of $f(G)$ does not belong to \mathcal{G} .

Observe that if j is fixed, then an interval F of size $m(j)$ belongs to exactly $2k$ images $f(G)$ with $|G| = \lfloor \frac{kn}{2k+1} \rfloor - j$. As mentioned above, for every $m > \frac{kn}{2k+1}$ there are exactly $2k$ values of j for which $m(j) = m$. Suppose there are M intervals of size m missing from \mathcal{G} . Then for any j with $m(j) = m$, \mathcal{G} can contain at most $2kM$ intervals of size $\lfloor \frac{kn}{2k+1} \rfloor - j$. Therefore, if

$$2k \sum_{j:m(j)=m} w_n \left(\left\lfloor \frac{kn}{2k+1} \right\rfloor - j \right) \leq w_n(m),$$

then the weight of all the n intervals of length m is larger than the weight of all the j -element intervals with $m(j) = m$ plus the weight of the m -element intervals in \mathcal{G} . In other words, to achieve largest possible weight, it is not worth putting any interval of length $\lfloor \frac{kn}{2k+1} \rfloor - j$ into \mathcal{G} . Proving this for every possible m would complete the proof of the upper bound on $w_n(\mathcal{G})$.

Now for $h < n/2$, we have $\frac{w_n(h)}{w_n(h-1)} = \frac{\binom{n}{h}}{\binom{n}{h-1}} = \frac{n-h+1}{h}$. So if $h < \frac{kn}{2k+1}$, then $\frac{w_n(h)}{w_n(h-1)} > \frac{k+1}{k}$.

Hence writing $z := \lfloor \frac{kn}{2k+1} \rfloor$, we have

$$\frac{w_n(m(j))}{w_n(z-j)} \geq \frac{w_n(z)}{w_n(z-j)} > \left(\frac{k+1}{k} \right)^j > 4k^2,$$

if $j \geq 2(k+1) \log(2k)$ as $(1 + \frac{1}{k})^{k+1} > e$ for all $k \leq 1$. This means that for any m such that $m(j) = m$ implies $j \geq 2(k+1) \log(2k)$, we have

$$\sum_{G \in \mathcal{G}: |G|=m} w_n(G) + \sum_{G \in \mathcal{G}: |G|=z-j, m(j)=m} w_n(G) \leq n \binom{n}{m}.$$

This implies $\sum_{G \in \mathcal{G}} w(G) \leq n \sum_{j \geq z - 2(k+1) \log(2k)} \binom{n}{j}$, as needed.

The second statement of the theorem follows as the first statement and Proposition 1.3 imply $\binom{n}{\leq \lfloor \frac{kn}{2k+1} \rfloor - (k+1) \log(2k)} \leq 2^n - \text{vex}(n, C_{2k+1}) \leq \binom{n}{\leq \lfloor \frac{kn}{2k+1} \rfloor}$, and for any $m \leq (1/2 - \varepsilon)n$ we have $\binom{n}{\leq m} \leq C_\varepsilon \binom{n}{m}$. \square

References

- [1] M. Alishahi, A. Taherkhani, Extremal G -free induced subgraphs of Kneser graphs. *Journal of Combinatorial Theory, Series A*, 159 (2018), 269-282.
- [2] N. Alon, A. Krech and T. Szabó. Turán's theorem in the hypercube. *SIAM Journal on Discrete Mathematics* 21 (2007), 66-72.
- [3] B. Bukh, Set Families with a Forbidden Subset. *The Electronic Journal of Combinatorics*, (2009) R142.

- [4] D. Ellis, M.-R. Ivan, I. Leader, Turán densities for daisies and hypercubes, arXiv:2401.16289
- [5] P. Erdős, On a lemma of Littlewood and Offord, *Bull. Amer. Math. Soc.* 51 (1945), 898–902.
- [6] P. Erdős, A problem on independent r -tuples, *Ann. Univ. Sci. Budapest* 8 (1965), 93–95.
- [7] P. Erdős, C. Ko, R. Rado, Intersection theorems for systems of finite sets. *Quart. J. Math. Oxford* 12, (1961) 313–320.
- [8] P. Frankl, A. Kupavskii, Families with no s pairwise disjoint sets, *J. Lond. Math. Soc.*, 95 (3) (2017), 875–894
- [9] P. Frankl, A. Kupavskii, New inequalities for families without k pairwise disjoint members, *J. Combin. Theory Ser. A*, 157 (2018), 427–434.
- [10] P. Frankl, A. Kupavskii, (2018). The Erdős Matching Conjecture and concentration inequalities. arXiv preprint arXiv:1806.08855.
- [11] P. Frankl, A. Kupavskii, Families of sets with no matching of sizes 3 and 4 *Eur. J. Comb.*, 75 (2019), 123–135.
- [12] Z. Füredi, M. Simonovits, The history of degenerate (bipartite) extremal graph problems. In *Erdős centennial (2013)* pp. 169-264. Berlin, Heidelberg: Springer Berlin Heidelberg.
- [13] D. Gerbner, N. Lemons, C. Palmer, B. Patkós, V. Szécsi, Almost intersecting families of sets *SIAM J. Discrete Math.* 26 (2012) 1657–1699.
- [14] D. Gerbner, A. Methuku, D.T. Nagy, B. Patkós, M. Vizer, Vertex Turán problems for the oriented hypercube. *Acta Universitatis Sapientiae, Mathematica*, 13(2) (2021), 356-366.
- [15] D. Gerbner, A. Methuku, D.T. Nagy, B. Patkós, M. Vizer, Stability Results for Vertex Turán Problems in Kneser Graphs. *The Electronic Journal of Combinatorics* (2019), P2-13.
- [16] D. Gerbner, B. Patkós, *Extremal finite set theory*. CRC Press, 2018.
- [17] J.R. Griggs, W.T. Li, Progress on poset-free families of subsets. *Recent trends in combinatorics*, (2016) 317-338.
- [18] J.R. Griggs, L. Lu, On families of subsets with a forbidden subposet. *Combinatorics, Probability and Computing*, 18(5) (2009), 731-748.
- [19] J. R. Johnson, J. Talbot. Vertex Turán problems in the hypercube. *Journal of Combinatorial Theory, Series A* 117 (2010), 454–465.
- [20] G.O.H. Katona, A simple proof of the Erdős-Chao Ko-Rado theorem, *Journal of Combinatorial Theory, Series B* 13(2), 183-184, 1972.
- [21] G.O.H. Katona, D.T. Nagy, Union-Intersecting Set Systems. *Graphs and Combinatorics* 31, 1507–1516 (2015).

- [22] D.J. Kleitman, Maximal number of subsets of a finite set no k of which are pairwise disjoint, *J. Combin. Theory*, 5 (1968), 157–163.
- [23] X. C. Li, R. X. Hao, (2022). The vertex Turán density in 3-ary n -cubes. *Discrete Applied Mathematics*, 311, 129-141.
- [24] L. Lu, On crown-free families of subsets. *Journal of Combinatorial Theory, Series A*, 126 (2014), 216–231.
- [25] A. Taherkhani, Size and Structure of Large (s, t) -Union Intersecting Families. *The Electronic Journal of Combinatorics*, (2022) P2-25.

3 Statements and Declarations

Research supported by NKFIH under grants FK132060 and KKP-133819.

The authors have no relevant financial or non-financial interests to disclose.