

WIGNER ANALYSIS OF FOURIER INTEGRAL OPERATORS WITH SYMBOLS IN THE SHUBIN CLASSES

ELENA CORDERO, GIANLUCA GIACCHI, LUIGI RODINO, AND MARIO VALENZANO

ABSTRACT. We study the decay properties of Wigner kernels for Fourier integral operators of types I and II. The symbol spaces that allow a nice decay of these kernels are the Shubin classes $\Gamma^m(\mathbb{R}^{2d})$, with negative order m . The phases considered are the so-called tame ones, which appear in the Schrödinger propagators. The related canonical transformations are allowed to be nonlinear. It is the nonlinearity of these transformations that are the main obstacles for nice kernel localizations when symbols are taken in the Hörmander's class $S_{0,0}^0(\mathbb{R}^{2d})$. Here we prove that Shubin classes overcome this problem and allow a nice kernel localization, which improves with the decreasing of the order m .

1. INTRODUCTION

The protagonist of this study is the Wigner distribution, one of the most popular time-frequency representations. It was introduced by Wigner in 1932 [35] in the framework of Quantum Mechanics and later applied to signal processing and time-frequency analysis by Ville, Cohen and many other authors, see, e.g., [3, 4, 34] and the textbooks [19, 25, 26].

Definition 1.1. Consider $f, g \in L^2(\mathbb{R}^d)$. The cross-Wigner distribution $W(f, g)$ is

$$(1) \quad W(f, g)(x, \xi) = \int_{\mathbb{R}^d} f\left(x + \frac{t}{2}\right) \overline{g\left(x - \frac{t}{2}\right)} e^{-2\pi i t \xi} dt, \quad (x, \xi) \in \mathbb{R}^{2d}.$$

If $f = g$ we write $Wf := W(f, f)$, the so-called Wigner distribution of f .

Wigner used the above representation to analyse the action of the Schrödinger propagators. We may extend the Wigner approach in [35] as follows: given a linear operator $T : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$, we consider an operator K on $\mathcal{S}(\mathbb{R}^{2d})$ such that

$$(2) \quad W(Tf, Tg) = KW(f, g), \quad f, g \in \mathcal{S}(\mathbb{R}^d).$$

Its integral kernel k is called the *Wigner kernel* of T :

$$(3) \quad W(Tf, Tg)(z) = \int_{\mathbb{R}^{2d}} k(z, w) W(f, g)(w) dw, \quad z \in \mathbb{R}^{2d}, \quad f, g \in \mathcal{S}(\mathbb{R}^d).$$

2010 *Mathematics Subject Classification.* 35S05, 35S30, 47G30, 42C15.

As an elementary example of the effectiveness of the Wigner distribution, consider the Schrödinger propagator T_τ , for a fixed time $\tau \in \mathbb{R}$, of the free particle equation

$$T_\tau f(x) = \int_{\mathbb{R}^d} e^{2\pi i(x\xi - \tau\xi^2)} \widehat{f}(\xi) d\xi, \quad x \in \mathbb{R}^d.$$

We have

$$W(T_\tau f)(x, \xi) = Wf(x - \tau\xi, \xi)$$

with Wigner kernel

$$(4) \quad k = \delta_{z - \chi(w)}, \quad w, z \in \mathbb{R}^{2d},$$

where, if we write $w = (y, \eta)$, then $\chi(y, \eta) = (y + \tau\eta, \eta)$. So we recapture exactly the inertial first Newton's law, from a probabilistic point of view. This striking result is due to the peculiar action of W on the phase $\Phi(x, \xi) = x\xi - \tau\xi^2$. It generalizes to quadratic $\Phi(x, \xi)$, corresponding to quadratic Hamiltonians and linear symplectic map χ in (4), see for example [24].

Our aim is to extend this analysis to more general operators, namely Fourier integral operators of the form

$$(5) \quad T_I f(x) = \int_{\mathbb{R}^d} e^{2\pi i\Phi(x, \xi)} \sigma(x, \xi) \widehat{f}(\xi) d\xi, \quad f \in \mathcal{S}(\mathbb{R}^d),$$

with phase Φ and symbol σ in suitable classes. A preliminary step was presented in [13], with T a pseudodifferential operator $\sigma(x, D)$, i.e., $\Phi(x, \xi) = x\xi$ in (5). The case of a quadratic Φ and a general σ was considered in [9] and in [8], where a generalization of (4) was obtained by combining a linear symplectic map χ with the kernel of a pseudodifferential operator.

In the present paper we focus on the case of nonlinear symplectic mappings χ corresponding to non-quadratic Φ , which we call *tame*, see Section 2 below for their definition.

As a counterpart of (4) we look for estimates of the type

$$(6) \quad |k(z, w)| \lesssim \frac{1}{\langle z - \chi(w) \rangle^{2N}},$$

in the spirit of the estimates for Gabor kernels, which have been widely investigated in the literature, classical references are [2, 11, 16, 17, 27, 28], see also [19, Chapter 5].

There are two obstructions to the validity of (6). The first, evident from (4) and also in the linear case, is that $k(z, w)$ is not point-wise defined for $z = \chi(w)$. This can be easily rephrased by a rescaling of regularity. The second obstruction is of deeper nature, and it concerns only the nonlinear symplectic map χ . In fact, it is well known that the Wigner transform may produce the so-called ghost frequencies. As observed in [9, 14], they are exactly preserved for Schrödinger propagators for linear χ , i.e., quadratic Φ , but this is not the case for nonlinear χ . Namely, highly

oscillating terms may appear in the expression of the kernel $k(z, w)$ outside the graph of $z = \chi(w)$.

As a first attempt for eliminating ghost frequencies and re-establishing the validity of (6), we shall consider in the sequel symbols σ of low order in Shubin classes [33]. Unluckily, this framework does not allow a direct application to Schrödinger equations, for which we address a future work, following a different smoothing procedure.

Let us outline the contents of the paper. Our starting point, in Section 3, will be the following *abstract* definition, in the lines of [16].

Definition 1.2. *Consider a tame symplectic diffeomorphism χ (cf. Definition 2.3 below). For $N \in \mathbb{N}_+$, $N > d$, we say that the operator K in (2) is in the class $FIO(\chi, N)$ if its Wigner kernel k in (3) satisfies, for $z = (z_1, z_2)$, $w = (w_1, w_2) \in \mathbb{R}^{2d}$,*

$$(7) \quad |k(z, w)| \lesssim \frac{1}{\langle z - \chi(w) \rangle^{2N}}.$$

Examples of operators which fall in the above class are pseudodifferential operators $\sigma(x, D)$ (the Kohn-Nirenberg form), defined by

$$(8) \quad \sigma(x, D)f(x) = \int_{\mathbb{R}^d} e^{2\pi i x \xi} \sigma(x, \xi) \hat{f}(\xi) d\xi,$$

with a symbol σ in the Shubin classes $\Gamma^m(\mathbb{R}^{2d})$, $m < -2(d + N)$, whose Wigner kernel k_σ satisfies

$$(9) \quad |k_\sigma(z, w)| \lesssim \frac{1}{\langle z - w \rangle^{2N}}.$$

Here $\chi = I$, the identity mapping, cf. Section 2 below. More generally, Fourier integral operators of type I (cf. (5)) and II, having symbols in the same Shubin classes above and *tame* canonical transformations, fall in the class above, as we shall show in Sections 4 and 5.

Let us state here the preliminary results of Section 3, which are the core of this study and may be collected as follows.

Theorem 1.3 (Properties of the class $FIO(\chi, N)$).

- (i) *Boundedness.* $T \in FIO(\chi, N)$ is bounded on $L^2(\mathbb{R}^d)$.
- (ii) *Algebra Property.* If $T_i \in FIO(\chi_i, N)$, $i = 1, 2$, then $T_1 T_2 \in FIO(\chi_1 \chi_2, N)$.
- (iii) If $T \in FIO(\chi, N)$ then its adjoint T^* is in $FIO(\chi^{-1}, N)$.

In Section 4 we shall show the Fourier integral operators of type I in (5), having symbols in suitable Shubin classes $\Gamma^m(\mathbb{R}^{2d})$ and tame phase functions are in the class $FIO(\chi, N)$.

The last Section 5 is devoted to the L^2 -adjoint of the FIO I in (5), which can be written explicitly in the form

$$T_{II}f(x) = \int_{\mathbb{R}^{2d}} e^{-2\pi i[\Phi(y,\xi)-x\xi]} \tau(y,\xi) f(y) dy d\xi, \quad f \in \mathcal{S}(\mathbb{R}^d).$$

Using tools from metaplectic Wigner distributions implemented in [8, 9] we are able to compute the Wigner kernel of the FIOs II above and prove that, under suitable assumptions on their symbols, they belong to $FIO(\chi, N)$ as well. We underline that these results are valid for the whole class of tame phase Φ defined in Subsection 2.3, of particular interest is the case Φ non quadratic which gives rise to nonlinear symplectic transformations χ , which were not treated in [8].

We believe that such theoretical study will pave the way to a better understanding of Wigner kernels for Fourier integral operators, with possible applications to dynamical versions of Hardy's uncertainty principles [21, 29, 36, 37, 38], see also the recent contribution [20].

2. PRELIMINARIES

Notation. We define $t^2 = t \cdot t$, $t \in \mathbb{R}^d$, and, similarly, $xy = x \cdot y$. The space $\mathcal{S}(\mathbb{R}^d)$ is the Schwartz class and $\mathcal{S}'(\mathbb{R}^d)$ its dual (the space of temperate distributions). The brackets $\langle f, g \rangle$ means the extension to $\mathcal{S}'(\mathbb{R}^d) \times \mathcal{S}(\mathbb{R}^d)$ of the inner product $\langle f, g \rangle = \int f(t)g(t)dt$ on $L^2(\mathbb{R}^d)$ (conjugate-linear in the second component).

A point in the phase space is denoted by $z = (x, \xi) \in \mathbb{R}^{2d}$. We call (time-frequency shift) the operators

$$(10) \quad \pi(z)f(t) = e^{2\pi i\xi t} f(t - x), \quad t \in \mathbb{R}^d.$$

$GL(d, \mathbb{R})$ denotes the group of real invertible $d \times d$ matrices.

2.1. The symplectic group $Sp(d, \mathbb{R})$, metaplectic operators and Wigner distributions. The standard symplectic matrix is

$$(11) \quad J = \begin{pmatrix} 0_{d \times d} & I_{d \times d} \\ -I_{d \times d} & 0_{d \times d} \end{pmatrix}.$$

The symplectic group is defined by

$$(12) \quad Sp(d, \mathbb{R}) = \{ \mathcal{A} \in GL(2d, \mathbb{R}) : \mathcal{A}^T J \mathcal{A} = J \},$$

where \mathcal{A}^T is the transpose of \mathcal{A} . We have $\det(\mathcal{A}) = 1$.

For $L \in GL(d, \mathbb{R})$ and $C \in Sym(2d, \mathbb{R})$, define:

$$(13) \quad \mathcal{D}_L := \begin{pmatrix} L^{-1} & 0_{d \times d} \\ 0_{d \times d} & L^T \end{pmatrix} \quad \text{and} \quad V_C := \begin{pmatrix} I_{d \times d} & 0 \\ C & I_{d \times d} \end{pmatrix}.$$

The matrices J , V_C , and \mathcal{D}_L generate the group $Sp(d, \mathbb{R})$.

The Schrödinger representation ρ of the Heisenberg group is given by

$$\rho(x, \xi; \tau) = e^{2\pi i \tau} e^{-\pi i \xi x} \pi(x, \xi),$$

for all $x, \xi \in \mathbb{R}^d$, $\tau \in \mathbb{R}$. For every $A \in Sp(d, \mathbb{R})$, $\rho_A(x, \xi; \tau) := \rho(A(x, \xi); \tau)$ defines another representation of the Heisenberg group that is equivalent to ρ , that is, there exists a unitary operator $\hat{A} : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ such that

$$(14) \quad \hat{A} \rho(x, \xi; \tau) \hat{A}^{-1} = \rho(A(x, \xi); \tau), \quad x, \xi \in \mathbb{R}^d, \tau \in \mathbb{R}.$$

This operator is not unique: if \hat{A}' is another unitary transformation satisfying (14), then $\hat{A}' = c\hat{A}$, for some $c \in \mathbb{C}$, with $|c| = 1$. The set $\{\hat{A} : A \in Sp(d, \mathbb{R})\}$ is a group under operator composition and has the metaplectic group $Mp(d, \mathbb{R})$ as subgroup. It is a realization of the two-fold cover of $Sp(d, \mathbb{R})$. The projection

$$(15) \quad \pi^{Mp} : Mp(d, \mathbb{R}) \rightarrow Sp(d, \mathbb{R})$$

is a group homomorphism with kernel $\ker(\pi^{Mp}) = \{-id_{L^2}, id_{L^2}\}$.

Here, if $\hat{A} \in Mp(d, \mathbb{R})$, the matrix A will be the unique symplectic matrix satisfying $\pi^{Mp}(\hat{A}) = A$. Some examples of metaplectic operators we will use in the following are detailed below.

Example 2.1. Consider the matrices J , \mathcal{D}_L and V_C defined in (11) and (13). Then, if we denoted by \mathcal{F} the Fourier transform,

- (i) $\pi^{Mp}(\mathcal{F}) = J$;
- (ii) if $\mathfrak{T}_L := |\det(L)|^{1/2} f(L \cdot)$, then $\pi^{Mp}(\mathfrak{T}_L) = \mathcal{D}_L$;

The relation between time-frequency shifts and metaplectic operators is the following:

$$(16) \quad \pi(\mathcal{A}z) = c_{\mathcal{A}} \hat{\mathcal{A}} \pi(z) \hat{\mathcal{A}}^{-1} \quad \forall z \in \mathbb{R}^{2d},$$

with a phase factor $c_{\mathcal{A}} \in \mathbb{C}$, $|c_{\mathcal{A}}| = 1$ (see, e.g., [23, 24]).

Metaplectic Wigner distributions. In the study of FIOs of type II we will use tools from the theory of metaplectic Wigner distributions. Here we list the basic elements for this study. For $\hat{A} \in Mp(2d, \mathbb{R})$, the *metaplectic Wigner distribution* associated to \hat{A} is defined as

$$(17) \quad W_{\mathcal{A}}(f, g) = \hat{\mathcal{A}}(f \otimes \bar{g}), \quad f, g \in L^2(\mathbb{R}^d).$$

The most important time-frequency representations are metaplectic Wigner distributions. The τ -Wigner distributions, $\tau \in \mathbb{R}$, defined by

$$(18) \quad W_{\tau}(f, g)(x, \xi) = \int_{\mathbb{R}^d} f(x + \tau t) \overline{g(x - (1 - \tau)t)} e^{-2\pi i \xi t} dt, \quad (x, \xi) \in \mathbb{R}^{2d},$$

for $f, g \in L^2(\mathbb{R}^d)$, are metaplectic Wigner distributions. The case $\tau = 1/2$ is the cross-Wigner distribution, defined in (1). τ -Wigner distributions are metaplectic Wigner distributions:

$$W_\tau(f, g) = \hat{A}_\tau(f \otimes \bar{g}),$$

with

$$(19) \quad A_\tau = \begin{pmatrix} (1-\tau)I_{d \times d} & \tau I_{d \times d} & 0_{d \times d} & 0_{d \times d} \\ 0_{d \times d} & 0_{d \times d} & \tau I_{d \times d} & -(1-\tau)I_{d \times d} \\ 0_{d \times d} & 0_{d \times d} & I_{d \times d} & I_{d \times d} \\ -I_{d \times d} & I_{d \times d} & 0_{d \times d} & 0_{d \times d} \end{pmatrix}.$$

In particular, we recapture the Wigner case when $\tau = 1/2$:

$$(20) \quad Wf = W_{1/2}(f, f) = \hat{A}_{1/2}(f \otimes \bar{f}), \quad f \in L^2(\mathbb{R}^d).$$

$\hat{A}_{1/2}$ can be split into the product

$$(21) \quad \hat{A}_{1/2} = \mathcal{F}_2 \mathfrak{L}_L,$$

with

$$L = \begin{pmatrix} I_{d \times d} & \frac{1}{2}I_{d \times d} \\ I_{d \times d} & -\frac{1}{2}I_{d \times d} \end{pmatrix}.$$

Hence

$$\hat{A}_{1/2}F(x, \xi) = \int_{\mathbb{R}^d} F(x + t/2, x - t/2) e^{-2\pi i \xi t} dt, \quad F \in \mathcal{S}(\mathbb{R}^{2d}),$$

and

$$\hat{A}_{1/2}^{-1} = \mathfrak{L}_{L^{-1}} \mathcal{F}_2^{-1},$$

where

$$L^{-1} = \begin{pmatrix} \frac{1}{2}I_{d \times d} & \frac{1}{2}I_{d \times d} \\ I_{d \times d} & -I_{d \times d} \end{pmatrix},$$

so that

$$(22) \quad \hat{A}_{1/2}^{-1}F(x, \xi) = \int_{\mathbb{R}^d} F(x/2 + \xi/2, y) e^{2\pi i(x-\xi)y} dy, \quad F \in \mathcal{S}(\mathbb{R}^{2d}).$$

2.2. Shubin and Hörmander classes [33], [31], [19]. In our study we shall consider the following weight functions

$$(23) \quad v_s(z) = \langle z \rangle^s = (1 + |z|^2)^{\frac{s}{2}}, \quad s \in \mathbb{R},$$

Definition 2.2. Fix $m \in \mathbb{R}$. The shubin class $\Gamma^m(\mathbb{R}^{2d})$ is the set of functions $a \in \mathcal{C}^\infty(\mathbb{R}^{2d})$ satisfying

$$|\partial_z^\alpha a(z)| \leq C_\alpha v_{m-|\alpha|}(z), \quad z \in \mathbb{R}^{2d}, \alpha \in \mathbb{Z}_+^{2d},$$

for a suitable constant $C_\alpha > 0$, where $v_s(z) = \langle z \rangle^s$ is defined in (23).

The Hörmander class $S_{0,0}^0(\mathbb{R}^{2d})$, consists of smooth functions σ on \mathbb{R}^{2d} such that

$$(24) \quad |\partial_x^\alpha \partial_\xi^\beta \sigma(x, \xi)| \leq c_{\alpha, \beta}, \quad \alpha, \beta \in \mathbb{N}^d, \quad x, \xi \in \mathbb{R}^d.$$

2.3. Tame phase functions and related canonical transformations.

Definition 2.3. *We follow the notation of [16, 8]. A real phase function $\Phi(x, \eta)$ is named tame if it satisfies the following properties:*

A1. $\Phi \in C^\infty(\mathbb{R}^{2d})$;

A2. For $z = (x, \xi) \in \mathbb{R}^{2d}$,

$$(25) \quad |\partial_z^\alpha \Phi(z)| \leq C_\alpha, \quad |\alpha| \geq 2;$$

A3. There exists $\delta > 0$:

$$(26) \quad |\det \partial_{x, \eta}^2 \Phi(x, \xi)| \geq \delta.$$

Solving the system

$$(27) \quad \begin{cases} y = \Phi_\eta(x, \eta), \\ \xi = \Phi_x(x, \eta), \end{cases}$$

with respect to (x, ξ) , one obtains a map χ

$$(28) \quad (x, \xi) = \chi(y, \eta),$$

with the following properties:

A4. $\chi : \mathbb{R}^{2d} \rightarrow \mathbb{R}^{2d}$ is a symplectomorphism (smooth, invertible, and preserves the symplectic form in \mathbb{R}^{2d} , i.e., $dx \wedge d\xi = dy \wedge d\eta$.)

A5. For $z = (y, \eta)$,

$$(29) \quad |\partial_z^\alpha \chi(z)| \leq C_\alpha, \quad |\alpha| \geq 1;$$

A6. There exists $\delta > 0$:

$$(30) \quad \left| \det \frac{\partial x}{\partial y}(y, \eta) \right| \geq \delta \quad \text{for } (x, \xi) = \chi(y, \eta).$$

Conversely, as it was observed in [16], to every transformation χ satisfying the three hypothesis above corresponds a tame phase Φ , uniquely determined up to a constant.

2.4. Properties of the Wigner Kernel. The Wigner kernel of a linear bounded operator $T : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$ was introduced and studied in [8]. We recall its definition and the properties useful for our framework.

Definition 2.4. *The Wigner kernel of $T : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$ linear and bounded operator is the distribution $k \in \mathcal{S}'(\mathbb{R}^{4d})$ satisfying*

$$(31) \quad \langle W(Tf, Tg), W(u, v) \rangle = \langle k, W(u, v) \otimes \overline{W(f, g)} \rangle, \quad f, g, u, v \in \mathcal{S}(\mathbb{R}^d).$$

Observe that if $k \in \mathcal{S}(\mathbb{R}^{4d})$ the integral formula (3) holds true. The results of Theorem 3.3 and 4.3 in [8] can be rephrased as follows:

Theorem 2.5. *Consider T as above and let $k_T \in \mathcal{S}'(\mathbb{R}^{2d})$ be its kernel. There exists a unique distribution $k \in \mathcal{S}'(\mathbb{R}^{4d})$ such that (31) holds. Hence, every bounded linear operator $T : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$ has a unique Wigner kernel. Furthermore,*

$$(32) \quad k = \mathfrak{T}_p W k_T,$$

with $\mathfrak{T}_p F(x, \xi, y, \eta) = F(x, y, \xi, -\eta)$.

In particular, if $T \in B(L^2(\mathbb{R}^d))$ has Wigner kernel k , then its adjoint $T^* \in B(L^2(\mathbb{R}^d))$ has Wigner kernel $k(z, w) = k(w, z)$, $z, w \in \mathbb{R}^{2d}$.

3. PROPERTIES OF $FIO(\chi, N)$

This section is devoted to prove Theorem 1.3 in the introduction. This requires several steps, developed in what follows.

Theorem 3.1. *An operator $T \in FIO(\chi, N)$, is bounded on $L^2(\mathbb{R}^d)$.*

Proof. For $f \in L^2(\mathbb{R}^d)$ we recall [19, Chapter 1] that the Wigner $Wf \in L^2(\mathbb{R}^{2d})$ and Moyal's identity $\|Wf\|_{L^2(\mathbb{R}^{2d})} = \|f\|_{L^2(\mathbb{R}^d)}^2$.

Using (3), Definition 1.2, for any $f \in L^2(\mathbb{R}^d)$,

$$\|Tf\|_{L^2(\mathbb{R}^d)}^2 = \|W(Tf)\|_{L^2(\mathbb{R}^{2d})},$$

and

$$\begin{aligned} \|W(Tf)\|_{L^2(\mathbb{R}^{2d})} &\lesssim \left\| \int_{\mathbb{R}^{2d}} \frac{1}{\langle z - \chi(w) \rangle^{2N}} Wf(w) dw \right\|_{L^2(\mathbb{R}^{2d})} \\ &\asymp \left\| \int_{\mathbb{R}^{2d}} \frac{1}{\langle \chi^{-1}(z) - w \rangle^{2N}} Wf(w) dw \right\|_{L^2(\mathbb{R}^{2d})} \\ &\lesssim \left\| \left(\frac{1}{\langle \cdot \rangle^{2N}} * Wf \right) (\chi^{-1}(z)) \right\|_{L^2(\mathbb{R}^{2d})} \\ &\lesssim \left\| \left(\frac{1}{\langle \cdot \rangle^{2N}} * Wf \right) (z) \right\|_{L^2(\mathbb{R}^{2d})} \\ &\leq \left\| \frac{1}{\langle \cdot \rangle^{2N}} \right\|_{L^1(\mathbb{R}^{2d})} \|Wf\|_{L^2(\mathbb{R}^{2d})} \\ &\leq C_N \|f\|_{L^2(\mathbb{R}^d)}^2, \end{aligned}$$

where in the last row we used Young's inequality (observe that $N > d$) and, in the last but one, the change of variables $z' = \chi^{-1}(z)$ which, for any $F \in L^2(\mathbb{R}^{2d})$,

$$\begin{aligned} \|F(\chi^{-1}\cdot)\|_{L^2(\mathbb{R}^{2d})}^2 &= \int_{\mathbb{R}^{2d}} |F(\chi^{-1}(z))|^2 dz = \int_{\mathbb{R}^{2d}} |F(z')|^2 \det |J\chi(z')| dz' \\ &\leq C_N \|F\|_{L^2(\mathbb{R}^{2d})}^2, \end{aligned}$$

by (29). Hence $\|Tf\|_{L^2(\mathbb{R}^d)} \leq \sqrt{C_N} \|f\|_{L^2(\mathbb{R}^d)}$, that is $T \in B(L^2(\mathbb{R}^d))$. \square

Theorem 3.2. *If $T_i \in FIO(\chi_i, N)$, $i = 1, 2$, then $T_1 T_2 \in FIO(\chi_1 \chi_2, N)$.*

Proof. Using the Wigner representation in (3) we can write

$$(33) \quad W(T_1 T_2 f, T_1 T_2 g)(z) = \int_{\mathbb{R}^{2d}} k_{I,1}(z, w) W(T_2 f, T_2 g)(w) dw,$$

where $k_{I,1}(z, w)$ is the Wigner kernel of the operator T_1 , satisfying (7) with symplectic transformation χ_1 . Similarly,

$$W(T_2 f, T_2 g)(w) = \int_{\mathbb{R}^{2d}} k_{I,2}(w, u) W(f, g)(u) du,$$

with $k_{I,2}(w, u)$ being the Wigner kernel of T_2 satisfying (7) with symplectic transformation χ_2 . Substituting the expression of $W(T_2 f, T_2 g)(w)$ in (33) we obtain

$$(34) \quad W(T_1 T_2 f, T_1 T_2 g)(z) = \int_{\mathbb{R}^{4d}} k_{I,1}(z, w) k_{I,2}(w, u) W(f, g)(u) du dw,$$

with $k_{I,i}(z, w)$ satisfying (7), $i = 1, 2$. Interchanging the integrals in (34) (observe that the assumptions of Fubini Theorem are satisfied) we can write

$$(35) \quad W(T_1 T_2 f, T_1 T_2 g)(z) = \int_{\mathbb{R}^{2d}} \left(\int_{\mathbb{R}^{2d}} k_{I,1}(z, w) k_{I,2}(w, u) dw \right) W(f, g)(u) du$$

so that the Wigner kernel $k_{I,12}$ of the product $T_1 T_2$ is given by

$$k_{I,12}(z, u) := \int_{\mathbb{R}^{2d}} k_{I,1}(z, w) k_{I,2}(w, u) dw.$$

Using the Wigner kernel's estimates in (7), we reckon

$$\begin{aligned}
|k_{I,12}(z, u)| &\leq \int_{\mathbb{R}^{2d}} |k_{I,1}(z, w)| |k_{I,2}(w, u)| dw \\
&\lesssim \int_{\mathbb{R}^{2d}} \frac{1}{\langle z - \chi_1(w) \rangle^{2N} \langle w - \chi_2(u) \rangle^{2N}} dw \\
&\succ \int_{\mathbb{R}^{2d}} \frac{1}{\langle z - \chi_1(w) \rangle^{2N} \langle \chi_1(w) - \chi_1 \chi_2(u) \rangle^{2N}} dw \\
&\lesssim \int_{\mathbb{R}^{2d}} \frac{1}{\langle z - \chi_1(w) \rangle^{2N} \langle \chi_1(w) - \chi_1 \chi_2(u) \rangle^{2N}} dw \\
&= \int_{\mathbb{R}^{2d}} \frac{1}{\langle \chi_1(w) - z \rangle^{2N} \langle \chi_1 \chi_2(u) - \chi_1(w) \rangle^{2N}} dw \\
&= \int_{\mathbb{R}^{2d}} \frac{1}{\langle w' - z \rangle^{2N} \langle \chi_1 \chi_2(u) - w' \rangle^{2N}} |\det J_{\chi_1^{-1}}(w)| dw'
\end{aligned}$$

where we used the change of variables $\chi_1(w) = w'$ so that $dw = |\det J_{\chi_1^{-1}}(w)| dw'$ since $|\det J_{\chi_1^{-1}}(w)| \leq C$ by (29), we obtain

$$\begin{aligned}
|k_{I,12}(z, u)| &\lesssim \int_{\mathbb{R}^{2d}} \frac{1}{\langle w' - z \rangle^{2N} \langle \chi_1 \chi_2(u) - w' \rangle^{2N}} dw' \\
&= \int_{\mathbb{R}^{2d}} \frac{1}{\langle v \rangle^{2N} \langle \chi_1 \chi_2(u) - z - v \rangle^{2N}} dv \\
&= (\langle \cdot \rangle^{-2N} * \langle \cdot \rangle^{-2N})(\chi_1 \chi_2(u) - z) \\
&\lesssim \frac{1}{\langle z - \chi_1 \chi_2(u) \rangle^{2N}}
\end{aligned}$$

where in the last row we used the weight convolution property $\langle \cdot \rangle^s * \langle \cdot \rangle^s \lesssim \langle \cdot \rangle^s$ for $s < -2d$ (observe $N > d$). Thus, we obtain the desired estimate

$$|k_{I,12}(z, u)| \lesssim \frac{1}{\langle z - \chi_1 \chi_2(u) \rangle^{2N}},$$

that is $T_1 T_2 \in FIO(\chi_1 \chi_2, N)$. □

Theorem 3.3. *If $T \in FIO(\chi, N)$, then $T^* \in FIO(\chi^{-1}, N)$.*

Proof. Theorem 3.1 gives that $T \in B(L^2(\mathbb{R}^d))$. Let k be integral kernel of T , then Theorem 2.5 says that the adjoint $T^* \in B(L^2(\mathbb{R}^d))$ has kernel \tilde{k} given by

$$\tilde{k}(z, w) = k(w, z).$$

This means it satisfies (cf. (7))

$$(36) \quad |\tilde{k}(z, w)| = |k(w, z)| \lesssim \frac{1}{\langle w - \chi(z) \rangle^{2N}}.$$

Since χ is a bi-Lipschitz transformation, $|w - \chi(z)| \asymp |z - \chi^{-1}(w)|$ so that $\langle w - \chi(z) \rangle^{2N} \asymp \langle z - \chi^{-1}(w) \rangle^{2N}$ and we obtain

$$(37) \quad |\tilde{k}(z, w)| \lesssim \frac{1}{\langle z - \chi^{-1}(w) \rangle^{2N}}.$$

Hence $T^* \in FIO(\chi^{-1}, N)$, as desired. \square

4. FIOS OF TYPE I

Here we focus on the analysis of Wigner kernels for FIOS of type I:

$$(38) \quad T_I f(x) = \int_{\mathbb{R}^d} e^{2\pi i \Phi(x, \xi)} \sigma(x, \xi) \hat{f}(\xi) d\xi, \quad f \in \mathcal{S}(\mathbb{R}^d).$$

Recall that the Schwartz' Kernel Theorem guarantees that every continuous linear operator $T : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$ can be expressed in the form $T = T_I$, with a given phase $\Phi(x, \xi)$ and symbol $\sigma(x, \xi)$ in $\mathcal{S}'(\mathbb{R}^{2d})$.

These operators have been widely investigated in the framework of PDEs, both from a theoretical and a numerical point of view; the literature is so huge that we cannot report all the results but limit to a very partial list of them, cf. [5, 6, 7, 15, 18, 22, 32].

If we assume that T is a continuous linear operator $\mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$ and χ satisfies conditions A4, A5, and A6 in Definition 2.3 then $T = T_{I, \Phi_\chi, \sigma}$ (FIO of type I), with symbol σ and phase Φ_χ .

A first result related to the Wigner kernel of a FIO I was obtained in [8, Theorem 5.8]. There, FIOS of type I with symbols in the Hörmander class $S_{0,0}^0(\mathbb{R}^{2d})$ were considered. Since $\Gamma^m(\mathbb{R}^{2d}) \subset S_{0,0}^0(\mathbb{R}^{2d})$ whenever $m \leq 0$, we can rephrase it in our context as follows.

Theorem 4.1. *Let T_I be a FIO of type I defined in (38) with symbol $\sigma \in \Gamma^m(\mathbb{R}^{2d})$, $m < 0$. For $f \in \mathcal{S}(\mathbb{R}^d)$,*

$$(39) \quad K(W(f, g))(x, \xi) = W(T_I f, T_I g)(x, \xi) = \int_{\mathbb{R}^{2d}} k_I(x, \xi, y, \eta) W(f, g)(y, \eta) dy d\eta,$$

with Wigner kernel k_I given by

$$(40) \quad k_I(x, \xi, y, \eta) = \int_{\mathbb{R}^{2d}} e^{2\pi i [\Phi_I(x, \eta, t, r) - (\xi t + r y)]} \sigma_I(x, \eta, t, r) dt dr,$$

and, for $x, \eta, t, r \in \mathbb{R}^d$,

$$(41) \quad \Phi_I(x, \eta, t, r) = \Phi\left(x + \frac{t}{2}, \eta + \frac{r}{2}\right) - \Phi\left(x - \frac{t}{2}, \eta - \frac{r}{2}\right),$$

$$(42) \quad \sigma_I(x, \eta, t, r) := \sigma\left(x + \frac{t}{2}, \eta + \frac{r}{2}\right) \overline{\sigma\left(x - \frac{t}{2}, \eta - \frac{r}{2}\right)}.$$

We have now all the tools to estimate the Wigner kernel of T_I .

Theorem 4.2. *Consider T_I the FIO of type I in (38). Fix $N \in \mathbb{N}$, $N > d$, and assume that the symbol $\sigma \in \Gamma^m(\mathbb{R}^{2d})$, with $m < -2(d+N)$. Let k_I be the associated Wigner kernel in (40). Then,*

$$(43) \quad |k_I(x, \xi, y, \eta)| \lesssim \frac{\langle (x, \eta) \rangle^{2N+m}}{\langle (x, \xi) - \chi(y, \eta) \rangle^{2N}}, \quad x, \xi, y, \eta \in \mathbb{R}^{2d}.$$

Proof. Since Φ is smooth, we can expand $\Phi\left(x + \frac{t}{2}, \eta + \frac{r}{2}\right)$ and $\Phi\left(x - \frac{t}{2}, \eta - \frac{r}{2}\right)$ into a Taylor series around (x, ξ) . Namely,

$$(44) \quad \Phi\left(x + \frac{t}{2}, \eta + \frac{r}{2}\right) = \Phi(x, \eta) + \frac{t}{2}\Phi_x(x, \eta) + \frac{r}{2}\Phi_\eta(x, \eta) + \Phi_2(x, \eta, t, r),$$

where the remainder Φ_2 is given by

$$(45) \quad \Phi_2(x, \eta, t, r) = \sum_{|\alpha|=2} \int_0^1 (1-\tau) \partial^\alpha \Phi\left(\left(x, \eta\right) + \tau\left(\frac{t}{2}, \frac{r}{2}\right)\right) d\tau \frac{(t, r)^\alpha}{2^3 \alpha!}.$$

Similarly,

$$(46) \quad \Phi\left(x - \frac{t}{2}, \eta - \frac{r}{2}\right) = \Phi(x, \eta) - \frac{t}{2}\Phi_x(x, \eta) - \frac{r}{2}\Phi_\eta(x, \eta) + \tilde{\Phi}_2(x, \eta, t, r),$$

with $\tilde{\Phi}_2$ defined as

$$(47) \quad \tilde{\Phi}_2(x, \eta, t, r) = \sum_{|\alpha|=2} \int_0^1 (1-\tau) \partial^\alpha \Phi\left(\left(x, \eta\right) - \tau\left(\frac{t}{2}, \frac{r}{2}\right)\right) d\tau \frac{(t, r)^\alpha}{2^3 \alpha!}.$$

Inserting the phase expansions above in (40) we obtain

$$(48) \quad k_I(x, \xi, y, \eta) = \int_{\mathbb{R}^{2d}} e^{-2\pi i[t \cdot (\xi - \Phi_x(x, \eta)) + r \cdot (y - \Phi_\eta(x, \eta))]} \tilde{\sigma}(x, \eta, t, r) dt dr$$

where $\tilde{\sigma}$ is defined as

$$(49) \quad \tilde{\sigma}(x, \eta, t, r) = e^{2\pi i[\Phi_2 - \tilde{\Phi}_2](x, \eta, t, r)} \sigma\left(x + \frac{t}{2}, \eta + \frac{r}{2}\right) \overline{\sigma\left(x - \frac{t}{2}, \eta - \frac{r}{2}\right)},$$

For $N \in \mathbb{N}$, $u = (t, r) \in \mathbb{R}^{2d}$, using the identity:

$$(1 - \Delta_u)^N e^{-2\pi i[(\xi - \Phi_x(x, \eta), y - \Phi_\eta(x, \eta)) \cdot (t, r)]} \\ = \langle 2\pi(\xi - \Phi_x(x, \eta), y - \Phi_\eta(x, \eta)) \rangle^{2N} e^{-2\pi i[(\xi - \Phi_x(x, \eta), y - \Phi_\eta(x, \eta)) \cdot (t, r)]},$$

we integrate by parts in (48) and obtain

$$k_I(x, \xi, y, \eta) = \frac{1}{\langle 2\pi(\xi - \Phi_x(x, \eta), y - \Phi_\eta(x, \eta)) \rangle^{2N}} \int_{\mathbb{R}^{2d}} e^{-2\pi i[(\xi - \Phi_x(x, \eta), y - \Phi_\eta(x, \eta)) \cdot (t, r)]} \\ \times (1 - \Delta_u)^N \tilde{\sigma}(x, \eta, t, r) dt dr.$$

The factor

$$(1 - \Delta_u)^N \tilde{\sigma}(z, u), \quad z = (x, \eta), \quad u = (t, r)$$

can be expressed as

$$e^{2\pi i[\Phi_2 - \tilde{\Phi}_2](z, u)} \sum_{|\alpha| + |\beta| + |\gamma| \leq 2N} C_{\alpha, \beta, \gamma} p(\partial_u^{|\alpha|}(\Phi_2 - \tilde{\Phi}_2)_z(u)) (\partial_u^\beta \sigma)(z + u/2) (\partial_u^\gamma \sigma)(z - u/2),$$

where $p(\partial_u^{|\alpha|}(\Phi_2 - \tilde{\Phi}_2)_z(u))$ is a polynomial made of derivatives w.r.t. u of $\Phi_2 - \tilde{\Phi}_2$ of order at most $|\alpha|$. By assumption,

$$|(\partial_u^\beta \sigma)(z + u/2) (\partial_u^\gamma \sigma)(z - u/2)| \lesssim \langle z + u/2 \rangle^{m - |\beta|} \langle z - u/2 \rangle^{m - |\gamma|},$$

which implies

$$|(1 - \Delta_u)^N \tilde{\sigma}(z, u)| \lesssim \sum_{|\alpha| + |\beta| + |\gamma| \leq 2N} \langle u/2 \rangle^{|\alpha|} \langle z - u/2 \rangle^{m - |\beta|} \langle z + u/2 \rangle^{m - |\gamma|} \\ \lesssim \sum_{|\beta| + |\gamma| \leq 2N} \langle u/2 \rangle^{2N - |\beta| - |\gamma|} \langle z - u/2 \rangle^{m - |\beta|} \langle z + u/2 \rangle^{m - |\gamma|} \\ \lesssim C_N \langle z - u/2 \rangle^{2N + m} \langle z + u/2 \rangle^{2N + m}.$$

Using the change of variables $u' = u/2 - z$, $du = 2^{2d} du'$,

$$\int_{\mathbb{R}^{2d}} \langle z - u/2 \rangle^{2N + m} \langle z + u/2 \rangle^{2N + m} du = 2^{2d} \int_{\mathbb{R}^{2d}} \langle u' \rangle^{2N + m} \langle (-2z) - u' \rangle^{2N + m} \lesssim \langle z \rangle^{2N + m}$$

where, for $v_s = \langle \cdot \rangle^s$, we used the weight convolution property $v_s * v_s \lesssim v_s$, for $s < 2d$, cf. [26, Lemma 11.1.1]. Hence,

$$\begin{aligned} |k_I(x, \xi, y, \eta)| &\leq \frac{1}{\langle 2\pi(\xi - \Phi_x(x, \eta), y - \Phi_\eta(x, \eta)) \rangle^{2N}} \int_{\mathbb{R}^{2d}} |(1 - \Delta_u)^N \tilde{\sigma}(x, \eta, t, r)| dt dr \\ &\lesssim \frac{\langle z \rangle^{2N+m}}{\langle 2\pi(\xi - \Phi_x(x, \eta), y - \Phi_\eta(x, \eta)) \rangle^{2N}} \\ &\asymp \frac{\langle z \rangle^{2N+m}}{\langle \chi_1(y, \eta) - x, \chi_2(y, \eta) - \xi \rangle^{2N}}. \end{aligned}$$

This gives the claim. \square

As a consequence,

Corollary 4.3. *Under the assumptions of Theorem 4.2, the estimate (7) holds true, hence $T_I \in FIO(\chi, N)$.*

Proof. It is an immediate consequence of Theorem 4.2, since $2N + m < 0$ so that $\langle (x, \eta) \rangle^{2N+m} \leq 1$, for every $x, \eta \in \mathbb{R}^d$. \square

5. FIOS OF TYPE II

In this section we focus on the L^2 -adjoint of a FIO of type I, which is a FIO of type II, written formally as

$$(50) \quad T_{II}f(x) = \int_{\mathbb{R}^{2d}} e^{-2\pi i[\Phi(y, \xi) - x\xi]} \tau(y, \xi) f(y) dy d\xi, \quad f \in \mathcal{S}(\mathbb{R}^d).$$

First, we shall work with symbols τ in the Hörmander class $S_{0,0}^0(\mathbb{R}^{2d})$, referring to [1] for their L^2 -boundedness.

Proposition 5.1. *Consider a FIO of type II as in (50), with $\tau \in S_{0,0}^0(\mathbb{R}^{2d})$. Then, for all $f, g \in \mathcal{S}(\mathbb{R}^d)$,*

$$(T_{II}f \otimes \bar{g})(x_1, x_2) = T_2(f \otimes \bar{g})(x_1, x_2),$$

$x = (x_1, x_2) \in \mathbb{R}^{2d}$, where T_2 is the FIO of type II given by

$$T_2F(x) = \int_{\mathbb{R}^{4d}} e^{-2\pi i[\Phi_2(y, \xi) - x\xi]} \tau_2(y, \xi) F(y) dy d\xi, \quad F \in \mathcal{S}(\mathbb{R}^{2d}),$$

$y = (y_1, y_2)$, $\xi = (\xi_1, \xi_2) \in \mathbb{R}^{2d}$, and Φ_2 is the tame phase on \mathbb{R}^{4d} given by

$$(51) \quad \Phi_2(y, \xi) = \Phi(y_1, \xi_1) + y_2 \xi_2;$$

whereas the symbol τ_2 is in $S_{0,0}^0(\mathbb{R}^{4d})$ and given by

$$(52) \quad \tau_2(y, \xi) = \tau(y_1, \xi_1) \otimes 1(y_2, \xi_2).$$

Proof. Let f, g be in $\mathcal{S}(\mathbb{R}^d)$. Using the Fourier inversion formula on g :

$$g(x_2) = \int_{\mathbb{R}^{2d}} g(y_2) e^{2\pi i(x_2 - y_2)\xi_2} dy_2 d\xi_2$$

we can write

$$\begin{aligned} (T_{II}f \otimes \bar{g})(x_1, x_2) &= \left(\int_{\mathbb{R}^{2d}} e^{-2\pi i[\Phi(y_1, \xi_1) - x_1 \xi_1]} \tau(y_1, \xi_1) f(y_1) dy_1 d\xi_1 \right) \overline{g(x_2)} \\ &= \int_{\mathbb{R}^{4d}} e^{-2\pi i[\Phi(y_1, \xi_1) + y_2 \xi_2 - (x_1 \xi_1 + x_2 \xi_2)]} f(y_1) \overline{g(y_2)} \tau(y_1, \xi_1) dy_1 dy_2 d\xi_1 d\xi_2. \end{aligned}$$

Observing that

$$x_1 \xi_1 + x_2 \xi_2 = x \xi,$$

we can write

$$\Phi(y_1, \xi_1) + y_2 \xi_2 = \Phi_2(y_1, y_2, \xi_1, \xi_2),$$

which is (51). Note that $\Phi_2 \in \mathcal{C}^\infty(\mathbb{R}^{4d})$ and satisfies (25) and (26), since

$$(53) \quad \partial_{y, \xi}^2 \Phi_2 = \begin{pmatrix} \partial_{y_1, y_1}^2 \Phi & 0_{d \times d} & \partial_{y_1, \xi_1}^2 \Phi & 0_{d \times d} \\ 0_{d \times d} & 0_{d \times d} & 0_{d \times d} & I_{d \times d} \\ \partial_{y_1, \xi_1}^2 \Phi & 0_{d \times d} & \partial_{\xi_1, \xi_1}^2 \Phi & 0_{d \times d} \\ 0_{d \times d} & I_{d \times d} & 0_{d \times d} & 0_{d \times d} \end{pmatrix}.$$

This means that Φ_2 is a tame phase function.

Finally, since both τ and the function constantly equal to 1 are in the Hörmander class $S_{0,0}^0(\mathbb{R}^{2d})$, it immediately follows that τ_2 belongs to $S_{0,0}^0(\mathbb{R}^{4d})$. \square

Using the same arguments as in the previous proposition, one can prove the issue below.

Proposition 5.2. *Under the same assumptions of Proposition 5.1, for all $f, g \in \mathcal{S}(\mathbb{R}^d)$,*

$$(f \otimes \overline{T_{II}g})(x_1, x_2) = T_2'(f \otimes \bar{g})(x_1, x_2),$$

$x = (x_1, x_2) \in \mathbb{R}^{2d}$, where the operator T_2' is the FIO of type II:

$$T_2'F(x) = \int_{\mathbb{R}^{4d}} e^{-2\pi i[\Phi_2'(y, \xi) - x \xi]} \tau_2'(y, \xi) F(y) dy d\xi, \quad F \in \mathcal{S}(\mathbb{R}^{2d}),$$

$y = (y_1, y_2)$, $\xi = (\xi_1, \xi_2) \in \mathbb{R}^{2d}$ and Φ_2' is the tame phase

$$\Phi_2'(y, \xi) = -\Phi(y_2, -\xi_2) + y_1 \xi_1,$$

whereas the symbol $\tau_2 \in S_{0,0}^0(\mathbb{R}^{4d})$ is given by

$$\tau_2'(y, \xi) = 1(y_1, \xi_1) \otimes \overline{\tau(y_2, -\xi_2)}.$$

Next, we study the composition of the FIOs T_2 and T_2' .

Proposition 5.3. *Consider the FIOs T_2 and T_2' defined in Propositions 5.1 and 5.2. Then their product T_2T_2' can be written as the following FIO of type II:*

$$(54) \quad T_2T_2'F(x) = \int_{\mathbb{R}^{4d}} e^{-2\pi i\Phi(y,\xi)-x\xi} \mathcal{T}(y,\xi) F(y) dy d\xi, \quad x \in \mathbb{R}^{2d},$$

for every $F \in \mathcal{S}(\mathbb{R}^{2d})$, with tame phase on \mathbb{R}^{4d} :

$$(55) \quad \Phi(y_1, y_2, \xi_1, \xi_2) = \Phi(y_1, \xi_1) - \Phi(y_2, -\xi_2)$$

and symbol

$$(56) \quad \mathcal{T}(y_1, y_2, \xi_1, \xi_2) = \tau(y_1, \xi_1) \overline{\tau(y_2, -\xi_2)} \in S_{0,0}^0(\mathbb{R}^{4d}).$$

Proof. Let $F \in \mathcal{S}(\mathbb{R}^{2d})$. We compute

$$\begin{aligned} T_2T_2'F(x_1, x_2) &= \int_{\mathbb{R}^{4d}} e^{-2\pi i[\Phi(y_1, \xi_1) + y_2\xi_2 - x_1\xi_1 - x_2\xi_2]} \tau(y_1, \xi_1) T_2'F(y_1, y_2) dy_1 dy_2 d\xi_1 d\xi_2 \\ &= \int_{\mathbb{R}^{4d}} e^{-2\pi i[\Phi(y_1, \xi_1) + y_2\xi_2 - x_1\xi_1 - x_2\xi_2]} \tau(y_1, \xi_1) \int_{\mathbb{R}^{4d}} e^{-2\pi i[-\Phi(z_2, -\eta_2) + z_1\eta_1 - y_1\eta_1 - y_2\eta_2]} \\ &\quad \times \overline{\tau(z_2, -\eta_2)} F(z_1, z_2) dz_1 dz_2 d\eta_1 d\eta_2 dy_1 dy_2 d\xi_1 d\xi_2 \\ &= \int_{\mathbb{R}^{8d}} e^{-2\pi i[\Phi(y_1, \xi_1) - \Phi(z_2, -\eta_2) + z_1\eta_1 + y_2\xi_2 - x_1\xi_1 - x_2\xi_2 - y_1\eta_1 - y_2\eta_2]} \\ &\quad \times \tau(y_1, \xi_1) \overline{\tau(z_2, -\eta_2)} F(z_1, z_2) dz d\eta dy dx \xi. \end{aligned}$$

Using the well-known formulae

$$\int_{\mathbb{R}^{2d}} e^{-2\pi i\eta_1(z_1 - y_1)} e^{-2\pi i\xi_2(y_2 - x_2)} d\eta_1 d\xi_2 dz_1 dy_2 = \delta_{y_1}(z_1) dz_1 \delta_{x_2}(y_2) dy_2,$$

we obtain

$$\begin{aligned} T_2T_2'F(x_1, x_2) &= \int_{\mathbb{R}^{4d}} e^{-2\pi i[\Phi(y_1, \xi_1) - \Phi(z_2, -\eta_2) - x_1\xi_1 - x_2\eta_2]} \tau(y_1, \xi_1) \overline{\tau(z_2, -\eta_2)} \\ &\quad \times F(y_1, z_2) dy_1 dz_2 d\xi_1 d\eta_2, \end{aligned}$$

which is (54). It is straightforward to check that the phase Φ in (55) is tame, that is, it satisfies the properties of Definition 2.3.

Since $\tau \in S_{0,0}^0(\mathbb{R}^d)$, then $\mathcal{T} \in S_{0,0}^0(\mathbb{R}^{4d})$. \square

Theorem 5.4. *Consider the type II FIO T_{II} in (50), with symbol $\tau \in S_{0,0}^0(\mathbb{R}^{2d})$ and tame phase Φ . Then*

$$W(T_{II}f, T_{II}g)(x, \xi) = \int_{\mathbb{R}^{2d}} k_{II}(x, \xi, s, z) W(f, g)(s, z) ds dz, \quad f, g \in \mathcal{S}(\mathbb{R}^d),$$

where

$$(57) \quad k_{II}(x, \xi, y, \eta) = \int_{\mathbb{R}^{2d}} e^{-2\pi i[\Phi(y+\frac{r}{2}, \xi+\frac{t}{2})-\Phi(y-\frac{r}{2}, \xi-\frac{t}{2})]} e^{2\pi i(tx+r\eta)} \\ \times \tau(y+\frac{r}{2}, \xi+\frac{t}{2}) \overline{\tau(y-\frac{r}{2}, \xi-\frac{t}{2})} dt dr.$$

Proof. Consider $f \in \mathcal{S}(\mathbb{R}^d)$ and use Proposition 5.3 and Remark 22 to compute

$$\begin{aligned} W(T_{II}f, T_{II}g)(x, \xi) &= \hat{A}_{1/2} T_2 T_2' \hat{A}_{1/2}^{-1} W(f, g)(x, \xi) \\ &= \int_{\mathbb{R}^d} (T_2 T_2' \hat{A}_{1/2}^{-1} W(f, g))(x + \frac{t}{2}, x - \frac{t}{2}) e^{-2\pi i \xi t} dt \\ &= \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^{4d}} e^{-2\pi i[\Phi(y_1, \eta_1) - \Phi(y_2, -\eta_2) - (x+t/2)\eta_1 - (x-t/2)\eta_2]} \tau(y_1, \eta_1) \overline{\tau(y_2, -\eta_2)} \right. \\ &\quad \times \left. (\hat{A}_{1/2}^{-1} W(f, g))(y_1, y_2) dy_1 dy_2 d\eta_1 d\eta_2 \right) e^{-2\pi i \xi t} dt \\ &= \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^{4d}} e^{-2\pi i[\Phi(y_1, \eta_1) - \Phi(y_2, -\eta_2) - (x+t/2)\eta_1 - (x-t/2)\eta_2]} \tau(y_1, \eta_1) \overline{\tau(y_2, -\eta_2)} \right. \\ &\quad \times \left. \left(\int_{\mathbb{R}^d} W(f, g)(y_1/2 + y_2/2, z) e^{2\pi i(y_1 - y_2)z} dz \right) dy_1 dy_2 d\eta_1 d\eta_2 \right) e^{-2\pi i \xi t} dt \\ &= \int_{\mathbb{R}^{6d}} e^{-2\pi i[\Phi(y_1, \eta_1) - \Phi(y_2, -\eta_2) - x\eta_1 - \frac{t}{2}\eta_1 - x\eta_2 + \frac{t}{2}\eta_2 - y_1 z + y_2 z + \xi t]} \tau(y_1, \eta_1) \overline{\tau(y_2, -\eta_2)} \\ &\quad \times W(f, g)(y_1/2 + y_2/2, z) dz dy_1 dy_2 d\eta_1 d\eta_2 dt. \end{aligned}$$

The change of variables $y_1/2 + y_2/2 = s$ gives

$$\begin{aligned} W(T_{II}f, T_{II}g)(x, \xi) &= 2^d \int_{\mathbb{R}^{6d}} e^{-2\pi i[\Phi(2s - y_2, \eta_1) - \Phi(y_2, -\eta_2) - x\eta_1 - \frac{t}{2}\eta_1 - x\eta_2 + \frac{t}{2}\eta_2 - (2s - y_2)z + y_2 z + \xi t]} \\ &\quad \times \tau(2s - y_2, \eta_1) \overline{\tau(y_2, -\eta_2)} W(f, g)(s, z) dz ds dy_2 d\eta_1 d\eta_2 dt. \end{aligned}$$

Next, observing that

$$\begin{aligned} \int_{\mathbb{R}^d} e^{-2\pi i(\frac{\eta_2}{2} - \frac{\eta_1}{2} + \xi)t} dt &= \int_{\mathbb{R}^d} e^{-2\pi i(\eta_2 - \eta_1 + 2\xi)\frac{t}{2}} dt = 2^d \int_{\mathbb{R}^d} e^{-2\pi i(\eta_2 - \eta_1 + 2\xi)t'} dt', \\ &= 2^d \int_{\mathbb{R}^d} e^{-2\pi i\eta_2 t'} M_{\eta_1 - 2\xi} 1(\eta_2) dt' = 2^d T_{\eta_1 - 2\xi} \hat{1}(\eta_2) \\ &= 2^d T_{\eta_1 - 2\xi} \delta(\eta_2). \end{aligned}$$

we reckon

$$\begin{aligned} W(T_{II}f, T_{II}g)(x, \xi) &= 2^{2d} \int_{\mathbb{R}^{4d}} e^{-2\pi i[\Phi(2s-y_2, \eta_1) - \Phi(y_2, 2\xi - \eta_1) + 2(\xi - \eta_1)x + 2(y_2 - s)z]} \\ &\quad \times \tau(2s - y_2, \eta_1) \overline{\tau(y_2, 2\xi - \eta_1)} W(f, g)(s, z) dz ds dy_2 d\eta_1 \\ &= \int_{\mathbb{R}^{2d}} k_{II}(x, \xi, s, z) W(f, g)(s, z) ds dz, \end{aligned}$$

where

$$\begin{aligned} k_{II}(x, \xi, s, z) &= 2^{2d} \int_{\mathbb{R}^{2d}} e^{-2\pi i[\Phi(2s-y_2, \eta_1) - \Phi(y_2, 2\xi - \eta_1) + 2(\xi - \eta_1)x + 2(y_2 - s)z]} \\ &\quad \times \tau(2s - y_2, \eta_1) \overline{\tau(y_2, 2\xi - \eta_1)} dy_2 d\eta_1. \end{aligned}$$

Next, we make the change of variables $s - y_2 = r/2$ and $\xi - \eta_1 = -t/2$ so that

$$\begin{aligned} k_{II}(x, \xi, s, z) &= \int_{\mathbb{R}^{2d}} e^{-2\pi i[\Phi(s+\frac{r}{2}, \xi+\frac{t}{2}) - \Phi(s-\frac{r}{2}, \xi-\frac{t}{2})]} e^{2\pi i(tx+rz)} \\ &\quad \times \tau(s + \frac{r}{2}, \xi + \frac{t}{2}) \overline{\tau(s - \frac{r}{2}, \xi - \frac{t}{2})} dt dr, \end{aligned}$$

which is (57). □

Theorem 5.5. *Consider T_{II} the FIO of type II in (50). Fix $N \in \mathbb{N}$, $N > d$, and assume that the symbol $\tau \in \Gamma^m(\mathbb{R}^{2d})$, with $m < -2(d + N)$. Let k_{II} be the associated Wigner kernel, given by (57). Then,*

$$(58) \quad |k_{II}(x, \xi, y, \eta)| \lesssim \frac{\langle (y, \xi) \rangle^{2N+m}}{\langle (y, \eta) - \chi(x, \xi) \rangle^{2N}}, \quad x, \xi, y, \eta \in \mathbb{R}^d.$$

Proof. We follow the pattern of the proof of Theorem 4.2. By (57) and using the Taylor expansions in (45) and (47) we reckon

$$(59) \quad k_{II}(x, \xi, y, \eta) = \int_{\mathbb{R}^{2d}} e^{2\pi i[r \cdot (\eta - \Phi_y(y, \xi)) + t \cdot (x - \Phi_\xi(y, \xi))]} \tilde{\tau}(y, \xi, r, t) dr dt,$$

where

$$\tilde{\tau}(y, \xi, r, t) = e^{-2\pi i[\Phi_2 - \tilde{\Phi}_2](y, \xi, r, t)} \times \tau(y + \frac{r}{2}, \xi + \frac{t}{2}) \overline{\tau(y - \frac{r}{2}, \xi - \frac{t}{2})},$$

and the reminders are given by:

$$\Phi_2(y, \xi, r, t) = \sum_{|\alpha|=2} \int_0^1 (1 - \tau) \partial^\alpha \Phi((y, \xi) + \tau(r, t)/2) d\tau \frac{(r, t)^\alpha}{2^3 \alpha!}$$

and

$$\tilde{\Phi}_2(y, \xi, r, t) = \sum_{|\alpha|=2} \int_0^1 (1 - \tau) \partial^\alpha \Phi((y, \xi) - \tau(r, t)/2) d\tau \frac{(r, t)^\alpha}{2^3 \alpha!}.$$

Again, for $N \in \mathbb{N}$ and setting $u = (r, t) \in \mathbb{R}^{2d}$, we have:

$$\begin{aligned} & (1 - \Delta_u)^N e^{2\pi i(\eta - \Phi_y(y, \xi), x - \Phi_\xi(y, \xi)) \cdot (r, t)} \\ &= \langle 2\pi(\eta - \Phi_y(y, \xi), x - \Phi_\xi(y, \xi)) \rangle^{2N} e^{2\pi i(\eta - \Phi_y(y, \xi), x - \Phi_\xi(y, \xi)) \cdot (r, t)}. \end{aligned}$$

Integrating by parts in (59), we get:

$$\begin{aligned} k_{II}(x, \xi, y, \eta) &= \frac{1}{\langle 2\pi(\eta - \Phi_y(y, \xi), x - \Phi_\xi(y, \xi)) \rangle^{2N}} \int_{\mathbb{R}^{2d}} e^{2\pi i(\eta - \Phi_y(y, \xi), x - \Phi_\xi(y, \xi)) \cdot (r, t)} \\ &\quad \times (1 - \Delta_u)^N \tilde{\tau}(y, \xi, r, t) dr dt. \end{aligned}$$

The same estimates of Theorem 4.2 yield to:

$$\begin{aligned} |k_{II}(x, \xi, y, \eta)| &\leq \frac{1}{\langle 2\pi(\eta - \Phi_y(y, \xi), x - \Phi_\xi(y, \xi)) \rangle^{2N}} \int_{\mathbb{R}^{2d}} |(1 - \Delta_u)^N \tilde{\tau}(y, \xi, r, t)| dr dt \\ &\asymp \frac{\langle (y, \xi) \rangle^{2N+m}}{\langle (y, \eta) - \chi(x, \xi) \rangle^{2N}}. \end{aligned}$$

□

From [13] we deduce

Corollary 5.6. *Under the assumptions of Theorem 5.5, the estimate (7) holds true, hence $T_{II} \in FIO(\chi, N)$.*

Proof. It follows from (58), since $2N + m < 0$ so that $\langle (y, \xi) \rangle^{2N+m} \leq 1$, for every $y, \xi \in \mathbb{R}^d$. □

ACKNOWLEDGEMENTS

The first three authors have been supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

REFERENCES

- [1] K. Asada and D. Fujiwara. On some oscillatory integral transformations in $L^2(\mathbf{R}^n)$. *Japan. J. Math. (N.S.)*, 4(2):299–361, 1978.
- [2] F. Bastianoni and E. Cordero. Characterization of smooth symbol classes by Gabor matrix decay. *J. Fourier Anal. Appl.*, 28(3), 2022. DOI:10.1007/s00041-021-09895-2.
- [3] L. Cohen, Generalized phase-space distribution functions, *J. Math. Phys.*, 7:781–786, 1966.
- [4] L. Cohen. *Time Frequency Analysis: Theory and Applications*, Prentice Hall, 1995.
- [5] E.J. Candés, L. Demanet and L. Ying. Fast computation of Fourier integral operators. *SIAM J. Sci. Comput.* 29(6): 2464–2493, 2007.
- [6] F. Concetti, G. Garello and J. Toft. Trace ideals for Fourier integral operators with non-smooth symbols II. *Osaka J. Math.*, 47(3):739–786, 2010.

- [7] F. Concetti and J. Toft. Trace ideals for Fourier integral operators with non-smooth symbols, “*Pseudo-Differential Operators: Partial Differential Equations and Time-Frequency Analysis*”, Fields Inst. Commun., Amer. Math. Soc., 52:255–264, 2007.
- [8] E. Cordero, G. Giacchi and L. Rodino. Wigner Representation of Schrödinger Propagators. *Submitted*. arXiv:2311.18383v2
- [9] E. Cordero, G. Giacchi and L. Rodino. Wigner Analysis of Operators. Part II: Schrödinger equations. *Submitted*. arXiv:2208.00505
- [10] E. Cordero, K. Gröchenig and F. Nicola. Approximation of Fourier integral operators by Gabor multipliers. *J. Fourier Anal. Appl.*, 18(4):661–684, 2012.
- [11] E. Cordero, F. Nicola and L. Rodino. Sparsity of Gabor representation of Schrödinger propagators. *Appl. Comput. Harmon. Anal.*, 26(3):357–370, 2009.
- [12] E. Cordero, F. Nicola and L. Rodino. Wave packet analysis of Schrödinger equations in analytic function spaces. *Adv. Math.*, 278:182–209, 2015
- [13] E. Cordero and N. Rodino. *Wigner Analysis of Operators. Part I: Pseudodifferential Operators and Wave Front Sets*. Appl. Comput. Harmon. Anal. 58 (2022) 85-123.
- [14] E. Cordero and N. Rodino. Characterization of Modulation Spaces By Symplectic Representations and Applications to Schrödinger Equations. *J. Funct. Anal.*, 284:109892, 2023.
- [15] E. Cordero, F. Nicola and L. Rodino. Time-frequency analysis of Fourier integral operators. *Commun. Pure Appl. Anal.*, 9(1):1–21, 2010.
- [16] E. Cordero, K. Gröchenig, F. Nicola and L. Rodino. Wiener algebras of Fourier integral operators. *J. Math. Pures Appl. (9)*, 99(2):219–233, 2013
- [17] E. Cordero, K. Gröchenig, F. Nicola and L. Rodino. Generalized Metaplectic Operators and the Schrödinger Equation with a Potential in the Sjöstrand Class, *J. Math. Phys.*, 55(8):081506, 17, 2014
- [18] E. Cordero, F. Nicola and L. Rodino, Sparsity of Gabor representation of Schrödinger propagators. *Appl. Comput. Harmon. Anal.*, 26(3):357–370, 2009.
- [19] E. Cordero and L. Rodino, *Time-Frequency Analysis of Operators*, De Gruyter Studies in Mathematics, 2020.
- [20] N.C. Dias, M. de Gosson, F. Luef and J.N. Prata. A metaplectic perspective of uncertainty principles in the Linear Canonical Transform domain. *Submitted*.
- [21] A. Fernández-Bertolin and E. Malinnikova. Dynamical versions of Hardy’s uncertainty principle: a survey. *Bull. Am. Math. Soc.* 58(3):357–375, 2021.
- [22] A. Córdoba and C. Fefferman. Wave packets and Fourier integral operators. *Comm. Partial Differential Equations*, 3(11):979–1005, 1978.
- [23] G. B. Folland. *Harmonic analysis in phase space*. Princeton Univ. Press, Princeton, NJ, 1989.
- [24] M. A. de Gosson. *Symplectic methods in harmonic analysis and in mathematical physics*, volume 7 of *Pseudo-Differential Operators. Theory and Applications*. Birkhäuser/Springer Basel AG, Basel, 2011.
- [25] M. de Gosson. *The Wigner Transform*. World Scientific Pub Co Inc, 2017.
- [26] K. Gröchenig. *Foundations of time-frequency analysis*. Birkhäuser Boston, Inc., Boston, MA, 2001.
- [27] K. Gröchenig. Time-Frequency Analysis of Sjöstrand’s Class. *Rev. Mat. Iberoamericana*, 22(2):703–724, 2006.
- [28] K. Gröchenig and Z. Rzeszotnik. Banach algebras of pseudodifferential operators and their almost diagonalization. *Ann. Inst. Fourier.* 58(7):2279-2314, 2008.

- [29] H. Knutsen. Notes on Hardy's uncertainty principle for the Wigner distribution and Schrödinger evolutions *Journal of Mathematical Analysis and Applications*, 525(1):127116, 2023.
- [30] S. G. Krantz and H. R. Parks, *The Implicit Function Theorem*, Birkhäuser Boston Inc, Boston, 2002.
- [31] B. Helffer. *Théorie spectrale pour des operateurs globalement elliptiques*. Astérisque, Société Mathématique de France, 1984.
- [32] L. Hörmander. Fourier integral operators I. *Acta Math.*, 127:79–183, 1971.
- [33] M.A. Shubin. *Pseudodifferential operators and spectral theory*. Springer-Verlag, Berlin, second edition, 2001.
- [34] J. Ville. Theorie et Applications de la notion de signal analytique. *Câbles et Transmissions*,2:61–74, 1948.
- [35] E. Wigner. *On the Quantum Correction for Thermodynamic Equilibrium*. *Phys. Rev.*, 40(5):749-759, 1932.
- [36] Z. C. Zhang. *Uncertainty Principle for the Free Metaplectic transformation*. Submitted.
- [37] Z. C. Zhang. *Linear Canonical Wigner Distribution Based Noisy LFM Signals Detection through the Output SNR Improvement Analysis*. Submitted.
- [38] Z. Zhang, Y. He. *Free Metaplectic Wigner Distribution: Definition and Heisenberg's Uncertainty Principles*.

UNIVERSITÀ DI TORINO, DIPARTIMENTO DI MATEMATICA, VIA CARLO ALBERTO 10, 10123 TORINO, ITALY

Email address: elena.cordero@unito.it

UNIVERSITÀ DI BOLOGNA, DIPARTIMENTO DI MATEMATICA, PIAZZA DI PORTA SAN DONATO 5, 40126 BOLOGNA, ITALY; UNIVERSITY OF LAUSANNE, SWITZERLAND; HES-SO SCHOOL OF ENGINEERING, RUE DE L'INDUSTRIE 21, SION, SWITZERLAND; CENTRE HOSPITALIER UNIVERSITAIRE VAUDOIS, SWITZERLAND

Email address: gianluca.giacchi2@unibo.it

UNIVERSITÀ DI TORINO, DIPARTIMENTO DI MATEMATICA, VIA CARLO ALBERTO 10, 10123 TORINO, ITALY

Email address: luigi.rodino@unito.it

UNIVERSITÀ DI TORINO, DIPARTIMENTO DI MATEMATICA, VIA CARLO ALBERTO 10, 10123 TORINO, ITALY

Email address: mario.valenzano@unito.it