

# Bayes-Optimal Fair Classification with Linear Disparity Constraints via Pre-, In-, and Post-processing

Xianli Zeng\*, Guang Cheng† and Edgar Dobriban‡

February 6, 2024

## Abstract

Machine learning algorithms may have disparate impacts on protected groups. To address this, we develop methods for Bayes-optimal fair classification, aiming to minimize classification error subject to given group fairness constraints. We introduce the notion of *linear disparity measures*, which are linear functions of a probabilistic classifier; and *bilinear disparity measures*, which are also linear in the group-wise regression functions. We show that several popular disparity measures—the deviations from demographic parity, equality of opportunity, and predictive equality—are bilinear.

We find the form of Bayes-optimal fair classifiers under a single linear disparity measure, by uncovering a connection with the Neyman-Pearson lemma. For bilinear disparity measures, Bayes-optimal fair classifiers become group-wise thresholding rules. Our approach can also handle multiple fairness constraints (such as equalized odds), and the common scenario when the protected attribute cannot be used at the prediction phase.

Leveraging our theoretical results, we design methods that learn fair Bayes-optimal classifiers under bilinear disparity constraints. Our methods cover three popular approaches to fairness-aware classification, via pre-processing (Fair Up- and Down-Sampling), in-processing (Fair Cost-Sensitive Classification) and post-processing (a Fair Plug-In Rule). Our methods control disparity directly while achieving near-optimal fairness-accuracy tradeoffs. We show empirically that our methods compare favorably to existing algorithms.

## Contents

|          |  |          |
|----------|--|----------|
| <b>1</b> | <b>Introduction</b>  | <b>2</b> |
| <b>2</b> | <b>Classification with a Protected Attribute</b>                   | <b>6</b> |
| 2.1      | Preliminaries  | 6        |
| <b>3</b> | <b>Generalized Neyman-Pearson Lemma and Fair Classification</b>    | <b>8</b> |
| 3.1      | Generalized Neyman-Pearson Lemma                                   | 8        |
| 3.2      | Fairness Measures with Linear and Bilinear Constraints             | 9        |
| <b>4</b> | <b>Fair Bayes-Optimal Classifiers and the Fair Pareto Frontier</b> | <b>9</b> |
| 4.1      | Form of Fair Bayes-Optimal Classifiers                             | 9        |
| 4.2      | Fairness-Accuracy Tradeoff: the Fair Pareto Frontier               | 11       |
| 4.3      | Extensions   | 12       |
| 4.3.1    | Multiple Fairness Constrains                                       | 12       |
| 4.3.2    | Cost-sensitive Risk  | 13       |
| 4.3.3    | Protected Attribute Not Available at the Prediction Phase          | 14       |

\*Department of Statistics and Data Science, University of Pennsylvania. zengx19911214@gmail.com.

†University of California, Los Angeles. guangcheng@ucla.edu.

‡Department of Statistics and Data Science, University of Pennsylvania. dobriban@wharton.upenn.edu.

|          |  |           |
|----------|--|-----------|
| <b>5</b> | <b>Bayes-Optimal Fair Classifiers via Pre-, In-, and Post-processing</b>       | <b>14</b> |
| 5.1      | Pre-processing: Fair Up- and Down-Sampling (FUDS)                              | 15        |
| 5.2      | In-Processing: Fair Cost-Sensitive Classification (FCSC)                       | 18        |
| 5.3      | Post-Processing: Fair Plug-in Thresholding Rule (FPIR)                         | 18        |
| <b>6</b> | <b>Empirical Experiments</b>   | <b>20</b> |
| 6.1      | Synthetic Datasets   | 20        |
| 6.2      | Empirical Data Analysis  | 21        |
| <b>7</b> | <b>Summary and Discussion</b>  | <b>24</b> |
| <b>A</b> | <b>Proofs</b>  | <b>29</b> |
| A.1      | Additional Lemmas  | 29        |
| A.2      | Proofs of Results from Section 3   | 32        |
| A.2.1    | Proof of Proposition 3.4   | 32        |
| A.3      | Proofs of Results from Section 4   | 33        |
| A.3.1    | Proof of Proposition 4.1   | 33        |
| A.3.2    | Proof of Theorem 4.2   | 34        |
| A.3.3    | Proof of Proposition 4.4   | 35        |
| A.3.4    | Proof of Theorem 4.6   | 35        |
| A.3.5    | Proof of Proposition 4.10  | 35        |
| A.4      | Proofs of Results from Section 5   | 36        |
| A.4.1    | Proof of Theorem 5.1   | 36        |
| A.4.2    | Proof of Theorem 5.3   | 37        |
| <b>B</b> | <b>Extensions</b>  | <b>37</b> |
| B.1      | Fair Bayes-optimal Classifier with Equalized Odds                              | 38        |
| B.1.1    | Proof of Proposition B.1   | 39        |
| B.1.2    | Proof of Theorem 4.7   | 40        |
| B.2      | Fair Bayes-optimal Classifier with a Multi-class Protected Attribute           | 44        |
| B.2.1    | Proof of Theorem 4.8   | 44        |
| B.3      | FUDS, FCSC and FPIR Algorithms for a Linear—But Not Bilinear—Disparity Measure | 45        |
| B.4      | Form of Fair Bayes-optimal Classifiers Without Distributional Assumptions      | 48        |
| <b>C</b> | <b>Bayes-optimal Classifiers for Synthetic Data</b>                            | <b>53</b> |
| C.1      | Demographic Parity   | 54        |
| C.2      | Equality of Opportunity  | 55        |
| C.3      | Predictive Equality  | 55        |

# 1 Introduction

Machine learning is increasingly deployed in algorithmic decision-making systems in high-stakes domains deeply impacting human lives, including in education (Tsai et al., 2020), finance (Ma et al., 2018), healthcare (Gupta and Mohammad, 2017), and judiciary systems (Angwin et al., 2016). While machine learning can improve decision-making efficiency, it has also surfaced many potential ethical risks.

One significant concern is algorithmic bias. Recent studies have revealed that without considering fairness, machine learning algorithms often make decisions that disadvantage vulnerable demographic groups, thereby exacerbating social injustice and potentially violating human rights (Angwin et al., 2016; Flores et al., 2016; Bhanot et al., 2021; Corbett-Davies et al., 2023).

In light of these risks, algorithmic fairness has attracted widespread attention from the public, government, and academia. Organizations like the White House (Executive Office of the President, 2016), and UNESCO (UNESCO, 2021) have all called for considering fairness when applying automated decision-making. In response, the emerging field of fair machine learning has seen significant development. An increasing amount of research focuses on defining quantitative fairness metrics for various applications (e.g.,

Table 1: Our contributions in comparison with prior theoretical work for Bayes-optimal classifier: [1]: Corbett-Davies et al. (2017); [2]: Menon and Williamson (2018); [3]: Chzhen et al. (2019); [4]: Jiang et al. (2020); [5]: Schreuder and Chzhen (2021); [6]: Wei et al. (2021) [7]: Chen et al. (2023); [8]: Xu and Strohmer (2023). The comparison is made based on scope of the theoretical framework, the fairness metrics considered, and whether theoretically optimal algorithms are proposed. Within each category, several criteria are considered. Our methods satisfy all desired criteria, while the prior works satisfy only some of them.

| References                              | [1] | [2] | [3] | [4] | [5] | [6] | [7] | [8] | This Work |
|---|-----|-----|-----|-----|-----|-----|-----|-----|-----------|
| <b>Scope of Theoretical Framework</b>   |     |     |     |     |     |     |     |     |           |
| Approximate Fairness                    |     | ✓   |     | ✓   |     | ✓   | ✓   | ✓   | ✓         |
| Explicit Form                           |     |     | ✓   | ✓   | ✓   |     |     | ✓   | ✓         |
| Multiple Constraints                    |     |     |     |     |     | ✓   | ✓   |     | ✓         |
| Pareto Analysis                         |     | ✓   |     |     |     |     |     | ✓   | ✓         |
| No Protected Attrib. $A$ at test time   |     | ✓   |     | ✓   |     | ✓   | ✓   |     | ✓         |
| <b>Fairness Metrics Considered</b>      |     |     |     |     |     |     |     |     |           |
| Demographic Parity                      | ✓   | ✓   |     | ✓   | ✓   | ✓   | ✓   | ✓   | ✓         |
| Equality of Opportunity                 | ✓   | ✓   | ✓   |     |     |     | ✓   |     | ✓         |
| Predictive Equality                     | ✓   |     |     |     |     |     |     |     | ✓         |
| Equalized Odds                          |     |     |     |     |     | ✓   | ✓   |     | ✓         |
| <b>Theoretically Optimal Algorithms</b> |     |     |     |     |     |     |     |     |           |
| Pre-processing                          |     |     |     |     |     | ✓   |     | ✓   | ✓         |
| In-processing                           |     |     |     | ✓   |     |     |     |     | ✓         |
| Post-processing                         | ✓   | ✓   | ✓   | ✓   | ✓   | ✓   | ✓   | ✓   | ✓         |

Calders et al., 2009; Dwork et al., 2012; Hardt et al., 2016; Zafar et al., 2017; Pleiss et al., 2017; Kusner et al., 2017; Zhang and Bareinboim, 2018; Narasimhan et al., 2020; Nandy et al., 2022, etc), designing algorithms that protect minority groups (e.g., Kamiran and Calders, 2012; Joseph et al., 2016; Zhang et al., 2018; Lahoti et al., 2019; Cho et al., 2020; Xian et al., 2023, etc), providing public software to help practitioners assess and improve fairness (e.g., Bellamy et al., 2019; Weerts et al., 2023, etc), and studying the theoretical underpinnings of algorithmic fairness (e.g., Corbett-Davies et al., 2017; Menon and Williamson, 2018; Schreuder and Chzhen, 2021; Zeng et al., 2022b, etc). We refer the readers to e.g., Mehrabi et al. (2021); Caton and Haas (2023), etc, for reviews of advances in fair machine learning.

In this paper, we develop methods for fair classification. In fair classification, the *fair Bayes-optimal classifier* is the method with the best possible accuracy under a given disparity constraint. This method serves as an ultimate “best possible” classifier. We consider the question of deriving Bayes-optimal classifiers under given group fairness constraints, which aim to equalize various quantities across protected groups.

Prior work by Corbett-Davies et al. (2017) showed that, for exact fairness with some specific disparity measures, fair Bayes-optimal classifiers are group-wise thresholding rules, but did not identify their exact form (see Table 1 for a summary of, and comparison with, related work). For the specific fairness criteria of demographic parity and equality of opportunity (see Section 2.1), Menon and Williamson (2018) characterized fair Bayes-optimal classifiers by a connection to cost-sensitive risks (again without an explicit form). Explicit forms of fair Bayes-optimal classifiers were derived for perfect demographic parity in Chzhen et al. (2019) and for perfect equality of opportunity in Schreuder and Chzhen (2021).

By considering the Wasserstein distance as the disparity measure, Jiang et al. (2020); Silvia et al. (2020); Xian et al. (2023) proved that the fair classification problem is equivalent to a Wasserstein-barycenter problem, and the perfectly fair Bayes-optimal classifier can be derived via optimal transport. Follow-up work by Xu and Strohmer (2023) further characterized the Pareto Frontier—i.e., the optimal fairness-accuracy tradeoff—using the Wasserstein geodesic. Rather than considering the fair Bayes-optimal classifier, Wei et al. (2021) considered the estimation of the Bayes-optimal score function and derived the optimal transformed score function that satisfies fairness constraints. In a selective classification framework, Rava et al. (2021) studied how to minimize outcomes without a decision, while equalizing the false selection rate among different protected groups. More recently, Chen et al. (2023) showed for certain disparity measures, Bayes-optimal fair classifiers can be derived by randomly flipping the output of the unconstrained Bayes-optimal classifier

(see Table 1).

Despite this progress, there is no systematic approach to derive explicit forms of Bayes-optimal fair classifiers for general disparity measures and arbitrary disparity levels. There is also no unified algorithmic framework to learn these classifiers from data.

In this paper, we propose a unified theoretical framework for deriving Bayes-optimal classifiers under various fairness measures, by leveraging a novel connection with the Neyman-Pearson argument for optimal hypothesis testing. We define a new notion of *linear disparity measures*, which are linear functions of a probabilistic classifier (See Definition 3.2), and show that finding the fair Bayes optimal classifier can then be recast as maximizing a linear functional subject to linear constraints. Since this is fundamentally similar to maximizing power subject to level constraints in optimal hypothesis testing, we can derive the explicit form of a fair Bayes-optimal classifier with any given level of disparity, for any given linear disparity measure.

We also introduce the notion of *bilinear disparity measures*, which are linear disparity measures that are also linear in the group-wise regression functions. We show that bilinear disparity measures include popular examples such as the deviations from demographic parity, equality of opportunity, predictive equality, and equalized odds. For bilinear disparity measures, Bayes-optimal fair classifiers become group-wise thresholding rules. We further show that our general theoretical framework can be extended to a wide range of scenarios (see Section 4.3): handling multiple fairness constraints (such as equalized odds, ensuring both equality of opportunity and predictive equality); and for the common scenario when the protected attribute cannot be used at the prediction phase.

In addition to deriving fair Bayes-optimal classifiers, we also investigate the *fair Pareto frontier*, which characterizes the optimal tradeoff between fairness and accuracy. This Pareto frontier comprises a set of fair Bayes-optimal classifiers. We study the tradeoff function that characterizes the best achievable misclassification rate for a given disparity level. We prove that for linear disparity measures, the tradeoff function is convex, highlighting that the marginal cost of fairness increases for smaller disparity levels.

In practice, Bayes-optimal classifiers need to be estimated from a finite dataset. For this, we develop methods covering three popular approaches to fairness-aware classification: pre-processing, in-processing, and post-processing (e.g., Caton and Haas, 2023).

- *Pre-processing methods* reduce biases implicit in the training data, and train classifiers on the debiased data. Examples include transformations (e.g., Feldman et al., 2015; Lum and Johndrow, 2016; Johndrow and Lum, 2019; Calmon et al., 2017, etc), fair representation learning (e.g., Zemel et al., 2013; Louizos et al., 2016; Creager et al., 2019, etc), and fair generative models (e.g., Xu et al., 2018; Sattigeri et al., 2019; Jang et al., 2021, etc). These methods are convenient to apply, as they do not change the training procedure. However, as argued for instance in Locatello et al. (2019), disparity could persist even after pre-processing.
- *In-processing algorithms* handle the fairness constraint during the training process. A commonly applied strategy is to incorporate fairness measures as a regularization term into the optimization objective (e.g., Goh et al., 2016; Zafar et al., 2019; Narasimhan, 2018; Celis et al., 2019; Cotter et al., 2019; Oneto et al., 2020; Cho et al., 2020, etc). However, fairness measures are often non-convex and even non-differentiable with respect to model parameters, so that scalable training is challenging. In the alternative approach of adversarial learning (e.g., Zhang et al., 2018; Wadsworth et al., 2018; Xu et al., 2019; Celis and Keswani, 2019, etc), the ability of the classifier to predict the protected attribute is minimized. Although this can achieve promising results through careful design, its training process can lack stability, as a min-max optimization problem is required to be solved (Cho et al., 2020). Other in-processing algorithms include domain-based training (Wang et al., 2020), where the protected attribute is explicitly encoded and its effect is mitigated.
- *Post-processing methods* fit classifiers to the training data in the standard way, and aim to mitigate disparities in the model output. The most frequently used post-processing algorithms include group-wise thresholding (e.g., Fish et al., 2016; Corbett-Davies et al., 2017; Valera et al., 2018; Menon and Williamson, 2018; Chzhen et al., 2019; Alabdulmohsin, 2020; Schreuder and Chzhen, 2021; Denis et al., 2021; Jang et al., 2022, etc) and post-processing via optimal transport (e.g., Jiang et al., 2020; Silvia et al., 2020; Xian et al., 2023; Xu and Strohmer, 2023, etc). In group-wise thresholding, after estimating the conditional probability of  $Y = 1$  for each protected group (i.e., the group-wise regression functions),

the method classifies  $\hat{Y} = 1$  when these probabilities exceed certain group-specific thresholds. On the other hand, the optimal transport-based method transforms these conditional probabilities into a fair Bayes-optimal classifier by employing optimal transport algorithms.

Motivated by our formulas for fair Bayes-optimal classifiers, we design three algorithms that mitigate algorithmic bias, via pre-processing (Fair Up- and Down-Sampling or FUDS), in-processing (Fair Cost-Sensitive Classification or FCSC) and post-processing (Fair Plug-In Rule or FPIR).

For the pre-processing algorithm FUDS, we characterize the perturbed data distribution for which the unconstrained Bayes-optimal classifier equals the fair Bayes-optimal classifier on the original target distribution. We prove that this perturbation can be obtained by adjusting the proportion of each demographic group defined by the label and protected attribute. This enables us to design an easy-to-use Fair Up- and Down-Sampling pre-processing algorithm.

For the in-processing algorithm FCSC, we observe that the fair Bayes-optimal classifier adjusts thresholds for each protected group to mitigate disparities. Leveraging that cost-sensitive classification achieves a similar effect, we show that the fair Bayes-optimal classifier can be achieved through a carefully designed group-wise cost-sensitive risk, leading to Fair Cost-Sensitive Classification (FCSC).

Finally, for the post-processing algorithm FPIR, since we derived the explicit form of fair Bayes-optimal classifiers, we can design a two-stage plug-in rule to estimate it. In the first stage, we estimate the feature-conditional probability of a positive outcome  $Y = 1$  for each protected group. In the second stage, the optimal fair decision boundary can be estimated by solving a one-dimensional search problem.

Compared to existing approaches, our proposed methods offer several advantages. First, they provably aim to learn optimal classifiers; controlling disparity with *minimal* impact on accuracy. Second, our methods allow for *direct* control over the level of disparity. This is beneficial for practitioners who seek to achieve a specific level of disparity. Third, our methods have a *strong* empirical performance, and compare favorably to existing methods in extensive numerical experiments.

We summarize our contributions below. We also provide a detailed comparison of our contributions with prior theoretical work for fair Bayes-optimal classifiers in Table 1.

1. **Unifying framework for fair Bayes-optimal classifiers with linear disparity measures:** We provide a unified framework for deriving Bayes-optimal classifiers under group fairness constraints. We introduce the notions of linear and bilinear disparity measures (Definitions 3.2 and 3.3) and show that several popular disparity measures—the deviations from demographic parity, equality of opportunity, predictive equality, and equalized odds—are bilinear (Proposition 3.4). We characterize Bayes-optimal classifiers under linear disparity measures (Theorem 4.2), by uncovering a connection with the Neyman-Pearson lemma. For bilinear disparity measures, Bayes-optimal fair classifiers become group-wise thresholding rules.
  2. **Extensions:** We illustrate that our approach can be extended to handle multiple fairness constraints (such as equalized odds (Theorem 4.7), which requires ensuring both equality of opportunity and predictive equality), and demographic parity with a multi-class protected attribute (Theorem 4.8). The result for equalized odds requires an intricate and lengthy argument, which unravels several fundamental properties of equality of opportunity and predictive equality. We also show how to handle cost-sensitive classification error (Corollary 4.9). Further, we derive Bayes-optimal fair classifiers for the common scenario when the protected attribute cannot be used at the prediction phase (Corollary 4.11).
- Finally, we give the form of Bayes-optimal fair classifiers under the general distributional assumptions that the features can belong to the decision boundary with nonzero probability; such as for discrete features. In this case, the optimal classifiers must be carefully randomized (Theorem B.9).
3. **Pareto frontier for fair classification and fairness-accuracy tradeoff:** We characterize the fair Pareto frontier that represents the optimal trade-off between accuracy and fairness. We show that the tradeoff function between misclassification and disparity is convex for linear disparity measures, indicating that the marginal cost of fairness increases as the level of disparity decreases.
  4. **Theoretically optimal fair classification via pre-, in-, and post-processing:** We propose pre-, in-, and post-processing algorithms aiming to recover the Bayes-optimal classifiers. Specifically

we introduce fair up-/down-sampling as a pre-processing method (Section 5.1), fair cost-sensitive classification as an in-processing method (Section 5.2), and a fair plug-in rule as a post-processing method (Section 5.3). Collectively, these methods establish a cohesive methodological framework for fair classification with linear disparity constraints.

5. **Empirical evaluation:** We evaluate our methods in numerical simulations and experiments on empirical datasets (Section 6). We compare with several existing methods, such as the disparate impact remover (Feldman et al., 2015), FAWOS (Salazar et al., 2021), KDE-based constrained optimization (Cho et al., 2020), adversarial training (Zhang et al., 2018), post-processing through flipping (Chen et al., 2023), and post-processing through optimal transport (Xian et al., 2023); on standard benchmark datasets. We observe that our methods compare favorably to existing methods. Our numerical results can be reproduced with the code provided at <https://github.com/XianliZeng/Bayes-Optimal-Fair-Classification>.

This paper significantly extends the results from our previous unpublished manuscript (Zeng et al., 2022a), introducing the notion of linear disparity measures, handling multiple constraints, and developing pre- and in-processing algorithms. Thus, this work supersedes Zeng et al. (2022a).

**Paper organization.** In Section 2.1, we review background concepts and notations for fair classification. Section 3 establishes the connection between the generalized Neyman-Pearson Lemma and fair Bayes-optimal classifiers. Our main results are presented in Section 4, providing explicit forms of fair Bayes-optimal classifiers under linear and bilinear group fairness measures. We characterize the fair Pareto frontier in Section 4.2. In Section 5, we propose methods for estimating fair Bayes-optimal classifiers through pre-, in- and post-processing methods. Simulation studies and empirical data analysis in Section 6 assess the finite sample performance of the proposed methods. We provide concluding remarks in Section 7. All proofs are in the supplementary material.

## 2 Classification with a Protected Attribute

In fair classification problems, two types of feature are observed: the usual feature  $X \in \mathcal{X}$  for some feature space  $\mathcal{X}$ , and the protected (or, sensitive) feature<sup>1</sup>  $A \in \mathcal{A} = \{0, 1\}$  with respect to which we aim to be fair. Here, we consider a binary classification problem with labels in  $\mathcal{Y} = \{0, 1\}$ . For example, in a credit lending setting,  $X$  may refer to common features such as education level and income,  $A$  may contain the race or gender of the individual and  $Y$  may correspond to the status of repayment or defaulting on a loan.

**Notation and Conventions.** For all  $a \in \mathcal{A}$ ,  $x \in \mathcal{X}$  and  $y \in \mathcal{Y}$ , we denote  $p_a := \mathbb{P}(A = a)$ ;  $p_{a,y} := \mathbb{P}(A = a, Y = y)$ ;  $\eta_a(x) := \mathbb{P}(Y = 1 \mid A = a, X = x)$ . Further, for all  $a \in \mathcal{A}$  and  $y \in \mathcal{Y}$ , we denote by  $\mathbb{P}_X(\cdot)$ ,  $\mathbb{P}_{X|A=a}(\cdot)$  and  $\mathbb{P}_{X|A=a, Y=y}(\cdot)$  the marginal distribution function of  $X$ , the conditional distribution function of  $X$  given  $A = a$ , and the conditional distribution of  $X$  given  $A = a, Y = y$ , respectively.

For two scalars  $a, b$ , we denote their maximum by  $\max\{a, b\}$  or  $a \vee b$ , and their minimum by  $\min\{a, b\}$  or  $a \wedge b$ . All quantities considered will be measurable with respect to appropriate sigma-algebras; and measurability may not always be mentioned in what follows.

### 2.1 Preliminaries

A *randomized classifier* outputs a prediction  $\hat{Y} \in \{0, 1\}$  with a certain probability based on the usual features  $X$  and the protected attribute  $A$ . Let  $\text{Bern}(p)$  be the Bernoulli distribution with success probability  $p \in [0, 1]$ , and let  $\mathcal{F}$  be the set of measurable functions  $f : \mathcal{X} \times \mathcal{A} \rightarrow [0, 1]$ . We denote by  $\hat{Y}_f = \hat{Y}_f(x, a) \in \{0, 1\}$  the prediction induced by the classifier  $f$ , which can be a random variable.

**Definition 2.1** (Randomized Classifier). *A randomized classifier  $f \in \mathcal{F}$  gives, for any  $x \in \mathcal{X}$  and  $a \in \mathcal{A}$ , the probability  $f(x, a)$  of predicting  $\hat{Y}_f = 1$  when observing  $X = x$  and  $A = a$ , i.e.,  $\hat{Y}_f \mid X = x, A = a \sim \text{Bern}(f(x, a))$ .*

<sup>1</sup>We consider a binary protected attribute here and extend our results to multi-class protected attributes in Section 4.3.



Without a fairness constraint, a *Bayes-optimal classifier* minimizes the misclassification rate, and is defined as any randomized classifier  $f^*$  satisfying  $f^* \in \underset{f \in \mathcal{F}}{\operatorname{argmin}} \mathbb{P}(Y \neq \hat{Y}_f)$ . The following classical result

characterizes Bayes-optimal classifiers in terms of the class-conditional probability functions  $\eta_a$  for which  $\eta_a(x) = \mathbb{P}(Y = 1 \mid A = a, X = x)$  for all  $x, a$  (see e.g., Devroye et al., 1996, etc). We denote by  $I(\cdot)$  the indicator function, which equals unity if its argument is true, and zero otherwise.

**Proposition 2.2.** *All Bayes-optimal classifiers  $f^* \in \mathcal{F}$  have the form  $f^*(x, a) = I(\eta_a(x) > 1/2) + \tau(x, a)I(\eta_a(x) = 1/2)$ , for all  $(x, a) \in \mathcal{X} \times \{0, 1\}$ , where  $\tau : \mathcal{X} \times \mathcal{A} \rightarrow [0, 1]$  is any measurable function.*

While Bayes-optimal classifiers are theoretically the best method, they depend on the usually unknown the class-conditional probability functions  $\eta_a$  of the population. However, Proposition 2.2 is still useful, because it suggests an approach to classification, by estimating the functions  $\eta_a$ , for all  $a$ , and defining the classifier in terms of the level sets of the estimates, see e.g., Audibert and Tsybakov (2007).

The Bayes-optimal classifier does not take fairness into account. To mitigate unfairness, a number of notions of parity have been considered, and we list below the ones we consider in this paper.

**Definition 2.3** (Demographic Parity (Calders et al., 2009)). *A classifier  $f$  satisfies demographic parity if its prediction  $\hat{Y}_f$  is probabilistically independent of the protected attribute  $A$ , so that  $\mathbb{P}_{X|A=1}(\hat{Y}_f = 1) = \mathbb{P}_{X|A=0}(\hat{Y}_f = 1)$ .*

**Definition 2.4** (Equality of Opportunity (Hardt et al., 2016)). *A classifier  $f$  satisfies equality of opportunity if it achieves the same true positive rate among protected groups:  $\mathbb{P}_{X|A=1, Y=1}(\hat{Y}_f = 1) = \mathbb{P}_{X|A=0, Y=1}(\hat{Y}_f = 1)$ .*

**Definition 2.5** (Predictive Equality (Corbett-Davies et al., 2017)). *A classifier  $f$  satisfies predictive equality if it achieves the same false positive rate among protected groups:  $\mathbb{P}_{X|A=1, Y=0}(\hat{Y}_f = 1) = \mathbb{P}_{X|A=0, Y=0}(\hat{Y}_f = 1)$ .*

**Definition 2.6** (Equalized Odds (Hardt et al., 2016)). *A classifier  $f$  satisfies equalized odds if it satisfies both equality of opportunity and predictive equality, achieving the same true positive rate and false positive rate among protected groups:  $\mathbb{P}_{X|A=1, Y=1}(\hat{Y}_f = 1) = \mathbb{P}_{X|A=0, Y=1}(\hat{Y}_f = 1)$ , and  $\mathbb{P}_{X|A=1, Y=0}(\hat{Y}_f = 1) = \mathbb{P}_{X|A=0, Y=0}(\hat{Y}_f = 1)$ .*

In applications, perfect fairness may require a large sacrifice of accuracy, and a “limited” disparate impact could be preferred. Following for instance Cho et al. (2020), we use the difference in the quantities that are equalized under perfect parity to measure disparity:

**Definition 2.7** (Disparity Measures). *We consider the following disparity measures: Demographic Disparity (DD), Disparity of Opportunity (DO), Predictive Disparity (PD), defined for probabilistic classifiers  $f$  as follows:*

$$\begin{aligned} \text{DD}(f) &= \mathbb{P}_{X|A=1}(\hat{Y}_f = 1) - \mathbb{P}_{X|A=0}(\hat{Y}_f = 1); \\ \text{DO}(f) &= \mathbb{P}_{X|A=1, Y=1}(\hat{Y}_f = 1) - \mathbb{P}_{X|A=0, Y=1}(\hat{Y}_f = 1); \\ \text{PD}(f) &= \mathbb{P}_{X|A=1, Y=0}(\hat{Y}_f = 1) - \mathbb{P}_{X|A=0, Y=0}(\hat{Y}_f = 1). \end{aligned} \tag{2.1}$$

Since equalized odds consists of two constraints, we will later use the maximum of the absolute values of the corresponding disparities as the measure of unfairness.

In the rest of this paper, we will use  $\text{Dis} : \mathcal{F} \rightarrow [0, 1]$  to refer to a generic disparity measure, such as any of the three measures from (2.1). Taking  $K \geq 1$  disparity measures  $\text{Dis}_k : \mathcal{F} \rightarrow [0, 1], k = 1, \dots, K$  into account, we say a classifier  $f$  satisfies  $\delta$ -disparity if  $\max_{k=1}^K |\text{Dis}_k(f)| \leq \delta$ . We next define the most accurate—equivalently, least inaccurate—classifiers that satisfy  $\delta$ -disparity.

**Definition 2.8** (Fair Bayes-optimal Classifier). *Consider any  $K \geq 1$  fairness measures  $\text{Dis}_k : \mathcal{F} \rightarrow [0, 1]$ ,  $k = 1, \dots, K$ . Then, a  $\delta$ -fair Bayes-optimal classifier  $f_{\text{Dis}, \delta}^*$  is defined by minimizing the misclassification error  $R(f) := \mathbb{P}(Y \neq \hat{Y}_f)$  over all classifiers that satisfy  $\delta$ -disparity:*

$$f_{\text{Dis}, \delta}^* \in \arg \min_{f \in \mathcal{F}} \left\{ R(f) : \max_{k=1}^K |\text{Dis}_k(f)| \leq \delta \right\}.$$

Just like for the unconstrained case, fair Bayes-optimal classifiers are theoretically the best method, but they do not directly lead to a feasible method. To be able to use classifiers inspired by fair Bayes-optimal ones, it would be helpful to know if there is a characterization similar to that from the unconstrained case in Proposition 2.2. We now turn to establishing this, by making a connection with the Neyman-Pearson lemma.

### 3 Generalized Neyman-Pearson Lemma and Fair Classification

#### 3.1 Generalized Neyman-Pearson Lemma

To derive Bayes-optimal classifiers under group fairness, we establish a connection with the Neyman-Pearson lemma (Neyman and Pearson, 1933); a theoretical result originally designed to derive most powerful tests with a given type I error. We use a slightly generalized version, which applies to optimization with linear constraints. For completeness and the reader's convenience, we present the full statement here; as this lemma is fundamental to our results.

**Lemma 3.1** (Generalized Neyman-Pearson Lemma (Lehmann and Romano, 2005; Shao, 2003)). *Let  $\phi_0, \phi_1, \dots, \phi_m$  be  $m + 1$  real-valued functions defined on a Euclidean space  $\mathcal{X}$ . Assume they are  $\nu$ -integrable for a  $\sigma$ -finite measure  $\nu$ . Let  $f^* \in \mathcal{F}$  be any function of the form*

$$f^*(x) = \begin{cases} 1, & \phi_0(x) > \sum_{i=1}^m c_i \phi_i(x); \\ \tau(x), & \phi_0(x) = \sum_{i=1}^m c_i \phi_i(x); \\ 0, & \phi_0(x) < \sum_{i=1}^m c_i \phi_i(x), \end{cases} \quad (3.1)$$

where  $0 \leq \tau(x) \leq 1$  for all  $x \in \mathcal{X}$ . For given constants  $t_1, \dots, t_m \in \mathbb{R}$ , let  $\mathcal{F}_{\leq}$  be the class of measurable functions  $f : \mathcal{X} \rightarrow \mathbb{R}$  satisfying

$$\int_{\mathcal{X}} f \phi_i d\nu \leq t_i, \quad i \in \{1, 2, \dots, m\}. \quad (3.2)$$

and  $\mathcal{F}_{=}$  be the set of functions in  $\mathcal{F}_{\leq}$  satisfying (3.2) with all inequalities replaced by equalities.

(1) If  $f^* \in \mathcal{F}_{=}$ , then

$$f^* \in \arg \max_{f \in \mathcal{F}_{=}} \int_{\mathcal{X}} f \phi_0 d\nu. \quad (3.3)$$

Moreover, if  $\nu(\{x : \phi_0(x) = \sum_{i=1}^m c_i \phi_i(x)\}) = 0$ , for all  $f' \in \arg \max_{f \in \mathcal{F}_{=}} \int_{\mathcal{X}} f \phi_0 d\nu$ ,  $f' = f^*$  almost everywhere with respect to  $\nu$ .

(2) Moreover, if  $c_i \geq 0$  for all  $i = 1, \dots, m$ , then

$$f^* \in \arg \max_{f \in \mathcal{F}_{\leq}} \int_{\mathcal{X}} f \phi_0 d\nu. \quad (3.4)$$

Moreover, if  $\nu(\{x : \phi_0(x) = \sum_{i=1}^m c_i \phi_i(x)\}) = 0$ , for all  $f' \in \arg \max_{f \in \mathcal{F}_{\leq}} \int_{\mathcal{X}} f \phi_0 d\nu$ , we have  $f'(x) = f^*(x)$  almost everywhere with respect to  $\nu$ .



## 3.2 Fairness Measures with Linear and Bilinear Constraints

To characterize Bayes-optimal fair classifiers, we want to find classifiers with the highest accuracy given a disparity level. Recalling the class-conditional regression functions such that for  $a \in \mathcal{A}$  and  $x \in \mathcal{X}$ ,  $\eta_a(x) = \mathbb{P}(Y = 1 \mid A = a, X = x)$ , the misclassification rate can be expressed as a linear functional of the classifier  $f$ , via

$$R(f) = \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, a)(1 - 2\eta_a(x))d\mathbb{P}_{X,A}(x, a) + C_{\mathbb{P}},$$

with  $C_{\mathbb{P}} = \int_{\mathcal{A}} \int_{\mathcal{X}} \eta_a(x)d\mathbb{P}_{X,A}(x, a)$ . As  $C_{\mathbb{P}}$  is independent of  $f$ , a  $\delta$ -fair Bayes-optimal classifier for a disparity measure  $\text{Dis}$  can be equivalently defined as

$$f_{\text{Dis},\delta}^* \in \arg \max_{f \in \mathcal{F}} \left\{ \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, a)(2\eta_a(x) - 1)d\mathbb{P}_{X,A}(x, a) : \text{Dis}(f) \leq \delta \right\}. \quad (3.5)$$

By taking  $\nu = \mathbb{P}_{X,A}$  and  $\phi_0(x, a) = 2\eta_a(x) - 1$  for all  $x, a$  in the objective from (3.4), we find that it reduces to the objective in (3.5); where the constraint set  $\mathcal{F}_{\leq}$  is to be specified.

As a result, the fair Bayes-optimal classifiers can be characterized by the generalized Neyman-Pearson lemma, *if the constraints are linear in the classifiers*. Motivated by this observation, we introduce the notion of *linear disparity measures*.

**Definition 3.2** (Linear Disparity Measure). *We call a disparity measure  $\text{Dis} : \mathcal{F} \rightarrow [0, 1]$  linear if for all  $\mathbb{P}$ , there is a weighting function  $w_{\text{Dis},\mathbb{P}} : \mathcal{X} \times \mathcal{A} \rightarrow \mathbb{R}$  such that for all  $f \in \mathcal{F}$ ,*

$$\text{Dis}(f) = \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, a)w_{\text{Dis},\mathbb{P}}(x, a)d\mathbb{P}_{X,A}(x, a). \quad (3.6)$$

Furthermore, we call a linear disparity measure *bilinear* if its weighting function is linear in the group-wise regression functions  $\eta_a$ :

**Definition 3.3** (Bilinear Disparity Measure). *A linear disparity measure  $\text{Dis} : \mathcal{F} \rightarrow [0, 1]$  is bilinear if for all  $\mathbb{P}$ , there is  $s_{\text{Dis},\mathbb{P},a}$  and  $b_{\text{Dis},\mathbb{P},a}$  depending on  $a \in \mathcal{A}$  such that for all  $x \in \mathcal{X}$ ,  $w_{\text{Dis},\mathbb{P}}(x, a) = s_{\text{Dis},\mathbb{P},a}\eta_a(x) + b_{\text{Dis},\mathbb{P},a}$ .*

Hereafter, we will omit the subscript  $\mathbb{P}$  in our notation for simplicity. We next show that the disparity measures from (2.1) are bilinear, with specific weighting functions.

**Proposition 3.4** (Classical Disparity Measures are Bilinear). *The disparity measures DD, DO, and PD from (2.1) are bilinear with weighting functions defined for all  $x, a$  by*

$$w_{\text{DD}}(x, a) = \frac{(2a - 1)}{p_a}; \quad w_{\text{DO}}(x, a) = \frac{(2a - 1)\eta_a(x)}{p_{a,1}}; \quad w_{\text{PD}}(x, a) = \frac{(2a - 1)(1 - \eta_a(x))}{p_{a,0}}. \quad (3.7)$$

*Thus, we have, for instance  $s_{\text{DD},a} = 0$  and  $b_{\text{DD},a} = (2a - 1)/p_a$  for all  $a$ ; the quantities  $s_{\text{Dis},a}$  and  $b_{\text{Dis},\mathbb{P},a}$  for the other disparity measures can be similarly read off from (3.7).*

Some of our results apply to linear disparity measures, and some to bilinear ones; thus we use both definitions. We will use (3.6) and Proposition 3.4 to develop a unified framework for deriving fair Bayes-optimal classifiers. For ease of exposition, we discuss a binary protected attribute and a single fairness constraint in the main text. However, our theory can also handle other scenarios, such as multiple fairness constraints and in particular a multi-class protected attribute. We briefly discuss these extensions in Section 4.3, and provide details in Appendix B.

## 4 Fair Bayes-Optimal Classifiers and the Fair Pareto Frontier

### 4.1 Form of Fair Bayes-Optimal Classifiers

In this section, we derive the explicit form of fair Bayes-optimal classifiers based on our theoretical framework. For simplicity, we present the results for a single linear disparity measure as per Definition 3.2, and discuss

cases with multiple constraints in Section 4.3.1. Furthermore, in the main text, we consider the setting where both random variables  $\eta_a(X)$  and  $w_{\text{Dis}}(X, a)$ , for  $a \in \{0, 1\}$ , have probability density functions over  $\mathcal{X}$ . This implies that the boundary sets of randomized classifiers are of measure zero, allowing us to only consider deterministic classifiers. We provide the—more involved—results applicable to the general case without this assumption in Appendix B.4.

Consider a linear disparity measure  $\text{Dis}$ . The expressions (3.5) and (3.6) motivate us to consider the following class of deterministic classifiers  $f_{\text{Dis},t}$  indexed by  $t \in \mathbb{R}$ , taking values

$$f_{\text{Dis},t}(x, a) = I\left(\eta_a(x) > \frac{1}{2} + \frac{t}{2}w_{\text{Dis}}(x, a)\right) \quad (4.1)$$

for all  $x \in \mathcal{X}$  and  $a \in \{0, 1\}$ . Indeed, by taking  $\phi_0(x, a) = 2\eta_a(x) - 1$  and  $\phi_1(x, a) = w_{\text{Dis}}(x, a)$  for all  $x, a$  in (3.3) and (3.2), respectively, the generalized Neyman-Pearson lemma suggests that a fair Bayes-optimal classifier is of the form  $f_{\text{Dis},t}$  for some  $t$ . To make this precise, it will be helpful to define a disparity function  $D_{\text{Dis}} : \mathbb{R} \rightarrow [-1, 1]$  measuring the disparity level of  $f_{\text{Dis},t}$  as a function of  $t$ , such that for all  $t \in \mathbb{R}$ :

$$\begin{aligned} D_{\text{Dis}}(t) &:= \text{Dis}(f_{\text{Dis},t}) = \int_{\mathcal{A}} \int_{\mathcal{X}} \left[ w_{\text{Dis}}(x, a) \cdot I\left(\eta_a(x) > \frac{1}{2} + \frac{t}{2}w_{\text{Dis}}(x, a)\right) \right] d\mathbb{P}_{X,A}(x, a) \\ &= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} \left[ w_{\text{Dis}}(x, a) \cdot I\left(\eta_a(x) > \frac{1}{2} + \frac{t}{2}w_{\text{Dis}}(x, a)\right) \right] d\mathbb{P}_{X|A=a}(x). \end{aligned} \quad (4.2)$$

The following proposition characterizes how the accuracy and disparity of  $f_{\text{Dis},t}$  depends on  $t$ , and will be a crucial stepping stone towards precisely characterizing the form of fair Bayes-optimal classifiers.

**Proposition 4.1** (Properties of Risk and Disparity). *Let  $f_{\text{Dis},t}$  and  $D_{\text{Dis}}$  be defined in (4.1) and (4.2), respectively. Then, as a function of  $t$ ,*

- (1) *the disparity  $D_{\text{Dis}}(t)$  is monotone non-increasing;*
- (2) *the misclassification error  $R(f_{\text{Dis},t})$  is monotone non-increasing on  $(-\infty, 0)$  and monotone non-decreasing on  $[0, \infty)$ .*

Now, consider finding a classifier  $f_{\text{Dis},t}$  that meets fairness constraints while minimizing the misclassification error. By Proposition 4.1, it is enough to minimize  $|t|$ ; indeed this can be seen considering the cases of  $|D_{\text{Dis}}(0)| \leq \delta$  (in which case the optimal  $t = 0$ ),  $D_{\text{Dis}}(0) < -\delta$  (in which case the optimal  $t$  is negative), and  $D_{\text{Dis}}(0) > \delta$  (in which case the optimal  $t$  is positive). This motivates us to define the function  $t_{\text{Dis}} : [0, \infty) \rightarrow \mathbb{R}$  as an “inverse function” of  $|D_{\text{Dis}}(t)|$  such that for all  $\delta \geq 0$ ,

$$t_{\text{Dis}}(\delta) = \arg \min_t \{|t| : |D_{\text{Dis}}(t)| \leq \delta\}. \quad (4.3)$$

With these preparations, we can provide the form of fair Bayes-optimal classifiers for linear disparity measures.

**Theorem 4.2** (Form of Fair Bayes-optimal Classifiers for Linear Disparity Measures). *For a given linear disparity measure  $\text{Dis}$  as per Definition 3.2, suppose that for  $a \in \{0, 1\}$ , both  $\eta_a(X)$  and  $w_{\text{Dis}}(X, a)$  have probability density functions on  $\mathcal{X}$ . Recalling  $f_{\text{Dis},t}$  from (4.1) and  $t_{\text{Dis}} : [0, \infty) \rightarrow \mathbb{R}$  from (4.3), for any  $\delta \geq 0$ ,  $f_{\text{Dis},t_{\text{Dis}}(\delta)}$  is a  $\delta$ -fair Bayes-optimal classifier as per Definition 2.8. This Bayes-optimal classifier  $f_{\text{Dis},\delta}^*$  of the form*

$$f_{\text{Dis},\delta}^*(x, a) := f_{\text{Dis},t_{\text{Dis}}(\delta)}(x, a) = I\left(\eta_a(x) > \frac{1}{2} + \frac{t_{\text{Dis}}(\delta)}{2}w_{\text{Dis}}(x, a)\right), \quad (4.4)$$

for all  $x, a$ . Furthermore, when the disparity measure is bilinear as per Definition 3.3, the above  $\delta$ -fair Bayes-optimal classifier simplifies to a group-wise thresholding rule, such that for all  $x, a$ ,

$$f_{\text{Dis},\delta}^*(x, a) = I\left(\eta_a(x) > \frac{1 + b_{\text{Dis},a} \cdot t_{\text{Dis}}(\delta)}{2 - s_{\text{Dis},a} \cdot t_{\text{Dis}}(\delta)}\right). \quad (4.5)$$

Thus, similarly to the unconstrained case from Proposition 2.2, the fair Bayes-optimal classifiers for bilinear disparity measures are group-wise thresholding rules. Clearly, the accuracy is maximized by predicting the more likely class in each group. Moreover, mitigating disparity over protected groups necessitates shifting the thresholds. This shift depends delicately on the disparity measure and the population distribution. By incorporating the expressions from Proposition 3.4, we conclude the following corollary for the common disparity measures.

**Corollary 4.3** (Bayes-optimal Classifiers for Three Common Disparity Measures). *For any  $\delta > 0$ , the group-wise thresholding rules such that for all  $x, a$ ,*

$$f_{\text{DD},\delta}^*(x, a) = I\left(\eta_a(x) > \frac{1}{2} + \frac{(2a-1)t_{\text{DD}}(\delta)}{p_a}\right); \quad f_{\text{DO},\delta}^*(x, a) = I\left(\eta_a(x) > \frac{p_{a,1}}{2p_{a,1} - (2a-1)t_{\text{DO}}(\delta)}\right);$$

$$f_{\text{PD},\delta}^*(x, a) = I\left(\eta_a(x) > \frac{p_{a,0} + (2a-1)t_{\text{PD}}(\delta)}{2p_{a,0} + (2a-1)t_{\text{PD}}(\delta)}\right),$$

with

$$t_{\text{DD}}(\delta) = \arg \min_t \left\{ |t| : \left| \sum_{a \in \{0,1\}} \int_X I\left(\eta_a(x) > \frac{1}{2} + \frac{(2a-1)t}{2p_a}\right) d\mathbb{P}_{X|A=a}(x) \right| \leq \delta \right\};$$

$$t_{\text{DO}}(\delta) = \arg \min_t \left\{ |t| : \left| \sum_{a \in \{0,1\}} \int_X I\left(\eta_a(x) > \frac{p_{a,1}}{2p_{a,1} - (2a-1)t}\right) d\mathbb{P}_{A=a,Y=1}(x) \right| \leq \delta \right\};$$

$$t_{\text{PD}}(\delta) = \arg \min_t \left\{ |t| : \left| \sum_{a \in \{0,1\}} \int_X I\left(\eta_a(x) > \frac{p_{a,0} + (2a-1)t}{2p_{a,0} + (2a-1)t}\right) d\mathbb{P}_{A=a,Y=0}(x) \right| \leq \delta \right\},$$

are  $\delta$ -fair Bayes-optimal classifiers under DD, DO and PD from (2.1), respectively.

## 4.2 Fairness-Accuracy Tradeoff: the Fair Pareto Frontier

Theorem 4.2 specifies fair Bayes-optimal classifiers for a given disparity level  $\delta$ . A core challenge is to set a suitable value for  $\delta$ , balancing accuracy and fairness. Identifying a suitable balance among multiple objectives can be viewed through the Pareto frontier. In the context of fair classification with a disparity measure  $\text{Dis}$ , we consider the following ordering relation on classifiers. A classifier  $f_1$  is *Pareto dominant* over another classifier  $f_2$ , if one of the following conditions is met:

- (1)  $R(f_1) < R(f_2)$  and  $\text{Dis}(f_1) \leq \text{Dis}(f_2)$ ,      or      (2)  $R(f_1) \leq R(f_2)$  and  $\text{Dis}(f_1) < \text{Dis}(f_2)$ .

A classifier  $f$  is considered *Pareto optimal* if it is not dominated by any other classifier. The collection of all such Pareto optimal classifiers constitutes the *fair Pareto frontier* (FPF). Formally, the FPF is the solution of the following vector objective optimization problem, with respect to the ordering defined above:

$$\arg \min_{f \in \mathcal{F}_\delta} (R(f), \text{Dis}(f)). \quad (4.6)$$

The following proposition characterizes the fair Pareto frontier:

**Proposition 4.4** (Fair Pareto Frontier). *Let  $t_{\text{Dis}}$  be defined in (4.2), and let  $\underline{t}_0 = \min(0, t_{\text{Dis}}(0))$  and  $\bar{t}_0 = \max(0, t_{\text{Dis}}(0))$ . Then the fair Pareto frontier FPF from (4.6) includes all classifiers  $f_{\text{Dis},t}$  for  $t \in [\underline{t}_0, \bar{t}_0]$ , i.e., we have*

$$\{f_{\text{Dis},t} : t \in [\underline{t}_0, \bar{t}_0]\} \subset \text{FPF}.$$

Moreover, for any  $f_{\text{FPF}} \in \text{FPF}$ , there is  $t \in [\underline{t}_0, \bar{t}_0]$  such that  $f_{\text{Dis},t}$  has the same classification error and disparity as  $f_{\text{FPF}}$ , i.e.,

$$R(f_{\text{Dis},t}) = R(f_{\text{FPF}}) \quad \text{and} \quad \text{Dis}(f_{\text{Dis},t}) = \text{Dis}(f_{\text{FPF}}).$$

Now, consider an equivalence relation “ $\sim$ ” between classifiers  $f_1, f_2 \in \mathcal{F}$  such that  $f_1 \sim f_2$  if and only if  $R(f_1) = R(f_2)$  and  $\text{Dis}(f_1) = \text{Dis}(f_2)$ . Then, for each equivalence class determined by “ $\sim$ ”, there is a minimizer of (4.6) over  $\{f_{\text{Dis},t} : t \in [\underline{t}_0, \bar{t}_0]\}$ . Hence, the classifiers  $f_{\text{Dis},t}$  fully characterize the Pareto frontier.

We next study the cost of fairness as a function of the disparity level. We define the following trade-off function that measures the best achievable performance for a given disparity level.

**Definition 4.5** (Tradeoff Function). *For a disparity measure  $\text{Dis}$  and any  $\delta \geq 0$ , let*

$$T(\delta) = \inf\{R(f) : \text{Dis}(f) \leq \delta\}. \quad (4.7)$$

From Proposition 4.4, it follows that for all  $\delta \geq 0$ ,  $T(\delta) = R(f_{\text{Dis},t_{\text{Dis}}(\delta)})$ . The tradeoff function plays a crucial role in characterizing the misclassification rate and disparity along the Pareto frontier. This function is monotone non-increasing by definition, as a more stringent fairness constraint cannot increase accuracy. The following proposition establishes that this tradeoff function is *convex* for linear disparity measures. Thus, the lower the disparity level, the larger the additional accuracy cost of fairness.

**Proposition 4.6.** *If the disparity measure  $\text{Dis}$  is linear, then the tradeoff function  $T$  is convex on  $[0, \text{Dis}(0)]$ .*

### 4.3 Extensions

In this section, we discuss extensions of our theoretical framework to three scenarios: (1) Multiple fairness constrains; (2) Cost-sensitive loss; and (3) The protected attribute  $A$  is excluded from the predictive features. We provide the insights and main results here and defer details to Appendix B.

#### 4.3.1 Multiple Fairness Constrains

In many applications, achieving fairness is a multifaceted challenge, needing simultaneous consideration of different fairness metrics. Additionally, there can be multiple protected attributes, which are often multi-class rather than binary. As an example, the U.S. Equal Employment Opportunity Commission recognizes 11 discrimination types, mostly with multi-class attributes. In such scenarios, managing multiple fairness constraints becomes crucial. To illustrate how our method can be extended to handle multiple fairness constraints, we consider two examples: (1) equalized odds, which requires ensuring *both* equality of opportunity *and* predictive equality, and (2) demographic parity with a multi-class protected attribute.

**Equalized Odds:** Recalling disparity of opportunity (DO) and predictive disparity (PD) from (2.1), a classifier  $f$  satisfies the  $\delta$ -parity constraint with respect to equalized odds from Definition 2.6 if

$$\max(|\text{DO}(f)|, |\text{PD}(f)|) \leq \delta.$$

The corresponding  $\delta$ -fair Bayes-optimal classifiers are then defined as

$$f_{\text{DEO},\delta}^* \in \arg \min_{f \in \mathcal{F}} \{R(f) : \max\{|\text{DO}(f)|, |\text{PD}(f)|\} \leq \delta\}.$$

By taking  $\phi_0, \phi_1, \phi_2$  such that for all  $x, a$ ,  $\phi_0(x, a) = 2\eta_a(x) - 1$ ,  $\phi_1(x, a) = (2a - 1)\eta_a(x)/p_{a,1}$ ,  $\phi_2(x, a) = (2a - 1)(1 - \eta_a(x))/p_{a,0}$  and applying the generalized Neyman Pearson lemma, after some intricate calculations, we can derive the following result.

**Theorem 4.7** (Bayes-optimal Classifiers for Equalized Odds). *For any  $\delta > 0$ , there is  $t_{\text{DEO},1}(\delta) \in (-p_{0,1}, p_{1,1})$  and  $t_{\text{DEO},2}(\delta) \in (-p_{1,0}, p_{0,0})$  such that a  $\delta$ -fair Bayes-optimal classifier takes values, for all  $x, a$ ,*

$$f_{\text{DEO}}^*(x, a) = I \left( \eta_a(x) > \frac{p_{a,1}p_{a,0} + (2a - 1)t_{\text{DEO},2}(\delta)p_{a,1}}{2p_{a,1}p_{a,0} + (2a - 1)(t_{\text{DEO},2}(\delta)p_{a,1} - t_{\text{DEO},1}(\delta)p_{a,0})} \right).$$

*The values of  $t_{\text{DEO},1}(\delta)$  and  $t_{\text{DEO},2}(\delta)$  are provided in (B.9) in Appendix B.1.*

The proof of Theorem 4.7 requires studying intricate properties of the DO and PD, as detailed in a series of lemmas (Proposition B.1, Lemmas B.2, B.3, B.4), as well discussing the seven cases that are involved in

the definitions of  $t_{\text{DEO},1}(\delta)$  and  $t_{\text{DEO},2}(\delta)$  in (B.9). Briefly, we define a candidate class of potentially Bayes-optimal classifiers, parametrized by two scalars  $t_1, t_2$ ; and show that all Bayes-optimal classifiers belong to this class in Lemma B.3. We show that the DO and DP of these classifiers is continuous and monotone non-increasing as a function of either of these arguments while holding the other one fixed (Proposition B.1). This leads to the definitions of  $t_{\text{DEO},1}(\delta)$  and  $t_{\text{DEO},2}(\delta)$  in (B.9); where the claim that they are well-defined is proved in Lemma B.4. Then, we show that the most efficient approach, in terms of minimizing changes to PD, to attain a particular level of DO involves solely varying the value of  $t_1$ . Conversely, when aiming to achieve a specific level of PD, the optimal strategy is to solely adjust the value of  $t_2$  (Lemma B.2). The proof of Theorem 4.7 follows by careful arguments leveraging all these properties.

**Demographic Parity with Multi-class Protected Attribute:** Next, we illustrate how our approach can be extended to handle demographic parity with a multi-class protected attribute. For a multi-class protected attribute  $a \in \mathcal{A} = \{1, 2, \dots, |\mathcal{A}|\}$ , we say that a classifier satisfies *perfect* demographic parity if the  $|\mathcal{A}| - 1$  linear constraints

$$\mathbb{P}(\widehat{Y}_f = 1 \mid A = a) = \mathbb{P}(\widehat{Y}_f = 1 \mid A = 1) \quad \text{for } a = 2, 3, \dots, |\mathcal{A}|$$

hold. A fair Bayes-optimal classifier is defined as

$$f_{\text{DD},|\mathcal{A}|}^* \in \arg \min_{f \in \mathcal{F}} \left\{ \mathbb{P}(Y \neq \widehat{Y}_f) : \sum_{a=2}^{|\mathcal{A}|} \left| \mathbb{P}(\widehat{Y}_f = 1 \mid A = a) - \mathbb{P}(\widehat{Y}_f = 1 \mid A = 1) \right| = 0 \right\}.$$

By leveraging the generalized Neyman-Pearson lemma, we can show the following result:

**Theorem 4.8** (Bayes-optimal Classifier for Demographic Parity with a Multi-class Protected Attribute). *Let for  $a \in \mathcal{A}$ ,  $p_a = \mathbb{P}(A = a)$ . Then, there are constants  $(t_{\text{DD},a})_{a=1}^{|\mathcal{A}|}$  with  $\sum_{a=1}^{|\mathcal{A}|} t_{\text{DD},a} = 0$ , such that, for all  $a \in \{2, \dots, |\mathcal{A}|\}$ ,*

$$\mathbb{P}_{X|A=a} \left( \eta_a(X) > \frac{1}{2} + \frac{t_{\text{DD},a}}{2p_a} \right) = \mathbb{P}_{X|A=1} \left( \eta_1(x) > \frac{1}{2} + \frac{t_{\text{DD},1}}{2p_1} \right). \quad (4.8)$$

Moreover, there is a fair Bayes-optimal classifier such that for all  $x, a$ ,

$$f_{\text{DD},t_{\text{DD},1},\dots,t_{\text{DD},|\mathcal{A}|}}(x, a) = I \left( \eta_a(x) > \frac{1}{2} + \frac{t_{\text{DD},a}}{2p_a} \right).$$

### 4.3.2 Cost-sensitive Risk

Cost-sensitive classification is useful when the consequences of false negatives and positives are unequally important, such as in certain medical applications. For a given cost parameter  $c \in [0, 1]$ , the cost-sensitive classification error of a classifier  $f$  is defined as:

$$R_c(f) = c \cdot \mathbb{P}(\widehat{Y}_f = 1, Y = 0) + (1 - c) \cdot \mathbb{P}(\widehat{Y}_f = 0, Y = 1). \quad (4.9)$$

When  $c = 1/2$ , cost-sensitive risk reduces to the usual mis-classification error.

Taking fairness into account, a  $\delta$ -fair Bayes-optimal classifier for the cost-sensitive risk is defined as  $f_{\text{Dis},\delta,c}^* \in \arg \min_{f \in \mathcal{F}_\delta} R_c(f)$ , or equivalently  $f_{\text{Dis},\delta,c}^* \in \arg \max_{f \in \mathcal{F}_\delta} \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, a) (\eta_a(x) - c) d\mathbb{P}_{X,A}(x, a)$ . Since this is a linear objective, our approach extends seamlessly to find fair Bayes-optimal classifiers.

**Corollary 4.9** (Fair Bayes-optimal Classifiers for a Cost-sensitive Risk). *For a given linear disparity measure Dis, suppose that for  $a \in \{0, 1\}$ , both  $\eta_a(X)$  and  $w_{\text{Dis}}(X, a)$  have probability density functions on  $\mathcal{X}$ . Let  $D_{\text{Dis},c} : \mathbb{R} \rightarrow [-1, 1]$  be defined for all  $t$  as*

$$D_{\text{Dis},c}(t) := \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} [w_{\text{Dis}}(x, a) \cdot I(\eta_a(x) > c + t w_{\text{Dis}}(x, a))] d\mathbb{P}_{X|A=a}(x),$$

and let  $t_{\text{Dis},c} : [0, \infty) \rightarrow \mathbb{R}$  be defined as, for all  $\delta \geq 0$ ,  $t_{\text{Dis},c}(\delta) = \arg \min_t \{|t| : |D_{\text{Dis},c}(t)| \leq \delta\}$ . Then, for any  $\delta > 0$ , there exists a  $\delta$ -fair Bayes-optimal classifier  $f_{\text{Dis},c}^*$  under cost-sensitive risk, such that for all  $x, a$ ,

$$f_{\text{Dis},c}^*(x, a) = I(\eta_a(x) > c + t_{\text{Dis},c} \cdot w_{\text{Dis},c}(x, a)),$$

### 4.3.3 Protected Attribute Not Available at the Prediction Phase

In the previous discussion, we assumed that the protected feature is available for training and prediction; and that it is allowed to use it in both phases. Although this is a common setting (e.g., Hardt et al. (2016); Corbett-Davies et al. (2017); Cho et al. (2020), etc.), in certain cases, there may be ethical or legal considerations that invalidate it. However, our framework can also be applied to fair classification when the protected attribute cannot be used at the prediction phase. When the protected attribute is not available for *training*, the problem is very different, and it may require inferring the unobserved protected attribute; this is beyond our scope.

When  $A$  is available for training but not for prediction/testing, the classifier we use at test-time must be defined on  $\mathcal{X}$  rather than on  $\mathcal{X} \times \mathcal{A}$ . In other words, a classifier is a measurable function  $f_X: \mathcal{X} \rightarrow [0, 1]$  with  $\widehat{Y}_{f_X} | X \sim \text{Bern}(f_X(X))$ . In this section, in addition to writing  $\eta_a(x) = \mathbb{P}(Y = 1 | A = a, X = x)$  for all  $x, a$ , we will further denote  $\eta^Y(x) = \mathbb{P}(Y = 1 | X = x)$  and  $\eta^A(x) = \mathbb{P}(A = 1 | X = x)$  for all  $x$ , the regression functions of  $Y$  and  $A$  on  $X$ , respectively.

We call a disparity measure  $\text{Dis}_X$  linear if (3.6) holds with  $(x, a)$  replaced by  $x$ , i.e., there is a weight function  $w_{X, \text{Dis}}: \mathcal{X} \rightarrow \mathbb{R}$  such that, for all  $f_X: \mathcal{X} \rightarrow [0, 1]$ ,

$$\text{Dis}_X(f_X) = \int_{\mathcal{X}} f_X(x) w_{X, \text{Dis}}(x) d\mathbb{P}_X(x). \quad (4.10)$$

We show that, when the protected attribute  $A$  is not used for prediction, the three common disparity measures from (2.1) are still linear.

**Proposition 4.10** (Common Disparity Measures are Linear when Not Using  $A$ ). *When the classifier depends only on  $X$  rather than on  $(X, A)$ , the disparity measures DD, DO, and PD from (2.1) are linear with weighting functions defined for all  $x$  by*

$$\begin{aligned} w_{\text{DD}}(x) &= \frac{\eta^A(x)}{p_1} - \frac{1 - \eta^A(x)}{p_0}; & w_{\text{DO}}(x) &= \frac{\eta_1(x)\eta^A(x)}{p_{1,1}} - \frac{\eta_0(x)(1 - \eta^A(x))}{p_{0,1}}; \\ w_{\text{PD}}(x) &= \frac{(1 - \eta_1(x))\eta^A(x)}{p_{1,0}} - \frac{(1 - \eta_0(x))(1 - \eta^A(x))}{p_{0,0}}. \end{aligned}$$

For a general disparity measure  $\text{Dis}$ , a  $\delta$ -fair Bayes-optimal classifier is defined as  $f_{X, \text{Dis}, \delta}^* \in \arg \min_{f_X: \mathcal{X} \rightarrow [0, 1]} \{R(f_X) : |\text{Dis}_X(f_X)| \leq \delta\}$ . We observe that the analysis of linear disparity measures also applies to the scenario, leading to the following result.

**Corollary 4.11** (Bayes-Optimal Classifiers for Linear Disparity Measures, Not Using  $A$ ). *When  $A$  is not used for prediction, for a given linear disparity measure  $\text{Dis}_X$  as defined in (4.10), suppose that both  $\eta^Y(X)$  and  $w_{X, \text{Dis}}(X)$  have probability density functions on  $\mathcal{X}$ . Let  $D_{X, \text{Dis}}: \mathbb{R} \rightarrow [-1, 1]$  be defined for all  $t$  as*

$$D_{X, \text{Dis}}(t) := \int_{\mathcal{X}} \left[ w_{X, \text{Dis}}(x) \cdot I \left( \eta^Y(x) > \frac{1}{2} + \frac{t}{2} w_{X, \text{Dis}}(x) \right) \right] d\mathbb{P}_X(x).$$

Let  $t_{X, \text{Dis}}: [0, \infty) \rightarrow \mathbb{R}$  be defined for all  $\delta \geq 0$  by

$$t_{X, \text{Dis}}(\delta) = \arg \min_t \{ |t| : |D_{X, \text{Dis}}(t)| \leq \delta \}. \quad (4.11)$$

Then, for any  $\delta \geq 0$ , there exists a  $\delta$ -fair Bayes-optimal classifier  $f_{X, \text{Dis}, \delta}^*$  taking the form, for all  $x \in \mathcal{X}$ ,

$$f_{X, \text{Dis}, \delta}^*(x) = I \left( \eta^Y(x) > \frac{1}{2} + \frac{t_{X, \text{Dis}}(\delta)}{2} w_{X, \text{Dis}}(x) \right).$$

## 5 Bayes-Optimal Fair Classifiers via Pre-, In-, and Post-processing

In this section, we propose a comprehensive set of algorithms for Bayes-optimal fair classification, based on pre-, in-, and post-processing. These algorithms aim to ensure fairness at different stages of the training



process. Each class of methods has strengths and limitations. Pre- and post-processing methods can be used without significant changes to standard classification methods, facilitating the use of widely-available software. However, these approaches might potentially clash with certain data protection regulations (Barocas and Selbst, 2016). On the other hand, in-processing methods are applicable whenever standard training workflows are, but may require modifying the fitting methods or optimization objectives.

We develop methods of each type aiming to recover the Bayes-optimal classifiers. Our methods include pre-processing via fair up-/down-sampling, in-processing via group-based cost-sensitive classification, and post-processing via plug-in estimation. Each method inherits the strengths mentioned above.

In this section, we consider bilinear disparity measures as per Definition 3.3 for simplicity. However, our methods are adaptable to the more general scenario of linear disparity measures, as shown in case studies in Section B.3 of the Appendix. Given a bilinear disparity measure characterized by  $s_{\text{Dis},a}$  and  $b_{\text{Dis},a}$  for  $a \in \mathcal{A}$ , we denote for  $t \in \mathbb{R}$  the threshold function

$$H_{\text{Dis},a}(t) = \frac{1 + t \cdot b_{\text{Dis},a}}{2 - t \cdot s_{\text{Dis},a}}. \quad (5.1)$$

Then, the disparity function  $D$  from (4.2) has values, for all  $t \in \mathbb{R}$ ,

$$D_{\text{Dis}}(t) = \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} [(s_{\text{Dis},a} \eta_a(x) + b_{\text{Dis},a}) \cdot I(\eta_a(x) > H_{\text{Dis},a}(t))] d\mathbb{P}_{X|A=a}(x). \quad (5.2)$$

In the following, we let the observed data points  $\mathcal{S}_n = \{(x_i, a_i, y_i)\}_{i=1}^n$  be an independent and identically distributed sample from  $\mathbb{P}$  over the domain  $\mathcal{X} \times \mathcal{A} \times \mathcal{Y}$ . Moreover, we separate the data according to the protected feature, letting for  $a \in \{0,1\}$ ,  $\mathcal{S}_{n,a} = \{(x_i, a_i, y_i) \in \mathcal{S}_n, a_i = a\}$ . For  $a \in \{0,1\}$ , the  $j$ -th element of  $\mathcal{S}_{n,a}$  is denoted as  $(x_{a,j}, a, y_{a,j})$ , for  $j \in [n_a]$ . We further separate the dataset  $\mathcal{S}_{n,a}$  into  $\mathcal{S}_{n,a,1}$  and  $\mathcal{S}_{n,a,0}$ , according to the label information. Moreover, we denote, for  $a, y \in \{0,1\}^2$ ,  $n_a = |\mathcal{S}_{n,a}|$  and  $n_{a,y} = |\mathcal{S}_{n,a,y}|$ .

## 5.1 Pre-processing: Fair Up- and Down-Sampling (FUDS)

Many pre-processing algorithms use heuristics to eliminate the impact of protected attributes on the common features  $X$  and the label  $Y$  (Lum and Johndrow, 2016; Xu et al., 2018). Nonetheless, pre-processing a dataset to satisfy a fairness criterion does not guarantee fair outcomes, see e.g., Eitan et al. (2022). To illustrate this, consider a transformed distribution  $(\tilde{X}, \tilde{A}, \tilde{Y})$ , where  $(\tilde{X}, \tilde{Y})$  is independent of  $A$ . A classifier  $f$  trained on a sample with the distribution  $(\tilde{X}, \tilde{Y})$  would be independent of  $A$ . Yet, during the testing phase, the classifier's output  $f(X)$  may still exhibit unfairness if  $X$  and  $A$  are dependent. This observation leads us to ask: *On what data distributions  $(\tilde{X}, \tilde{A}, \tilde{Y}) \sim \tilde{\mathbb{P}}$  can we learn classifiers without a fairness constraint and recover a fair Bayes-optimal classifier for the original distribution  $(X, A, Y) \sim \mathbb{P}$ ?*

We propose an answer by designing a joint distribution for  $(\tilde{A}, \tilde{Y})$ , while keeping the conditional distribution of  $\tilde{X}$  given  $\tilde{A}$  and  $\tilde{Y}$  unchanged from that of  $X$  given  $A$  and  $Y$ . We will consider distributions  $\tilde{\mathbb{P}}^t$  depending on a scalar  $t \in \mathbb{R}$ . For the distribution  $\tilde{\mathbb{P}}^t$  of  $(\tilde{X}, \tilde{A}, \tilde{Y})$ , we denote for all  $x, a, y$ ,  $\tilde{p}_{a,y}^t = \tilde{\mathbb{P}}^t(\tilde{A} = a, \tilde{Y} = y)$ , denote  $\tilde{\eta}_a^t(x) = \tilde{\mathbb{P}}^t(\tilde{Y} = 1 | \tilde{A} = a, \tilde{X} = x)$ , and denote by  $\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=a, \tilde{Y}=y}^t(\cdot)$  the marginal distribution of  $\tilde{X}$  given  $\tilde{A} = a$  and  $\tilde{Y} = y$ .

**Theorem 5.1** (Bayes-Optimal Fair Up- and Down-Sampling). *Suppose that for  $a \in \{0,1\}$ , both  $\eta_a(X)$  and  $w_{\text{Dis}}(X, a)$  have probability density functions on  $\mathcal{X}$ . For  $t \in \mathbb{R}$ , let  $\tilde{\mathbb{P}}^t$  be the distribution satisfying that for all  $(x, a, y) \in \mathcal{X} \times \{0,1\}^2$ ,  $\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=a, \tilde{Y}=y}^t(x) = \mathbb{P}_{X|A=a, Y=y}(x)$ , and, with  $H_{\text{Dis},a}$  from (5.1),*

$$\tilde{p}_{a,y}^t = \tilde{\mathbb{P}}^t(\tilde{A} = a, \tilde{Y} = y) = c_a[(1 - H_{\text{Dis},a}(t))y + H_{\text{Dis},a}(t)(1 - y)]p_{a,y}. \quad (5.3)$$

Here  $c_1$  and  $c_0$  are such that  $\tilde{p}_{11}^t + \tilde{p}_{10}^t + \tilde{p}_{01}^t + \tilde{p}_{00}^t = 1$ . Then, the unconstrained Bayes-optimal classifier for  $\tilde{\mathbb{P}}^t$  is a  $|D_{\text{Dis}}(t)|$ -fair Bayes-optimal classifier for  $\mathbb{P}$ . Thus, for a given  $\delta \geq 0$ , any Bayes-optimal classifier for  $\tilde{\mathbb{P}}^{t_{\text{Dis}}(\delta)}$  is a  $\delta$ -fair Bayes-optimal classifier for  $\mathbb{P}$ .

Theorem 5.1 designs a fairness-inducing distribution by multiplying the probabilities of each group with a group-wise factor, as per (5.3). As a result,  $(1 - H_{\text{Dis},a}(t))y + H_{\text{Dis},a}(t)(1 - y)$  can be viewed as a fairness-inducing factor for the group with  $A = a$  and  $Y = y$ . As can be seen from (5.2), a large threshold  $H_{\text{Dis},a}(t)$  for the group with  $A = a$  indicates that the fair Bayes-optimal classifier specifies a higher acceptance standard for group  $A = a$ . In other words, the unconstrained classifier has a higher acceptance rate on the subgroup with  $A = a$ . To avoid this disparity, we need to generate more data with  $Y = 0$  and less data with  $Y = 1$ , for the group with  $A = a$ ; reducing  $\eta_a(x) = \mathbb{P}(Y = 1|X = x, A = a)$ . For this reason, we call our approach Fair Up- and Down-Sampling (FUDS).

---

**Algorithm 1:** Fair Up-/Down-Sampling (FUDS)

---

**Input:** Step size  $\alpha > 0$ ; Disparity Level  $\delta \geq 0$ ; Error tolerance level  $\varepsilon > 0$ ; Dataset  $S = S_1 \cup S_0$  with  $S = \{x_i, a_i, y_i\}_{i=1}^n$ ,  $S_1 = \{x_{1,i}, y_{1,i}\}_{i=1}^{n_1}$  and  $S_0 = \{x_{0,i}, y_{0,i}\}_{i=1}^{n_0}$ ;  $n_{a,y} = \#\{a_i = a, y_i = y\}$ .

---

**Disparity Estimation sub-routine: Construct  $\hat{f}_t^{\text{FUDS}}$  and estimate  $\text{Dis}(\hat{f}_t^{\text{FUDS}})$  for a given threshold parameter  $t$ :**

1. For all  $a$ , estimate  $\hat{w}_{\text{Dis}}$  and  $\hat{H}_{\text{Dis},a}$  using the expressions in (3.7) and (5.1).
  2. For all  $a, y$ , let  $\hat{p}_{a,y}^t$  be as in (5.4).
  3. Apply up- and down-sampling to generate a new dataset  $S_t$  with proportions  $\hat{p}_{a,y}^t$  for all  $a, y$ .
  4. Fit classifier  $\hat{f}_t^{\text{FUDS}}$  on the dataset  $S_t$  using any method.
  5. Estimate the disparity level of  $\hat{f}_t^{\text{FUDS}}$  as in (5.5).
- 

**Estimate the  $\delta$ -fair Bayes-optimal classifier:**

Run Disparity Estimation sub-routine with  $t = 0$ .

**if**  $|\widehat{\text{Dis}}^{\text{FUDS}}(0)| \leq \delta$  **then**  
  |  $\hat{t}_{\text{Dis}}(\delta) = 0$ .

**else**

**if**  $\widehat{\text{Dis}}^{\text{FUDS}}(0) > \delta$  **then**

    |  $\delta' = \delta$ ;  $t_{\min} = 0$ ,  $t_{\max} = 1$ .

**else**

    |  $\delta = -\delta$ ;  $t_{\min} = -1$ ,  $t_{\max} = 0$ .

**end**

**while**  $t_{\max} - t_{\min} > \varepsilon$  **do**

    |  $t = (t_{\max} + t_{\min})/2$ ;

    Run Disparity Estimation sub-routine with current  $t$ .

**if**  $\widehat{\text{Dis}}^{\text{FUDS}}(t) > \delta$  **then**

      |  $t_{\max} = t$

**else**

      |  $t_{\min} = t$

**end**

**end**

$\hat{t}_{\text{Dis}}(\delta) = t$ .

**end**

**Output:**  $\hat{f}_{\text{Dis},\delta} = \hat{f}_{\hat{t}_{\text{Dis}}(\delta)}^{\text{FUDS}}$ .

---

While the sampling probabilities from (5.3) depend on unknown population parameters, we can estimate them from the training data. Then, we can generate approximately fair datasets using up- and down-sampling; see Algorithm 1. According to Theorem 5.1, for each value of  $t$ , there is a fair Bayes-optimal classifier for *some* level— $|\text{Dis}(t)|$ —of disparity. Therefore, by adjusting  $t$ , we can create a range of data distributions, each leading to a different point on the fair Pareto frontier as defined in (4.6). When we need a dataset with a predetermined disparity level  $\delta$ , we can estimate the threshold parameter  $t_{\text{Dis}}(\delta)$  using an iterative method for updating  $t$ , see Algorithm 1.

Specifically, for a given  $t$ , we estimate  $f_t^{\text{FUDES}}$  and its disparity level as follows:

(1) For all  $x, a, y, t$ , estimate  $p_{a,y}$  by  $n_{a,y}/n$  and estimate  $w_{\text{Dis}}(x, a)$  and  $H_{\text{Dis},a}(t)$  from (3.7) and (5.1) by plug-in estimation. Taking demographic disparity as an example, we let for all  $x, a$  and for all  $t \in \mathbb{R}$ ,

$$\widehat{w}_{\text{DD}}(x, a) = \frac{n}{n_{a,1} + n_{a,0}}, \quad \text{and} \quad \widehat{H}_{\text{DD},a}(t) = \frac{1}{2} + \frac{(2a-1)nt}{n_{a,1} + n_{a,0}}.$$

(2) Estimate the group-wise proportions, for  $(a, y) \in \{0, 1\}^2$ ,

$$\widehat{p}_{a,y}^t = \widehat{c}_a[(1 - \widehat{H}_{\text{Dis},a}(t))y + \widehat{H}_{\text{Dis},a}(t)(1 - y)] \cdot n_{a,y}/n. \quad (5.4)$$

Here  $\widehat{c}_1$  and  $\widehat{c}_2$  are such that  $\widehat{p}_{11}^t + \widehat{p}_{10}^t + \widehat{p}_{01}^t + \widehat{p}_{00}^t = 1$ .

(3) Apply up- and down- sampling to generate a new dataset  $S_t = \{\tilde{x}_i, \tilde{a}_i, \tilde{y}_i\}_{i=1}^n$  with  $\tilde{n}_{a,y}^t = \#\{\tilde{a}_i = a, \tilde{y}_i = y\} = \lfloor n \cdot \widehat{p}_{a,y}^t \rfloor$ . The precise sampling method is presented in the paragraphs below.

(4) Fit a classifier  $\widehat{f}_t^{\text{FUDES}}$  to the dataset  $S_t$  using any method, such as logistic regression or deep neural nets.

(5) Estimate the disparity level of  $\widehat{f}_t^{\text{FUDES}}$  for the given  $t$ :

$$\widehat{\text{Dis}}^{\text{FUDES}}(t) = \frac{1}{n_1} \sum_{j=1}^{n_1} \widehat{f}_t^{\text{FUDES}}(x_{1,j}, 1) \widehat{w}_{\text{Dis}}(x_{1,j}, 1) - \frac{1}{n_0} \sum_{j=1}^{n_0} \widehat{f}_t^{\text{FUDES}}(x_{0,j}, 0) \widehat{w}_{\text{Dis}}(x_{0,j}, 0). \quad (5.5)$$

Following this,  $t$  can be adjusted using the bisection method for solving  $D_{\text{Dis}}(t) = 0$ , based on the estimating equation  $\widehat{\text{Dis}}^{\text{FUDES}}(t) = 0$ . This is grounded in our observation that, as per Proposition 4.1, at a population level, the disparity function  $\text{Dis}(f_t^{\text{FUDES}})$  is monotone non-increasing in  $t$ .

Performing independent random resampling of the original dataset  $S_n$  in each iteration can lead to significant variability in the datapoints—and also in the induced classifiers. To address this issue, we propose a sampling strategy aimed at reducing variability.

Consider the generated dataset  $\widetilde{S}_n^{t'}$  for a given threshold  $t'$ , and write it as  $\widetilde{S}_n^{t'} = \cup_{a,y \in \{0,1\}^2} \widetilde{S}_{n,a,y}^{t'}$ , where for all  $a, y$ ,  $\widetilde{S}_{n,a,y}^{t'}$  represents the set of points with  $\tilde{a}_i = a$  and  $\tilde{y}_i = y$ . Let  $t$  be the next threshold, and let  $\tilde{n}_{a,y}^{t'} = \lfloor n \cdot \widehat{p}_{a,y}^{t'} \rfloor$  and  $\tilde{n}_{a,y}^t = \lfloor n \cdot \widehat{p}_{a,y}^t \rfloor$  for all  $a, y$ . Our strategy considers the following two cases, for all  $a, y$ :

- If  $\tilde{n}_{a,y}^t \geq \tilde{n}_{a,y}^{t'}$ , we randomly select  $\tilde{n}_{a,y}^t - \tilde{n}_{a,y}^{t'}$  data points from  $S_{n,a}$  to add to  $\widetilde{S}_{n,a,y}^{t'}$ , forming  $\widetilde{S}_{n,a,y}^t$ .
- If  $\tilde{n}_{a,y}^t < \tilde{n}_{a,y}^{t'}$ , we randomly select  $\tilde{n}_{a,y}^t$  data points from  $\widetilde{S}_{n,a,y}^{t'}$  to form  $\widetilde{S}_{n,a,y}^t$ .

Our simulation studies indicate that this strategy offers greater stability than independent random sampling in each iteration, leading to a more favorable fairness-accuracy tradeoff and reduced variance in the empirical performance.

Another challenge is posed by computational complexity. As discussed above, we fit a classifier at each iteration and then estimate its disparity level. For larger datasets and more complex models, this procedure can become time-consuming. A faster method is to adjust  $t$  while fitting each classifier, fine-tuning the optimal sampling ratio concurrently with model training. We only need to revise step (4): for instance, when training via an iterative optimization-based method, rather than finishing the entire training procedure, we only train for a small number of steps—or, epochs—in each iteration to obtain a classifier  $\widehat{f}_t^{\text{FUDES}}$ . The classifier obtained in the each iteration is used as the starting point in the next iteration.

## 5.2 In-Processing: Fair Cost-Sensitive Classification (FCSC)

In this section, we propose an in-processing fair classification method. Our objective is to identify a risk function whose unconstrained minimizer is the fair Bayes-optimal classifier. The key observation is that fair Bayes-optimal classifiers adjust thresholds for each protected group, which is also known to occur in cost-sensitive classification.

Recall the cost-sensitive risk  $R_c$  from (4.9). An unconstrained Bayes-optimal classifier for this cost-sensitive risk, denoted as  $f_c^*$ , is a classifier that minimizes  $R_c(f)$ , i.e.,  $f_c^* \in \arg \min_{f \in \mathcal{F}} R_c(f)$ . All such classifiers can be represented as  $f_c^*(x, a) = I(\eta_a(x) > c)$  for all  $x, a$ , (see e.g., Elkan, 2001). Drawing inspiration from this connection, we propose a *group-wise* cost-sensitive classification approach. We aim to shift the thresholds used for the protected groups in opposite directions to reduce disparity.

**Definition 5.2** (Fair Cost-sensitive Risk). *Recall  $H_{\text{Dis},a}$ ,  $a \in \mathcal{A}$  from (5.1) and let  $c_{a,y}(t) = (1 - 2y)H_{\text{Dis},a}(t) + y$  for all  $t, a, y$ . For  $t \in \mathbb{R}$ , define the following fair cost-sensitive risk of a classifier  $f$ :*

$$R_t^{\text{FCSC}}(f) = \sum_{a \in \{0,1\}} \sum_{y \in \{0,1\}} c_{a,y}(t) \mathbb{P}(\hat{Y}_f = 1 - y, Y = y, A = a).$$

*Fair cost-sensitive classification* (FCSC) minimizes the fair cost-sensitive risk from Definition 5.2:

$$f_t^{\text{FCSC}} = \arg \min_{f \in \mathcal{F}_\delta} R_t^{\text{FCSC}}(f). \quad (5.6)$$

Our key result here is that fair cost-sensitive classification is Bayes-optimal.

**Theorem 5.3** (Fair Cost-sensitive Classification is Bayes-Optimal). *Letting  $t_0^* = t_{\text{Dis}}(0) = \arg \min_t \{|t| : |D_{\text{Dis}}(t)| = 0\}$ , we have, for  $t \in [\min(t_0^*, 0), \max(t_0^*, 0)]$ , with  $D_{\text{Dis}}(t)$  defined in (4.2), that  $f_t^{\text{FCSC}}$  from (5.6) is a  $|D_{\text{Dis}}(t)|$ -fair Bayes-optimal classifier for  $\mathbb{P}$ . In particular,  $f_{t_{\text{Dis}}(\delta)}^{\text{FCSC}}$  is a  $\delta$ -fair Bayes-optimal classifier for  $\mathbb{P}$ .*

Theorem 5.3 shows that carefully designed group-wise cost-sensitive classification can lead to a fair Bayes-optimal classifier. Similar to the fair sampling method from Section 5.1, one can generate a fairness-accuracy tradeoff curve by using FCSC and varying  $t$ . To achieve a specific disparity level  $\delta$ , an updating method similar to that from Algorithm 1 can be used to estimate  $t_{\text{Dis}}(\delta)$ ; see Algorithm 2.

Similar to the FUDS algorithm, the computational efficiency of FCSC can be improved by adjusting the threshold parameter  $t$  and the associated loss function after a few training epochs (rather than after the whole training process in step 3).

## 5.3 Post-Processing: Fair Plug-in Thresholding Rule (FPIR)

Finally, we consider a post-processing algorithm that modifies the model output to control the disparity. Given the near-explicit form of Bayes-optimal classifiers from Theorem 4.2, after fitting any classifier  $\hat{f}$ , we can estimate the group-wise thresholds. Such an approach has been proposed in a more limited case for demographic parity and equality of opportunity in Menon and Williamson (2018). In contrast, our framework is applicable to *any* bilinear disparity measure.

First, we can use any method to estimate the feature-conditional probability of  $Y = 1$  for each protected group. Second, we estimate the threshold for each protected group. One can either vary the value of  $t$  to generate fairness-accuracy tradeoff curves or solve the estimated one-dimensional fairness equation for a specific disparity level.

Specifically, let  $\hat{\eta}_a$  be an estimator of  $\eta_a$ , and consider the following plug-in rule for all  $x, a$ :

$$\hat{f}_t^{\text{FPIR}}(x, a) = I(\hat{\eta}_a(x) > \hat{H}_{\text{Dis},a}),$$

where  $\hat{H}_{\text{Dis},a}$  is estimated using the expression in (5.1). Now, our goal is to construct an estimate  $\hat{t}_\delta$  such that  $\hat{f}_{\hat{t}_\delta}$  approximately satisfies the fairness constraint. We define  $\widehat{\text{Dis}}(t)$ , an estimator of the disparity function

---

**Algorithm 2:** Fair Cost-Sensitive Classification (FCSC)
 

---

**Input:** Step size  $\alpha > 0$ ; Disparity Level  $\delta \geq 0$ , Error tolerance level  $\varepsilon > 0$ ; Dataset  $S = S_1 \cup S_0$  with  $S = \{x_i, a_i, y_i\}_{i=1}^n$ ,  $S_1 = \{x_{1,i}, y_{1,i}\}_{i=1}^{n_1}$  and  $S_0 = \{x_{0,i}, y_{0,i}\}_{i=1}^{n_0}$ ;  $n_{a,y} = \#\{a_i = a, y_i = y\}$ ; Step size  $\alpha \geq 0$ .

**Disparity Estimation sub-routine: Construct  $\hat{f}_t^{\text{FCSC}}$  and estimate  $\widehat{\text{Dis}}(\hat{f}_t^{\text{FCSC}})$  for a given threshold parameter  $t$ :**

1. For all  $a$ , estimate  $\hat{w}_{\text{Dis}}$  and  $\hat{H}_{\text{Dis},a}$  by plug-in estimation using the expressions in (3.7) and (5.1).
  2. Denote  $\hat{c}_{a,y}(t) = (1 - 2y)\hat{H}_{\text{Dis},a}(t) + y$ , for all  $t, a, y$ .
  3. Use any cost-sensitive classification method to fit  $\hat{f}_t^{\text{FCSC}}(\cdot, 1)$  on  $S_1$  and  $\hat{f}_t^{\text{FCSC}}(\cdot, 0)$  on  $S_0$ .
  4. Estimate the disparity level of  $\hat{f}_t^{\text{FCSC}}$  as in (5.5) with  $\hat{f}_t^{\text{FCSC}}$  instead of  $\hat{f}_t^{\text{FUDS}}$ .
- 

**Estimate the fair Pareto frontier:**

Run Disparity Estimation sub-routine with  $t = 0$ .

**if**  $\widehat{\text{Dis}}^{\text{FCSC}}(0) > 0$  **then**

  |  $\alpha' = \alpha$

**else**

  |  $\alpha' = -\alpha$

**end**

**while**  $\widehat{\text{Dis}}^{\text{FCSC}}(t) \cdot \widehat{\text{Dis}}^{\text{FCSC}}(0) > 0$  **do**

  |  $t = t + \alpha'$ .

  | Run Disparity Estimation sub-routine with current  $t$ .

**end**

**Output:**  $\{\hat{f}_t^{\text{FCSC}}\}_{t=0,1,2,\dots}$

**Estimate the  $\delta$ -fair Bayes-optimal classifier:**

Run Disparity Estimation sub-routine with  $t = 0$ . **if**

  |  $\widehat{\text{Dis}}^{\text{FCSC}}(0) \leq \delta$  **then**

    |  $\hat{t}_{\text{Dis}}(\delta) = 0$ .

**else**

**if**  $\widehat{\text{Dis}}^{\text{FCSC}}(0) > \delta$  **then**

    |  $\delta' = \delta$ ;  $t_{\min} = 0$ ,  $t_{\max} = 1$ .

**else**

    |  $\delta = -\delta$ ;  $t_{\min} = -1$ ,  $t_{\max} = 0$ .

**end**

**while**  $t_{\max} - t_{\min} > \varepsilon$  **do**

    |  $t = (t_{\max} + t_{\min})/2$ ;

    | Run Disparity Estimation sub-routine with current  $t$ .

**if**  $\widehat{\text{Dis}}^{\text{FCSC}}(t) > \delta$  **then**

      |  $t_{\max} = t$

**else**

      |  $t_{\min} = t$

**end**

**end**

    |  $\hat{t}_{\text{Dis}}(\delta) = t$ .

**end**

**Output:**  $\hat{f}_{\text{Dis},\delta} = \hat{f}_{\hat{t}_{\text{Dis}}(\delta)}^{\text{FUDS}}$ .

---

$D_{\text{Dis}}(t)$ , as in (5.5) with  $\hat{f}_t^{\text{FPIR}}$  instead of  $\hat{f}_t^{\text{FUDS}}$ , where  $\hat{w}_{\text{Dis}}$  is estimated using the plug-in estimators  $\hat{p}_{a,y} = n_{a,y}/n$  in (3.7). We then follow the definition of  $t_{\text{Dis}}(\delta)$  in (4.3) to estimate it by

$$\hat{t}_{\text{Dis}}(\delta) = \arg \min_t \{ |t| : |\widehat{\text{Dis}}(t)| \leq \delta \}.$$

Since  $\widehat{\text{Dis}}(t)$  is monotone non-increasing as a function of  $t$ ,  $\hat{t}_{\text{Dis}}(\delta)$  can be estimated via the bisection method, as summarized in Algorithm 3. Our final FPIR estimator of the fair Bayes-optimal classifier is  $\hat{f}_{\hat{t}_{\text{Dis}}(\delta)}^{\text{FPIR}}$ .

To summarize, we have proposed fair classifiers via pre-, in-, and post-processing. Our comprehensive methodology offers a broad framework for reducing algorithmic disparities. Compared to existing methods, our algorithms have two main advantages. First, all three methods are Bayes-optimal at the population level. This proves theoretically that they aim to address bias with a minimal sacrifice of accuracy. Second, in all three methods, the level of maximal disparity can be precisely specified. In contrast, most current methods either exclusively target perfect fairness ( $\delta = 0$ ) or use a tuning parameter that indirectly controls the disparity level, such as a Lagrange multiplier in constrained optimization.

---

**Algorithm 3:** Fair Plug-in Rule (FPIR)

---

**Input:** Step size  $\alpha > 0$ ; Disparity level  $\delta \geq 0$ ; Error tolerance level  $\varepsilon > 0$ ; Dataset  $S = S_1 \cup S_0$  with  $S = \{x_i, a_i, y_i\}_{i=1}^n$ ,  $S_1 = \{x_{1,i}, y_{1,i}\}_{i=1}^{n_1}$  and  $S_0 = \{x_{0,i}, y_{0,i}\}_{i=1}^{n_0}$ ;  $n_{a,y} = \#\{a_i = a, y_i = y\}$ .

**Step 1:** Construct estimate  $\hat{\eta}_a$  of  $\eta_a$  using any approach, for all  $a$ .

**Step 2:** Estimate the the fair Pareto frontier or the  $\delta$ -fair Bayes-optimal classifier:

**Disparity Estimation sub-routine:** Construct  $\hat{f}_t^{\text{FPIR}}$  and estimate  $\widehat{\text{Dis}}(\hat{f}_t^{\text{FPIR}})$  with threshold parameter  $t$ :

1. For all  $a$ , estimate  $\hat{w}_{\text{Dis}}$  and  $\hat{H}_{\text{Dis},a}$  by plug-in estimation using the expressions in (3.7) and (5.1).
  2. Let  $\hat{f}_t^{\text{FPIR}}$  be defined by  $\hat{f}_t^{\text{FPIR}}(x, a) = I(\hat{\eta}_a(x) > \hat{H}_{\text{Dis},a}(t))$  for all  $x, a$ .
  3. Evaluate the disparity level of  $\hat{f}_t^{\text{FPIR}}$  as in (5.5) with  $\hat{f}_t^{\text{FPIR}}$  instead of  $\hat{f}_t^{\text{FUDES}}$ .
- 

**Estimate the fair Pareto frontier:**

Run Disparity Estimation sub-routine with  $t = 0$ .

**if**  $\widehat{\text{Dis}}^{\text{FPIR}}(0) > 0$  **then**

  |  $\alpha' = \alpha$

**else**

  |  $\alpha' = -\alpha$

**end**

**while**  $\widehat{\text{Dis}}^{\text{FPIR}}(t) \cdot \widehat{\text{Dis}}^{\text{FPIR}}(0) > 0$  **do**

  |  $t = t + \alpha'$ .

  | Run Disparity Estimation sub-routine with current  $t$ .

**end**

**Output:**  $\{\hat{f}_t^{\text{FPIR}}\}_{t=0,1,2,\dots}$ .

**Estimate the  $\delta$ -fair Bayes-optimal classifier:**

Run Disparity Estimation sub-routine with  $t = 0$ .

**if**  $|\widehat{\text{Dis}}^{\text{FPIR}}(0)| \leq \delta$  **then**

  |  $\hat{t}_{\text{Dis}}(\delta) = 0$ .

**else**

**if**  $\widehat{\text{Dis}}^{\text{FPIR}}(0) > \delta$  **then**

    |  $\delta' = \delta$ ;  $t_{\min} = 0$ ,  $t_{\max} = 1$ .

**else**

    |  $\delta = -\delta$ ;  $t_{\min} = -1$ ,  $t_{\max} = 0$ .

**end**

**while**  $t_{\max} - t_{\min} > \varepsilon$  **do**

    |  $t = (t_{\max} + t_{\min})/2$ ;

    | Run Disparity Estimation sub-routine with current  $t$ .

**if**  $\widehat{\text{Dis}}^{\text{FPIR}}(t) > \delta$  **then**

      |  $t_{\max} = t$

**else**

      |  $t_{\min} = t$

**end**

**end**

  |  $\hat{t}_{\text{Dis}}(\delta) = t$ .

**end**

**Output:**  $\hat{f}_{\text{Dis},\delta} = \hat{f}_{\hat{t}_{\text{Dis}}(\delta)}^{\text{FPIR}}$ .

---

## 6 Empirical Experiments

### 6.1 Synthetic Datasets

In this section, we conduct simulation studies to illustrate the numerical performance of our proposed methods. We consider a label-conditional normal distribution, for which the fair Bayes-optimal classifier has a simple closed form.

**Data-generating process.** For a positive integer dimension  $p$ , let  $X = (X_1, X_2, \dots, X_p)^\top \in \mathbb{R}^p$  be the usual features,  $A \in \{0, 1\}$  be the protected attribute and  $Y \in \{0, 1\}$  be the label. Recalling the notations from the end of Section 2, we generate  $A$  and  $Y$  according to the probabilities  $p_{1,1} = 0.49$ ,  $p_{1,0} = 0.3$ ,  $p_{0,1} = 0.12$  and  $p_{0,0} = 0.18$ . Conditional on  $A = a$  and  $Y = y$ , we generate  $X$  from a multivariate Gaussian distribution  $\mathcal{N}(\mu_{a,y}, \sigma^2 I_p)$ , where  $I_p$  is the  $p$ -dimensional identity covariance matrix, and  $\sigma^2$  controls the variability of the feature entries. The entries of  $\mu_{a,y}$  are sampled from  $\mu_{ay,j} \sim \text{Unif}(0, 1)$ ,  $j = 1, \dots, p$ , where  $\text{Unif}(0, 1)$  is the uniform distribution over  $[0, 1]$ . In this model,  $\eta_a$  has a closed form, and we can use it to find the fair Bayes-optimal classifier, see Appendix C. Intriguingly, under this model, group-wise thresholding



rules are linear in  $x$ .

**Experimental setting.** To evaluate our FUDS, FCSC and FPIR algorithms, we randomly generate 10,000 training data points and 5,000 test data points. Since the group-wise thresholding rules are linear in  $x$ , we use logistic regression in our methods. All experiments are conducted in python and we use the scikit-learn package to train the logistic regression model. This package allows us to specify label weights for cost-sensitive classification, and thus works seamlessly for FCSC. In our experiments, we consider four fairness metrics: demographic parity, equality of opportunity, predictive equality and overall accuracy equality.

We first evaluate the three algorithms with various pre-determined levels of disparity. We present the simulation results in Table 4. We observe that FPIR controls the disparity level at the pre-determined value, as desired. We further present the fairness-accuracy tradeoff curve of fair Bayes-optimal classifiers (the fair Pareto frontier from Section 4.2) and our three methods in Figure 1. Our classifiers closely track the fair Pareto frontier.

## 6.2 Empirical Data Analysis

To further illustrate our proposed methods, we compare our method with strong baseline methods on standard datasets.

**Data Description.** We consider the benchmark datasets AdultCensus, COMPAS, and LawSchool and adopt the same data pre-processing pipeline as in Cho et al. (2020).

- (1) AdultCensus: In the AdultCensus dataset, the target variable  $Y$  is whether the income of an individual is more than \$50,000. Age, marriage status, education level, and other related variables are included in  $X$ , and the protected attribute  $A$  refers to gender.
- (2) COMPAS: In the COMPAS dataset,  $Y$  indicates whether or not a criminal will re-offend. Here  $X$  includes prior criminal records, age, and an indicator of misdemeanor. The protected attribute  $A$  is the race of an individual, “white-vs-non-white”.

Table 2: Numerical simulation results showing classification accuracies and levels of disparity of the true fair Bayes-optimal classifier and the three proposed methods (FUDS, FCSC, FPIR) using logistic regression. The reported results are averages over 5,000 test data points, with standard deviations shown in parentheses.

| DEMOGRAPHIC PARITY      |       |               |               |               |               |               |               |
|-------------------------|-------|---------------|---------------|---------------|---------------|---------------|---------------|
| THEORETICAL             |       | FUDS          |               | FCSC          |               | FPIR          |               |
| $\delta$                | ACC   | DD            | ACC           | DD            | ACC           | DD            | ACC           |
| 0.00                    | 0.727 | 0.013 (0.010) | 0.722 (0.007) | 0.013 (0.010) | 0.722 (0.007) | 0.013 (0.010) | 0.722 (0.007) |
| 0.10                    | 0.736 | 0.100 (0.017) | 0.735 (0.007) | 0.100 (0.017) | 0.736 (0.007) | 0.100 (0.017) | 0.736 (0.007) |
| 0.20                    | 0.746 | 0.200 (0.018) | 0.746 (0.007) | 0.200 (0.018) | 0.746 (0.006) | 0.200 (0.018) | 0.746 (0.006) |
| 0.30                    | 0.753 | 0.300 (0.017) | 0.753 (0.006) | 0.300 (0.017) | 0.753 (0.006) | 0.300 (0.017) | 0.753 (0.006) |
| EQUALITY OF OPPORTUNITY |       |               |               |               |               |               |               |
| THEORETICAL             |       | FUDS          |               | FCSC          |               | FPIR          |               |
| $\delta$                | ACC   | DD            | ACC           | DD            | ACC           | DD            | ACC           |
| 0.00                    | 0.731 | 0.017 (0.013) | 0.730 (0.007) | 0.017 (0.013) | 0.731 (0.007) | 0.017 (0.013) | 0.731 (0.007) |
| 0.10                    | 0.745 | 0.101 (0.022) | 0.744 (0.007) | 0.101 (0.022) | 0.744 (0.007) | 0.101 (0.023) | 0.744 (0.007) |
| 0.20                    | 0.753 | 0.201 (0.024) | 0.752 (0.006) | 0.202 (0.024) | 0.753 (0.006) | 0.202 (0.024) | 0.753 (0.006) |
| 0.30                    | 0.757 | 0.302 (0.025) | 0.756 (0.006) | 0.302 (0.025) | 0.757 (0.006) | 0.302 (0.025) | 0.757 (0.006) |
| PREDICTIVE EQUALITY     |       |               |               |               |               |               |               |
| THEORETICAL             |       | FUDS          |               | FCSC          |               | FPIR          |               |
| $\delta$                | ACC   | DD            | ACC           | DD            | ACC           | DD            | ACC           |
| 0.00                    | 0.743 | 0.020 (0.015) | 0.743 (0.006) | 0.020 (0.015) | 0.743 (0.006) | 0.020 (0.015) | 0.743 (0.007) |
| 0.10                    | 0.751 | 0.099 (0.025) | 0.750 (0.006) | 0.099 (0.025) | 0.750 (0.006) | 0.099 (0.025) | 0.750 (0.006) |
| 0.20                    | 0.756 | 0.199 (0.024) | 0.755 (0.006) | 0.199 (0.025) | 0.755 (0.006) | 0.199 (0.025) | 0.755 (0.006) |
| 0.30                    | 0.758 | 0.297 (0.023) | 0.757 (0.006) | 0.296 (0.024) | 0.757 (0.006) | 0.297 (0.024) | 0.757 (0.006) |




Figure 1: Numerical simulation results showing the fairness-accuracy tradeoff the fair Bayes-optimal classifier (fair Pareto frontier) and the three proposed methods (FUDS, FCSC, FPIR) using logistic regression.

(3) LawSchool: The target variable  $Y$  in the LawSchool dataset is whether a student gets admitted to law school. Here  $X$  includes the LSAT scores, undergraduate GPA and more. The protected attribute  $A$  we consider is the race, “white-vs-non-white”.

**Algorithms Considered.** We select six methods from the literature, comprising two pre-processing, in-processing, and post-processing methods each.

(1) Disparate Impact Remover (DIR, Feldman et al. (2015)):

DIR is a pre-processing method that aims to modify the value of  $Y$  to  $\tilde{Y}$ , ensuring that the probability of  $\tilde{Y} = 1$  is equal across different protected groups, specifically,  $\mathbb{P}(\tilde{Y} = 1|A = 1) = \mathbb{P}(\tilde{Y} = 1|A = 0)$ . The key idea of DIR is to align the cumulative distribution functions of  $\eta_1(X)$  and  $\eta_0(X)$  while maximizing similarity between the transformed and original datasets.

(2) FAWOS (Salazar et al., 2021):

FAWOS is a pre-processing method that uses data augmentation to achieve fairness. Unlike our FUDS method that adjusts the sizes of all groups, FAWOS applies SMOTE (Chawla et al., 2002)—a popular synthetic data augmentation method for unbalanced classification problems—to increase the number of data points in the positive unprivileged group ( $A = 0, Y = 1$ ). The number of data points generated is  $N = \alpha_F \times (n_{11}n_{00}/n_{10} - n_{01})$ , with  $\alpha_F$  tuned to control the accuracy-fairness tradeoff.

(3) KDE-based constrained optimization (KDE, Cho et al. (2020)):

KDE-based optimization is an in-processing technique that uses kernel density estimation (KDE) to approximate fairness constraints. The proposed estimator of disparity is a differentiable function with respect to the model parameters. KDE-based optimization uses this estimated disparity as a regularizer for empirical risk minimization.

(4) Adversarial training (ADV, Zhang et al. (2018)):

Adversarial training is an in-processing method that involves simultaneously training two models: (1) a classifier to predict the label and (2) an adversary that attempts to predict protected attributes from the classifier’s outputs. This dual-training process encourages the classifier to learn predictable representations that are devoid of biases related to protected attributes, thereby promoting fairness in its predictions.

(5) Post-processing through flipping (PPF, Chen et al. (2023)):

Chen et al. (2023) showed that a Bayes-optimal fair classifier can be obtained by flipping the output of the unconstrained classifier and proved that the flipping probability satisfies an estimation equation. Based on these observations, they designed a post-processing method that first estimates the flipping probability of each output and then estimates the fair Bayes-optimal classifier.

(6) Post-processing through optimal transport (PPOT, Xian et al. (2023)):

Xian et al. (2023) proved that the fair classification problem with bounded demographic parity is equivalent to a Wasserstein-barycenter problem, and the fair Bayes-optimal classifier is given by the composition of the

Table 3: Training details for the empirical datasets.

| DATASET     | BATCH | TRAINING | STARTING      | LEARNING RATE |
|-------------|-------|----------|---------------|---------------|
|             | SIZE  | EPOCHS   | LEARNING RATE | DECAY FACTOR  |
| ADULTCENSUS | 512   | 200      | 1E-1          | 0.98          |
| COMPAS      | 2048  | 500      | 5E-4          | NONE          |
| LAWSCHOOL   | 2048  | 200      | 2E-4          | NONE          |

Bayes-optimal score function  $\eta_a$  and the optimal transport map from the Wasserstein-barycenter problem. They proposed a post-processing algorithm to estimate the fair Bayes-optimal classifier from the estimated score functions  $\hat{\eta}_a$ .

**Experimental Setting:** We follow the training settings from Cho et al. (2020). For all datasets and methods, a three-layer fully connected neural network with 32 hidden neurons is trained with the Adam optimizer using  $(\beta_1, \beta_2) = (0.9, 0.999)$ , the default hyperparameters.

For adversarial training (Zhang et al., 2018), we further use a linear classifier as the discriminator. The training batch size, training epochs, starting learning rate, and learning rate decay factor for the three datasets are summarized in Table 3. Let  $n_{\text{epoch}}$  represent the total number of training epochs. For the FUDS and FCSC methods, we initially pre-train the model by training the neural network for  $n_{\text{epoch}}/4$  epochs. Subsequently, we apply the bisection method as described in Algorithms 1 and 2 to update  $t$  after every  $n_{\text{epoch}}/20$  epochs. Through this approach, we update  $t$  a total of 15 times, resulting in a tolerance level of  $2^{-15} \sim 3 \times 10^{-5}$ . For adversarial training, we first pre-train the classifier over  $n_{\text{epoch}}/4$  epochs and then update the weights of both the classifier and discriminator in the later  $3n_{\text{epoch}}/4$  epochs.

For all datasets and methods, we repeat the experiments 50 times with different random seeds. For each dataset and random seed, we randomly split the data into training and test sets (with a 70%-30% split). As a result, the randomness of the experiments arises from the stochasticity of the train-test set split, the neural network initialization, and the minibatch selection during the optimization.

**Simulation Results for Our Methods:** We first evaluate the FUDS, FCSC, and FPIR algorithms with various pre-determined levels of disparity. We present the simulation results in Table 4. We observe that the proposed three methods control the disparity level at the pre-determined values, as desired. We then compare the fairness-accuracy tradeoff of the proposed methods to the baseline.

**Experimental Setting for Controlling Disparity:** In our proposed methods—FUDS, FCSC, and

Table 4: Empirical data results showing the DD of FUDS, FCSC and FPIR with predetermined unfairness levels.

| ADULTCENSUS |          |               |               |               |               |               |
|-------------|----------|---------------|---------------|---------------|---------------|---------------|
| METHODS     | $\delta$ | 0.00          | 0.04          | 0.08          | 0.12          | 0.16          |
| FUDS        | DD       | 0.008 (0.005) | 0.038 (0.012) | 0.075 (0.012) | 0.121 (0.011) | 0.154 (0.010) |
| FCSC        |          | 0.017 (0.020) | 0.036 (0.012) | 0.079 (0.011) | 0.121 (0.008) | 0.159 (0.007) |
| FPIR        |          | 0.006 (0.004) | 0.040 (0.007) | 0.082 (0.011) | 0.122 (0.014) | 0.158 (0.007) |
| COMPAS      |          |               |               |               |               |               |
| METHODS     | $\delta$ | 0.00          | 0.06          | 0.12          | 0.18          | 0.24          |
| FUDS        | DD       | 0.030 (0.024) | 0.056 (0.029) | 0.111 (0.037) | 0.169 (0.036) | 0.227 (0.035) |
| FCSC        |          | 0.034 (0.027) | 0.087 (0.047) | 0.130 (0.041) | 0.180 (0.045) | 0.229 (0.036) |
| FPIR        |          | 0.030 (0.022) | 0.062 (0.034) | 0.116 (0.037) | 0.177 (0.040) | 0.233 (0.033) |
| LAWSCHOOL   |          |               |               |               |               |               |
| METHODS     | $\delta$ | 0.000         | 0.0016        | 0.032         | 0.48          | 0.64          |
| FUDS        | DD       | 0.007 (0.005) | 0.018 (0.007) | 0.034 (0.007) | 0.048 (0.006) | 0.063 (0.006) |
| FCSC        |          | 0.007 (0.005) | 0.021 (0.008) | 0.036 (0.007) | 0.050 (0.008) | 0.065 (0.007) |
| FPIR        |          | 0.004 (0.003) | 0.016 (0.005) | 0.032 (0.005) | 0.047 (0.005) | 0.063 (0.005) |

FPIR—along with the two other post-processing methods, the level of disparity is directly controlled. We set the range from zero to the empirical DD of the unconstrained classifier. For the DIR method, as suggested by Feldman et al. (2015), a “partial remover” tool is employed. This tool features a parameter  $\lambda_D$ , which varies from zero (indicating the original dataset) to 1 (indicating perfect distribution alignment between  $\eta_1(X)$  and  $\eta_0(X)$ ). We adjust  $\lambda_D$  from 0.05 to 0.95 to explore the fairness-accuracy tradeoff.

In FAWOS, we vary  $\alpha_F$  from zero to two. Higher values of  $\alpha_F$  lead to an excess of data points in the positive unprivileged group, potentially causing reverse discrimination. KDE-based constrained optimization balances fairness and accuracy by adjusting a tuning parameter that control the ratio between empirical loss and fairness regularization. This tuning parameter  $\lambda_K$  is varied from 0.05 to 0.95. Finally, in adversarial training, the tradeoff is controlled by altering the parameter  $\alpha_A$  that controls the gradient of the discriminator. We vary this parameter from zero to three, as we empirically find that, similar to FAWOS, a larger  $\alpha_A$  would result in reverse discrimination.

**Simulation Results for Controlling Disparity:** Figure 2 presents the fairness-accuracy tradeoff with respect to Demographic Disparity (DD) evaluated across three datasets. In the plot, each point corresponds to a specific tuning parameter. Among the three pre-processing methods, our FUDS algorithm exhibits the most favorable fairness-accuracy tradeoff and allows for direct control of disparity.

In contrast, while the DIR method shows satisfactory accuracy, it falls short in achieving perfect fairness. This is because, although DIR ensures equal probabilities of positive outcomes across different groups in the transformed dataset—i.e.,  $\mathbb{P}(\tilde{Y} = 1|A = 1) = \mathbb{P}(\tilde{Y} = 1|A = 0)$ —this does not automatically ensure that a fair model is learned. The third method, FAWOS, effectively manages disparity, but its heuristic-based sampling strategy strays from the ideal proportion we identified. As a result, compared to FUDS, FAWOS compromises more on accuracy at all levels of disparity.

For the three in-processing methods, FCSC outperforms the KDE and ADV methods on the COMPAS dataset and demonstrates comparable results on the other two datasets. In comparison, the proposed FCSC method directly controls disparity levels, making it more practical and user-friendly in applications. The KDE method experiences a sudden drop in accuracy when the disparity level approaches zero. This issue may be caused by its use of a Huber surrogate loss to address the non-differentiability of the absolute value function at  $\delta = 0$ . Compared with FCSC and KDE-based optimization, adversarial training exhibits inferior performance and fails to achieve near-perfect fairness. This shortcoming could be attributed to the challenges associated with minimax optimization, which often leads to unstable training processes.

Finally, we observe that the three post-processing methods present comparable performance in balancing fairness and accuracy. This similarity arises because they all aim to estimate the fair Bayes-optimal classifier. However, FPIR can be viewed as a more direct approach.

In summary, the three methods we proposed demonstrate better or comparable performance compared to existing methods. Additionally, our methods provide the ability to flexibly adjust the level of disparity. This flexibility is particularly beneficial in scenarios where a particular level of disparity is required.

## 7 Summary and Discussion

In this paper, we study the statistical and methodological underpinnings of fair classification problems. We establish a unified framework for deriving fair Bayes-optimal classifiers. We introduce the notion of linear and bilinear disparity measures, and find Bayes-optimal classifiers under these constraints. Our framework can be applied in a wide range of scenarios, including various disparity measures, cost-sensitive loss functions, and multiple fairness constraints while allowing for direct control of the disparity level. Additionally, we study the Pareto frontier for fair classification and prove that the tradeoff function is convex for linear group fairness measures.

Based on our theoretical results, we further design pre-, in-, and post-processing methods that handle algorithmic bias. The proposed methods aim to recover the optimal fairness-accuracy tradeoff. Moreover, they allow for direct control of the disparity level.

Our work suggests several promising directions for further theoretical and algorithmic development. From the theoretical perspective, the fair Bayes-optimal classifier only characterizes the best solution at the population level, and the finite-sample properties need to be characterized. From the algorithmic perspective, it is of interest to improve our algorithms. For example, for our pre-processing method, we only considered

fair sampling. However, with the desired proportions in hand, we can generate fair synthetic data using more advanced conditional generative models, such as conditional GANs and conditional diffusion models.

Furthermore, our analysis supposes the availability of protected attributes during training. However, the collection and use of sensitive information is often restricted. For instance, under the General Data Protection Regulations in the European Union, acquiring sensitive personal information requires consent from individuals. This motivates studying methods for protected attributes only partially available during training. Recently, Chai et al. (2022) proposed addressing this challenge through knowledge distillation, which generates soft labels for protected attributes. Exploring the integration of this technique into our framework presents an intriguing avenue for future research.




Figure 2: Fairness-accuracy tradeoff on the AdultCensus (Panel (A)), COMPAS (Panel (B)) and LawSchool (Panel (C)) datasets.

## Acknowledgements

This work was supported in part by ARO W911NF-23-1-0296, NSF 2031895, NSF 2046874, ONR N00014-21-1-2843, and the Sloan Foundation.

## References

- I. Alabdulmohsin. Fair classification via unconstrained optimization. *arXiv preprint arXiv:2005.14621*, 2020.
- J. Angwin, J. Larson, S. Mattu, and L. Kirchner. Machine Bias. There’s software used across the country to predict future criminals. And it’s biased against blacks. ProPublica, May 2016. URL <https://www.propublica.org/article/machine-bias-risk-assessments-in-criminal-sentencing>.
- J.-Y. Audibert and A. B. Tsybakov. Fast learning rates for plug-in classifiers. *The Annals of Statistics*, 35(2):608–633, 2007.
- S. Barocas and A. D. Selbst. Big data’s disparate impact. *California Law Review*, 104(3):671–732, 2016.
- R. K. E. Bellamy, K. Dey, M. Hind, S. C. Hoffman, S. Houde, K. Kannan, P. Lohia, J. Martino, S. Mehta, A. Mojsilovic, S. Nagar, K. N. Ramamurthy, J. T. Richards, D. Saha, P. Sattigeri, M. Singh, K. R. Varshney, and Y. Zhang. AI Fairness 360: An extensible toolkit for detecting and mitigating algorithmic bias. *IBM Journal of Research and Development*, 63(4/5):4:1–4:15, 2019.
- K. Bhanot, M. Qi, J. S. Erickson, I. Guyon, and K. P. Bennett. The problem of fairness in synthetic healthcare data. *Entropy*, 23(9):1165, 2021.
- T. Calders, F. Kamiran, and M. Pechenizkiy. Building classifiers with independency constraints. In *2009 IEEE International Conference on Data Mining Workshops*, pages 13–18. IEEE, 2009.
- F. Calmon, D. Wei, B. Vinzamuri, K. Natesan Ramamurthy, and K. R. Varshney. Optimized pre-processing for discrimination prevention. In *Advances in Neural Information Processing Systems*. Curran Associates, Inc., 2017.
- S. Caton and C. Haas. Fairness in machine learning: A survey. *ACM Computing Surveys*, 2023.
- L. E. Celis and V. Keswani. Improved adversarial learning for fair classification. *arXiv preprint arXiv:1901.10443*, 2019.
- L. E. Celis, L. Huang, V. Keswani, and N. K. Vishnoi. Classification with fairness constraints: A meta-algorithm with provable guarantees. In *Proceedings of the Conference on Fairness, Accountability, and Transparency*, pages 319–328. Association for Computing Machinery, 2019.
- J. Chai, T. Jang, and X. Wang. Fairness without demographics through knowledge distillation. In *Advances in Neural Information Processing Systems*, pages 19152–19164. Curran Associates, Inc., 2022.
- N. V. Chawla, K. W. Bowyer, L. O. Hall, and W. P. Kegelmeyer. SMOTE: Synthetic minority over-sampling technique. *Journal of Artificial Intelligence Research*, 16(1):321–357, 2002.
- W. Chen, Y. Klochkov, and Y. Liu. Post-hoc bias scoring is optimal for fair classification. *arXiv preprint arXiv:2310.05725*, 2023.
- J. Cho, G. Hwang, and C. Suh. A fair classifier using kernel density estimation. In *Advances in Neural Information Processing Systems*, pages 15088–15099. Curran Associates, Inc., 2020.
- E. Chzhen, C. Denis, M. Hebiri, L. Oneto, and M. Pontil. Leveraging labeled and unlabeled data for consistent fair binary classification. In *Advances in Neural Information Processing Systems*. Curran Associates, Inc., 2019.
- S. Corbett-Davies, E. Pierson, A. Feller, S. Goel, and A. Huq. Algorithmic decision making and the cost of fairness. In *Proceedings of the 23rd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, pages 797–806. Association for Computing Machinery, 2017.
- S. Corbett-Davies, J. D. Gaebler, H. Nilforoshan, R. Shroff, and S. Goel. The measure and mismeasure of fairness. *Journal of Machine Learning Research*, 24(312):1–117, 2023.
- A. Cotter, H. Jiang, M. Gupta, S. Wang, T. Narayan, S. You, and K. Sridharan. Optimization with non-differentiable constraints with applications to fairness, recall, churn, and other goals. *Journal of Machine Learning Research*, 20(172):1–59, 2019.
- E. Creager, D. Madras, J.-H. Jacobsen, M. Weis, K. Swersky, T. Pitassi, and R. Zemel. Flexibly fair representation learning by disentanglement. In *Proceedings of the 36th International Conference on Machine Learning*, pages 1436–1445. PMLR, 2019.
- C. Denis, R. Elie, M. Hebiri, and F. Hu. Fairness guarantee in multi-class classification. *arXiv preprint arXiv:2109.13642*, 2021.
- L. Devroye, L. Györfi, and G. Lugosi. *A Probabilistic Theory of Pattern Recognition*. Springer New York, NY, 1996.
- C. Dwork, M. Hardt, T. Pitassi, O. Reingold, and R. Zemel. Fairness through awareness. In *Proceedings of the 3rd Innovations in Theoretical Computer Science Conference*, page 214–226. Association for Computing Machinery, 2012.



- Y. Eitan, N. Cavaglione, M. Arbel, and S. Cohen. Fair synthetic data does not necessarily lead to fair models. In *NeurIPS 2022 Workshop on Synthetic Data for Empowering ML Research*, 2022.
- C. Elkan. The foundations of cost-sensitive learning. In *Proceedings of the 17th International Joint Conference on Artificial Intelligence - Volume 2*, pages 973–978. Morgan Kaufmann Publishers Inc., 2001.
- Executive Office of the President. Big data: A report on algorithmic systems, opportunity, and civil rights. The White House, 2016. URL [https://obamawhitehouse.archives.gov/sites/default/files/microsites/ostp/2016\\_0504\\_data\\_discrimination.pdf](https://obamawhitehouse.archives.gov/sites/default/files/microsites/ostp/2016_0504_data_discrimination.pdf).
- M. Feldman, S. A. Friedler, J. Moeller, C. Scheidegger, and S. Venkatasubramanian. Certifying and removing disparate impact. In *Proceedings of the 21th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, pages 259–268. Association for Computing Machinery, 2015.
- B. Fish, J. Kun, and Á. D. Lelkes. A confidence-based approach for balancing fairness and accuracy. In *Proceedings of the 2016 SIAM International Conference on Data Mining*, pages 144–152. SIAM, 2016.
- A. W. Flores, K. Bechtel, and C. T. Lowenkamp. False positives, false negatives, and false analyses: A rejoinder to Machine Bias: “There’s software used across the country to predict future criminals. And it’s biased against blacks”. *Federal Probation*, 80(2):38–46, 2016.
- G. Goh, A. Cotter, M. Gupta, and M. P. Friedlander. Satisfying real-world goals with dataset constraints. In *Advances in Neural Information Processing Systems*. Curran Associates, Inc., 2016.
- M. Gupta and Q. Mohammad. Advances in AI and ML are reshaping healthcare. TechCrunch, Mar. 2017. URL <https://techcrunch.com/2017/03/16/advances-in-ai-and-ml-are-reshaping-healthcare/>.
- M. Hardt, E. Price, E. Price, and N. Srebro. Equality of opportunity in supervised learning. In *Advances in Neural Information Processing Systems*. Curran Associates, Inc., 2016.
- T. Jang, F. Zheng, and X. Wang. Constructing a fair classifier with generated fair data. In *Proceedings of the AAAI Conference on Artificial Intelligence*, pages 7908–7916, 2021.
- T. Jang, P. Shi, and X. Wang. Group-aware threshold adaptation for fair classification. In *Proceedings of the AAAI Conference on Artificial Intelligence*, pages 6988–6995, 2022.
- R. Jiang, A. Pacchiano, T. Stepleton, H. Jiang, and S. Chiappa. Wasserstein fair classification. In *Proceedings of The 35th Uncertainty in Artificial Intelligence Conference*, pages 862–872. PMLR, 2020.
- J. E. Johndrow and K. Lum. An algorithm for removing sensitive information: Application to race-independent recidivism prediction. *The Annals of Applied Statistics*, 13(1):189–220, 2019.
- M. Joseph, M. Kearns, J. H. Morgenstern, and A. Roth. Fairness in learning: Classic and contextual bandits. In *Advances in Neural Information Processing Systems*. Curran Associates, Inc., 2016.
- F. Kamiran and T. Calders. Data preprocessing techniques for classification without discrimination. *Knowledge and Information Systems*, 33:1–33, 2012.
- M. J. Kusner, J. Loftus, C. Russell, and R. Silva. Counterfactual fairness. In *Advances in Neural Information Processing Systems*. Curran Associates, Inc., 2017.
- P. Lahoti, K. P. Gummadi, and G. Weikum. iFair: Learning individually fair data representations for algorithmic decision making. In *2019 IEEE 35th International Conference on Data Engineering (ICDE)*, pages 1334–1345. IEEE, 2019.
- E. L. Lehmann and J. P. Romano. *Testing Statistical Hypotheses*. Springer New York, NY, 3rd edition, 2005.
- F. Locatello, G. Abbati, T. Rainforth, S. Bauer, B. Schölkopf, and O. Bachem. On the fairness of disentangled representations. In *Advances in Neural Information Processing Systems*. Curran Associates, Inc., 2019.
- C. Louizos, K. Swersky, Y. Li, M. Welling, and R. S. Zemel. The variational fair autoencoder. In *4th International Conference on Learning Representations (ICLR)*, 2016.
- K. Lum and J. Johndrow. A statistical framework for fair predictive algorithms. *arXiv preprint arXiv:1610.08077*, 2016.
- X. Ma, J. Sha, D. Wang, Y. Yu, Q. Yang, and X. Niu. Study on a prediction of P2P network loan default based on the machine learning LightGBM and XGboost algorithms according to different high dimensional data cleaning. *Electronic Commerce Research and Applications*, 31:24–39, 2018.
- N. Mehrabi, F. Morstatter, N. Saxena, K. Lerman, and A. Galstyan. A survey on bias and fairness in machine learning. *ACM Computing Surveys*, 54(6):1–35, 2021.
- A. K. Menon and R. C. Williamson. The cost of fairness in binary classification. In *Proceedings of the 1st Conference on Fairness, Accountability and Transparency*, pages 107–118. PMLR, 2018.
- P. Nandy, C. DiCiccio, D. Venugopalan, H. Logan, K. Basu, and N. El Karoui. Achieving fairness via post-processing in web-scale recommender systems. In *Proceedings of the 2022 ACM Conference on Fairness, Accountability, and Transparency*, page 715–725. Association for Computing Machinery, 2022.
- H. Narasimhan. Learning with complex loss functions and constraints. In *Proceedings of the Twenty-First International Conference on Artificial Intelligence and Statistics*, pages 1646–1654. PMLR, 2018.

- H. Narasimhan, A. Cotter, M. Gupta, and S. Wang. Pairwise fairness for ranking and regression. In *Proceedings of the AAAI Conference on Artificial Intelligence*, pages 5248–5255, 2020.
- J. Neyman and E. S. Pearson. On the problem of the most efficient tests of statistical hypotheses. *Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character*, 231:289–337, 1933.
- L. Oneto, M. Donini, and M. Pontil. General fair empirical risk minimization. In *2020 International Joint Conference on Neural Networks (IJCNN)*, pages 1–8. IEEE, 2020.
- G. Pleiss, M. Raghavan, F. Wu, J. Kleinberg, and K. Q. Weinberger. On fairness and calibration. In *Advances in Neural Information Processing Systems*. Curran Associates, Inc., 2017.
- B. Rava, W. Sun, G. M. James, and X. Tong. A burden shared is a burden halved: A fairness-adjusted approach to classification. *arXiv preprint arXiv:2110.05720*, 2021.
- H. Royden and P. Fitzpatrick. *Real Analysis*. Pearson, 4th edition, 2010.
- T. Salazar, M. S. Santos, H. Araújo, and P. H. Abreu. FAWOS: Fairness-aware oversampling algorithm based on distributions of sensitive attributes. *IEEE Access*, 9:81370–81379, 2021.
- P. Sattigeri, S. C. Hoffman, V. Chenthamarakshan, and K. R. Varshney. Fairness GAN: Generating datasets with fairness properties using a generative adversarial network. *IBM Journal of Research and Development*, 63(4/5): 3:1–3:9, 2019.
- N. Schreuder and E. Chzhen. Classification with abstention but without disparities. In *Proceedings of the Thirty-Seventh Conference on Uncertainty in Artificial Intelligence*, pages 1227–1236. PMLR, 2021.
- J. Shao. *Mathematical Statistics*. Springer New York, NY, 2nd edition, 2003.
- C. Silvia, J. Ray, S. Tom, P. Aldo, J. Heinrich, and A. John. A general approach to fairness with optimal transport. In *Proceedings of the AAAI Conference on Artificial Intelligence*, pages 3633–3640, 2020.
- S.-C. Tsai, C.-H. Chen, Y.-T. Shiao, J.-S. Ciou, and T.-N. Wu. Precision education with statistical learning and deep learning: A case study in Taiwan. *International Journal of Educational Technology in Higher Education*, 17(12): 1–13, 2020.
- UNESCO. Recommendation on the ethics of artificial intelligence. UNESCO, 2021. URL <https://unesdoc.unesco.org/ark:/48223/pf0000380455>.
- I. Valera, A. Singla, and M. Gomez Rodriguez. Enhancing the accuracy and fairness of human decision making. In *Advances in Neural Information Processing Systems*. Curran Associates, Inc., 2018.
- C. Wadsworth, F. Vera, and C. Piech. Achieving fairness through adversarial learning: An application to recidivism prediction. *arXiv preprint arXiv:1807.00199*, 2018.
- Z. Wang, K. Qinami, I. C. Karakozis, K. Genova, P. Nair, K. Hata, and O. Russakovsky. Towards fairness in visual recognition: Effective strategies for bias mitigation. In *2020 IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, pages 8916–8925. IEEE, 2020.
- H. Weerts, M. Dudík, R. Edgar, A. Jalali, R. Lutz, and M. Madaio. Fairlearn: Assessing and improving fairness of AI systems. *Journal of Machine Learning Research*, 24(257):1–8, 2023.
- D. Wei, K. N. Ramamurthy, and F. Calmon. Optimized score transformation for consistent fair classification. *Journal of Machine Learning Research*, 22(258):1–78, 2021.
- R. Xian, L. Yin, and H. Zhao. Fair and optimal classification via post-processing. In *Proceedings of the 40th International Conference on Machine Learning*, pages 37977–38012. PMLR, 2023.
- D. Xu, S. Yuan, L. Zhang, and X. Wu. FairGAN: Fairness-aware generative adversarial networks. In *2018 IEEE International Conference on Big Data (Big Data)*, pages 570–575. IEEE, 2018.
- D. Xu, Y. Wu, S. Yuan, L. Zhang, and X. Wu. Achieving causal fairness through generative adversarial networks. In *Proceedings of the Twenty-Eighth International Joint Conference on Artificial Intelligence, IJCAI-19*, pages 1452–1458. International Joint Conferences on Artificial Intelligence Organization, 2019.
- S. Xu and T. Strohmer. Fair data representation for machine learning at the Pareto frontier. *Journal of Machine Learning Research*, 24(331):1–63, 2023.
- M. B. Zafar, I. Valera, M. Rodriguez, K. Gummadi, and A. Weller. From parity to preference-based notions of fairness in classification. In *Advances in Neural Information Processing Systems*. Curran Associates, Inc., 2017.
- M. B. Zafar, I. Valera, M. Gomez-Rodriguez, and K. P. Gummadi. Fairness constraints: A flexible approach for fair classification. *Journal of Machine Learning Research*, 20(75):1–42, 2019.
- R. Zemel, Y. Wu, K. Swersky, T. Pitassi, and C. Dwork. Learning fair representations. In *Proceedings of the 30th International Conference on Machine Learning*, pages 325–333. PMLR, 2013.
- X. Zeng, E. Dobriban, and G. Cheng. Bayes-optimal classifiers under group fairness. *arXiv preprint arXiv:2202.09724*, 2022a.
- X. Zeng, E. Dobriban, and G. Cheng. Fair Bayes-optimal classifiers under predictive parity. In *Advances in Neural Information Processing Systems*, pages 27692–27705. Curran Associates, Inc., 2022b.

B. H. Zhang, B. Lemoine, and M. Mitchell. Mitigating unwanted biases with adversarial learning. In *Proceedings of the 2018 AAAI/ACM Conference on AI, Ethics, and Society*, pages 335–340. Association for Computing Machinery, 2018.

J. Zhang and E. Bareinboim. Fairness in decision-making - The causal explanation formula. In *Proceedings of the AAAI Conference on Artificial Intelligence*, 2018.

## Appendix

**Additional notation and conventions.** In this appendix, we use some additional notation. For a real-valued function  $f$  defined on  $[a, b)$  for some  $a < b$ , we denote by  $\lim_{x \rightarrow a^+} f(x)$  the limit from the right of  $f$  at  $a$ , if it exists. Similarly, if  $f$  is defined on  $(b, a]$  for  $b < a$ , we denote by  $\lim_{x \rightarrow a^-} f(x)$  the limit from the left of  $f$  at  $a$ , if it exists. For an interval  $[a, b]$ , and scalars  $c \in \mathbb{R}$ ,  $d > 0$ , we denote  $c + d[a, b] = [c + da, c + db]$ . For a classifier  $f$ , we denote  $\text{Acc}(f) = 1 - R(f)$ . When needed, we define  $0/0 := 0$ .

## A Proofs

In Sections A.1 to A.4, we present the proofs of our theoretical results from the main text, except for results in Section 4.3.1 (Theorem 4.7 and Theorem 4.8). We provide an independent Appendix B to discuss the interesting extension of our theoretical and methodological framework. We first introduce several technical lemmas that are essential for proving our theoretical results (Section A.1).

### A.1 Additional Lemmas

**Lemma A.1** (Risk and Accuracy as Linear Functionals of Classifiers). *For any classifier  $f : \mathcal{X} \times \mathcal{A} \rightarrow [0, 1]$ , we have*

$$R(f) = \sum_{a \in \{0,1\}} p_a \int (1 - 2\eta_a(x)) f(x, a) d\mathbb{P}_{X|A=a}(x) + \sum_{a \in \{0,1\}} p_a \int \eta_a(x) d\mathbb{P}_{X|A=a}(x), \quad (\text{A.1})$$

and

$$\text{Acc}(f) = \sum_{a \in \{0,1\}} p_a \int (2\eta_a(x) - 1) f(x, a) d\mathbb{P}_{X|A=a}(x) + \sum_{a \in \{0,1\}} p_a \int (1 - \eta_a(x)) d\mathbb{P}_{X|A=a}(x), \quad (\text{A.2})$$

*Proof.* By definition,  $\widehat{Y}$  is conditionally independent of  $Y$  given  $X$  and  $A$ . Thus,

$$\begin{aligned} \mathbb{P}(\widehat{Y}_f = 1, Y = 0 \mid X = x, A = a) &= f(x, a)(1 - \eta_a(x)), \\ \mathbb{P}(\widehat{Y}_f = 1, Y = 0 \mid X = x, A = a) &= \eta_a(x)(1 - f(x, a)). \end{aligned}$$

This implies that

$$\begin{aligned} R(f) &= \mathbb{P}(Y \neq \widehat{Y}_f) = \sum_{a \in \mathcal{A}} p_a \mathbb{P}(Y \neq \widehat{Y}_f \mid A = a) \\ &= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} \left( \mathbb{P}(\widehat{Y}_f = 1, Y = 0 \mid X = x, A = a) + \mathbb{P}(\widehat{Y}_f = 1, Y = 0 \mid X = x, A = a) \right) d\mathbb{P}_{X|A=a}(x) \\ &= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} (f(x, a)(1 - \eta_a(x)) + \eta_a(x)(1 - f(x, a))) d\mathbb{P}_{X|A=a}(x) \\ &= \sum_{a \in \{0,1\}} p_a \int (1 - 2\eta_a(x)) f(x, a) d\mathbb{P}_{X|A=a}(x) + \sum_{a \in \{0,1\}} p_a \int (\eta_a(x)) d\mathbb{P}_{X|A=a}(x). \end{aligned}$$

Moreover,

$$\begin{aligned}
\text{Acc}(f) &= 1 - R(f) = 1 - \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} (f(x,a)(1 - \eta_a(x)) + \eta_a(x)(1 - f(x,a))) d\mathbb{P}_{X|A=a}(x) \\
&= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} (1 - \eta_a(x) - (1 - 2\eta_a(x))f(x,a)) d\mathbb{P}_{X|A=a}(x) \\
&= \sum_{a \in \{0,1\}} p_a \int (2\eta_a(x) - 1) f(x,a) d\mathbb{P}_{X|A=a}(x) + \sum_{a \in \{0,1\}} p_a \int (1 - \eta_a(x)) d\mathbb{P}_{X|A=a}(x).
\end{aligned}$$

□

**Lemma A.2** (Characterizing Bayes-Optimal Classifiers for Linear Disparity Measures). *For  $t \in \mathbb{R}$  and  $\delta \geq 0$ , let  $f_{\text{Dis},t}$ ,  $D_{\text{Dis}}(t)$  and  $t_{\text{Dis}}(\delta)$  be defined in (4.1), (4.2) and (4.3), respectively. Then, using the convention that  $0/0 = 0$ , for any fixed  $t$ , we have*

$$f_{\text{Dis},t} = \arg \min_{f \in \mathcal{F}} \left\{ R(f) : \frac{t \cdot \text{Dis}(f)}{|t|} \leq \frac{t \cdot D_{\text{Dis}}(t)}{|t|} \right\}.$$

Moreover, for all classifiers  $f' \in \arg \min_{f \in \mathcal{F}} \{R(f) : t \cdot \text{Dis}(f)/|t| \leq t \cdot D_{\text{Dis}}(t)/|t|\}$ ,  $f' = f_{\text{Dis},t}$  almost surely with respect to  $\mathbb{P}_{X,A}$ . In addition, if  $t \in [\min(0, t_{\text{Dis}}(0)), \max(0, t_{\text{Dis}}(0))]$ ,

$$f_{\text{Dis},t} = \arg \min_{f \in \mathcal{F}} \{R(f) : |\text{Dis}(f)| \leq |D_{\text{Dis}}(t)|\}.$$

*Proof.* This result is a consequence of the generalized Neyman-Pearson lemma. If  $t = 0$ , the result follows since  $f_{\text{Dis},0}$  is the unconstrained Bayes-optimal classifier. When  $t \neq 0$ , let, for all  $x, a$ ,  $\phi_0(x, a) = 2\eta_1(x, a) - 1$ ,  $\phi_1(x, a) = w_{\text{Dis}}(x, a)$  and for all classifiers  $f$ , and  $t \in \mathbb{R}$ ,  $\overline{\text{Dis}}_t(f) = t \cdot \text{Dis}(f)/|t|$ . We have, for all  $x, a$ ,

$$f_{\text{Dis},t}(x, a) = I(\phi_0(x, a) > t\phi_1(x, a)) = I\left(\phi_0(x, a) > |t| \frac{t\phi_1(x, a)}{|t|}\right).$$

Moreover, by Lemma A.1 and (3.6), we can write  $\text{Acc}(f)$  and  $\overline{D}_t(f)$  as

$$\begin{aligned}
\text{Acc}(f) &= \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, a) \phi_0(x, a) d\mathbb{P}_{X,A}(x, a) + \int_{\mathcal{A}} \int_{\mathcal{X}} (1 - \eta_a(x)) d\mathbb{P}_{X,A}(x, a); \\
\overline{\text{Dis}}_t(f) &= \frac{t}{|t|} \text{Dis}(f) = \int_{\mathcal{X} \times \mathcal{A}} f(x, a) \frac{t\phi_1(x, a)}{|t|} d\mathbb{P}_{X,A}(x, a).
\end{aligned}$$

Define:

$$\mathcal{F}_{t,=} = \left\{ f : \overline{\text{Dis}}_t(f) = \frac{tD_{\text{Dis}}(t)}{|t|} \right\}; \mathcal{F}_{t,|\cdot|, \leq} = \left\{ f : |\overline{\text{Dis}}_t(f)| \leq \frac{tD_{\text{Dis}}(t)}{|t|} \right\}; \text{ and } \mathcal{F}_{t, \leq} = \left\{ f : \overline{\text{Dis}}_t(f) \leq \frac{tD_{\text{Dis}}(t)}{|t|} \right\}.$$

It is clear that  $f_{\text{Dis},t} \in \mathcal{F}_{t,=} \subset \mathcal{F}_{t, \leq}$ . Since  $|t| \geq 0$ , by the generalized Neyman-Pearson lemma (Lemma 3.1),

$$f_{\text{Dis},t} \in \arg \max_{f \in \mathcal{F}_{t, \leq}} \text{Acc}(f),$$

and, since  $\mathbb{P}_{X|A=a}(\eta_a(X) = 1/2 + t_{\text{Dis}}(\delta)w_{\text{Dis}}(X, a)/2) = 0$  when both  $\eta_a(X)$  and  $w_{\text{Dis}}(X, a)$  are continuous random variables with respect to  $\mathbb{P}_{X|A=a}$ , for all  $f' \in \arg \max_{f \in \mathcal{F}_{t, \leq}} \text{Acc}(f)$ ,  $f' = f_{\text{Dis},t}$  almost surely with respect to  $\mathbb{P}_{X,A}$ .

Now, suppose further that  $t \in [\min(0, t_{\text{Dis}}(0)), \max(0, t_{\text{Dis}}(0))]$ ,  $tD(0) \geq 0$ . By result (1) of proposition 4.1,  $D_{\text{Dis}}(t)$  is monotone non-increasing with respect to  $t$ . By the definition of  $t_{\text{Dis}}(\delta)$  in (4.3), we have  $t_{\text{Dis}}(0) \geq 0$  when  $D_{\text{Dis}}(0) \geq 0$  and  $t_{\text{Dis}}(0) \leq 0$  when  $D_{\text{Dis}}(0) \leq 0$ . In both cases, we have  $t_{\text{Dis}}(0) \cdot D_{\text{Dis}}(0) \geq 0$ . This further implies  $t \cdot D_{\text{Dis}}(0) \geq 0$  and consequently,  $f_{\text{Dis},t} \in \mathcal{F}_{t,=} \subset \mathcal{F}_{t,|\cdot|, \leq} \subset \mathcal{F}_{t, \leq}$ . Then,

$$\max_{f \in \mathcal{F}_{t, \leq}} \text{Acc}(f) = \text{Acc}(f_{\text{Dis},t}) \leq \max_{f \in \mathcal{F}_{t,|\cdot|, \leq}} \text{Acc}(f) \leq \max_{f \in \mathcal{F}_{t, \leq}} \text{Acc}(f).$$

Thus, we can conclude that

$$f_t = \arg \max_{f \in \mathcal{F}_{t, |\cdot|, \leq}} \text{Acc}(f) = \arg \min_{f \in \mathcal{F}} \left\{ R(f) : |\text{Dis}(f)| \leq \frac{t \cdot D_{\text{Dis}}(t)}{|t|} \right\}.$$

□

**Lemma A.3** (Properties of the Tradeoff function). *For any  $\delta \geq 0$ , let  $t_{\text{Dis}}(\delta)$  and  $T(\delta)$  be defined in (4.2) and (4.7), respectively. Let for  $a \in \{0, 1\}$ ,  $\eta_a(X)$  and  $w_{\text{Dis}}(X, a)$  be continuous random variables when  $X \sim \mathbb{P}_{X|A=a}$ , for  $a \in \{0, 1\}$ . Then, for  $\min(0, D_{\text{Dis}}(0)) \leq \delta_1 < \delta_2 \leq \max(0, D_{\text{Dis}}(0))$ ,*

$$t_{\text{Dis}}(\delta_2)(\delta_2 - \delta_1) \leq T(\delta_1) - T(\delta_2) \leq t_{\text{Dis}}(\delta_1)(\delta_2 - \delta_1). \quad (\text{A.3})$$

*Proof.* As  $D_{\text{Dis}}(t)$  is monotone non-increasing and continuous in  $t \in \mathbb{R}$ , it follows that  $t_{\text{Dis}}(\delta)$  is strictly decreasing in  $\delta$  on  $[0, |D_{\text{Dis}}(0)|]$ . Then, we have  $t_{\text{Dis}}(\delta_1) > t_{\text{Dis}}(\delta_2)$ . For  $a \in \{0, 1\}$ , we define

$$\mathcal{T}_{a,+} = \left\{ x \in \mathcal{X}, w_{\text{Dis}}(x, a) > 0, t_{\text{Dis}}(\delta_2) \leq \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} < t_{\text{Dis}}(\delta_1) \right\},$$

and

$$\mathcal{T}_{a,-} = \left\{ x \in \mathcal{X}, w_{\text{Dis}}(x, a) < 0, t_{\text{Dis}}(\delta_2) \leq \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} < t_{\text{Dis}}(\delta_1) \right\}.$$

By definition,

$$\begin{aligned} & f_{t_{\text{Dis}}(\delta_1)}(x, a) - f_{t_{\text{Dis}}(\delta_2)}(x, a) \\ &= I \left( \eta_a(x) > \frac{1}{2} + \frac{t_{\text{Dis}}(\delta_1)}{2} w_{\text{Dis}}(x, a) \right) - I \left( \eta_a(x) > \frac{1}{2} + \frac{t_{\text{Dis}}(\delta_2)}{2} w_{\text{Dis}}(x, a) \right) \\ &= \begin{cases} -I \left( t_{\text{Dis}}(\delta_2) \leq \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} < t_{\text{Dis}}(\delta_1) \right), & w_{\text{Dis}}(x, a) > 0; \\ I \left( t_{\text{Dis}}(\delta_2) \leq \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} < t_{\text{Dis}}(\delta_1) \right), & w_{\text{Dis}}(x, a) < 0; \\ 0, & w_{\text{Dis}}(x, a) = 0, \end{cases} \end{aligned}$$

In addition, we have, on  $\mathcal{T}_{a,+}$ ,

$$t_{\text{Dis}}(\delta_2) \cdot w_{\text{Dis}}(x, a) \leq 2\eta_a(x) - 1 \leq t_{\text{Dis}}(\delta_1) \cdot w_{\text{Dis}}(x, a),$$

and on  $\mathcal{T}_{a,-}$ ,

$$t_{\text{Dis}}(\delta_1) \cdot w_{\text{Dis}}(x, a) \leq 2\eta_a(x) - 1 \leq t_{\text{Dis}}(\delta_2) \cdot w_{\text{Dis}}(x, a).$$

Then, by (A.1) from Lemma A.1,

$$\begin{aligned} T(\delta_1) - T(\delta_2) &= R(f_{t_{\text{Dis}}(\delta_1)}) - R(f_{t_{\text{Dis}}(\delta_2)}) \\ &= \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{X}} (1 - 2\eta_a(x)) (f_{t_{\text{Dis}}(\delta_1)}(x, a) - f_{t_{\text{Dis}}(\delta_2)}(x, a)) d\mathbb{P}_{X|A=a}(x) \\ &= \sum_{a \in \mathcal{A}} p_a \int_{\{w_{\text{Dis}}(x, a) > 0\}} (2\eta_a(x) - 1) I \left( t_{\text{Dis}}(\delta_2) \leq \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} < t_{\text{Dis}}(\delta_1) \right) d\mathbb{P}_{X|A=a}(x) \\ &\quad - \sum_{a \in \mathcal{A}} p_a \int_{\{w_{\text{Dis}}(x, a) < 0\}} (2\eta_a(x) - 1) I \left( t_{\text{Dis}}(\delta_2) \leq \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} < t_{\text{Dis}}(\delta_1) \right) d\mathbb{P}_{X|A=a}(x) \\ &= \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{T}_{a,+}} (2\eta_a(x) - 1) d\mathbb{P}_{X|A=a}(x) - \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{T}_{a,-}} (2\eta_a(x) - 1) d\mathbb{P}_{X|A=a}(x) \\ &\leq t_{\text{Dis}}(\delta_1) \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{T}_{a,+}} w_{\text{Dis}}(x, a) d\mathbb{P}_{X|A=a}(x) - t_{\text{Dis}}(\delta_1) \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{T}_{a,-}} w_{\text{Dis}}(x, a) d\mathbb{P}_{X|A=a}(x) \\ &= -t_{\text{Dis}}(\delta_1) \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{X} \cap \{w_{\text{Dis}}(x, a) > 0\}} w_{\text{Dis}}(x, a) \cdot (f_{t_{\text{Dis}}(\delta_1)}(x, a) - f_{t_{\text{Dis}}(\delta_2)}(x, a)) d\mathbb{P}_{X|A=a}(x) \end{aligned}$$

$$\begin{aligned}
& -t_{\text{Dis}}(\delta_1) \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{X} \cap \{w_{\text{Dis}}(x,a) < 0\}} w_{\text{Dis}}(x,a) \cdot (f_{t_{\text{Dis}}(\delta_1)}(x,a) - f_{t_{\text{Dis}}(\delta_2)}(x,a)) d\mathbb{P}_{X|A=a}(x) \\
& = -t_{\text{Dis}}(\delta_1) \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{X}} w_{\text{Dis}}(x,a) \cdot (f_{t_{\text{Dis}}(\delta_1)}(x,a) - f_{t_{\text{Dis}}(\delta_2)}(x,a)) d\mathbb{P}_{X|A=a}(x) \\
& = -t_{\text{Dis}}(\delta_1) (D(f_{t_{\text{Dis}}(\delta_1)}) - D(f_{t_{\text{Dis}}(\delta_2)})) = t_{\text{Dis}}(\delta_1)(\delta_2 - \delta_1).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
T(\delta_1) - T(\delta_2) & = \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{T}_{a,+}} (2\eta_a(x) - 1) d\mathbb{P}_{X|A=a}(x) - \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{T}_{a,+}} (2\eta_a(x) - 1) d\mathbb{P}_{X|A=a}(x) \\
& \geq t_{\text{Dis}}(\delta_1) \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{T}_{a,+}} w_{\text{Dis}}(x,a) d\mathbb{P}_{X|A=a}(x) - t_{\text{Dis}}(\delta_2) \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{T}_{a,-}} w_{\text{Dis}}(x,a) d\mathbb{P}_{X|A=a}(x) \\
& = -t_{\text{Dis}}(\delta_2) \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{X} \cap \{w_{\text{Dis}}(x,a) > 0\}} w_{\text{Dis}}(x,a) \cdot (f_{t_{\text{Dis}}(\delta_1)}(x,a) - f_{t_{\text{Dis}}(\delta_2)}(x,a)) d\mathbb{P}_{X|A=a}(x) \\
& \quad - t_{\text{Dis}}(\delta_2) \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{X} \cap \{w_{\text{Dis}}(x,a) < 0\}} w_{\text{Dis}}(x,a) \cdot (f_{t_{\text{Dis}}(\delta_1)}(x,a) - f_{t_{\text{Dis}}(\delta_2)}(x,a)) d\mathbb{P}_{X|A=a}(x) \\
& = -t_{\text{Dis}}(\delta_2) \sum_{a \in \mathcal{A}} p_a \int_{\mathcal{X}} w_{\text{Dis}}(x,a) \cdot (f_{t_{\text{Dis}}(\delta_1)}(x,a) - f_{t_{\text{Dis}}(\delta_2)}(x,a)) d\mathbb{P}_{X|A=a}(x) \\
& = -t_{\text{Dis}}(\delta_2) (D(f_{\delta_1}^*) - D(f_{\delta_2}^*)) = t_{\text{Dis}}(\delta_2)(\delta_2 - \delta_1).
\end{aligned}$$

We conclude (A.3). □

## A.2 Proofs of Results from Section 3

### A.2.1 Proof of Proposition 3.4

By Bayes' theorem,

$$\eta_a(x) = \mathbb{P}(Y = 1 | X = x, A = a) = \frac{\mathbb{P}(A = a, Y = 1) d\mathbb{P}_{X|A=a, Y=1}(x)}{\mathbb{P}(A = a) d\mathbb{P}_{X|A=a}(x)} = \frac{p_{a,1} d\mathbb{P}_{X|A=a, Y=1}(x)}{p_a d\mathbb{P}_{X|A=a}(x)},$$

and

$$p_a d\mathbb{P}_{X|A=a}(x) = p_{a,1} d\mathbb{P}_{X|A=a, Y=1}(x) + p_{a,0} d\mathbb{P}_{X|A=a, Y=0}(x).$$

It follows that

$$\frac{d\mathbb{P}_{X|A=a, Y=1}(x)}{d\mathbb{P}_{X|A=a}(x)} = \frac{p_a \eta(x)}{p_{a,1}}; \quad \text{and} \quad \frac{d\mathbb{P}_{X|A=a, Y=0}(x)}{d\mathbb{P}_{X|A=a}(x)} = \frac{p_a(1 - \eta(x))}{p_{a,0}}.$$

Let  $f$  be any classifier. Its DD, DO and PD can be expressed in turn as

$$\begin{aligned}
\text{DD}(f) & = \mathbb{P}(\hat{Y}_f = 1 | A = 1) - \mathbb{P}(\hat{Y}_f = 1 | A = 0) \\
& = \int_{\mathcal{X}} f(x, 1) d\mathbb{P}_{X|A=1}(x) - \int_{\mathcal{X}} f(x, 0) d\mathbb{P}_{X|A=0}(x) \\
& = \sum_{a \in \{0,1\}} p_a \left( \int_{\mathcal{X}} \left[ \frac{I(a=1)}{p_1} f(x,a) - \frac{I(a=0)}{p_0} f(x,a) \right] d\mathbb{P}_{X|A=a}(x) \right) \\
& = \int_{\mathcal{A}} \int_{\mathcal{X}} f(x,a) \left( \frac{2a-1}{p_a} \right) d\mathbb{P}_{X|A=a} d\mathbb{P}_A(a) \\
& = \int_{\mathcal{A}} \int_{\mathcal{X}} f(x,a) (2a-1) \left( \frac{1}{p_a} \right) d\mathbb{P}_{X,A}(x,a); \tag{A.4}
\end{aligned}$$

$$\begin{aligned}
\text{DO}(f) &= \mathbb{P}(\widehat{Y}_f = 1|A = 1, Y = 1) - \mathbb{P}(\widehat{Y}_f = 1|A = 0, Y = 1) \\
&= \int_{\mathcal{X}} f(x, 1) d\mathbb{P}_{X|A=1, Y=1}(x) - \int_{\mathcal{X}} f(x, 0) d\mathbb{P}_{X|A=0, Y=1}(x) \\
&= \sum_{a \in \{0,1\}} p_a \left( \int_{\mathcal{X}} \left[ \frac{I(a=1)}{p_1} f(x, a) - \frac{I(a=0)}{p_0} f(x, a) \right] d\mathbb{P}_{X|A=a, Y=1}(x) \right) \\
&= \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, a) \left( \frac{2a-1}{p_a} \right) \frac{d\mathbb{P}_{X|A=a, Y=1}(x)}{d\mathbb{P}_{X|A=a}} d\mathbb{P}_{X|A=a} d\mathbb{P}_A(a) \\
&= \int_{\mathcal{X} \times \mathcal{A}} f(x, a) (2a-1) \left( \frac{\eta_a(x)}{p_{a,1}} \right) d\mathbb{P}_{X,A}(x, a); \tag{A.5}
\end{aligned}$$

$$\begin{aligned}
\text{PD}(f) &= \mathbb{P}(\widehat{Y}_f = 1|A = 1, Y = 0) - \mathbb{P}(\widehat{Y}_f = 1|A = 0, Y = 0) \\
&= \int_{\mathcal{X}} f(x, 1) d\mathbb{P}_{X|A=1, Y=0}(x) - \int_{\mathcal{X}} f(x, 0) d\mathbb{P}_{X|A=0, Y=0}(x) \\
&= \sum_{a \in \{0,1\}} p_a \left( \int_{\mathcal{X}} \left[ \frac{I(a=1)}{p_1} f(x, a) - \frac{I(a=0)}{p_0} f(x, a) \right] d\mathbb{P}_{X|A=a, Y=0}(x) \right) \\
&= \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, a) \left( \frac{2a-1}{p_a} \right) \frac{d\mathbb{P}_{X|A=a, Y=0}(x)}{d\mathbb{P}_{X|A=a}} d\mathbb{P}_{X|A=a} d\mathbb{P}_A(a) \\
&= \int_{\mathcal{X} \times \mathcal{A}} f(x, a) (2a-1) \left( \frac{1-\eta_a(x)}{p_{a,0}} \right) d\mathbb{P}_{X,A}(x, a). \tag{A.6}
\end{aligned}$$

Thus, Proposition 3.4 follows.

### A.3 Proofs of Results from Section 4

#### A.3.1 Proof of Proposition 4.1

For (1), since, for  $a \in \{0, 1\}$ , both  $\eta_a(X)$  and  $w_{\text{Dis}}(X, a)$  are continuous random variable given  $A = a$ , we have that for  $a \in \{0, 1\}$ ,  $t \mapsto \mathbb{P}_{X|A=a}(\eta_a(X) > 1/2 + tw_{\text{Dis}}(X, a)/2)$  is a continuous function. Thus,  $t \mapsto D_{\text{Dis}}(t)$  is continuous. Now, define, for  $a \in \{0, 1\}$ ,

$$G_{a,+} = \{x \in \mathcal{X}, w_{\text{Dis}}(x, a) > 0\}; \quad G_{a,0} = \{x \in \mathcal{X}, w_{\text{Dis}}(x, a) = 0\} \quad G_{a,-} = \{x \in \mathcal{X}, w_{\text{Dis}}(x, a) < 0\}.$$

Letting  $t_1 < t_2$ , we have for  $a \in \{0, 1\}$  and  $x \in \mathcal{X}$ ,

$$f_{\text{Dis}, t_1}(x, a) - f_{\text{Dis}, t_2}(x, a) = \begin{cases} I\left(t_1 < \frac{2\eta_a(x)-1}{w_{\text{Dis}}(x, a)} \leq t_2\right), & x \in G_{a,+}; \\ -I\left(t_1 < \frac{2\eta_a(x)-1}{w_{\text{Dis}}(x, a)} \leq t_2\right), & x \in G_{a,-}; \\ 0, & \text{otherwise.} \end{cases} \tag{A.7}$$

It thus follows that,

$$\begin{aligned}
D_{\text{Dis}}(t_1) - D_{\text{Dis}}(t_2) &= \text{Dis}(f_{\text{Dis}, t_1}) - \text{Dis}(f_{\text{Dis}, t_2}) = \int_{\mathcal{A}} \int_{\mathcal{X}} [f_{\text{Dis}, t_1}(x, a) - f_{\text{Dis}, t_2}(x, a)] w_{\text{Dis}}(x, a) d\mathbb{P}_{X,A}(x, a) \\
&= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} [f_{\text{Dis}, t_1}(x, a) - f_{\text{Dis}, t_2}(x, a)] w_{\text{Dis}}(x, a) d\mathbb{P}_{X|A=a} \\
&= \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} I\left(t_1 < \frac{2\eta_a(x)-1}{w_{\text{Dis}}(x, a)} \leq t_2\right) w_{\text{Dis}}(x, a) d\mathbb{P}_{X|A=a} \\
&\quad - \sum_{a \in \{0,1\}} p_a \int_{G_{a,-}} I\left(t_1 < \frac{2\eta_a(x)-1}{w_{\text{Dis}}(x, a)} \leq t_2\right) w_{\text{Dis}}(x, a) d\mathbb{P}_{X|A=a} \geq 0.
\end{aligned}$$

The last inequality holds since the indicator function is non-negative, and  $w_{\text{Dis}}(x, a)$  is positive on  $G_{a,+}$  and negative on  $G_{a,-}$ . Thus,  $t \mapsto D_{\text{Dis}}(t)$  is a monotone non-increasing function.



For (2), when  $t_1 < t_2 < 0$ , it follows that on  $G_{a,+}$ ,

$$(1 - 2\eta_a(x))I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) \geq -t_2 w_{\text{Dis}}(x, a)I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) \geq 0,$$

and, on  $G_{a,-}$ ,

$$(2\eta_a(x) - 1)I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) \geq t_2 w_{\text{Dis}}(x, a)I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) \geq 0.$$

Then, by Lemma A.1 and (A.7),

$$\begin{aligned} R(f_{\text{Dis}, t_1}) - R(f_{\text{Dis}, t_2}) &= \int_{\mathcal{A}} \int_{\mathcal{X}} (2\eta_a(x) - 1) [f_{\text{Dis}, t_1}(x, a) - f_{\text{Dis}, t_2}(x, a)] d\mathbb{P}_{X, A}(x, a) \\ &= \sum_{a \in \{0, 1\}} p_a \int_{\mathcal{X}} (2\eta_a(x) - 1) [f_{\text{Dis}, t_1}(x, a) - f_{\text{Dis}, t_2}(x, a)] d\mathbb{P}_{X|A=a}(x, a) \\ &= \sum_{a \in \{0, 1\}} p_a \int_{G_{a,+}} (1 - 2\eta_a(x))I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) d\mathbb{P}_{X|A=a}(x, a) \\ &\quad + \sum_{a \in \{0, 1\}} p_a \int_{G_{a,-}} (2\eta_a(x) - 1)I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) d\mathbb{P}_{X|A=a}(x, a) \geq 0. \end{aligned}$$

Thus,  $t \mapsto R(f_{\text{Dis}, t})$  is monotone non-increasing on  $(-\infty, 0)$ .

On the other hand, when  $0 \leq t_1 < t_2$ , we have on  $G_{a,+}$  that

$$(1 - 2\eta_a(x))I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) \leq -t_1 w_{\text{Dis}}(x, a)I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) \leq 0,$$

and, on  $G_{a,-}$  that

$$(2\eta_a(x) - 1)I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) \leq t_1 w_{\text{Dis}}(x, a)I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) \leq 0.$$

Then, by Lemma A.1 and (A.7),

$$\begin{aligned} R(f_{\text{Dis}, t_1}) - R(f_{\text{Dis}, t_2}) &= \int_{\mathcal{A}} \int_{\mathcal{X}} (2\eta_a(x) - 1) [f_{\text{Dis}, t_1}(x, a) - f_{\text{Dis}, t_2}(x, a)] d\mathbb{P}_{X, A}(x, a) \\ &= \sum_{a \in \{0, 1\}} p_a \int_{\mathcal{X}} (2\eta_a(x) - 1) [f_{\text{Dis}, t_1}(x, a) - f_{\text{Dis}, t_2}(x, a)] d\mathbb{P}_{X|A=a}(x, a) \\ &= \sum_{a \in \{0, 1\}} p_a \int_{G_{a,+}} (1 - 2\eta_a(x))I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) d\mathbb{P}_{X|A=a}(x, a) \\ &\quad + \sum_{a \in \{0, 1\}} p_a \int_{G_{a,-}} (2\eta_a(x) - 1)I\left(t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2\right) d\mathbb{P}_{X|A=a}(x, a) \leq 0. \end{aligned}$$

Thus,  $t \mapsto R(f_{\text{Dis}, t})$  is monotone non-decreasing on  $[0, \infty)$ . This shows that  $t \mapsto R(f_{\text{Dis}, t})$  is non-increasing in  $|t|$ .

### A.3.2 Proof of Theorem 4.2

We analyze the following three cases separately: (1)  $|\text{Dis}(0)| \leq \delta$ , (2)  $\text{Dis}(0) > \delta$  and (3)  $\text{Dis}(0) < -\delta$ . Since the proof for case (3) is analogous to case (2), we omit the discussion of case (3).

Case (1):  $|\text{Dis}(0)| \leq \delta$ . In this case, the unconstrained Bayes-optimal classifier satisfies the fairness constraint. As a result, we have  $t_{\text{Dis}}(\delta) = 0$  and a  $\delta$ -fair Bayes-optimal classifier is given by  $f_{\text{Dis}, \delta}^*(x, a) = I(\eta_a(x) > 1/2)$ , for all  $x, a$ .

Case (2):  $\text{Dis}(0) > \delta$ . When, for  $a \in \{0, 1\}$ ,  $\eta_a(X)$  has probability density function on  $\mathcal{X}$ ,  $D_{\text{Dis}}(t)$  is a continuous non-increasing function on  $\mathbb{R}$ , and thus we have  $D_{\text{Dis}}(t_{\text{Dis}}(\delta)) = \delta$ . Moreover,  $\text{Dis}(0) > \delta$  indicates  $t_{\text{Dis}}(\delta) > 0$ . Then, by Lemma A.2,

$$f_{\text{Dis}, t_{\text{Dis}}(\delta)} = \arg \min_{f \in \mathcal{F}} \left\{ R(f) : |\text{Dis}(f)| \leq \frac{t \cdot D_{\text{Dis}}(t_{\text{Dis}}(\delta))}{|t|} \right\} = \arg \min_{f \in \mathcal{F}} \{ R(f) : |\text{Dis}(f)| \leq \delta \}.$$

This finishes the proof.

### A.3.3 Proof of Proposition 4.4

Without loss of generality, we assume  $D_{\text{Dis}}(0) \geq 0$ . In this case, we have  $t_0 = 0$  and  $\bar{t}_0 = t_{\text{Dis}}(0)$ . By Lemma A.2, we have

$$\{f_{\text{Dis}, t} : t \in [0, t_{\text{Dis}}(0)]\} \subset \text{FPF}.$$

On the other hand, since  $f_{\text{Dis}, 0} \in \arg \min_{f \in \mathcal{F}} \{R(f)\}$ , and by Lemma A.2,  $f_{\text{Dis}, t_{\text{Dis}}(0)} \in \arg \min_{f \in \mathcal{F}} \{R(f) : \text{Dis}(f) \leq 0\}$ , we have, for  $f_1$  with  $\text{Dis}(f) < 0$  and  $f_2$  with  $\text{Dis}(f) > \delta$ ,

$$R(f_{\text{Dis}, t_{\text{Dis}}(0)}) \leq R(f_1), |\text{Dis}(f_{\text{Dis}, t_{\text{Dis}}(0)})| = 0 < |\text{Dis}(f_1)|,$$

and

$$R(f_0) \leq R(f_2), |\text{Dis}(f_0)| = D_{\text{Dis}}(0) < |\text{Dis}(f_2)|.$$

Thus, for any  $f_{\text{FPF}} \in \text{FPF}$ , we have

$$0 \leq \text{Dis}(f_{\text{FPF}}) \leq \text{Dis}(0).$$

Moreover, by the definition of the fair Pareto frontier, we have

$$R(f_{\text{FPF}}) = \min_{f \in \mathcal{F}} \{R(f) : \text{Dis}(f) \leq \text{Dis}(f_{\text{FPF}})\}.$$

Since  $t \mapsto D_{\text{Dis}}(t)$  is a continuous monotone non-increasing function on  $[0, t_{\text{Dis}}(0)]$  with  $D_{\text{Dis}}(t_{\text{Dis}}(0)) = 0$ , there exists a  $t \in [0, t_{\text{Dis}}(0)]$  such that  $\text{Dis}(f_{\text{Dis}, t}) = D_{\text{Dis}}(t) = \text{Dis}(f_{\text{FPF}})$ . By Lemma A.2,

$$R(f_{\text{Dis}, t}) = \min_{f \in \mathcal{F}} \{R(f) : \text{Dis}(f) = \text{Dis}(f_{\text{FPF}})\}.$$

In conclusion, there exists a  $t \in [0, t_{\text{Dis}}(0)]$  such that,

$$R(f_{\text{Dis}, t}) = R(f_{\text{FPF}}) \quad \text{and} \quad \text{Dis}(f_{\text{Dis}, t}) = \text{Dis}(f_{\text{FPF}}).$$

### A.3.4 Proof of Theorem 4.6

Let  $0 \leq \delta_1 < \delta_2 \leq |D_{\text{Dis}}(0)|$  and  $\lambda \in [0, 1]$ , and denote  $\delta_\lambda = \lambda\delta_1 + (1 - \lambda)\delta_2$ . As  $t_{\text{Dis}}(\delta)$  is strictly decreasing on  $[0, |D_{\text{Dis}}(0)|]$ , we have  $t_{\text{Dis}}(\delta_1) > t_{\text{Dis}}(\delta_\lambda) > t_{\text{Dis}}(\delta_2)$ . By Lemma A.3,

$$\begin{aligned} T(\delta_\lambda) &= \lambda T(\delta_\lambda) + (1 - \lambda)T(\delta_\lambda) = \lambda T(\delta_1) + (1 - \lambda)T(\delta_2) + \lambda(T(\delta_\lambda) - T(\delta_1)) + (1 - \lambda)(T(\delta_\lambda) - T(\delta_2)) \\ &\leq \lambda T(\delta_1) + (1 - \lambda)T(\delta_2) - \lambda t_{\text{Dis}}(\delta_\lambda)(\delta_\lambda - \delta_1) + (1 - \lambda)t_{\text{Dis}}(\delta_\lambda)(\delta_2 - \delta_\lambda) = \lambda T(\delta_1) + (1 - \lambda)T(\delta_2). \end{aligned}$$

The proof is thus completed.

### A.3.5 Proof of Proposition 4.10

By Bayes' theorem, we have for all  $x, y, a$  that

$$\mathbb{P}(A = a \mid X = x) = \frac{p_a d\mathbb{P}_{X|A=a}(x)}{d\mathbb{P}_X(x)},$$

and

$$\mathbb{P}(Y = y \mid A = a, X = x)\mathbb{P}(A = a \mid X = x) = \mathbb{P}(A = a, Y = y \mid X = x) = \frac{p_{a,y} d\mathbb{P}_{X|A=a,Y=y}(x)}{d\mathbb{P}_X(x)}.$$

It then follows that

$$\begin{aligned}
\frac{d\mathbb{P}_{X|A=1}(x)}{d\mathbb{P}_X(x)} &= \frac{\mathbb{P}(A=1|X=x)}{\mathbb{P}(A=1)} = \frac{\eta^A(x)}{p_1}; \\
\frac{d\mathbb{P}_{X|A=0}(x)}{d\mathbb{P}_X(x)} &= \frac{\mathbb{P}(A=0|X=x)}{\mathbb{P}(A=0)} = \frac{1-\eta^A(x)}{p_0}; \\
\frac{d\mathbb{P}_{X|A=1,Y=1}(x)}{d\mathbb{P}_X(x)} &= \frac{\mathbb{P}(Y=1|A=1,X=x)\mathbb{P}(A=1|X=x)}{p_{1,1}} = \frac{\eta_1(x)\eta^A(x)}{p_{1,1}}; \\
\frac{d\mathbb{P}_{X|A=1,Y=0}(x)}{d\mathbb{P}_X(x)} &= \frac{\mathbb{P}(Y=0|A=1,X=x)\mathbb{P}(A=1|X=x)}{p_{1,0}} = \frac{(1-\eta_1(x))\eta^A(x)}{p_{1,0}}; \\
\frac{d\mathbb{P}_{X|A=0,Y=1}(x)}{d\mathbb{P}_X(x)} &= \frac{\mathbb{P}(Y=1|A=0,X=x)\mathbb{P}(A=0|X=x)}{p_{0,1}} = \frac{\eta_0(x)(1-\eta^A(x))}{p_{0,1}}; \\
\frac{d\mathbb{P}_{X|A=0,Y=0}(x)}{d\mathbb{P}_X(x)} &= \frac{\mathbb{P}(Y=0|A=0,X=x)\mathbb{P}(A=0|X=x)}{p_{0,0}} = \frac{(1-\eta_0(x))(1-\eta^A(x))}{p_{0,0}}.
\end{aligned}$$

Let  $f$  be any classifier. Its DD, DO and PD can be expressed in turn as

$$\begin{aligned}
\text{DD}(f) &= \mathbb{P}(\widehat{Y}_f = 1 | A = 1) - \mathbb{P}(\widehat{Y}_f = 1 | A = 0) \\
&= \int_{\mathcal{X}} f(x) d\mathbb{P}_{X|A=1,Y=1}(x) - \int_{\mathcal{X}} f(x) d\mathbb{P}_{X|A=0,Y=1}(x) \\
&= \int_{\mathcal{X}} \left[ f(x) \left( \frac{d\mathbb{P}_{X|A=1}(x)}{d\mathbb{P}_X(x)} - \frac{d\mathbb{P}_{X|A=0}(x)}{d\mathbb{P}_X(x)} \right) \right] d\mathbb{P}_X(x) \\
&= \int_{\mathcal{X}} f(x, a) \left( \frac{\eta^A(x)}{p_1} - \frac{1-\eta^A(x)}{p_0} \right) d\mathbb{P}_X(x); \\
\text{DO}(f) &= \mathbb{P}(\widehat{Y}_f = 1 | A = 1, Y = 1) - \mathbb{P}(\widehat{Y}_f = 1 | A = 0, Y = 1) \\
&= \int_{\mathcal{X}} f(x) d\mathbb{P}_{X|A=1}(x) - \int_{\mathcal{X}} f(x) d\mathbb{P}_{X|A=0}(x) \\
&= \int_{\mathcal{X}} \left[ f(x) \left( \frac{d\mathbb{P}_{X|A=1,Y=1}(x)}{d\mathbb{P}_X(x)} - \frac{d\mathbb{P}_{X|A=0,Y=1}(x)}{d\mathbb{P}_X(x)} \right) \right] d\mathbb{P}_X(x) \\
&= \int_{\mathcal{X}} f(x, a) \left( \frac{\eta_1(x)\eta^A(x)}{p_{1,1}} - \frac{\eta_0(x)(1-\eta^A(x))}{p_{0,1}} \right) d\mathbb{P}_X(x); \\
\text{PD}(f) &= \mathbb{P}(\widehat{Y}_f = 1 | A = 1, Y = 0) - \mathbb{P}(\widehat{Y}_f = 1 | A = 0, Y = 0) \\
&= \int_{\mathcal{X}} f(x) d\mathbb{P}_{X|A=1,Y=0}(x) - \int_{\mathcal{X}} f(x) d\mathbb{P}_{X|A=0,Y=0}(x) \\
&= \int_{\mathcal{X}} \left[ f(x) \left( \frac{d\mathbb{P}_{X|A=1,Y=0}(x)}{d\mathbb{P}_X(x)} - \frac{d\mathbb{P}_{X|A=0,Y=0}(x)}{d\mathbb{P}_X(x)} \right) \right] d\mathbb{P}_X(x) \\
&= \int_{\mathcal{X}} f(x, a) \left( \frac{(1-\eta_1(x))\eta^A(x)}{p_{1,0}} - \frac{(1-\eta_0(x))(1-\eta^A(x))}{p_{0,0}} \right) d\mathbb{P}_X(x).
\end{aligned}$$

Thus, Proposition 4.10 follows.

## A.4 Proofs of Results from Section 5

### A.4.1 Proof of Theorem 5.1

By Lemma A.2, we only need to prove that, for  $t \in [t_0, \bar{t}_0]$ , the unconstrained Bayes-optimal classifier for  $\widetilde{\mathbb{P}}^t$  is the same as  $f_{\text{Dis},t}$ . By Bayes' theorem, for all  $x, a$ ,

$$\widetilde{\eta}_a^t(x) = \frac{\widetilde{p}_{a,1}^t d\widetilde{\mathbb{P}}_{\widetilde{X}|\widetilde{A}=a,\widetilde{Y}=1}^t(x)}{\widetilde{p}_{a,1}^t d\widetilde{\mathbb{P}}_{\widetilde{X}|\widetilde{A}=a,\widetilde{Y}=1}^t(x) + \widetilde{p}_{a,0}^t d\widetilde{\mathbb{P}}_{\widetilde{X}|\widetilde{A}=a,\widetilde{Y}=0}^t(x)}$$

Then, the unconstrained Bayes-optimal classifier for  $\tilde{\mathbb{P}}^t$  is

$$\begin{aligned}
f_t^{\text{FUDS}}(x, a) &= I\left(\tilde{\eta}_a^t(x) > \frac{1}{2}\right) = I\left(\frac{\tilde{p}_{a,1}^t d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=a, \tilde{Y}=1}^t(x)}{\tilde{p}_{a,1}^t d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=a, \tilde{Y}=1}^t(x) + \tilde{p}_{a,0}^t d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=a, \tilde{Y}=0}^t(x)} > \frac{1}{2}\right) \\
&= I\left(\frac{c_a(1 - H_{\text{Dis},a}(t))p_{a,y}d\mathbb{P}_{X|A=a, Y=1}(x)}{c_a(1 - H_{\text{Dis},a}(t))p_{a,y}d\mathbb{P}_{X|A=a, Y=1}(x) + c_a H_{\text{Dis},a}(t)p_{a,y}d\mathbb{P}_{X|A=a, Y=0}(x)} > \frac{1}{2}\right) \\
&= I\left((1 - H_{\text{Dis},a}(t))p_{a,1}d\mathbb{P}_{X|A=a, Y=1}(x) > H_{\text{Dis},a}(t)p_{a,0}d\mathbb{P}_{X|A=a, Y=0}(x)\right) \\
&= I\left(p_{a,y}d\mathbb{P}_{X|A=a, Y=1}(x) > H_{\text{Dis},a}(t)\left[p_{a,1}d\mathbb{P}_{X|A=a, Y=1}(x) + p_{a,0}d\mathbb{P}_{X|A=a, Y=0}(x)\right]\right) \\
&= I\left(\eta_a(x) > H_{\text{Dis},a}(t)\right) = I\left(\eta_a(x) > \frac{1}{2} + \frac{t}{2}w_{\text{Dis}}(x, a)\right) = f_{\text{Dis},t}(x, a).
\end{aligned}$$

This finishes the proof.

#### A.4.2 Proof of Theorem 5.3

Again, we only need to prove that for  $t \in [\underline{t}_0, \bar{t}_0]$ ,  $f_t^{\text{FCSC}} = f_{\text{Dis},t}$ . By definition,

$$\begin{aligned}
R_t^{\text{FCSC}}(f) &= \sum_{a \in \{0,1\}} \left[ H_{\text{Dis},a}(t) \cdot \mathbb{P}(\hat{Y}_f = 1, Y = 0, A = a) + (1 - H_{\text{Dis},a}(t)) \cdot \mathbb{P}(\hat{Y}_f = 0, Y = 1, A = a) \right] \\
&= \sum_{a \in \{0,1\}} p_a \left[ H_{\text{Dis},a}(t) \cdot \mathbb{P}(\hat{Y}_f = 1, Y = 0 | A = a) + (1 - H_{\text{Dis},a}(t)) \cdot \mathbb{P}(\hat{Y}_f = 0, Y = 1 | A = a) \right] \\
&= \sum_{a \in \{0,1\}} p_a H_{\text{Dis},a}(t) \cdot \int_{\mathcal{X}} \mathbb{P}(\hat{Y}_f = 1, Y = 0 | A = a, X = x) d\mathbb{P}_{X|A=a}(x) \\
&\quad + \sum_{a \in \{0,1\}} p_a (1 - H_{\text{Dis},a}(t)) \cdot \int_{\mathcal{X}} \mathbb{P}(\hat{Y}_f = 0, Y = 1 | X = x, A = a) d\mathbb{P}_{X|A=a}(x) \\
&= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} [H_{\text{Dis},a}(t)f(x, a)(1 - \eta_a(x)) + (1 - H_{\text{Dis},a}(t))(1 - f(x, a))\eta_a(x)] d\mathbb{P}_{X|A=a}(x) \\
&= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} [H_{\text{Dis},a}(t)f(x, a) - \eta_a(x) - H_{\text{Dis},a}(t)\eta_a(x) - \eta_a(x)f(x, a)] d\mathbb{P}_{X|A=a} \\
&= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} (H_{\text{Dis},a}(t) - \eta_a) f(x, a) d\mathbb{P}_{X|A=a} - \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} (1 + H_{\text{Dis},a}(t)) \eta_a(x) d\mathbb{P}_{X|A=a}.
\end{aligned}$$

Note that the second term does not depend on  $f$  and that the first term is minimized by taking  $f(x, a) = 1$  if  $\eta_a(x) > H_{\text{Dis},a}(t)$  and  $f(x, a) = 0$  if  $\eta_a \leq H_{\text{Dis},a}(t)$ . We can thus conclude that for all  $x, a$ ,

$$f_t^{\text{FCSC}}(x, a) = I(\eta_a(x) > H_{\text{Dis},a}(t)) = f_{\text{Dis},t}(x, a).$$

This finishes the proof.

## B Extensions

In this section, we discuss some interesting extensions of our theoretical and methodological framework, including (1) fair Bayes-optimal classifier with equalized odds; (2) fair Bayes-optimal classifier with multi-class protected attribute; (3) fair Bayes-optimal classifier with no distributional assumptions; (4) FUDS and FCSC algorithms with linear disparities (protected attribute  $A$  is excluded from predictive attribute.)

## B.1 Fair Bayes-optimal Classifier with Equalized Odds

In this section, we provide the detailed characterization of the fair Bayes-optimal classifier under equalized odds (EO), as defined in Theorem 4.7. Recall that, for any  $\delta > 0$ , a  $\delta$ -fair Bayes-optimal classifier is defined as

$$f_{\text{DEO},\delta}^* = \arg \min \{R(f) : \max\{|\text{DO}(f)|, |\text{PD}(f)|\} \leq \delta\},$$

with DO and PD defined in (2.1) and written more explicitly as

$$\text{DO}(f) = \int_{\mathcal{X} \times \mathcal{A}} f(x, a) \frac{(2a-1)\eta_a(x)}{p_{a,1}} d\mathbb{P}_{X,A}(x, a); \quad (\text{B.1})$$

$$\text{PD}(f) = \int_{\mathcal{X} \times \mathcal{A}} f(x, a) \frac{(2a-1)(1-\eta_a(x))}{p_{a,0}} d\mathbb{P}_{X,A}(x, a). \quad (\text{B.2})$$

The fairness constraint  $\max\{|\text{DO}(f)|, |\text{PD}(f)|\} \leq \delta$  consists of four linear inequalities on  $f$ :

$$\text{DO}(f) \leq \delta, \quad -\text{DO}(f) \leq \delta, \quad \text{PD}(f) \leq \delta \quad \text{and} \quad -\text{PD}(f) \leq \delta.$$

or

$$\begin{aligned} \int_{\mathcal{X} \times \mathcal{A}} f(x, a) \frac{(2a-1)\eta_a(x)}{p_{a,1}} d\mathbb{P}_{X,A}(x, a) &\leq \delta; & \int_{\mathcal{X} \times \mathcal{A}} f(x, a) \frac{(1-2a)\eta_a(x)}{p_{a,1}} d\mathbb{P}_{X,A}(x, a) &\leq \delta; \\ \int_{\mathcal{X} \times \mathcal{A}} f(x, a) \frac{(2a-1)(1-\eta_a(x))}{p_{a,0}} d\mathbb{P}_{X,A}(x, a) &\leq \delta; & \int_{\mathcal{X} \times \mathcal{A}} f(x, a) \frac{(1-2a)(1-\eta_a(x))}{p_{a,0}} d\mathbb{P}_{X,A}(x, a) &\leq \delta. \end{aligned}$$

Then, the generalized Neyman-Pearson lemma motivates us to consider classifiers of the form, for  $(c_1, c_2, c_3, c_4) \in \mathbb{R}^4$ ,  $x \in \mathcal{X}$  and  $a \in \{0, 1\}$ ,

$$f_{c_1, c_2, c_3, c_4}(x, a) = I \left( 2\eta_a(x) - 1 > c_1 \frac{(2a-1)\eta_a(x)}{p_{a,1}} - c_2 \frac{(2a-1)\eta_a(x)}{p_{a,1}} + c_3 \frac{(2a-1)(1-\eta_a(x))}{p_{a,0}} - c_4 \frac{(2a-1)(1-\eta_a(x))}{p_{a,0}} \right). \quad (\text{B.3})$$

Let  $\mathcal{T} = [-p_{0,1}, p_{1,1}] \times [-p_{1,0}, p_{0,0}] \setminus \{(-p_{0,1}, p_{0,0}), (-p_{1,0}, p_{1,1})\}$ . Take  $t_1 = c_1 - c_2 \in \mathbb{R}$  and  $t_2 = c_3 - c_4 \in \mathbb{R}$  and define, for  $a \in \{0, 1\}$ ,  $T_a : \mathcal{T} \rightarrow [0, 1]$  for all  $(t_1, t_2) \in \mathcal{T}$  by

$$T_a(t_1, t_2) = \frac{p_{a,1}p_{a,0} + t_2p_{a,1}}{2p_{a,1}p_{a,0} + (2a-1)(t_2p_{a,1} - t_1p_{a,0})}. \quad (\text{B.4})$$

Then, (B.3) can be simplified to, for all  $x, a$ ,

$$f_{\text{Dis}, t_1, t_2}(x, a) = I \left( 2\eta_a(x) - 1 > t_1 \frac{(2a-1)\eta_a(x)}{p_{a,1}} + t_2 \frac{(2a-1)(1-\eta_a(x))}{p_{a,0}} \right) = I(\eta_a(x) > T_a(t_1, t_2)). \quad (\text{B.5})$$

We then define the disparity functions  $D_{\text{DO}}: \mathcal{T} \rightarrow [-1, 1]$  and  $D_{\text{PD}}: \mathcal{T} \rightarrow [-1, 1]$  to measure the DO and PD of  $f_{\text{Dis}, t_1, t_2}$ , for all  $t_1, t_2 \in \mathcal{T}$ , as

$$D_{\text{DO}}(t_1, t_2) = \text{DO}(f_{\text{Dis}, t_1, t_2}) = \int_{\mathcal{X} \times \mathcal{A}} I(\eta_a(x) > T_a(t_1, t_2)) \frac{(1-2a)\eta_a(x)}{p_{a,1}} d\mathbb{P}_{X,A}(x, a); \quad (\text{B.6})$$

$$D_{\text{PD}}(t_1, t_2) = \text{PD}(f_{\text{Dis}, t_1, t_2}) = \int_{\mathcal{X} \times \mathcal{A}} I(\eta_a(x) > T_a(t_1, t_2)) \frac{(2a-1)(1-\eta_a(x))}{p_{a,0}} d\mathbb{P}_{X,A}(x, a). \quad (\text{B.7})$$

**Proposition B.1** (Continuity and Monotonicity of DO and PD).

(1) For fixed  $t_2 \in (-p_{1,0}, p_{0,0})$ , both  $t \mapsto D_{\text{DO}}(t, t_2)$  and  $t \mapsto D_{\text{PD}}(t, t_2)$  are continuous and monotone non-increasing. Moreover,  $\text{DO}(-p_{0,1}, t_2) \geq 0$ ,  $\text{DO}(p_{1,1}, t_2) \leq 0$ ,  $\text{PD}(-p_{0,1}, t_2) \geq 0$  and  $\text{PD}(p_{1,1}, t_2) \leq 0$ .

(2) For fixed  $t_1 \in (-p_{0,1}, p_{1,1})$ , both  $t \mapsto D_{\text{DO}}(t_1, t)$  and  $t \mapsto D_{\text{PD}}(t_1, t)$  are continuous and monotone non-increasing. Moreover  $D_{\text{DO}}(t_1, -p_{1,0}) \geq 0$ ,  $D_{\text{DO}}(t_1, p_{0,0}) \leq 0$ ,  $D_{\text{PD}}(t_1, -p_{1,0}) \geq 0$  and  $D_{\text{PD}}(t_1, p_{0,0}) \leq 0$ .

Next, we define the following quantities, which are useful in determining the optimal thresholds  $t_1, t_2$  for the  $\delta$ -fair Bayes-optimal classifier. For any  $\delta > 0$ , define:  $\acute{t}_{\text{DO}} : \mathbb{R}_+ \rightarrow [-p_{0,1}, p_{1,1}]$  and  $\acute{t}_{\text{PD}} : \mathbb{R}_+ \rightarrow [-p_{1,0}, p_{0,0}]$  by

$$\acute{t}_{\text{DO}}(\delta) = \arg \min_t \{|t|, |D_{\text{DO}}(t, 0)| \leq \delta\}, \quad \acute{t}_{\text{PD}}(\delta) = \arg \min_t \{|t|, |\text{PD}(0, t)| \leq \delta\}. \quad (\text{B.8})$$

By Proposition B.1 with  $t_1 = t_2 = 0$ , these quantities are well-defined.

Now, we define  $t_{\text{DEO},1} : [0, \infty) \rightarrow \mathbb{R}$  and  $t_{\text{DEO},2} : [0, \infty) \rightarrow \mathbb{R}$ , the thresholds of the  $\delta$ -fair Bayes-optimal classifiers in Theorem 4.7, for any  $\delta \geq 0$ , as follows; where the claim that they are well-defined is proved in Lemma B.4:

$$(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) \quad (\text{B.9})$$

$$= \begin{cases} (0, 0), & (1) |D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta))| \leq \delta \text{ and } |D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0)| \leq \delta; \\ (\acute{t}_{\text{DO}}(\delta), 0), & (2) |D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta))| > \delta \text{ and } |D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0)| \leq \delta; \\ (0, \acute{t}_{\text{PD}}(\delta)), & (3) |D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta))| \leq \delta \text{ and } |D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0)| > \delta; \\ \text{solve}\{(D_{\text{DO}}(t_1, t_2), D_{\text{PD}}(t_1, t_2)) = (\delta, \delta)\}, & (4) D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta)) > \delta \text{ and } D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0) > \delta; \\ \text{solve}\{(D_{\text{DO}}(t_1, t_2), D_{\text{PD}}(t_1, t_2)) = (\delta, -\delta)\}, & (5) D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta)) > \delta \text{ and } D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0) < -\delta; \\ \text{solve}\{(D_{\text{DO}}(t_1, t_2), D_{\text{PD}}(t_1, t_2)) = (-\delta, \delta)\}, & (6) D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta)) < -\delta \text{ and } D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0) > \delta; \\ \text{solve}\{(D_{\text{DO}}(t_1, t_2), D_{\text{PD}}(t_1, t_2)) = (-\delta, -\delta)\}, & (7) D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta)) < -\delta \text{ and } D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0) < -\delta. \end{cases}$$

The seven cases above are mutually exclusive and collectively exhaustive. The first three cases, where at least one of  $t_{\text{DEO},1}$  and  $t_{\text{DEO},2}$  equals zero, correspond to (1) the unconstrained Bayes-optimal classifier, (2) the  $\delta$ -fair Bayes-optimal classifier under equality of opportunity, and (3) the  $\delta$ -fair Bayes-optimal classifier under predictive equality. These satisfy the  $\delta$ -parity constraint for equalized odds. For the remaining four cases, both  $t_{\text{DEO},1}$  and  $t_{\text{DEO},2}$  are carefully selected to satisfy the hard constraint that for  $a \in \{\text{DO}, \text{PD}\}$ ,  $D_a(t_{\text{DEO},1}, t_{\text{DEO},2})$  equals either  $\delta$  or  $-\delta$ .

### B.1.1 Proof of Proposition B.1

We prove claim (1), and claim (2) follows by similar arguments. For any  $0 \leq c_1 \leq c_2$ , we have that  $x \mapsto c_1/(c_2 + x)$  is monotone non-increasing on  $(-c_2, \infty)$  and  $x \mapsto c_1/(c_2 - x)$  is monotone non-decreasing on  $(-\infty, c_2)$ . Let, for  $a \in \{0, 1\}$ ,  $0 \leq c_{1,a} = p_{a,1}p_{a,0} + t_2p_{a,1} < 2p_{a,1}p_{a,0} + t_2p_{a,1} = c_{2,a}$ . Fixing  $t_2 \in (-p_{1,0}, p_{0,0})$ , on one hand, we have for  $-p_{0,1} \leq t_{1,1} \leq t_{1,2} \leq p_{1,1}$  that  $t_{1,1}p_{1,0} \leq t_{1,2}p_{1,0} \leq p_{1,1}p_{1,0} < 2p_{1,1}p_{1,0} + t_2p_{1,1} = c_{2,1}$ . Thus, since  $t_{1,1} \leq t_{1,2}$ ,

$$T_1(t_{1,1}, t_2) = \frac{p_{1,1}p_{1,0} + t_2p_{1,1}}{2p_{1,1}p_{1,0} + t_2p_{1,1} - t_{1,1}p_{1,0}} \leq \frac{p_{1,1}p_{1,0} + t_2p_{1,1}}{2p_{1,1}p_{1,0} + t_2p_{1,1} - t_{1,2}p_{1,0}} \leq T_1(t_{1,2}, t_2).$$

On the other hand,  $t_{1,2}p_{0,0} \geq t_{1,1}p_{0,0} \geq -p_{0,1}p_{0,0} > t_2p_{0,1} - 2p_{0,1}p_{0,0} = -c_{2,0}$ . Thus

$$T_0(t_{1,1}, t_2) = \frac{p_{0,1}p_{0,0} - t_2p_{0,1}}{2p_{0,1}p_{0,0} - t_2p_{0,1} + t_{1,1}p_{0,0}} \geq \frac{p_{0,1}p_{0,0} - t_2p_{0,1}}{2p_{0,1}p_{0,0} - t_2p_{0,1} + t_{1,2}p_{0,0}} = T_0(t_{1,2}, t_2).$$

Then, by definition, we have

$$\begin{aligned} D_{\text{DO}}(t_{1,1}, t_2) &= \mathbb{P}_{A=1, Y=1}(\eta_1(X) > T_1(t_{1,1}, t_2)) - \mathbb{P}_{A=0, Y=1}(\eta_0(X) > T_0(t_{1,1}, t_2)) \\ &\geq \mathbb{P}_{A=1, Y=1}(\eta_1(X) > T_1(t_{1,2}, t_2)) - \mathbb{P}_{A=0, Y=1}(\eta_0(X) > T_0(t_{1,2}, t_2)) = D_{\text{DO}}(t_{1,1}, t_2)(t_{1,2}, t_2); \end{aligned}$$

and

$$\begin{aligned} D_{\text{PD}}(t_{1,1}, t_2) &= \mathbb{P}_{A=1, Y=0}(\eta_1(X) > T_1(t_{1,1}, t_2)) - \mathbb{P}_{A=0, Y=0}(\eta_0(X) > T_0(t_{1,1}, t_2)) \\ &\geq \mathbb{P}_{A=1, Y=0}(\eta_1(X) > T_1(t_{1,2}, t_2)) - \mathbb{P}_{A=0, Y=0}(\eta_0(X) > T_0(t_{1,2}, t_2)) = D_{\text{PD}}(t_{1,1}, t_2)(t_{1,2}, t_2). \end{aligned}$$

Thus, for fixed  $t_2 \in (-p_{1,0}, p_{0,0})$ , both  $t \mapsto D_{\text{DO}}(t, t_2)$  and  $t \mapsto D_{\text{PD}}(t, t_2)$  are monotone non-increasing on  $[-p_{0,1}, p_{1,1}]$ . Continuity follows since, for  $a \in \{0, 1\}$ ,  $t \mapsto T_a(t, t_2)$  is continuous and  $\eta_a(X)$  has density function on  $\mathcal{X}$ .

Moreover, since for any fixed  $t_2 \in (-p_{1,0}, p_{0,0})$ ,  $T_1(p_{1,1}, t_2) = T_0(-p_{0,1}) = 1$ , we have,

$$\begin{aligned}
& \text{DO}(-p_{0,1}, t_2) = \mathbb{P}_{A=1, Y=1}(\eta_1(X) > T_1(-p_{0,1}, t_2)) - \mathbb{P}_{A=0, Y=1}(\eta_0(X) > 1) \\
& = \mathbb{P}_{A=1, Y=1}(\eta_1(X) > T_1(-p_{0,1}, t_2)) \geq 0; \\
& \text{DO}(p_{1,1}, t_2) = \mathbb{P}_{A=1, Y=1}(\eta_1(X) > 1) - \mathbb{P}_{A=0, Y=1}(\eta_0(X) > T_0(p_{1,1}, t_2)) \\
& = -\mathbb{P}_{A=0, Y=1}(\eta_0(X) > T_0(p_{1,1}, t_2)) \leq 0; \\
& \text{PD}(-p_{0,1}, t_2) = \mathbb{P}_{A=1, Y=0}(\eta_1(X) > T_1(-p_{0,1}, t_2)) - \mathbb{P}_{A=0, Y=0}(\eta_0(X) > 1) \\
& = \mathbb{P}_{A=1, Y=2-j}(\eta_1(X) > T_1(-p_{0,1}, t_2)) \geq 0; \\
& \text{PD}(p_{1,1}, t_2) = \mathbb{P}_{A=1, Y=0}(\eta_1(X) > 1) - \mathbb{P}_{A=0, Y=0}(\eta_0(X) > T_0(p_{1,1}, t_2)) \\
& = -\mathbb{P}_{A=0, Y=2-j}(\eta_0(X) > T_0(p_{1,1}, t_2)) \leq 0.
\end{aligned}$$

### B.1.2 Proof of Theorem 4.7

The following lemmas are useful in proving Theorem 4.7. The first result characterizes the relation between  $D_{\text{DO}}$  and  $D_{\text{PD}}$ .

**Lemma B.2** (Relation between  $D_{\text{DO}}$  and  $D_{\text{PD}}$ ). *Let  $t, t_1$  and  $t_2$  be three real numbers.*

- (1) *If  $t_2 \geq 0$  and  $D_{\text{DO}}(t_1, t_2) = D_{\text{DO}}(t, 0)$ , we have  $D_{\text{PD}}(t_1, t_2) \leq D_{\text{PD}}(t, 0)$ .*
- (2) *If  $t_2 \leq 0$  and  $D_{\text{DO}}(t_1, t_2) = D_{\text{DO}}(t, 0)$ , we have  $D_{\text{PD}}(t_1, t_2) \geq D_{\text{PD}}(t, 0)$ .*
- (3) *If  $t_1 \geq 0$  and  $D_{\text{PD}}(t_1, t_2) = D_{\text{PD}}(0, t)$ , we have  $D_{\text{DO}}(t_1, t_2) \leq D_{\text{DO}}(0, t)$ .*
- (4) *If  $t_1 \leq 0$  and  $D_{\text{PD}}(t_1, t_2) = D_{\text{PD}}(0, t)$ , we have  $D_{\text{DO}}(t_1, t_2) \geq D_{\text{DO}}(0, t)$ .*

*Proof of Lemma B.2.* For conciseness, we show the proofs for claim (1) Claims (2) through (4) can be verified using the same approach as for claim (1).

Recall  $f_{\text{Dis}, t_1, t_2}$  and  $f_{\text{Dis}, t, 0}$  from (B.5). By Lemma A.2 adapted to equality of opportunity,  $f_{\text{Dis}, t, 0}$  is Bayes-optimal with respect to DO, and as  $f_{\text{Dis}, t_1, t_2}$  is another classifier,

$$R(f_{\text{Dis}, t, 0}) = \min_{f \in \mathcal{F}} \left\{ R_f : \frac{t \cdot \text{DO}(f)}{|t|} \leq \frac{t \cdot \text{DO}(t, 0)}{|t|} \right\} \leq R(f_{\text{Dis}, t_1, t_2}).$$

Moreover, since  $\mathbb{P}_{X|A=a}(\eta_a(X) = T_a(t_1, t_2)) = 0$  when  $\eta_a(X)$  is a continuous random variable with respect to  $\mathbb{P}_{X|A=a}$ , for all  $f' \in \arg \min_{f \in \mathcal{F}} \{ R_f : t \cdot \text{DO}(f)/|t| \leq t \cdot \text{DO}(t, 0)/|t| \}$ ,  $f'(x, a) = f_{\text{Dis}, t, 0}(x, a)$  almost surely with respect to  $\mathbb{P}_{X, A}$ .

We have two cases: (i)  $R(f_{\text{Dis}, t, 0}) = R(f_{\text{Dis}, t_1, t_2})$  and (ii)  $R(f_{\text{Dis}, t, 0}) < R(f_{\text{Dis}, t_1, t_2})$ .

(i) When  $R(f_{\text{Dis}, t, 0}) = R(f_{\text{Dis}, t_1, t_2})$ , we have  $f_{\text{Dis}, t_1, t_2}(x, a) = f_{\text{Dis}, t, 0}(x, a)$  almost surely with respect to  $\mathbb{P}_{X, A}$ . As a result,

$$D_{\text{PD}}(t_1, t_2) - D_{\text{PD}}(t, 0) = \int_{\mathcal{X} \times \mathcal{A}} (f_{\text{Dis}, t_1, t_2}(x, a) - f_{\text{Dis}, t, 0}(x, a)) \frac{(2a-1)(1-\eta_a(x))}{p_{a,0}} d\mathbb{P}_{X, A}(x, a) = 0.$$

(ii) When  $R(f_{\text{Dis}, t, 0}) < R(f_{\text{Dis}, t_1, t_2})$ , we first notice that since  $t_2 \geq 0$ , by Lemma B.3,

$$f_{\text{Dis}, t_1, t_2} \in \arg \min_{f \in \mathcal{F}} \left\{ R(f) : \frac{t_1 \cdot \text{DO}(f)}{|t_1|} \leq \frac{t_1 \cdot D_{\text{DO}}(t_1, t_2)}{|t_1|}, \text{PD}(f) \leq D_{\text{PD}}(t_1, t_2) \right\}. \quad (\text{B.10})$$

This implies  $D_{\text{PD}}(t_1, t_2) < D_{\text{PD}}(t, 0)$ . Otherwise,  $f_{\text{Dis}, t, 0}$  satisfies  $\text{DO}(f_{\text{Dis}, t, 0}, 0) = D_{\text{DO}}(t_1, t_2)$ ,  $\text{PD}(f_{\text{Dis}, t, 0}) \leq D_{\text{PD}}(t, 0) \leq D_{\text{PD}}(t_1, t_2)$  and  $R(f_{\text{Dis}, t, 0}) < R(f_{\text{Dis}, t_1, t_2})$ , which contradicts (B.10).

In both cases, we have  $D_{\text{PD}}(t_1, t_2) \leq D_{\text{PD}}(t, 0)$ , finishing the proof.  $\square$

The next result characterizes Bayes-optimal classifiers under equalized odds.



**Lemma B.3** (Characterizing  $f_{\text{Dis},t_1,t_2}$  as Bayes-optimal Classifiers under Equalized Odds). For  $(t_1, t_2) \in \mathbb{R}^2$  and  $\delta \geq 0$ , let  $f_{\text{Dis},t_1,t_2}$ ,  $D_{\text{DO}}(t)$ ,  $D_{\text{PD}}(t)$  be defined in (B.5), (B.6) and (B.7), respectively. Then, recalling we use the convention that  $0/0 = 0$ ,

$$f_{\text{Dis},t_1,t_2} \in \arg \min_{f \in \mathcal{F}} \left\{ R(f) : \frac{t_1 \cdot \text{DO}(f)}{|t_1|} \leq \frac{t_1 \cdot D_{\text{DO}}(t_1, t_2)}{|t_1|}, \frac{t_2 \cdot \text{PD}(f)}{|t_2|} \leq \frac{t_2 \cdot D_{\text{PD}}(t_1, t_2)}{|t_2|} \right\}.$$

Moreover, for all

$$f' \in \arg \min_{f \in \mathcal{F}} \{ R(f) : t_1 \cdot \text{DO}(f)/|t_1| \leq t_1 \cdot D_{\text{DO}}(t_1, t_2)/|t_1|, t_2 \cdot \text{PD}(f)/|t_2| \leq t_2 \cdot D_{\text{PD}}(t_1, t_2)/|t_2| \},$$

we have  $f' = f_{\text{Dis},t_1,t_2}$  almost surely with respect to  $\mathbb{P}_{X,A}$ . In particular, if  $t_1 \cdot D_{\text{DO}}(t_1, t_2) \geq 0$  and  $t_2 \cdot D_{\text{PD}}(t_1, t_2) \geq 0$ ,

$$f_{\text{Dis},t_1,t_2} = \arg \min_{f \in \mathcal{F}} \{ R(f) : |\text{DO}(f)| \leq |D_{\text{DO}}(t_1, t_2)|, |\text{PD}(f)| \leq |D_{\text{PD}}(t_1, t_2)| \}.$$

*Proof of Lemma B.3.* Similar to Lemma A.2, this result is a consequence of the generalized Neyman-Pearson lemma. If either  $t_1 = 0$  or  $t_2 = 0$ , the result is an application of Lemma A.2. Now, we assume  $t_1 \neq 0$  and  $t_2 \neq 0$ . Let, for all  $x, a$ ,  $\phi_0(x, a) = 2\eta_1(x, a) - 1$ ,  $\phi_1(x, a) = (2a - 1)\eta_a(x)/p_{a,1}$  and  $\phi_2(x, a) = (2a - 1)(1 - \eta_a(x))/p_{a,0}$ . Also, let, for all classifiers  $f$ , and  $t \in \mathbb{R}$ ,  $\overline{\text{DO}}_t(f) = t \cdot \text{DO}(f)/|t|$  and  $\overline{\text{PD}}_t(f) = t \cdot \text{PD}(f)/|t|$ . We have, for  $(t_1, t_2) \in \mathbb{R}^2$  and all  $x, a$ , that with  $f_{\text{Dis},t_1,t_2}$  from (B.5),

$$f_{\text{Dis},t_1,t_2}(x, a) = I(\phi_0(x, a) > t_1\phi_1(x, a) + t_2\phi_2(x, a)) = I\left(\phi_0(x, a) > |t_1| \frac{t_1\phi_1(x, a)}{|t_1|} + |t_2| \frac{t_2\phi_2(x, a)}{|t_2|}\right).$$

Moreover, by Lemma A.1 and (B.1), we can write  $\text{Acc}(f)$ ,  $\overline{\text{DO}}_t(f)$  and  $\overline{\text{PD}}_t(f)$  as

$$\begin{aligned} \text{Acc}(f) &= \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, a) \phi_0(x, a) d\mathbb{P}_{X,A}(x, a) + \int_{\mathcal{A}} \int_{\mathcal{X}} (1 - \eta_a(x)) d\mathbb{P}_{X,A}(x, a); \\ \overline{\text{DO}}_{t_1}(f) &= \frac{t_1}{|t_1|} \text{DO}(f) = \int_{\mathcal{X} \times \mathcal{A}} f(x, a) \frac{t_1\phi_1(x, a)}{|t_1|} d\mathbb{P}_{X,A}(x, a); \\ \overline{\text{PD}}_{t_2}(f) &= \frac{t_2}{|t_2|} \text{PD}(f) = \int_{\mathcal{X} \times \mathcal{A}} f(x, a) \frac{t_2\phi_2(x, a)}{|t_2|} d\mathbb{P}_{X,A}(x, a). \end{aligned}$$

Define the sets of classifiers

$$\begin{aligned} \mathcal{F}_{t_1,t_2,=} &= \left\{ f : \overline{\text{DO}}_{t_1}(f) = \frac{t_1 \cdot D_{\text{DO}}(t_1, t_2)}{|t_1|}, \overline{\text{PD}}_{t_2}(f) = \frac{t_2 \cdot D_{\text{PD}}(t_1, t_2)}{|t_2|} \right\}; \\ \mathcal{F}_{t_1,t_2,|\cdot|,\leq} &= \left\{ f : |\overline{\text{DO}}_{t_1}(f)| \leq |D_{\text{DO}}(t_1, t_2)|, |\overline{\text{PD}}_{t_2}(f)| \leq |D_{\text{PD}}(t_1, t_2)| \right\}; \\ \mathcal{F}_{t_1,t_2,\leq} &= \left\{ f : \overline{\text{DO}}_{t_1}(f) \leq \frac{t_1 \cdot D_{\text{DO}}(t_1, t_2)}{|t_1|}, \overline{\text{PD}}_{t_2}(f) \leq \frac{t_2 \cdot D_{\text{PD}}(t_1, t_2)}{|t_2|} \right\}. \end{aligned}$$

As  $f_{\text{Dis},t_1,t_2} \in \mathcal{F}_{t_1,t_2,=}$ , by the generalized Neyman-Pearson lemma (Lemma 3.1),

$$f_{\text{Dis},t_1,t_2} \in \arg \max_{f \in \mathcal{F}_{t_1,t_2,=}} \text{Acc}(f).$$

Moreover, since  $|t_1| \geq 0$  and  $|t_2| \geq 0$ , we have

$$f_{\text{Dis},t_1,t_2} \in \arg \max_{f \in \mathcal{F}_{t_1,t_2,\leq}} \text{Acc}(f) = \arg \min_{f \in \mathcal{F}} \left\{ R(f) : \frac{t_1 \text{DO}(f)}{|t_1|} \leq \frac{t_1 D_{\text{DO}}(t_1, t_2)}{|t_1|}, \frac{t_2 \text{PD}(f)}{|t_2|} \leq \frac{t_2 D_{\text{PD}}(t_1, t_2)}{|t_2|} \right\}.$$

In addition, since  $\mathbb{P}_{X|A=a}(\eta_a(X) = T_a(t_1, t_2)) = 0$  when  $\eta_a(X)$  is a continuous random variable with respect to  $\mathbb{P}_{X|A=a}$ , for all  $f' \in \arg \max_{f \in \mathcal{F}_{t_1,t_2,\leq}} \text{Acc}(f)$ ,  $f' = f_{\text{Dis},t_1,t_2}$  almost surely with respect to  $\mathbb{P}_{X,A}$ . Furthermore, if  $t_1 \cdot D_{\text{DO}}(t_1, t_2) \geq 0$  and  $t_2 \cdot D_{\text{PD}}(t_1, t_2) \geq 0$ , we have  $f_{\text{Dis},t_1,t_2} \in \mathcal{F}_{t_1,t_2,|\cdot|,\leq} \subset \mathcal{F}_{t_1,t_2,=}$ . Thus

$$f_{\text{Dis},t_1,t_2} = \arg \min_{f \in \mathcal{F}} \{ R(f) : |\text{DO}(f)| \leq |D_{\text{DO}}(t_1, t_2)|, |\text{PD}(f)| \leq |D_{\text{PD}}(t_1, t_2)| \}.$$

This finishes the proof.  $\square$

Next, we show that  $t_{\text{DEO},1}(\delta)$  and  $t_{\text{DEO},2}(\delta)$  are well defined in cases (4) to (7). For conciseness, we consider case (4), and the arguments for the other cases are similar.

**Lemma B.4** (The Quantities  $t_{\text{DEO},1}(\delta)$  and  $t_{\text{DEO},2}(\delta)$  Are Well-Defined). *Let  $\acute{t}_{\text{DO}}(\delta)$  and  $\acute{t}_{\text{PD}}(\delta)$  be defined in (B.8), Suppose that, for  $a \in \{0,1\}$ ,  $\eta_a(X)$  has a density function on  $\mathcal{X}$ . Then, for fixed  $\delta \geq 0$ , when  $D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta)) > \delta$  and  $D_{\text{PD}}(0, \acute{t}_{\text{DO}}(\delta)) > \delta$ , we have  $\acute{t}_{\text{DO}}(\delta) > 0$ ,  $\acute{t}_{\text{PD}}(\delta) > 0$ , and there exist  $(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) \in [0, \acute{t}_{\text{DO}}(\delta)] \times [0, \acute{t}_{\text{PD}}(\delta)]$  such that  $(D_{\text{DO}}(t_1, t_2), D_{\text{PD}}(t_1, t_2)) = (\delta, \delta)$ .*

*Proof of Lemma B.4.* If  $\acute{t}_{\text{DO}}(\delta) \leq 0$ , then by the definition of  $\acute{t}_{\text{DO}}(\delta)$  in (B.8), we have  $D_{\text{DO}}(0, 0) \leq \delta$  and  $\text{DO}(\acute{t}_{\text{DO}}(\delta), 0) \leq \delta$ . Moreover, by Proposition B.1,  $D_{\text{PD}}(0, 0) \geq D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0) > \delta$ . Thus, it follows that  $\acute{t}_{\text{PD}}(\delta) > 0$ . By applying Proposition B.1 again,  $D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta)) \leq D_{\text{DO}}(0, 0) \leq \delta$ , which is a contradiction. As a result, we have  $\acute{t}_{\text{DO}}(\delta) > 0$  and  $\text{DO}(\acute{t}_{\text{DO}}(\delta), 0) = \delta$ . On the other hand, since  $D_{\text{DO}}(0, p_{0,0}) \leq \delta$ , we have  $\acute{t}_{\text{PD}}(\delta) < p_{0,0}$ . Similarly, we can show  $0 < \acute{t}_{\text{DO}}(\delta) < p_{1,1}$  and  $D_{\text{PD}}(0, \acute{t}_{\text{PD}}(\delta)) = \delta$ .

Now, for any  $t_2 \in (0, \acute{t}_{\text{PD}}(\delta))$ , we have  $D_{\text{DO}}(0, t_2) \geq D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta)) > \delta$  and  $\text{DO}(p_{1,1}, t_2) \leq \delta$ . By the continuity of  $D_{\text{DO}}(t, t_2)$  as a function of  $t$ , there exists  $t_1 \in (0, p_{1,1})$  such that  $D_{\text{DO}}(t_1, t_2) = \delta$ . We then define  $Q_\delta : [0, \acute{t}_{\text{PD}}(\delta)] \rightarrow (0, p_{1,1})$  for all  $t_2 \in (0, p_{1,1})$  as

$$Q_\delta(t_2) = \inf_{t \in (0, p_{1,1})} \{D_{\text{DO}}(t, t_2) < \delta\}.$$

Clearly, we have  $Q_\delta(0) \leq \acute{t}_{\text{DO}}(\delta)$  and, since  $D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta)) > \delta$ , we have  $Q_\delta(\acute{t}_{\text{PD}}(\delta)) > 0$ .

Now, we consider the function  $P$  defined on  $[0, \acute{t}_{\text{PD}}(\delta)]$   $t_2 \mapsto \mathbb{P}(t_2) = \text{PD}(Q_\delta(t_2), t_2)$ . Then we have  $\mathbb{P}(0) = D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0) > \delta$  and  $\mathbb{P}(\acute{t}_{\text{PD}}(\delta)) = \text{PD}(Q_\delta(\acute{t}_{\text{PD}}(\delta)), \acute{t}_{\text{PD}}(\delta)) \leq D_{\text{PD}}(0, \acute{t}_{\text{PD}}(\delta)) = \delta$ . Thus, by the continuity of  $D_{\text{PD}}$ , there exists  $t_{\text{DEO},2}(\delta) \in [0, \acute{t}_{\text{PD}}(\delta)]$  such that  $\text{PD}(Q_\delta(t_{\text{DEO},2}(\delta)), t_{\text{DEO},2}(\delta)) = \delta$ . Setting  $t_{\text{DEO},1}(\delta) = Q_\delta(t_{\text{DEO},2}(\delta))$ , we have  $\text{DO}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) = D_{\text{PD}}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) = \delta$ . Moreover, since  $t_{\text{DEO},2}(\delta) \leq \acute{t}_{\text{PD}}(\delta)$  and  $D_{\text{PD}}(0, \acute{t}_{\text{PD}}(\delta)) = D_{\text{PD}}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) = \delta$ , we have  $t_{\text{DEO},1}(\delta) \geq 0$ . On the other hand, since  $t_{\text{DEO},2}(\delta) \geq 0$  and  $\text{DO}(\acute{t}_{\text{DO}}(\delta), 0) = D_{\text{PD}}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) = \delta$ , we have  $t_{\text{DEO},1}(\delta) \leq \acute{t}_{\text{DO}}(\delta)$ . This finishes the proof.  $\square$

We proceed with the proof of Theorem 4.7.

*Proof of Theorem 4.7.* We demonstrate the result in cases (1), (2), (4), and (5). The results in other cases can be verified similarly: case (3) is analogous to case (2), case (6) is analogous to case (5), and case (7) is analogous to case (4).

The following facts follow directly from the definitions of  $\acute{t}_{\text{DO}}(\delta)$  and  $\acute{t}_{\text{PD}}(\delta)$  in (B.8), and Proposition B.1:

Fact (1). For  $j \in \{\text{DO}, \text{PD}\}$ ,  $\delta \mapsto \acute{t}_j(\delta)$  is monotone non-decreasing.

Fact (2). For  $j \in \{\text{DO}, \text{PD}\}$ ,  $\acute{t}_j > 0$  if and only if  $D_j(0, 0) > \delta$ . Further,  $\acute{t}_j < 0$  if and only if  $D_j(0, 0) < -\delta$ ;

Fact (3). If  $D_{\text{DO}}(0, 0) > \delta$ ,  $\text{DO}(\acute{t}_{\text{PD}}(\delta), 0) = \delta$ ; If  $D_{\text{DO}}(0, 0) < -\delta$ , then  $\text{DO}(\acute{t}_{\text{PD}}(\delta), 0) = -\delta$ . Further,  $D_{\text{PD}}(0, 0) > \delta$ , then  $D_{\text{PD}}(0, \acute{t}_{\text{PD}}(\delta)) = \delta$ . Finally, if  $D_{\text{PD}}(0, 0) < -\delta$ , then  $D_{\text{PD}}(0, \acute{t}_{\text{PD}}(\delta)) = -\delta$ ;

We now study the cases mentioned for Theorem 4.7.

Case (1).  $|D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0)| \leq \delta$  and  $|D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta))| \leq \delta$ .

In this scenario, we have  $t_{\text{DEO},1}(\delta) = t_{\text{DEO},2}(\delta) = 0$  and, for all  $x, a$ ,  $f_{\text{Dis},0,0}(x, a) = f_{\text{Dis}, t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)}(x, a) = I(\eta_a(x) > 1/2)$  is the unconstrained Bayes-optimal classifier. As a result, we only need to prove  $|D_{\text{DO}}(0, 0)| \leq \delta$  and  $|D_{\text{PD}}(0, 0)| \leq \delta$ .

If  $\acute{t}_{\text{DO}}(\delta) > 0$ , then we have  $D_{\text{DO}}(0, 0) > \delta$ . By  $|D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta))| \leq \delta$  and Proposition (B.1), we conclude  $\acute{t}_{\text{PD}}(\delta) < 0$ , which leads to  $D_{\text{PD}}(0, 0) < -\delta$ . Finally, since  $\acute{t}_{\text{DO}}(\delta) > 0$ , we see  $\text{PD}(\acute{t}_{\text{DO}}(\delta), 0) \leq D_{\text{PD}}(0, 0) < -\delta$ , which contradicts  $|D_{\text{PD}}(0, 0)| \leq \delta$ . Thus, we can conclude that  $\acute{t}_{\text{DO}}(\delta) \leq 0$ .

With analogous arguments, we can also demonstrate that  $\acute{t}_{\text{DO}}(\delta) \geq 0$ . Then, we must have  $\acute{t}_{\text{DO}}(\delta) = 0$ . As a result,  $|D_{\text{DO}}(0, 0)| \leq \delta$  and  $|D_{\text{PD}}(0, 0)| \leq \delta$ .

Case (2).  $|D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta))| > \delta$  and  $|D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0)| \leq \delta$ .

In this scenario, we have  $t_{\text{DEO},1} = \acute{t}_{\text{DO}}(\delta) = \arg \min_t \{|t| : |\text{DO}| \leq \delta\}$  and  $t_{\text{DEO},2} = 0$ . On one hand, by Theorem 4.2 adapted for equality of opportunity, we have

$$R(f_{\text{Dis}, t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)}) = R(f_{\text{Dis}, \acute{t}_{\text{DO}}(\delta), 0}) = \min_{f \in \mathcal{F}} \{R(f) : |\text{DO}(f)| \leq \delta\}.$$

On the other hand, given that  $|D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0)| \leq \delta$ , we have

$$f_{\text{Dis}, t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)} = f_{\text{Dis}, \acute{t}_{\text{DO}}(\delta), 0} \in \{f \in \mathcal{F} : \max(|\text{DO}(f)|, |\text{PD}(f)|) \leq \delta\} \subset \{f \in \mathcal{F} : \text{DO}(f) \leq \delta\}.$$

It follows that

$$R(f_{\text{Dis}, t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)}) \geq \min_{f \in \mathcal{F}} \{R(f) : \max(|\text{DO}(f)|, |\text{PD}(f)|) \leq \delta\} \geq \min_{f \in \mathcal{F}} \{R(f) : \text{DO}(f) \leq \delta\}.$$

Then, we can conclude that

$$f_{\text{Dis}, t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)} \in \arg \min_{f \in \mathcal{F}} \{R(f) : \max(|\text{DO}(f)|, |\text{PD}(f)|) \leq \delta\}.$$

Case (4)  $D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta)) > \delta$  and  $D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0) > \delta$

In this case,  $(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta))$  is the solution of the equation  $(D_{\text{DO}}(t_1, t_2), D_{\text{PD}}(t_1, t_2)) = (\delta, \delta)$ .

Following the proof of Lemma B.4, we find that  $\acute{t}_{\text{DO}}(\delta) > 0$  and  $\acute{t}_{\text{PD}}(\delta) > 0$ . This further implies  $\text{DO}(\acute{t}_{\text{DO}}(\delta), 0) = \text{PD}(0, \acute{t}_{\text{DO}}(\delta)) = \delta$ . By claims (2) and (4) of Lemma B.2, we can conclude that  $t_{\text{DEO},1}(\delta) \geq 0$  and  $t_{\text{DEO},2}(\delta) \geq 0$ . Otherwise, if  $t_{\text{DEO},2}(\delta) < 0$ , since  $D_{\text{DO}}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) = \delta = \text{DO}(\acute{t}_{\text{DO}}(\delta), 0)$ , we would have  $D_{\text{PD}}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) \geq \text{PD}(\acute{t}_{\text{DO}}(\delta), 0) > \delta$ , which contradicts  $D_{\text{PD}}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) = \delta$ . A similar argument applies to  $t_{\text{DEO},1}(\delta)$ .

Now, we have  $t_{\text{DEO},1}(\delta) \cdot D_{\text{DO}}(t_{\text{DEO},1}, t_{\text{DEO},2}) = t_{\text{DEO},1}(\delta) \cdot \delta \geq 0$  and  $t_{\text{DEO},2}(\delta) \cdot D_{\text{PD}}(t_{\text{DEO},1}, t_{\text{DEO},2}) = t_{\text{DEO},2}(\delta) \cdot \delta \geq 0$ . By Lemma B.3,

$$f_{\text{Dis}, t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)} \in \arg \min_{f \in \mathcal{F}} \{R(f) : \max(|\text{DO}(f)|, |\text{PD}(f)|) \leq \delta\}.$$

Case (5)  $D_{\text{DO}}(0, \acute{t}_{\text{PD}}(\delta)) > \delta$  and  $D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0) < -\delta$ .

In this case,  $(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta))$  is the solution of the equation  $(D_{\text{DO}}(t_1, t_2), D_{\text{PD}}(t_1, t_2)) = (\delta, -\delta)$ .

We consider two cases: (i)  $\acute{t}_{\text{DO}}(\delta) > 0$  and (ii)  $\acute{t}_{\text{DO}}(\delta) \leq 0$ .

- (i) When  $\acute{t}_{\text{DO}}(\delta) > 0$ , we have  $D_{\text{DO}}(0, 0) > \delta$  and  $\text{DO}(\acute{t}_{\text{DO}}(\delta), 0) = \delta$ . Then, from result (1) of Lemma B.2, it follows that  $t_{\text{DEO},1}(\delta) > 0$  and  $t_{\text{DEO},2}(\delta) < 0$ . Otherwise, on one hand, if  $t_{\text{DEO},2}(\delta) \geq 0$ , since  $\text{DO}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) = \delta = \text{DO}(\acute{t}_{\text{DO}}(\delta), 0)$ , we have  $\text{PD}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) \leq \text{PD}(\acute{t}_{\text{DO}}(\delta), 0) < -\delta$ , which is a contradiction. On the other hand, if  $t_{\text{DEO},1}(\delta) \leq 0$ , we have  $t_{\text{DEO},1}(\delta) \leq 0 \leq \acute{t}_{\text{DO}}(\delta)$ . Then, by the monotonicity properties of  $t \mapsto D_{\text{DO}}$  from Proposition B.1 and the fact that  $\text{DO}(\acute{t}_{\text{DO}}(\delta), 0) = \delta = \text{DO}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta))$ , we have  $t_{\text{DEO},2}(\delta) \geq 0$ . We again reach a contradiction.
- (ii) When  $\acute{t}_{\text{DO}}(\delta) \leq 0$ , the argument is similar. By Proposition B.1, we have  $D_{\text{PD}}(0, 0) \leq D_{\text{PD}}(\acute{t}_{\text{DO}}(\delta), 0) < -\delta$ , which implies  $\acute{t}_{\text{PD}}(\delta) < 0$  and  $D_{\text{PD}}(0, \acute{t}_{\text{PD}}(\delta)) = -\delta$ . Then, by claim (4) of Lemma B.2, we have  $t_{\text{DEO},1}(\delta) > 0$  and  $t_{\text{DEO},2}(\delta) < 0$ . Otherwise, on one hand, if  $t_{\text{DEO},1}(\delta) \leq 0$ , since  $\text{PD}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) = -\delta = \text{PD}(0, \acute{t}_{\text{PD}}(\delta))$ , we would have  $\text{DO}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)) \geq \text{DO}(0, \acute{t}_{\text{PD}}(\delta)) > \delta$ , which is a contradiction. On the other hand, if  $t_{\text{DEO},2}(\delta) \geq 0$ , we would have  $t_{\text{DEO},2}(\delta) \geq 0 \geq \acute{t}_{\text{DO}}(\delta)$ . Then, by the monotonicity properties of  $t \mapsto D_{\text{DO}}$  from Proposition B.1 and the fact that  $\text{PD}(0, \acute{t}_{\text{PD}}(\delta)) = -\delta = \text{PD}(t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta))$ , we conclude  $t_{\text{DEO},1}(\delta) \leq 0$ ; also a contradiction.

In both cases, we have  $t_{\text{DEO},1}(\delta) \cdot D_{\text{DO}}(t_{\text{DEO},1}, t_{\text{DEO},2}) = t_{\text{DEO},1}(\delta) \cdot \delta \geq 0$  and  $t_{\text{DEO},2}(\delta) \cdot D_{\text{PD}}(t_{\text{DEO},1}, t_{\text{DEO},2}) = t_{\text{DEO},2}(\delta) \cdot (-\delta) \geq 0$ . By Lemma B.3, we conclude that

$$f_{\text{Dis}, t_{\text{DEO},1}(\delta), t_{\text{DEO},2}(\delta)} \in \arg \min_{f \in \mathcal{F}} \{R(f) : \max(|\text{DO}(f)|, |\text{PD}(f)|) \leq \delta\}.$$

The proof of Theorem 4.7 is complete. □

## B.2 Fair Bayes-optimal Classifier with a Multi-class Protected Attribute

In this section, we consider demographic parity with a multi-class protected attribute. We assume that  $A \in \mathcal{A} = \{1, 2, \dots, |\mathcal{A}|\}$  for some integer  $|\mathcal{A}| > 2$ . Our theoretical results concern perfect fairness. Under perfect fairness, we have  $|\mathcal{A}| - 1$  equality constraints, and the fair Bayes-optimal classifier can be derived using the Neyman-Pearson lemma. However, for approximate fairness for a multi-class protected attribute, it is unknown ahead of time how many constraints become equalities, and how many are strict inequalities. A careful analysis of these two types of constraints is required, and we leave this to future work. Recall that, a classifier satisfies demographic parity if

$$\mathbb{P}(\widehat{Y}_f = 1 \mid A = a) = \mathbb{P}(\widehat{Y}_f = 1 \mid A = 1), \quad \text{for } a = 2, \dots, \mathcal{A}.$$

Similar to the expression of DD in (A.4), these constraints can be expressed as:

$$\int_{\mathcal{A}} \int_{\mathcal{X}} f(x, \tilde{a}) \frac{I(\tilde{a} = a)}{p_a} - \frac{I(\tilde{a} = 1)}{p_1} d\mathbb{P}_{X,A}(x, \tilde{a}) \quad \text{for } a = 2, \dots, \mathcal{A}.$$

Take  $\phi_0(x, \tilde{a}) = 2\eta_{\tilde{a}}(x)$  and  $\phi_a(x, \tilde{a}) = I(\tilde{a} = a)/p_a - I(\tilde{a} = 1)/p_1$  for  $a = 2, \dots, |\mathcal{A}|$ , the generalized Neyman-Pearson lemma suggests us to consider the group-wise thresholding rule, for all  $(x, \tilde{a}) \in \mathcal{X} \times \{0, 1\}$ ,

$$f(x, \tilde{a}, t_2, \dots, t_{\mathcal{A}}) = I \left( \phi_0(x, \tilde{a}) > \sum_{a=2}^{|\mathcal{A}|} t_a \phi_a(x, \tilde{a}) \right).$$

This further leads to our formal Theorem 4.8 in Section 4.3.1

### B.2.1 Proof of Theorem 4.8

We first demonstrate the existence of thresholds  $\{t_{\text{Dis},a}\}_{a=1}^{|\mathcal{A}|}$  that satisfy the condition required in Theorem 4.8.

**Lemma B.5** (Existence of Thresholds). *Suppose that, for all  $a \in \mathcal{A}$ ,  $\eta_a(X)$  has a density function on  $\mathcal{X}$ . Then, there exist  $\{t_{\text{Dis},a}\}_{a=1}^{|\mathcal{A}|}$  such that  $\sum_{a=1}^{|\mathcal{A}|} t_{\text{Dis},a} = 0$ , and for all  $a \in \{2, \dots, |\mathcal{A}|\}$ , (4.8) holds.*

*Proof of Lemma B.5.* Define the functions  $\underline{Q}_a, \overline{Q}_a : [0, 1] \rightarrow \mathbb{R}$ , such that for  $s \in [0, 1]$ ,

$$\begin{aligned} \underline{Q}_a(s) &= \sup \left\{ t : \mathbb{P}_{X|A=a} \left( \eta_a(X) > \frac{1}{2} + \frac{t}{p_a} \right) > s \right\}; \\ \overline{Q}_a(s) &= \sup \left\{ t : \mathbb{P}_{X|A=a} \left( \eta_a(X) > \frac{1}{2} + \frac{t}{p_a} \right) \geq s \right\}. \end{aligned}$$

By definition, we have the following:

- Both  $\underline{Q}_a$  and  $\overline{Q}_a$  are strictly monotonically decreasing, as  $\eta_a(X)$  has a density function;
- $\underline{Q}_a$  is right-continuous and  $\overline{Q}_a$  is left-continuous;
- For all  $s_0 \in [0, 1]$ ,  $\lim_{s \rightarrow s_0^-} \overline{Q}_a(s) = \lim_{s \rightarrow s_0^-} \underline{Q}_a(s)$  and  $\lim_{s \rightarrow s_0^+} \underline{Q}_a(s) = \lim_{s \rightarrow s_0^+} \overline{Q}_a(s)$  with  $\underline{Q}_a(s_0) \leq \overline{Q}_a(s_0)$ ;
- For all  $t \in [\underline{Q}_a(s), \overline{Q}_a(s)]$ ,  $\mathbb{P}_{X|A=a}(\eta(X) > 1/2 + \frac{t}{2p_a}) = s$ .

Now, we set

$$s^* = \sup \left\{ s : \sum_{a=1}^{|\mathcal{A}|} \underline{Q}_a(s) > 0 \right\} = \sup \left\{ s : \sum_{a=1}^{|\mathcal{A}|} \overline{Q}_a(s) > 0 \right\}.$$

By the right-continuity of  $\sum_{a=1}^{|\mathcal{A}|} \underline{Q}_a(s)$  and the left-continuity of  $\sum_{a=1}^{|\mathcal{A}|} \overline{Q}_a(s)$ , we have

$$\sum_{a=1}^{|\mathcal{A}|} \underline{Q}_a(s^*) \leq 0 \leq \sum_{a=1}^{|\mathcal{A}|} \overline{Q}_a(s^*).$$

Letting

$$t_{\text{Dis},a} = \frac{\sum_{a=1}^{|\mathcal{A}|} \bar{Q}_a(s^*)}{\sum_{a=1}^{|\mathcal{A}|} \bar{Q}_a(s^*) - \sum_{a=1}^{|\mathcal{A}|} \underline{Q}_a(s^*)} \underline{Q}_a(s^*) - \frac{\sum_{a=1}^{|\mathcal{A}|} \underline{Q}_a(s^*)}{\sum_{a=1}^{|\mathcal{A}|} \bar{Q}_a(s^*) - \sum_{a=1}^{|\mathcal{A}|} \underline{Q}_a(s^*)} \bar{Q}_a(s^*),$$

we have  $\sum_{a=1}^{|\mathcal{A}|} t_{\text{Dis},a} = 0$ . Moreover, as  $t_{\text{Dis},a} \in [\underline{Q}_a(s), \bar{Q}_a(s)]$  for all  $a \in \mathcal{A}$ , we have, for all  $a \in \mathcal{A}$ ,

$$\mathbb{P}_{X|A=a} \left( \eta_a(X) > 1/2 + \frac{t_{\text{Dis},a}}{2p_a} \right) = s^*.$$

This completes the proof.  $\square$

We now proceed to the proof of Theorem 4.8.

*Proof of Theorem 4.8.* Recall that, for  $(x, \tilde{a}) \in \mathcal{X} \times \{0, 1\}$ , we denote, for  $a = 2, \dots, |\mathcal{A}|$ ,  $\phi_0(x, \tilde{a}) = 2\eta_{\tilde{a}}(x) - 1$ ; and  $\phi_a(x, \tilde{a}) = I(\tilde{a} = a)/p_a - I(\tilde{a} = 1)/p_1$ . Since  $t_{\text{Dis},1} = -\sum_{a=2}^{|\mathcal{A}|} t_{\text{Dis},a}$ , We have, for all  $x, \tilde{a}$ ,

$$\begin{aligned} f_{t_{\text{Dis},1}, \dots, t_{\text{Dis},|\mathcal{A}|}}^*(x, \tilde{a}) &= I \left( \eta_{\tilde{a}}(x) > \frac{1}{2} + \frac{t_{\text{Dis},\tilde{a}}}{2p_{\tilde{a}}} \right) = I \left( 2\eta_{\tilde{a}}(x) - 1 > \frac{t_{\tilde{a}}}{p_{\tilde{a}}} \right) \\ &= I \left( 2\eta_{\tilde{a}}(x) - 1 > \sum_{a=2}^{|\mathcal{A}|} t_a \left( \frac{I(\tilde{a} = a)}{p_a} - \frac{I(\tilde{a} = 1)}{p_1} \right) \right) = I \left( \phi_0(x, \tilde{a}) > \sum_{t=2}^{|\mathcal{A}|} t_a \phi_a(x, \tilde{a}) \right). \end{aligned}$$

Moreover, for any classifier  $f$ , we can write its misclassification rate and corresponding disparity measures ( $\mathbb{P}_{X|A=a}(\hat{Y}_f = 1) - \mathbb{P}_{X|A=1}(\hat{Y}_f = 1)$ ) as

$$R(f) = \mathbb{P}(\hat{Y}_f \neq Y) = - \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, \tilde{a}) \phi_0(x, \tilde{a}) d\mathbb{P}_{X,A}(x, \tilde{a}) + \int_{\mathcal{A}} \int_{\mathcal{X}} \eta_{\tilde{a}}(x) d\mathbb{P}_{X,A}(x, \tilde{a})$$

and, for  $a = 2, \dots, |\mathcal{A}|$ ,

$$\mathbb{P}_{X|A=a}(\hat{Y}_f = 1) - \mathbb{P}_{X|A=1}(\hat{Y}_f = 1) = \int_{\mathcal{A}} \int_{\mathcal{X}} \phi_a(x, \tilde{a}) f(x, \tilde{a}) d\mathbb{P}_{X,A}(x, \tilde{a}). \quad (\text{B.11})$$

Denote

$$\mathcal{F}_{D,=} = \left\{ f : \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, \tilde{a}) \left( \frac{I(\tilde{a} = a)}{p_a} - \frac{I(\tilde{a} = 1)}{p_1} \right) d\mathbb{P}_{X,A}(x, \tilde{a}) = 0, \quad a = 2, 3, \dots, |\mathcal{A}| \right\},$$

By the construction of  $\{t_{\text{Dis},a}\}_{a=1}^{|\mathcal{A}|}$ , we have  $f_{t_{\text{Dis},1}, \dots, t_{\text{Dis},|\mathcal{A}|}}^* \in \mathcal{F}_{D,=}$ . Then, by the generalized Neyman Pearson lemma,

$$f_{t_{\text{Dis},1}, \dots, t_{\text{Dis},|\mathcal{A}|}}^* \in \arg \max_{f \in \mathcal{F}_{D,=}} \left\{ \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, \tilde{a}) \phi_0(x, \tilde{a}) d\mathbb{P}_{X,A}(x, \tilde{a}) \right\} = \arg \min_{f \in \mathcal{F}_{D,=}} R(f),$$

which completes the proof.  $\square$

### B.3 FUDS, FCSC and FPIR Algorithms for a Linear—But Not Bilinear—Disparity Measure

In this section, we illustrate the applicability of our proposed methods to scenarios with linear—rather than bilinear—constraints, through detailed case studies. We focus on demographic parity when the protected attribute,  $A$ , is unavailable during prediction. As outlined in Proposition 4.10, the fairness constraint we consider here is linear, but not bilinear. Based on Corollary 4.11 and proposition 4.10, a  $\delta$ -fair Bayes-optimal classifier under this scenario is given by:

$$f_{X,\text{DD},\delta}^*(x) = I \left( 2\eta^Y(x) - 1 > t_{X,\text{DD}}(\delta) \left( \frac{\eta^A(x)}{p_1} - \frac{1 - \eta^A(x)}{p_0} \right) \right)$$

$$= I \left( \eta^Y(x) > \frac{1}{2} + \frac{t_{X,DD}(\delta)(\eta^A(x) - p_1)}{2p_1p_0} \right),$$

with

$$t_{X,DD}(\delta) = \arg \min_t \left\{ |t| : \left| \int_{\mathcal{X}} \left[ \left( \frac{\eta^A(x) - p_1}{p_1p_0} \right) \cdot I \left( \eta^Y(x) > \frac{1}{2} + \frac{t(\eta^A(x) - p_1)}{2p_1p_0} \right) \right] d\mathbb{P}_X(x) \right| \leq \delta \right\}. \quad (\text{B.12})$$

The following two theorems, proved below, parallel Theorems 5.1 and 5.3, providing a theoretical underpinning for our Fair Up-/Down-Sampling and Fair Cost-Sensitive Classification methods.

**Theorem B.6** (Bayes-Optimal FUDS for Demographic Parity with an Unavailable Protected Attribute). *For  $\delta \geq 0$ , let  $t_{X,DD}(\delta)$  be defined in (B.12). Let  $\tilde{\mathbb{P}}^\delta$  be the distribution satisfying  $\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=a,\tilde{Y}=y}^\delta(x) = \mathbb{P}_{X|A=a,Y=y}(x)$  for all  $x$ , and for  $(a, y) \in \{0, 1\}^2$ ,*

$$\tilde{p}_{a,y}^\delta = \tilde{\mathbb{P}}^\delta(\tilde{A} = a, \tilde{Y} = y) = c_{DD}(\delta) \left( 1 + \frac{(2a-1)(1-2y)t_{X,DD}(\delta)}{p_a} \right) p_{a,y}. \quad (\text{B.13})$$

where  $c_{DD}(\delta)$  is the constant such that  $\tilde{p}_{11}^\delta + \tilde{p}_{10}^\delta + \tilde{p}_{01}^\delta + \tilde{p}_{00}^\delta = 1$ . Let  $\tilde{\eta}^{Y,\delta}(x) = \tilde{\mathbb{P}}^\delta(\tilde{Y} = 1 | \tilde{X} = x)$  for all  $x$ . Then, the unconstrained Bayes-optimal classifier for  $\tilde{\mathbb{P}}^\delta$  is a  $\delta$ -fair Bayes-optimal classifier for  $\mathbb{P}$ .

With  $t_{X,DD}(\delta)$  defined in (B.12), consider the following fair cost-sensitive risk:

$$R_{t_{X,DD}(\delta)}^{\text{FCSC}}(f) = \sum_{a \in \{0,1\}} \sum_{y \in \{0,1\}} \left( \frac{1}{2} + \frac{(2a-1)(1-2a)t_{X,DD}(\delta)}{2p_a} \right) \mathbb{P}(\hat{Y}_f = 1 - y, Y = y, A = a),$$

and define the cost-sensitive classifier

$$f_{t_{X,DD}(\delta)}^{X,\text{FCSC}} \in \arg \min R_{t_{X,DD}(\delta)}^{\text{FCSC}}(f).$$

**Theorem B.7** (Bayes-Optimal FCSC for Demographic Parity with an Unavailable Protected Attribute). *For any  $\delta \geq 0$ , we have that  $f_{t_{X,DD}(\delta)}^{X,\text{FCSC}}$  is a  $\delta$ -fair Bayes-optimal classifier for  $\mathbb{P}$ .*

Based on these two theorems, algorithms analogous to those from Section 5 can be designed for the case that  $A$  is not used at test time.

*Proof of Theorem B.6.* By Bayes' theorem, we have for all  $x$  that

$$\begin{aligned} \eta^Y(x) &= \mathbb{P}(Y = 1 | X = x) = \mathbb{P}(Y = 1, A = 1 | X = x) + \mathbb{P}(Y = 1, A = 0 | X = x) \\ &= \frac{p_{1,1} d\mathbb{P}_{X|A=1,Y=1}(x)}{d\mathbb{P}_X(x)} + \frac{p_{0,1} d\mathbb{P}_{X|A=0,Y=1}(x)}{d\mathbb{P}_X(x)} \\ &= \frac{p_{1,1} d\mathbb{P}_{X|A=1,Y=1}(x) + p_{0,1} d\mathbb{P}_{X|A=0,Y=1}(x)}{p_{1,1} d\mathbb{P}_{X|A=1,Y=1}(x) + p_{1,0} d\mathbb{P}_{X|A=1,Y=0}(x) + p_{0,1} d\mathbb{P}_{X|A=0,Y=1}(x) + p_{0,0} d\mathbb{P}_{X|A=0,Y=0}(x)}. \end{aligned}$$

Similarly, we have for all  $x$  that

$$\begin{aligned} \tilde{\eta}^{Y,\delta}(x) &= \tilde{\mathbb{P}}^\delta(\tilde{Y} = 1 | \tilde{X} = x) = \tilde{\mathbb{P}}^\delta(\tilde{Y} = 1, \tilde{A} = 1 | \tilde{X} = x) + \tilde{\mathbb{P}}^\delta(\tilde{Y} = 1, \tilde{A} = 0 | \tilde{X} = x) \\ &= \frac{\tilde{p}_{1,1}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=1,\tilde{Y}=1}^\delta(x) + \tilde{p}_{0,1}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=0,\tilde{Y}=1}^\delta(x)}{\tilde{p}_{1,1}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=1,\tilde{Y}=1}^\delta(x) + \tilde{p}_{1,0}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=1,\tilde{Y}=0}^\delta(x) + \tilde{p}_{0,1}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=0,\tilde{Y}=1}^\delta(x) + \tilde{p}_{0,0}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=0,\tilde{Y}=0}^\delta(x)}. \end{aligned}$$

and

$$\begin{aligned} \eta^A(x) &= \mathbb{P}(A = 1 | X = x) = \mathbb{P}(A = 1, Y = 1 | X = x) + \mathbb{P}(A = 1, Y = 0 | X = x) \\ &= \frac{p_{1,1} d\mathbb{P}_{X|A=1,Y=1}(x)}{d\mathbb{P}_X(x)} + \frac{p_{1,0} d\mathbb{P}_{X|A=1,Y=0}(x)}{d\mathbb{P}_X(x)} \end{aligned}$$

$$= \frac{p_{1,1}d\mathbb{P}_{X|A=1,Y=1}(x) + p_{1,0}d\mathbb{P}_{X|A=1,Y=0}(x)}{p_{1,1}d\mathbb{P}_{X|A=1,Y=1}(x) + p_{1,0}d\mathbb{P}_{X|A=1,Y=0}(x) + p_{0,1}d\mathbb{P}_{X|A=0,Y=1}(x) + p_{0,0}d\mathbb{P}_{X|A=0,Y=0}(x)}.$$

Then, by our construction, for all  $x$ ,

$$\begin{aligned} f_{X,DD,t_{X,DD}(\delta)}^{\text{FUDES}}(x) &= I\left(\tilde{\eta}^{Y,\delta}(x) > \frac{1}{2}\right) \\ &= I\left(\frac{\tilde{p}_{1,1}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=1,\tilde{Y}=1}^\delta(x) + \tilde{p}_{0,1}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=0,\tilde{Y}=1}^\delta(x)}{\tilde{p}_{1,1}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=1,\tilde{Y}=1}^\delta(x) + \tilde{p}_{1,0}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=1,\tilde{Y}=0}^\delta(x) + \tilde{p}_{0,1}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=0,\tilde{Y}=1}^\delta(x) + \tilde{p}_{0,0}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=0,\tilde{Y}=0}^\delta(x)} > \frac{1}{2}\right) \\ &= I\left(\tilde{p}_{1,1}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=1,\tilde{Y}=1}^\delta(x) + \tilde{p}_{0,1}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=0,\tilde{Y}=1}^\delta(x) > \tilde{p}_{1,0}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=1,\tilde{Y}=0}^\delta(x) + \tilde{p}_{0,0}^\delta d\tilde{\mathbb{P}}_{\tilde{X}|\tilde{A}=0,\tilde{Y}=0}^\delta(x)\right) \\ &= I\left(c_{\text{DD}}(\delta) \cdot \left(1 - \frac{t_{X,DD}(\delta)}{p_1}\right) p_{1,1}\mathbb{P}_{X|A=1,Y=1}(x) + c_{\text{DD}}(\delta) \cdot \left(1 + \frac{t_{X,DD}(\delta)}{p_0}\right) p_{0,1}\mathbb{P}_{X|A=0,Y=1}(x)\right. \\ &> c_{\text{DD}}(\delta) \cdot \left(1 + \frac{t_{X,DD}(\delta)}{p_1}\right) p_{1,0}\mathbb{P}_{X|A=1,Y=0}(x) + c_{\text{DD}}(\delta) \cdot \left(1 - \frac{t_{X,DD}(\delta)}{p_0}\right) p_{0,0}\mathbb{P}_{X|A=0,Y=0}(x)\left. \right). \quad (\text{B.14}) \end{aligned}$$

Denote  $g_{a,y}(x) = p_{a,y}d\mathbb{P}_{X|A=a,Y=y}(x)$ , (B.14) further equals,

$$\begin{aligned} &I\left(g_{1,1}(x) + g_{0,1}(x) - g_{1,0}(x) - g_{0,0}(x) > \frac{t_{X,DD}(\delta)}{p_1}(g_{1,1}(x) + g_{1,0}(x)) - \frac{t_{X,DD}(\delta)}{p_0}(g_{0,1}(x) + g_{0,0}(x))\right) \\ &= I\left(2(g_{1,1}(x) + g_{0,1}(x)) - (g_{1,1}(x) + g_{0,1}(x) + g_{1,0}(x) + g_{0,0}(x))\right. \\ &> \left.\left(\frac{t_{X,DD}(\delta)}{p_1} + \frac{t_{X,DD}(\delta)}{p_0}\right)(g_{1,1}(x) + g_{1,0}(x)) - \frac{t_{X,DD}(\delta)}{p_0}(g_{1,1}(x) + g_{1,0}(x) + g_{0,1}(x) + g_{0,0}(x))\right) \\ &= I\left(\frac{2(g_{1,1}(x) + g_{0,1}(x))}{g_{1,1}(x) + g_{0,1}(x) + g_{1,0}(x) + g_{0,0}(x)} > \frac{t_{X,DD}(\delta)(g_{1,1}(x) + g_{1,0}(x))}{p_1 p_0 (g_{1,1}(x) + g_{0,1}(x) + g_{1,0}(x) + g_{0,0}(x))} + 1 - \frac{t_{X,DD}(\delta)}{p_0}\right) \\ &= I\left(2\eta^Y(x) > \frac{t_{X,DD}(\delta)}{p_1 p_0} \eta^A(x) + 1 - \frac{t_{X,DD}(\delta)}{p_0}\right) = I\left(\eta^Y(x) > \frac{1}{2} + \frac{t_{X,DD}(\delta)(\eta^A(x) - p_1)}{2p_1 p_0}\right) = f_{X,DD,\delta}^*(x). \end{aligned}$$

This completes the proof.  $\square$

*Proof of Theorem B.7.* Denoting for all  $a, y$ ,  $c_{a,y}(\delta) = 1/2 + (2a - 1)(1 - 2y)/(2p_a)$ , we have,

$$\begin{aligned} R_{t_{X,DD}(\delta)}^{\text{FCSC}}(f) &= \sum_{a \in \{0,1\}} \left[ c_{a,0}(\delta) \cdot \mathbb{P}(\hat{Y}_f = 1, Y = 0, A = a) + c_{a,1}(\delta) \cdot \mathbb{P}(\hat{Y}_f = 0, Y = 1, A = a) \right] \\ &= \sum_{a \in \{0,1\}} p_a \left[ c_{a,0}(\delta) \cdot \mathbb{P}(\hat{Y}_f = 1, Y = 0 | A = a) + c_{a,1}(\delta) \cdot \mathbb{P}(\hat{Y}_f = 0, Y = 1 | A = a) \right] \\ &= \sum_{a \in \{0,1\}} p_a c_{a,0}(\delta) \cdot \int_{\mathcal{X}} \mathbb{P}(\hat{Y}_f = 1, Y = 0 | A = a, X = x) d\mathbb{P}_{X|A=a}(x) \\ &+ \sum_{a \in \{0,1\}} p_a c_{a,1}(\delta) \cdot \int_{\mathcal{X}} \mathbb{P}(\hat{Y}_f = 0, Y = 1 | X = x, A = a) d\mathbb{P}_{X|A=a}(x). \end{aligned}$$

This further equals

$$\begin{aligned} &\sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} [c_{a,0}(\delta)f(x)(1 - \eta_a(x)) + c_{a,1}(\delta)(1 - f(x))\eta_a(x)] d\mathbb{P}_{X|A=a}(x) \\ &= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} [c_{a,0}(\delta)f(x) - c_{a,0}f(x)\eta_a(x) + c_{a,1}(\delta)\eta_a(x) - c_{a,1}(\delta)f(x)\eta_a(x)] d\mathbb{P}_{X|A=a}(x) \\ &= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} (c_{a,0}(\delta) - (c_{a,1}(\delta) + c_{a,0}(\delta))\eta_a(x)) f(x) d\mathbb{P}_{X|A=a} + \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} c_{a,1}(\delta)\eta_a(x) d\mathbb{P}_{X|A=a}(x) \end{aligned}$$



$$\begin{aligned}
&= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} (c_{a,0}(\delta) - \eta_a(x)) f(x) d\mathbb{P}_{X|A=a} + \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} c_{a,1}(\delta) \eta_a(x) d\mathbb{P}_{X|A=a}(x) \\
&= \int_{\mathcal{X}} \left( \sum_{a \in \{0,1\}} p_a (c_{a,0}(\delta) - \eta_a(x)) \frac{d\mathbb{P}_{X|A=a}(x)}{d\mathbb{P}_X(x)} \right) f(x) \mathbb{P}_X(x) + \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} c_{a,1}(\delta) \eta_a(x) d\mathbb{P}_{X|A=a}(x) \\
&= \int_{\mathcal{X}} \left( \sum_{a \in \{0,1\}} (c_{a,0}(\delta) - \eta_a(x)) \eta^A(x) \right) f(x) \mathbb{P}_X(x) + \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} c_{a,1}(\delta) \eta_a(x) d\mathbb{P}_{X|A=a}(x).
\end{aligned}$$

The second term does not depend on  $f$  and the first term can be expressed as:

$$\begin{aligned}
&\sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} (c_{a,0}(\delta) - \eta_a(x)) f(x) d\mathbb{P}_{X|A=a} = \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} (c_{a,0}(\delta) - \eta_a(x)) f(x) \frac{d\mathbb{P}_{X|A=a}(x)}{d\mathbb{P}_X(x)} d\mathbb{P}_X(x) \\
&= \int_{\mathcal{X}} \sum_{a \in \{0,1\}} [(c_{a,0}(\delta) - \eta_a(x)) \mathbb{P}(A = a|X = x)] f(x) d\mathbb{P}_X(x) \\
&= \int_{\mathcal{X}} \sum_{a \in \{0,1\}} [(c_{a,0}(\delta) \mathbb{P}(A = a|X = x) - \mathbb{P}(Y = 1, A = a|X = x))] f(x) d\mathbb{P}_X(x) \\
&= \int_{\mathcal{X}} [(c_{1,0}(\delta) \eta^A(x) + c_{0,0}(\delta)(1 - \eta^A(x))) - \eta^Y(x)] f(x) d\mathbb{P}_X(x) \\
&= \int_{\mathcal{X}} \left[ \left( \left( \frac{1}{2} + \frac{t_{X,DD}(\delta)}{2p_1} \right) \eta^A(x) + \left( \frac{1}{2} - \frac{t_{X,DD}(\delta)}{2p_0} \right) (1 - \eta^A(x)) \right) - \eta^Y(x) \right] f(x) d\mathbb{P}_X(x) \\
&= \int_{\mathcal{X}} \left[ \frac{1}{2} + \frac{t_{X,DD}(\eta^A(x) - p_1)}{2p_1p_0} - \eta^Y(x) \right] f(x) d\mathbb{P}_X(x).
\end{aligned}$$

Clearly, this quantity is minimized by taking  $f(x) = 1$  if  $\eta^Y(x) > 1/2 + t_{X,DD}(\delta) (\eta^A(x) - p_1)/(2p_1p_0)$  and  $f(x) = 0$  if  $\eta^Y(x) \leq 1/2 + t_{X,DD}(\delta) (\eta^A(x) - p_1)/(2p_1p_0)$ . We can thus conclude that for all  $x, a$ ,

$$f_{X,t_{X,DD}(\delta)}^{\text{FCSC}}(x) = I \left( \eta_a(x) > \frac{1}{2} + \frac{t_{X,DD}(\eta^A(x) - p_1)}{2p_1p_0} \right) = f_{X,DD,\delta}^*(x).$$

This finishes the proof.  $\square$

## B.4 Form of Fair Bayes-optimal Classifiers Without Distributional Assumptions

In our main text, we assume that, for  $a \in \{0,1\}$ ,  $\eta_a(X)$  has a probability density function on  $\mathcal{X}$ . In this section, we consider the more general case with no distributional assumptions on features. Without this assumption, the boundary case (where the features are exactly on the “fair” boundary) does not necessarily have zero probability. In this case, the optimal classifiers must be carefully randomized.

To handle this technical difficulty, we consider the randomized classifier. Let  $t \in \mathbb{R}$  and  $\tau : \mathcal{X} \times \mathcal{A} \rightarrow [0,1]$  be a measurable function. For  $x \in \mathcal{X}$  and  $a \in \{0,1\}$ , we define  $f_{\text{Dis},t,\tau}$  as,

$$f_{\text{Dis},t,\tau}(x, a) = I \left( \eta_a(x) > \frac{1}{2} + \frac{t}{2} w_{\text{Dis}}(x, a) \right) + \tau(x, a) I \left( \eta_a(x) = \frac{1}{2} + \frac{t}{2} w_{\text{Dis}}(x, a) \right). \quad (\text{B.15})$$

We further define the functions  $D_{\text{Dis},\min} : \mathbb{R} \rightarrow \mathbb{R}$  and  $D_{\text{Dis},\max} : \mathbb{R} \rightarrow \mathbb{R}$  to measure the minimal and maximal disparities of  $f_{\text{Dis},t,\tau}(x, a)$ , such that for all  $t \in \mathbb{R}$  and  $\tau : \mathcal{X} \times \{0,1\} \rightarrow [0,1]$ :

$$D_{\text{Dis},\min}(t) = \min_{\tau} \{\text{Dis}(f_{\text{Dis},t,\tau})\} \quad \text{and} \quad D_{\text{Dis},\max}(t) = \max_{\tau} \{\text{Dis}(f_{\text{Dis},t,\tau})\}.$$

We note that  $D_{\text{Dis},\min}(0)$  and  $D_{\text{Dis},\max}(0)$  are, respectively, the infimum and supremum of the disparity over all unconstrained Bayes-optimal classifiers. Moreover,  $D_{\text{Dis},\min}$  and  $D_{\text{Dis},\max}$  satisfy the following monotonicity properties.

**Proposition B.8** (Monotonicity of  $D_{\text{Dis},\min}$  and  $D_{\text{Dis},\max}$ ). *As functions of  $t$ , both  $D_{\text{Dis},\min}$  and  $D_{\text{Dis},\max}$  are monotone non-increasing.*

The proofs of all results are presented later in this section. For any  $\delta > 0$ , define the following quantity, which can be viewed as an “inverse” of the disparity functions:

$$t_{\text{Dis}}(\delta) = \arg \min_t \left\{ |t| : \inf_{\tau} |\text{Dis}(f_{\text{Dis},t,\tau})| \leq \delta \right\} = \begin{cases} \inf \{t : D_{\text{Dis},\min}(t) \leq \delta\}, & D_{\text{Dis},\min}(t) > \delta; \\ \sup \{t : D_{\text{Dis},\max}(t) \geq -\delta\}, & D_{\text{Dis},\max}(t) < -\delta; \\ 0, & \text{otherwise.} \end{cases} \quad (\text{B.16})$$

Again, we want to find the  $t$  with minimal  $|t|$  such that, for some  $\tau$ ,  $f_{\text{Dis},t,\tau}$  satisfies the fairness constraint. Here, we identify three different cases: (1)  $\delta \geq \max(D_{\text{Dis},\min}(t), -D_{\text{Dis},\max}(t))$ , (2)  $\delta < D_{\text{Dis},\min}(t)$ , and (3)  $\delta < -D_{\text{Dis},\max}(t)$ . These three cases are disjoint and cover all possible scenarios. Clearly, either case (2) or case (3) is true when case (1) does not hold. Furthermore, since  $\delta \geq 0$  and  $D_{\text{Dis},\min}(t) \leq D_{\text{Dis},\max}(t)$  by definition, cases (2) and (3) cannot occur simultaneously. We obtain the following result.

**Theorem B.9** (Fair Bayes-optimal Classifiers Without Distributional Assumptions). *For any  $\delta \geq 0$ , there is a measurable function  $\tau_{\text{Dis},\delta} : \mathcal{X} \times \{0, 1\} \rightarrow [0, 1]$ , such that  $f_{\text{Dis},\delta}^* = f_{t_{\text{Dis}}(\delta), \tau_{\text{Dis},\delta}}$  is a  $\delta$ -fair Bayes-optimal classifier. Here,  $\tau_{\text{Dis},\delta}$  is determined to satisfy the following constraints:*

(1). When  $D_{\text{Dis},\min}(t) > \delta$ ,

$$\text{Dis}(f_{\text{Dis},t_{\text{Dis}}(\delta), \tau_{\text{Dis},\delta}}) = \delta.$$

(2). When  $D_{\text{Dis},\max}(t) < -\delta$ ,

$$\text{Dis}(f_{\text{Dis},t_{\text{Dis}}(\delta), \tau_{\text{Dis},\delta}}) = -\delta.$$

(3). When  $\delta \geq \max(D_{\text{Dis},\min}(t), -D_{\text{Dis},\max}(t))$ ,

$$|\text{Dis}(f_{\text{Dis},t_{\text{Dis}}(\delta), \tau_{\text{Dis},\delta}})| \leq \delta.$$

To specify a particular form for  $\tau_{\text{Dis},\delta}$  in Theorem B.9, we first introduce the following proposition, which offers a more comprehensive understanding of  $D_{\text{Dis},\min}$  and  $D_{\text{Dis},\max}$ . Recall that for any function  $f : t \mapsto f(t)$ , we denote its left-hand limit and right-hand limit at point  $t \in \mathbb{R}$  as  $\lim_{t' \rightarrow t^-} f(t')$  and  $\lim_{t' \rightarrow t^+} f(t')$ , respectively.

**Proposition B.10.** *As functions of  $t$ ,*

(1) *For  $G_{a,+}$  and  $G_{a,-}$  in (B.17), we have, for all  $t \in \mathbb{R}$ , and all  $x, a$*

$$\tau_{\text{Dis},\min}(x, a) := I((x, a) \in G_{a,-}) \in \arg \min_{\tau} \{\text{Dis}(f_{\text{Dis},t,\tau})\};$$

$$\tau_{\text{Dis},\max}(x, a) := I((x, a) \in G_{a,+}) \in \arg \max_{\tau} \{\text{Dis}(f_{\text{Dis},t,\tau})\}.$$

*Moreover,*

$$\begin{aligned} D_{\text{Dis},\min}(t) &= \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} > t \right) \right] d\mathbb{P}_{X|A=a}(x) \\ &\quad + \sum_{a \in \{0,1\}} p_a \int_{G_{a,-}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t \right) \right] d\mathbb{P}_{X|A=a}(x); \\ D_{\text{Dis},\max}(t) &= \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \geq t \right) \right] d\mathbb{P}_{X|A=a}(x) \\ &\quad + \sum_{a \in \{0,1\}} p_a \int_{G_{a,-}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} < t \right) \right] d\mathbb{P}_{X|A=a}(x). \end{aligned}$$

(2) *We have that  $D_{\text{Dis},\min}$  is right-continuous and  $D_{\text{Dis},\max}$  is left-continuous.*

(3) We have, for  $t \in \mathbb{R}$ ,

$$\lim_{t' \rightarrow t^-} D_{\text{Dis},\min}(t') = D_{\text{Dis},\max}(t).$$

Now, we provide a specific choice of  $\tau_{\text{Dis},\delta}$ . Let  $\tau_0$  be the identically zero function with  $\tau_0(x, a) := 0$ , for all  $x, a$ , and for  $t \in \mathbb{R}$ ,  $D_{\text{Dis},0}(t) = \text{Dis}(f_{\text{Dis},t,\tau_0})$ . Clearly, for all  $t$ ,  $D_{\text{Dis},\min}(t) \leq D_{\text{Dis},0}(t) \leq D_{\text{Dis},\max}(t)$ . We consider three cases in order.

(1). When  $D_{\text{Dis},\min}(t) > \delta$ , we have  $t_{\text{Dis}}(\delta) = \inf \{t : D_{\text{Dis},\min}(t) \leq \delta\}$ . Since  $D_{\text{Dis},\min}$  is right-continuous, we have  $D_{\text{Dis},\min}(t) \leq \delta \leq \lim_{t' \rightarrow t^-} D_{\text{Dis},\min}(t') = D_{\text{Dis},\max}(t')$ . We can take, for all  $x \in \mathcal{X}$  and  $a \in \{0, 1\}$ ,

$$\tau_{\text{Dis},\delta}(x, a) = \begin{cases} \frac{[D_{\text{Dis},0}(t_{\text{Dis}}(\delta)) - \delta] \cdot I((x, a) \in G_{a,-})}{D_{\text{Dis},0}(t_{\text{Dis}}(\delta)) - D_{\text{Dis},\min}(t_{\text{Dis}}(\delta))}, & D_{\text{Dis},\min}(t_{\text{Dis}}(\delta)) \leq \delta \leq D_{\text{Dis},0}(t_{\text{Dis}}(\delta)); \\ \frac{[\delta - D_{\text{Dis},0}(t_{\text{Dis}}(\delta))] \cdot I((x, a) \in G_{a,+})}{D_{\text{Dis},\max}(t_{\text{Dis}}(\delta)) - D_{\text{Dis},0}(t_{\text{Dis}}(\delta))}, & D_{\text{Dis},0}(t_{\text{Dis}}(\delta)) < \delta \leq D_{\text{Dis},\max}(t_{\text{Dis}}(\delta)). \end{cases}$$

(2). When  $D_{\text{Dis},\max}(t) < -\delta$ , we have  $t_{\text{Dis}}(\delta) = \sup \{t : D_{\text{Dis},\max}(t) \geq -\delta\}$ . Since  $D_{\text{Dis},\max}$  is left-continuous, we have  $D_{\text{Dis},\min}(t) = \lim_{t' \rightarrow t^+} D_{\text{Dis},\max}(t') \leq -\delta \leq D_{\text{Dis},\max}(t)$ . We can take, for all  $x \in \mathcal{X}$  and  $a \in \{0, 1\}$ ,

$$\tau_{\text{Dis},\delta}(x, a) = \begin{cases} \frac{[D_{\text{Dis},0}(t_{\text{Dis}}(\delta)) + \delta] \cdot I((x, a) \in G_{a,-})}{D_{\text{Dis},0}(t_{\text{Dis}}(\delta)) - D_{\text{Dis},\min}(t_{\text{Dis}}(\delta))}, & D_{\text{Dis},\min}(t_{\text{Dis}}(\delta)) \leq -\delta \leq D_{\text{Dis},0}(t_{\text{Dis}}(\delta)); \\ \frac{[-\delta - D_{\text{Dis},0}(t_{\text{Dis}}(\delta))] \cdot I((x, a) \in G_{a,+})}{D_{\text{Dis},\max}(t_{\text{Dis}}(\delta)) - D_{\text{Dis},0}(t_{\text{Dis}}(\delta))}, & D_{\text{Dis},0}(t_{\text{Dis}}(\delta)) < -\delta \leq D_{\text{Dis},\max}(t_{\text{Dis}}(\delta)). \end{cases}$$

(3). When  $\delta \geq \max(D_{\text{Dis},\min}(t), -D_{\text{Dis},\max}(t))$ , since  $D_{\text{Dis},0} \leq D_{\text{Dis},0} \leq D_{\text{Dis},0}$ , we have  $D_{\text{Dis},\min} \leq \delta$  when  $\delta < D_{\text{Dis},0}$  and  $D_{\text{Dis},\max} \geq -\delta$  when  $\delta < -D_{\text{Dis},0}$ . Thus, we can take, for all  $x \in \mathcal{X}$  and  $a \in \{0, 1\}$ ,

$$\tau_{\text{Dis},\delta}(x, a) = \begin{cases} \frac{[D_{\text{Dis},0}(t_{\text{Dis}}(\delta)) - \delta] \cdot I((x, a) \in G_{a,-})}{D_{\text{Dis},0}(t_{\text{Dis}}(\delta)) - D_{\text{Dis},\min}(t_{\text{Dis}}(\delta))}, & \delta < D_{\text{Dis},0}(t_{\text{Dis}}(\delta)); \\ \frac{[-\delta - D_{\text{Dis},0}(t_{\text{Dis}}(\delta))] \cdot I((x, a) \in G_{a,+})}{D_{\text{Dis},\max}(t_{\text{Dis}}(\delta)) - D_{\text{Dis},0}(t_{\text{Dis}}(\delta))}, & \delta < -D_{\text{Dis},0}(t_{\text{Dis}}(\delta)); \\ 0; & \text{Otherwise.} \end{cases}$$

We now present the proofs of the results discussed in this section.

*Proof of Proposition B.8.* We define, for  $a \in \{0, 1\}$ ,

$$G_{a,+} = \{x \in \mathcal{X}, w_{\text{Dis}}(x, a) > 0\}; G_{a,0} = \{x \in \mathcal{X}, w_{\text{Dis}}(x, a) = 0\}; G_{a,-} = \{x \in \mathcal{X}, w_{\text{Dis}}(x, a) < 0\}. \quad (\text{B.17})$$

Let  $t_1 < t_2$ , as well as  $\tau_1 : \mathcal{X} \times \{0, 1\} \rightarrow [0, 1]$  and  $\tau_2 : \mathcal{X} \times \{0, 1\} \rightarrow [0, 1]$ . For  $x \in \mathcal{X}$  and  $a \in \{0, 1\}$  with  $w_{\text{Dis}}(x, a) \neq 0$ , we let  $q_{\text{Dis}}(x, a) = (2\eta_a(x) - 1)/w_{\text{Dis}}(x, a)$ . Then, we have, for all  $x, a$ ,

$$\begin{aligned} & f_{\text{Dis},t_1,\tau_1,\tau_0,1}(x, a) - f_{\text{Dis},t_2,\tau_1,\tau_0,2}(x, a) \\ &= \begin{cases} I(t_1 < q_{\text{Dis}}(x, a) \leq t_2) + \tau_{a,1}(x, a)I(q_{\text{Dis}}(x, a) = t_1) - \tau_{a,2}(x, a)I(q_{\text{Dis}}(x, a) = t_2), & x \in G_{a,+}; \\ -I(t_1 \leq q_{\text{Dis}}(x, a) < t_2) + \tau_{a,1}(x, a)I(q_{\text{Dis}}(x, a) = t_1) - \tau_{a,2}(x, a)I(q_{\text{Dis}}(x, a) = t_2), & x \in G_{a,-}; \\ (\tau_{a,1}(x, a) - \tau_{a,2}(x, a))I(\eta_a(x) = \frac{1}{2}), & x \in G_{a,0}; \end{cases} \\ &\geq \begin{cases} I\left(t_1 < \frac{2\eta_a(x)-1}{w_{\text{Dis}}(x,a)} < t_2\right), & x \in G_{a,+}; \\ -I\left(t_1 \leq \frac{2\eta_a(x)-1}{w_{\text{Dis}}(x,a)} \leq t_2\right), & x \in G_{a,-}; \\ (\tau_{a,1} - \tau_{a,2})I\left(\eta_a(x) = \frac{1}{2}\right), & x \in G_{a,0}. \end{cases} \end{aligned} \quad (\text{B.18})$$

It thus follows that

$$D_{\text{Dis},\min}(t_1) - D_{\text{Dis},\min}(t_2) = \inf_{\tau_1} \text{Dis}(f_{\text{Dis},t_1,\tau_1}) - \inf_{\tau_2} \text{Dis}(f_{\text{Dis},t_2,\tau_2})$$

$$\begin{aligned}
&\geq \inf_{\tau_1, \tau_2} \int_{\mathcal{A}} \int_{\mathcal{X}} [f_{\text{Dis}, t_1, \tau_1}(x, a) - f_{\text{Dis}, t_2, \tau_2}(x, a)] w_{\text{Dis}}(x, a) d\mathbb{P}_{X, A}(x, a) \\
&= \inf_{\tau_1, \tau_2} \left[ \sum_{a \in \{0, 1\}} p_a \int_{\mathcal{X}} [f_{\text{Dis}, t_1}(x, a) - f_{\text{Dis}, t_2}(x, a)] w_{\text{Dis}}(x, a) d\mathbb{P}_{X|A=a} \right] \\
&\geq \sum_{a \in \{0, 1\}} p_a \int_{G_{a,+}} I \left( t_1 < \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} < t_2 \right) w_{\text{Dis}}(x, a) d\mathbb{P}_{X|A=a} \\
&\quad - \sum_{a \in \{0, 1\}} p_a \int_{G_{a,-}} I \left( t_1 \leq \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t_2 \right) w_{\text{Dis}}(x, a) d\mathbb{P}_{X|A=a} \\
&\quad + \sum_{a \in \{0, 1\}} p_a \int_{G_{a,=}} I \left( \eta_a(x) = \frac{1}{2} \right) \cdot 0 d\mathbb{P}_{X|A=a} \geq 0.
\end{aligned}$$

The last inequality holds since the indicator function is non-negative, and  $w_{\text{Dis}}$  is positive on  $G_{a,+}$  and negative on  $G_{a,-}$ . Using a similar argument, we can verify that  $D_{\text{Dis}, \max}$  is also monotonically non-increasing.  $\square$

*Proof of Theorem B.9.* We analyze the following three cases: (1)  $\delta \geq \max\{D_{\text{Dis}, \min}(0), -D_{\text{Dis}, \max}(0)\}$ , (2)  $D_{\text{Dis}, \min}(0) > \delta$  and (3)  $D_{\text{Dis}, \max}(0) < -\delta$ . Since the proof for case (3) is analogous to that for case (2), we omit the discussion of case (3).

Case 1:  $\delta \geq \max\{D_{\text{Dis}, \min}(0), -D_{\text{Dis}, \max}(0)\}$ . In this case, there exists at least one unconstrained Bayes-optimal classifier satisfying the fairness constraint. As a result, we have  $t_{\text{Dis}}(\delta) = 0$  and the value of function  $\tau_{\text{Dis}, \delta}$  has no effect on the excess risk of the classifier. Thus, we only need to choose one such that

$$\left| \mathbb{P} \left( \widehat{Y}_{f_\delta^*} = 1 \mid A = 1 \right) - \mathbb{P} \left( \widehat{Y}_{f_\delta^*} = 0 \mid A = 1 \right) \right| \leq \delta.$$

Case 2:  $D_{\text{Dis}, \min}(0) > \delta$ . By the definition of  $t_{\text{Dis}}(\delta)$  in (B.16) and Proposition B.8, we have  $t_{\text{Dis}}(\delta) > 0$ . Let, for all  $x, a$ ,  $\phi_0(x, a) = 2\eta_1(x, a) - 1$ ,  $\phi_1(x, a) = w_{\text{Dis}}(x)$ . By Lemma A.1 and (3.6), we can write  $\text{Acc}(f)$  and  $D_{\text{Dis}}(f)$  as

$$\begin{aligned}
\text{Acc}(f) &= \int_{\mathcal{A}} \int_{\mathcal{X}} f(x, a) \phi_0(x, a) d\mathbb{P}_{X, A}(x, a) + \int_{\mathcal{A}} \int_{\mathcal{X}} (1 - \eta_a(x)) d\mathbb{P}_{X, A}(x, a); \\
\text{Dis}(f) &= \int_{\mathcal{X} \times \mathcal{A}} f(x, a) \phi_1(x, a) d\mathbb{P}_{X, A}(x, a).
\end{aligned}$$

Define

$$\mathcal{F}_= = \{f : \text{Dis}(f) = \delta\}; \mathcal{F}_{|\cdot|, \leq} = \{f : |\text{Dis}(f)| \leq \delta\}; \text{ and } \mathcal{F}_{\leq} = \{f : \text{Dis}(f) \leq \delta\}.$$

By our construction, we have  $f_{\text{Dis}, t_{\text{Dis}}(\delta), \tau_{\text{Dis}, \delta}} \in \mathcal{F}_= \subset \mathcal{F}_{|\cdot|, \leq} \subset \mathcal{F}_{\leq}$ . Since  $t_{\text{Dis}}(\delta) \geq 0$ , by the generalized Neyman-Pearson lemma (Lemma 3.1),

$$\max_{f \in \mathcal{F}_{\leq}} \text{Acc}(f) = \text{Acc}(f_{\text{Dis}, t_{\text{Dis}}(\delta), \tau_{\text{Dis}, \delta}}) \leq \max_{f \in \mathcal{F}_{|\cdot|, \leq}} \text{Acc}(f) \leq \max_{f \in \mathcal{F}_{\leq}} \text{Acc}(f).$$

Thus, we can conclude that

$$f_{\text{Dis}, t_{\text{Dis}}(\delta), \tau_{\text{Dis}, \delta}} = \arg \max_{f \in \mathcal{F}_{|\cdot|, \leq}} \text{Acc}(f) = \arg \min_{f \in \mathcal{F}} \{R(f) : |\text{Dis}(f)| \leq \delta\}.$$

The proof is thus completed.  $\square$

*Proof of Proposition B.10.* We only prove the results for  $D_{\text{Dis}, \min}$  and similar arguments apply to the result for  $D_{\text{Dis}, \max}(t)$ . For (1), by definition, we have

$$\text{Dis}(f_{\text{Dis}, t, \tau}) = \sum_{a \in \{0, 1\}} p_a \int_{\mathcal{X}} [w_{\text{Dis}}(x, a) f_{\text{Dis}, t, \tau}(x, a)] d\mathbb{P}_{X|A=a}(x)$$

$$\begin{aligned}
&= \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} \left[ w_{\text{Dis}}(x, a) I \left( \eta_a(x) > \frac{1}{2} + \frac{t}{2} w_{\text{Dis}}(x, a) \right) \right] d\mathbb{P}_{X|A=a}(x) \\
&\quad + \sum_{a \in \{0,1\}} p_a \int_{\mathcal{X}} \left[ w_{\text{Dis}}(x, a) \tau(x, a) I \left( \eta_a(x) = \frac{1}{2} + \frac{t}{2} w_{\text{Dis}}(x, a) \right) \right] d\mathbb{P}_{X|A=a}(x) \\
&= \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \eta_a(x) > \frac{1}{2} + \frac{t}{2} w_{\text{Dis}}(x, a) \right) \right] d\mathbb{P}_{X|A=a}(x) \\
&\quad + \sum_{a \in \{0,1\}} p_a \int_{G_{a,-}} \left[ w_{\text{Dis}}(x, a) I \left( \eta_a(x) > \frac{1}{2} + \frac{t}{2} w_{\text{Dis}}(x, a) \right) \right] d\mathbb{P}_{X|A=a}(x) \\
&\quad + \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) \tau(x, a) I \left( \eta_a(x) = \frac{1}{2} + \frac{t}{2} w_{\text{Dis}}(x, a) \right) \right] d\mathbb{P}_{X|A=a}(x) \\
&\quad + \sum_{a \in \{0,1\}} p_a \int_{G_{a,-}} \left[ w_{\text{Dis}}(x, a) \tau(x, a) I \left( \eta_a(x) = \frac{1}{2} + \frac{t}{2} w_{\text{Dis}}(x, a) \right) \right] d\mathbb{P}_{X|A=a}(x). \tag{B.19}
\end{aligned}$$

Since the indicator function is always non-negative, the infimum of (B.19) with respect to  $\tau : \mathcal{X} \times \{0, 1\} \rightarrow [0, 1]$  is achieved by taking  $\tau_{\text{Dis}, \min}(x, a) = I((x, a) \in G_{a,-})$  for all  $x, a$ . Moreover, we have,

$$\begin{aligned}
D_{\text{Dis}, \min}(t) &= \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \eta_a(x) > \frac{1}{2} + \frac{t}{2} w_{\text{Dis}}(x, a) \right) \right] d\mathbb{P}_{X|A=a}(x) \\
&\quad + \sum_{a \in \{0,1\}} p_a \int_{G_{a,-}} \left[ w_{\text{Dis}}(x, a) I \left( \eta_a(x) \geq \frac{1}{2} + \frac{t}{2} w_{\text{Dis}}(x, a) \right) \right] d\mathbb{P}_{X|A=a}(x) \\
&= \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} > t \right) \right] d\mathbb{P}_{X|A=a}(x) \\
&\quad + \sum_{a \in \{0,1\}} p_a \int_{G_{a,-}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t \right) \right] d\mathbb{P}_{X|A=a}(x).
\end{aligned}$$

For (2), denote

$$\begin{aligned}
D_{\text{Dis}, +, >}(t) &= \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} > t \right) \right] d\mathbb{P}_{X|A=a}(x) \\
D_{\text{Dis}, -, \leq}(t) &= \sum_{a \in \{0,1\}} p_a \int_{G_{a,-}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \leq t \right) \right] d\mathbb{P}_{X|A=a}(x).
\end{aligned}$$

We study the left- and right-hand limit of  $D_{\text{Dis}, +, >}$  at a point  $t \in \mathbb{R}$  as follows. Let  $(t_n)_{n=1}^{\infty}$  be monotone decreasing sequence tending to zero as  $n \rightarrow \infty$ . For fixed  $t \in \mathbb{R}$  and  $a \in \{0, 1\}$ , we consider the sets

$$\begin{aligned}
J_{a,-,t} &= \left\{ (x, a) : \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \geq t \right\}; & J_{n,a,-,t} &= \left\{ (x, a) : \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} > t - t_n \right\}; \\
J_{a,+,t} &= \left\{ (x, a) : \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} > t \right\}; & J_{n,a,+,t} &= \left\{ (x, a) : \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} > t + t_n \right\}.
\end{aligned}$$

and define the following functions on  $\mathcal{X} \times \{0, 1\}$ . For all  $x, a$ , let,

$$\begin{aligned}
j_{n,a,-,t}(x, a) &= w_{\text{Dis}}(x, a) I(G_{a,+} \cap J_{n,a,-,t}); & j_{a,-,t}(x, a) &= w_{\text{Dis}}(x, a) I(G_{a,+} \cap J_{a,-,t}); \\
j_{n,a,+,t}(x, a) &= w_{\text{Dis}}(x, a) I(G_{a,+} \cap J_{n,a,+,t}); & j_{a,+,t}(x, a) &= w_{\text{Dis}}(x, a) I(G_{a,+} \cap J_{a,+,t}).
\end{aligned}$$

It follows that

$$j_{1,a,-,t}(x, a) \geq j_{2,a,-,t}(x, a) \geq j_{3,a,-,t}(x, a) \geq \dots \quad \text{with} \quad \lim_{n \rightarrow \infty} j_{n,a,-,t}(x, a) = j_{a,-,t}(x, a);$$

$$j_{1,a,+,t}(x, a) \leq j_{2,a,+,t}(x, a) \leq j_{3,a,+,t}(x, a) \leq \dots \quad \text{with} \quad \lim_{n \rightarrow \infty} j_{n,a,+,t}(x, a) = j_{a,+,t}(x, a).$$

Then, by the monotone convergence theorem (Royden and Fitzpatrick, 2010),

$$\begin{aligned} \lim_{t' \rightarrow t^-} D_{\text{Dis},+,>}(t') &= \lim_{n \rightarrow \infty} D_{\text{Dis},+,>}(t - t_n) \\ &= \lim_{n \rightarrow \infty} \left[ \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} > t - t_n \right) \right] d\mathbb{P}_{X|A=a}(x) \right] \\ &= \sum_{a \in \{0,1\}} \left\{ p_a \lim_{n \rightarrow \infty} \left[ \int_{\mathcal{X}} j_{n,a,-,t}(x, a) d\mathbb{P}_{X|A=a}(x) \right] \right\} = \sum_{a \in \{0,1\}} \left\{ p_a \int_{\mathcal{X}} j_{a,-,t}(x, a) d\mathbb{P}_{X|A=a}(x) \right\} \\ &= \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \geq t \right) \right] d\mathbb{P}_{X|A=a}(x). \end{aligned}$$

Similarly, we can verify that

$$\begin{aligned} \lim_{t' \rightarrow t^+} D_{\text{Dis},+,>}(t') &= \lim_{n \rightarrow \infty} D_{\text{Dis},+,>}(t + t_n) \\ &= \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} > t \right) \right] d\mathbb{P}_{X|A=a}(x) = D_{\text{Dis},+,>}(t). \end{aligned}$$

Thus,  $D_{\text{Dis},+,>}$  is right-continuous and for  $t \in \mathbb{R}$ ,

$$\lim_{t' \rightarrow t^-} D_{\text{Dis},+,>}(t') = \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \geq t \right) \right] d\mathbb{P}_{X|A=a}(x).$$

With the similar arguments above, we can show that  $D_{\text{Dis},-,\leq}$  is also right-continuous, moreover, for  $t \in \mathbb{R}$

$$\lim_{t' \rightarrow t^-} D_{\text{Dis},-,\leq}(t') = \sum_{a \in \{0,1\}} p_a \int_{G_{a,-}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} < t \right) \right] d\mathbb{P}_{X|A=a}(x).$$

As a consequence,  $D_{\text{Dis},\min} = D_{\text{Dis},+,>} + D_{\text{Dis},-,\leq}$  is right-continuous.

Finally, (3) holds since, for  $t \in \mathbb{R}$ ,

$$\begin{aligned} \lim_{t' \rightarrow t^+} D_{\text{Dis},\min}(t') &= \lim_{t' \rightarrow t^+} D_{\text{Dis},+,>}(t') + \lim_{t' \rightarrow t^+} D_{\text{Dis},-,\leq}(t') \\ &= \sum_{a \in \{0,1\}} p_a \int_{G_{a,+}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} \geq t \right) \right] d\mathbb{P}_{X|A=a}(x) \\ &\quad + \sum_{a \in \{0,1\}} p_a \int_{G_{a,-}} \left[ w_{\text{Dis}}(x, a) I \left( \frac{2\eta_a(x) - 1}{w_{\text{Dis}}(x, a)} < t \right) \right] d\mathbb{P}_{X|A=a}(x) = D_{\text{Dis},\max}(t). \end{aligned}$$

This finishes the proof.  $\square$

## C Bayes-optimal Classifiers for Synthetic Data

In this section, we derive the  $\delta$ -fair Bayes-optimal classifiers for our synthetic model used in Section 6.1. In particular, we consider the following data distribution of  $(X, A, Y)$ : for  $(a, y) \in \{0, 1\}^2$ , we let  $\mathbb{P}(A = a, Y = y) = p_{a,y}$  and  $X|A = a, Y = y \sim \mathcal{N}(\mu_{a,y}, \sigma^2 I_p)$ .

Denote, for all  $x$ , by  $g_{a,y}(x) = (2\pi)^{-p/2} \sigma^{-p} \exp(-\|x - \mu_{a,y}\|^2 / (2\sigma^2))$  the conditional probability density function of  $X$  given  $A = a$  and  $Y = y$ . We have

$$\eta_a(x) = \mathbb{P}(Y = 1 | X = x, A = a) = \frac{p_{a,1} g_{a,1}(x)}{p_{a,1} g_{a,1}(x) + p_{a,0} g_{a,0}(x)}$$

$$= \frac{p_{a,1} \exp(-\frac{1}{2\sigma^2} \|x - \mu_{a,1}\|^2)}{p_{a,1} \exp(-\frac{1}{2\sigma^2} \|x - \mu_{a,1}\|^2) + p_{a,0} \exp(-\frac{1}{2\sigma^2} \|x - \mu_{a,0}\|^2)}.$$

In the following, for  $a \in \{0, 1\}$ , we denote  $\Delta_{\mu,a} = \mu_{a,1} - \mu_{a,0}$ . Moreover, for  $t \in [0, 1]$  and  $a \in \{0, 1\}$ , we denote,  $q_a(t) = tp_{a,0}/((1-t)p_{a,1})$ . Then, we have, for  $(a, y) \in \{0, 1\}^2$ ,

$$\begin{aligned} \mathbb{P}_{X|A=a, Y=y}(\eta_a(X) > t) &= \mathbb{P}_{X|A=a, Y=y} \left( \frac{p_{a,1}g_{a,1}(x)}{p_{a,1}g_{a,1}(x) + p_{a,0}g_{a,0}(x)} > t \right) \\ &= \mathbb{P}_{X|A=a, Y=y} \left( (1-t)p_{a,1}g_{a,1}(x) > tp_{a,0}g_{a,0}(x) \right) = \mathbb{P}_{X|A=a, Y=y} (g_{a,1}(x) > q_a(t)g_{a,0}(x)) \\ &= \mathbb{P}_{X|A=a, Y=y} \left( \exp\left(-\frac{1}{2\sigma^2} \|x - \mu_{a,1}\|^2\right) > q_a(t) \cdot \exp\left(-\frac{1}{2\sigma^2} \|x - \mu_{a,0}\|^2\right) \right) \\ &= \mathbb{P}_{X|A=a, Y=y} \left( \|x - \mu_{a,0}\|^2 - \|x - \mu_{a,1}\|^2 > 2\sigma^2 \log(q_a(t)) \right) \\ &= \mathbb{P}_{X|A=a, Y=y} \left( 2x^\top \Delta_{\mu,a} > 2\sigma^2 \log(q_a(t)) + \|\mu_{a,1}\|^2 - \|\mu_{a,0}\|^2 \right) \\ &= \mathbb{P}_{Z \sim \mathcal{N}(0, I_p)} \left( 2(\sigma \cdot Z + \mu_{a,y})^\top \Delta_{\mu,a} > 2\sigma^2 \log(q_a(t)) + \|\mu_{a,1}\|^2 - \|\mu_{a,0}\|^2 \right). \end{aligned}$$

From the definition of the distribution of  $(X, A, Y)$ , it follows that this further equals

$$\begin{aligned} &\mathbb{P}_{Z \sim \mathcal{N}(0, I_p)} \left( \sigma \cdot Z^\top \Delta_{\mu,a} > \sigma^2 \log(q_a(t)) + \frac{\|\mu_{a,1}\|^2 - \|\mu_{a,0}\|^2 - 2\mu_{a,y}^\top \Delta_{\mu,a}}{2} \right) \\ &= \mathbb{P}_{Z \sim \mathcal{N}(0, I_p)} \left( Z^\top \Delta_{\mu,a} > \sigma \log(q_a(t)) + (1-2y) \frac{\|\Delta_{\mu,a}\|^2}{2\sigma} \right) \\ &= \mathbb{P}_{Z' \sim \mathcal{N}(0, 1)} \left( Z' > \frac{\sigma \log(q_a(t))}{\|\Delta_{\mu,a}\|} + (1-2y) \frac{\|\Delta_{\mu,a}\|}{2\sigma} \right) = 1 - \Phi \left( \frac{\sigma \log(q_a(t))}{\|\Delta_{\mu,a}\|} + (1-2y) \frac{\|\Delta_{\mu,a}\|}{2\sigma} \right), \end{aligned}$$

where  $\Phi(\cdot)$  is the cumulative distribution function of the standard normal distribution. Hereafter, we denote, for  $t \in [0, 1]$ , and  $(a, y) \in \{0, 1\}^2$ ,

$$\psi_{a,y}(t) = \Phi \left( \frac{\sigma \log(q_a(t))}{\|\Delta_{\mu,a}\|} + (1-2y) \frac{\|\Delta_{\mu,a}\|}{2\sigma} \right).$$

Next, we derive  $\delta$ -fair Bayes-optimal classifiers under demographic parity, equality of opportunity and predictive parity.

## C.1 Demographic Parity

For demographic parity, the weight function is given by, for all  $(x, a)$ ,  $w_{DD}(x, a) = 1/p_a$ . This time, the disparity function DD take values, for  $t \in \mathcal{T}_{DD} := [-\min(p_1, p_0), \min(p_1, p_0)]$ ,

$$\begin{aligned} \text{DD}(t) &= \sum_{a \in \{0,1\}} (2a-1) \mathbb{P}_{X|A=a} \left( \eta_a(X) > \frac{1}{2} + \frac{t}{2p_a} \right) \\ &= \sum_{a \in \{0,1\}} \sum_{y \in \{0,1\}} \left[ (2a-1) \mathbb{P}_{X|A=a} \left( Y = y, \eta_a(X) > \frac{1}{2} + \frac{t}{2p_a} \right) \right] \\ &= \sum_{a \in \{0,1\}} \sum_{y \in \{0,1\}} \left[ (2a-1) \mathbb{P}(Y = y|A = a) \cdot \mathbb{P}_{X|A=a, Y=y} \left( \eta_a(X) > \frac{1}{2} + \frac{t}{2p_a} \right) \right] \\ &= \sum_{a \in \{0,1\}} \sum_{y \in \{0,1\}} \left[ \frac{(2a-1)p_{a,y}}{p_a} \left( 1 - \psi_{a,y} \left( \frac{1}{2} + \frac{t}{2p_a} \right) \right) \right] \\ &= -\frac{p_{1,1}}{p_1} \psi_{1,1} \left( \frac{p_1+t}{2p_1} \right) - \frac{p_{1,0}}{p_1} \psi_{1,0} \left( \frac{p_1+t}{2p_1} \right) + \frac{p_{0,1}}{p_0} \psi_{0,1} \left( \frac{p_0-t}{2p_0} \right) + \frac{p_{0,0}}{p_0} \psi_{0,0} \left( \frac{p_0-t}{2p_0} \right). \end{aligned}$$



By Corollary 4.3, a  $\delta$ -fair Bayes-optimal classifier is, for all  $x, a$ ,

$$f_{\text{DD},\delta}^*(x, a) = I\left(\eta_a(x) > \frac{1}{2} + \frac{t_{\text{DD}}(\delta)}{p_a}\right),$$

with  $t_{\text{DD}}(\delta) = \arg \min_{t \in \mathcal{T}_{\text{DD}}} \{ |t| : \text{DD}(t) \leq \delta \}$ . Moreover, we have,

$$\begin{aligned} R(f_{\text{DD},\delta}^*) &= \mathbb{P}\left(\widehat{Y}_{f_{\text{DD},\delta}^*} \neq Y\right) = \sum_{y \in \{0,1\}} \mathbb{P}\left(\widehat{Y}_{f_{\text{DD},\delta}^*} = y, Y = 1 - y\right) \\ &= \sum_{a \in \{0,1\}} \sum_{y \in \{0,1\}} p_{a,y} \cdot \mathbb{P}_{X|A=a, Y=y}\left(\widehat{Y}_{f_{\text{DD},\delta}^*} = 1 - y\right) \\ &= \sum_{a \in \{0,1\}} \left[ p_{a,1} \cdot \mathbb{P}_{A=a, Y=1}\left(\eta_a(X) \leq \frac{1}{2} + \frac{t}{2p_a}\right) + p_{a,0} \cdot \mathbb{P}_{A=a, Y=0}\left(\eta_a(X) > \frac{1}{2} + \frac{t}{2p_a}\right) \right] \\ &= p_{1,0} + p_{0,0} + p_{1,1}\psi_{1,1}\left(\frac{p_{1,1} + t}{2p_1}\right) - p_{1,0}\psi_{1,0}\left(\frac{p_{1,1} + t}{2p_1}\right) + p_{0,1}\psi_{0,1}\left(\frac{p_{0,1} - t}{2p_0}\right) - p_{0,0}\psi_{0,0}\left(\frac{p_{0,1} - t}{2p_0}\right). \end{aligned}$$

## C.2 Equality of Opportunity

For equality of opportunity, the weight function is given for all  $(x, a)$ , by  $w_{\text{DO}}(x, a) = \eta_a(x)/p_{a,1}$ . The disparity function DO take values, for  $t \in \mathcal{T}_{\text{DO}} := [-p_{0,1}, p_{1,1}]$ ,

$$\begin{aligned} D_{\text{DO}}(t) &= \sum_{a \in \{0,1\}} (2a - 1) \mathbb{P}_{A=a, Y=1}\left(\eta_a(X) > \frac{1}{2} + \frac{t\eta_a(X)}{p_{a,1}}\right) \\ &= \sum_{a \in \{0,1\}} \left[ (2a - 1) \mathbb{P}_{A=a, Y=1}\left(\eta_a(X) > \frac{p_{a,1}}{2p_{a,1} - t}\right) \right] \\ &= p_{1,1} \left(1 - \psi_{1,1}\left(\frac{p_{1,1}}{2p_{1,1} - t}\right)\right) - p_{0,1} \left(1 - \psi_{0,1}\left(\frac{p_{0,1}}{2p_{0,1} + t}\right)\right). \end{aligned}$$

By Corollary 4.3, a  $\delta$ -fair Bayes-optimal classifier is, for all  $x, a$ ,

$$f_{\text{DO},\delta}^*(x, a) = I\left(\eta_a(x) > \frac{p_{a,1}}{2p_{a,1} - t_{\text{DO}}(\delta)}\right),$$

with  $t_{\text{DO}}(\delta) = \arg \min_{t \in \mathcal{T}_{\text{DO}}} \{ |t| : D_{\text{DO}}(t) \leq \delta \}$ . Similarly, we have,

$$\begin{aligned} R(f_{\text{DO},\delta}^*) &= \mathbb{P}\left(\widehat{Y}_{f_{\text{DO},\delta}^*} \neq Y\right) = \sum_{y \in \{0,1\}} \mathbb{P}\left(\widehat{Y}_{f_{\text{DO},\delta}^*} = y, Y = 1 - y\right) \\ &= p_{1,0} + p_{0,0} + p_{1,1}\psi_{1,1}\left(\frac{p_{1,1}}{2p_{1,1} - t}\right) - p_{1,0}\psi_{1,0}\left(\frac{p_{1,1}}{2p_{1,1} - t}\right) + p_{0,1}\psi_{0,1}\left(\frac{p_{0,1}}{2p_{0,1} + t}\right) - p_{0,0}\psi_{0,0}\left(\frac{p_{0,1}}{2p_{0,1} + t}\right). \end{aligned}$$

## C.3 Predictive Equality

For predictive equality, the weight function is given by, for all  $(x, a)$ ,  $w_{\text{PD}}(x, a) = (1 - \eta_a(x))/p_{a,0}$ . This time, the disparity function PD take values, for  $t \in \mathcal{T}_{\text{PD}} := [-p_{1,0}, p_{0,0}]$ ,

$$\begin{aligned} D_{\text{PD}}(t) &= \sum_{a \in \{0,1\}} (2a - 1) \mathbb{P}_{A=a, Y=0}\left(\eta_a(X) > \frac{1}{2} + \frac{t(1 - \eta_a(X))}{p_{a,0}}\right) \\ &= \sum_{a \in \{0,1\}} \left[ (2a - 1) \mathbb{P}_{A=a, Y=0}\left(\eta_a(X) > \frac{p_{a,0} + t}{2p_{a,0} + t}\right) \right] \\ &= p_{1,0} \left(1 - \psi_{1,0}\left(\frac{p_{1,0} + t}{2p_{1,0} + t}\right)\right) - p_{0,0} \left(1 - \psi_{0,0}\left(\frac{p_{0,0} - t}{2p_{0,0} - t}\right)\right). \end{aligned}$$

By Corollary 4.3, a  $\delta$ -fair Bayes-optimal classifier is, for all  $x, a$ ,

$$f_{\text{PD},\delta}^*(x, a) = I\left(\eta_a(x) > \frac{p_{a,0} + t_{\text{PD}}(\delta)}{2p_{a,0} + t_{\text{PD}}(\delta)}\right),$$

with  $t_{\text{PD}}(\delta) = \arg \min_{t \in \mathcal{T}_{\text{PD}}} \{|t| : D_{\text{PD}}(t) \leq \delta\}$ . Moreover,

$$\begin{aligned} R(f_{\text{PD},\delta}^*) &= \mathbb{P}\left(\widehat{Y}_{f_{\text{PD},\delta}^*} \neq Y\right) = \sum_{y \in \{0,1\}} \mathbb{P}\left(\widehat{Y}_{f_{\text{PD},\delta}^*} = y, Y = 1 - y\right) \\ &= p_{1,0} + p_{0,0} + p_{1,1}\psi_{1,1}\left(\frac{p_{1,0} + t}{2p_{1,0} + t}\right) - p_{1,0}\psi_{1,0}\left(\frac{p_{1,0} + t}{2p_{1,0} + t}\right) + p_{0,1}\psi_{0,1}\left(\frac{p_{0,0} - t}{2p_{0,0} - t}\right) - p_{0,0}\psi_{0,0}\left(\frac{p_{0,0} - t}{2p_{0,0} - t}\right). \end{aligned}$$