

Independent set reconfiguration in H -free graphs*

Valentin Bartier¹, Nicolas Bousquet¹, and Moritz Mühenthaler²

¹Univ. Lyon, Université Lyon 1, CNRS, LIRIS UMR CNRS 5205, F-69621, Lyon, France

²G-SCOP, Université Grenoble-Alpes, Grenoble, France

Abstract

Given a graph G and two independent sets of G , the independent set reconfiguration problem asks whether one independent set can be transformed into the other by moving a single vertex at a time, such that at each intermediate step we have an independent set of G . We study the complexity of this problem for H -free graphs under the token sliding and token jumping rule. Our contribution is twofold. First, we prove a reconfiguration analogue of Alekseev’s theorem, showing that the problem is PSPACE-complete unless H is a path or a subdivision of the claw. We then show that under the token sliding rule, the problem admits a polynomial-time algorithm if the input graph is fork-free.

1 Introduction

For some nominal combinatorial problem, such as the satisfiability problem or the independent set problem, a corresponding *reconfiguration problem* asks the following question: Given two solutions to some instance of the nominal problem, can one be transformed into the other by a sequence of allowed modifications, such that each intermediate solution is feasible? Such problems appear in many domains, such as combinatorial puzzles (Rubik’s cube, Sokoban), motion planning, or in the context of phylogenetic trees, see [vdH13, Nis18] for an overview of recent results. The complexity of reconfiguration problems on graphs has received considerable attention recently, for example the reconfiguration of vertex covers [MNR⁺17, MNRS18], dominating sets [HIM⁺16, SMN16, BDO21], colorings [BP16, BHI⁺20, DF21, BDK⁺23], and independent sets [BKW14, BBD⁺21]. The natural complexity class of reconfiguration problems is PSPACE and we may ask under which conditions we can do better, that is, under which conditions a reconfiguration problem admits a polynomial-time algorithm. We consider this question for the reconfiguration of independent sets in graphs that are H -free, that is, graphs that do not contain a fixed graph H as an induced subgraph.

Our study of independent set reconfiguration in H -free graphs is largely motivated by a similar line of research on the nominal problem of finding a maximum independent set (MIS) in H -free graphs. The input is an H -free graph G and the task is to find an independent set of G of maximum size. Despite roughly 40 years of effort, a complete classification of the complexity is not known. Alekseev showed that the problem remains NP-hard for a large class of graphs H .

Theorem 1 ([Ale82]). *The maximum independent set problem on H -free graphs is NP-hard, unless H is a path, the claw, or a subdivision of the claw.*

Let P_k denote the path graph on k vertices and see Figure 1 for illustrations of P_4 , the claw and the fork. Minty [Min80] and Sbihi [Sbi80] showed independently that a maximum independent set of a claw-free graph can be computed in polynomial time. Twenty years later, Alekseev generalized this result to fork-free graphs [Ale04]. Noticing that a maximum independent set can be computed in polynomial time P_4 -free graphs, the smallest interesting case left open by Theorem 1 is $H = P_5$. The complexity of this problem remained open for three decades, but recently, polynomial-time algorithms were obtained for $H = P_5$ [LVV14] and $H = P_6$ [GKPP22]. The complexity remains open for $H = P_\ell$ for $\ell \geq 7$.

*The authors are supported by ANR project GrR (ANR-18-CE40-0032).




Figure 1: Small graphs of interest.

We follow the line of results above and investigate the complexity of INDEPENDENT SET RECONFIGURATION (ISR) in H -free graphs. The ISR problem has been introduced in [IDH⁺11] and it has been studied widely since, see e.g., [BKW14, IKO⁺14b, IKO14a, DDF⁺15, FHO15, BB17, BBD⁺21]. An instance of ISR is a graph G and two independent sets I and J of G of the same size, which are considered as *tokens* placed on the vertices of G . The question is whether I can be transformed into J by a sequence of *moves*, such that in each intermediate step the tokens form an independent set. Two main types of moves have been considered: *token jumping*, where a token can move to any other vertex of the graph (provided that it is not adjacent to a token), and *token sliding*, where a token can slide along an edge to a neighboring vertex. We denote by TOKEN JUMPING and TOKEN SLIDING and the corresponding variants of ISR, where the only allowed moves are token jumping and token sliding, respectively. Both problems are equivalent for maximum independent sets and are known to be PSPACE-complete, even for planar graphs of maximum degree three [HD05, IDH⁺11]. Polynomial-time algorithms are known for example on trees [DDF⁺15], interval graphs [BB17], bipartite permutation and bipartite distance hereditary graphs [FHO15].

We prove the following reconfiguration analogue of Alekseev’s theorem for ISR in H -free graphs.

Theorem 2. *TOKEN SLIDING and TOKEN JUMPING in H -free graphs are PSPACE-complete, unless H is a path, the claw, or a subdivision of the claw.*

It is known that TOKEN SLIDING and TOKEN JUMPING admit polynomial-time algorithm on P_4 -free graphs [Bon16, KMM12] and claw-free graphs [BKW14]. We generalize the results for TOKEN SLIDING, showing that the problem admits a polynomial-time algorithm in fork-free graphs.

Theorem 3. *TOKEN SLIDING in fork-free graphs admits a polynomial-time algorithm.*

Let us note that the complexity of TOKEN JUMPING on fork-free graphs remains open. Furthermore, the smallest graph H for which the complexity of TOKEN SLIDING in H -free graphs is open is $H = P_5$.

Overview of the proofs. In order to prove Theorem 2, we use similar ideas to those that Alekseev used to prove Theorem 1. Let G_t be the graph obtained from a graph $G = (V, E)$ by subdividing each edge of G exactly t times. The main idea is to relate maximum independent sets of a graph G with those of G_t . In [Ale82] Alekseev showed that $\alpha(G_{2t}) = \alpha(G) + t|E|$, where $\alpha(G)$ denotes the size of a maximum independent set of G . We may therefore assume that the input graph has arbitrarily large girth and that vertices of degree at least three are arbitrarily far apart. Theorem 1 then follows from the fact that MIS is NP-hard on graph of maximum degree three. To prove Theorem 2, we associate each maximum independent set of G in a canonical way with a maximum independent set of G_{2t} . Let I and J be two maximum independent sets of G and let I' and J' be the corresponding canonical independent sets of G_{2t} . We show that there exists a transformation from I to J if and only if there exists a transformation from I' to J' . Combining this with the fact that TOKEN SLIDING and TOKEN JUMPING are PSPACE-complete on graphs of maximum degree three [HD05, IDH⁺11] yields Theorem 2.

We then show that TOKEN SLIDING in H -free admits a polynomial-time algorithm (Theorem 3). Let G be a fork-free graph and I and J be independent sets of G of the same size. We notice that a vertex v of G is

irrelevant for TOKEN SLIDING if no independent set reachable from I contains v . Throughout the algorithm, we reduce the graph by deleting irrelevant vertices. For instance, any vertex with at least three tokens in its neighborhood is irrelevant. In general, we distinguish two cases, depending on whether I and J are maximum or not. In the first case, we provide a simple reduction to TOKEN SLIDING in claw-free graphs. We show that if I is maximum then any induced claw of G contains precisely one token on a degree-one vertex of the claw. This means that the center of any claw is irrelevant and may be removed from the graph. We thus obtain an equivalent instance on a claw-free graph. Notice that by the equivalence of TOKEN SLIDING and TOKEN JUMPING for maximum independent sets, this also yields a polynomial-time algorithm for TOKEN JUMPING in claw-free graphs if the independent sets are maximum.

The case that I and J are not maximum is more involved. We observe that, after removing irrelevant vertices, each connected component the graph induced by the vertices I and J is either a path or a cycle. Sliding tokens on induced paths is easy, so it remains to deal with the cycles and isolated vertices. Similar obstructions occur for alternating cycles in MATCHING RECONFIGURATION [IDH⁺11] and cycles induced by the symmetric difference of independent sets of claw-free graphs [BKW14]). We show that by sliding tokens along an I -augmenting path (I is not maximum), we may assume that G contains an I -free vertex, that is, a vertex f , such that $I + f$ is an independent set. We then show that, for any vertex c of a cycle in the graph induced by the symmetric difference of the two independent sets, either we can transform I into $I - c + f$ or find an irrelevant vertex, which can be removed (Theorem 14). In the former case, this allows us to move the remaining tokens on the cycle to their target positions in J . To prove Theorem 14, we first show that we may assume that every connected component of G is prime with respect to the modular decomposition. Notice that the modular decomposition was used “in disguise” in the form of a *cotree* in the polynomial-time algorithm for ISR in cographs (P_4 -free graphs) in [Bon16]. In order to move the token on c to the free vertex f , we imagine sliding it on a shortest path P from c to f . Then c may be blocked by other tokens on P or in the neighborhood of P . Using a structural characterization of claws in fork-free prime graphs due to Brandstädt et al. [BHV04], we show we may move c and every vertex that is blocking c forward on the path, one by one in reverse order, or find an irrelevant vertex (see in particular Lemma 18).

Related work. The complexity of independent set reconfiguration has been widely studied in the literature, for example in particular graph classes [DDF⁺15, BB17, FHO15] and in the context of parameterized complexity [Nis18, IKO⁺20]. For H -free graphs we are aware of only two results: Bonsma [Bon16] showed that TOKEN SLIDING is solvable in polynomial on in P_4 -free graphs. In [BKW14], Bonsma et al. showed that TOKEN SLIDING and TOKEN JUMPING admit a polynomial-time algorithm on claw-free graphs [BKW14]. In the case that more than one graph is forbidden (as induced subgraph), Kamiński et al. [KMM12] showed that TOKEN JUMPING admits a polynomial-time algorithm on even-hole free graphs. Interestingly, the complexity of MIS is open for this graph class. The complexity of TOKEN SLIDING and TOKEN JUMPING can be very different, for example, Lokshtanov and Mouawad [LM19] showed that TOKEN JUMPING is NP-complete on bipartite graphs (which are odd-hole free), whereas TOKEN SLIDING remains PSPACE-complete. Similarly, TOKEN JUMPING admits a linear-time algorithm on split graphs, while TOKEN SLIDING is PSPACE-complete on this graph class [BKL⁺21]. The reader is referred to [Bar21] for a complete and recent bibliography on the topic. ISR has also been studied from the point of view of parameterized complexity, where the parameter is for example the size of the independent sets or the length of the reconfiguration sequence. The reader is referred to [BMNS22] for an overview of results in this direction. Many other reconfiguration problems have been studied when an induced subgraph is forbidden, especially from a structural point of view. For single vertex recoloring for instance, a complete dichotomy theorem was recently obtained by Belavadi, Cameron, Merkel in [BCM23], who characterized exactly for which graphs H , $G(G, \chi(G) + 1)$ is connected for every H -free graph G (where $\chi(G)$ is the chromatic number of G).

Organization. Section 2 contains useful notation and definitions. In Section 3, we prove the reconfiguration analogue of Alekseev’s theorem (Theorem 2). We discuss irrelevant vertices of fork-free graphs in Section 4. We propose a polynomial-time algorithm for TOKEN SLIDING for maximum independent sets in fork-free graphs in Section 5 and for non-maximum independent sets in Section 6. Section 7 concludes the

paper with a discussion of open questions.

2 Preliminaries

For two sets U, W we denote by $U \Delta W$ their symmetric difference $(U \setminus W) \cup (W \setminus U)$. Graphs in this paper are undirected and finite. Let $G = (V, E)$ be a graph. A set $I \subseteq V$ is *independent* in G if the vertices in I are pairwise non-adjacent in G . We denote by $\alpha(G)$ the maximal size of an independent set of G . For a vertex $v \in V$ we denote by $N(v)$ the *neighborhood* of v , that is, the set of vertices that are adjacent to v in G . For $t \geq 1$ we denote by P_t the path graph on t vertices. A vertex of degree one of P_t is called *endvertex* and a vertex of degree two of P_t is called *internal vertex*. Given graph G , the graph \overline{G} is the *complement graph* of G and is obtained by replacing every edge of G by a non-edge en vice-versa.

A *claw* is a graph on four vertices that has one vertex of degree three (the *center*) and three vertices of degree one (the *leaves*). A *fork* is a graph obtained from a claw by subdividing one edge, that is, by replacing an edge by a path on three vertices. The *center* of a fork F is the vertex of F of degree 3. For $t \geq 1$, the *t-subdivision* of G is the graph obtained from G by replacing each edge uv of G with a copy of P_{t+2} and identifying its endvertices with u and v . We say that a *t-subdivision* of G is *even* if t is even.

Modular decomposition. A set $U \subseteq V$ of vertices is called *module* if for each vertex $v \in V \setminus U$, the vertex v is either adjacent to every vertex in U or v is non-adjacent to every vertex in U . A module $U \subseteq V$ is *non-trivial* if $U \neq \emptyset$ and $U \neq V$. By recursively dividing a connected graph into maximal modules we may compute in linear time its *modular decomposition* [MS99]. The modular decomposition of a graph has been introduced by Gallai in [Gal67]. Using this decomposition, many graph problems that are hard in general can be solved efficiently on certain classes of graphs [Möh85, MR84]. For instance, the modular decomposition is used in the polynomial-time algorithm for computing a maximum-weight independent set in a fork-free graph [LM08]. In the context of reconfiguration, the modular decomposition of cographs (that are P_4 -free graphs) has been used (implicitly) in order to obtain a polynomial-time algorithm for TOKEN JUMPING on cographs [Bon16]¹.

Independent Set Reconfiguration. In the following we define two adjacency relations on the independent sets of G that correspond to three natural reconfiguration steps on independent sets: *token sliding* (TS) and *token jumping* (TJ). Let $I, J \subseteq V$ be two independent sets of G . We say that I and J are TJ-adjacent ($I \leftrightarrow_{\text{TJ}} J$) if there is precisely one vertex in I that is not in J and vice versa. Furthermore, we say that I and J are TS-adjacent ($I \leftrightarrow_{\text{TS}} J$) if $I \leftrightarrow_{\text{TJ}} J$ and the vertex u in $I \setminus J$ and the vertex v in $J \setminus I$ are adjacent in G . For each rule $R \in \{\text{TS}, \text{TJ}, \}$ we say that I and J admit a *R-reconfiguration sequence* I_0, I_1, \dots, I_ℓ and write $I \rightsquigarrow_R J$ if $I = I_0$, $J = I_\ell$ and for $0 \leq i < \ell$ we have $I_i \leftrightarrow_R I_{i+1}$. If it is clear from the context we may omit the rule R . Given a graph G and two independent sets $I, J \subseteq V(G)$, the problem TOKEN SLIDING (resp., TOKEN JUMPING) asks whether I and J admit a TS-reconfiguration sequence (resp., TJ-reconfiguration sequence).

Note that TOKEN SLIDING and TOKEN JUMPING are equivalent on maximum independent sets since, if a vertex can jump on a non-adjacent vertex, the independent set is indeed not maximum.

3 A reconfiguration analogue of Alekseev’s theorem

In this section we prove Theorem 2, a reconfiguration analogue of Alekseev’s Theorem 1). In the following, let $G = (V, E)$ be a graph and G_t be the t -subdivision of G for some even $t \geq 1$. For any independent set I of G , we can construct in a canonical way an independent set I' of G_t by placing for each edge uv of G on the internal vertices of the copy of P_{t+2} in G_t that corresponds to uv exactly $t/2$ additional tokens. The independent set I' is called the *extension* of I . Alekseev proved the following:

¹Note that the algorithm is given in the TAR-model which is proven to be equivalent to the TJ-model for independent set reconfiguration.

Theorem 4 ([Ale82]). *The graph G has a maximum independent set of size k if and only if G_t has maximum independent set of size $k + t \cdot |E(G)|/2$.*

Theorem 4 implies immediately that for any maximum independent set I of G , the extension I' of I in G is also maximum. Our goal in this section is to adapt Theorem 4 to TOKEN SLIDING and TOKEN JUMPING as follows.

Theorem 5. *Let I and J be two maximum independent sets of G and let $R \in \{\text{TS}, \text{TJ}\}$. Then $I \rightsquigarrow_R J$ with respect to G if and only if $I' \rightsquigarrow_R J'$ with respect to G_t .*

The rest of this section is devoted to prove Theorem 5. Recall that TOKEN SLIDING and TOKEN JUMPING are PSPACE-complete in graphs of maximum degree three [IDH⁺11, HD05]. By considering subdivisions of G , we may assume that G has arbitrarily high girth. Therefore, just as Alekseev's Theorem 4 implies Theorem 1, it suffices to prove Theorem 5 to obtain Theorem 2. The remainder of this section is devoted to the proof of Theorem 5. Observe that by Theorem 4 the independent sets I' and J' of Theorem 5 are maximum independent sets of G_t . Recall that for maximum independent sets, TOKEN SLIDING and TOKEN JUMPING are equivalent², so it suffices to consider the TS version.

For an edge uv of G let $S_{uv} = \{s_{uv}^1, s_{uv}^2, \dots, s_{uv}^t\}$ be the set of internal vertices of the copy of P_{t+2} that replaces the edge uv in the construction of G_t . We assume that s_1 is adjacent to u and s_t is adjacent to v and that for $1 < i < t$, the vertex s_i is adjacent to s_{i+1} and s_{i-1} . A *left-move* on S_{uv} (assuming that we have an order between u and v) is a move of a token on s_i with $i > 1$ to s_{i-1} , provided that s_{i-1} is not adjacent to a token.

Claim 1. *Let uv be an edge of G and let I' be a maximum independent set of G_t . If $\{u, v\} \subseteq I'$ then $|I' \cap S_{uv}| = (t - 2)/2$ and $|I' \cap S_{uv}| = t/2$ otherwise.*

Proof. Starting from I' , apply left-moves to S_{uv} until no longer possible and let I'' be the resulting independent set. Since no left-move can be applied to S_{uv} , if $\{u, v\} \subseteq I''$ we have that for $1 \leq i < t$ even, there is a token of I'' on s_i . Since t is even and I' is maximum, there are precisely $(t - 2)/2$ tokens in $S_{uv} \cap I''$. Since we only moved tokens on S_{uv} to obtain I'' , the same must hold for I' . On the other hand, suppose that $\{u, v\} \not\subseteq I''$ and (without loss of generality) that $u \notin I''$. Since no left-move can be applied to S_{uv} , any token of I'' is on some s_i with $1 \leq i \leq t$ odd. Since t is even and I'' is maximum, there must be a token on each s_i for $1 \leq i < t$ odd and thus there are exactly $t/2$ tokens of I'' on S_{uv} , and the same holds for I' . \square

Let G_t be the t -extension of $G = (V, E)$. By abuse of notation, we will denote by V the subset of vertices of G_t corresponding to vertices of G .

Using Claim 1, we show that any two maximum independent sets of G_t that agree on V admit a TS-reconfiguration sequence, and hence also a TJ-reconfiguration sequence.

Claim 2. *Let I' and J' be two maximum independent sets of G_t such that $I' \cap V = J' \cap V$. Then $I' \rightsquigarrow_{\text{TS}} J'$.*

Proof. Let uv be an edge of G and let $T := I' \cap V = J' \cap V$. If $\{u, v\} \subseteq T$ we have $I' \cap S_{uv} = J' \cap S_{uv} = \{s_{uv}^2, s_{uv}^4, \dots, s_{uv}^{t-2}\}$ by Claim 1. If $|\{u, v\} \cap T| = 1$, suppose without loss of generality that $u \notin T$. Then Claim 1 ensures that $I' \cap S_{uv} = J' \cap S_{uv} = \{s_{uv}^1, s_{uv}^3, \dots, s_{uv}^{t-1}\}$. Finally, if $\{u, v\} \cap T = \emptyset$, Claim 1 ensures that $|T| = t/2$. By applying any possible valid left-move on S_{uv} for both I' and J' we obtain independent sets I'' and J'' , respectively, such that $I'' \cap S_{uv} = J'' \cap S_{uv} = \{s_{uv}^1, s_{uv}^3, \dots, s_{uv}^{t-1}\}$. Notice that $T = I'' \cap V = J'' \cap V$. It is then sufficient to repeat this process for any edge of G that satisfies $\{u, v\} \cap T = \emptyset$ to obtain a TS-reconfiguration sequence from I' to J' . \square

The following claim, which ensures that a single reconfiguration step in G can be adapted in G_t , will immediately imply the “if”-direction of Theorem 5. Indeed, it suffices to apply the following claim to each step of the transformation to get the conclusion.

²If a token can jump to a non-neighbor, then the independent set is indeed not maximum.

Claim 3. Let I_1, I_2 be two maximum independent sets of G such that $I_1 \leftrightarrow_{\text{TS}} I_2$ and let I'_1 and I'_2 be the extensions of I_1 and I_2 in G_t . Then $I'_1 \rightsquigarrow_{\text{TS}} I'_2$.

Proof. Since $I_1 \leftrightarrow_{\text{TS}} I_2$ let $e = uv$ be the edge such that $I_1 \Delta I_2 = \{u, v\}$ with $u \in I_1$ and $v \in I_2$. We show that there exists a reconfiguration sequence $I'_1 \rightsquigarrow_{\text{TS}} I'_2$ in G_t .

First note that, except for u , there is no token on w in I'_1 for every neighbor w of v distinct from u (since $I_2 = I_1 + v - u$ is an independent set). So by Claim 2, $I'' = (I'_1 \setminus S_{vw}) \cup \{s_{vw}^2, s_{vw}^4, \dots, s_{vw}^t\}$ is a maximum independent set of G_t that can be reached from I'_1 . So, for every $w \in N(v) \setminus u$, we can assume that there is no token on s_{vw}^1 . Let us denote by I''_1 the resulting independent set. We can slide in I''_1 the token s_{uv}^t of I''_1 to v . Finally, it suffices to slide the token from s_{uv}^i to s_{uv}^{i+1} for every i such that $s_{uv}^i \in I''_1$ (in decreasing order) and then slide the token from u to s_{uv}^1 in order to obtain I'_2 from I'_1 . We conclude that $I'_1 \rightsquigarrow_{\text{TS}} I'_2$. \square

Let us now prove the converse direction of Theorem 5. We start with the following lemma:

Claim 4. Let I' be a maximum independent set of G_t . Then there are no three vertices $u, v, w \in I' \cap V$, such that $uv, vw \in E(G)$.

Proof. Assume for a contradiction that there are three vertices $u, v, w \in I' \cap V$ such that $uv, vw \in E(G)$. By Claim 1 and 2, we can transform I' into an independent J such that $J \cap S_{uv} = \{s_{uv}^2, s_{uv}^4, \dots, s_{uv}^{t-2}\}$ and $J \cap S_{vw} = \{s_{vw}^3, s_{vw}^5, \dots, s_{vw}^{t-1}\}$. But then we can slide the token on v to s_{uv}^t and obtain an independent I'' such that s_{vw}^1 has no token in its neighborhood in G_t , which contradicts the maximality of I' . \square

A graph G' is an induced subgraph of G if there exists a subset of vertices $V' \subseteq V$ such that the graph with vertex V' where uv is an edge if and only if uv is an edge of G . Let I' be a maximum independent set of G_t . By the previous claim, the graph $G[I' \cap V]$ consists of isolated vertices and edges. Let $v(I')$ (resp, $e(I')$) denote the number of isolated vertices (resp., isolated edges) of $G[I' \cap V]$.

Claim 5. $v(I') + e(I') = \alpha(G)$.

Proof. Notice that since I' is a maximum independent set of G_t it has size $\alpha(G) + t \cdot |E(G)|/2$ by Theorem 4. Let uv be an edge of G . By Claim 1, we have $|S_{uv} \cap I'| = \frac{t-2}{2}$ if $\{u, v\} \subseteq I'$ and $|S_{uv} \cap I'| = \frac{t}{2}$ otherwise. It follows that

$$\begin{aligned} \alpha(G) + t \cdot |E(G)|/2 &= |I'| \\ &= v(I') + \left(\sum_{\substack{uv \in E(G): \\ |\{u, v\} \cap I'| \leq 1}} \frac{t}{2} \right) + 2e(I') + \left(\sum_{\substack{uv \in E(G) \\ : u, v \in I'}} \frac{t-2}{2} \right) \\ &= v(I') + e(I') + t \cdot |E(G)|/2 \end{aligned}$$

from which we conclude that $v(I') + e(I') = \alpha(G)$. \square

To complete the proof of Theorem 4, consider a TS-reconfiguration sequence $I'_1 \leftrightarrow_{\text{TS}} I'_2 \leftrightarrow_{\text{TS}} \dots \leftrightarrow_{\text{TS}} I'_\ell$ of maximum independent sets of G_t such that $I' = I'_1$ and $J' = I'_\ell$, where I' (resp., J') is the extension of I (resp. J) in G_t . Notice that $G[I' \cap V]$ and $G[J' \cap V]$ contain no edge. And by Claim 5, we can associate to each independent set I' an induced subgraph consisting of $\alpha(G)$ isolated vertices or edges. We obtain from I' a maximum independent set I_i of G by picking all isolated vertices and exactly one endpoint of each edge of $G[I'_i \cap V]$, say the first vertex with respect to an arbitrary fixed ordering of V . To conclude the proof it is enough to show that, for every $i \in \{1, 2, \dots, \ell - 1\}$, we have $I_i \rightsquigarrow_{\text{TS}} I_{i+1}$.

Claim 6. $|(I'_i \cap V) \Delta (I'_{i+1} \cap V)| \leq 1$.

Proof. Since $I'_i \leftrightarrow_{\text{TS}} I'_{i+1}$ there exists an edge xy such that $I'_{i+1} = (I'_i \setminus x) \cup y$. Note that at most one vertex in $x, y \in V$ since no edge of G_t has both endpoints in V . We distinguish three cases:

1. $x, y \notin V$. Then $\{x, y\} \subseteq S_{vw}$ for some edge vw of G and therefore $|(I'_i \cap V) \Delta (I'_{i+1} \cap V)| = 0 \leq 1$.
2. $x \in V$. Then $y \in S_{vw}$ for some edge vw of G and therefore $|(I'_i \cap V) \Delta (I'_{i+1} \cap V)| = 1$.
3. $y \in V$. Then $x \in S_{vw}$ for some edge vw of G and therefore $|(I'_i \cap V) \Delta (I'_{i+1} \cap V)| = 1$.

□

By claims 5 and 6, we just have to distinguish the following cases to complete the proof of the converse direction:

1. $e(I'_i) = e(I'_{i+1})$. In this case we have $I_i = I_{i+1}$ and we are done.
2. $e(I'_i) = e(I'_{i+1}) + 1$. Let uv be the edge of G that is not present in $G[I'_{i+1} \cap V]$ and assume without loss of generality that $u \in I_{i+1}$. It follows that I'_{i+1} is obtained from I'_i by sliding a token from v to S_{uv}^p . Then either $u \in I_i$, so we have $I_i = I_{i+1}$ or $v \in I_i$ and we have $I_i \leftrightarrow_{\text{TS}} I_{i+1}$.
3. $e(I'_i) = e(I'_{i+1}) - 1$. The same argument can be applied by exchanging I'_i and I'_{i+1} .

So, for every $1 \leq i < \ell$, we have $I_i \leftrightarrow_{\text{TS}} I_{i+1}$. And then $I \leftrightarrow_{\text{TS}} J$ which completes the proof.

4 Irrelevant vertices in fork free graphs

In the following, let $G = (V, E)$ be a fork-free graph and I and J be two independent sets of G . We will identify vertices of G that are irrelevant in any transformation from I to J in the sense that they cannot contain a token. We show that vertices adjacent to at least two tokens are irrelevant if they satisfy a certain neighborhood-condition. As a consequence, all vertices adjacent to at least three tokens are useless and can be safely deleted. We say that a set $X \subseteq V$ of vertices is *locally blocked* (with respect to I) if any vertex in X is adjacent to at least two tokens in I . We are interested in locally blocked vertices that remain locally blocked under any token move. Let $X \subseteq V$ be a locally blocked subset of vertices. We say that the set $B_X := N(X) \cap I$ is *X -blocking*: no token can slide from B_X to X , since there is another token in B_X blocking it.

Let $u \in V$ and $X \subseteq V - u$. A vertex v of $G - (X + u)$ is an *X -twin* of u if uv is an edge and $N(v) \cap X = N(u) \cap X$. A set $X \subseteq V$ of vertices *permanently blocked* if it is locally blocked (with respect to I) and for each independent set I' such that $I \leftrightarrow_{\text{TS}} I'$ and for each vertex u in the blocking set $B_X = N(X) \cap I'$, any I' -free neighbor v of u is an X -twin of u . (It means that if we move, in I' a token in the neighborhood of X from u to another vertex v then the neighborhood in u and v in X are the same). We first introduce the following reduction rule, which states that permanently blocked vertices can be safely removed from the graph.

Reduction Rule Z Let $X \subseteq V$ be permanently blocked. Return $(G - X, I, J)$ if $X \cap J = \emptyset$ and a trivial No-instance otherwise.

A reduction rule is *safe* if applying it yields an equivalent instance.

Lemma 6. *Reduction Rule Z is safe.*

Proof. First notice that for any independent set I' such that $I \leftrightarrow_{\text{TS}} I'$, the set X is locally blocked with respect to I' , since whenever a token $u \in B_X$ moves to an I' -free neighbor v then v is an X -neighbor of u . In particular, no independent set reachable from I contains a token in X .

If $J \cap X \neq \emptyset$ then (G, I, J) is a No-instance, since for any independent set reachable from I , the set X is locally blocked. Suppose that $J \cap X = \emptyset$ and that $I \leftrightarrow_{\text{TS}} J$ with respect to G . Since no independent set I' reachable from I in G contains a token in X , we have $I \leftrightarrow_{\text{TS}} J$ with respect to G if and only if $I \leftrightarrow_{\text{TS}} J$ with respect to $G - X$. □

At this point it is not quite clear how to use Reduction Rule Z algorithmically. We will later prove that certain subsets of vertices are permanently blocked (and can be found in polynomial time). For now, we show that any vertex of G adjacent to at least three tokens of I is permanently blocked and can thus be safely deleted according to Lemma 6.

Lemma 7. *Let $c \in V$ be a vertex that is adjacent to exactly $k \geq 3$ tokens in I and let I' be an independent set such that $I' \rightsquigarrow_{\text{TS}} I$. Then c is adjacent to exactly k tokens of I' .*

Proof. Let J' be an independent set such that $I \rightsquigarrow_{\text{TS}} J'$. Suppose for a contradiction that c is adjacent to $k' \neq k$ tokens of J' . In a reconfiguration sequence from I to J' , let I' be the first independent set such that c is adjacent to a number of tokens of I' different from k . Suppose first that $|N(c) \cap I'| < k$. Let $x \in N(c)$ be the token that moved to some neighbor x' to obtain I' . Since $N(c)$ contains $k \geq 3$ tokens in any independent set preceding I' in the sequence, we have that I' contains at least two tokens $y, z \in N(c)$ different from x . Since $x', y, z \in I'$, we have that x' must be adjacent to c , otherwise $\{x', x, c, y, z\}$ induces a fork. But then $|N(c) \cap I'| = k$, a contradiction. Now suppose that $|N(c) \cap I'| > k$. Then some token $x' \notin N(c)$ moved to a vertex $x \in N(c)$. Let $y, z \in N(c) \cap I'$ be two tokens different from x . But then $\{x', x, c, y, z\}$ induces a fork, a contradiction. \square

Lemma 7 immediately implies the following sufficient condition for a vertex to be permanently blocked.

Observation 1. *Let $c \in V$ be a vertex that is adjacent to at least three tokens in I . Then c is permanently blocked.*

From the safeness of Reduction Rule Z and the previous observation we conclude that the following Reduction Rule is safe.

Reduction Rule A. Let $c \in V$ such that c is adjacent to at least three tokens in I . Return $(G - c, I, J)$ if $c \notin J$, otherwise return a trivial No-instance.

We say that G is I -reduced if Reduction Rule A is not applicable with respect to G and I . That is, G contains no vertex that is adjacent to at least three I -tokens. Notice that by applying Reduction Rule A exhaustively, we may obtain from G in polynomial time a fork-free graph G' that is I -reduced and J -reduced. The resulting equivalent instance is either (G', I, J) or a trivial No-instance. The following observations are consequences of Lemma 7.

Observation 2. *If G is I -reduced then G is I' -reduced for every independent set I' such that $I \rightsquigarrow_{\text{TS}} I'$.*

Observation 3. *Let G be I -reduced and let I' be an independent set such that $I \rightsquigarrow_{\text{TS}} I'$. Then for any vertex v of G we have $|N(v) \cap I'| \leq 2$.*

5 Maximum independent set reconfiguration in fork-free graphs

As a warm-up, let us prove the following theorem.

Theorem 8. *TOKEN SLIDING restricted to maximum independent sets admits a polynomial-time algorithm.*

To do so, we provide a simple reduction of Maximum Independent Set Reconfiguration in fork-free graphs to the same problem in claw-free graphs, which can be decided in polynomial time [BKW14]. In the following, let $G = (V, E)$ be a fork-free graph and I and J be two maximum independent sets of G . Note that it suffices to consider the token sliding rule only since I and J are maximum. By Lemma 6 and Observation 1 we may assume that G is both I -reduced and J -reduced. Therefore, G contains no induced claw that contains three tokens of I . Since I is maximum, the following stronger property holds.

Lemma 9. *Let $\{c, x, y, z\}$ induce a claw in G with center c . Then $|\{x, y, z\} \cap I| = 1$.*

Proof. Since I is maximum and G is I -reduced, we have $1 \leq |\{c, x, y, z\} \cap I| \leq 2$. Assume for a contradiction that $|\{c, x, y, z\} \cap I| = 2$ and let without loss of generality $\{c, x, y, z\} \cap I = \{x, y\}$.

Since I is maximum, the vertex z must have a neighbor t in I and since c has at most two neighbors in I , we have $t \notin N(c)$. It follows that $\{c, x, y, z, t\}$ induces a fork in G , contradiction.

So it remains to show that $c \notin I$. Assume for a contradiction that $c \in I$. Since I is maximum, there are at least three tokens in the neighborhood of $\{x, y, z\}$: otherwise we add $\{x, y, z\}$ to I and delete $N(\{x, y, z\})$ from I to obtain a larger independent set. Let $u, v \in I$ be distinct from c in the neighborhood of $\{x, y, z\}$. If u (resp., v) is adjacent to only one vertex in $\{x, y, z\}$ then $\{c, x, y, z, u\}$ (resp., $\{c, x, y, z, v\}$) induces a fork. Therefore, the vertices u and v have a common neighbor in $\{x, y, z\}$, say x . But then x has three neighbors in I , which contradicts our assumption that G is I -reduced. \square

By the previous lemma, no maximum independent set of G has a token on the center of any induced claw of G . Therefore, the following reduction rule is safe.

Reduction Rule MIS. Let $\{c, x, y, z\}$ induce a claw in G with center c . Return $(G - c, I, J)$.

Proof of Theorem 8. The induced claws of G can be enumerated in polynomial time (by brute-force), so Reduction Rule MIS can be applied exhaustively in polynomial time. We obtain an equivalent instance (G', I, J) of TOKEN SLIDING such that the graph G' is claw-free. It remains to apply the algorithm from [BKW14] on (G', I, J) to decide the initial instance (G, I, J) . Notice that any reconfiguration sequence from I to J in G' is a reconfiguration sequence from I to J in G' . \square

6 Non-maximum Token Sliding in fork-free graphs

We provide a polynomial-time algorithm for TOKEN SLIDING on fork-free graphs, generalizing the algorithm for claw-free graphs given in [BKW14]. In the following, let $G = (V, E)$ be a fork-free graph and let $I, J \subseteq V$ be two independent sets of G of the same size. If I and J are maximum, we may apply Theorem 8 and are done. Otherwise, we show that we may assume that each connected component of G is prime (Section 6.1). The algorithm then proceeds as follows. We consider the symmetric difference of the two independent sets and notice that each connected component of $G[I \Delta J]$ is a path or an (even) cycle, each of which can be treated separately. Paths are easy to deal with while cycles require more work. In order to deal with the cycles we use an idea that has been used for MATCHING RECONFIGURATION in [IDH⁺11]: we push tokens along a path from the cycle to a vertex that is I -free (that is, it has no neighbor in I), which allows us to move the remaining tokens on the cycle to their target positions. Showing that pushing the tokens along such a path is possible is non-trivial and is discussed in Section 6.2. It turns out that if we get stuck at any point then there is a permanently blocked vertex that can be removed from the graph. The complete algorithm is presented in Section 6.3.

6.1 Reduction to prime graphs

We show that we may assume without loss of generality that the input graph G is prime with respect to the modular decomposition. First, we observe that if a module of a graph contains at least two tokens then the instance induced by the vertices of this module can be treated independently since no token will be able to leave the module at any step.

Observation 4. *Let M be a module of G such that $\ell := |M \cap I|$ and I' be such that $I \rightsquigarrow_{\text{TS}} I'$. If $\ell \geq 2$, we have $|M \cap I'| = \ell$ and if $\ell \leq 1$ we have $|M \cap I'| \leq 1$.*

By Observation 4, if a non-trivial module M contains at least two tokens of I and J , respectively, then all the vertices in the neighborhood of M can be deleted. Otherwise, if $G[M]$ is connected and contains at most one token of I and J , respectively, then all vertices of M are equivalent with respect to token sliding, so the vertices of M can be identified. However, the situation is slightly more complicated if M is not connected since a token might have to leave the module to reach its target position. In that case we show that either

a token in M can move to a different connected component of $G[M]$ in exactly two moves, or it can never leave its connected component.

Lemma 10. *Assume that G is I -reduced, let M be a non-trivial module of G , and let C_1, C_2, \dots, C_ℓ be the connected components of $G[M]$. Suppose that $\ell \geq 2$ and $M \cap I = \{v\} \subseteq V(C_1)$. Then exactly one of the following holds:*

- *The vertex v has a neighbor $c \in V \setminus M$ such that $N(c) \cap I = \{v\}$.*
- *$N(M) \setminus M$ is permanently blocked. In particular, we have $|I' \cap V(C_1)| = 1$ for every independent set I' of G such that $I \rightsquigarrow_{\text{TS}} I'$.*

Proof. If v has a neighbor $c \in V(G) \setminus M$ such that $N(c) \cap I = \{v\}$ then we can slide the token on v to c to obtain an independent set $I' \leftrightarrow_{\text{TS}} I$ that satisfies $|I' \cap C_1| = 0$.

If v has no such neighbor, let us prove that $N(M) \setminus M$ is permanently blocked. Since G is I -reduced and by the assumptions on v , each vertex u of $N(M)$ has exactly two neighbors in I and in any independent set reachable from I contains at most two neighbors of u . Assume for a contradiction that there exists a vertex $u \in N(M)$ and an independent set I' such that $I \rightsquigarrow_{\text{TS}} I'$ and $|N(u) \cap I'| = 1$. Chose I' such that the reconfiguration sequence from I to I' is shortest and let J' be the independent set before I' in the sequence. By Observation 4, we have $|J' \cap N(u)| = 1$. Let $z \in (N(u) \cap J') \setminus M$. To obtain I' from J' , the token on z slides to $z' \notin N(u)$. Let $a, b \in N(u) \cap M$ such that a and b are in different components of M . But then $\{u, a, b, z, z'\}$ induces a fork, a contradiction. \square

Suppose that M is a module of G such that that $M \cap I$ and $M \cap J$ have size at most one. We denote by $(G, I, J)_{/M}$ the instance of TOKEN SLIDING where M has been replaced by a single vertex m and the independent set I (resp., J) contains m if $I \cap M$ (resp., $J \cap M$) has size one³. We say that an instance of (G, I, J) of TOKEN SLIDING is *balanced* if for every non-trivial module M of G satisfying $|M \cap I| = |M \cap J| = 1$, the unique vertex in $M \cap I$ and the unique vertex in $M \cap J$ are in the same connected component of $G[M]$. By applying the next reduction rule exhaustively, we obtain either a trivial No-instance or a balanced instance.

Reduction Rule B. Assume there is a non-trivial module M of G such that $M \cap I = \{u\}$, $M \cap J = \{v\}$ and u and v are in two distinct components of $G[M]$. If there is a vertex $c \in V \setminus M$ such that $N(c) \cap I = \{u\}$, return $(G, I, J)_{/M}$. Otherwise, return a trivial No-instance.

Lemma 11. *Reduction Rule B is safe.*

Proof. Let M be non-trivial module of G such that $M \cap I = \{u\}$, $M \cap J = \{v\}$ and u and v are in two distinct components of $G[M]$. If there is a vertex $c \in V \setminus M$ such that $N(c) \cap I = \{u\}$, we may move the token on u to c and then to v , to obtain an independent set $I' \rightsquigarrow_{\text{TS}} I$. Since any vertex of M has the same neighborhood in $V \setminus M$, the module M can contain at most one token in any independent set reachable from I . Since $v \in I' \cap J$, we may safely identify all vertices in M .

On the other hand, assume that there is no vertex $c \in V \setminus M$ such that $N(c) \cap I = \{u\}$. Let C be the vertex set of the connected component of $G[M]$ containing u . By Lemma 10, for any independent set I' such that $I' \rightsquigarrow_{\text{TS}} I$, we have $|I' \cap C| = 1$, but $|J \cap C| = 0$. Therefore (G, I, J) is a No-instance. \square

Assume that the instance (G, I, J) of TOKEN SLIDING is balanced. Then Observation 4 tells us that if the input graph contains a non-trivial module the graph can be either reduced or split into connected components that can be treated independently. We now formalize this observation by introducing further reduction rules that are then applied exhaustively under the assumption that Reduction Rule B is not applicable. Each connected component of the resulting graph will be prime.

Reduction Rule D Let M be a non-trivial module of G such that $|M \cap I| \leq 1$. If $|M \cap J| \leq 1$ then return $(G, I, J)_{/M}$, otherwise return a trivial No-instance.

³We still denote by I and J the independent sets by abuse of notations

Reduction Rule E Let M be a non-trivial module of G such that $|M \cap I| \geq 2$. If $|M \cap J| \neq |M \cap I|$ return a trivial No-instance, otherwise return $(G - N(M), I, J)$.

Observe that if G is prime then none of the reduction rules B, D, E are applicable.

Lemma 12. *Suppose that Reduction Rule B is not applicable. Then reduction rules D and E are safe.*

Proof. Suppose first that Reduction Rule D is applicable and let $\{u\} = M \cap I$. By Observation 4 and Lemma 10, if $|M \cap J| > 1$ then (G, I, J) is a No-instance; so assume that $\{v\} = M \cap J$. Given that Reduction Rule B is not applicable, the tokens on u and v are in the same connected component of M , so we may slide the token on u to v . Notice that in any independent set reachable from I and J , the module M contains at most one token. Therefore, we may contract the module M to a single vertex to obtain the instance $(G, I, J)_{/M}$ which is equivalent to (I, J, M) .

Now suppose that Reduction Rule E is applicable. By Observation 4 and Lemma 10, if $I \cap M$ and $J \cap M$ do not have the same size then I and J do not admit a TS-reconfiguration sequence. Otherwise, by Lemma 10, no token of M can leave the module, so we can treat $G[M]$ and $G[V \setminus N[M]]$ independently. \square

We now prove we may consider without loss of generality instances of TOKEN SLIDING on fork-free graphs that are prime.

Theorem 13. *Let (G, I, J) be an instance of TOKEN SLIDING such that G is fork-free. We may compute in polynomial time from (G, I, J) an equivalent instance (G', I', J') of TOKEN SLIDING on a fork-free graph G' that is I' -reduced and J' -reduced, such that each connected component of G' is prime.*

Proof. We apply Reductions Rules A,B,D,E as long as we can (in this order) to obtain in polynomial time the instance (G', I', J') of TOKEN SLIDING. Assume for contradiction that there is some connected component C of G' contains a non-trivial module M . If M contains more than one token then Reduction Rule E can be applied. Otherwise, Reduction Rule D can be applied, a contradiction. Furthermore, (G', I', J') and (G, I, J) are equivalent by the safeness of the reduction rules (see Observation 1 and Lemmas 11 and 12). Finally, G' is fork-free, since fork-free graphs are closed under taking induced subgraphs and contracting modules. \square

6.2 Reaching I-free vertices

In the following, let $G = (V, E)$ be a fork-free graph and let I be an independent set of G such that G is I -reduced (Reduction Rule A is not applicable). Our goal in this section is to prove the following theorem.

Theorem 14. *Let $v \in I$ and let u be an I -free vertex of G . Then $J := I - v + u$ is an independent set of G . If G is prime then there is a polynomial-time algorithm that computes a reconfiguration sequence $I \leftrightarrow_{\text{TS}} J$ or finds a permanently blocked vertex of G .*

First, we provide the intuition behind the proof. Consider a shortest path P from u to v in G : Either we can directly slide step by step a token along the path from v to u , or some other tokens $a_1, a_2, \dots, a_t \in I$ are in the (closed) neighborhood of the path P and therefore prevent v from moving to u . Let a_1, a_2, \dots, a_t be ordered by increasing distance to v . The idea is to move a_t to v and then for i for $t - 1$ down to 1, move a_i to a_{i+1} . Finally, u may move to a_1 . The resulting independent set is precisely $J = I - v + u$, so $I \leftrightarrow_{\text{TS}} J$. However, two tokens in $\{a_1, \dots, a_t\}$ might block each other (if both have the same neighbors on the path). We first prove that if P has length at least three then this is impossible. We use a characterization of claws in prime graphs due to Brandstädt et al. [BHV04] in order to handle the case where P has length two. In this case we may not be able to directly reach J , but then some vertex of G is permanently blocked and can therefore be deleted safely by Lemma 6.

Let $u, v \in V$ and let $P = u_1, \dots, u_\ell$ be a $u - v$ path in G . The *leftmost* (resp. *rightmost*) neighbor of $x \in N(V(P))$ on P is the vertex of $V(P) \cap N(x)$ with the lowest (resp. highest) index.

Lemma 15. *Let u be an I -free vertex of G and let $v \in I$. Let P be a shortest path from u to v in G . If P has length at least three then the neighbor of u in P has at most one neighbor in I .*




Figure 2: An illustration in the two cases of the proof of Lemma 15. The dashed lines on the left indicate non-edges and the red squares indicate tokens.

Proof. Let $P = u_1, \dots, u_\ell$ be a shortest $u - v$ path in G such that $\ell \geq 4$. Suppose for a contradiction that u_2 has two neighbors in I (it cannot have more than two since G is I -reduced). There are two cases to consider.

Case 1: $u_3 \in I$ and u_2 has a neighbor $w \in N(V(P))$ in I (see Figure 2 (left)). Notice that u, u_3 , and w are pairwise non-adjacent since u is I -free and $u_3, w \in I$. Since $\ell \geq 4$, the edge wu_4 exists, otherwise $\{u_2, u_1, u_3, w, u_4\}$ induces a fork. Since $v, w \in I$, we have $u_4 \neq v$, so $\ell \geq 5$. If $u_5 = v$, then u_4 has three neighbors in I , a contradiction to our assumption that G is I -reduced, so $\ell \geq 6$. Then either wu_5 or wu_6 is an edge of G , so P cannot be a shortest $u - v$ path, a contradiction.

Case 2: u_2 has two neighbors $w, x \in I$ that are not on P (see Figure 2 (right)). Let u_i, u_j be the rightmost neighbors of w and x on P respectively. First assume that $i = j$. Then $u_j \neq v$ since $v \in I$ and $u_{j+1} \neq v$ since G is I -reduced, so $\ell \geq i + 2$. But then $\{u_i, w, x, u_{i+1}, u_{i+2}\}$ induces a fork, a contradiction. We therefore assume without loss of generality that $j > i \geq 2$. If $j > 3$ then $\{u_2, u_1, w, x, u_j\}$ induces a fork. If $j = 3$ then w has no other neighbor on P but u_2 (since w is non-adjacent to u and u_i). Then we can slide the token on x to u_3 : if not, then another token on a vertex z must prevent this move. But then $\{u_2, u_1, w, u_3, z\}$ induces a fork. We may therefore slide the token on x to u_3 and the previous case applies. \square

Lemma 16. *Let u be an I -free vertex of G , $v \in I$, and let P a be shortest path $u - v$ path in G . If P has length at least three then no two vertices of $N[V(P)] \cap I$ can have the same leftmost neighbor on P .*

Proof. Let $P = u_1, u_2, \dots, u_\ell$, such that $\ell \geq 4$. Suppose for a contradiction that there exists $w, x \in N[V(P)] \cap I$ that have the same leftmost neighbor $u_i \in P$. By Lemma 15, we have $i > 2$. Notice that u_{i-2} and u_i are non-adjacent. Since u_i is the leftmost neighbor of w and x , we have that $\{w, x, u_i, u_{i-1}, u_{i-2}\}$ induces a fork. \square

The two previous lemmas allow us to deal with the case where a shortest path from $v \in I$ to $u \notin N[I]$ has length at least three. The following theorem, due to Brandstädt et al., will be useful in order to deal with the case where a shortest $u - v$ path has length two:

Theorem 17 ([BHV04]). *Let G be a fork-free prime graph which contains a claw induced by the vertices $\{c, u, v, w\}$. Then one of the graphs H_1, \dots, H_5 shown in Figure 3 is an induced subgraph of G .*

We use this characterization of the prime extensions of a claw in a fork-free graph to prove that if we are given a claw containing two tokens, we can either “rotate” the tokens around the center of the claw locally without moving any other token or that the center of the claw is permanently blocked and can thus be safely deleted from the graph by Lemma 6.

Lemma 18. *Let I be an independent set of a fork-free prime graph G such that G is I -reduced. Let $\{c, u, v, w\}$ induce a claw in G with center c such that $\{u, v\} \subseteq I$. Then exactly one of the following holds:*

1. *There exists a reconfiguration sequence from I to $I - v + w$ and from I to $I - u + v$ such that only the tokens initially on u and v are moved and such a sequence can be found in polynomial time, or*




Figure 3: The fork-free prime expansions of a claw. The claws considered in Lemma 18 are marked by thick edges. The graph H_5 is shown three times, once for each of its induced claws.

2. the vertex c is permanently blocked.

Proof. We first consider the case where the induced subgraph of G containing the claw $\{c, u, v, w\}$ is H_5 shown in Figure 3. Notice that the graphs H_1, \dots, H_4 are subgraphs of H_5 . In the following we will use two arguments: First, we show that certain edges must exist since otherwise there is an induced fork. Second, we show that certain edges cannot be present since Reduction Rule A is not applicable. In almost all cases, if these arguments apply to H_5 they also apply to H_1, \dots, H_4 , so it suffices to treat H_5 . Finally, there is an automorphism between the first, the second and the third claw in H_5 shown in Figure 3. That is, the second and the third claw can be obtained from the first by renaming vertices (preserving adjacencies and non-adjacencies), so exactly the same arguments apply as for the first claw of H_5 . Let I be an independent set of G containing the vertices u and v . The lemma follows from the following claims.

Claim 7. *We can move the token v of I to w via y .*

Proof. We show that neither w nor y have a token in their neighborhood that is preventing us from moving the token on v to w . Suppose that y has a neighbor y' containing a token. Then c and x are non-adjacent to y' since Reduction Rule A is not applicable. Furthermore, w and y' are non-adjacent since otherwise $\{y'w'c'u'v\}$ induces a fork. But then $\{w, y, y', x, u\}$ induces a fork if xy is present (H_2, H_4, H_5) and $\{w, y, y', v, x\}$ induces a fork otherwise (H_1 and H_3). Therefore y' does not exist. Now suppose that w has a neighbor w' that contains a token. Then c and w' are non-adjacent since Reduction Rule A is not applicable. But then $\{w', w, c, u, v\}$ induces a fork, so the token w' does not exist. Therefore, we may move the token on v to w via y . \diamond

Claim 8. *Either we can move the token w of $I - v + w$ to v via y or the vertex c is permanently blocked.*

Proof. We try to move the token on w to v via y . The vertex v has no neighbor v' containing a token if two tokens are on u and w , since otherwise $\{u, c, w, v, v'\}$ induces a fork. Therefore, we can move w to v via y unless y has a neighbor y' with a token on it. In this case c is non-adjacent to y' since Reduction Rule A is not applicable. If x is adjacent to y (H_2, H_4, H_5) then x is also adjacent to y' , since otherwise $\{u, x, y, w, y'\}$ induces a fork. Otherwise, if x is non-adjacent to y (H_1 and H_3) then x is adjacent to y' , otherwise $\{x, v, y, w, y'\}$ induces a fork. It follows that $X = \{c, x, y\}$ is locally blocked, i.e., each vertex of X has precisely two tokens in $B_X = \{u, w, y'\}$ in its neighborhood. In order to apply Lemma 6, we show that if we slide any token $d \in B_X$ to a vertex d' then d' is an X -twin of d .




Figure 4: An illustration of the situation in the proof of Theorem 14. The red squares represent the initial positions of the tokens. The dashed lines represent non-edges.

- Suppose that $u \in B_X$ slides to a neighbor u' . Then y and u' are non-adjacent by Observation 2 and v and u' are non-adjacent since otherwise $\{w, y, z, v, u'\}$ induces a fork. Furthermore, c and u' are adjacent since otherwise $\{v, c, w, u, u'\}$ induces a fork. Finally, x and u' are adjacent since otherwise $\{u', u, x, v, y'\}$ induces a fork. Therefore, the vertex u' is an X -neighbor of u .
- Suppose that $w \in B_X$ slides to a neighbor w' . Then x and w' are non-adjacent (otherwise $\{u, x, y', w', w\}$ induces a fork) and v and w' are non-adjacent (otherwise $\{u, x, z, v, w'\}$ induces a fork). Then c and w' are adjacent (otherwise $\{v, c, u, w, w'\}$ induces a fork) and w' and y are adjacent (otherwise $\{u, c, w, w', y\}$ induces a fork). Therefore, the vertex w' is an X -neighbor of w .
- Suppose that $y' \in B_X$ slides to a neighbor y'' . Then v and y'' are non-adjacent (otherwise $\{u, c, w, v, y''\}$ induces a fork) and c and y'' are non-adjacent (by Observation 2). Furthermore, y and y'' are adjacent (otherwise $\{w, y, v, y', y''\}$ induces a fork) and x and y'' are adjacent (otherwise $\{u, x, v, y', y''\}$ induces a fork). Therefore, the vertex y'' is an X -neighbor of w .

◇

The next two claims follow from the previous ones by noticing that renaming the vertices u and w as well as x and y is an automorphism of H_5 .

Claim 9. *We can move the token v of $I - u + w$ to u via x .*

Claim 10. *Either we can move the token on u of $I - v + w$ to v via x or the vertex c is permanently blocked.*

□

We now have all the ingredients to prove Theorem 14.

Proof of Theorem 14. Let I be an independent set of G such that Reduction Rule A is not applicable and let u be a vertex of G that is I -free. Let $v \in I$ and $J := I - v + u$. Since u is I -free, J is an independent set of G . We show that $I \rightsquigarrow_{\text{TS}} J$. Let $P := u_1, \dots, u_\ell$ be a shortest u - v path in G . Since u is I -free, we have $\ell \geq 3$. If $\ell = 3$ then either we can directly slide from v to u through u_2 or there is a token on $z \in N(u_2)$. In this case, the vertices $\{u_2, u, v, z\}$ induce a claw and by Lemma 18, we can either “rotate” the tokens on the claw to reach J , or $u = u_1$ is permanently blocked. So we may assume that $\ell \geq 4$. By Lemma 16, the leftmost neighbors on P of the I -tokens in the neighborhood of P are distinct. Let $\{a_1, \dots, a_t\} := N[P] \cap I$ with respective leftmost neighbors u_{i_1}, \dots, u_{i_t} on P , such that $i_j < i_{j+1}$ for all j . We show by induction on i that there exists a sequence of slides that moves the token initially on a_{i+1} to a_i for $1 < i < t$ in increasing order, where $a_0 := u$.

Case $i=1$. Slide the token on a_1 to a_0 via u_{i_1} and then slide from u_{i_1} to u along P . This is possible since the other tokens on $N[P]$ lie on a_2, \dots, a_t , which have no neighbors in u_0, \dots, u_{i_1} by the definition of leftmost neighbors.

Induction step. Let $1 < j < t$ and assume that the tokens initially on a_1, \dots, a_j have moved respectively to a_0, \dots, a_{j-1} . We show that the token on a_{j+1} can slide to a_j . The tokens in $N[P]$ currently are on $\{a_0, \dots, a_t\} \setminus \{a_j\}$.

We first slide the token on a_{j+1} to $u_{i_{j+1}}$. By the definition of leftmost neighbors, none of a_{j+2}, \dots, a_t is adjacent to $u_{i_{j+1}}$. Furthermore, none of a_0, \dots, a_{j-2} is adjacent to $u_{i_{j+1}}$ since P is a shortest u - v path. Therefore, the only token that may prevent us from sliding a_{j+1} to $u_{i_{j+1}}$ is on a_{j-1} , namely if the edge $a_{j-1}u_{i_{j+1}}$ is present. Assume for a contradiction that this is the case. Then $u_{i_{j-1}}$ is at distance two from $u_{i_{j+1}}$ in P since P is a shortest path. By Lemma 16, the leftmost neighbors of a_1, \dots, a_t are distinct, so $u_{i_{j-1}}, u_{i_j}, u_{i_{j+1}}$ are three consecutive vertices of P . See Figure 4 for an illustration. Since u_{i_j} is a leftmost neighbor of a_j , we have that a_j and $u_{i_{j-1}}$ are non-adjacent. Since a_{j-1} and a_{j+1} are adjacent to $u_{i_{j+1}}$ and Reduction Rule A is not applicable ($a_{j-1}, a_j, a_{j+1} \in I$), we have that a_j is non-adjacent to $u_{i_{j+1}}$. But then $\{u_{i_{j-1}}, u_{i_j}, a_j, u_{i_{j+1}}, a_{j+1}\}$ induces a fork, so a_{j-1} is non-adjacent to $u_{i_{j+1}}$. We may therefore slide the token on a_{j+1} to $u_{i_{j+1}}$.

If $a_{i_{j+1}}$ is adjacent to a_j , we can slide the token on $a_{i_{j+1}}$ to its destination u_j . Otherwise, the only obstruction that prevents us from moving the token on $a_{i_{j+1}}$ to $a_{i_{j+1}-1}$ is the presence of the edge $a_{j-1}a_{i_{j+1}-1}$. But then u_{i_j}, u_j , and $u_{i_{j+1}-1} = u_{i_{j+1}}$ are three consecutive vertices of P since P is a shortest path. Then $u_{i_{j+1}}$ is adjacent to a_j , since otherwise $\{u_{i_{j-1}}, u_{i_j}, a_j, u_{i_{j+1}}, u_{i_{j+1}}\}$ induces a fork. But then $\{a_{j-1}, u_{i_j}, a_j, u_{i_{j+1}}, u_{i_j}, a_{j+1}\}$ induces a fork. Therefore, we can slide the token on $u_{i_{j+1}} = u_{i_{j+2}}$ to $u_{i_{j+1}}$. To conclude, observe that $u_{i_{j+1}}$ is adjacent to a_j , since otherwise $\{u_{i_{j-1}}, u_{i_j}, a_j, u_{i_{j+1}}, u_{i_{j+1}}\}$ induces a fork. Therefore, we can slide the token on $u_{i_{j+2}}$ to its destination a_j .

Therefore, $I \rightsquigarrow_{\text{TS}} I - v + u$ or G contains a permanently blocked vertex. In order to reach $I - v + u$ from I , we move tokens along a shortest u - v path P to their destination as described above, which can be done in polynomial time. \square

6.3 A polynomial-time algorithm for Token Sliding in fork-free graphs

Using the reductions from the previous sections we provide a polynomial-time algorithm for TOKEN SLIDING in fork-free graphs. A *complex* is a complete bipartite graph minus a matching. We recall the following characterization of bipartite fork-free graphs due to Alekseev:

Theorem 19 ([Ale04]). *A bipartite and connected fork-free graph G is either a path, a cycle or a complex.*

In the following, let G be a fork-free graph and I and J be two independent sets of G . Since the symmetric difference of two independent sets induces a bipartite graph, $I\Delta J$ is a collection of disjoint paths, cycles, complex and isolated vertices. The following Lemma shows that we can reduce to the case where it does not contain any complex that is not a cycle or a path:

Lemma 20. *If G is I -reduced and J -reduced then $I\Delta J$ is a collection of disjoint paths, cycles and isolated vertices.*

Proof. Since Reduction Rule A is not applicable, each vertex of I has at most two neighbors in J and vice-versa. Therefore, each connected component of $G[I\Delta J]$ is a cycle or a path. \square

Let C be a non-trivial connected component of $G[I\Delta J]$. Let $I' := (I \setminus V(G)) \cup (J \cap V(C))$. We say that C can be *resolved* if $I \rightsquigarrow_{\text{TS}} I'$. That is, C can be resolved if there exists a reconfiguration sequence that replaces on C the tokens of I with the tokens of J . By abuse of notation, we say that an isolated vertex u of $G[I \setminus J]$ can be resolved if there exists a reconfiguration sequence that slides u to an isolated vertex of $G[J \setminus I]$. Note that if C is a path it can be resolved simply by sliding tokens one after the other along the path. If C is a cycle we proceed as follows. If there exists a free vertex u with respect to the current independent set I , we pick an arbitrary token on $C \cap I$ and slide it to v or find a permanently blocked vertex according to Theorem 14. In the former case we slide the remaining tokens on $C \cap I$ to $C \cap J$ just as for paths and then slide the token on v back to C (again according to Theorem 14). We can suppose that I

is not maximum since otherwise the reduction to claw-free graphs of Section 5 applies. If I is maximal, we show next that we can always “free” a vertex to resolve a cycle as explained before.

We first need a few definitions. Let G be a graph and A be an independent set of G . An independent set X of $G - A$ is called *magnifier of A* if the independent set $(A \cup X) \setminus (N(X) \cap A)$ is strictly larger than A and the corresponding subgraph $G[X \cup (N(X) \cap A)]$ is called *A -augmenting graph* [Ale04]. Notice that augmenting graphs are bipartite. Augmenting graphs generalize augmenting paths and are important for computing maximum independent sets.

Lemma 21. *If G is I -reduced and I is maximal but not maximum then there exists an I -augmenting path.*

Proof. Since I is not maximum, there exists an I -augmenting graph H and since I is maximal, H has at least two vertices. Since G is fork-free, Theorem 19 ensures that H is either a path or a complex. Since furthermore Reduction Rule A is not applicable, we have that H is an I -augmenting path. \square

We use the following special case of the result of Gerber et al., who showed that I -augmenting paths can be computed in polynomial time in graphs with no induced skew stars.

Theorem 22 ([GHL06]). *There is a polynomial-time algorithm that finds an I -augmenting path in G if it exists.*

We now have everything at hand to show how to resolve cycles of the symmetric difference in polynomial time.

Lemma 23. *Assume that G is prime, I -reduced and J -reduced, and that I and J are not maximum. Then we can find in polynomial-time either a C -resolving sequence or a permanently blocked vertex of G .*

Proof. There are two cases to consider. First, suppose that I is not maximal. Then there exists a vertex $u \notin N[I]$. Pick an arbitrary token on v in C . Since u is I -free, Theorem 14 ensures that we can find in polynomial time either a locally blocked vertex or a sequence of slides from I to $A := I - u + v$. In the later case, we can first apply the sequence from I to A and then move the remaining tokens on $C \cap A$ to $C \cap J$. Let A' denote the obtained independent set: since Reduction Rule A is not applicable with respect to I and $I \rightsquigarrow_{\text{TS}} A'$, we have that Reduction Rule A is not applicable with respect to A' . Furthermore, there exists a vertex $u' \in C$ that is A' -free. We can therefore apply Theorem 14 to find in polynomial time either a sequence of slides from $A' - u + u'$ from A' , thus resolving the cycle C , or a locally blocked vertex of G .

Now suppose that I is maximal, so by Lemma 21, there must exist a I -augmenting path. By Theorem 22, such a path $v_1, u_1, \dots, u_k, v_k$ can be found in polynomial time, where u_1, \dots, u_k denote the vertices of the path that are in I . After sliding for $1 \leq i < k$ in increasing order the token on u_i to v_i , the vertex v_k has no token in its neighborhood. Therefore, the previous case applies and we may then slide the token of the chain back to their initial position \square

We now have everything we need to prove the main result of this section.

Theorem 24. *TOKEN SLIDING in fork-free graphs admits a polynomial-time algorithm.*

Proof. Let (G, I, J) be an instance of TOKEN SLIDING such that G is fork-free. We may assume that I and J are not maximum, since otherwise Theorem 8 applies. This can be tested in polynomial time by the algorithm given in [Ale04]. Otherwise, we invoke Theorem 13 and obtain in polynomial time an equivalent instance (G', I', J') , such that G' is fork-free and each connected component is prime (or (G', I', J') is a trivial No-instance).

Since (G', I, J) is a Yes-instance if and only if $(C, I \cap V(C), J \cap V(C))$ is a Yes-instance for each connected component C of G' , we may assume that G' is connected. By Lemma 20, the graph $G'[I \Delta J]$ is a collection of disjoint paths, cycles and isolated vertices. If $G'[I \Delta J]$ contains a path P we return the sequence sliding the tokens from $I \cap P$ to $J \cap P$ one by one. If $G'[I \Delta J]$ contains two isolated vertices $u \in I$ and $v \in J$, Theorem 14 ensures that we can either find a sequence from I to $I - u + v$ or a permanently blocked vertex in polynomial time. If $G'[I \Delta J]$ contains a cycle C , by Lemma 23, we can either find a resolving sequence for C or a permanently blocked vertex. If at any step we find a permanently blocked vertex, we can delete it from G' by Lemma 6 and restart the algorithm. Since furthermore the reduction rules A, B, D, E can be applied exhaustively in polynomial time, the overall running time of the algorithm is polynomial. \square

7 Conclusion

We showed `TOKEN SLIDING` and `TOKEN JUMPING` remain `PSPACE`-complete on H -free graphs, unless H is a path, a claw, or a subdivision of the claw. Furthermore, we showed that `TOKEN SLIDING` admits a polynomial-time algorithm on fork-free graphs. The overall goal is the same as for the nominal problem, namely to obtain a complete classification of the complexity of `ISR` on H -free graphs. In this direction, an interesting open question is whether `TOKEN SLIDING` admits a polynomial-time algorithm on fork-free graphs if the independent sets are not maximum. There are two main reasons why our approach for `TOKEN SLIDING` does not work for `TOKEN JUMPING`. First, we cannot assume that tokens stay in their respective modules, so the modular decomposition seems less useful. Secondly, the irrelevant vertices for `TOKEN SLIDING` may not be irrelevant for `TOKEN JUMPING`, for example, vertices that are adjacent to three tokens. This means in particular that in order to resolve a cycle, it is not sufficient to consider augmenting chains, but it may be necessary to deal with large complexes induced by the symmetric difference of the independent sets. A second interesting question is whether `ISR` admits a polynomial-time algorithm on P_5 -free graphs. As a first step, the case of maximum independent sets could be considered.

References

- [Ale82] Vladimir E Alekseev. The effect of local constraints on the complexity of determination of the graph independence number. *Combinatorial-algebraic methods in applied mathematics*, pages 3–13, 1982.
- [Ale04] Vladimir E Alekseev. Polynomial algorithm for finding the largest independent sets in graphs without forks. *Discrete Applied Mathematics*, 135(1-3):3–16, January 2004.
- [Bar21] Valentin Bartier. *Combinatorial and Algorithmic aspects of Reconfiguration. (Aspects combinatoires et algorithmiques de la Reconfiguration)*. PhD thesis, Grenoble Alpes University, France, 2021.
- [BB17] Marthe Bonamy and Nicolas Bousquet. Token Sliding on Chordal Graphs. In Hans L. Bodlaender and Gerhard J. Woeginger, editors, *Graph-Theoretic Concepts in Computer Science*, volume 10520, pages 127–139. Springer International Publishing, Cham, 2017. Series Title: Lecture Notes in Computer Science.
- [BBD⁺21] Valentin Bartier, Nicolas Bousquet, Clément Dallard, Kyle Lomer, and Amer E. Mouawad. On girth and the parameterized complexity of token sliding and token jumping. *Algorithmica*, 83(9):2914–2951, 2021.
- [BCM23] Manoj Belavadi, Kathie Cameron, and Owen Merkel. Reconfiguration of vertex colouring and forbidden induced subgraphs, 2023.
- [BDK⁺23] Marthe Bonamy, Oscar Defrain, Tereza Klimosová, Aurélie Lagoutte, and Jonathan Narboni. On Vizing’s edge colouring question. *J. Comb. Theory, Ser. B*, 159:126–139, 2023.
- [BDO21] Marthe Bonamy, Paul Dorbec, and Paul Ouvrard. Dominating sets reconfiguration under token sliding. *Discret. Appl. Math.*, 301:6–18, 2021.
- [BHI⁺20] Marthe Bonamy, Marc Heinrich, Takehiro Ito, Yusuke Kobayashi, Haruka Mizuta, Moritz Mühlenhaller, Akira Suzuki, and Kunihiro Wasa. Diameter of colorings under Kempe changes. *Theor. Comput. Sci.*, 838:45–57, 2020.
- [BHV04] Andreas Brandstädt, Chinh T. Hoàng, and Jean-Marie Vanherpe. On minimal prime extensions of a four-vertex graph in a prime graph. *Discret. Math.*, 288(1-3):9–17, 2004.
- [BKL⁺21] Rémy Belmonte, Eun Jung Kim, Michael Lampis, Valia Mitsou, Yota Otachi, and Florian Sikora. Token sliding on split graphs. *Theory Comput. Syst.*, 65(4):662–686, 2021.

- [BKW14] Paul S. Bonsma, Marcin Kamiński, and Marcin Wrochna. Reconfiguring independent sets in claw-free graphs. In *Algorithm Theory - SWAT 2014 - 14th Scandinavian Symposium and Workshops*, volume 8503 of *Lecture Notes in Computer Science*, pages 86–97. Springer, 2014.
- [BMNS22] Nicolas Bousquet, Amer E. Mouawad, Naomi Nishimura, and Sebastian Siebertz. A survey on the parameterized complexity of the independent set and (connected) dominating set reconfiguration problems. *CoRR*, abs/2204.10526, 2022.
- [Bon16] Paul S. Bonsma. Independent set reconfiguration in cographs and their generalizations. *J. Graph Theory*, 83(2):164–195, 2016.
- [BP16] Nicolas Bousquet and Guillem Perarnau. Fast recoloring of sparse graphs. *Eur. J. Comb.*, 52:1–11, 2016.
- [DDF⁺15] Erik D. Demaine, Martin L. Demaine, Eli Fox-Epstein, Duc A. Hoang, Takehiro Ito, Hirotaka Ono, Yota Otachi, Ryuhei Uehara, and Takeshi Yamada. Linear-time algorithm for sliding tokens on trees. *Theor. Comput. Sci.*, 600:132–142, 2015.
- [DF21] Zdenek Dvorák and Carl Feghali. A Thomassen-type method for planar graph recoloring. *Eur. J. Comb.*, 95:103319, 2021.
- [FHO15] Eli Fox-Epstein, Duc A. Hoang, Yota Otachi, and Ryuhei Uehara. Sliding token on bipartite permutation graphs. In *Algorithms and Computation - 26th International Symposium, ISAAC 2015, Nagoya, Japan, December 9-11, 2015, Proceedings*, pages 237–247, 2015.
- [Gal67] Tibor Gallai. Transitiv orientierbare Graphen. *Acta Mathematica Hungarica*, 18(1-2):25–66, 1967.
- [GHL06] Michael U. Gerber, Alain Hertz, and Vadim V. Lozin. Augmenting chains in graphs without a skew star. *J. Comb. Theory, Ser. B*, 96(3):352–366, 2006.
- [GKPP22] Andrzej Grzesik, Tereza Klimosová, Marcin Pilipczuk, and Michal Pilipczuk. Polynomial-time algorithm for maximum weight independent set on p_6 -free graphs. *ACM Trans. Algorithms*, 18(1):4:1–4:57, 2022.
- [HD05] Robert A. Hearn and Erik D. Demaine. PSPACE-completeness of sliding-block puzzles and other problems through the nondeterministic constraint logic model of computation. *Theor. Comput. Sci.*, 343(1-2):72–96, 2005.
- [vdH13] Jan van den Heuvel. The complexity of change. In Simon R. Blackburn, Stefanie Gerke, and Mark Wildon, editors, *Surveys in Combinatorics 2013*, volume 409 of *London Mathematical Society Lecture Note Series*, pages 127–160. Cambridge University Press, 2013.
- [HIM⁺16] Arash Haddadan, Takehiro Ito, Amer E. Mouawad, Naomi Nishimura, Hirotaka Ono, Akira Suzuki, and Youcef Tebbal. The complexity of dominating set reconfiguration. *Theor. Comput. Sci.*, 651:37–49, 2016.
- [IDH⁺11] Takehiro Ito, Erik D. Demaine, Nicholas J. A. Harvey, Christos H. Papadimitriou, Martha Sideri, Ryuhei Uehara, and Yushi Uno. On the complexity of reconfiguration problems. *Theor. Comput. Sci.*, 412(12-14):1054–1065, 2011.
- [IKO14a] Takehiro Ito, Marcin Kamiński, and Hirotaka Ono. Fixed-parameter tractability of token jumping on planar graphs. In *Algorithms and Computation - 25th International Symposium, ISAAC 2014*, volume 8889 of *Lecture Notes in Computer Science*, pages 208–219. Springer, 2014.
- [IKO⁺14b] Takehiro Ito, Marcin Kamiński, Hirotaka Ono, Akira Suzuki, Ryuhei Uehara, and Katsuhisa Yamanaka. On the parameterized complexity for token jumping on graphs. In *Theory and Applications of Models of Computation - 11th Annual Conference, TAMC. Proceedings*, volume 8402 of *Lecture Notes in Computer Science*, pages 341–351. Springer, 2014.

- [IKO⁺20] Takehiro Ito, Marcin Kamiński, Hirotaka Ono, Akira Suzuki, Ryuhei Uehara, and Katsuhisa Yamanaka. Parameterized complexity of independent set reconfiguration problems. *Discret. Appl. Math.*, 283:336–345, 2020.
- [KMM12] Marcin Kamiński, Paul Medvedev, and Martin Milanič. Complexity of independent set reconfigurability problems. *Theor. Comput. Sci.*, 439:9–15, 2012.
- [LM08] Vadim V. Lozin and Martin Milanič. A polynomial algorithm to find an independent set of maximum weight in a fork-free graph. *Journal of Discrete Algorithms*, 6(4):595–604, December 2008.
- [LM19] Daniel Lokshтанov and Amer E. Mouawad. The Complexity of Independent Set Reconfiguration on Bipartite Graphs. *ACM Transactions on Algorithms*, 15(1):1–19, January 2019.
- [LVV14] Daniel Lokshтанov, Martin Vatshelle, and Yngve Villanger. Independent set in P_5 -free graphs in polynomial time. In *Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2014*, pages 570–581. SIAM, 2014.
- [Min80] George J. Minty. On maximal independent sets of vertices in claw-free graphs. *J. Comb. Theory, Ser. B*, 28(3):284–304, 1980.
- [MNR⁺17] Amer E. Mouawad, Naomi Nishimura, Venkatesh Raman, Narges Simjour, and Akira Suzuki. On the parameterized complexity of reconfiguration problems. *Algorithmica*, 78(1):274–297, 2017.
- [MNRS18] Amer E. Mouawad, Naomi Nishimura, Venkatesh Raman, and Sebastian Siebertz. Vertex cover reconfiguration and beyond. *Algorithms*, 11(2):20, 2018.
- [Möh85] Rolf H. Möhring. Algorithmic aspects of the substitution decomposition in optimization over relations, set systems and boolean functions. *Annals of Operations Research*, 4:195–225, 1985.
- [MR84] Rolf H. Möhring and Franz J. Radermacher. Substitution decomposition for discrete structures and connections with combinatorial optimization. In *North-Holland mathematics studies*, volume 95, pages 257–355. Elsevier, 1984.
- [MS99] Ross M. McConnell and Jeremy P. Spinrad. Modular decomposition and transitive orientation. *Discrete Mathematics*, 201(1-3):189–241, 1999.
- [Nis18] Naomi Nishimura. Introduction to reconfiguration. *Algorithms*, 11(4):52, 2018.
- [Sbi80] Najiba Sbihi. Algorithme de recherche d’un stable de cardinalité maximum dans un graphe sans étoile. *Discrete Mathematics*, 29(1):53–76, 1980.
- [SMN16] Akira Suzuki, Amer E. Mouawad, and Naomi Nishimura. Reconfiguration of dominating sets. *Journal of Combinatorial Optimization*, 32(4):1182–1195, Nov 2016.