

# A Note on Rounding Matchings in General Graphs

Aditi Dudeja\*

Department of Computer Science, University of Salzburg

## Abstract

In this note, we revisit the rounding algorithm of Wajc [Waj20]. Wajc gave a fully-adaptive randomized algorithm that rounds a dynamic fractional matching in an *unweighted bipartite graph* to an integral matching of nearly the same value in  $O(\text{poly}(\log n, 1/\varepsilon))$  update time. We give show that the guarantees of this algorithm hold for *general graphs* as well. Additionally, we show useful properties of this subroutine which have applications in rounding weighted fractional matchings.

In the dynamic matching problem, the graph undergoes edge insertions and deletions, and the algorithm is required to maintain a good approximation to the maximum matching. The goal is to optimize the update time which is the time required to cope with a single edge insertion or deletion. Several dynamic algorithms for approximate maximum matching proceed by computing a fractional matching [BHI15, BHN16, BCH20, BHN17, BK23] and then rounding it [ACC<sup>+</sup>18, Waj20, BK21, Kis22, BKS<sup>+</sup>23]. Fractional matchings have several interesting advantages. They are not only computationally easier to compute, but also certain types of fractional matchings are robust to adversarial edge insertions and deletions [BGS20, ABD22, CST23, JJST22]. Thus, there has been a lot of focus on designing efficient rounding algorithms.

## 1 Background on Dynamic Rounding of Fractional Matching

Arar, Chechik, Cohen, Stein, and Wajc [ACC<sup>+</sup>18] gave a rounding procedure based on independent sampling: given a dynamic unweighted fractional matching  $\vec{x}$ , sample every edge with a probability proportional to  $x(e)$  and include it in the subgraph  $S$ . The resulting subgraph  $S$  has the property that  $\mu(S) \geq (1 - \varepsilon) \cdot \sum_{e \in E} x(e)$ , and is sparse. However, it is only robust to an oblivious adversary. Subsequently, [Waj20] gave a dependent version of the above sampling scheme, and showed that the resulting sparsifier was now robust to an adaptive adversary. This rounding procedure, combined with existing fractional matching results, then gave a slew of fully-adaptive dynamic matching algorithms, with varying approximation ratios and update times. However, his analysis only demonstrated that *arbitrary* fractional matchings in bipartite graphs can be rounded to integral matchings of nearly the same value. In another work, Bhattacharya and Kiss [BK21] gave a deterministic rounding procedure for fractional matchings in bipartite graphs.

Subsequently, Bhattacharya, Kiss, Sidford, and Wajc [BKS<sup>+</sup>23] showed that the sampling approach of [ACC<sup>+</sup>18] can be made robust to output-adaptive adversaries. They also gave deterministic rounding algorithms for bipartite graphs, and for certain types of structured matchings in general graphs. Very recently, Chen, Sidford, and Tu [CST23] showed that the deterministic rounding procedures of [BKS<sup>+</sup>23] and [ACC<sup>+</sup>18] also work for general graphs. Curiously, the randomized sparsifier of [ACC<sup>+</sup>18] also has some additional nice properties, which have applications for rounding weighted matchings. The deterministic sparsifier doesn't seem to share these additional nice properties. In particular, [ACC<sup>+</sup>18] show the following theorem for dense weighted graphs:

\*aditi.dudeja@plus.ac.at. This work is supported by Austrian Science Fund (FWF): P 32863-N. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 947702).

**Result 1** (Informal [CST23]). There is an output-adaptive randomized algorithm that maintains  $(1 - \varepsilon)$ -approximate maximum weight matching in decremental dense weighted graphs with an update time of  $O(\text{poly}(\log n, \frac{1}{\varepsilon}))$ .

Prior to this, all known weighted matching algorithms either had a dependence of  $(1/\varepsilon)^{O(1/\varepsilon)}$ , or had a significantly worse approximation ratio than their unweighted counterparts. A crucial step in obtaining Result 1, is the rounding algorithm of [ACC<sup>+</sup>18] and [BKS<sup>+</sup>23]. This is also the reason why this algorithm is not fully-adaptive.

In their paper, [CST23] left open the question of analysing Wajc’s algorithm, since it is a dependent and consequently, easier to sample from version of [ACC<sup>+</sup>18]. We show that Wajc’s algorithm also works for general graphs, and additionally show the fully adaptive version of Result 1. In particular, we show the following two results:

**Result 2** (Informal). There is a fully-adaptive randomized dynamic rounding algorithm for fractional matchings in general graphs with amortized update time  $O(\text{poly}(\log n, 1/\varepsilon))$  update time.

**Result 3** (Informal). There is a fully adaptive randomized algorithm that maintains  $(1 - \varepsilon)$ -approximate maximum weight matching in decremental dense weighted graphs with an update time of  $O(\text{poly}(\log n, 1/\varepsilon))$ .

## 2 General Graph Rounding

We start by defining some notation, then we state Wajc’s algorithm. We emphasize that the algorithm remains the same. We merely show that the analysis extends to the case of general (possibly non-bipartite) graphs as well.

**Notation** Given a graph  $G$ , we will use  $\mu(G)$  to denote the size of the maximum matching in  $G$ . Sometimes, we will consider weighted graphs, and in this case, we use  $\text{mwm}(G)$  to denote the weight of the maximum weight matching. We will use  $\vec{x}$  to denote a fractional matching. Given  $\vec{x}$ , we define

$$E_i = \{e \mid x(e) \in [(1 + \varepsilon)^{-i}, (1 + \varepsilon)^{-i+1}]\} \tag{1}$$

$$G_i = (V, E_i) \tag{2}$$

For a vertex  $v$ , we define  $E_v$  to be the set of edges incident on the vertex  $v$ . Additionally, given a graph  $G$  we have the following notation, which we borrow from [CST23]:

$$\begin{aligned} \mathcal{O} &= \{B \subseteq V \mid |B| \text{ is odd}\} \\ \mathcal{O}_\varepsilon &= \{B \subseteq V \mid |B| \text{ is odd and } |B| \leq 1/\varepsilon\} \\ \mathcal{P}_G &= \left\{ \vec{x} \in \mathbb{R}_{\geq 0}^E \mid \sum_{e \in E_v} x(e) \leq 1 \right\} \\ \mathcal{M}_G &= \left\{ \vec{x} \in \mathcal{P}_G \mid \sum_{e \in G[B]} x(e) \leq \frac{|B| - 1}{2}, \forall B \in \mathcal{O} \right\} \\ \mathcal{M}_{G,\varepsilon} &= \left\{ \vec{x} \in \mathcal{P}_G \mid \sum_{e \in G[B]} x(e) \leq \frac{|B| - 1}{2}, \forall B \in \mathcal{O}_\varepsilon \right\} \end{aligned}$$

We will also often use the following fact about matchings.

**Fact 4** ([ABD22]). Suppose  $\vec{x} \in \mathcal{M}_{G,\varepsilon}$ , then  $\frac{\vec{x}}{(1+\varepsilon)} \in \mathcal{M}_G$ .

**Observation 5.** Ignoring the set of edges  $E_{\text{low}} = \{e \in E \mid x(e) \leq \frac{\varepsilon^2}{n^2}\}$  does not affect  $\vec{x}$  by more than a value  $\varepsilon$ . Consequently, in the notation in (1), we can assume that  $i \in [2 \cdot \log \frac{n}{\varepsilon}]$ .

---

**Algorithm 1** SPARSIFICATION( $G, \vec{x}, \varepsilon$ )

---

- 1: **for**  $i = 1, \dots, \lceil 2 \cdot \log \frac{n}{\varepsilon} \rceil$  **do**
  - 2:     Edge color the graph  $G_i$  using  $2 \cdot \lceil (1 + \varepsilon)^i \rceil$  many colors. ▷ Coloring Phase
  - 3:     Sample  $\min \{2 \cdot \lceil d \cdot (1 + \varepsilon) \rceil, 2 \cdot \lceil (1 + \varepsilon)^i \rceil\}$  colors from  $G_i$ . Denote sampled edges by  $S_i$ . ▷ Sampling Phase
  - 4: **end for**
  - 5: Output  $S = \cup_i S_i$ .
- 

The sparsification procedure is stated in Algorithm 1. We first start by giving the following property of this random process, which is a paraphrasing of Lemma 3.2 from [Waj20].

**Observation 6.** [Waj20] Suppose  $d \geq \frac{\log n}{\varepsilon^2}$ , for an edge  $e$ , we have,

$$\frac{\min \{1, x(e) \cdot d\}}{(1 + \varepsilon)^2} \leq \Pr(e \in S) \leq \min \{1, x(e) \cdot d\} \cdot (1 + \varepsilon).$$

Additionally, if  $x(e) > 1/d$ , then  $\Pr(e \in S) = 1$ .

**Definition 7.** For an edge  $e$ , we denote  $\mathbb{1}_e$  to be the indicator random variable for the event  $e \in S$ .

**Definition 8.** Observe that the Coloring Phase splits up the graphs  $G_i$ , and consequently  $G$  into monochromatic matchings. Let  $M_1, \dots, M_l$  be these matchings, where  $l \leq 2 \cdot \lceil d \cdot (1 + \varepsilon) \rceil \cdot \lceil 2 \cdot \log \frac{n}{\varepsilon} \rceil$ . For each of these matchings we associate an indicator random variable  $\mathbb{1}_{M_i}$  which takes value 1 if  $M_i \in S$  and 0 otherwise.

**Lemma 9.** ([DP09]) Suppose  $b_1, b_2, \dots, b_r$  be any set of  $r$  elements. Consider the random process of sampling  $k \leq r$  objects without replacement from these  $r$  elements. Let  $X_i$  denote the indicator variable of  $b_i$  being included in the sample. Then,  $\{X_i\}_{i \in [r]}$  are negatively associated random variables.

**Observation 10.** Consider a vertex  $v$ , then the random variables  $\{\mathbb{1}_e\}_{e \in E_v}$  are negatively associated random variables.

We also have the following observation, which

**Observation 11.** Consider any odd set  $B$ , note that the Coloring Phase splits up  $G_i[B]$  and consequently,  $G[B]$  into monochromatic matchings of size at most  $\frac{|B|-1}{2}$ . Moreover, all edges on a given matching  $M$  are roughly the same weight. Let  $M'$  be any such matching in  $G_i$  for  $i \in \{1, 2, \dots, \lceil 2 \cdot \log \frac{n}{\varepsilon} \rceil\}$ , and  $\mathbb{1}_{M'}$  is an indicator random variable which takes value 1 if  $M' \in S$  in the Sampling Phase.

**Lemma 12.** Consider any odd set  $B$ , and let  $M'_1, \dots, M'_j$  be the matchings  $G[B]$  has been split up into after the Coloring Phase. Then,  $\{\mathbb{1}_{M'_j}\}_{M'_j \in G[B]}$  are negatively associated random variables.

*Proof.* Consider  $G_l$  and recall Lemma 9. The set of colors, and consequently the matchings correspond to the set of elements we are sampling from in Lemma 9. Let  $M_1, \dots, M_k$  be the matchings associated with each color (these matchings can be potentially empty, if the colors are not used). In the Coloring Phase, we are sampling a subset of these matchings without replacement. Thus, the random variables,  $\{\mathbb{1}_{M_i}\}_{i \in [k]}$  are negatively associated by Lemma 9. Since subsets of negatively associated random variables are also negatively associated, we have that  $\{\mathbb{1}_{M_i} \mid M_i \neq \emptyset\}_{i \in [k]}$  are also negatively associated. Next, the independent union of negatively associated random variables are also negatively associated. Thus, we have,  $\mathcal{R} = \{\mathbb{1}_M \mid M \neq \emptyset\}_{M \in G}$  are negatively associated as well.

Finally, consider an odd set  $B$ , and let  $M'_1, \dots, M'_p$  be the matchings  $G[B]$  is split up into. Note that the indicator random associated with these matchings are identical to distinct random variables in  $\mathcal{R}$ . Consequently, we have,  $\{\mathbb{1}_{M'_i}\}_{M'_i \in G[B]}$  are negatively associated as well.  $\square$

Next, we have the following lemma, which is an analog of Observation 6 for the case of matchings, and in fact directly follows from Observation 6.

**Observation 13.** Suppose  $M \in G_i[B]$ , and let  $e \in M$ , then we have,

$$\frac{d \cdot x(M)}{(1 + \varepsilon) \cdot |M|} \leq \Pr(M \in S) \leq (1 + \varepsilon)^2 \cdot \frac{d \cdot x(M)}{|M|},$$

if  $d < (1 + \varepsilon)^{i-1}$ , and  $\Pr(M \in S) = 1$  otherwise.

**Lemma 14** (Chernoff Bounds [DP09]). Suppose  $X_1, \dots, X_m$  are  $m$  negatively associated random variables with range  $[0, b]$  each for some  $b \geq 1$ . Let  $X := \sum_{i=1}^m X_i$  and  $\mu_L \leq \mathbb{E}[X] \leq \mu_H$ . Then, for any  $\delta > 0$ ,

$$\Pr(X > (1 + \delta) \cdot \mu_H) \leq \exp\left(-\frac{\delta^2 \cdot \mu_H}{(3 + \delta) \cdot b}\right) \quad \text{and} \quad \Pr(X < (1 - \delta) \cdot \mu_L) \leq \exp\left(-\frac{\delta^2 \cdot \mu_L}{(2 + \delta) \cdot b}\right)$$

**Lemma 15.** Suppose  $\vec{x} \in \mathcal{M}_G$  is input to SPARSIFICATION(), and  $d \geq \frac{\log n}{\varepsilon^4}$ , then, with high probability, there is a vector  $\vec{y} \subseteq \text{supp}(S)$  such that:

- a) For all vertices  $v \in V$ ,  $(1 - \varepsilon) \cdot \sum_{e \in E_v} x(e) - \varepsilon \leq \sum_{e \in E_v} y(e) \leq (1 - \varepsilon) \cdot \sum_{e \in E_v} x(e) + \varepsilon$ .
- b) For all odd sets  $B$  with  $|B| \leq 1/\varepsilon$ ,

$$(1 - \varepsilon) \cdot \sum_{e \in G[B]} x(e) - \frac{\varepsilon \cdot (|B| - 1)}{2} \leq \sum_{e \in G[B]} y(e) \leq (1 - \varepsilon) \cdot \sum_{e \in G[B]} x(e) + \frac{\varepsilon \cdot |B|}{2}.$$

Moreover, if  $\sum_{e \in E} x(e) \geq (1 - \varepsilon) \cdot \mu(G)$ , then with a constant probability, we have, that  $\vec{y}$  satisfies:

$$\sum_{e \in E} y(e) \geq (1 - \varepsilon) \cdot \sum_{e \in E} x(e). \quad (3)$$

The above lemma essentially states that with a constant probability there is a fractional matching  $\vec{y} \subseteq \text{supp}(S)$  with  $\sum_{e \in E} y(e) \geq (1 - \varepsilon) \cdot \sum_{e \in E} x(e)$  satisfying small odd set constraints.

*Proof.* We first define a flow  $\vec{y}$  on the support of  $S$ . Then, we show a) and b) are satisfied for  $\vec{y}$  with high probability. Finally, we conditioned on a) and b), we show that  $\sum_{e \in E} y(e) \geq (1 - \varepsilon) \cdot \sum_{e \in E} x(e)$  with a constant probability. Let  $S_v = \{e \ni v \mid x(e) < \frac{1}{d}\}$ . We now define the flow as follows, for all  $v \in V$ ,  $e \in S_v$ ,

$$y(e) = \frac{1}{d} \cdot \mathbb{1}_e \cdot (1 - 4\varepsilon)$$

On the other hand, for  $e \notin S_v$ , we let  $y(e) = x(e)$ . We first show a). Consider an edge  $e \in S_v$ , from Observation 6, we have,

$$x(e) \cdot (1 - 3\varepsilon) \leq \mathbb{E}[y(e)] \leq x(e) \cdot (1 - 2\varepsilon)$$

Consequently, for a vertex  $v$ , we have,

$$(1 - 3\varepsilon) \cdot \sum_{e \in S_v} x(e) \leq \mathbb{E}\left[\sum_{e \in S_v} y(e)\right] \leq (1 - 2\varepsilon) \cdot \sum_{e \in S_v} x(e)$$

Note that to show concentration, it is sufficient to consider the set of edges  $S_v$ . This is because of Observation 6. Next, observe that for  $e \in S_v$ ,  $y(e) \leq \frac{1}{d}$ . Finally, by Observation 10, we have, that  $\{y(e)\}_{e \in S_v}$  are negatively associated. Thus, by Chernoff bound (Lemma 14), we have, the following upper tail bound by picking  $\mu_H = (1 - 2\varepsilon) \cdot \sum_{e \in S_v} x(e)$ , and  $\delta = \frac{\varepsilon}{(1 - 2\varepsilon) \cdot \sum_{e \in S_v} x(e)}$ ,

$$\Pr\left(\sum_{e \in S_v} y(e) \geq (1 - 2\varepsilon) \cdot \sum_{e \in S_v} x(e) + \varepsilon\right) \leq \exp\left(-\frac{\left(\frac{\varepsilon}{(1 - 2\varepsilon) \cdot \sum_{e \in S_v} x(e)}\right)^2 \cdot \left((1 - 2\varepsilon) \cdot \sum_{e \in S_v} x(e)\right)}{\left(3 + \frac{\varepsilon}{(1 - 2\varepsilon) \cdot \sum_{e \in S_v} x(e)}\right) \cdot \frac{\varepsilon^4}{\log n}}\right)$$

$$= O\left(\frac{1}{n^{1/\varepsilon}}\right)$$

Similarly, substituting  $\mu_L = (1 - 3\varepsilon) \cdot \sum_{e \in S_v} x(e)$  and  $\delta = \frac{\varepsilon}{(1-3\varepsilon) \cdot \sum_{e \in S_v} x(e)}$ ,

$$\begin{aligned} \Pr\left(\sum_{e \in S_v} y(e) \leq (1 - 3\varepsilon) \cdot \sum_{e \in S_v} x(e) - \varepsilon\right) &\leq \exp\left(-\frac{\left(\frac{\varepsilon}{(1-3\varepsilon) \cdot \sum_{e \in S_v} x(e)}\right)^2 \cdot \left((1 - 3\varepsilon) \cdot \sum_{e \in S_v} x(e)\right)}{\left(2 + \frac{\varepsilon}{(1-3\varepsilon) \cdot \sum_{e \in S_v} x(e)}\right) \cdot \frac{\varepsilon^4}{\log n}}\right) \\ &= O\left(\frac{1}{n^{1/\varepsilon}}\right) \end{aligned}$$

Finally, since  $\sum_{e \in E_v \setminus S_v} y(e) = (1 - 4\varepsilon) \cdot \sum_{e \in E_v \setminus S_v} x(e)$  for all  $v \in V$ , we can conclude that with high probability,

$$(1 - 4\varepsilon) \cdot \sum_{e \in E_v} x(e) - \varepsilon \leq \sum_{e \in E_v} y(e) \leq (1 - 3\varepsilon) \cdot \sum_{e \in E_v} x(e) - \varepsilon$$

We now show *b*). Consider an odd set  $B$ , and let  $M'_1, \dots, M'_k$  denote the set of matchings  $G[B]$  is split up into. As before, we can consider only matchings in  $G_i$ , where  $d < (1 + \varepsilon)^{i-1}$  due to Observation 13.

$$\mathbb{E}\left[\sum_{e \in G[B]} y(e)\right] = \mathbb{E}\left[\sum_{i=1}^k y(M'_i)\right] \leq \sum_{i=1}^k x(M'_i) \cdot (1 - 4\varepsilon) \leq \sum_{e \in G[B]} x(e) \cdot (1 - 4\varepsilon)$$

Similarly, we have,

$$\mathbb{E}\left[\sum_{e \in G[B]} y(e)\right] = \mathbb{E}\left[\sum_{i=1}^k y(M'_i)\right] \geq \sum_{e \in G[B]} x(e) \cdot (1 - 5\varepsilon)$$

Additionally, observe that  $y(M'_i)$  can take value at most  $\frac{|M'_i|}{d} \leq \frac{|B|-1}{2 \cdot d}$ . Thus, using  $\mu_H = \sum_{e \in G[B]} x(e) \cdot (1 - 4\varepsilon)$ ,  $\delta = \frac{\varepsilon \cdot (|B|-1)}{\sum_{e \in G[B]} x(e) \cdot (1-4\varepsilon)}$ , we have,

$$\begin{aligned} \Pr\left(\sum_{e \in G[B]} y(e) \geq \sum_{e \in G[B]} x(e) \cdot (1 - 4\varepsilon) + \frac{\varepsilon \cdot (|B|-1)}{2}\right) &\leq \exp\left(-\frac{\left(\frac{-\varepsilon^2 \cdot (|B|-1)^2}{4 \cdot \left(\sum_{e \in G[B]} x(e) \cdot (1-4\varepsilon)\right)^2}\right) \cdot (1 - 4\varepsilon) \cdot \sum_{e \in G[B]} x(e)}{\left(3 + \frac{\varepsilon}{\sum_{e \in G[B]} x(e) \cdot (1-3\varepsilon)}\right) \cdot \frac{|B|-1}{2 \cdot d}}\right) \\ &= O\left(\frac{1}{n^{1/\varepsilon^2}}\right). \end{aligned}$$

Using the fact that  $\sum_{e \in G[B]} x(e) \leq \frac{|B|-1}{2}$  and  $d \geq \log n / \varepsilon^4$ , we have the second equality. Finally, we union bounding over all odd sets of size at most  $\frac{1}{\varepsilon}$ , we have the upper bound. For the lower bound, consider  $\mu_L = (1 - 5\varepsilon) \cdot \sum_{e \in G[B]} x(e)$ , and  $\delta = \frac{\varepsilon \cdot (|B|-1)}{(1-4\varepsilon) \sum_{e \in G[B]} x(e)}$ ,

$$\begin{aligned} \Pr\left(\sum_{e \in G[B]} y(e) \geq \sum_{e \in G[B]} x(e) \cdot (1 - 4\varepsilon) - \frac{\varepsilon \cdot (|B|-1)}{2}\right) &\leq \exp\left(-\frac{\left(\frac{-\varepsilon^2 \cdot (|B|-1)^2}{4 \cdot \left(\sum_{e \in G[B]} x(e) \cdot (1-4\varepsilon)\right)^2}\right) \cdot (1 - 4\varepsilon) \cdot \sum_{e \in G[B]} x(e)}{\left(3 + \frac{\varepsilon}{\sum_{e \in G[B]} x(e) \cdot (1-3\varepsilon)}\right) \cdot \frac{|B|-1}{2 \cdot d}}\right) \\ &\leq O\left(\frac{1}{n^{1/\varepsilon^2}}\right). \end{aligned}$$

Taking a union bound over all odd sets of size at most  $\frac{1}{\varepsilon}$ , we have *b*). We now move on to proving Equation (3). First, let  $\mathcal{E}_{b,c}$  denote the event that *a*) and *b*) occur. We have,

$$\Pr(\mathcal{E}_{b,c}) \geq 1 - O\left(\frac{1}{n^{1/\varepsilon^2}}\right).$$

Additionally, we have,

$$\mathbb{E} \left[ \sum_{e \in E} y(e) \right] \geq (1 - 4\varepsilon) \cdot \sum_{e \in E} x(e).$$

By law of total probability, we have,

$$\begin{aligned} \mathbb{E} \left[ \sum_{e \in E} y(e) \mid \mathcal{E}_{b,c} \right] \cdot \Pr(\mathcal{E}_{b,c}) + \mathbb{E} \left[ \sum_{e \in E} y(e) \mid \neg \mathcal{E}_{b,c} \right] \cdot \Pr(\neg \mathcal{E}_{b,c}) &\geq (1 - 4\varepsilon) \cdot \sum_{e \in E} x(e) \\ \mathbb{E} \left[ \sum_{e \in E} y(e) \mid \mathcal{E}_{b,c} \right] + n^2 \cdot O\left(\frac{1}{n^{1/\varepsilon}}\right) &\geq (1 - 4\varepsilon) \cdot \sum_{e \in E} x(e) \end{aligned}$$

Thus, we have,  $\mathbb{E} \left[ \sum_{e \in E} y(e) \mid \mathcal{E}_{b,c} \right] \geq (1 - 5\varepsilon) \cdot \sum_{e \in E} x(e)$ . Let  $\vec{x}^* \in \mathcal{M}_G$  be any optimum fractional matching. Conditioned on  $\mathcal{E}_{b,c}$ , we have  $\vec{y} \in \mathcal{M}_{G,\varepsilon}$ . Thus, from Fact 4, we have,  $\sum_{e \in E} y(e) \leq (1 + \varepsilon) \cdot \sum_{e \in E} x^*(e)$ . Therefore, the random variable  $(1 + \varepsilon) \cdot \sum_{e \in E} x^*(e) - y(e)$  is non-negative, and moreover, by our assumption that  $\sum_{e \in E} x(e) \geq (1 - \varepsilon) \cdot \sum_{e \in E} x^*(e)$ , we have,

$$\mathbb{E} \left[ (1 + \varepsilon) \cdot \sum_{e \in E} x^*(e) - y(e) \mid \mathcal{E}_{b,c} \right] \leq 4\varepsilon \cdot \sum_{e \in E} x^*(e)$$

Thus, by Markov inequality, we have,

$$\Pr \left( (1 + \varepsilon) \cdot \sum_{e \in E} x^*(e) - y(e) \geq 8\varepsilon \cdot \sum_{e \in E} x^*(e) \mid \mathcal{E}_{b,c} \right) \leq \frac{1}{2}$$

This implies,

$$\Pr \left( \sum_{e \in E} y(e) \leq (1 - 8\varepsilon) \cdot \sum_{e \in E} x^*(e) \mid \mathcal{E}_{b,c} \right) \leq \frac{1}{2}$$

Thus, the total probability of either [a](#)), [b](#)), or Equation (3) not happening is upper bounded as follows:

$$\Pr(\neg \mathcal{E}_{b,c}) + \Pr \left( \sum_{e \in E} y(e) \leq (1 - 8\varepsilon) \cdot \sum_{e \in E} x^*(e) \mid \mathcal{E}_{b,c} \right) \leq \frac{1}{2} + O\left(\frac{1}{n^{1/\varepsilon^2}}\right).$$

This proves our lemma.  $\square$

**Corollary 16.** Let  $\vec{x} \in \mathcal{M}_G$  be such that  $\sum_{e \in E} x(e) \geq (1 - \varepsilon) \cdot \mu(G)$ . Let  $S$  be the output of SPARSIFICATION( $G, \vec{x}, \varepsilon$ ). Then,  $\mu(S) \geq (1 - \varepsilon) \cdot \sum_{e \in E} x(e)$ .

*Proof.* Note that by Equation (3), we know that there is  $\vec{y} \subseteq \text{supp}(S)$  such that  $\sum_{e \in E} y(e) \geq (1 - \varepsilon) \cdot \sum_{e \in E} x(e)$ . Additionally, from Lemma 15a) we know that for every vertex  $v$ ,

$$\sum_{e \in E_v} y(e) \leq (1 - \varepsilon) \cdot \sum_{e \in E_v} x(e) + \varepsilon \leq 1$$

Finally, we know that for any odd set  $B$  with  $|B| \leq 1/\varepsilon$ ,

$$\sum_{e \in G[B]} y(e) \leq (1 - 2\varepsilon) \cdot \sum_{e \in G[B]} x(e) + \frac{\varepsilon \cdot |B|}{2} \leq (1 - 2\varepsilon) \cdot \frac{|B| - 1}{2} + \frac{\varepsilon \cdot (|B| - 1)}{2} \leq (1 - \varepsilon) \cdot \frac{|B| - 1}{2}$$

Thus, we have,  $\frac{\vec{y}}{1 + \varepsilon} \in \mathcal{M}_G$ .  $\square$

Next, we show two properties of the Algorithm 1 that [Waj20] gives in his paper.

**Observation 17.** [Waj20] Let  $\vec{x} \in \mathcal{M}_G$  be a fractional matching of  $G$ , and let  $S$  be the output of SPARSIFICATION( $G, \vec{x}, \varepsilon$ ). Then,  $|E(H)| = O(d \cdot \mu(G) \cdot \log \frac{n}{\varepsilon})$ .

**Observation 18.** [Waj20] Given a dynamic fractional matching  $\vec{x}$ , Algorithm 1 maintains a sparsifier  $S$  of  $\vec{x}$  satisfying Lemma 15 with update time  $O(\text{poly}(\log n, 1/\varepsilon))$ . Moreover, the guarantees of Algorithm 1 hold against an adaptive adversary.

**Lemma 19.** [DP14] There is a static algorithm, `STATIC-MATCH()` that computes a  $(1 - \varepsilon)$ -approximation to the maximum weight matching in  $O(m \cdot 1/\varepsilon \cdot \log 1/\varepsilon)$  time.

**Theorem 20.** There is a randomized dynamic rounding algorithm for general graphs with amortized update time  $O(\text{poly}(\log n, 1/\varepsilon))$ . The algorithm is robust to adaptive adversaries.

*Proof.* Suppose  $\vec{x}$  is the initial fractional matching with  $\sum_{e \in E} x(e) \geq (1 - \varepsilon) \cdot \text{mwm}(G)$ , then Algorithm 1 initializes the sparsifier in  $O(m \cdot \text{poly}(\log n, 1/\varepsilon))$  total time, and outputs the graph  $S$ . From Observation 17, we know that  $|E(S)| = O(d \cdot \mu(G) \cdot \log \frac{n}{\varepsilon})$ . Thus, we can run a static matching algorithm (say, [DP14]) to get a matching  $M$  with  $|M| \geq (1 - \varepsilon) \cdot \mu(S)$ . By Lemma 15 and Fact 4, we have,

$$\mu(S) \geq (1 - \varepsilon)^2 \sum_{e \in E} x(e) \geq (1 - \varepsilon)^2 \cdot \mu(G)$$

Now, we have to argue how the rounding algorithm deals with updates to the input, that is, the fractional matching. Suppose the input fractional matching  $\vec{x}$  is modified. Then, note that by Observation 18, in  $O(\text{poly}(\log n, 1/\varepsilon))$  time, we can update the sparsifier  $S$  to reflect the change in the fractional matching. As for the integral matching  $M$ , if the deleted edge  $e$  is in  $M$ , then we remove it from  $M$ . If at any point,  $|M| \leq (1 - 2 \cdot \varepsilon) \cdot \sum_{e \in E} x(e)$ , then we recompute  $M$  by running [DP14] on  $S$ . The amortized time now, will be  $O(\text{poly}(\log n, 1/\varepsilon))$ , since we waited for at least  $\varepsilon \cdot \mu(G)$  updates before recomputing the matching. The runtime of [DP14] is  $O(d \cdot \mu(G) \cdot \text{poly}(\log n, 1/\varepsilon))$  by Observation 17. Finally, Algorithm 1 is robust to adaptive adversaries, and [DP14] is a deterministic algorithm. Consequently, the rounding algorithm is robust to adaptive adversaries.  $\square$

### 3 Application

In this section, our goal will be to show the following theorem.

**Theorem 21.** There is a randomized fully adaptive algorithm maintaining a  $(1 - \varepsilon)$ -approximate integral weighted matching in a decremental graph with amortized update time  $\tilde{O}(\varepsilon^{-41} + \frac{n^2}{m} \cdot \varepsilon^{-6})$ .

In their work, [CST23] proved the same theorem, but with guarantees against an output-adaptive adversary. In this section, we will strengthen this to a fully adaptive adversary. Our main tool will be as in the case of [CST23], to use the *entropy regularized matching*. Their proof strategy was as follows:

1. They first consider the sparsification algorithm  $\mathcal{A}$  of [ACC<sup>+</sup>18], which is used to round **unweighted fractional matchings**. They showed that  $\mathcal{A}$  output a sparsifier  $S$  which was also a degree sparsifier. A degree sparsifier is a subgraph, which satisfies a stronger version of Definition 22. This algorithm is robust against output-adaptive adversaries.
2. Next, they considered the *modified entropy regularized matching*  $\vec{x}$  (see Definition 34) and showed the following:
  - (a) We have,  $\sum_{e \in E} x(e) \cdot w(e) \geq (1 - \varepsilon) \cdot \text{mwm}(G)$ .
  - (b) When  $\vec{x}$  is fed into  $\mathcal{A}$ , then the sparsifier  $S$  output by the algorithm not only preserves most of its weight.
  - (c) The fractional matching  $\vec{x}$  can be computed in  $O(m \cdot \text{poly}(\log n, 1/\varepsilon))$  time.

They combine all of these facts to show the output adaptive version of Theorem 21.

We follow a similar strategy: we show the sparsifier output by Algorithm 1 is a weak-degree sparsifier. We consider *entropy regularized matching* problem, and show that when  $\vec{x}$  is fed into SPARSIFICATION(), then most of its weight is preserved. Now, since SPARSIFICATION() is fully-adaptive, we are able to prove a stronger version of the corresponding theorem of [CST23]. In the subsequent section, we will start by giving the definition of a weak degree sparsifier and show SPARSIFICATION() outputs a graph satisfying these properties.

### 3.1 Weak $\varepsilon$ -degree Sparsifier

We consider the modified definition of a degree sparsifier from [CST23].

**Definition 22.** [Weak  $\varepsilon$ -degree sparsifier] Suppose  $\vec{x}$  is a fractional matching of  $G$ , then  $H$  is an  $s$ -sparse, weak  $\varepsilon$ -degree sparsifier of  $G$  if  $|H| \leq s \cdot \sum_{e \in E} x(e)$ , and moreover, there exists a fractional matching  $x^{(H)} \in \mathcal{M}_G$  supported on  $H$ , such that:

1. For all  $v \in V$ , we have,  $x^H(v) \geq x(v) - \varepsilon$ ,
2. For all odd sets  $B \in \mathcal{O}_G^\varepsilon$ , we have,  $\sum_{e \in G[B]} x^H(e) \geq \sum_{e \in G[B]} x(e) - \varepsilon \cdot \frac{|B|-1}{2}$ , and
3. If  $\sum_{e \in E} x(e) \geq (1 - \varepsilon) \cdot \mu(G)$ , then  $\sum_{e \in E} x^H(e) \geq (1 - \varepsilon) \cdot \sum_{e \in E} x(e)$ ,

**Corollary 23.** From Lemma 15, we can deduce that if  $\vec{x}$  is input to SPARSIFICATION() with parameter  $\varepsilon$  and  $d \geq \frac{\log n}{\varepsilon^4}$ , then the output  $S$  is a  $O(\log n)$ -sparse, weak  $\varepsilon$ -degree sparsifier of  $G$  with probability at least 0.5.

In the subsequent section we define the fractional matching problem we will be solving.

### 3.2 Entropy Regularized Matching

The next part is, to show the properties of **entropy regularized matching problem**. Note that [CST23] considered the **modified entropy regularized matching problem**, given in Definition 34. We consider a different version due to the fact that our sparsifier is now weaker, so to prove the required guarantees, we need to solve a slightly different version of the problem than [CST23].

**Definition 24.** The *entropy regularized matching* problem is the problem of finding a fractional matching  $\vec{x} \in \mathcal{M}_{G,\varepsilon}$ , maximizing the following objective function:

$$g(\vec{x}) = \sum_{e \in E} x(e) \cdot w(e) + \delta \cdot \sum_{e \in E} w(e) \cdot x(e) \cdot \log \frac{\gamma}{w(e) \cdot x(e)}$$

**Remark 25.** Note that [CST23] solve the same objective function as in Definition 24, but over the polytope  $\mathcal{M}_G$ .

The next observation is a modified version of Lemma 4.8 in [CST23].

**Observation 26.** If  $\delta \leq \varepsilon/8 \cdot \log n W$  and  $\text{mwm}(G) \leq \gamma \leq \text{mwm}(G) \cdot n \cdot W$ , then and suppose  $\vec{x}^* \in \mathcal{M}_{G,\varepsilon}$  is the optimal solution to  $g$ , then  $g(\vec{x}^*) \leq (1 + \varepsilon) \cdot \text{mwm}(G)$ .

Let  $Z_{E,\gamma}^\delta$  denote the optimal value of the entropy regularized matching over  $\mathcal{M}_{G,\varepsilon}$ . Moreover, let  $\vec{x}^* \in \mathcal{M}_{G,\varepsilon}$  be the unique matching realizing this value. Let  $w^* = \sum_{e \in E} w(e) \cdot x^*(e)$ . We have the following lemma, which is an adaptation of Lemma 6.23 of [CST23].

**Lemma 27.** For an edge  $e = (u, v)$  let  $\mathcal{O}_\varepsilon^e = \{B \in \mathcal{O}_\varepsilon \mid e \in G[B]\}$ . There exist a pair of vertex duals  $\vec{y}$  and odd set duals  $\vec{z}$  such that for every edge  $e$ , we have,

$$x^*(e) = 2^{\frac{1}{\delta} - 1 - \frac{s_e}{\delta \cdot w(e)} + \log \frac{\gamma}{w(e)}},$$

where  $s_e = y_u + y_v + \sum_{B \in \mathcal{O}_\varepsilon^e} z_B$  for  $e = (u, v)$ . Additionally, the optimal objective value  $Z_{E,\gamma}^\delta$  satisfies

$$Z_{E,\gamma}^\delta = \delta \cdot \sum_{e \in E} w(e) \cdot x^*(e) + \sum_{v \in V} y_v + \sum_{B \in \mathcal{O}_\varepsilon} z_B \cdot \frac{|B| - 1}{2}.$$



*Proof.* We consider the Lagrangian of the problem defined in Definition 24:

$$L(\vec{x}, \vec{y}, \vec{z}, \vec{r}) = g(\vec{x}) + \sum_{v \in V} y_v \cdot (1 - \sum_{e \ni v} x(e)) + \sum_{B \in \mathcal{O}_\varepsilon} z_B \cdot \left( \left\lfloor \frac{|B|}{2} \right\rfloor - \sum_{e \in G[B]} x(e) \right) + \sum_{e \in E} x(e) \cdot r_e.$$

Suppose  $(\vec{y}, \vec{z}, \vec{r})$  is the optimal solution of the dual of the problem defined in Definition 24. Then by the KKT Stationarity Condition, we have, that the optimal primal solution is:

$$x^*(e) = 2^{\frac{1}{\delta} - 1 - \frac{s_e}{\delta \cdot w(e)} + \frac{r_e}{\delta \cdot w(e)} + \log \frac{\gamma}{w(e)}}$$

Since  $x^*(e) > 0$  for all  $e$ , using complementary slackness, we have  $r_e = 0$  for all  $e$ . This proves the first part of the claim. Additionally, the *entropy regularized matching problem* satisfies Slater's condition, and consequently has strong duality. Therefore,  $Z_{E,\gamma}^\delta = L(\vec{x}^*, \vec{y}, \vec{z}, \vec{r})$ . By substituting  $(\vec{x}^*, \vec{y}, \vec{z}, \vec{r})$  into  $L$ , we have,

$$Z_{E,\gamma}^\delta = \delta \cdot \sum_{e \in E} w(e) \cdot x^*(e) + \sum_{v \in V} y_v + \sum_{B \in \mathcal{O}_\varepsilon} z_B \cdot \frac{|B| - 1}{2}$$

□

Similarly, we have the following corollary, again adapted from [CST23].

**Corollary 28.** It holds that  $Z_{E,\gamma}^\delta \geq \sum_{v \in V} y_v + \sum_{B \in \mathcal{O}_\varepsilon} z_B \cdot \frac{|B| - 1}{2}$ .

The next observation of [CST23] states that the optimal dual solution  $(\vec{y}, \vec{z})$  corresponding to entropy regularized matching over  $\mathcal{M}_{G,\varepsilon}$  is also a fractional cover for all edges. Given a fractional matching  $\vec{x}$ , we define  $E_\varepsilon(x) = \{e \in E \mid x(e) \geq \frac{\varepsilon}{3n}\}$ .

**Lemma 29.** Let  $\varepsilon > 0$ , let  $w^* \leq \gamma \leq m \cdot w^*$ , and suppose  $\delta \leq \frac{\varepsilon}{8 \cdot \log\left(\frac{n^4 \cdot W}{\varepsilon}\right)}$ . Let  $\vec{y}, \vec{z}$  be the optimal dual solution to the entropy regularized matching problem over  $\mathcal{M}_{G,\varepsilon}$ . Then, for every edge  $e \in E$ ,  $s_e \geq (1 - \varepsilon) \cdot w(e)$ . Moreover, for every edge  $e \in E_\varepsilon(x^*)$ ,  $s_e \leq (1 + \varepsilon) \cdot w(e)$ .

*Proof.* To prove the first part, we invoke Lemma 27, and the fact that  $x^*(e) \leq 1$ , we have,

$$\begin{aligned} \frac{1}{\delta} - 1 - \frac{s_e}{\delta \cdot w(e)} + \log \frac{\gamma}{w(e)} &\leq 0 \\ w(e) - \delta \cdot w(e) + \delta \cdot w(e) \cdot \log \frac{\gamma}{w(e)} &\leq s_e \\ w(e) \cdot (1 - \varepsilon) &\leq s_e \end{aligned}$$

The last inequality follows from the fact that  $\gamma \geq w(e)$ . For  $e \in E_\varepsilon(x^*)$ , we have:

$$\begin{aligned} \log \frac{\varepsilon}{3n} &\leq \frac{1}{\delta} - 1 - \frac{s_e}{\delta \cdot w(e)} + \log \frac{\gamma}{w(e)} \\ s_e &\leq w(e) \cdot (1 - \delta) + \delta \cdot w(e) \log \frac{\gamma}{w(e)} - \delta \cdot w(e) \cdot \log \frac{\varepsilon}{n} \\ &\leq (1 + \varepsilon) \cdot w(e) \end{aligned}$$

This proves our claim. □

Similar to [CST23] did for **modified entropy regularization problem**, we prove the following lemma for the unmodified version.

**Lemma 30.** Let  $\varepsilon > 0$ . Consider any fractional matching  $\vec{x} \in \mathcal{M}_{G,\varepsilon}$ ,  $w^* \leq \gamma \leq m \cdot w^*$ , and  $\delta \leq \frac{\varepsilon}{8 \log\left(\frac{n^4 \cdot W}{\varepsilon}\right)}$ , we have,

$$\sum_{e \in E_\varepsilon(x)} x(e) \cdot w(e) \geq \sum_{e \in E} x(e) \cdot w(e) - \varepsilon \cdot Z_{E,\gamma}^\delta$$

*Proof.* For edges in  $E \setminus E_\varepsilon(x)$ , we have,

$$\begin{aligned}
(1 - \varepsilon) \sum_{e \in E \setminus E_\varepsilon(x)} w(e) \cdot x(e) &\leq \sum_{e \in E \setminus E_\varepsilon(x)} s_e \cdot x(e) \\
&\text{(From Lemma 29)} \\
&= \sum_{v \in V} y_v \cdot \sum_{e \in E_v \setminus E_\varepsilon(x)} x(e) + \sum_{B \in \mathcal{O}_\varepsilon} z_B \cdot \sum_{e \in G[B] \setminus E_\varepsilon(x)} x(e) \\
&\leq \varepsilon \cdot \sum_{v \in V} y_v + \varepsilon \cdot \sum_{B \in \mathcal{O}_\varepsilon} z_B \cdot \frac{|B| - 1}{2} \\
&\leq \varepsilon \cdot Z_{E, \gamma}^\delta \\
&\text{(From Corollary 28)}
\end{aligned}$$

The second last inequality follows from the following line of reasoning:

$$\begin{aligned}
\sum_{e \in E_v \setminus E_\varepsilon(x)} x(e) &\leq \sum_{e \in E_v \setminus E_\varepsilon(x)} \frac{\varepsilon}{n} \leq \varepsilon, \text{ and} \\
\sum_{e \in G[B] \setminus E_\varepsilon(x)} x(e) &\leq \frac{\varepsilon}{n} \cdot |B| \cdot \frac{|B| - 1}{2} \leq \varepsilon \cdot \frac{|B| - 1}{2}
\end{aligned}$$

This shows our claim.  $\square$

**Lemma 31.** Suppose  $\varepsilon > 0$ ,  $w^* \leq \gamma \leq m \cdot \gamma$ , and  $\delta \leq \frac{\varepsilon}{\log\left(\frac{n^4 \cdot W}{\varepsilon}\right)}$ , suppose  $\vec{x} \in \mathcal{M}_{G, \varepsilon}$  with,

$$\sum_{e \in E} w(e) \cdot |x(e) - x^*(e)| \leq \varepsilon \cdot Z_{E, \gamma}^\mu,$$

for any subset  $E' \subset E$  such that,

$$\sum_{e \in E'} w(e) \cdot x(e) \geq (1 - \varepsilon) \cdot \sum_{e \in E} w(e) \cdot x(e),$$

then suppose  $\vec{f}$  is  $\vec{x}$  restricted to  $E_\varepsilon(x)$  and let  $S$  be the output of SPARSIFICATION( $\vec{f}, \varepsilon, d$ ). Then,  $S$  satisfies,

$$\text{mwm}(S) \geq (1 - \varepsilon) \cdot \sum_{e \in E'} w(e) \cdot x(e).$$

*Proof.* Let  $S$  denote the output of SPARSIFICATION( $\vec{y}, \varepsilon, d$ ). Then, by Corollary 23, we know that  $\vec{x}^S$  is a  $O(\log n)$ -sparse, weak  $\varepsilon$ -degree sparsifier. Then, we want to show the following:

$$\sum_{e \in E' \cap E_\varepsilon(x)} w(e) \cdot x^S(e) \geq (1 - \varepsilon) \cdot \sum_{e \in E'} w(e) \cdot x(e)$$

Let  $\vec{x}^* \in \mathcal{M}_{G, \varepsilon}$  be the optimal for the problem defined in Definition 24. Then, we have the following two observations from Lemma 30:

$$\begin{aligned}
\sum_{e \in E' \setminus E_\varepsilon(x)} w(e) \cdot x(e) &\leq \varepsilon \cdot Z_{E', \gamma}^\delta \leq 2 \cdot \varepsilon \cdot Z_{E, \gamma}^\delta \\
\sum_{e \in E' \setminus E_\varepsilon(x^*)} w(e) \cdot x^*(e) &\leq \varepsilon \cdot Z_{E', \gamma}^\delta \leq 2 \cdot \varepsilon \cdot Z_{E, \gamma}^\delta
\end{aligned}$$

Using the premise of the lemma, we have,

$$\begin{aligned}
\sum_{e \in E' \setminus E_\varepsilon(x^*)} w(e) \cdot x(e) &\leq \sum_{e \in E' \setminus E_\varepsilon(x^*)} w(e) \cdot x^*(e) + \sum_{e \in E' \setminus E_\varepsilon(x^*)} w(e) \cdot |x(e) - x^*(e)| \\
&\leq 3 \cdot \varepsilon \cdot Z_{E', \gamma}^\delta
\end{aligned}$$

Consequently, we have,

$$\sum_{e \in E' \setminus E_\varepsilon(\vec{x}^*) \cap E_\varepsilon(\vec{x})} w(e) \cdot x(e) \leq 4 \cdot \varepsilon \cdot Z_{E', \gamma}^\delta. \quad (4)$$

We now focus on proving the main statement mentioned above. We define  $E'_\varepsilon(\vec{x}^*) := E_\varepsilon(\vec{x}^*) \cap E'$ , and similarly we define  $E'_\varepsilon(\vec{x})$ . Additionally, define  $E'_\varepsilon(v) = E_v \cap E'_\varepsilon(\vec{x}) \cap E'_\varepsilon(\vec{x}^*)$ , and  $G'_\varepsilon[B] = G[B] \cap E'_\varepsilon(\vec{x}) \cap E'_\varepsilon(\vec{x}^*)$ .

$$\begin{aligned} \sum_{e \in E'} w(e) \cdot x^S(e) &\geq \sum_{e \in E'_\varepsilon(\vec{x}) \cap E'_\varepsilon(\vec{x}^*)} w(e) \cdot x^S(e) \\ &\geq (1 - \varepsilon) \cdot \sum_{e \in E'_\varepsilon(\vec{x}) \cap E'_\varepsilon(\vec{x}^*)} s_e \cdot x^S(e) \\ &\text{(From Lemma 29)} \\ &\geq (1 - \varepsilon) \cdot \sum_{v \in V} y_v \cdot \sum_{e \in E'_\varepsilon(v)} x^S(e) + \sum_{B \in \mathcal{O}_\varepsilon} z_B \cdot \sum_{e \in G'_\varepsilon[B]} x^S(e) \\ &\geq (1 - \varepsilon) \cdot \left( \sum_{v \in V} y_v \sum_{e \in E'_\varepsilon(v)} x(e) + \sum_{B \in \mathcal{O}_\varepsilon} z_B \sum_{e \in G'_\varepsilon[B]} x(e) \right) - \varepsilon \cdot \left( \sum_{v \in V} y_v + \sum_{B \in \mathcal{O}_\varepsilon} z_B \cdot \frac{|B| - 1}{2} \right) \\ &\text{(From Corollary 23 and Definition 22)} \\ &\geq (1 - \varepsilon) \cdot \sum_{e \in E'_\varepsilon(\vec{x}) \cap E'_\varepsilon(\vec{x}^*)} w(e) \cdot x(e) - \varepsilon \cdot Z_{E, \gamma}^\delta \\ &\text{(From Corollary 28)} \\ &\geq (1 - \varepsilon) \cdot \sum_{e \in E'} w(e) \cdot x(e) - 3 \cdot \varepsilon \cdot Z_{E, \gamma}^\delta \\ &\text{(From Equation (4))} \\ &\geq (1 - 3\varepsilon) \cdot \sum_{e \in E'} w(e) \cdot x(e) \end{aligned}$$

□

**Fact 32.** Suppose  $\vec{z} \in \mathcal{M}_{G, \varepsilon}$ , and  $\vec{x}^*$  be the optimal solution to the *entropy regularized matching problem*. Then,

$$g(\vec{z}) \leq g(\vec{x}^*) - \frac{\delta}{\text{mwm}(G)} \cdot \left( \sum_{e \in E} w(e) \cdot |z(e) - x^*(e)| \right)^2.$$

**Lemma 33.** Suppose  $\vec{y} \in \mathcal{M}_{G, \varepsilon}$  is a  $(1 - \varepsilon)$ -approximate solution to the **entropy regularization problem**, and  $\vec{x}^* \in \mathcal{M}_{G, \varepsilon}$  is the optimal solution to the **entropy regularization problem**. Then,

$$\sum_{e \in E} w(e) \cdot |y(e) - x^*(e)| \leq \varepsilon \cdot Z_{E, \gamma}^\delta$$

*Proof.* Observe that we have,  $g(\vec{y}) \geq (1 - \varepsilon) \cdot g(\vec{x}^*)$ . Consequently, we have,  $g(\vec{x}^*) - g(\vec{y}) \leq \varepsilon \cdot Z_{E, \gamma}^\delta$ . Then, combining the fact that  $Z_{E, \gamma}^\delta \geq \text{mwm}(G)$  and Fact 32, we have,

$$\begin{aligned} \varepsilon \cdot Z_{E, \gamma}^\delta &\geq \frac{\delta}{\text{mwm}(G)} \cdot \left( \sum_{e \in E} w(e) \cdot |z(e) - x^*(e)| \right)^2 \\ \varepsilon \cdot Z_{E, \gamma}^\delta &\geq \sum_{e \in E} w(e) \cdot |z(e) - x^*(e)| \end{aligned}$$

□

Thus, from Lemma 33 and Lemma 30, we have concluded the proof that the entropy regularized fractional matching when fed into SPARSIFICATION() outputs a sparsifier  $S$  that preserves most of its weight.

### 3.3 From Approximate Entropy Regularized Matching to Decremental Matching

In this section, show how to get decremental matching algorithms from **approximate entropy regularized matching**. The proof is almost identical to that given by [CST23]. We just state it here for completeness. We start with the following definition, used by [CST23]:

**Definition 34.** The **modified entropy regularized matching** problem is the problem of finding a fractional matching  $\vec{x} \in \mathcal{M}_G$ , maximizing the following objective function:

$$g(\vec{x}) = \sum_{e \in E} x(e) \cdot w(e) + \delta \cdot \sum_{e \in E} w(e) \cdot x(e) \cdot \log \frac{\gamma}{w(e) \cdot x(e)}$$

Let  $Y_{E,\gamma}^\delta$  denote the optimal solution to this problem.

**Observation 35.** We have,  $Y_{E,\gamma}^\delta \geq Z_{E,\gamma}^\delta \cdot (1 - \varepsilon)$ .

*Proof.* Suppose  $\vec{x}^* \in \mathcal{M}_{G,\varepsilon}$  realizes  $Z_{E,\gamma}^\delta$ . From Fact 4, we have,  $\vec{z} = \frac{\vec{x}^*}{1+\varepsilon} \in \mathcal{M}_G$ . Thus, we have,

$$Y_{E,\gamma}^\delta \geq g(\vec{z}) \geq (1 - \varepsilon) \cdot g(\vec{x}^*) \geq Z_{E,\gamma}^\delta \cdot (1 - \varepsilon).$$

This proves the observation.  $\square$

The next lemma shows that we can compute an approximate solution to the modified entropy maximization problem in  $O(m \cdot \text{poly}(\log n, 1/\varepsilon))$  time.

**Lemma 36.** [CST23] For any  $\varepsilon = \Omega(1/\sqrt{n})$ , there is a randomized algorithm ENTREGMATCHING() that give a  $(1 - \varepsilon)$ -approximation to the **modified entropy regularized matching problem** in  $O(m \cdot \varepsilon^{-6} + n \cdot \varepsilon^{-13})$  time.

Using the above lemma, we will show the following theorem:

**Theorem 21.** There is a randomized fully adaptive algorithm maintaining a  $(1 - \varepsilon)$ -approximate integral weighted matching in a decremental graph with amortized update time  $\tilde{O}(\varepsilon^{-41} + \frac{n^2}{m} \cdot \varepsilon^{-6})$ .

We also need the following lemma, which states that the solution to the *modified entropy regularized matching* is a  $(1 - \varepsilon)$ -approximation to  $\text{mwm}(G)$ .

**Lemma 37** ([CST23]). For  $\mu \leq \frac{\varepsilon}{\log n \cdot W}$  and  $\text{mwm}(G) \leq \gamma \leq n \cdot W \cdot \text{mwm}(G)$ , let  $\vec{x}^*$  be the optimal solution to the *modified entropy regularized matching* problem, then we have,

$$\sum_{e \in E} w(e) \cdot x^*(e) \geq (1 - \varepsilon) \cdot \text{mwm}(G).$$

Similarly, suppose  $\vec{x}$  is a  $(1 - \varepsilon)$ -approximate solution to the *modified entropy regularized matching* problem, then we have,

$$\sum_{e \in E} w(e) \cdot x(e) + \varepsilon \cdot \text{mwm}(G) \geq (1 - \varepsilon) \cdot g(\vec{x}^*) \geq (1 - \varepsilon) \cdot \sum_{e \in E} w(e) \cdot x^*(e) \geq (1 - \varepsilon)^2 \cdot \text{mwm}(G)$$

In order to prove Theorem 21, we state the main algorithm, and then show its correctness and runtime.

**Lemma 38.** Algorithm 2 maintains an integral matching  $M$  at all times such that  $w(M) \geq (1 - \varepsilon) \cdot \text{mwm}(G)$ .

*Proof.* Note that from Lemma 36, we can conclude that the fractional matching  $\vec{y}$  output by ENTREGMATCH( $G, \varepsilon$ ) has the property that:

$$g(\vec{y}) \geq (1 - \varepsilon) \cdot g(\vec{y}^*),$$

where  $\vec{y}^* \in \mathcal{M}_G$  is the optimal solution to the **modified entropy regularization** problem. This satisfies the premise of Lemma 37. Consequently, from that we can deduce:

$$\sum_{e \in E} w(e) \cdot y(e) \geq (1 - \varepsilon)^2 \cdot \text{mwm}(G).$$

---

**Algorithm 2** DECREMENTAL MATCHING

---

**Input:**  $G$ , weights  $w$ , and a precision parameter  $\varepsilon > 0$ 

```
1: procedure INITIALIZE( $G, E$ )
2:   Initialize  $G \leftarrow G_0$  ▷  $G$  refers to the current graph, and  $G_0$  is the initial graph.
3:   Initialize  $E \leftarrow E_0$  ▷  $E$  refers to the current edge set, and  $E_0 = E(G_0)$ .
4:   COUNTERM ← 0 ▷ Counts the total weight deleted from  $M$ 
5:   COUNTERX ← 0 ▷ Counts the total weight deleted from  $\vec{x}$ 
6: end procedure
7: procedure REBUILD
8:    $\vec{y} \leftarrow \text{ENTREGMATCH}(G, \varepsilon)$ 
9:   Let  $\vec{x}$  be  $\vec{y}$  restricted to  $E_\varepsilon(\vec{y})$ .
10:   $\mu^* \leftarrow \sum_{e \in E} w(e) \cdot x(e)$  ▷  $\mu^* \geq (1 - \varepsilon) \cdot \text{mwm}(G)$ 
11: end procedure
12: procedure ROUND
13:   $S \leftarrow \text{SPARSIFICATION}(\vec{x}, \varepsilon)$ 
14:   $M \leftarrow \text{STATIC-MATCH}(S, \varepsilon)$ 
15: end procedure
16: procedure DELETION( $e$ )
17:   $E \leftarrow E \setminus e$ 
18:  if  $e \in \text{supp}(\vec{x})$  then
19:    Delete  $e$  from  $\text{supp}(\vec{x})$ 
20:    COUNTERX ← COUNTERX +  $w(e) \cdot x(e)$ 
21:  end if
22:  if COUNTERX  $\geq \varepsilon \cdot \mu^*$  then
23:    REBUILD()
24:  end if
25:  if  $e \in M$  then
26:     $M \leftarrow M \setminus \{e\}$ 
27:    COUNTERM ← COUNTERM + 1
28:  end if
29:  if COUNTERM  $\geq \varepsilon \cdot \mu^*$  then
30:    COUNTERM ← 0
31:    ROUND()
32:  end if
33: end procedure
```

---

Note that  $\vec{x}$  is  $\vec{y}$  restricted to  $E_\varepsilon(\vec{y})$ . Therefore, from Lemma 30 and Observation 26, we can then conclude the following:

$$\sum_{e \in E} w(e) \cdot x(e) \geq \sum_{e \in E} w(e) \cdot y(e) - \varepsilon \cdot Z_{E,\gamma}^\delta \geq (1 - 3\varepsilon) \cdot \text{mwm}(G)$$

Additionally, note that the algorithm processes deletions, and rebuilds when  $\sum_{e \in E} w(e) \cdot x(e)$  has dropped by a  $(1 - \varepsilon)$  factor. Thus, each time the algorithm is maintaining a fractional matching  $\vec{x}$  such that  $\sum_{e \in E} w(e) \cdot x(e) \geq (1 - \varepsilon)^3 \cdot \text{mwm}(G)$ . Next, the algorithm runs SPARSIFICATION( $\vec{x}, \varepsilon$ ) to compute a sparsifier  $S$ . Observe  $\vec{y}$  satisfies the premise of Lemma 33. Consequently, we have, Observation 35,

$$\sum_{e \in E} w(e) \cdot |y(e) - x^*(e)| \leq \varepsilon \cdot Z_{E,\gamma}^\delta$$

Note that we feed  $\vec{x}$ , which is  $\vec{y}$  restricted to  $E_\varepsilon(\vec{y})$  into SPARSIFICATION(). Thus, from Lemma 31, we have,

$$\text{mwm}(S) \geq (1 - \varepsilon) \cdot \sum_{e \in E} w(e) \cdot y(e)$$

Therefore, from Lemma 19 we can conclude that the matching  $M$  output by STATIC-MATCH( $S, \varepsilon$ ) has  $w(M) \geq (1 - \varepsilon) \cdot \text{mwm}(S) \geq (1 - \varepsilon)^2 \cdot \sum_{e \in E} w(e) \cdot y(e) \geq (1 - \varepsilon)^3 \cdot \text{mwm}(G)$ . Here, Lemma 37 gives us the last inequality. This shows the lemma.  $\square$

Now, we show the runtime, and for that we need the following lemma.

**Lemma 39.** ([CST23]) If ENTREGMATCH() returns a  $(1 - \varepsilon)$  approximate solution to  $g(\vec{x})$ , then the REBUILD() will be called at most  $O(\text{poly}(\log n, 1/\varepsilon))$  times before  $\text{mwm}(G)$  drops from at least  $\mu^*$  to at most  $\mu^* \cdot (1 - \varepsilon)$ .

**Lemma 40.** Algorithm 2 has an amortized updated time of  $\tilde{O}\left(\text{poly}(1/\varepsilon) + \frac{n^2}{m} \cdot \text{poly}(1/\varepsilon)\right)$ .

*Proof.* First, observe the runtime of each of the procedures and each time they are invoked:

1. For the procedure REBUILD(), we can conclude from Lemma 36 that each time it is run it takes time  $\tilde{O}(m \cdot \text{poly}(\log n, 1/\varepsilon))$ . This procedure is called each time COUNTERX increases to  $\varepsilon \cdot \mu^*$  or  $\sum_{e \in E^*} w(e) \cdot x(e)$  has dropped by  $(1 - \varepsilon)$  amount. From Lemma 39, we can conclude REBUILD( $\vec{x}, \varepsilon$ ) is called at most  $O(\text{poly}(\log n, 1/\varepsilon))$  times. Thus, the total runtime of this procedure over  $m$  deletions is at most  $O(m \cdot \text{poly}(\log n, 1/\varepsilon))$ . That is, till the matching weight drops from  $\mu^*$  to 1.
2. Next we consider the procedure ROUND(). Note that from Observation 17 we can conclude that  $|E(S)| = O(n \cdot \text{poly}(\log n, 1/\varepsilon))$ . Moreover, the time taken to maintain it is  $O(\text{poly}(\log n, 1/\varepsilon))$ . Thus, from Lemma 19 we can conclude that STATIC-MATCH( $S, \varepsilon$ ) and consequently, the procedure ROUND() take time  $O(n \cdot \text{poly}(\log n, 1/\varepsilon))$  each time they are invoked. This is invoked each time COUNTERM drops by  $\varepsilon \cdot \mu^*$  amount. Since  $x(e) \geq \varepsilon/n$  for all  $e \in E$ , this also translates into the adversary deleting a weight of at least  $\varepsilon^2 \cdot \mu^*/n$  from the fractional matching  $\vec{x}$ . Consequently, till before the next time REBUILD() is invoked, ROUND() is invoked  $O(n/\varepsilon^2)$  times. Since the total number of times REBUILD() is called is at most  $O(\text{poly}(\log n, 1/\varepsilon))$ , this implies that the total number of times ROUND() is called is at most  $O(n \cdot \text{poly}(\log n, 1/\varepsilon))$ , and therefore, the total runtime contributed by this procedure is  $O(n^2 \cdot \text{poly}(\log n, 1/\varepsilon))$ .
3. The other steps in the DELETION() procedure take  $O(1)$  time per edge to implement. So the total time over all updates is at most  $O(m)$ .

This concludes the proof.  $\square$

## 4 Acknowledgements

Thank you to David Wajc for numerous helpful email exchanges. Thank you to Rishabh Dudeja, Madhusudhan Raman and Aditya Potukuchi for encouragement.

## References

- [ABD22] Sepehr Assadi, Aaron Bernstein, and Aditi Dudeja. Decremental matching in general graphs. In Mikolaj Bojanczyk, Emanuela Merelli, and David P. Woodruff, editors, *49th International Colloquium on Automata, Languages, and Programming, ICALP 2022, July 4-8, 2022, Paris, France*, volume 229 of *LIPICs*, pages 11:1–11:19. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2022.
- [ACC<sup>+</sup>18] Moab Arar, Shiri Chechik, Sarel Cohen, Cliff Stein, and David Wajc. Dynamic matching: Reducing integral algorithms to approximately-maximal fractional algorithms. In Ioannis Chatzigiannakis, Christos Kaklamanis, Dániel Marx, and Donald Sannella, editors, *45th International Colloquium on Automata, Languages, and Programming, ICALP 2018, July 9-13, 2018, Prague, Czech Republic*, volume 107 of *LIPICs*, pages 7:1–7:16. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2018.
- [BCH20] Sayan Bhattacharya, Deeparnab Chakrabarty, and Monika Henzinger. Deterministic dynamic matching in  $O(1)$  update time. *Algorithmica*, 82(4):1057–1080, 2020.
- [BGS20] Aaron Bernstein, Maximilian Probst Gutenberg, and Thatchaphol Saranurak. Deterministic decremental reachability, scc, and shortest paths via directed expanders and congestion balancing. In *2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)*, pages 1123–1134. IEEE, 2020.
- [BHI15] Sayan Bhattacharya, Monika Henzinger, and Giuseppe F. Italiano. Deterministic fully dynamic data structures for vertex cover and matching. In *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA '15*, page 785–804, USA, 2015. Society for Industrial and Applied Mathematics.
- [BHN16] Sayan Bhattacharya, Monika Henzinger, and Danupon Nanongkai. New deterministic approximation algorithms for fully dynamic matching. In *Proceedings of the Forty-Eighth Annual ACM Symposium on Theory of Computing, STOC '16*, page 398–411, New York, NY, USA, 2016. Association for Computing Machinery.
- [BHN17] Sayan Bhattacharya, Monika Henzinger, and Danupon Nanongkai. Fully dynamic approximate maximum matching and minimum vertex cover in  $O(\log^3 n)$  worst case update time. In Philip N. Klein, editor, *Proceedings of the Twenty-Eighth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2017, Barcelona, Spain, Hotel Porta Fira, January 16-19*, pages 470–489. SIAM, 2017.
- [BK21] Sayan Bhattacharya and Peter Kiss. Deterministic rounding of dynamic fractional matchings. In Nikhil Bansal, Emanuela Merelli, and James Worrell, editors, *48th International Colloquium on Automata, Languages, and Programming, ICALP 2021, July 12-16, 2021, Glasgow, Scotland (Virtual Conference)*, volume 198 of *LIPICs*, pages 27:1–27:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [BK23] Joakim Blikstad and Peter Kiss. Incremental  $(1-\epsilon)$ -approximate dynamic matching in  $o(\text{poly}(1/\epsilon))$  update time. In Inge Li Gørtz, Martin Farach-Colton, Simon J. Puglisi, and Grzegorz Herman, editors, *31st Annual European Symposium on Algorithms, ESA 2023, September 4-6, 2023, Amsterdam, The Netherlands*, volume 274 of *LIPICs*, pages 22:1–22:19. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2023.
- [BKSW23] Sayan Bhattacharya, Peter Kiss, Aaron Sidford, and David Wajc. Near-optimal dynamic rounding of fractional matchings in bipartite graphs. *CoRR*, abs/2306.11828, 2023.
- [CST23] Jiale Chen, Aaron Sidford, and Ta-Wei Tu. Entropy regularization and faster decremental matching in general graphs. *CoRR*, abs/2312.09077, 2023.

- [DP09] Devdatt P. Dubhashi and Alessandro Panconesi. *Concentration of Measure for the Analysis of Randomized Algorithms*. Cambridge University Press, 2009.
- [DP14] Ran Duan and Seth Pettie. Linear-time approximation for maximum weight matching. *Journal of the ACM (JACM)*, 61(1):1–23, 2014.
- [JJST22] Arun Jambulapati, Yujia Jin, Aaron Sidford, and Kevin Tian. Regularized box-simplex games and dynamic decremental bipartite matching. In Mikolaj Bojanczyk, Emanuela Merelli, and David P. Woodruff, editors, *49th International Colloquium on Automata, Languages, and Programming, ICALP 2022, July 4-8, 2022, Paris, France*, volume 229 of *LIPICs*, pages 77:1–77:20. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2022.
- [Kis22] Peter Kiss. Deterministic dynamic matching in worst-case update time. In Mark Braverman, editor, *13th Innovations in Theoretical Computer Science Conference, ITCS 2022, January 31 - February 3, 2022, Berkeley, CA, USA*, volume 215 of *LIPICs*, pages 94:1–94:21. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2022.
- [Waj20] David Wajc. Rounding dynamic matchings against an adaptive adversary. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing*, pages 194–207, 2020.