

Groups generated by pattern avoiding permutations

Marilena Barnabei*

P.A.M.

Università di Bologna, 40126, ITALY

marilena.barnabei@unibo.it

Niccolò Castronuovo

Liceo “A. Einstein”, Rimini, 47923, ITALY

castronuovoniccolo@gmail.com

Matteo Silimbani

Istituto Comprensivo “E. Rosetti”, Forlimpopoli, 47034, ITALY

matteosilimbani@icrosetti.istruzioneer.it

Abstract

We study groups generated by sets of pattern avoiding permutations. In the first part of the paper we prove some general results concerning the structure of such groups. In the second part we carry out a case-by-case analysis of groups generated by permutations avoiding few short patterns.

Keywords: Pattern avoiding permutation, permutation group, generated group.

MSC2020: 05A05, 20B05 (primary); 20B10 (secondary).

1 Introduction

The symmetric group S_n over n elements is ubiquitous in mathematics, indeed, it is studied both from a group-theoretical and a combinatorial point of view.

*corresponding author

In the first case the elements of S_n are thought as bijections acting on the set $\{1, 2, \dots, n\}$ and are written as sequences of cycles, in the second case they are considered as words containing all the symbols from 1 to n once and are written in one-line notation as the sequence of such symbols.

The group-theoretical study of the symmetric group embodies the wide fields of representation theory of the symmetric group (see, e.g., [19]) and permutation groups (see, e.g., [7]). The combinatorics of permutations (see, e.g., [5]) includes many topics among whom one of the most recently developed is the study of pattern avoiding permutations (see, e.g., [15]).

The two aspects are rarely considered together, due to the different nature of the two approaches. Papers that deal with both the perspectives are, for instance, [1, 4, 6, 14, 16, 17].

In this paper we consider the problem of determining the group generated by a set of pattern avoiding permutations. This topic has never been investigated, to the best of our knowledge.

We begin with introducing some notations and presenting some preliminary results (Sections 2 and 3).

Then, given a set of patterns T , we consider the subset $S_n(T)$ of the symmetric group S_n consisting of permutations avoiding each pattern in T , and turn our attention to the group G_n generated by $S_n(T)$.

When dealing with such groups some questions naturally arise. First of all, under which conditions the sequence $(G_n)_{n \geq 0}$ is eventually constant and which kind of groups arise?

In Section 4 we find a set of patterns such that $(G_n)_{n \geq 0}$ is eventually constant and equal to an assigned symmetric group S_k . In Section 5 we show that the alternating group can not be obtained in this way and that the only non-trivial abelian groups that arise in this way, are \mathbb{Z}_2 and $\mathbb{Z}_2 \times \mathbb{Z}_2$. The proof of such results take advantage of a characterization of those sets of patterns such that the sequence (G_n) has polynomial growth.

A further question is under which conditions $G_n = S_n$ for every sufficiently large n .

In Section 6 we single out a fairly general condition under which this happens. Sections 7 and 8 are devoted to the analysis of the groups generated by $S_n(T)$ when T consists of few patterns or patterns of small length. In the last section we present some open problems.

2 Notations and definitions

Firstly we recall some notations from group theory. We denote by

- S_n the symmetric group, i.e., the set of permutations of $[n] = \{1, 2, \dots, n\}$, with the operation of composition, and by S the union of all these sets,

$$S = \bigcup_{n \geq 0} S_n,$$

- A_n the alternating group, i.e., the subgroup of S_n of even permutations,
- \mathbb{Z}_n the cyclic group with n elements,
- D_n the dihedral group with $2n$ elements, i.e., the group of symmetries of the regular n -agon.

Recall that every element π of S_n can be written either as a product of cycles, or in one-line notation $\pi = \pi_1 \pi_2 \dots \pi_n$ as the sequence of images of the elements $1, 2, \dots, n$. The presence or not of the brackets $(,)$ will make clear which notation we are using. We write products from right to left, that is, if $\pi, \sigma \in S_n$, $\pi\sigma$ is the permutation obtained by applying σ first.

Given a set $A \subseteq S_n$, the symbol $\langle A \rangle$ will denote the group generated by A .

We will use the following notation for special elements of S_n .

- the identity permutation will be denoted by id_n ,

$$\text{id}_n = 1 \ 2 \ \dots \ n,$$

- the decreasing permutation will be denoted by ψ_n ,

$$\psi_n = n \ n - 1 \ \dots \ 2 \ 1 = \prod_{i=1}^{\lfloor \frac{n}{2} \rfloor} (i, n + 1 - i).$$

We will use the symbols id and ψ when the size is clear from the context.

If $\pi \in S_n$ we will say that π has *length* n and write $|\pi| = n$. A permutation π such that $\pi = \pi^{-1}$ is called an *involution*.

The *reverse*, the *complement* and *reverse-complement* of the permutation π are the permutations $\pi^r = \pi\psi$, $\pi^c = \psi\pi$ and $\psi^r\pi = \psi\pi\psi$, respectively.

A permutation $\pi \in S_n$ *contains the pattern* $\tau \in S_k$ if there are indices i_1, i_2, \dots, i_k such that the subsequence $\pi_{i_1} \pi_{i_2} \dots \pi_{i_k}$ is order isomorphic to τ , and we write $\pi \geq \tau$.

Otherwise we say that π *avoids the pattern* τ .

The set of permutations avoiding all the patterns in the set $T = \{\tau_1, \tau_2, \dots\}$ is denoted by $S_n(T)$ or, with a slight abuse of notation, by $S_n(\tau_1, \tau_2, \dots)$.

We set $S(T) = \cup_{n \geq 0} S_n(T)$.

Notice that the set S with the containment relation \leq is a poset. A downset in this order is called a *permutation class* or simply a *class*. Given a class \mathcal{C} , let $\mathcal{C}_n = \mathcal{C} \cap S_n$ for every $n \geq 0$. A class can be specified by its *basis*, the minimal permutations not in the class. If T is the basis of a class \mathcal{C} , then $\mathcal{C} = S(T)$. Observe that the basis of a class can be infinite, since the poset (S, \leq) contains infinite antichains.

3 Preliminary results

Now we state some basic Lemmas that will be useful in the following.

Lemma 3.1. *Let T and T' be two subsets of S , with $T \subseteq T'$. Then $\langle S_n(T') \rangle \subseteq \langle S_n(T) \rangle$.*

Proof. Immediate consequence of the definition of pattern containment. \square

Lemma 3.2. *Let $\sigma_1, \dots, \sigma_k$ and τ_1, \dots, τ_k be permutations of any length such that $\sigma_i \leq \tau_i$ for every i . Then $\langle S_n(\sigma_1, \dots, \sigma_k) \rangle \subseteq \langle S_n(\tau_1, \dots, \tau_k) \rangle$.*

Proof. This fact is a straightforward consequence of the following inclusion: $S_n(\sigma_1, \dots, \sigma_k) \subseteq S_n(\tau_1, \dots, \tau_k)$. \square

Given a set of permutations A , let A^r be the set $\{\sigma^r \mid \sigma \in A\}$. The sets A^c , A^{rc} , A^{-1} are defined similarly. Notice that $S_n(T)^r = S_n(T^r)$, and the same is true for the complement, the reverse-complement and the inverse map. Moreover, since $(\pi^r)^{-1} = (\pi^{-1})^c$,

$$(S_n(T^r))^{-1} = S_n((T^r)^{-1}) = S_n((T^{-1})^c).$$

A similar identity holds with r and c interchanged.

Lemma 3.3. *Let $T \subseteq S$ be a set of permutations of any length. Then*

$$\langle S_n(T) \rangle = \langle S_n(T^{-1}) \rangle = \langle S_n(T) \cup S_n(T^{-1}) \rangle,$$

and

$$\langle S_n(T) \rangle \cong \langle S_n(T^{rc}) \rangle.$$

Proof. Given $\pi \in S_n$, $\pi^{rc} = \psi\pi\psi$. This fact implies the second assertion. The first assertion is trivial. \square

Lemma 3.4. *Let $T \subseteq S$ be a set of permutations of any length. If $\psi_k \notin T$ for every $k \geq 1$, then*

$$\langle S_n(T) \rangle = \langle S_n(T) \cup S_n(T^r) \cup S_n(T^c) \cup S_n(T^{rc}) \rangle.$$

Proof. Notice that if $\psi_k \notin T$ for every $k \geq 1$, then $\psi_n \in S_n(T)$.

Since $\sigma^r = \sigma\psi$, $\sigma^c = \psi\sigma$ and $\sigma^{rc} = \psi\sigma\psi$, the assertion follows. □

We recall a well-known result that presents some families of generating sets for the group S_n .

Lemma 3.5. *The following sets generate S_n , for every $n \geq 1$.*

- $\{(j, j + 1), 1 \leq j \leq n - 1\}$.
- $\{(1, j), 2 \leq j \leq n\}$.
- $\{(a, b), (1, 2, \dots, n)\}$ if $b - a$ and n are coprime.

In the following we will use the notion of *direct product* and *semidirect product* of groups (we refer to [12] for definitions and notations). In particular we will exploit the following result stated and proved in [12, p. 69].

Lemma 3.6. *If G is a group, G is the semidirect product of its subgroups N and H if and only if the following three conditions are verified*

- $N \trianglelefteq G$,
- $G = NH$,
- $N \cap H = \{id\}$.

Finally we recall a lemma which expresses a connection between the notion of pattern containment and of permutation group. See [17] for the proof.

Lemma 3.7. *Let G be any subgroup of a symmetric group S_k . Then $S_n(S_k \setminus G)$ is a subgroup of S_n .*

4 Generating a fixed symmetric group S_k .

First of all we ask the following question.

Question 4.1. *Let G be any finite group. Is it possible to find a subset T of S and a positive integer n such that $\langle S_n(T) \rangle = G$?*

Notice that if T can depend on n , then the answer to the previous question is trivially true for every G . In fact, if G is a finite group, by Cayley's theorem there exists n such that there is an isomorphic copy \widehat{G} of G into S_n . Then

$$G \cong \langle S_n(S_n \setminus \widehat{G}) \rangle = S_n(S_n \setminus \widehat{G}).$$

If the set of patterns T must be independent of n the question is more subtle.

The next theorem shows that the question can be answered affirmatively for $G = S_k$, for any k .

Theorem 4.2. *We have*

$$\langle S_n(132, 231, 321, k12 \dots k-2 \ k-1) \rangle \cong S_{k-1} \quad \forall n \geq k.$$

Proof. The set $S_n(132, 231, 321, k12 \dots k-2 \ k-1)$ contains precisely the permutations id and $t123 \dots t-1 \ t+1 \dots n-1 \ n$ for $2 \leq t \leq k-1$. In fact, if π is in $S_n(132, 231, 321, k12 \dots k-2 \ k-1)$, the first symbol $t = \pi_1$ must be smaller or equal to $k-1$ in order to avoid the patterns 321 and $k12 \dots k-2 \ k-1$. If $t = 1$, $\pi = \text{id}$ because the other symbols must be in increasing order, since π avoids 132. If $t \geq 2$, the second symbol π_2 must be equal to 1 in order to avoid 231 and 321. All the other symbols must be in increasing order, in order to avoid 132.

In particular $S_n(132, 231, 321, k12 \dots k-2 \ k-1)$ consists of the identity and the cycles $(1, 2)$, $(1, 3, 2)$, $(1, 4, 3, 2)$, \dots $(1 \ k-1 \ k-2 \dots 2)$. Hence, it generates a group isomorphic to S_{k-1} . □

5 Alternating groups and abelian groups

In this section we consider a refinement of Question 4.1.

Question 5.1. *Let G be any finite group. Is it possible to find a subset T of S and a positive integer n such that $\langle S_n(T) \rangle = G$ for every $n \geq n_0$?*

We give a negative answer to the previous question for some families of groups.

We need some preliminary results. Firstly notice that, if $\langle S_n(T) \rangle$ is isomorphic to a fixed group G for every $n \geq n_0$, the cardinality of $S_n(T)$ must be bounded above by the order of G .

The set of pattern avoiding permutations with limited growth rate are well studied. See e.g. [2, 3, 10, 11, 13, 18, 22, 21], to cite only a few.

Now we recall some results about classes of permutations with limited growth rate. First of all we introduce some definitions, following [10].

An *interval* in a permutation is a sequence of contiguous entries whose values form an interval of natural numbers. A *monotone interval* is an interval in which the entries are increasing or decreasing. Given a permutation σ of length m and nonempty permutations $\alpha_1, \dots, \alpha_m$, the *inflation* of σ by

$\alpha_1, \dots, \alpha_m$ is the permutation $\pi = \sigma[\alpha_1, \dots, \alpha_m]$ obtained by replacing the i -th entry of σ by an interval that is order isomorphic to α_i , while maintaining the relative order of the intervals themselves. For example,

$$3142[1, 321, 1, 12] = 6\ 321\ 7\ 45.$$

Notice that we allow inflations by the empty permutation.

A *peg permutation* $\tilde{\rho}$ is a permutation ρ where some elements are labeled by the symbol $+$ and some other elements are labeled by the symbol $-$. For example $\tilde{\rho} = 3^+1^-24^-$ is a peg permutation whose underlying (non-pegged) permutation is 3124.

The *grid class* of the peg permutation $\tilde{\rho}$, where $\rho = \rho_1 \dots \rho_n$, denoted by $\text{Grid}(\tilde{\rho})$, is the set of all permutations which may be obtained inflating ρ by monotone intervals of type determined by the signs of $\tilde{\rho}$: ρ_i may be inflated by an increasing (resp., decreasing) possibly empty interval if ρ_i is labeled with a $+$ (resp., $-$) while it may only be inflated by a single entry (or the empty permutation) if ρ_i is not labeled.

For example the permutation 45621387 is an element of the grid class $\text{Grid}(3^+1^-24^-)$.

Given a set \tilde{P} of peg permutations, we denote the union of their corresponding grid classes by

$$\text{Grid}(\tilde{P}) = \cup_{\tilde{\rho} \in \tilde{P}} \text{Grid}(\tilde{\rho}).$$

We will need the following result ([10, Thm. 1.3], which in turn is a consequence of some results in [2] and in [11]).

Theorem 5.2. *For a permutation class \mathcal{C} the following are equivalent:*

1. *The cardinality $|\mathcal{C}_n|$ is a polynomial in n , for sufficiently large n .*
2. *$|\mathcal{C}_n| < F_n$ for some n , where F_n is the n -th Fibonacci numbers.*
3. *$\mathcal{C} = \text{Grid}(\tilde{P})$, where \tilde{P} is a finite set of peg permutations.*

Example 5.3. Consider the class $\mathcal{C} = S(123, 231, 312)$. As we will see in the proof of Theorem 7.1,

$$S(123, 231, 312) = \{k\ k-1 \dots 2\ 1\ n\ n-1 \dots k+1, \quad 1 \leq k \leq n\}.$$

The class \mathcal{C} can be written as $\mathcal{C} = \text{Grid}(\tilde{P})$, where \tilde{P} contains only the peg permutation 1^-2^- .

An immediate consequence of the previous Theorem is the following Proposition that describes the structure of permutation classes with growth rate bounded above by a constant.

Proposition 5.4. *For a permutation class \mathcal{C} the following are equivalent.*

1. *For every $n \geq 1$, $|\mathcal{C}_n| < K$ where K is a constant independent of n .*
2. *$\mathcal{C} = \text{Grid}(\tilde{P})$, where \tilde{P} is a finite set of peg permutations each of which has at most one labeled element.*
3. *For sufficiently large n , $|\mathcal{C}_n| = D$, where D is a constant independent of n .*

Proof. [1. \implies 2.] If for every $n \geq 1$, $|\mathcal{C}_n| < K$, where K is a constant independent of n , the growth rate of $|\mathcal{C}_n|$ is polynomial (the constant polynomial K). Hence, by the previous Theorem, $\mathcal{C} = \text{Grid}(\tilde{P})$ where \tilde{P} is a finite set of peg permutations. Suppose that in \tilde{P} there is a peg permutation $\tilde{\rho}$ with more than one element labeled. Let a and b be two labeled elements in $\tilde{\rho}$. We can inflate a and b by increasing or decreasing sequences of arbitrary length. Hence $\text{Grid}(\tilde{P}) \cap S_n$ would have at least $n + 1$ elements of the form $t_1 t_2 \dots t_k s_1 s_2 \dots s_{n-k}$, with $0 \leq k \leq n$, where $t_1 t_2 \dots t_k$ is the monotone sequence obtained by inflating a , $s_1 s_2 \dots s_{n-k}$ is the monotone sequence obtained by inflating b , and all the other elements of ρ have been inflated by the empty interval. This is a contradiction, because the cardinality of $\text{Grid}(\tilde{P}) \cap S_n$ is bounded above by the constant K .

[2. \implies 3.]

If $\mathcal{C} = \text{Grid}(\tilde{P})$ where \tilde{P} is a finite set of peg permutations each of which has at most one labeled element, then $\mathcal{C}_n = \text{Grid}(\tilde{P}) \cap S_n$ has a constant number of elements, for n sufficiently large. In fact, let $\tilde{\rho} = a_1 \dots a_r b^+ c_1 \dots c_s$ an element of \tilde{P} . If $n \geq r + s$, such an element gives rise to exactly 2^{r+s} in \mathcal{C}_n depending on how many of the a_i 's or c_j 's are inflated by the empty permutation. The same holds when the element b has label $-$.

[3. \implies 1.] Trivial. □

Example 5.5. Consider the class $\mathcal{C} = S(132, 312, 321, 2314)$.

It is easily seen that

$$S_n(132, 312, 321, 2314) = \{2\ 3 \dots n\ 1, \quad 2\ 1\ 3 \dots n, \quad \text{id}_n\}$$

for every $n \geq 3$.

Hence $|\mathcal{C}_n| = 3$ for $n \geq 3$. The class \mathcal{C} can be written as $\mathcal{C} = \text{Grid}(\tilde{P})$ where $\tilde{P} = \{213^+, 2^+1\}$.

Notice also that the group generated by $S_n(132, 312, 321, 2314)$ is S_n by Lemma 3.5.

Theorem 5.6. *Let $T \subseteq S$ and let G be a finite group. Suppose that there exists n_0 in \mathbb{N} such that $\langle S_n(T) \rangle = G$ for every $n \geq n_0$.*

Then $S(T) = \text{Grid}(\tilde{P})$, where every peg permutation in \tilde{P} is either unlabeled, or of the form

$$\tilde{\rho} = a_1 \dots a_r (r+1)^+ b_1 \dots b_s,$$

or

$$\tilde{\rho} = b_1 \dots b_s (r+1)^- a_1 \dots a_r,$$

where, for every i and j , $a_i < r+1$ and $b_j > r+1$.

In particular, for n sufficiently large, every $\pi \in S_n(T)$ is either of the form

$$\pi = 123[\tau, \text{id}, \sigma]$$

or of the form

$$\pi = 321[\sigma, \psi, \tau]$$

where $\tau \leq a_1 \dots a_r$, $\sigma \leq b_1 \dots b_s$ and $|\tau| + |\text{id}| + |\sigma| = n$.

Proof. Since $\langle S_n(T) \rangle = G$ for every $n \geq n_0$, the cardinality of $S_n(T)$ is bounded above by a constant $|G| = K$ independent of n . By the previous corollary, each element of the class $S(T)$ can be obtained inflating a peg permutation with at most one labeled element.

Let $\tilde{\rho}$ be a peg permutation with exactly one labeled element.

- Suppose

$$\tilde{\rho} = a_1 \dots a_r c^+ b_1 \dots b_s,$$

where c is the only labeled element, and suppose that there exists $1 \leq i \leq r$ with $a_i > c$. Then, inflating such a_i by the permutation 1 , every other a_j and b_j by the empty permutation and c by the increasing permutation (id), we obtain the element $n \ 1 \ 2 \ \dots \ n-1 \in S_n(T)$ for every n . But this is a cycle of order n . Hence $\langle S_n(T) \rangle$ contains at least n elements and its cardinality is not constant.

Similarly, if there exists $1 \leq i \leq s$ with $b_i < c$, the set $S_n(T)$ contains the cycle $(1, 2, 3, \dots, n)$ of order n .

- Suppose otherwise that

$$\tilde{\rho} = b_1 \dots b_s c^- a_1 \dots a_r,$$

where c is the only labeled element, and suppose that there exists $1 \leq i \leq s$ with $b_i < c$. Then, inflating such b_i by the empty permutation or by the permutation 1, every other b_j and a_j by the empty permutation and c by a decreasing permutation (ψ), we obtain that the elements ψ_n and $\alpha = 1 \ n \ n-1 \ \dots \ 2 \in S_n(T)$ for every n . But $\alpha\psi_n = (1, 2, \dots, n)$, a cycle of order n , and hence $\langle S_n(T) \rangle$ contains at least n elements and its cardinality would not be constant.

Similarly, if there exists $1 \leq i \leq r$ with $a_i > c$, the set $S_n(T)$ contains the permutations ψ_n and $\beta = n-1 \ n-2 \ \dots \ 1 \ n$ whose composition $\psi_n\beta = (1, 2, \dots, n)$ is a cycle of order n .

This concludes the proof of the first assertion.

Now we consider the second assertion. If n is greater than the maximal length of every unlabeled permutation in \tilde{P} , every element in $S_n(T)$ can be obtained by inflating a peg permutation of one of the two forms above. Consider

$$\tilde{\rho} = a_1 \dots a_r (r+1)^+ b_1 \dots b_s.$$

Let τ be a pattern in $a_1 \dots a_r$ and σ be a pattern in $b_1 \dots b_s$. Now inflate every element of τ and σ by the permutation 1, the element $r+1$ by an increasing permutation and every other element by the empty permutation. In this way, we get the permutation $\pi = 123[\tau, \text{id}, \sigma]$.

The permutation $\pi = 321[\sigma, \text{id}, \tau]$ can be obtained similarly from a peg permutation of the form

$$\tilde{\rho} = b_1 \dots b_s (r+1)^- a_1 \dots a_r.$$

□

Example 5.7. Consider the class $S(123, 132, 231, 3214)$. As we will see in Theorem 8.2, $\langle S_n(123, 132, 231, 3214) \rangle \cong (S_3 \times S_3) \rtimes \mathbb{Z}_2$, for every $n \geq 6$, a group of order 72.

The permutations in $S_n(123, 132, 231, 3214)$, with $n \geq 3$ are ψ ,

$$\alpha = n \ n-1 \ n-2 \ \dots \ 4 \ 2 \ 1 \ 3$$

and

$$\beta = n \ n-1 \ n-2 \ \dots \ 4 \ 3 \ 1 \ 2.$$

Hence $S(T) = \text{Grid}(\tilde{P})$, where \tilde{P} consists only of the peg permutation 4^-213 .

Now we are in position to answer Question 5.1 for every abelian G .

Theorem 5.8. *Let $T \subseteq S$ and let G be a finite, nontrivial abelian group. If there exists n_0 in \mathbb{N} such that $\langle S_n(T) \rangle \cong G$ for every $n \geq n_0$, then $G \cong \mathbb{Z}_2$ or $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.*

Proof. Suppose that $S_n(T)$ generates a fixed abelian group G for every $n \geq n_0$.

By Proposition 5.4 and by Theorem 5.6, $\cup_{n \geq 0} S_n(T) = \text{Grid}(\tilde{P})$, where \tilde{P} is a finite set of peg permutations each of which is either unlabeled, or of the form

$$\tilde{\rho} = a_1 \dots a_r (r+1)^+ b_1 \dots b_s,$$

or

$$\tilde{\rho} = b_1 \dots b_s (r+1)^- a_1 \dots a_r,$$

where, for every i and j , $a_i < r+1$ and $b_j > r+1$.

Case 1 Suppose that in $\text{Grid}(\tilde{P})$ there exists a peg permutation

$$\tilde{\rho} = a_1 \dots a_r (r+1)^+ b_1 \dots b_s.$$

- If $a_1 \dots a_r$ contains a subsequence a_x, a_y, a_z order isomorphic to 132, then, inflating a_x, a_y and a_z by the permutation 1, every other a_i and b_j with the empty permutation and c with an increasing permutation, we obtain in $S_n(T)$ the permutation $1\ 3\ 2\ 4\ 5 \dots n = (2, 3)$.
Inflating a_y and a_z by the permutation 1, every other a_i and b_j with the empty permutation and c with an increasing permutation we obtain in $S_n(T)$ the permutation $2\ 1\ 3\ 4 \dots n = (1, 2)$. But this is impossible since $(1, 2)$ and $(2, 3)$ do not commute.
- Reasoning as above, if $a_1 \dots a_r$ contains the pattern 321 we get the cycles $(1, 2)$ and $(1, 3)$ that do not commute.
- If $a_1 \dots a_r$ contains the pattern 231 we get the cycles $(1, 2)$ and $(1, 2, 3)$ that do not commute.
- If $a_1 \dots a_r$ contains the pattern 312 we get the cycles $(1, 2)$ and $(1, 3, 2)$ that do not commute.

As a consequence, $a_1 \dots a_r \in S_r(132, 231, 312, 321)$ and hence either

$$a_1 \dots a_r = \text{id}_r,$$

or

$$a_1 \dots a_r = 2\ 1\ 3\ 4 \dots r.$$

Now we turn our attention to the suffix $b_1 \dots b_s$.

- If $b_1 \dots b_s$ contains the subsequence b_x, b_y, b_z order isomorphic to 321, then, inflating b_x, b_y and b_z by the permutation 1, every other a_i and b_j with the empty permutation and c with an increasing permutation, we obtain in $S_n(T)$ the transposition $(n-2, n)$.
Inflating b_y and b_z by the permutation 1, every other a_i and b_j by the empty permutation and c by an increasing permutation we obtain in $S_n(T)$ the transposition $(n-1, n)$. But this is impossible since $(n-2, n)$ and $(n-1, n)$ do not commute.
- Similarly, if $b_1 \dots b_s$ contains the pattern 231 we get the cycles $(n-1, n)$ and $(n-2, n-1, n)$ that do not commute.
- If $b_1 \dots b_s$ contains the pattern 213 we get the cycles $(n-1, n)$ and $(n-2, n-1)$ that do not commute.
- If $b_1 \dots b_s$ contains the pattern 312 we get the cycles $(n-1, n)$ and $(n-2, n, n-1)$ that do not commute.

As a consequence the suffix $b_1 \dots b_s$ of $\tilde{\rho}$ is a permutation of the symbols $\{r+2, \dots, r+s+1\}$ that avoids 213, 231, 312, and 321 and hence it is easily seen that $b_1 \dots b_s$ is either the identity or the transposition that exchanges the two last symbols.

We deduce that the peg permutation $\tilde{\rho}$ is one of the following

$$1^+, 213^+, 1^+32, 213^+54.$$

Case 2 Suppose that in $\text{Grid}(\tilde{P})$ there exists a peg permutation

$$\tilde{\rho} = b_1 \dots b_s (r+1)^- a_1 \dots a_r.$$

Inflating every a_i and every b_j by the empty permutation and $r+1$ by a decreasing permutation, we get the permutation $\psi_n \in S_n(T)$. Notice that every other permutation α in $S_n(T)$ must commute with ψ_n because we are supposing that $S_n(T)$ generates a commutative group. But $\alpha\psi_n = \psi_n\alpha$ is equivalent to $\alpha = \alpha^{rc}$. Hence every element of $S_n(T)$ must be equal to its reverse-complement.

Now consider once more the peg permutation

$$\tilde{\rho} = b_1 \dots b_s (r+1)^- a_1 \dots a_r.$$

If $b_1 \dots b_s$ contains the pattern 12 realized by the elements b_x and b_y (with $x < y$), inflating b_x and b_y by the permutation 1, every other b_i

and a_j by the empty permutation and $r + 1$ by a decreasing permutation, we get the permutation $\alpha = n-1 \ n \ n-2 \ n-3 \ \dots \ 1 \in S_n(T)$. This is impossible since $\alpha \neq \alpha^{rc}$. As a consequence $b_1 \dots b_s$ and, similarly, $a_1 \dots a_r$ must be the decreasing permutations and hence the peg permutation $\tilde{\rho}$ is equivalent to the peg permutation 1^- .

Case 1 and **Case 2** prove that $S(T) = \text{Grid}(\tilde{P})$, where \tilde{P} contains either unlabeled permutations or elements from the set of peg permutation

$$\{1^-, 1^+, 213^+, 1^+32, 213^+54\}.$$

It is easily checked that if \tilde{P} contains 1^- then \tilde{P} does not contain 213^+ , 1^+32 , 213^+54 , since G is commutative. For every possible choice of \tilde{P} we can only get either $G \cong \mathbb{Z}_2$ or $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ for n large.

This concludes the proof. \square

Now we consider Question 5.1 when G is an alternating group.

Theorem 5.9. *Let $T \subseteq S$ and let $G = A_m$, $m \geq 1$, be an alternating group. If there exists n_0 in \mathbb{N} such that $\langle S_n(T) \rangle \cong G$ for every $n \geq n_0$, then $m \leq 2$.*

Proof. Suppose that $S_n(T)$ generates a fixed alternating group $G = A_m$ for every $n \geq n_0$.

Proposition 5.4 and Theorem 5.6 imply that $\cup_{n \geq 0} S_n(T) = \text{Grid}(\tilde{P})$ where \tilde{P} is a finite set of peg permutations each of which is either unlabeled, or of the form

$$\tilde{\rho} = a_1 \dots a_r (r + 1)^+ b_1 \dots b_s,$$

or

$$\tilde{\rho} = b_1 \dots b_s (r + 1)^- a_1 \dots a_r,$$

where, for every i and j , $a_i < r + 1$ and $b_j > r + 1$.

If one peg permutation of the second kind appears in \tilde{P} , inflating each a_i and b_j by the empty permutation and $r + 1$ by a decreasing permutation we get $\psi_n \in S_n(T)$ for every n . But in this case it is impossible that $S_n(T)$ generates a fixed alternating group A_m for sufficiently large n , because ψ_n is odd if n is odd. Hence every labeled peg permutation in \tilde{P} is of the first kind.

Let

$$\tilde{\rho} = a_1 \dots a_r (r + 1)^+ b_1 \dots b_s$$

be one of the labeled peg permutation in \tilde{P} .

Suppose that $a_1 \dots a_r$ contains the pattern 21, realized by the elements a_x and a_y (with $x < y$). Then, inflating a_x and a_y by the permutation 1,

every other a_i and b_j by the empty permutation and $r+1$ by the identity, we get the permutation $2\ 1\ 3\ \dots\ n = (1, 2) \in S_n(T)$. This is impossible, since this permutation is odd. Hence $a_1 \dots a_r = \text{id}_r$.

We can show similarly that $b_1 \dots b_s$ must be the increasing permutation id_s .

As a consequence $\tilde{\rho}$ is equivalent to 1^+ , the set $S_n \cap \text{Grid}(\tilde{P})$ contains only id_n for sufficiently large n and $\langle S_n(T) \rangle = \{\text{id}_n\} = A_1 = A_2$. \square

6 Sets T such that $\langle S_n(T) \rangle = S_n$.

From now on we determine the groups $\langle S_n(T) \rangle$ for particular sets of patterns T .

First of all we state a rather general result that allows us to determine sets $T \subseteq S$ such that $\langle S_n(T) \rangle = S_n$.

Theorem 6.1. *Let $T \subseteq S$. If either*

$$T \cap \{k\ 2\ 3\ \dots\ k-1\ 1, \quad k\ 1\ 2\ \dots\ k-1, \quad 2\ 3\ \dots\ k\ 1, \quad \text{id}_k \mid k \geq 1\} = \emptyset,$$

or

$$T \cap \{\text{id}_k, \quad (r-1, \quad r) \mid 2 \leq r \leq k \text{ and } k \geq 1\} = \emptyset,$$

or

$$T \cap \{2\ 3\ \dots\ k\ 1, \quad \text{id}_k, \quad 2\ 1\ 3\ \dots\ k \mid k \geq 1\} = \emptyset,$$

or

$$T \cap \{1\ \dots\ k-2\ k\ k-1, \quad \text{id}_k, \quad 2\ 3\ \dots\ k\ 1 \mid k \geq 1\} = \emptyset,$$

then $\langle S_n(T) \rangle = S_n$ for every n .

In particular,

- i. if $T \subseteq S_3$ and $\langle S_n(T) \rangle \neq S_n$, then T shares at least an element with each one of the following sets*

$$\{123, 231, 312, 321\}, \{123, 132, 213\}, \{123, 213, 231\}, \text{ and} \\ \{123, 132, 231\}.$$

- ii. if $T \subseteq S_4$ and $\langle S_n(T) \rangle \neq S_n$, then T shares at least an element with each one of the following sets*

$$\{1234, 2341, 4123, 4231\}, \{1234, 1243, 1324, 2134\}, \\ \{1234, 2134, 2341\} \text{ and } \{1234, 1243, 2341\}.$$

Proof. If $n < k$ the theorem is trivially true. Assume that $n > k$. Note that

- the patterns of length k contained in at least one elementary transposition in S_n are id_k and $(r-1, r)$ for $2 \leq r \leq k$,
- the patterns of length k contained either in the cycle $(1, n) \in S_n$ or in $(1, 2, \dots, n) \in S_n$ are $k-2 \ 3 \dots k-1 \ 1$, $k-1 \ 2 \dots k-1$, $1 \ 2 \ 3 \dots k-1$, id_k ,
- the patterns of length k contained either in the cycle $(1, 2) \in S_n$ or in $(1, 2, \dots, n) \in S_n$ are $2 \ 3 \dots k-1$, id_k , $2 \ 1 \ 3 \dots k$, and
- the patterns of length k contained either in the cycle $(n-1, n) \in S_n$ or in $(1, 2, \dots, n) \in S_n$ are $2 \ 3 \dots k-1 \ 1$, id_k , $1 \ 2 \dots k-2 \ k \ k-1$.

The assertion follows by Lemma 3.5.

If $k = n$ the proof is similar. □

7 Groups generated by $S_n(T)$ with $T \subseteq S$ and $|T| \leq 3$.

In this section we consider the group generated by the set $S_n(T)$, where $T \subseteq S$ and $|T| \leq 3$.

We begin with the analysis of $\langle S_n(T) \rangle$ when T consists of three patterns of length 3.

Theorem 7.1. 1. $\langle S_n(123, 321, \tau) \rangle = \{\text{id}_n\}$, for every $n > 4$ and for every $\tau \in S_3$.

2. $\langle S_n(132, 213, 321) \rangle \cong \mathbb{Z}_n$,

3. $\langle S_n(123, 231, 312) \rangle \cong D_n$, for every $n \geq 3$.

4. $\langle S_n(T) \rangle = S_n$ for every other set T of three patterns of length three.

Proof. 1. The set of generators is trivially empty for $n > 4$.

2. The set of generators (which turns out to be itself a group by Lemma 3.7) consists of the cycle $(1, 2, \dots, n)$ and its powers.

3. The set $S_n(123, 231, 312)$ contains only n elements whose one-line notation is:

$$k \ k-1 \ \dots \ 2 \ 1 \ n \ n-1 \ \dots \ k+1, \quad 1 \leq k \leq n.$$

This means that ψ and

$$\alpha = 1 \ n \ n-1 \ \dots \ 3 \ 2$$

belong to this set. Observe that

$$\alpha^2 = \psi^2 = (\alpha\psi)^n = \text{id},$$

hence

$$\langle \alpha, \psi \rangle \cong D_n.$$

It is easily seen that the other elements in $S_n(123, 231, 312)$ are of the form $(\alpha\psi)^j\alpha$ for some j .

4. By Lemma 3.4 we have

$$\langle S_n(123, 132, 213) \rangle = \langle S_n(123, 132, 213) \cup S_n(231, 312, 321) \rangle,$$

and $\langle S_n(231, 312, 321) \rangle = S_n$ by Theorem 6.1[i.].

By Lemma 3.3 we have

$$\begin{aligned} \langle S_n(123, 213, 231) \rangle &\cong \langle S_n(123, 213, 312) \rangle \cong \langle S_n(123, 132, 312) \rangle \cong \\ &\langle S_n(123, 132, 231) \rangle, \end{aligned}$$

and

$$\begin{aligned} \langle S_n(132, 312, 321) \rangle &\cong \langle S_n(132, 231, 321) \rangle \cong \langle S_n(213, 231, 321) \rangle \cong \\ &\langle S_n(213, 312, 321) \rangle, \end{aligned}$$

by Lemma 3.4 we have

$$\begin{aligned} \langle S_n(123, 213, 231) \rangle &= \\ \langle S_n(123, 213, 231) \cup S_n(132, 312, 321) \cup S_n(213, 231, 321) \cup S_n(123, 132, 312) \rangle, \end{aligned}$$

and $\langle S_n(132, 312, 321) \rangle = S_n$ by Theorem 6.1[i.].

By Lemma 3.3 we have

$$\langle S_n(213, 231, 312) \rangle \cong \langle S_n(132, 231, 312) \rangle,$$

and

$$\langle S_n(132, 213, 231) \rangle \cong \langle S_n(132, 213, 312) \rangle,$$

by Lemma 3.4 we have

$$\langle S_n(213, 231, 312) \rangle = \langle S_n(132, 213, 231) \rangle =$$

$$\langle S_n(213, 231, 312) \cup S_n(312, 132, 213) \cup S_n(132, 213, 231) \cup S_n(132, 231, 312) \rangle.$$

This last generating set contains the transposition $(1, 2)$ (an element of $S_n(132, 231, 312)$) and the cycle $(1, 2, \dots, n)$ (an element of $S_n(132, 213, 312)$). Hence the generated group is S_n by Lemma 3.5. □

As a consequence we can characterize all the groups $\langle S_n(T) \rangle$ where T contains no more than three arbitrary patterns.

Corollary 7.2. *Let $T \subseteq S$ be a set of patterns, with $|T| \leq 3$.*

1. *If $\{\text{id}_r, \psi_s\} \subseteq T$, for some $r, s \geq 1$, then $\langle S_n(T) \rangle = \{\text{id}_n\}$ for $n > (r-1)(s-1)$.*
2. *If for every $r, s \geq 1$ $\{\text{id}_r, \psi_s\} \not\subseteq T$, $T \neq \{132, 213, 321\}$ and $T \neq \{123, 231, 312\}$, then $\langle S_n(T) \rangle = S_n$.*

Proof. The first assertion is trivial since the set $S_n(T)$ is empty for n sufficiently large when $\{\text{id}_r, \psi_s\} \subseteq T$. More precisely, every permutation of length $n > (r-1)(s-1)$ must contain a monotonically increasing subsequence of length r or a monotonically decreasing subsequence of length s , by the Erdős-Szekeres Theorem [8].

To prove the second assertion, recall that the only sets T of three patterns of length three such that $\langle S_n(T) \rangle \neq S_n$ are $\{132, 213, 321\}$ and $\{123, 231, 312\}$, by Theorem 7.1. Hence, the second assertion follows from Theorem 7.1 and from Lemmas 3.1 and 3.2 in all cases, except for the following ones.

- **Case 1** T consists of three (distinct) patterns τ_1, τ_2 and τ_3 with $\tau_1 \geq 132$, $\tau_2 \geq 213$ and $\tau_3 \geq 321$, with at least one of these patterns of length strictly greater than 3.
- **Case 2** T consists of three (distinct) patterns τ_1, τ_2 and τ_3 with $\tau_1 \geq 123$, $\tau_2 \geq 231$ and $\tau_3 \geq 312$, with at least one of these patterns of length strictly greater than 3.

If Case 1 occurs, we can assume that $\tau_3 \neq 321$, otherwise Lemma 3.2 and Theorem 7.1 would imply that $\langle S_n(T) \rangle = S_n$.

Notice that $\langle S_n(132, 213, 4321) \rangle = S_n$ because the set of generators contains the cycles $(1, 2, \dots, n)$ and $(1, n)$, which generate S_n by Lemma 3.5.

Now the fact that $S_n(132, 213, \tau_3)$ with $\tau_3 > 4321$ generates S_n follows by Lemma 3.2.

If Case 2 occurs, we can assume that $\tau_1 \neq 123$, otherwise Lemma 3.2 and Theorem 7.1 would imply that $\langle S_n(T) \rangle = S_n$.

Notice that

$$\begin{aligned} \langle S_n(231, 312, 1234) \rangle = \\ \langle S_n(231, 312, 1234) \cup S_n(132, 213, 4321) \cup S_n(231, 312, 1234) \rangle \end{aligned}$$

by Lemma 3.4. This last group is S_n because $S_n(132, 213, 4321)$ contains the permutations $(1, n)$ and $(1, 2, \dots, n)$, which generate S_n by Lemma 3.5.

Now the fact that $S_n(\tau_1, 231, 312)$ with $\tau_1 > 1234$ generates S_n follows by Lemma 3.2. □

8 Groups generated by $S_n(T)$ with $T \subseteq S_3 \cup S_4$ and $|T| \geq 4$.

In this section we begin the study of the groups $\langle S_n(T) \rangle$ where T has cardinality greater than three. We select some of the most meaningful cases.

We consider the group generated by $S_n(T)$ with $T \subseteq S_3$. The only cases not already covered by Theorem 7.1 and Corollary 7.2 are those with $|T| \geq 4$. The following theorem deals with all these cases.

Theorem 8.1. 1. $\langle S_n(\{123, 321\} \cup T') \rangle = \{\text{id}\}$, for every $n > 4$, where T' is an arbitrary subset of S .

2. $\langle S_n(213, 231, 312, 321) \rangle \cong \langle S_n(132, 231, 312, 321) \rangle \cong \langle S_n(132, 213, 231, 312) \rangle \cong \mathbb{Z}_2$,
3. $\langle S_n(123, 132, 213, 231) \rangle \cong \langle S_n(123, 132, 213, 312) \rangle \cong D_4$,
4. $\langle S_n(132, 213, 312, 321) \rangle \cong \langle S_n(132, 213, 231, 321) \rangle \cong \mathbb{Z}_n$,
5. $\langle S_n(123, 213, 231, 312) \rangle \cong \langle S_n(123, 132, 231, 312) \rangle \cong D_n$,
6. $\langle S_n(123, 132, 213, 231, 312) \rangle \cong \mathbb{Z}_2$,
7. $\langle S_n(132, 213, 231, 312, 321) \rangle \cong \{\text{id}_n\}$.

Proof. 1. The set of generators is trivially empty for $n > 4$.

2. In each case, we have that the set of generators consists only of the identity and an involution.
3. The set $S_n(123, 132, 213, 231)$ contains only ψ and $(1, 2)\psi$ (whose period is 4), hence $\langle S_n(123, 132, 213, 231) \rangle = D_4$.

4. The set $S_n(132, 213, 312, 321)$ contains only the identity and the cycle $(1, 2, 3, \dots, n)$.
5. The set $S_n(123, 213, 231, 312)$ contains only ψ and $\alpha = 1 \ n \ n - 1 \ \dots \ 3 \ 2$. As seen in the proof of Theorem 7.1, these two permutations generate the dihedral D_n .
6. The only element in the generating set is ψ_n .
7. The only element in the generating set is id_n .

□

Now we consider the case of three patterns τ_1, τ_2, τ_3 in S_3 and one pattern τ_4 in S_4 . We suppose also that τ_4 avoids each of τ_1, τ_2 and τ_3 , otherwise we fall back in the previous cases.

- Theorem 8.2.**
1. $\langle S_n(123, 132, 213, 4231) \rangle \cong \langle S_n(132, 213, 231, 4123) \rangle \cong \langle S_n(132, 231, 312, 3214) \rangle \cong D_4$ for every $n \geq 4$.
 2. $\langle S_n(123, 132, 231, 3214) \rangle \cong \langle S_n(132, 213, 231, 1234) \rangle \cong (S_3 \times S_3) \rtimes \mathbb{Z}_2$ for every $n \geq 6$.
 3. $\langle S_n(123, 231, 312, 1432) \rangle \cong \langle S_n(123, 231, 312, 2143) \rangle \cong \langle S_n(132, 213, 231, 4312) \rangle \cong \langle S_n(132, 231, 312, 2134) \rangle \cong D_n$ for every $n \geq 2$.
 4. $\langle S_n(132, 213, 321, 2341) \rangle \cong \langle S_n(132, 213, 321, 3412) \rangle \cong \mathbb{Z}_n$ for every $n \geq 1$.
 5. $\langle S_n(132, 231, 312, 4321) \rangle \cong \langle S_n(132, 231, 321, 4123) \rangle \cong S_3$ for every $n \geq 1$.
 6. $\langle S_n(231, 312, 321, 1243) \rangle \cong \langle S_n(231, 312, 321, 2134) \rangle \cong S_4$ for every $n \geq 1$.
 7. $\langle S_n(231, 312, 321, 1324) \rangle = \mathbb{Z}_2 \times \mathbb{Z}_2 \cong D_2$ for every $n \geq 4$.
 8. $\langle S_n(123, 132, 213, 4312) \rangle \cong (((((\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_3) \rtimes \mathbb{Z}_2) \rtimes \mathbb{Z}_3) \rtimes \mathbb{Z}_2) \rtimes \mathbb{Z}_2$ for every $n \geq 8$.

9. Let $\tau_1, \tau_2, \tau_3 \in S_3$ and $\tau_4 \in S_4$, and suppose that the set $\{\tau_1, \tau_2, \tau_3, \tau_4\}$ is not obtained from one of the set in the preceding cases either by the application of the reverse-complement map, or of the inverse map, or their composition. Suppose also that τ_4 avoids τ_1, τ_2 and τ_3 . Then, for sufficiently large n , $\langle S_n(\tau_1, \tau_2, \tau_3, \tau_4) \rangle$ is either S_n , or the trivial group $\{\text{id}_n\}$.

Proof. 1. We prove only that $\langle S_n(123, 132, 213, 4231) \rangle \cong D_4$, the other cases are analogous. Since a permutation in the generating set must avoid 123, 132, 213 and 4231, the only elements of this set are ψ, α, β and γ , where

$$\alpha = n-1 \quad n \quad n-2 \quad n-3 \dots 4 \ 3 \ 1 \ 2 = (1, n-1)(2, n) \prod_{j=3}^{\lfloor \frac{n}{2} \rfloor + 1} (j, n-j+1),$$

$$\beta = n \quad n-1 \quad n-2 \quad n-3 \dots 4 \ 3 \ 1 \ 2 = (1, n, 2, n-1) \prod_{j=3}^{\lfloor \frac{n}{2} \rfloor + 1} (j, n-j+1),$$

and

$$\gamma = n-1 \quad n \quad n-2 \quad n-3 \dots 4 \ 3 \ 2 \ 1 = (1, n-1, 2, n) \prod_{j=3}^{\lfloor \frac{n}{2} \rfloor + 1} (j, n-j+1).$$

Note that $\gamma = \beta^{-1}$, $\psi = \beta\alpha\beta$, the order of β is four, the order of α and $\alpha\beta = (1, 2)$ is two, and the group generated by ψ, α, β and γ is D_4 .

2. We prove only that $\langle S_n(123, 132, 231, 3214) \rangle \cong (S_3 \times S_3) \rtimes \mathbb{Z}_2$.

Since a permutation in the generating set must avoid 123, 132, 231 and 3214, the only elements of this set are ψ, α and β , where

$$\alpha = n \quad n-1 \quad n-2 \dots 4 \ 2 \ 1 \ 3 = (1, n, 3, n-2, 2, n-1) \prod_{j=4}^{\lfloor \frac{n}{2} \rfloor + 1} (j, n-j+1),$$

and

$$\beta = n \quad n-1 \quad n-2 \dots 4 \ 3 \ 1 \ 2 = (1, n, 2, n-1)(3, n-2) \prod_{j=4}^{\lfloor \frac{n}{2} \rfloor + 1} (j, n-j+1).$$

Notice that ψ can be eliminated from the set of generators, since

$$\psi = (\beta^{-1}\alpha^{-1})^3\beta^{-1}.$$

Now, the subgroup $\langle \alpha, \beta \rangle$ of S_n is clearly isomorphic to the subgroup $G = \langle \widehat{\alpha}, \widehat{\beta} \rangle$ of S_6 where $\widehat{\alpha} = (1, 6, 3, 4, 2, 5)$ and $\widehat{\beta} = (1, 6, 2, 5)(3, 4)$. Consider now the subgroups N and H of S_6 defined by

$$N = \langle (4, 6, 5), (2, 3)(4, 5), (1, 3, 2)(4, 5, 6), (1, 6, 3, 4, 2, 5) \rangle$$

and

$$H = \langle (5, 6) \rangle.$$

Long but trivial calculations show that N and H are subgroups of G , with $N \trianglelefteq G$, every $g \in G$ can be written as $g = kh$, where $k \in N$ and $h \in H$, and $N \cap H = \{id\}$. By Lemma 3.6, we have $G \cong N \rtimes H$.

Moreover, it is easy to show that $N \cong S_3 \times S_3$. In fact, if we identify $S_3 \times S_3$ with the subgroup of S_6 generated by the cycles $(1, 2, 3)$, $(1, 2)$, $(4, 5, 6)$ and $(4, 5)$, the following map f yields an isomorphism between N and $S_3 \times S_3$.

$$f((4, 6, 5)) = (1, 2, 3)(4, 6, 5), \quad f((2, 3)(4, 5)) = (1, 2)(4, 6),$$

$$f((1, 3, 2)(4, 5, 6)) = (1, 2, 3), \quad \text{and} \quad f((1, 6, 3, 4, 2, 5)) = (1, 3, 2)(4, 5).$$

3. We prove only that $\langle S_n(123, 231, 312, 1432) \rangle \cong D_n$.

The generating set is easily seen to contain only the permutations ψ , α and β , where

$$\alpha = n - 1 \quad n - 2 \quad \dots \quad 2 \quad 1 \quad n = \prod_{j=1}^{\lfloor \frac{n}{2} \rfloor + 1} (j, n - j)$$

and

$$\beta = n - 2 \quad n - 3 \quad \dots \quad 2 \quad 1 \quad n \quad n - 1 = (n - 1, n) \prod_{j=1}^{\lceil \frac{n}{2} \rceil} (j, n - j - 1).$$

Notice that β is a superfluous generator since $\beta = \alpha\psi\alpha$, and that $\psi\alpha = (1, 2, 3, \dots, n)$ has order n . The assertion follows.

4. We consider only $\langle S_n(132, 213, 321, 2341) \rangle$.

The only elements of the generating set different from the identity permutation are

$$\alpha = n \quad 1 \quad 2 \quad \dots \quad n - 1 = (1, n, n - 1, \dots, 2)$$

and

$$\beta = n - 1 \quad n \ 1 \ \dots \ n - 2.$$

Notice that β is composed by two cycles of length $\frac{n}{2}$ if n is even and by one cycle of length n if n is odd. In both cases $\beta = \alpha^2$, hence the generated group is \mathbb{Z}_n .

5. This is a particular case of Theorem 4.2.
6. The only elements in the generating set are the identity, $(1, 2)$, $(2, 3)$ and $(1, 2)(3, 4)$ in the first case, and the identity, $(n-1, n)$, $(n-2, n-1)$ and $(n-3, n-2)(n-1, n)$ in the second case.
7. Given a permutation in the generating set, the symbols $n-1$ and n must occupy the last and the second last position, otherwise the permutation contains 312 or 321. Similarly, the symbols 1 and 2 must occupy the first or the second position. The remaining symbols must be in increasing order since the permutation must avoid 1324.

Hence the set $S_n(231, 321, 312, 1324)$ contains only the identity, the elementary transpositions $(1, 2)$ and $(n-1, n)$, and the involution $(1, 2)(n-1, n)$, which generate $\mathbb{Z}_2 \times \mathbb{Z}_2$.

8. Consider $\langle S_n(123, 132, 213, 4312) \rangle$, with $n \geq 4$. The generating set contains the permutations ψ_n , α , β and γ , where

$$\alpha = n - 1 \quad n \quad n - 2 \quad n - 3 \quad \dots \ 1,$$

$$\beta = n - 1 \quad n \quad n - 3 \quad n - 2 \quad n - 4 \quad n - 5 \quad \dots \ 1,$$

and

$$\gamma = n \quad n - 2 \quad n - 1 \quad n - 3 \quad n - 4 \quad n - 5 \quad \dots \ 1.$$

One can easily realize that the groups $\langle S_n(123, 132, 213, 4312) \rangle$ for $n \geq 8$ are all isomorphic. Using GAP [9] we verified that $\langle S_8(123, 132, 213, 4312) \rangle$ is isomorphic to the group of order 1152 appearing in the statement.

9. Let $T \in S$ be a set consisting of three (distinct) patterns of length three, τ_1, τ_2 and τ_3 , and one pattern τ_4 of length four. Assume that τ_4 does not contain any of the other elements of T . Suppose that $\langle S_n(T) \rangle \neq S_n$. Recall that by Theorem 6.1 for $\langle S_n(T) \rangle$ to be different from S_n , T must intersect each of the following sets

$$A = \{123, 231, 312, 321, 1234, 2341, 4123, 4231\},$$

$$B = \{123, 132, 213, 1234, 1243, 1324, 2134\},$$

$$C = \{123, 213, 231, 1234, 2134, 2341\}$$

and

$$D = \{123, 132, 231, 1234, 1243, 2341\}.$$

Case 1:

If T does not contain neither ψ_3 nor ψ_4 , by Lemma 3.4 and Lemma 3.3, we have $\langle S_n(T) \rangle = \langle S_n(T) \cup S_n(T^r) \cup S_n(T^c) \cup S_n(T^{rc}) \cup S_n(T^{-1}) \cup S_n((T^r)^{-1}) \cup S_n((T^c)^{-1}) \cup S_n((T^{rc})^{-1}) \rangle$.

Hence, each of the sets T^* , where $*$ denotes any of $1, r, c, rc, -1$, or of their compositions, must intersect each of the sets A, B, C and D or, equivalently, T must intersect each of the sets A^*, B^*, C^* and D^* .

A long but trivial computation that we realized with SageMath [20] shows that the sets T with these characteristics are 26. Previous considerations and results allow us to reduce ourselves to the following three sets

$$\{132, 231, 312, 1234\}, \{123, 132, 231, 4312\}, \{123, 213, 231, 4312\}.$$

- Let $G = \langle S_n(132, 231, 312, 1234) \rangle$. Notice that the generating set contains only

$$\alpha = n - 2 \quad n - 3 \quad \dots \quad 1 \quad n - 1 \quad n,$$

$$\beta = n - 1 \quad n - 2 \quad \dots \quad 1 \quad n,$$

and ψ_n .

Hence in G we have the elements

$$(\beta^r)^{-1} = 2 \ 3 \ \dots \ n \ 1 = (1, 2, \dots, n),$$

and

$$\psi \beta^{-1} \psi \beta \alpha \psi = (1, 2),$$

which implies $G = S_n$ by Lemma 3.5.

- Let $G = \langle S_n(123, 213, 231, 4312) \rangle$. Notice that the generating set contains only

$$\alpha = n \quad n - 2 \quad n - 3 \quad \dots \quad 1 \quad n - 1,$$

$$\beta = n - 1 \ n - 2 \ n - 3 \ \dots 1 \ n,$$

and ψ_n . The generated group thus contains

$$(\alpha\beta)^{rc} = (1, 2)$$

and $\beta^c = (1, 2, \dots, n)$, which generate S_n by Lemma 3.5.

- Let $G = \langle S_n(123, 132, 231, 4312) \rangle$. Notice that the generating set contains only

$$\alpha = n \ 1 \ n - 2 \ n - 3 \ \dots 2,$$

$$\beta = 1 \ n \ n - 2 \ n - 3 \ \dots 2$$

and ψ_n . The generated group thus contains

$$(\beta\alpha)^{rc} = (1, 2)$$

and $\beta^r = (1, 2, \dots, n)$ which generate S_n by Lemma 3.5.

Case 2:

Now we consider the cases when T contains ψ_3 or ψ_4 .

If $\sigma_4 = \psi_4$, ψ_3 is not in T since σ_4 does not contain any of σ_1 , σ_2 and σ_3 . As recalled above, if $\langle S_n(T) \rangle \neq S_n$, T must intersect each of the sets A, B, C and D . We can also assume that T does not contain id_3 , otherwise the generated group would be trivial for large n .

The only sets T with all these properties are

$$\{132, 213, 312, 4321\}, \{132, 213, 231, 4321\},$$

$$\{132, 231, 312, 4321\}, \{213, 231, 312, 4321\}.$$

The third one of these sets has been already considered in the previous items of the Theorem and the last one is the reverse-complement of the third one. The second case is the inverse of the first one. So we can consider only the first case.

Let $G = \langle S_n(132, 213, 312, 4321) \rangle$. Notice that the generating set contains only

$$\alpha = (1, 2, \dots, n),$$

$$\beta = 3 \ 4 \ \dots \ n \ 2 \ 1,$$

and id_n .

The generated group thus contains

$$\alpha^2\beta^{-1} = (1, 2),$$

and coincides with S_n by Lemma 3.5.

Finally, we consider the case when $\psi_3 \in T$. In this case we can suppose that

- $\sigma_4 \neq \psi_4, \text{id}_4$,
- T does not contain id_3 ,
- σ_4 does not contain any other σ_i 's,
- T intersects the sets A, B, C , and D defined above,
- T intersects the sets A^{-1}, B^{-1}, C^{-1} and D^{-1} (since $\langle S_n(T) \rangle = \langle S_n(T^{-1}) \rangle$).

If these hypothesis are not satisfied the generated group is either $\{\text{id}_n\}$ or S_n .

There are 14 of such sets T , but 13 of them have been already considered.

The remaining case is

$$\{213, 312, 321, 1243\}.$$

Hence let $G = \langle S_n(213, 312, 321, 1243) \rangle$. Notice that the generating set contains only

$$\begin{aligned} \alpha &= 2\ 3\ 4\ \dots\ n\ 1, \\ \beta &= 1\ 3\ 4\ \dots\ n\ 2 \end{aligned}$$

and id_n . The generated group thus contains

$$\beta\alpha^{-1} = (1, 2)$$

thus is S_n by Lemma 3.5.

This concludes the proof. □

Now we consider the groups generated by sets $S_n(T)$ where $T \subseteq S_4$ and $|T| = 4$.

Theorem 8.3. *Let $T \subseteq S_4$ with $|T| = 4$. Suppose that T does not contain ψ_4 and that $T \neq \{1234, 1432, 3214, 4312\}$ and $T \neq \{1234, 1432, 3214, 4231\}$. Then $\langle S_n(T) \rangle = S_n$.*

Proof. Let $T \subseteq S_4 \setminus \{4321\}$ with $|T| = 4$. Suppose that $S_n(T)$ generates a group different from S_n .

Recall from Theorem 6.1.ii that for $\langle S_n(T) \rangle$ to be different from S_n , T must intersect each of the following sets

$$A = \{1234, 2341, 4123, 4231\},$$

$$B = \{1234, 1243, 1324, 2134\},$$

$$C = \{1234, 2134, 2341\}$$

and

$$D = \{1234, 1243, 2341\}.$$

Since $4321 \notin T$, by Lemma 3.4 and Lemma 3.3, $\langle S_n(T) \rangle = \langle S_n(T) \cup S_n(T^r) \cup S_n(T^c) \cup S_n(T^{rc}) \cup S_n(T^{-1}) \cup S_n((T^r)^{-1}) \cup S_n((T^c)^{-1}) \cup S_n((T^{rc})^{-1}) \rangle$.

We are assuming that $\langle S_n(T) \rangle$ is not S_n . Hence each one of the sets T^* , where $*$ denotes any of $1, r, c, rc, -1$ or of their compositions, must intersect each of the sets A, B, C and D .

A long but trivial computation that we carried out with the aid of Sage-Math shows that the only sets $T \subseteq S_4 \setminus \{4321\}$ with $|T| = 4$ that intersect each of the sets A^*, B^*, C^* and D^* are

$$\{1234, 3214, 3421, 4312\}, \{1234, 1324, 3421, 4312\},$$

$$\{1234, 1432, 3421, 4312\}, \{1234, 1432, 3214, 4312\},$$

$$\{1234, 1432, 3214, 3421\}, \{1234, 1432, 3214, 4231\}.$$

Notice that

$$S_n(123, 213, 312, 3421) \subseteq S_n(1234, 3214, 3421, 4312)$$

and the set on the left generates S_n by Theorem 8.2. Hence

$$\langle S_n(1234, 3214, 3421, 4312) \rangle = S_n.$$

Similarly,

$$S_n(123, 132, 312, 3421) \subseteq S_n(1234, 1324, 3421, 4312),$$

and

$$S_n(123, 132, 312, 3421) \subseteq S_n(1234, 1432, 3421, 4312),$$

and

$$S_n(123, 132, 213, 3421) \subseteq S_n(1234, 1432, 3214, 3421),$$

hence $S_n(1234, 1324, 3421, 4312)$, $S_n(1234, 1432, 3421, 4312)$ and $S_n(1234, 1432, 3214, 3421)$ generate S_n .

Using SageMath it is possible to verify that $S_n(1234, 1432, 3214, 4312)$ does not generate S_n for sufficiently large n ($n > 20$).

In fact, for $n \geq 9$, the last $n-9$ elements of any permutation in $S_n(1234, 1432, 3214, 4312)$ are the smallest ones in decreasing order and the set $S_n(1234, 1432, 3214, 4312)$ has always cardinality 145. To explain this fact, notice that given $\pi \in S_n(1234, 1432, 3214, 4312)$, with $n \geq 9$,

- the position of the element 1 is greater than or equal to $n-4$, otherwise π would contain 1234 or 1432.
- Similarly the element 2 has position greater than or equal to $n-5$.
- The element 2 precedes the element 1. In fact, if the element 2 follows the element 1, all the elements before 1 must be in increasing order because π avoids 4312. But this is impossible since π avoids also 1234.
- For the same reason 3 precedes 2. Hence we have that the elements 321 form a decreasing sequence.
- The element 1 is in last position, otherwise we would have an occurrence of 3214.

Iterating this process we conclude that the last $n-9$ elements of π are the smallest ones in decreasing order. Notice that the same conclusion does not hold for the first nine elements of the permutation. As an example consider the following permutation in $S_{21}(1234, 1432, 3214, 4312)$,

$$19\ 20\ 21\ 14\ 13\ 16\ 15\ 18\ 17\ 12\ 11\ 10\ 9\ 8\ 7\ 6\ 5\ 4\ 3\ 2\ 1.$$

As a consequence, if $n \geq 9$, there is a bijection between $S_n(1234, 1432, 3214, 4312)$ and $S_{n+1}(1234, 1432, 3214, 4312)$.

For $n = 2k + 1$, $k \geq 10$, the element $k + 1$ turns out to be fixed in any permutation in this set and hence also in the generated group. Similarly for $n = 2k$, $k \geq 10$, the elements k and $k + 1$ are in position $k + 1$ and k respectively. Hence every element in the generated group fixes the set $\{k, k + 1\}$.

It can be shown similarly that neither $S_n(1234, 1432, 3214, 4231)$ generates S_n for sufficiently large n . In fact, it possible to prove that, given $\pi \in S_n(1234, 1432, 3214, 4231)$ then, for $n \geq 13$, $\pi_7\pi_8 \dots \pi_{n-6} = n-6 \ n-7 \dots 7$. □

9 Open problems

We conclude the paper with some open problems.

Firstly, suppose that the sequence $(S_n(T))_{n \geq 0}$ is eventually constant. What kind of groups arise in this way? For example, is it possible to obtain the dihedral group D_k , for $k > 4$?

Another problem is that of completing the classification of $\langle S_n(T) \rangle$ when T consists of four or more patterns. In particular, we wonder whether there exists a general procedure to determine the group $\langle S_n(T) \rangle$ for any finite set of patterns T .

References

- [1] Michael H. Albert, Robert E. L. Aldred, Mike D. Atkinson, Hans van Ditmarsch, Chris C. Handley, Derek A. Holton, and D. J. McCaughan. Compositions of pattern restricted sets of permutations. *Australas. J Comb.*, 37:43–56, 2007.
- [2] Michael H. Albert, Mike D. Atkinson, Mathilde Bouvel, Nik Ruvskuc, and Vincent Vatter. Geometric grid classes of permutations. *Transactions of the American Mathematical Society*, 365:5859–5881, 2013.
- [3] Michael H. Albert, Mike D. Atkinson, and Robert Brignall. Permutation classes of polynomial growth. *Annals of Combinatorics*, 11:249–264, 2006.
- [4] Mike D. Atkinson and Robert Beals. Permutation involvement and groups. *Quarterly Journal of Mathematics*, 52:415–421, 2001.
- [5] M. Bóna. *Combinatorics of Permutations*. Discrete mathematics and its applications. CRC Press, 3rd edition, 2022.
- [6] Amanda Burcroff and Colin Defant. Pattern-avoiding permutation powers. *Discret. Math.*, 343:112017, 2020.
- [7] P.J. Cameron. *Permutation Groups*. London Mathematical Society Student Texts. Cambridge University Press, 1999.
- [8] P. Erdős and G. Szekeres. A combinatorial problem in geometry. *Compositio Mathematica*, 2:463–470, 1935.
- [9] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.12.2*, 2022.

- [10] Cheyne Homberger and Vincent Vatter. On the effective and automatic enumeration of polynomial permutation classes. *J. Symb. Comput.*, 76:84–96, 2016.
- [11] Sophie Huczynska and Vincent Vatter. Grid classes and the Fibonacci dichotomy for restricted permutations. *Electron. J. Comb.*, 13, 2006.
- [12] I.M. Isaacs. *Finite Group Theory*. Graduate studies in mathematics. American Mathematical Society, 2008.
- [13] Tomás Kaiser and Martin Klazar. On growth rates of closed permutation classes. *Electron. J. Comb.*, 9, 2003.
- [14] Mark Karpilovskij. Composability of permutation classes. *Electron. J. Comb.*, 26(1):#1.41, 2019.
- [15] S. Kitaev. *Patterns in Permutations and Words*. Monographs in Theoretical Computer Science. Springer, 2011.
- [16] Erkkö Lehtonen. Permutation groups arising from pattern involvement. *Journal of Algebraic Combinatorics*, 52:251 – 298, 2020.
- [17] Erkkö Lehtonen, Reinhard Pöschel, and Gábor Czédli. Permutation groups, pattern involvement, and galois connections. *Acta Scientiarum Mathematicarum*, 83:355–375, 2017.
- [18] Jay Pantone and Vincent Vatter. Growth rates of permutation classes: Categorization up to the uncountability threshold. *Israel Journal of Mathematics*, 236:1 – 43, 2020.
- [19] B.E. Sagan. *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*, volume 203 of *Graduate Texts in Mathematics*. Springer New York, 2nd edition, 2001.
- [20] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 9.8)*, 2023. <https://www.sagemath.org>.
- [21] Vincent Vatter. Permutation classes of every growth rate above 2.48188. *Mathematika*, 56:182–192, 2010.
- [22] Vincent Vatter. Small permutation classes. *Proceedings of the London Mathematical Society*, 103:879 – 921, 2011.