EXAMPLES OF RELATIVE QUANTUM COHOMOLOGY

KAI HUGTENBURG AND SARA B. TUKACHINSKY

Abstract. We compute the quantum cohomology relative to a Lagrangian submanifold in some complete intersections. For quadric hypersurfaces, we also give a full computation of the genus zero open Gromov-Witten invariants.

CONTENTS

1. Introduction

Let (X, ω) be a symplectic manifold of dimension $2n$, and let $L \subset X$ be a relatively spin Lagrangian submanifold. In [\[12\]](#page-43-1), the relative quantum cohomology $QH(X, L)$ is defined under certain conditions, for example, if L is a real cohomology sphere. The associativity of the relative quantum product is shown to be equivalent to the open and closed WDVV equations.

This paper gives explicit computations of the small relative quantum ring structure for some complete intersections, as well as an explicit treatment and a full computation of all open Gromov-Witten invariants of the quadric, as follows.

We start by formulating the results for Beauville-type [\[1\]](#page-43-2) complete intersections. Let $X \subset \mathbb{C}P^{n+r}$ with $n \geq 3$ be a complete intersection of degree (d_1, \ldots, d_r) . The first Chern class of X is given by

$$
c_1(X) = n + 1 - \sum_{j=1}^{r} (d_j - 1).
$$

For the purposes of this section, assume that $n > 2\sum_{j=1}^{r} (d_j - 1) - 1$, meaning,

$$
2c_1(X) > n + 1. \t\t(1)
$$

Date: Feb. 2024.

²⁰²⁰ Mathematics Subject Classification. 53D45, 14N35 (Primary) 53D37, 53D12 (Secondary).

Throughout, we use the notation

$$
\Upsilon := \prod_{j=1}^r d_j^{d_j}, \qquad \delta := \sum_{j=1}^r (d_j - 1), \qquad d := \prod_{j=1}^r d_j.
$$

Let $v_1, \ldots, v_p \in H^n(X; \mathbb{R})$ be the primitive classes. For all $j, k \in \{1, \ldots, p\}$, write

$$
g_{jk} := \int_X v_j \wedge v_k.
$$

Finally, we will consider the following polynomials in the formal variables a, b, c :

$$
r(a) := a^{n+1} - q\Upsilon \cdot a^{\delta},
$$

$$
r_{jk}(a, b, c) := bc - \frac{1}{d}g_{jk} \cdot (a^n - q\Upsilon \cdot a^{\delta - 1}).
$$

The following is immediate from the main theorem of $[1]$. Here u is a variable that corresponds to the hyperplane class in $H^2(X; \mathbb{R})$, and v_1, \ldots, v_p , correspond to the primitive classes in $Hⁿ(X; \mathbb{R})$. Then the absolute quantum cohomology is given by

$$
QH^*(X) \simeq \mathbb{R}[[q]][u, v_1, \dots, v_p]/(r(u), r_{jk}(u, v_j, v_k), uv_j)_{j,k \in \{1, \dots, p\}}.
$$
\n(2)

Let $L \subset X$ be a Lagrangian submanifold such that L is homeomorphic to $Sⁿ$. In particular, $\bar{\omega} =$ Id and we take the unique relative spin structure with $w_{\rm s} = 0$. In view of Lemma [2.1](#page-6-1) below, we generalize the assumption [\(1\)](#page-0-1) to

$$
\mu > n + 1. \tag{3}
$$

Taking x to be a variable that corresponds to the hyperplane class in $\hat{H}^2(X, L; \mathbb{R})$ and let w_1, \ldots, w_P , correspond to the primitive classes in $\hat{H}^n(X, L; \mathbb{R})$. In particular, $\{w_j\}_{j=1}^P$ map to $\{v_j\}_{j=1}^p$ under $\widehat{H}^*(X, L; \mathbb{R}) \to H^*(X)$.

Consider the case $[L] \neq 0 \in H_n(X; \mathbb{R})$. Then $P = p - 1$, and we may assume without loss of generality that v_p corresponds to the Poincaré dual of [L]. Then we prove the following.

Theorem 1. If $[L] \neq 0$, then the relative quantum cohomology is given by

$$
QH^*(X, L) \simeq \mathbb{R}[[q]][x, w_1, \ldots, w_{p-1}]/(r(x), r_{jk}(x, w_j, w_k), xw_j)_{j,k \in \{1, \ldots, p-1\}}.
$$

The relative quantum cohomology injects into the absolute one via

$$
\begin{aligned} \mathfrak{a}: QH^*(X, L) &\longrightarrow QH^*(X) \\ q &\mapsto q, \quad x \mapsto u, \quad w_j \mapsto v_j. \end{aligned}
$$

Remark 1.1. One can verify directly by inspecting the relations ideal that the formula for $QH^*(X, L)$ indeed defines a subalgebra. Similarly, in the next theorem we get a a rank-1 extension of the algebra $QH^*(X)$. This is a general phenomenon, explained in more detail in Sections [2.2](#page-6-2) and [2.9](#page-12-0) below.

Now consider the case $[L] = 0 \in H_n(X;\mathbb{R})$. Then $P = p$, and $\hat{H}^*(X,L;\mathbb{R})$ has an additional generator to which we associate the variable y . Then we prove the following.

Theorem 2. If $[L] = 0$, then the relative quantum cohomology is given by

$$
QH^*(X, L) \simeq \mathbb{R}[[q]][x, w_1, \dots, w_p, y]/(r(x), r_{jk}(x, w_j, w_k), xw_j, y^2, yx, yw_j)_{j,k \in \{1, \dots, p\}}.
$$

The relative quantum cohomology surjects onto the absolute one via

$$
\mathfrak{a}: QH^*(X, L) \longrightarrow QH^*(X)
$$

 $q \mapsto q, \quad x \mapsto u, \quad w_j \mapsto v_j, \quad y \mapsto 0.$

For our first example computed explicitly, consider $(X, L) = (\mathbb{C}P^n, \mathbb{R}P^n)$. It is shown in [\[5,](#page-43-3) [9\]](#page-43-4) that the absolute quantum cohomology of $\mathbb{C}P^n$ is given by the ring $\mathbb{R}[[q]][u]/(u^{n+1}-q)$. To allow the absolute quantum cohomology to interact with the relative quantum cohomology, we extend the Novikov ring by adjoining $q^{1/2}$, and we twist the signs of quantum cohomology by the class in $H^2(\mathbb{C}P^n;\mathbb{Z}/2\mathbb{Z})$ determined by the relative spin structure on $\mathbb{R}P^n$. Thus we take

$$
QH^{*}(\mathbb{C}P^{n}) \simeq \mathbb{R}[[q^{1/2}]][u]/(u^{n+1}-(-1)^{\frac{n+1}{2}}q).
$$

Theorem 3. The relative quantum cohomology of the odd-dimensional projective space is given by

$$
QH^*(\mathbb{C}P^n, \mathbb{R}P^n) \simeq \mathbb{R}[[q^{1/2}]][x, y] \big/ I
$$

with

$$
I = (y^2 - 2q^{1/2}y, \quad x^{n+1} - (-1)^{\frac{n+1}{2}}q - (-1)^{\frac{n-1}{2}}\frac{1}{2}q^{1/2}y, \quad xy = 0).
$$

The relative quantum cohomology surjects onto the absolute one via

$$
\mathfrak{a}: QH^*(\mathbb{CP}^n, \mathbb{RP}^n) \longrightarrow QH^*(\mathbb{CP}^n)
$$

$$
q^{1/2} \mapsto q^{1/2}, \quad x \mapsto u, \quad y \mapsto 0.
$$

Remark 1.2. Note that the case $(\mathbb{CP}^n, \mathbb{RP}^n)$ is not covered by the Beauville-style complete intersection theorem, since $\mu = n + 1$. This indeed makes the relations in the quantum ring substantially different: While in Theorem [2](#page-2-0) the additional generator y gave rise to an independent nilpotent component, here y satisfies non-trivial relations, and in particular changes non-trivially the relation for x^{n+1} .

Further, we carry out direct, definition-based computations of the small relative quantum cohomology in the special case when X is a quadric hypersurfaces and $L \simeq S^n$ is the real locus with respect to a particular real structure for arbitrary $n > 1$. Again, we take the unique relative spin structure with $w_s = 0$. This family of examples, while in some ways straightforward, offers a comprehensive overview of the principles at play. Indeed, the Lagrangian submanifold of the quadric is homologically nontrivial if and only if the quadric is even dimensional. Direct calculations in the two cases, of being homologically trivial or nontrivial, require the different tools of the wall-crossing formula and the open-closed relation, respectively, proven in [\[12\]](#page-43-1) and cited below as Theorems [14](#page-11-0) and [15.](#page-11-1) More concretely, we get the following results.

Let X be a quadric hypersurface in \mathbb{CP}^{n+1} with odd n given by $\sum_{j=0}^{n} z_j^2 - z_{n+1}^2 = 0$. It is shown in [\[4\]](#page-43-5) for $n = 3$ and in [\[1\]](#page-43-2) for $n \geq 3$ that the absolute quantum cohomology of X is given by

$$
QH^*(X) \simeq \mathbb{R}[[q]][u]/(u^{n+1} - 4qu).
$$

See also Proposition [6.1](#page-26-0) below.

Theorem 4. Let $L \subset X$ be the Lagrangian submanifold given by the real locus of complex conjugation. The relative quantum cohomology is given by

$$
QH^*(X,L) \simeq \mathbb{R}[[q]][x,y]/I
$$

with

$$
I = (x^{n+1} - 4qx, xy, y^2).
$$

The relative quantum cohomology surjects onto the absolute one via

$$
\mathfrak{a}: QH^*(X, L) \longrightarrow QH^*(X)
$$

$$
q \mapsto q, \quad x \mapsto u, \quad y \mapsto 0.
$$

Now, let X be a quadric hypersurface in \mathbb{CP}^{n+1} with even n given by $\sum_{j=0}^{n} z_j^2 - z_{n+1}^2 = 0$. As shown in [\[1\]](#page-43-2) for $n \geq 4$ and revisited in Proposition [7.2](#page-32-0) below,

the absolute quantum cohomology of X is given by

$$
QH^*(X) \simeq \mathbb{R}[[q]][u, v]/I,
$$

with

$$
I = (uv, v2 - (-1)n/2+14q - (-1)n/2un).
$$

Theorem 5. Let $L \subset X$ be the Lagrangian submanifold given by the real locus of complex conjugation. The relative quantum cohomology is given by

$$
QH^*(X,L) \simeq \mathbb{R}[[q]][x]/(x^{n+1} - 4qx).
$$

The relative quantum cohomology injects into the absolute one via

$$
\begin{aligned} \mathfrak{a}: QH^*(X,L) &\longrightarrow QH^*(X) \\ q &\mapsto q, \quad x\mapsto u. \end{aligned}
$$

Lastly, let X be the quadric hypersurface in \mathbb{CP}^3 given by $\sum_{j=0}^2 z_j^2 - z_3^2 = 0$. As shown in [\[3,](#page-43-6) Example 7.2, the absolute quantum cohomology of X is given by $\mathbb{R}[[q_1, q_2]][l, l^*]/(l^2 - q_2, l^{*2} - q_1)$. As with the example of the projective space, we need to modify the Novikov ring in order to allow the absolute quantum cohomology to interact with the relative one. As explained in Section [8,](#page-34-0) this is done by identifying q_1 and q_2 . Thus we take

$$
QH^*(X) \simeq \mathbb{R}[[q]][l, l^*]/(l^2 - q, \quad l^{*2} - q).
$$

Changing basis to $u = l + l^*$ and $v = l - l^*$ gives

$$
QH^*(X) \simeq \mathbb{R}[[q]][u, v]/(u^2 + v^2 - 4q, uv),
$$

Which agrees with the result obtained for the general even-dimensional quadric hypersurfaces.

Theorem 6. Let $L \subset X$ be the Lagrangian submanifold given by the real locus of complex conjugation. The relative quantum cohomology is given by

$$
QH^*(X,L) \simeq \mathbb{R}[[q]][x]/(x^3 - 4qx).
$$

The relative quantum cohomology injects into the absolute one via

$$
\mathfrak{a}: QH^*(X, L) \longrightarrow QH^*(X)
$$

$$
q \mapsto q, \quad x \mapsto u.
$$

For each of the above explicit examples we verify that all the closed and open Gromov-Witten invariants are computable via WDVV-based recursions. This is akin to spelling out the big quantum product. The computability of the closed invariants of the odd dimensional quadric is given by [\[5,](#page-43-3)[9\]](#page-43-4), while the computation of the open invariants is new. For the even dimensional quadric, even the closed invariants require special care, since the cohomology of the ambient manifold is not generated by divisors. We prove the following:

Theorem 7. Let X be a quadric hypersurface in \mathbb{CP}^{n+1} with odd n given by $\sum_{j=0}^{n} z_j^2$ – $z_{n+1}^2 = 0$. Let $L \cong S^n$ be the real locus of X. The open Gromov-Witten invariants of (X, L) are entirely determined by the open WDVV equations, the axioms of OGW, the wall-crossing formula Theorem [14,](#page-11-0) the closed Gromov-Witten invariants of X, and $\text{OGW}_{1,3} =$ $(-1)^{\frac{(n-2)(n-3)}{2}}$ 2.

We observe an interesting property of the open Gromov-Witten invariants of the odd dimensional quadrics, which follows from a general result regarding compatibility of the open WDVV equations and sign changes (see Proposition [2.10\)](#page-12-1).

Proposition 1.3. For the odd dimensional quadrics, suppose that we change the sign of the initial value $\text{OGW}_{1,3}$, then all the other OGW only change by a sign computed in Proposition [6.9.](#page-30-0)

We include some sample values of open Gromov-Witten invariants for various odd quadrics (tables [1](#page-4-0) and [2\)](#page-5-0).

Theorem 8. Let X be a quadric hypersurface in \mathbb{CP}^{n+1} with even n given by $\sum_{j=0}^{n} z_j^2 - z_{n+1}^2 =$ 0. Let $L \cong S^n$ be the real locus of X. The open Gromov-Witten invariants of (X, L) are entirely determined by the axioms of OGW, the closed Gromov-Witten invariants of X, and the open-closed relation Theorem [15.](#page-11-1) Moreover, the closed Gromov-Witten invariants of X are entirely determined by the axioms of GW, the WDVV equations, and $GW_1(\Delta_{n-1}, \Delta_n)$ 4.

Theorem 9. Let X be a quadric hypersurface in \mathbb{CP}^3 given by $\sum_{j=0}^3 z_j^2 - z_4^2 = 0$. Let $L \cong S^2$ be the real locus of X. The open Gromov-Witten invariants of (X, L) are entirely determined by the axioms of OGW , the closed Gromov-Witten invariants of X, and the openclosed relation Theorem [15.](#page-11-1) Moreover, the closed Gromov-Witten invariants of X are entirely determined by the axioms of GW, the WDVV equations, and the basic closed Gromov-Witten invariants (see Lemma [8.1\)](#page-36-1).

We include some sample values of open Gromov-Witten invariants for various odd quadrics (tables [3](#page-5-1) and [4\)](#page-5-2).

TABLE 3. Values of $OGW_{\beta,1}^4(\Gamma_2^{\otimes l_2} \otimes \Gamma_3^{\otimes l_3})$ with $l_2 = 4\beta - 1 - 2l_3$.

1.1. Acknowledgments. The authors would like to thank Jake Solomon for essentially collaborating on parts of this manuscript, particularly Section [5](#page-18-0) and the appendix. The first author would like to thank Nick Sheridan for suggesting the project of computing open Gromov-Witten invariants for odd-dimensional quadrics. The first author was partly supported by the Royal Society Research Grant for Research Fellows: RGFnR1n181009ERC, the ERC Starting Grant 850713 – HMS and EPSRC Grant EP/W015749/1. The second author was partly supported by NSF grant No. DMS-1638352, ISF grant No. 2793/21, and the Colton Foundation.

2. Background

2.1. Setting and notation. Let (X,ω) be closed symplectic manifold of dimension $2n$, J an ω -tame almost complex structure, and $L \subset X$ a Lagrangian submanifold with a relative spin structure $\boldsymbol{\mathfrak{s}}$. Recall that $\boldsymbol{\mathfrak{s}}$ determines a class

$$
w_{\mathfrak{s}} \in H^2(X; \mathbb{Z}/2\mathbb{Z})
$$

such that $w_2(TL) = i^*w_{\mathfrak{s}}$. By abuse of notation we think of $w_{\mathfrak{s}}$ as a homomorphism $w_{\mathfrak{s}}$: $H_2(X;\mathbb{Z}) \to \mathbb{Z}/2\mathbb{Z}$. Let

$$
\varpi: H_2(X; \mathbb{Z}) \longrightarrow H_2(X, L; \mathbb{Z})
$$

be the map from the long exact sequence of the pair (X, L) . Let

$$
\mu: H_2(X,L;\mathbb{Z}) \longrightarrow \mathbb{Z}
$$

be the Maslov index as defined in [\[2\]](#page-43-7). The following is well known.

Lemma 2.1. We have $\mu(\varpi(\beta)) = 2c_1(\beta)$.

2.2. Bases. Consider the subcomplex of differential forms on X consisting of those with trivial integral on L,

$$
\widehat{A}^*(X, L) := \left\{ \eta \in A^*(X) \mid \int_L \eta = 0 \right\},\
$$

and write $\hat{H}^*(X, L; \mathbb{R}) := H^*(\hat{A}^*(X, L), d)$. This version of relative cohomology fits in the long exact sequence

$$
\widehat{H}^*(X, L; \mathbb{R}) \xrightarrow{\rho} H^*(X; \mathbb{R})
$$
\n
$$
\mathbb{R}[-n],
$$
\n(4)

where

$$
i_{\mathbb{R}}:H^*(X;\mathbb{R})\longrightarrow\mathbb{R}
$$

is the map given by integration over L. If $[L] \neq 0 \in H_n(X;\mathbb{R})$, the long exact sequence [\(4\)](#page-6-3) becomes a short exact sequence

$$
0 \longrightarrow \widehat{H}^*(X, L) \stackrel{\rho}{\longrightarrow} H^*(X) \stackrel{i_{\mathbb{R}}}{\longrightarrow} \mathbb{R}[-n] \longrightarrow 0. \tag{5}
$$

If $[L] = 0$, the sequence becomes

$$
0 \longrightarrow \mathbb{R}[-n-1] \stackrel{y[-1]}{\longrightarrow} \widehat{H}^*(X, L; \mathbb{R}) \stackrel{\rho}{\longrightarrow} H^*(X; \mathbb{R}) \longrightarrow 0. \tag{6}
$$

Set $M := \text{rk } H^*(X; \mathbb{R}) - 1$, $N := \text{rk } \tilde{H}^*(X, L; \mathbb{R}) - 1$, and $K = \min\{M, N\}$. In particular, $K + 1 = \max\{M, N\}.$ Choose homogeneous bases

$$
\Delta_0, \dots, \Delta_M \in H^*(X; \mathbb{R}) \quad \text{and} \quad \Gamma_0, \dots, \Gamma_N \in \widehat{H}^*(X, L; \mathbb{R})
$$

such that $\rho(\Gamma_j) = \Delta_j$ for $j = 0, \ldots, K$, and if $N > M$ also $\rho(\Gamma_N) = 0$. In particular, $\deg \Delta_j = \deg \Gamma_j$ for $j \leq K$. Additionally, set

$$
\Gamma_{\diamond} := y(1) \in \widehat{H}^*(X, L; \mathbb{R}) \quad \text{and} \quad \Delta_L := PD([L]) \in H^*(X; \mathbb{R}). \tag{7}
$$

Note that exactly one of the two does not vanish, depending on whether $[L] \in H_*(X;\mathbb{R})$ is trivial or not. If $|L| = 0$, we assume without loss of generality that the bases are chosen so that $\Gamma_N = \Gamma_{\diamond}$. For convenience, we define $\Gamma_M := 0$ if $M > N$ and $\Delta_N := 0$ if $N > M$.

Remark 2.2. By [\[13,](#page-43-8) Lemma 5.14], $\hat{H}^*(X, L; \mathbb{R})$ is isomorphic to the standard (reduced) relative cohomology when $H^*(L;\mathbb{R}) \simeq H^*(S^n;\mathbb{R})$. In particular, this is true in all the cases considered in the current paper.

2.3. Rings. Define a Novikov coefficient ring

$$
\Lambda = \left\{ \sum_{i=0}^{\infty} a_i T^{\beta_i} \middle| a_i \in \mathbb{R}, \beta_i \in H_2(X, L; \mathbb{Z}), \omega(\beta_i) \ge 0, \lim_{i \to \infty} \omega(\beta_i) = \infty \right\}.
$$

Grading on Λ is defined by declaring T^{β} to be of degree $\mu(\beta)$.

Let $\Gamma_0, \ldots, \Gamma_N$, and $\Delta_0, \ldots, \Delta_M$, be the bases from Section [2.2.](#page-6-2) Let t_0, \ldots, t_{K+1} , be formal variables with degrees $\deg t_j = 2 - \Delta_j$ for $j \leq M$ and $\deg t_j = 2 - \deg \Gamma_j$ for $j \leq N$. Let s be a formal variable with degree deg $s = 1 - n$. We will use Δ_j , Γ_j , for interior constraints of closed and open Gromov-Witten invariants, respectively, while the variable s will correspond to boundary point constraints for open invariants.

Define graded-commutative rings

$$
R := \Lambda[[t_0,\ldots,t_N,s]], \qquad Q := \Lambda[[t_0,\ldots,t_N]], \qquad Q_X := \Lambda[[t_0,\ldots,t_M]].
$$

Define

$$
\rho^*: Q_X \longrightarrow Q
$$

to be the Λ algebra homomorphism determined by

$$
\rho^*(t_j) = \begin{cases} t_j, & j \le K, \\ 0, & j > K. \end{cases}
$$

The ring R comes with a valuation

$$
\nu: R \longrightarrow \mathbb{R},
$$

given by

$$
\nu\left(\sum_{j=0}^{\infty}a_jT^{\beta_j}s^k\prod_{i=0}^N t_i^{l_{ij}}\right)=\inf_{\substack{j\\a_j\neq 0}}\left(\omega(\beta_j)+k+\sum_{i=0}^N l_{ij}\right),
$$

and similarly for Q and Q_X . For $\Upsilon \in \{R, Q, Q_X\}$, write $H^*(X, \Upsilon)$ and $\widetilde{H}^*(X, L; \Upsilon)$ for $H^*(X;\mathbb{R})\otimes_{\mathbb{R}} \Upsilon$ and $\hat{H}^*(X,L;\mathbb{R})\otimes_{\mathbb{R}} \Upsilon$, respectively. Here the tensor is completed with respect to the valuation on Υ.

Remark 2.3. Comparing with the notation convention of [\[12\]](#page-43-1), throughout the paper we take $\Pi = H_2(X, L; \mathbb{Z})$ and $U = H^*(X)$. Furthermore, here we write R, Q, and Q_X instead of R_W , Q_W , and Q_U , respectively; we write $QH^*(X)$ for $QH_U^*(X)$ and $QH^*(X, L)$ for $QH_U(X, L)$.

Remark 2.4. In [\[12\]](#page-43-1), we used two different Novikov coefficient rings, Λ and Λ_c . This was essential in the construction of a cone complex that provides a natural setting for definitions and proofs in open Gromov-Witten theory, where all cases can be treated uniformly. The current paper, however, has more limited aims, which allows us to simplify exposition by removing Λ_c .

2.4. Closed GW. The genus zero closed Gromov-Witten invariants are a collection of maps

$$
GW_{\beta}: H^*(X; \mathbb{R})^{\otimes l} \longrightarrow \mathbb{R}, \qquad \beta \in H_2(X; \mathbb{Z}), \quad l \in \mathbb{Z}_{\geq 0}.
$$

Intuitively, for $A_1, \ldots, A_l \in H^*(X; \mathbb{R})$, the invariant $\text{GW}_\beta(A_1, \ldots, A_l)$ is the algebraic count of J-holomorphic spheres of degree β that pass through chains representing the Poincaré duals of the classes A_j .

These invariants can be combined into the potential function $\Phi \in Q_X$, which can be written explicitly in coordinates as follows. Let $\Delta_j \in H^*(X;\mathbb{R})$ be the basis from Section [2.2](#page-6-2) associated with the variables t_0, \ldots, t_M . Then

$$
\Phi = \Phi(\{t_i\}_{i=0}^M) = \sum_{\substack{\beta \in H_2(X;\mathbb{Z}) \\ r_i \ge 0}} (-1)^{w_{\mathfrak{s}}(\beta)} \frac{T^{\varpi(\beta)} \prod_{i=0}^M t_{M-i}^{r_{M-i}}}{\prod_{i=0}^M r_i!} \text{GW}_{\beta}(\otimes_{i=0}^M \Delta_i^{\otimes r_i}).
$$

The non-standard sign $(-1)^{w_s(\beta)}$ allows the closed Gromov-Witten invariants encoded in Φ to interact with the open Gromov-Witten invariants of the Lagrangian L, which depend on the relative spin structure s. See in particular Theorem [13](#page-10-0) below.

As shown in, e.g., [\[5,](#page-43-3) [7,](#page-43-9) [9\]](#page-43-4), the Gromov-Witten invariants have the following properties.

Theorem 10 (Closed Gromov-Witten axioms). The invariants GW of X satisfy the following axioms. Let $A_j \in H^*(X; \mathbb{R})$ for $j = 1, \ldots, l$.

(a) (Degree) $\text{GW}_{\beta}(A_1, \ldots, A_l) = 0$ unless

$$
2n - 6 + 2c_1(\beta) + 2l = \sum_{j=1}^{l} |A_j|.
$$
 (8)

(b) (Unit / Fundamental class)

$$
GW_{\beta}(1, A_1, \dots, A_{l-1}) = \begin{cases} \int_X A_1 \wedge A_2, & (\beta, l) = (\beta_0, 3), \\ 0, & otherwise. \end{cases}
$$
(9)

 (c) (Zero)

$$
GW_{\beta_0}(A_1,\ldots,A_l) = \begin{cases} \int_X A_1 \wedge A_2 \wedge A_3, & l = 3, \\ 0, & otherwise. \end{cases}
$$
 (10)

(d) (Divisor) If $|A_l| = 2$, then

$$
GW_{\beta}(A_1,\ldots,A_l) = \int_{\beta} A_l \cdot GW_{\beta}(A_1,\ldots,A_{l-1}).
$$
\n(11)

In addition, the potential Φ satisfies the Witten-Dijkgraaf-Verlinde-Verlinde (WDVV) equations, given in [\[5,](#page-43-3) Theorem-Definition 4.5] and [\[8,](#page-43-10) Theorem 3]. To formulate them, we need some additional notation. Set

$$
g_{ij} := \int_X \Delta_i \wedge \Delta_j,
$$

and define $(g^{ij})_{i,j=0}^M$ to be the matrix inverse to $(g_{ij})_{i,j=0}^M$.

Theorem 11 (Closed WDVV equations). Let $u, v, w \in \{0, ..., M\}$. Then,

$$
\sum_{m,l=0}^{M} \partial_{t_u} \partial_{t_v} \partial_{t_m} \Phi \cdot g^{ml} \cdot \partial_{t_l} \partial_{t_w} \partial_{t_y} \Phi = (-1)^{|\Delta_u|(|\Delta_v| + |\Delta_w|)} \sum_{m,l=0}^{M} \partial_{t_v} \partial_{t_w} \partial_{t_m} \Phi \cdot g^{ml} \cdot \partial_{t_l} \partial_{t_u} \partial_{t_y} \Phi. (12)
$$

2.5. Open GW. In many situations, genus zero open Gromov-Witten invariants are defined in [\[12,](#page-43-1) [13\]](#page-43-8). For the purposes of this paper, we focus on the situation when $H^*(L;\mathbb{R}) \simeq$ $H^*(S^n; \mathbb{R})$, which includes all the examples discussed here. The genus zero open Gromov-Witten invariants are a collection of maps

$$
\mathrm{OGW}_{\beta,k} : \widehat{H}^*(X, L; \mathbb{R})^{\otimes l} \to \mathbb{R}, \qquad \beta \in H_2(X, L; \mathbb{Z}), \quad k, l \in \mathbb{Z}_{\geq 0}.
$$

Intuitively, for $A_1, \ldots, A_l \in \dot{H}^*(X, L; \mathbb{R})$, the invariant $\mathrm{OGW}_{\beta,k}(A_1, \ldots, A_l)$ counts configurations of J-holomorphic disks that collectively have degree β and pass through k points in L and chains in $X \setminus L$ representing the Poincaré duals of A_j ; which disks join into such configurations is determined by a bounding chain. See [\[13,](#page-43-8) Section 1.2.4] for more details. The open Gromov-Witten invariants can be arranged in the open superpotential $\Omega \in \mathbb{R}$, which can be written explicitly in coordinates as follows. Let $\Gamma_j \in \hat{H}^*(X, L; \mathbb{R})$ be the basis from Section [2.2](#page-6-2) associated with the variables t_0, \ldots, t_N , and think of the additional variable s as corresponding to the Poincaré dual of a point in $Hⁿ(L;\mathbb{R})$. Then the superpotential $\Omega \in R$ is given by

$$
\Omega = \Omega(s, \{t_i\}_{i=0}^N) = \sum_{\substack{\beta \in \Pi \\ k \ge 0 \\ r_i \ge 0}} \frac{T^{\beta} s^k \prod_{i=0}^N t_{N-i}^{r_{N-i}}}{k! \prod_{i=0}^N r_i!} \text{OGW}_{\beta, k}(\otimes_{i=0}^N \Gamma_i^{\otimes r_i}). \tag{13}
$$

By design, if $k = 0$ and $\beta \in \text{Im }\varpi$ then $\text{OGW}_{\beta,k} \equiv 0$. See [\[13,](#page-43-8) Remark 1.8]. In the case when $[L] = 0 \in H_n(X;\mathbb{R})$, enhanced invariants

$$
\overline{\text{OGW}}_{\beta,k} : \widehat{H}^*(X,L;\mathbb{R})^{\otimes l} \to \mathbb{R}, \qquad \beta \in H_2(X,L;\mathbb{Z}), \quad k, l \in \mathbb{Z}_{\geq 0},
$$

are defined in [\[12\]](#page-43-1), such that

$$
\text{OGW}_{\beta,k}(\eta_1,\ldots,\eta_l) = \begin{cases} 0, & k = 0 \text{ and } \beta \in \text{Im}\,\varpi, \\ \overline{\text{OGW}}_{\beta,k}(\eta_1,\ldots,\eta_l), & \text{otherwise.} \end{cases}
$$
(14)

These can be arranged into the enhanced superpotential

$$
\overline{\Omega}(s, \{t_i\}_{i=0}^N) = \sum_{\substack{\beta \in \Pi \\ k \ge 0 \\ r_i \ge 0}} \frac{T^{\beta} s^k \prod_{i=0}^N t_{N-i}^{r_{N-i}}}{k! \prod_{i=0}^N r_i!} \overline{\text{OGW}}_{\beta,k}(\otimes_{i=0}^N \Gamma_i^{\otimes r_i}).
$$
\n(15)

When $k = 0$ and $\beta \in \text{Im}\,\varpi$, the enhanced invariants $\overline{\text{OGW}}_{\beta,k}$ do not necessarily vanish. However, their values depend on an additional piece of data: A homomorphism

$$
P_{\mathbb{R}} : \widehat{H}^*(X, L; \mathbb{R}) \longrightarrow \text{Coker}(i_{\mathbb{R}})
$$
\n
$$
(16)
$$

that is a left inverse to the map $\bar{y}: \mathrm{Coker} i_{\mathbb{R}} \to \hat{H}^*(X, L; \mathbb{R})$ induced by y from the long exact sequence [\(4\)](#page-6-3). When $[L] \neq 0$, note that Coker $(i_{\mathbb{R}}) = 0$ and therefore $P_{\mathbb{R}} \equiv 0$. When $[L] = 0$, a choice of $P_{\mathbb{R}}$ is equivalent to a choice of a splitting in [\(6\)](#page-6-4). Another perspective on the choice of $P_{\mathbb{R}}$ is explained in [\[12,](#page-43-1) Remark 4.12]: It is equivalent to the choice of a homology class $V \in H_{n+1}(X, L; \mathbb{R})$ such that $\partial V = [L].$

The open Gromov-Witten invariants satisfy the following properties analogous to the closed ones. For OGW this is [\[13,](#page-43-8) Theorem 4], and for OGW this is [\[12,](#page-43-1) Proposition 4.19].

Theorem 12 (Open Gromov-Witten axioms). The invariants \overline{OGW} of (X, L) satisfy the following axioms. Let $A_j \in W$ for $j = 1, \ldots, l$.

(a) (Degree) $\overline{\text{OGW}}_{\beta,k}(A_1,\ldots,A_l)=0$ unless

$$
n - 3 + \mu(\beta) + k + 2l = kn + \sum_{j=1}^{l} |A_j|.
$$
 (17)

(b) (Unit / Fundamental class)

$$
\overline{\text{OGW}}_{\beta,k}(1, A_1, \dots, A_{l-1}) = \begin{cases} -1, & (\beta, k, l) = (\beta_0, 1, 1), \\ P_{\mathbb{R}}(A_1), & (\beta, k, l) = (\beta_0, 0, 2), \\ 0, & \text{otherwise.} \end{cases}
$$
(18)

 (c) (Zero)

$$
\overline{\text{OGW}}_{\beta_0,k}(A_1,\ldots,A_l) = \begin{cases}\n-1, & (k,l) = (1,1) \text{ and } A_1 = 1, \\
P_{\mathbb{R}}(A_1 \cup A_2), & (k,l) = (0,2), \\
0, & \text{otherwise.} \n\end{cases}
$$
\n(19)

(d) (Divisor) If $|A_l| = 2$, then

$$
\overline{\text{OGW}}_{\beta,k}(A_1,\ldots,A_l) = \int_{\beta} A_l \cdot \overline{\text{OGW}}_{\beta,k}(A_1,\ldots,A_{l-1}).
$$
\n(20)

The invariants OGW satisfy the same axioms with $P_{\mathbb{R}}$ replaced with 0.

The superpotential Ω satisfies an open version of the WDVV equations. To formulate it, we need some additional notation. Recall that $M = \text{rk } H^*(X;\mathbb{R}) - 1, N = \text{rk } \tilde{H}^*(X,L;\mathbb{R}) - 1$, and $K = min{M, N}$. In particular, $K = \text{rk Ker}(P_{\mathbb{R}}) - 1$. Possibly modifying our choice of $\{\Gamma_j\}_{j=0}^K$ from Section [2.2,](#page-6-2) we can assume that $\{\Gamma_j\}_{j=0}^K \subset \text{Ker}(P_\mathbb{R})$ is a basis. Set

$$
g_{ij} := \int_X \Delta_i \wedge \Delta_j, \qquad i, j = 0, \dots, M,
$$

and define $(g^{ij})_{i,j=0}^M$ to be the matrix inverse to $(g_{ij})_{i,j=0}^M$. Below is the statement of [\[12,](#page-43-1) Corollary 1.6].

Theorem 13 (Open WDVV equations for point-like bounding chains). Let $u, v, w \in \{0, \ldots, N\}$. If $[L] = 0$, then

$$
\sum_{\substack{0 \le l \le K \\ 0 \le m \le M}} \partial_{t_u} \partial_{t_l} \overline{\Omega} \cdot g^{lm} \cdot \rho^* \partial_{t_m} \partial_{t_w} \partial_{t_v} \Phi - \partial_{t_u} \partial_s \overline{\Omega} \cdot \partial_{t_w} \partial_{t_v} \overline{\Omega} =
$$
\n
$$
= \sum_{\substack{0 \le l \le M \\ 0 \le m \le K}} \rho^* \partial_{t_u} \partial_{t_w} \partial_{t_l} \Phi \cdot g^{lm} \cdot \partial_{t_m} \partial_{t_v} \overline{\Omega} - \partial_{t_u} \partial_{t_w} \overline{\Omega} \cdot \partial_{t_v} \partial_s \overline{\Omega}, \quad (21)
$$

and

$$
\sum_{\substack{0 \le l \le K \\ 0 \le m \le M}} \partial_s \partial_{t_l} \overline{\Omega} \cdot g^{lm} \cdot \rho^* \partial_{t_m} \partial_{t_w} \partial_{t_v} \Phi - \partial_s^2 \overline{\Omega} \cdot \partial_{t_w} \partial_{t_v} \overline{\Omega} = -\partial_s \partial_{t_w} \overline{\Omega} \cdot \partial_{t_v} \partial_s \overline{\Omega}. \tag{22}
$$

If $[L] \neq 0$, then equation [\(21\)](#page-10-1) holds with $\overline{\Omega}$ replaced by Ω .

2.6. Interplay. In addition to the standard Gromov-Witten-type axioms, we have two phenomena unique to the open setup. The first is a wall-crossing property, which allows us to trade boundary constraints for interior ones if L is homologically trivial in X . This is $[12,$ Theorem 6.

Theorem 14 (Wall-crossing). Suppose $[L] = 0$. The invariants \overline{OGW} satisfy

$$
\overline{\text{OGW}}_{\beta,k+1}(\eta_1,\ldots,\eta_l) = -\overline{\text{OGW}}_{\beta,k}(\Gamma_{\diamond},\eta_1,\ldots,\eta_l). \tag{23}
$$

The second relates open invariants with closed ones when L is homologically nontrivial in X. This is [\[12,](#page-43-1) Corollary 1.5]:

Theorem 15 (Open-closed relation). Suppose that $[L] \neq 0$. Let $\eta_1, \ldots, \eta_l \in \tilde{H}^*(X, L)$. If $k \geq 2$, or if $k \geq 1$ and $\beta \notin \text{Im}\,\varpi$, then

$$
\overline{\mathrm{OGW}}_{\beta,k}(\eta_1,\ldots,\eta_l)=0.
$$

Moreover, for $\beta \in \text{Im}(\varpi)$ and $\eta \in H^*(X; \mathbb{R})$ such that $\int_L \eta = 1$,

$$
\overline{\text{OGW}}_{\beta,1}(\eta_1,\ldots\eta_l) = \sum_{\substack{\hat{\beta}\in H_2(X;\mathbb{Z})\\ \varpi(\hat{\beta})=\beta}} (-1)^{n+1+w_{\mathfrak{s}}(\hat{\beta})} \text{GW}_{\hat{\beta}}(\eta,\Delta_L,\rho(\eta_1),\ldots,\rho(\eta_l)).\tag{24}
$$

2.7. Sign changes of the open GW. We also observe the following result about compatibility of the open Gromov-Witten axioms (Theorem [12\)](#page-10-2) and the open WDVV equations (Theorem [13\)](#page-10-0) with specific sign changes. When all open Gromov-Witten invariants are determined from certain initial values by the axioms and the open WDVV equations, this sometimes allows us to deduce how the invariants change when the initial values change sign. See Proposition [6.9.](#page-30-0) For the cleanest statements, we need to consider invariants with interior constraints only from $\ker(P_{\mathbb{R}})$

Definition 2.5. Restrict the invariants \overline{OGW} (or OGW when $[L] \neq 0$) to invariants

$$
\overline{\mathrm{OGW}}_{\beta, k}' : \mathrm{ker}(P_\mathbb{R})^{\otimes l} \to \mathbb{R}
$$

and consider the restricted superpotential

$$
\overline{\Omega}'(s, \{t_i\}_{i=0}^{N-1}) = \sum_{\substack{\beta \in \Pi \\ k \ge 0 \\ r_i \ge 0}} \frac{T^{\beta} s^k \prod_{i=0}^{N-1} t_{N-i}^{r_{N-i}}}{k! \prod_{i=0}^{N-1} r_i!} \overline{\text{OGW}}_{\beta,k}(\otimes_{i=0}^{N-1} \Gamma_i^{\otimes r_i}).
$$
\n(25)

Equivalently, $\overline{\Omega}' := \overline{\Omega}|_{t_N=0}$.

The following is clear from the definition:

Lemma 2.6. The invariants $\overline{\text{OGW}}'$ and OGW' satisfy the open Gromov-Witten axioms with $P_{\mathbb{R}}$ replaced with 0.

We also have:

Lemma 2.7. The restricted superpotentials $\overline{\Omega}'$ and Ω' still satisfy the open WDVV equations.

Proof. In the open WDVV equations, the summation indices run over $0 \leq l \leq K$ and $0 \leq m \leq M$ (as any integral $\bar{f}_X \Gamma_{\circ} \Gamma_j = 0$ for any j). This ensures that the equations still hold after restriction to $t_N = 0$.

Definition 2.8. For an arbitrary integer r consider the sign-changed invariants

$$
\overline{\text{OGW}}_{\beta,k}^{\circ} := (-1)^{\frac{k-1}{r}} \, \overline{\text{OGW}}_{\beta,k}',
$$

with a similar definition with OGW in place of OGW. We then also form the sign-changed superpotentials $\overline{\Omega}^{\circ}$ and Ω° by using the invariants $\overline{\text{OGW}}^{\circ}$ and $\overline{\text{OGW}}^{\circ}$ in their definitions.

Remark 2.9. Of course, in order for the invariants to be real numbers (or even rational) the above only makes sense if $\overline{\text{OGW}}_{\beta,k} = 0$ for all k such that $r \nmid k-1$. We will ignore this issue for now.

Finally, we can state the new result:

Proposition 2.10. For any choice of r, the invariants \overline{OGW}° and OGW° still satisfy the open Gromov-Witten axioms, and the superpotentials $\overline{\Omega}^{\circ}$ and Ω° still satisfy the open WDVV equations.

Proof. The first statement holds as the only non-zero invariants featuring in the open Gromov-Witten axioms have $k = 1$. For the second statement, consider the coefficient of s^k in Equation [\(21\)](#page-10-1). The first term on the left-hand side has its sign changed by $(-1)^{\frac{k-1}{r}}$ compared to the equation for $\overline{\Omega}$. The second term has it changed by $(-1)^{\frac{k_1}{r}}(-1)^{\frac{k_2-1}{r}} = (-1)^{\frac{k-1}{r}}$ as required. Similar reasoning holds for the right-side of the equation, so that the overall equality is maintained. The same holds for Equation [\(22\)](#page-11-2). \Box

2.8. Closed quantum cohomology. Write $QH^*(X) := H^*(X;\Lambda)$. The small quantum product

$$
: QH^(X) \otimes QH^*(X) \longrightarrow QH^*(X)
$$

was first introduced in the physics literature [\[6,](#page-43-11) [14\]](#page-43-12), and given a mathematical definition in $[5, 8, 9]$ $[5, 8, 9]$ $[5, 8, 9]$. The version we use in the current work is given by the formula

$$
\Delta_v * \Delta_u = \sum_{\substack{0 \le l,m \le M \\ d \in H_2(\overline{X};\mathbb{Z})}} (-1)^{w_s(d)} T^{\varpi(d)} \operatorname{GW}_d(\Delta_v, \Delta_u, \Delta_l) \cdot g^{lm} \cdot \Delta_m, \qquad 0 \le u, v \le M. \tag{26}
$$

This product is well known [\[5,](#page-43-3) [8,](#page-43-10) [9\]](#page-43-4) to be associative and graded commutative.

2.9. Relative quantum cohomology. The small relative quantum product is defined analogously, though the definition is a bit more delicate. Write $QH^*(X, L) := \hat{H}^*(X, L; \Lambda)$. Then the product 1

$$
\mathfrak{a}: QH^*(X, L) \otimes QH^*(X, L) \longrightarrow QH^*(X, L)
$$

 1 1 Read $\mathop{\mathtt{m}}$ as "mem".

is given by

$$
\mathbf{D}(\Gamma_v, \Gamma_u) = \sum_{\substack{0 \le m \le K \\ 0 \le l \le M \\ d \in H_2(X, L; \mathbb{Z})}} (-1)^{w_s(d)} T^{\varpi(d)} \text{GW}_d(\Delta_v, \Delta_u, \Delta_l) \cdot g^{lm} \cdot \Gamma_m +
$$
\n
$$
+ \sum_{d \in H_2(X, L; \mathbb{Z})} T^d \overline{\text{OGW}}_{d,0}(\Gamma_v, \Gamma_u) \cdot \Gamma_o, \qquad 0 \le u, v \le N.
$$
\n(27)

In the special case $[L] \neq 0$, since $\Gamma_{\diamond} = 0$, the expression simplifies to

$$
\mathbf{D}(\Gamma_v, \Gamma_u) = \sum_{\substack{0 \le m \le K \\ 0 \le l \le M \\ d \in H_2(X;\mathbb{Z})}} (-1)^{w_{\mathfrak{s}}(d)} T^{\varpi(d)} \, \mathrm{GW}_d(\Delta_v, \Delta_u, \Delta_l) \cdot g^{lm} \cdot \Gamma_m, \qquad 0 \le u, v \le N.
$$

The product \hat{p} is shown in [\[12\]](#page-43-1) to be associative and graded commutative. Though in [12] the definition of α is based on a chain-level construction, the above explicit formulas follow from [\[12,](#page-43-1) Lemma 5.5].

Remark 2.11. The definition of γ given in [\[12\]](#page-43-1) is free of choices. The explicit expression [\(27\)](#page-13-0) a priori appears to depend on the choice of $P_{\mathbb{R}}$ through the definition of $\overline{\text{OGW}}$, but this cancels with the dependence on $P_{\mathbb{R}}$ through the condition that $\{\Gamma_j\}_{j=0}^K \subset \text{Ker}(P_{\mathbb{R}})$ is a basis.

Lemma 2.12. The map ρ from [\(4\)](#page-6-3) induces an algebra homomorphism

$$
\mathfrak{a}: QH^*(X, L) \longrightarrow QH^*(X).
$$

Proof. This follows from equation (53) of [\[12\]](#page-43-1).

2.10. Big closed quantum cohomology. The product $*$ can be deformed by $\Delta = \sum_{j=0}^{M} t_j \Delta_j$, resulting in the big closed quantum product. More explicitly, set $QH_{big}^*(X) := QH^*(X) \otimes Q$. Then

$$
*^{big}: QH^*_{big}(X) \otimes QH^*_{big}(X) \longrightarrow QH^*_{big}(X)
$$

is given by

$$
\Delta_v *^{big} \Delta_u = \sum_{\substack{0 \le l,m \le M \\ d \in H_2(X;\mathbb{Z}) \\ p \ge 0}} (-1)^{w_s(d)} T^{\varpi(d)} \frac{1}{p!} \, \mathrm{GW}_d(\Delta^{\otimes p} \otimes \Delta_v \otimes \Delta_u \otimes \Delta_l) \cdot g^{lm} \cdot \Delta_m, \qquad 0 \le u, v, \le M.
$$

This product is well known [\[5,](#page-43-3) [8,](#page-43-10) [9\]](#page-43-4) to be associative and graded commutative.

2.11. Big relative quantum cohomology. The product α can be deformed by $\Gamma =$ $\sum_{j=0}^{N} t_j \Gamma_j$, resulting in the big relative quantum product. More explicitly, set $QH_{big}^*(X, L) :=$ $QH^*(X,L) \otimes Q$. The product

$$
\mathbf{a}^{big}: QH_{big}^*(X, L) \otimes QH_{big}^*(X, L) \longrightarrow QH_{big}^*(X, L)
$$

$$
\qquad \qquad \Box
$$

is given by

$$
\mathbf{a}^{big}(\Gamma_v, \Gamma_u) = \sum_{\substack{0 \le m \le K \\ 0 \le l \le M \\ d \in H_2(X, L; \mathbb{Z}) \\ p \ge 0}} (-1)^{w_s(d)} T^{\varpi(d)} \frac{1}{p!} \text{GW}_d(\Delta^{\otimes p} \otimes \Delta_v \otimes \Delta_u \otimes \Delta_l) \cdot g^{lm} \cdot \Gamma_m + \sum_{\substack{0 \le l \le M \\ p \ge 0}} T^d \frac{1}{p!} \overline{\text{OGW}}_{d,0}(\Gamma^{\otimes p} \otimes \Gamma_v \otimes \Gamma_u) \cdot \Gamma_\diamond, \qquad 0 \le u, v \le N.
$$

When $[L] \neq 0$ the formula simplifies to

$$
\mathbf{a}^{big}(\Gamma_v, \Gamma_u) = \sum_{\substack{0 \le m \le K \\ 0 \le l \le M \\ d \in H_2(X;\mathbb{Z}) \\ p \ge 0}} (-1)^{w_s(d)} T^{\varpi(d)} \frac{1}{p!} \, \mathrm{GW}_d(\Delta^{\otimes p} \otimes \Delta_v \otimes \Delta_u \otimes \Delta_l) \cdot g^{lm} \cdot \Gamma_m,
$$
\n
$$
0 \le u, v \le N.
$$

The product \mathbf{a}^{big} is shown in [\[12\]](#page-43-1) to be associative and graded commutative. Though in [12] the definition of p^{big} is based on a chain-level construction, the above explicit formulas follow from [\[12,](#page-43-1) Lemma 5.5].

Remark 2.13. The definition of p^{big} given in [\[12\]](#page-43-1) is free of choices. The explicit expression [\(27\)](#page-13-0) a priori appears to depend on the choice of $P_{\mathbb{R}}$ through the definition of $\overline{\text{OGW}}$, but this cancels with the dependence on $P_{\mathbb{R}}$ through the condition that $\{\Gamma_j\}_{j=0}^K \subset \text{Ker}(P_{\mathbb{R}})$ is a basis.

Lemma 2.14. The map ρ from [\(4\)](#page-6-3) induces an algebra homomorphism

$$
\mathfrak{a}_{big}: QH^*_{big}(X,L) \longrightarrow QH^*_{big}(X).
$$

Proof. This follows from equation (53) of [\[12\]](#page-43-1).

3. Complete intersections

In this section, we prove Theorems [1](#page-1-0) and [2.](#page-2-0)

3.1. Closed quantum cohomology. We choose a basis $\{\Delta_j\}_{j=0}^M$ of $H^*(X;\mathbb{R})$ as follows. Let $\Delta_1 \in H^2(X;\mathbb{R})$ be the hyperplane class, namely, writing ω_{FS} for the Fubini-Study form on \mathbb{CP}^{n+r} and taking

$$
\iota: X \hookrightarrow \mathbb{CP}^{n+r} \tag{28}
$$

to be the inclusion, we take $\Delta_1 = [\iota^* \omega_{FS}] \in H^2(X; \mathbb{R})$. Let $\Delta_j := \Delta_1^{\wedge j}$ for $j \in \{0, 1, \ldots, n\}$. Let $\{\Delta_j\}_{j=n+1}^M$ be generators of the primitive cohomology such if $[L] \neq 0$ then $\Delta_M = PD([L])$. Further, assume without loss of generality that Δ_j for $j > n$ are chosen so that $\{\Delta_j\}_{j=0}^K$ span $\text{Ker}(i_{\mathbb{R}})$. Denote the number of primitive basis elements by

$$
p:=M-n.
$$

To choose a basis for the relative cohomology $\hat{H}^*(X, L; \mathbb{R})$, and subsequently computing the quantum cohomology, we have to discuss the cases of $[L]$ being trivial and non-trivial separately.

3.2. L is homologically non-trivial. We choose a basis $\{\Gamma_j\}_{j=0}^N$ of $\hat{H}^*(X, L; \mathbb{R})$ as follows. Let $\Gamma_1 \in H^2(X, L; \mathbb{R})$ be the hyperplane class, namely, for ι the inclusion from [\(28\)](#page-14-1), we set $\Gamma_1 = [\iota^* \omega_{FS}] \in \widehat{H}^2(X, L; \mathbb{R})$. For $j \in \{0, 1, \ldots, n\}$, let $\Gamma_j := \Gamma_1^{\wedge j}$. In particular, $\rho(\Gamma_j) = \Delta_j$. Let $\{\Gamma_j\}_{j=n+1}^N$ be generators of the primitive part of $\tilde{H}^*(X, L; \mathbb{R})$, namely, homogeneous classes independent of $\Gamma_{n/2}$ (when n is even) such that $\rho(\Gamma_j) = \Delta_j$ for $j = n+1, \ldots, N$. Note that when $[L] \neq 0$ we have $N = M - 1$, and recall that $\Delta_M = PD([L])$.

Proof of Theorem [1.](#page-1-0) Lemma [2.12,](#page-13-1) exact sequence [\(4\)](#page-6-3) and the assumption that L is homologically non-trivial, imply that $QH^*(X, L)$ is the subalgebra of $QH(X, L)$ given by $\ker(i_{\mathbb{R}}) \otimes \mathbb{R}[[q]].$

3.3. L is homologically trivial. We choose a basis $\{\Gamma_j\}_{j=0}^N$ of $\hat{H}^*(X, L; \mathbb{R})$ as follows. Let $\Gamma_1 \in H^2(X, L; \mathbb{R})$ be the hyperplane class, namely, for ι the inclusion from [\(28\)](#page-14-1), we set $\Gamma_1 = [\iota^* \omega_{FS}] \in \widehat{H}^2(X, L; \mathbb{R})$. For $j \in \{0, 1, \ldots, n\}$, let $\Gamma_j := \Gamma_1^{\wedge j}$. In particular, $\rho(\Gamma_j) = \Delta_j$. Let $\{\Gamma_j\}_{j=n+1}^{N-1}$ be generators of the primitive part of $\tilde{H}^*(X,L;\mathbb{R})$, namely, homogeneous classes independent of $\Gamma_{n/2}$ (when n is even) such that $\rho(\Gamma_j) = \Delta_j$ for $j = n+1, \ldots, N-1$. Note that when $[L] = 0$ we have $N - 1 = M$. Finally, set $\Gamma_N := \Gamma_{\infty}$ from [\(7\)](#page-7-0).

Lemma 3.1. For any homogeneous $a_1, a_2 \in \hat{H}^*(X, L)$ with $|a_1| + |a_2| \leq 2n + 2$, we have $\overline{\text{OGW}}_{\beta,0}(a_1,a_2)=0.$

Proof. By the degree axiom [\(17\)](#page-10-3), the value of $\overline{OGW}_{\beta,0}(a_1, a_2)$ can only be nonzero if

$$
n + 1 + \mu(\beta) = |a_1| + |a_2| \le 2n + 2.
$$

But then $\mu(\beta) \leq n+1$, contradicting the assumption [\(3\)](#page-1-1) and Lemma [2.1.](#page-6-1)

Proof of Theorem [2.](#page-2-0) Let $a \in \{y, x, w_1, \ldots, w_p\}$. Since y is trivial in absolute cohomology, the GW term in equation [\(27\)](#page-13-0) vanishes in $\mathfrak{g}(y, a)$ and we obtain

$$
\mathbf{D}(y,a) = \sum_{\beta \in H_2(X,L;\mathbb{Z})} T^{\beta} \overline{\text{OGW}}_{\beta,0}(y,a) \cdot y.
$$

By Lemma [3.1,](#page-15-0) the right-hand side vanishes. Thus,

$$
ya = 0
$$
 for $a \in \{y, x, w_1, \dots, w_p\}.$ (29)

Recall the short exact sequence [\(6\)](#page-6-4). Let

$$
\sigma: H^*(X; \mathbb{R}) \longrightarrow \widehat{H}^*(X, L; \mathbb{R})
$$

be the section of ρ induced by $P_{\mathbb{R}}$ from [\(16\)](#page-9-0). In particular, $\sigma(u) = x$ and $\sigma(v_j) = w_j$. For

$$
(a_1, a_2) \in \{(x, x^j) \mid 1 \le j \le n\} \cup \{(x, w_j), (w_j, w_k) \mid j, k \in \{1, \ldots, p\}\},\
$$

equation [\(27\)](#page-13-0) and Lemma [3.1](#page-15-0) give

$$
\mathbf{a}(a_1, a_2) = \sum_{\substack{0 \le l,m \le K=M\\ d \in \mathbb{Z}_{\ge 0}}} T^d \operatorname{GW}_d(\rho(a_1), \rho(a_2), \Delta_l) \cdot g^{lm} \cdot \Gamma_m + \sum_{d \in \mathbb{Z}_{\ge 0}} T^d \overline{\operatorname{OGW}}_{d,0}(a_1, a_2) \cdot \Gamma_{\diamond}
$$

\n
$$
= \sum_{\substack{0 \le l,m \le K=M\\ d \in \mathbb{Z}_{\ge 0}}} T^d \operatorname{GW}_d(\rho(a_1), \rho(a_2), \Delta_l) \cdot g^{lm} \cdot \Gamma_m
$$

\n
$$
= \sigma(\rho(a_1) * \rho(a_2)). \tag{30}
$$

By induction, we obtain

$$
x^{i} = \sigma(u^{i}), \qquad i = 1, \dots, n. \tag{31}
$$

Combining equations (30) , (31) , and (2) , we obtain

$$
x^{n+1} = \mathbf{D}(x, x^n) = \sigma(u * u^n) = \sigma(q\Upsilon u^\delta) = q\Upsilon \sigma(u^\delta) = q\Upsilon x^\delta. \tag{32}
$$

Similarly,

$$
w_j w_k = \sigma(v_j v_k) = \frac{1}{d} g_{jk} \cdot (\sigma(u^n - q\Upsilon \cdot u^{\delta - 1})) = \frac{1}{d} g_{jk} \cdot (x^n - q\Upsilon \cdot x^{\delta - 1}).
$$
 (33)

Lastly, it follows directly from [\(30\)](#page-15-1) and [\(2\)](#page-1-2) that

$$
xw_j = 0, \qquad j = 1, \dots, p.
$$

Lemma [2.12](#page-13-1) and equation [\(2\)](#page-1-2) imply that $1, x, \ldots, x^n, w_1, \ldots, w_p, y$, generate $QH(X, L)$ as a A-module. From the associativity of α it follows that [\(29\)](#page-15-2), [\(32\)](#page-16-2), and [\(33\)](#page-16-3), are the only non-trivial relations in $QH(X, L)$.

4. Projective space

4.1. **Topological preliminaries.** In this section, $(X, L) = (\mathbb{CP}^n, \mathbb{RP}^n)$ for odd $n \geq 3$. We take $\omega = \omega_{FS}$ the Fubini-Study form. We equip $\mathbb{RP}^n \subset \mathbb{CP}^n$ with the relative spin structure that gives

$$
OGW_{1,2} = 2,
$$

as explained in [\[12,](#page-43-1) Lemma 6.1]. We write $\Gamma_j = [\omega^j] \in \hat{H}^*(X, L; \mathbb{R})$ and $\Delta_j = [\omega^j] \in$ $H^*(X;\mathbb{R})$, for $j = 0,\ldots,n$. Then $\{\Delta_j\}_{j=0}^n$ form a basis of $H^*(X;\mathbb{R})$. Since $[L] = 0 \in$ $H_n(X;\mathbb{Z})$, by short exact sequence [\(6\)](#page-6-4), the classes $\{\Gamma_j\}_{j=0}^n\cup \{\Gamma_{\diamond}=y(1)\}\$ form a basis of $\hat{H}^*(X,L;\mathbb{R}).$

Lemma 4.1. (a) $H_2(X, L; \mathbb{Z}) \simeq \mathbb{Z}$. (b) The map $\varpi : H_2(X;\mathbb{Z}) \to H_2(X,L;\mathbb{Z})$ is given by multiplication by 2.

Identify $H_2(X, L; \mathbb{Z})$ with $\mathbb Z$ so that the non-negative integers correspond to classes $\beta \in$ $H_2(X, L; \mathbb{Z})$ with $\omega(\beta) \geq 0$, and similarly for $H_2(X; \mathbb{Z})$. It follows from Lemma [4.1](#page-16-4) that for any $\beta \in H_2(X, L; \mathbb{Z})$ and $\hat{\beta} \in H_2(X; \mathbb{Z})$, we have

$$
\int_{\beta} \Gamma_1 = \frac{1}{2}\beta, \qquad \int_{\hat{\beta}} \Delta_1 = \hat{\beta}.
$$
\n(34)

These values will be used when applying the divisor axioms [\(20\)](#page-10-4) and [\(11\)](#page-8-0).

Lemma 4.2. The relevant characteristic classes of (X, L) are as follows:

(a) For all $\hat{\beta} \in H_2(X;\mathbb{Z})$ we have

$$
c_1(\hat{\beta}) = (n+1)\hat{\beta}, \qquad w_{\mathfrak{s}}(\hat{\beta}) \equiv \frac{n+1}{2}\hat{\beta} \pmod{2}.
$$

(b) For all $\beta \in H_2(X, L; \mathbb{Z})$, we have $\mu(\beta) = (n+1)\beta$. 17

For $i, j \in \{0, \ldots, n\}$, the pairing on absolute cohomology is given by

$$
g_{ij} = \int_X \Delta_i \wedge \Delta_j = \delta_{i,n-j},\tag{35}
$$

where $\delta_{k,l}$ is the Kronecker delta. Therefore the inverse matrix $(g^{ij})_{i,j} := (g_{ij})_{i,j}^{-1}$ is also given by $g^{ij} = \delta_{i,n-j}$.

4.2. **Absolute quantum ring.** Using equation (35) , Lemmas [4.1](#page-16-4) and [4.2](#page-16-5) (a) , and the defi-nition of the quantum product [\(26\)](#page-12-4), for $i, j \in \{0, ..., n\}$, we obtain

$$
\Delta_i * \Delta_j = \sum_{\substack{0 \le m \le n \\ d \ge 0}} (-1)^{\frac{n+1}{2} \cdot d} T^{2d} \, \mathrm{GW}_d(\Delta_i, \Delta_j, \Delta_m) \Delta_{n-m}.
$$

It is shown in [\[5,](#page-43-3) Section 5.3] and [\[9,](#page-43-4) Example 8.5] that, over the coefficient ring $\mathbb{R}[[q]]$, the absolute quantum cohomology of the projective space is given by $\mathbb{R}[[q]][x]/(x^{n+1}-q)$. Here, x corresponds to Δ_1 and q corresponds to $T^{\varpi(1)} = T^2$. For our purposes, we modify the setup as follows. First, we extend the coefficient ring by adjoining $q^{1/2}$, which corresponds to $T = T^1$, thereby obtaining the coefficient ring $\mathbb{R}[[q^{1/2}]] \simeq \Lambda$. Further, we twist $\text{GW}_{\hat{\beta}}$ by the sign $(-1)^{w_s(\hat{\beta})} = (-1)^{\frac{n+1}{2}\hat{\beta}}$. Thus, we have the following.

Proposition 4.3. The absolute quantum cohomology is given by

$$
QH^*(X) \simeq \mathbb{R}[[q^{1/2}]][x]/(x^{n+1}-(-1)^{\frac{n+1}{2}}q).
$$

4.3. Relative quantum ring. Using equation [\(35\)](#page-17-0), Lemmas [4.1](#page-16-4) and [4.2](#page-16-5)[\(a\),](#page-16-6) and the defi-nition of the relative quantum product [\(27\)](#page-13-0), for $i, j \in \{0, ..., n\} \cup \{\infty\}$, we obtain

$$
\mathbf{D}(\Gamma_i, \Gamma_j) = \sum_{\substack{0 \le m \le n \\ d \ge 0}} (-1)^{\frac{n+1}{2} \cdot d} T^{2d} \, \mathrm{GW}_d(\Delta_i, \Delta_j, \Delta_m) \Gamma_{n-m} + \sum_{d \ge 0} T^d \, \overline{\mathrm{OGW}}_{d,0}(\Gamma_i, \Gamma_j) \Gamma_{\diamond}. \tag{36}
$$

The following is included in [\[12,](#page-43-1) Theorem 10].

Lemma 4.4.
$$
OGW_{1,2} = 2
$$
, $OGW_{1,1}(\Gamma_{n+1}) = 0$, $OGW_{1,0}(\Gamma_n) = (-1)^{\frac{n-1}{2}}$.

Proof of Theorem [3.](#page-2-1) By the degree axiom for GW (8) and \overline{OGW} (17) , all contributions with $d \geq 2$ to the sums in [\(36\)](#page-17-1) are zero. Consider now invariants of the form $\mathrm{OGW}_{d,0}(\Gamma_i, \Gamma_j)$, constituting the coefficients in the second sum in [\(36\)](#page-17-1). For $d = 0$, the zero axiom [\(19\)](#page-10-5) implies that the only nonzero values are when $i = \infty$ and $j = 0$ or vice versa. For $d = 1$, the degree axiom [\(17\)](#page-10-3) implies that nonzero values are only possible when $|\Gamma_i| + |\Gamma_j| = 2n + 2$.

For $i, j \in \{0, \ldots, n\}$, this means there is potentially contribution if $j = n+1-i$, that is, an invariant of the type $\mathrm{OGW}_{1,0}(\Gamma_i,\Gamma_{n+1-i})$. By Lemma [4.4,](#page-17-2) the wall crossing theorem [\(23\)](#page-11-3), zero axiom [\(19\)](#page-10-5), and divisor axiom [\(20\)](#page-10-4), the other possibly nontrivial values of $\text{OGW}_{d,0}(\Gamma_i, \Gamma_j)$ for $d = 0, 1$, are explicitly computed to be

$$
\overline{OGW}_{0,0}(\Gamma_{\diamond}, 1) = 1,
$$

\n
$$
\overline{OGW}_{1,0}(\Gamma_{\diamond}, \Gamma_{\diamond}) = \overline{OGW}_{1,2} = 2,
$$

\n
$$
\overline{OGW}_{1,0}(\Gamma_{\diamond}, \Gamma_{\frac{n+1}{2}}) = -\overline{OGW}_{1,1}(\Gamma_{\frac{n+1}{2}}) = 0,
$$

\n
$$
\overline{OGW}_{1,0}(\Gamma_1, \Gamma_n) = \frac{1}{2}\overline{OGW}_{1,0}(\Gamma_n) = \frac{1}{2} \cdot (-1)^{\frac{n-1}{2}}.
$$

This immediately implies that

$$
\textbf{D}(\Gamma_1,\Gamma_\diamond)=0.
$$

Further, combining the above computation of $\mathrm{OGW}_{d,0}(\Gamma_i, \Gamma_j)$ with the standard computation of the closed small quantum product, we deduce the following relations:

$$
\begin{aligned} &\mathbf{D}(\Gamma_\diamond,1)=\Gamma_\diamond,\\ &\mathbf{D}(\Gamma_\diamond,\Gamma_\diamond)=2T\Gamma_\diamond,\\ &\mathbf{D}(\Gamma_\diamond,\Gamma_{\frac{n+1}{2}})=0,\\ &\mathbf{D}(\Gamma_1,\Gamma_n)=(-1)^{\frac{n+1}{2}}T^2+(-1)^{\frac{n-1}{2}}\frac{1}{2}T\Gamma_\diamond,\\ &\mathbf{D}(\Gamma_i,\Gamma_j)=\Gamma_{i+j},\qquad i,j\in\{0,\ldots,n\},\quad i+j\leq n. \end{aligned}
$$

From the associativity of **p** we see that $\overline{OGW}_{1,0}(\Gamma_i,\Gamma_j) = \overline{OGW}_{1,0}(\Gamma_1,\Gamma_n)$, which does not generate new relations. Taking q, x, y , to be the variables corresponding to $T^2, \Gamma_1, \Gamma_\diamond$, respectively, we get the desired result.

Remark 4.5. As noted above, it follows from the associativity of Δ that

$$
\mathbf{D}(\Gamma_{\frac{n+1}{2}}, \Gamma_{\frac{n+1}{2}}) = \mathbf{D}(\Gamma_1, \Gamma_n) = (-1)^{\frac{n+1}{2}} T^2 + (-1)^{\frac{n-1}{2}} \frac{1}{2} T \Gamma_{\diamond}.
$$

This implies that

$$
\overline{\text{OGW}}_{1,0}(\Gamma_{\frac{n+1}{2}}, \Gamma_{\frac{n+1}{2}}) = (-1)^{\frac{n-1}{2}} \frac{1}{2}.
$$

This corrects a sign error made in Section 1.3.11 of [\[12\]](#page-43-1).

4.4. Full computation of invariants.

Proposition 4.6. All GW invariants are determined by the WDVV equations and the initial value $\text{GW}_1(\Delta_n^{\otimes 2}) = 1$. All OGW invariants are determined by the OWDVV equations, the wall-crossing theorem [\(23\)](#page-11-3), the closed invariants, and the initial value $\text{OGW}_{1,2} = 2$.

Proof. The closed case is given in [\[5,](#page-43-3) Claim 5.22] and [\[8,](#page-43-10) Corollary 1]. The open case is $|12$, Theorem 10.

5. The quadric – preliminaries

5.1. The singular cohomology.

5.1.1. Notation. For the rest of this paper, we compute the relative quantum cohomology and open Gromov-Witten invariants of (X, L) where $X \subset \mathbb{CP}^{n+1}$, $n > 1$, is the quadric hypersurface and L is as follows. Keeping in mind that all quadric hypersurfaces are isomorphic over C, we can take X to be given by $\sum_{j=0}^{n} z_j^2 - z_{n+1}^2 = 0$. We take L to be the fixed locus of complex conjugation $\vec{z} \mapsto \vec{\bar{z}}$. In particular, $L \simeq S^n$. We consider the symplectic form ω given by the restriction of the Fubini-Study form ω_{FS} from $\mathbb{C}P^{n+1}$. Let $r : X \hookrightarrow \mathbb{C}\mathbb{P}^{n+1}$ be the inclusion and let

$$
r^*: H^*(\mathbb{CP}^{n+1}; \mathbb{Z}) \to H^*(X; \mathbb{Z})
$$

be the induced restriction. Let $\check{h} \in H^*(\mathbb{CP}^n;\mathbb{Z})$ denote the Poincaré dual to a hyperplane $\mathcal{H} \subset \mathbb{C}P^n$. In other words, $\check{h} = [\omega_{FS}]$. Let

$$
h=r^*\check{h}.
$$

Let $p \in H^n(X;\mathbb{Z})$ denote the Poincaré dual of L.

Let $m = \lfloor n/2 \rfloor$ and let $m^* = \lfloor n/2 \rfloor$, so $m + m^* = n$. Let $\mathcal{L} \subset X$ be a linear subspace of the maximal possible dimension. For concreteness, choose homogeneous coordinates w_0, \ldots, w_{n+1} , on $\mathbb{C}P^{n+1}$ such that X is given by the equation

$$
w_0 w_1 + \dots + w_{n-1} w_n + w_{n+1}^2 = 0, \qquad \text{when } n \text{ is odd, and}
$$

$$
w_0 w_1 + \dots + w_n w_{n+1} = 0, \qquad \text{when } n \text{ is even.}
$$
 (37)

Take

$$
\mathcal{L} = \{w_{2j} = 0 \, | \, j = 0, \dots, m\} \subset \mathbb{C}P^{n+1}.
$$

Let $l \in H^{2m}(X;\mathbb{Z})$ be the Poincaré dual of \mathcal{L} . When n is odd, we will also use the linear subspace $\mathcal{L}^{\star} \subset X$ of maximal possible dimension given by

$$
\mathcal{L}^* = \{ w_{2j} = 0 \mid j = 0, \dots, m-1 \} \cap \{ w_{n+1} = 0 \} \subset \mathbb{C}P^{n+1},
$$
\n(38)

and the Poincaré dual cohomology class $l^* \in H^{2m}(X;\mathbb{Z})$.

5.1.2. Cohomology ring. It is well known that the singular cohomology of X is described as follows.

Proposition 5.1. Equip the quadric hypersurface X with its complex orientation. The ring structure of $H^*(X;\mathbb{Z})$ is determined as follows. For n odd,

$$
H^*(X; \mathbb{Z}) \simeq \mathbb{Z}[h, l]/(h^m - 2l, l^2).
$$

For n and m even,

$$
H^*(X;\mathbb{Z}) \simeq \mathbb{Z}[h,l]/(h^{m+1} - 2hl, l^2 - h^m l).
$$

For n even and m odd,

$$
H^*(X; \mathbb{Z}) \simeq \mathbb{Z}[h, l]/(h^{m+1} - 2hl, l^2).
$$

For all n, the Poincaré pairing of $H^*(X;\mathbb{Z})$ is determined by $\int_X h^{m^*} l = 1$.

Proposition 5.2. As a group, the cohomology of X is given by

$$
H^*(X;\mathbb{Z}) = \bigoplus_{j=0}^{2n} H^j(X;\mathbb{Z}), \qquad H^j(X;\mathbb{Z}) \simeq \begin{cases} 0, & j \neq n, \ j \text{ is odd}, \\ \mathbb{Z}, & j \neq n, \ j \text{ is even}, \\ 0, & j = n, \ n \text{ is odd}, \\ \mathbb{Z} \oplus \mathbb{Z}, & j = n, \ n \text{ is even}. \end{cases}
$$

It is generated by $\{h^{j}\}_{j=0}^{m^{*}} \cup \{h^{j}l\}_{j=0}^{m^{*}}$.

Lemma 5.3. We have

(a) $\int_X h^n = 2$.

(b)
$$
\int_X h^{m^*} l = 1.
$$

(c) When n or m is odd, $l^2 = 0$. When n and m are even, $\int_X l^2 = 1$.

Proof. (a) By Bezout's theorem $\int_X h^n = \mathcal{H}^n \cdot X = 2$.

(b) Since a codimension m^* linear subspace of $\mathcal L$ is a single point, we have

$$
\int_X h^{m^*} l = \mathcal{H}^{m^*} \cdot \mathcal{L} = 1.
$$

(c) First, consider the case that n is odd, so $m > n/2$. Consequently,

$$
|l^2| = 4m > 2n = \dim_{\mathbb{R}} X
$$

and $l^2=0$.

Now consider the case *n* is even, so $m = n/2$. Choose homogeneous coordinates w_0, \ldots, w_{n+1} , on $\mathbb{C}P^{n+1}$ such that X is given by equation [\(37\)](#page-19-0) and let $\mathcal L$ be as in [\(5.1.1\)](#page-19-0). Consider the maximal linear subspace $\mathcal{L}' \subset X$ given by

$$
\mathcal{L}' = \{w_{2j+1} = 0 \mid j = 0, ..., m\} \subset \mathbb{C}P^{n+1}, \text{ when } m \text{ is odd, and}
$$

$$
\mathcal{L}' = \{w_n = 0, w_{2j+1} = 0 \mid j = 0, ..., m-1\} \subset \mathbb{C}P^{n+1}, \text{ when } m \text{ is even.}
$$

We claim that $\mathcal L$ and $\mathcal L'$ are homologous in X. Indeed, we can define a family of maximal linear subspaces $\mathcal{L}_t \subset X$, $t \in [0, 1]$, by

$$
\mathcal{L}_t = \{tw_3 - (1-t)w_0 = 0 = (1-t)w_2 + tw_1, w_{2j} = 0 | j = 2,...,m\}.
$$

Then $\mathcal{L}_0 = \mathcal{L}$ and

$$
\mathcal{L}_1 = \{w_1 = w_3 = w_4 = \cdots = w_n = 0\} \subset \mathbb{C}P^{n+1}.
$$

Proceeding similarly with w_k, \ldots, w_{k+3} instead of w_0, \ldots, w_3 , for $k = 4, 8, \ldots$ successively, we conclude that $\mathcal L$ and $\mathcal L'$ are homologous as desired. Observe that $\mathcal L \cap \mathcal L' = \emptyset$ when m is odd, but $\mathcal L$ intersects $\mathcal L'$ transversally at a single point when m is even. Thus,

$$
\mathcal{L} \cdot \mathcal{L} = \mathcal{L} \cdot \mathcal{L}' = \begin{cases} 0, & m \text{ odd,} \\ 1, & m \text{ even.} \end{cases}
$$

The desired formula for l^2 follows.

 \Box

5.1.3. Involution and maximal linear subspaces. Let $\phi: X \to X$ be the involution given by

$$
\phi([z_0 : \dots : z_{n+1}]) = [z_0 : \dots : -z_{n+1}]. \tag{39}
$$

Recall the definition of \mathcal{L}^* and l^* from Section [5.1.1.](#page-18-1)

Lemma 5.4. For n even, we have $\phi^*l = l^*$.

Proof. In the coordinates w_i of equation [\(37\)](#page-19-0), we have

$$
\phi([w_0:\cdots:w_n:w_{n+1}])=[w_0:\cdots:w_{n+1}:w_n].
$$

So, $\phi(\mathcal{L}) = \mathcal{L}^*$

Lemma 5.5. For n even, we have $h^m = l + l^* \in H^{2m}(X)$.

Proof. The class h^m is Poincaré dual to the intersection

$$
\{w_0 = 0\} \cap \{w_2 = 0\} \cap \cdots \cap \{w_{n-2} = 0\} \cap X = \mathcal{L} + \mathcal{L}^*.
$$

.
1990 - Johann Stein, Amerikaansk politiker († 1905)

 \Box

5.1.4. The fundamental class of the Lagrangian.

Lemma 5.6. Let W be a symplectic manifold of dimension 4w, let $K \subset W$ be a Lagrangian homology sphere, and let $k \in H^{2w}(W)$ be the Poincaré dual of K. Then $\int_W k^2 = (-1)^w 2$.

Proof. Fix an orientation on K and thus on $TK \simeq T^*K$. Equip TW with the canonical symplectic orientation. Let N_K denote the normal bundle of $K \subset W$. The short exact sequence

$$
0\to TK\to TW\to N_K\to 0
$$

induces an orientation on N_K . Explicitly, choose an oriented basis $u_1, \ldots, u_{2w} \in T_xK$ and choose $v_1, \ldots, v_{2w} \in T_xW$ such that $u_1, \ldots, u_{2w}, v_1, \ldots, v_{2w}$ form a symplectic basis. That is,

$$
\omega_p(u_i, u_j) = 0 = \omega(v_i, v_j), \qquad \omega(u_i, v_j) = \delta_{ij}.
$$

Then $u_1 \wedge v_1 \wedge \ldots \wedge u_{2w} \wedge v_{2w}$ is the symplectic orientation of T_xW . Let $\bar{v}_1, \ldots, \bar{v}_{2w}$, be the images of v_1, \ldots, v_{2w} , in $(N_K)_x$. Then $(-1)^w\overline{v}_1 \wedge \ldots \wedge \overline{v}_{2w}$ is the induced orientation of $(N_K)_x$. Indeed,

$$
u_1 \wedge v_1 \wedge \ldots \wedge u_{2w} \wedge v_{2w} = (-1)^{\frac{2w(2w-1)}{2}} u_1 \wedge \ldots \wedge u_{2w} \wedge v_1 \wedge \ldots \wedge v_{2w},
$$

$$
v_{2w-1}
$$

and $(-1)^{\frac{2w(2w-1)}{2}} = (-1)^w$.

The symplectic form determines a bundle isomorphism $N_K \to T^*K$. Explicitly, with u_1^*,\ldots,u_{2w}^* , the dual basis of u_1,\ldots,u_{2w} , the isomorphism is given by $\bar{v}_i\mapsto u_i^*$. In particular, the isomorphism has sign $(-1)^w$. Therefore,

$$
\int_{W} k^{2} = [K] \cdot [K] = \int_{K} e(N_{K}) =
$$
\n
$$
= (-1)^{w} \int_{K} e(T^{*}K) = (-1)^{w} \chi(K) = (-1)^{w} \chi(S^{2w}) = (-1)^{w} 2.
$$

Let $p \in H^n(X;\mathbb{Z})$ denote the Poincaré dual of L.

Lemma 5.7. When n is odd, $p = 0$. When n is even, $p = \pm(h^m - 2l) = \mp(l - l^*)$ with the sign depending on the orientation of L.

Proof. When *n* is odd, Proposition [5.2](#page-19-1) gives $Hⁿ(X) = 0$, so $p = 0$. When *n* is even, since L is Lagrangian, we know that

$$
\int_{X} p \cup h^{m} = \int_{X} p \cup [\omega]^{m} = \int_{L} \omega^{m} = 0.
$$
\n(40)

By Lemma [5.6,](#page-21-0) we have

$$
\int_{X} p^{2} = (-1)^{n/2} 2.
$$
\n(41)

By Proposition [5.2,](#page-19-1) $rk H^{n}(X;\mathbb{Z}) = 2$. So, equations [\(40\)](#page-21-1) and [\(41\)](#page-21-2) suffice to determine p up to sign. Using Lemma [5.3,](#page-19-2) one verifies these equations for $p = \pm (h^m - 2l)$. The equality $\pm(h^m - 2l) = \pm(l - l^*)$ follows from Lemma [5.5.](#page-20-0) **Proposition 5.8.** The de Rham cohomology of X is given by

$$
H^*(X) \simeq \mathbb{R}[h] \big/ (h^{n+1})
$$

if n is odd and

$$
H^*(X) \simeq \mathbb{R}[h, p] / (h^{n+1}, \quad hp, \quad p^2 - (-1)^{n/2} h^n)
$$

if n is even. The Poincaré pairing is determined by $\int_X h^n = 2$.

Proof. This follows from Proposition [5.1](#page-19-3) and Lemma [5.7.](#page-21-3) \Box

5.1.5. Relative cohomology ring. Let $\hat{A}^*(L)$ be the reduced de Rham complex on L and let $\hat{H}^*(L)$ the associated reduced cohomology; specifically, for $j > 0$ we take $\hat{A}^j(L) = A^j(L)$, and for $j = 0$ we take $A^{j}(L) = A^{0}(L)/\mathbb{R}$, where \mathbb{R} is thought of as the subspace of constant zero-forms. Define the associated relative complex $\hat{A}^*(X, L)$ by the short exact sequence

$$
0 \to \widetilde{A}^*(X, L) \longrightarrow A^*(X) \longrightarrow \widetilde{A}^*(L) \to 0.
$$

Its cohomology $\overline{H}^*(X, L)$ then fits in the long exact sequence

$$
\widetilde{H}^*(X, L) \xrightarrow{a} H^*(X) \tag{42}
$$
\n
$$
\widetilde{H}^*(L). \tag{42}
$$

Let $\tilde{u} \in \tilde{H}^{n+1}(X,L)$ be the image under c of the class with integral 1 in $\tilde{H}^{n}(L)$. Let $\tilde{h} =$ $[r^*\omega_{FS}] \in \tilde{H}^2(X,L).$

Proposition 5.9. As \mathbb{R} algebras, when n is odd

$$
\widetilde{H}^*(X,L) \simeq \mathbb{R}[\tilde{h},\tilde{u}]/(\tilde{h}^{n+1},\tilde{u}^2,\tilde{u}\tilde{h}).
$$

When n is even,

$$
\widetilde{H}^*(X,L) \simeq \mathbb{R}[\widetilde{h}]/(\widetilde{h}^{n+1}).
$$

Proof. By definition, we have $a(\tilde{h}) = h$. Since $L \simeq S^n$, it follows that $\widetilde{H}^*(L)$ is one dimensional and concentrated in degree n. For the case n odd, observe that $b = 0$ for degree reasons, so c is injective and a is an isomorphism except in degree $n + 1$. By exactness, $\tilde{u} \in \text{Im}(c)$ implies $a(\tilde{u}) = 0$. Since $a(\tilde{u}\tilde{h}) = a(\tilde{u})a(\tilde{h}) = 0$, we deduce that $\tilde{u}\tilde{h} = 0$. Also, $\tilde{u}^2 = 0$ for degree reasons. The claim follows from Proposition [5.8.](#page-22-0)

For the case *n* even, the integral of $b(p) \in H^*(L)$ is $(-1)^{n/2}2$ by Lemma [5.6.](#page-21-0) In particular, $b(p) \neq 0$, so b is surjective and $c = 0$. So, the long exact sequence [\(42\)](#page-22-1) implies that $\overline{H}^*(X, L)$ is the kernel of b, which is the subalgebra of $\widetilde{H}^*(X)$ generated by \widetilde{h} . So, the claim follows by Proposition [5.8.](#page-22-0)

Recalling the long exact sequence [\(4\)](#page-6-3), let $\hat{u} = y(1) \in \hat{H}^{n+1}(X)$. Let $\hat{h} = [r^*\omega_{FS}] \in$ $\hat{H}^2(X,L).$

Proposition 5.10. As \mathbb{R} -vector spaces, when n is odd,

$$
\widehat{H}^*(X,L) \simeq \mathbb{R}[\hat{h},\hat{u}]/(\hat{h}^{n+1},\hat{u}^2,\hat{u}\hat{h}).
$$

When n is even,

$$
\widehat{H}^*(X,L) \simeq \mathbb{R}[\widehat{h}]\big/(\widehat{h}^{n+1}).
$$

Proof. Since $L = Sⁿ$, Lemma 5.14 of [\[13\]](#page-43-8) asserts that the inclusion of underlying complexes induces an isomorphism $\widetilde{H}^*(X,L) \stackrel{\sim}{\to} \widehat{H}^*(X,L)$. So, the claim follows from Proposition [5.9.](#page-22-2) П

Remark 5.11. Proposition [5.10](#page-23-0) gives a natural splitting of the exact sequence [\(6\)](#page-6-4). Indeed, a section of ρ is given by mapping $h^j \in H^*(X)$ to $\hat{h}^j \in \hat{H}^*(X)$ for $j = 0, \ldots, n$.

5.2. Additional topological properties.

Lemma 5.12. We have $H_2(X, L; \mathbb{Z}) \simeq \mathbb{Z}$. Moreover, the map $\varpi : H_2(X; \mathbb{Z}) \to H_2(X, L; \mathbb{Z})$ from the long exact sequence of the pair (X, L) is an isomorphism if $n > 2$ and a surjection if $n=2$.

Identify $H_2(X, L; \mathbb{Z})$ with $\mathbb Z$ so that the non-negative integers correspond to classes $\beta \in$ $H_2(X, L; \mathbb{Z})$ with $\omega(\beta) \geq 0$. When $n > 2$, we also identify the non-negative integers with classes $\beta \in H_2(X;\mathbb{Z})$ with $\omega(\beta) > 0$. Then $\varpi = \mathrm{Id}_{\mathbb{Z}}$. When $n = 2$, Lemma [5.7](#page-21-3) implies that the inclusion $H_2(L; \mathbb{Z}) \hookrightarrow H_2(X; \mathbb{Z})$ is given by $a \mapsto \pm(a, -2a)$. Exactness then gives $\varpi(a, b) = 2a + b.$

Remark 5.13. Using the identifications above, we can write explicitly the integrals of divisors, analogously to equation [\(34\)](#page-16-7). Namely, by Proposition [5.10,](#page-23-0) for any $\beta \in H_2(X, L; \mathbb{Z})$ we have

$$
\int_{\beta} \hat{h} = \beta.
$$

By Proposition [5.2,](#page-19-1) when $n > 2$ we have

$$
\int_{\hat{\beta}} h = \hat{\beta}, \qquad \forall \hat{\beta} \in H_2(X; \mathbb{Z}),
$$

and when $n = 2$ we have

$$
\int_{(\beta_1,\beta_2)} h = \beta_1, \quad \int_{(\beta_1,\beta_2)} l = \beta_2, \qquad \forall (\beta_1,\beta_2) \in H_2(X;\mathbb{Z}).
$$

As in Section [4,](#page-16-0) these values will be used when applying the divisor axioms [\(20\)](#page-10-4) and [\(11\)](#page-8-0).

Lemma 5.14. The relevant characteristic classes of (X, L) are as follows:

(a) For all $\hat{\beta} \in H_2(X;\mathbb{Z})$ we have $c_1(\hat{\beta}) = n\hat{\beta}$. (b) For all $\beta \in H_2(X, L; \mathbb{Z})$, we have $\mu(\beta) = 2n\beta$. (c) $w_{\rm s} = 0$.

5.3. Basic values for GW and OGW invariants. Here we compute the basic values required to deduce the quantum ring structure on the absolute and relative cohomologies. In this section we assume $n > 3$. For the case $n = 2$ see Section [8.](#page-34-0) In particular, we identify $H_2(X;\mathbb{Z})$ and $H_2(X, L;\mathbb{Z})$ with \mathbb{Z} , so that positive-energy classes correspond to $\mathbb{Z}_{>0}$.

For $\beta = 0$, the invariant $\text{GW}_{\beta}(h^i, h^j, h^m)$ does not vanish if and only if $i + j + m = n$, in which case by Lemma [5.3](#page-19-2)[\(a\)](#page-19-4) we have $GW_0(h^i, h^j, h^m) = 2$.

Lemma 5.15. $\text{GW}_1(h^{m^*}l, h^{m^*-1}l) = 1$.

Proof. By Lemma [5.3](#page-19-2)[\(b\),](#page-19-5) the question translates to the number of lines in X though one point and one line. Any point $x \in X$ and line $\ell \subset X$ determine a plane in \mathbb{CP}^{n+1} . This plane intersects the hypersurface X in a degree 2 curve C . On the other hand, C contains the line ℓ . Therefore, C is the union of two lines, $\ell \cup \ell'$. In other words, there is a unique line ℓ' in X passing through x that intersects ℓ . Thus, $GW_1(h^{m^*}l, h^{m^*-1}l) = 1$.

Corollary 5.16. $\text{GW}_1(h^n, h^{n-1}) = 2^2 \cdot \text{GW}_1(h^{m^{*}l}, h^{m^{*}-1}l) = 4.$

Lemma 5.17. For i, j, k, such that $i + j + k = 2n$, $i \leq j \leq k$, we have

$$
GW_1(h^i, h^j, h^k) = \begin{cases} 0, & (i, j, k) = (0, n, n), \\ 4, & i > 0, k = n, \\ 8, & i > 0, k < n. \end{cases}
$$

Note that the condition $i + j + k = 2n$ is required by the degree axiom [\(8\)](#page-8-1).

Proof. For $i = 0$, we apply the unit axiom [\(9\)](#page-8-2) to get the first case:

$$
\mathrm{GW}_1(h^0, h^n, h^n) = 0.
$$

For $i = 1$, we apply the divisor axiom [\(11\)](#page-8-0) and Corollary [5.16](#page-24-0) to get

$$
GW1(h, hn-1, hn) = GW1(hn-1, hn) = 4.
$$

Now assume $i > 1$. Take the coefficient of T^1 in the equation [\(12\)](#page-9-1) with $u = h^{i-1}, v =$ $h, w = h^j$, and $y = h^k$, to get

$$
GW_1(h^i, h^j, h^k) + GW_1(h^{i-1}, h^{j+k}) = GW_1(h^{i-1}, h^{j+1}, h^k) + GW_1(h^j, h^{i+k-1}).
$$

The summand $GW_1(h^{i-1}, h^{j+k})$ contributes nothing: By the degree axiom [\(8\)](#page-8-1), it can only be nonzero if $j + k = n$ and $i - 1 = n - 1$, but in this case $i = n > j$, contradicting our assumption.

The summand $GW_1(h^j, h^{i+k-1})$ satisfies

GW₁
$$
(h^j, h^{i+k-1}) = \begin{cases} 4, & (i, j, k) = (2, n - 1, n - 1), \\ 0, & otherwise, \end{cases}
$$

for the following reason. By the degree axiom [\(8\)](#page-8-1), $GW_1(h^j, h^{i+k-1})$ can only be nonzero if $\{j, i + k - 1\} = \{n - 1, n\}$. Since we assumed $i > 1$, we have $i + k - 1 > k \geq j$, so $(j, i + k - 1) = (n - 1, n)$. In this case $k = n + 1 - i$. Since $i > 1$ we get $k < n$, and since $k \ge j = n - 1$, we get $k = n - 1$ and $i = 2$.

Thus, if $(i, j, k) = (2, n - 1, n - 1)$, we have

$$
GW_1(h^i, h^j, h^k) = GW_1(h^2, h^{n-1}, h^{n-1}) = GW_1(h, h^n, h^{n-1}) + GW_1(h^{n-1}, h^n) = 8,
$$

₂₅

and if $(i, j, k) \neq (2, n - 1, n - 1)$ with $i > 1$, we have

$$
GW1(hi, hj, hk) = GW1(hi-1, hj+1, hk).
$$
\n(43)

Thus, if $k = n$ then $i + j = n$, and inductively,

$$
GW_1(h^i, h^{n-i}, h^n) = GW_1(h^{i-(i-1)}, h^{(n-i)+(i-1)}, h^n) = GW_1(h, h^{n-1}, h^n) = 4.
$$

If $k < n$, we would again like to apply equation [\(43\)](#page-25-1) repeatedly. In order to do so, consider the transformation

$$
\psi:\mathbb{Z}^{\oplus 3}\longrightarrow\mathbb{Z}^{\oplus 3}
$$

defined by

$$
\psi(a, b, c) := \begin{cases} (a - 1, b + 1, c), & b + 1 \leq c, \\ (a - 1, c, b + 1), & b + 1 > c. \end{cases}
$$

Note that ψ preserves the sum of the components, and takes non-decreasing triples to nondecreasing ones. Since $k < n$, there exists $q \in \mathbb{Z}_{\geq 0}$ such that $\psi^q(i,j,k)$ has the form $\psi^q(i,j,k) = (\nu, \mu, n-1)$. Then $\mu = n+1-\nu, \mu \leq n-1$, and consequently $\nu \geq 2$. Taking $q' := \nu - 2$, we get

$$
\psi^{q+q'}(i,j,m) = \psi^{q'}(\nu,\mu,n-1) = (\nu - (\nu - 2), (n+1-\nu) + (\nu - 2), n-1) = (2, n-1, n-1).
$$

Thus,

$$
GW_1(h^i, h^j, h^k) = GW_1(h^2, h^{n-1}, h^{n-1}) = 8.
$$

Lemma 5.18. OG $W_{1,1}(h^n) = 2$.

Proof. Apply ∂_s to equation [\(22\)](#page-11-2) with $v = h$ and $w = h^{n-1}$, set $t_j = s = 0$ and take the coefficient of T^1 . This gives

$$
\frac{1}{2} GW_0(h, h^{n-1}, 1) OGW_{1,1}(h^n) + \frac{1}{2} GW_1(h, h^{n-1}, h^n) OGW_{0,1}(1) = 0.
$$

Apply the zero axiom [\(10\)](#page-8-3) and Lemma [5.3](#page-19-2)[\(a\)](#page-19-4) to the first summand, and apply the divisor [\(11\)](#page-8-0) and zero [\(19\)](#page-10-5) axioms and Corollary [5.16](#page-24-0) to the second summand, to get

$$
OGW_{1,1}(h^n) - 2 = 0.
$$

6. The quadric – odd dimension

6.1. Absolute quantum ring. In this section, we assume $n > 1$ is odd. We work with the basis

$$
\Delta_j := h^j, \qquad j = 0, \dots, n,
$$

of $H^*(X;\mathbb{R})$. By Lemma [5.3](#page-19-2)[\(a\),](#page-19-4) the Poincaré pairing on $H^*(X;\mathbb{R})$ is given by

$$
g_{ij} = \int_X \Delta_i \wedge \Delta_j = \delta_{i,n-j} \cdot 2,
$$

26

where $\delta_{k,l}$ is the Kronecker delta, and the inverse matrix $(g^{ij})_{i,j} := (g_{ij})_{i,j}^{-1}$ is given by $g^{ij}=\delta_{i,n-j}\cdot\frac{1}{2}$ $\frac{1}{2}$. Thus, the quantum product is given by

$$
\Delta_i * \Delta_j = \sum_{\substack{0 \le m \le n \\ d \ge 0}} \frac{1}{2} T^d \, \mathrm{GW}_d(\Delta_i, \Delta_j, \Delta_m) \Delta_{n-m}, \qquad i, j \in \{0, \ldots, n\}.
$$

Proposition 6.1. The absolute quantum cohomology is given by

$$
QH^*(X) \simeq \mathbb{R}[[q]][x]/(x^{n+1} - 4qx).
$$

Here, as in the case of \mathbb{CP}^n , the variable q corresponds to $T^{\varpi(\hat{\beta}_1)}$ where $\hat{\beta}_1$ is the class of a line generating $H_2(X;\mathbb{Z})$, and x corresponds to a divisor generating $H^2(X)$. Note that for the quadric, $T^{\varpi(\hat{\beta}_1)} = T^1 = T$.

Proof. If $i + j < n$, use the degree axiom [\(8\)](#page-8-1) and the zero axiom [\(10\)](#page-8-3) to get

$$
\Delta_i * \Delta_j = \sum_{d,m} \frac{1}{2} T^d \, \text{GW}_d(\Delta_i, \Delta_j, \Delta_m) \Delta_{n-m}
$$

$$
= \sum_m \frac{1}{2} \cdot \text{GW}_0(\Delta_i, \Delta_j, \Delta_m) \Delta_{n-m}
$$

$$
= \frac{1}{2} \cdot \left(\int_X \Delta_i \wedge \Delta_j \wedge \Delta_{n-i-j} \right) \cdot \Delta_{i+j}
$$

$$
= \Delta_{i+j}.
$$

By the associativity of \ast , for $k < n$ we therefore have

$$
\Delta_1^{*k} = \Delta_k.
$$

Next, we compute $\Delta_1^{*n} = \Delta_1^{*(n-1)} * \Delta_1$, as follows. The degree axiom [\(8\)](#page-8-1) implies that $GW_d(\Delta_{n-1}, \Delta_1, \Delta_m)$ can only have nonzero values for $d = 0, 1$. By the zero axiom [\(10\)](#page-8-3), the case $d = 0$ contributes $GW_0(\Delta_{n-1}, \Delta_1, \Delta_0) = 2$, as before. For $d = 1$, again by the degree axiom [\(8\)](#page-8-1), the only nonzero case is when $m = n$. By the divisor axiom [\(11\)](#page-8-0), the value is

$$
GW_1(\Delta_{n-1}, \Delta_1, \Delta_n) = \int_{\beta_1} \Delta_1 \cdot GW_1(\Delta_{n-1}, \Delta_n) = GW_1(\Delta_{n-1}, \Delta_n) = 4.
$$

So,

$$
\Delta_1^{*n} = \frac{1}{2} \operatorname{GW}_0(\Delta_{n-1}, \Delta_1, \Delta_0) \Delta_n + \frac{1}{2} T \operatorname{GW}_1(\Delta_{n-1}, \Delta_1, \Delta_n) \Delta_0 = \Delta_n + 2T.
$$

Thus, we see that $\{\Delta_1^{*j}\}$ ^{*j} $\prod_{i=0}^{j}$ generate the Q_X -module $QH^*(X)$. To complete the computation of the ring structure, we need to compute powers of Δ_1 higher than n. By the degree axiom [\(8\)](#page-8-1) and the zero axiom [\(10\)](#page-8-3), we have

$$
\Delta_n * \Delta_1 = \frac{1}{2} T \operatorname{GW}_1(\Delta_n, \Delta_1, \Delta_{n-1}) \Delta_1 = \frac{1}{2} T \cdot 4 \Delta_1 = 2T \Delta_1.
$$

Therefore,

$$
\Delta_1^{*(n+1)} = \Delta_1^{*n} * \Delta_1 = (\Delta_n + 2T) * \Delta_1 = 2T\Delta_1 + 2T\Delta_1 = 4T\Delta_1.
$$

Taking the variables q, x, to correspond to T, Δ_1 , respectively, we get the desired result. \Box

6.2. Relative quantum ring. In this section, we work with the following basis. Let $\Gamma_1 =$ $[r^*\omega_{FS}] \in \widehat{H}^*(X, L; \mathbb{R})$ and set $\Gamma_j = \Gamma_1^{\hat{\wedge}j}$ for $j = 0, \ldots, n$. Let $\Gamma_{\diamond} = y(1)$ where y is the map from [\(4\)](#page-6-3). Then $\{\Gamma_j\}_{j=0}^n\cup \{\Gamma_{\diamond}\}\$ is a basis of $\hat{H}^*(X, L; \mathbb{R})$, and for Δ_j of Section [6.1](#page-25-2) we have

$$
\rho(\Gamma_j) = \Delta_j, \qquad j = 0, \dots, n.
$$

For convenience, we also define $\Delta_{\diamond} := 0 \in H^*(X; \mathbb{R})$.

For $i, j \in \{0, \ldots, n\} \cup \{\infty\}$, we have

$$
\mathbf{D}(\Gamma_i, \Gamma_j) = \sum_{\substack{0 \le m \le n \\ d \ge 0}} \frac{1}{2} T^d \, \mathrm{GW}_d(\Delta_i, \Delta_j, \Delta_m) \Gamma_{n-m} + \sum_{d \ge 0} T^d \, \overline{\mathrm{OGW}}_{d,0}(\Gamma_i, \Gamma_j) \Gamma_{\diamond}.
$$

Proof of Theorem [4.](#page-3-0) By the degree axiom [\(17\)](#page-10-3) of OGW, the only nonzero values possible for $\mathrm{OGW}_{d,0}(\Gamma_i, \Gamma_j)$ are when $d = 0, 1$. When $d = 0$, by the zero axiom [\(19\)](#page-10-5) of OGW and the wall-crossing theorem [\(23\)](#page-11-3), the only nontrivial value is

$$
\mathrm{OGW}_{0,0}(\Gamma_\diamond,\Gamma_0)=-\mathrm{OGW}_{0,1}(\Gamma_0)=1.
$$

When $d = 1$, by the degree axiom [\(17\)](#page-10-3) nonzero values are only possible when $|\Gamma_i| + |\Gamma_j|$ $3n + 1$. If, without loss of generality, $j = \diamond$, we use the wall-crossing theorem [\(23\)](#page-11-3) and Lemma [5.18](#page-25-3) to compute

$$
OGW_{1,0}(\Gamma_n, \Gamma_\diamond) = -OGW_{1,1}(\Gamma_n) = -2.
$$

If $i, j \in \{0, \ldots, n\}$, take the coefficient of T in the first OWDVV equation with $u = t_i, w = t_1$, and $v = t_{j-1}$:

$$
OGW_{1,0}(\Gamma_i, \Gamma_j) = OGW_{1,0}(\Gamma_{i+1}, \Gamma_{j-1}).
$$

In the case $i = n$ we interpret the right-hand side by setting $\Gamma_{n+1} := 0$, and the identity still holds. Note that $2 \cdot ((i+1) + (j-1)) = 2 \cdot (i+j) = 3n+1$. By induction, we increase the degree of the first constraint i (while decreasing the second) until it reaches n and conclude $\mathrm{OGW}_{1,0}(\Gamma_i, \Gamma_j) = 0.$

Since ρ is an isomorphism when restricted to $W = Span(\Gamma_0, \ldots, \Gamma_n)$, we can summarize

$$
\mathbf{D}(\Gamma_i, \Gamma_j) = \begin{cases} \Gamma_{\diamond}, & \{i, j\} = \{\diamond, 0\}, \\ -2T\Gamma_{\diamond}, & \{i, j\} = \{\diamond, n\}, \\ (\rho|_W)^{-1}(\Delta_i * \Delta_j), & otherwise. \end{cases}
$$

Taking q, x, y, to be the variables corresponding to $T, \Gamma_1, \Gamma_\diamond$, respectively, we get the desired result.

6.3. Full computation of invariants. In this section we prove Theorem [7.](#page-4-1)

In Appendix [A,](#page-36-0) we compute the following value.

Lemma 6.2. The unique spin structure on L is such that $\text{OGW}_{1,3} = (-1)^{\frac{(n-2)(n-3)}{2}}2$.

Let $n = 2r + 1$. From Lemma [5.17](#page-24-1) we have another initial value:

Lemma 6.3. GW₁($\Delta_{r+1}, \Delta_{r+1}, \Delta_{2r}$) = 8.

We are now ready to prove computability lemmas.

Lemma 6.4. All open Gromov-Witten invariants $\text{OGW}_{\beta,k}(\gamma_j)$ with $j \neq 1, \diamond$ can be computed from closed Gromov-Witten invariants and open Gromov-Witten invariants of strictly lower degree.

Proof. Consider the invariant $\text{OGW}_{\beta,k}(\Gamma_i)$. If $j = 0$, the invariant vanishes unless $\beta = 0$, in which case the value is determined by the OGW axioms. Thus we can assume $j \geq 2$. Note that for degree reasons we must have $k \geq 1$. We can then consider the open WDVV equation [\(22\)](#page-11-2) with $v = 1$, $w = j - 1$ in degree β after applying ∂_s^{k-1} :

$$
GW_0(\Delta_1, \Delta_{j-1}, \Delta_{n-j})\n\text{OGW}_{\beta,k}(\Gamma_j)g^{n-j,j}\n+ \sum_{\substack{\beta_1+\beta_2=\beta\\ \beta_2<\beta\\i=0,\dots,n}}\n\text{GW}_{\beta_1}(\Delta_1, \Delta_{j-1}, \Delta_i)\n\text{OGW}_{\beta_2,k}(\Gamma_{n-i})g^{n-i,i}
$$
\n
$$
(44)
$$

$$
-\sum_{\substack{\beta_1+\beta_2=\beta\\ \beta_1,\beta_2<\beta\\i=1,\dots,k-1}} {k-1 \choose i} \text{OGW}_{\beta_1,1+k-i} \text{OGW}_{\beta_2,i}(\Gamma_1,\Gamma_{j-1})\tag{45}
$$

$$
= -\sum_{\substack{\beta_1+\beta_2=\beta\\ \beta_1,\beta_2<\beta\\i=0,\dots,k-1}} {k-1 \choose i} \text{OGW}_{\beta_1,1+i}(\Gamma_1) \text{OGW}_{\beta_2,k-i}(\Gamma_{j-1})
$$
(46)

Here in line [\(44\)](#page-28-0), we have $\beta_2 < \beta$, as the only (possibly) non-zero term with $\beta_2 = \beta$ involves the term we are trying to compute. In line [\(45\)](#page-28-1) both $\beta_1, \beta_2 < \beta$, as there are no nontrivial OGW_{0,k} with either 0 or 2 interior constraints, which follows from the choice of $P_{\mathbb{R}}$ and the degree-zero property. Finally in line [\(46\)](#page-28-2), we have $\beta_1, \beta_2 < \beta$ as the the invariant with $\beta_2 = 0$ that could appear $\text{OGW}_{0,k-i}(\Gamma_{j-1})$ vanishes by the energy-zero property, as $j > 1$. As $GW_0(\Delta_1, \Delta_{i-1}, \Delta_{n-j}) = 2$, we have expressed $\text{OGW}_{\beta,k}(\Gamma_i)$ in terms of closed Gromov-Witten invariants and open Gromov-Witten invariants of strictly lower degree. \Box

Lemma 6.5. An open Gromov-Witten invariant $\text{OGW}_{\beta,k}(\Gamma_{j_1}, \ldots, \Gamma_{j_l})$ with $l \geq 2$ can be computed from closed Gromov-Witten invariants, open Gromov-Witten invariants of smaller degree, and invariants of the form $\text{OGW}_{\beta,k'}(\Gamma_{j_1-1},\Gamma_J)$ for $k' \leq k$ and some index set J.

Proof. By reordering the inputs, we may assume $j_1 \leq j_2 \leq \cdots \leq j_l$. Consider the open WDVV equation [\(21\)](#page-10-1) with $u = j_2, v = j_1 - 1, w = 1$, and take the coefficient of $T^{\beta}t_{j_3} \dots t_{j_l}$ after applying ∂_s^k . Set $I = (j_3, \ldots, j_l)$. This yields:

$$
\mathrm{GW}_0(\Gamma_1, \Gamma_{j_1-1}, \Gamma_{n-j_1}) \mathrm{OGW}_{\beta,k}(\Gamma_{j_1}, \dots, \Gamma_{j_l}) g^{n-j_1, j_1} \tag{47}
$$

$$
+\sum_{\substack{\beta_1+\beta_2=\beta\\ \beta_2\leq \beta}} \text{GW}_{\beta_1}(\Gamma_1, \Gamma_{j_1-1}, \Gamma_{I_1}, \Gamma_i) \text{OGW}_{\beta_2, k}(\Gamma_{n-i}, \Gamma_{j_2}, \Gamma_{I_2}) g^{n-i,i}
$$
(48)

$$
-\sum_{\substack{j=0,\ldots,n\\ \beta_1+\beta_2=\beta\\ \beta_1,\beta_2<\beta}}^{I_1\sqcup I_2=I} {k \choose i} \text{OGW}_{\beta_1,1+i}(\Gamma_{j_2},\Gamma_{I_1}) \text{OGW}_{\beta_2,k-i}(\Gamma_{j_1-1},\Gamma_1,\Gamma_{I_2})
$$
(49)

$$
I_{1 \sqcup I_{2}=I}^{I_{1} \sqcup I_{2}=I}
$$

= GW₀($\Gamma_{j_2}, \Gamma_1, \Gamma_{n-j_2-1}$) OGW _{β, k} ($\Gamma_{j_2+1}, \Gamma_{j_1-1}, \Gamma_I$) g^{n-j_2-1, j_2+1} (50)
29

+
$$
\sum_{\substack{\beta_1+\beta_2=\beta\\ \beta_2\leq \beta_{-}}} GW_{\beta_1}(\Gamma_{j_2}, \Gamma_1, \Gamma_{I_1}, \Gamma_i) \text{OGW}_{\beta_2, k}(\Gamma_{n-i}, \Gamma_{j_1-1}, \Gamma_{I_2}) g^{n-i,i}
$$
(51)

$$
I_{i \sqcup I_2=I}^{I_1 \sqcup I_2=I} \n-\sum_{\substack{\beta_1+\beta_2=\beta \\ \beta_1,\beta_2<\beta \\ I_1 \sqcup I_2=I \\ i=0,\dots,k}} \binom{k}{i} \text{OGW}_{\beta_1,1+i}(\Gamma_{j_1-1}\Gamma_{I_1}) \text{OGW}_{\beta_2,k-i}(\Gamma_{j_2},\Gamma_1,\Gamma_{I_2}) \n\tag{52}
$$

Here in line [\(48\)](#page-28-3), we have $\beta_2 < \beta$, as the only (possibly) non-zero term with $\beta_2 = \beta$ involves the term we are trying to compute, line [\(47\)](#page-28-4). For lines [\(49\)](#page-28-5) and [\(52\)](#page-29-0), we need $\beta_1, \beta_2 > 0$, as the OGW have at least one interior constraint on Γ_{j_1-1} or Γ_{j_2} , and by assumption $j_1, j_2 > 1$, so that by the degree zero axiom, there are no contributions from these terms in degree 0. This implies that $\beta_1, \beta_2 < \beta$ for lines [\(49\)](#page-28-5) and [\(52\)](#page-29-0). Line [\(51\)](#page-29-1) does not contain terms with $\beta_2 = \beta$, as these appear in the line we have singled out [\(50\)](#page-28-6) We have thus reduced the invariant $\mathrm{OGW}_{\beta,k}(\Gamma_{j_1},\ldots,\Gamma_{j_l})$ to terms of lower degree, or of the same degree, but with the degree of one of the interior constraints decreased by 2. \Box

Using the above two lemmas we can compute:

Lemma 6.6. The closed Gromov-Witten invariants, together with Lemmas [6.4](#page-28-7) and [6.5](#page-28-8) allow us to compute: $\text{OGW}_{1,1}(\Gamma_{r+1}, \Gamma_{r+1}) = 2$.

Lemma 6.7. All open Gromov-Witten invariants with no interior constraints, except for $\rm{OGW}_{1,3}$, can be computed from invariants of lower degree, and invariants of equal degree, but with at least one interior constraint of degree greater than 2.

Proof. Consider the open WDVV equation [\(22\)](#page-11-2), with $v = w = r + 1$, and apply ∂_s^{k-1} , taking the coefficient of $T^{\beta+1}$. We find:

$$
GW1(\Delta_{r+1}, \Delta_{r+1}, \Delta_{2r})\text{OGW}_{\beta,k}(\Gamma_1)g^{2r,1}
$$
\n(53)

$$
+\sum_{\beta_1+\beta_2=\beta+1} GW_{\beta_1}(\Gamma_{r+1},\Gamma_{r+1},\Gamma_i)\text{OGW}_{\beta_2,k}(\Gamma_{n-i})g^{i,n-i}
$$
(54)

$$
\rho_2 < \beta
$$
\n
$$
= (k-1) \text{OGW}_{\beta,k} \text{OGW}_{1,1}(\Gamma_{r+1}, \Gamma_{r+1}) \tag{55}
$$

$$
-\sum_{\substack{\beta_1+\beta_2=\beta+1\\ \beta_1<\beta}} \binom{k-1}{i} \text{OGW}_{\beta_1,1+k-i} \text{OGW}_{\beta_2,i}(\Gamma_{r+1},\Gamma_{r+1})\tag{56}
$$

$$
= - \sum_{\substack{\beta_1 + \beta_2 = \beta + 1 \\ \beta_1, \beta_2 \leq \beta \\ i = 0, \dots, k-1}} \binom{k-1}{i} \text{OGW}_{\beta_1, 1+i}(\Gamma_{r+1}) \text{OGW}_{\beta_2, k-i}(\Gamma_{r+1}) \tag{57}
$$

For line [\(54\)](#page-29-2), note there are no contributions with $\beta_2 = \beta + 1$, as $GW_0(\Gamma_{r+1}, \Gamma_{r+1}, \Gamma_i) = 0$ for any i, by the degree property. There are also no contributions with $\beta_2 = \beta$, as the only $GW_1(\Gamma_{r+1}, \Gamma_{r+1}, \Gamma_i)$ that contributes is the one with $i = 2r$, which is the term in the line above it. Secondly for line [\(56\)](#page-29-3) there are no contributions with β_1 or $\beta_2 = \beta + 1$, as then the other term is $\text{OGW}_{0,\star}$ with either 0 or 2 interior marked points, which thus vanishes by the degree zero property. Also, there only contribution with $\beta_1 = \beta$ has been singled out in the line above it. Finally for [\(57\)](#page-29-4) there are no contributions with β_1 or $\beta_2 = \beta + 1$, as then the other term is $\text{OGW}_{0,\star}$ with 1 interior marked point on Γ_{r+1} . As $r+1>0$, this vanishes by the degree zero property. By the previous lemma, we can combine lines [\(53\)](#page-29-5) and [\(55\)](#page-29-6) to: $2(2\beta - k + 1)$ OGW_{β,k}.

By the degree property of the OGW, we have $2n\beta = k(n-1)+3-n$, and thus $n(2\beta-k+1) =$ $3 - k \neq 0$ unless $k = 3$ and $\beta = 1$. The invariant OGW_{1,3} is precisely the invariant we have computed already, thus the above equations determine $\text{OGW}_{\beta,k}$ in terms of invariants of lower degree, or of equal degree, with at least 1 interior constraint.

Finally, we combine the above information to prove Theorem [7.](#page-4-1)

Proof of theorem [7.](#page-4-1) Let $\mathrm{OGW}_{\beta,k}(\Gamma_I)$ be an invariant we want to compute. By the divisor axiom, we may assume $1 \notin I$. By Theorem [14](#page-11-0) we may assume $\diamond \notin I$. Furthermore, the number of boundary constraints k is determined by β , I and the degree axiom of OGW. We now introduce a total ordering on the set $\{(\beta,I)\in\mathbb{N}\times\{0,2,3,\ldots,n\}^{|I|}\}$ of parameters for open Gromov-Witten invariants. Order the degrees β by the natural ordering on N (under the identification $H_2(X, L) \cong \mathbb{Z}$. Order the index sets I as follows: the element $I = \emptyset$ is the largest element. Then, if $|I| \neq \emptyset$ order such I by |I| under the natural ordering on $\mathbb{N}_{>0}$. Assume that I is ordered so that $i_1 \leq i_2 \leq \cdots \leq i_{|I|}$. If two sets I, I' have the same size, say $I < I'$ if I appears first in the lexicographic ordering on $(N \setminus \{1\})^{|I|}$. The total order on tuples (β, I) is then given by the lexicographic order. This total order satisfies the well-ordering principle, so we can apply induction. The three lemmas [6.4,](#page-28-7) [6.5](#page-28-8) and [6.7](#page-29-7) precisely show that the open WDVV equations allow us to compute the invariant $\text{OGW}_{\beta,k}(\Gamma_I)$ in terms of invariants strictly lower in the ordering on (β, I, k) . Except for the case OGW_{1,3} which we calculated separately. The result thus follows by induction. \Box

We now consider how the invariant change if instead we run the above induction process with $OGW_{1,3} = -2$. We first have the following observation:

Lemma 6.8. For $I \subset \{0, 1, ..., n\}^l$ and k even, we have: $OGW_{\beta,k}(\Gamma_I) = 0$.

Proof. The result holds for $\beta = 0$. Then apply the above recursion process, noting that always at least one of the terms in the products has an even number of boundary constraints. \Box

We then show:

Proposition 6.9. Suppose that instead we run the above induction process with the initial value $\overline{\text{OGW}}_{1,3} = -2$, then for any $I \subset \{0, 1, \ldots, n\}^l$, the resulting invariant $\overline{\text{OGW}}_{\beta,k}(\Gamma_I)$ changes by the sign $(-1)^{\frac{k-1}{2}}$ and the invariant $\text{OGW}_{\beta,k}(\Gamma_{\diamond}^{\otimes r},\Gamma_I)$ changes by the sign $(-1)^{\frac{k+r-1}{2}}$.

Proof. Consider the sign changed invariants \overline{OGW}° with $r = 2$ as in Definition [2.8.](#page-12-5) As a reminder, these are defined by $\overline{\text{OGW}}_{\beta,k}^{\circ} = (-1)^{\frac{k-1}{2}} \overline{\text{OGW}}_{\beta,k}$. By Proposition [2.10,](#page-12-1) the invariants \overline{OGW}° still satisfy the open WDVV equations and the open Gromov-Witten axioms. As these, together with the initial value, were the only inputs for the above induction, the result follows. For invariants with inputs Γ_{\diamond} the result follows from the first statement and the wall-crossing formula [14.](#page-11-0)

7. The quadric – even dimension

7.1. Relative quantum ring. In this section, we assume $n > 2$ is even. For $n = 2$ see Section [8.](#page-34-0) We work with the basis

$$
\Gamma_j := [r^* \omega_{FS}]^j, \qquad j = 0, \dots, n,
$$

of $H^*(X, L; \mathbb{R})$. We also write $\Delta_j := h^j \in H^*(X; \mathbb{R})$. Note that in the exact sequence [\(5\)](#page-6-5) we have $\rho(\Gamma_j) = \Delta_j$. By Lemma [5.3](#page-19-2)[\(a\),](#page-19-4) the Poincaré pairing on Span $(\{\Delta_j\}_{j=0}^n)$ is given by

$$
g_{ij} = \int_X \Delta_i \wedge \Delta_j = \delta_{i,n-j} \cdot 2,
$$

where $\delta_{k,l}$ is the Kronecker delta, and the inverse matrix $(g^{ij})_{i,j} := (g_{ij})_{i,j}^{-1}$ is given by $g^{ij} = \delta_{i,n-j} \cdot \frac{1}{2}$ $\frac{1}{2}$. Thus, the relative quantum product is given by

$$
\mathbf{D}(\Gamma_i, \Gamma_j) = \sum_{\substack{0 \le m \le n \\ d \ge 0}} \frac{1}{2} T^d \, \mathrm{GW}_d(\Delta_i, \Delta_j, \Delta_m) \Gamma_{n-m}, \qquad i, j \in \{0, \ldots, n\}.
$$

Proof of Theorem [5.](#page-3-1) By degree axiom [\(8\)](#page-8-1), the only nonzero values of $\text{GW}_d(\Delta_i, \Delta_j, \Delta_m)$ occur for $d \leq 2$. For $d = 2$, the value is only nonzero if $i = j = m = n$. We will not need to compute $\text{GW}_2(\Delta_n^{\otimes 3})$ explicitly, although we deduce it below in Remark [7.1.](#page-32-1) For $d=0$, the zero axiom [\(10\)](#page-8-3) says the only nontrivial values are when $i + j + m = n$, in which case $GW_0(\Delta_i, \Delta_j, \Delta_m) = 2$ by Section [5.3.](#page-24-2)

For $d = 1$, the degree axiom [\(8\)](#page-8-1) says the only nontrivial values occur when $i + j + m = 2n$. By Lemma [5.17,](#page-24-1) the resulting values are

$$
GW_1(\Delta_i, \Delta_j, \Delta_m) = \begin{cases} 0, & (i, j, m) = (0, n, n), \\ 4, & i > 0, m = n, \\ 8, & i > 0, m < n. \end{cases}
$$

Thus, we have

$$
\mathbf{D}(\Gamma_i, \Gamma_j) = \begin{cases} \Gamma_{i+j}, & i+j < n, \\ 2T + \Gamma_n, & i+j = n, \\ 2T\Gamma_{i+j-n}, & n < i+j < 2n. \end{cases}
$$

To compute the case $i = j = n$, use the associativity of **p**. For short, we rite Γ_1^{pk} for the k -fold product of Γ_1 under Δ . So,

$$
\mathbf{a}(\Gamma_n, \Gamma_n) = \mathbf{a}(\Gamma_1^{2n} - 2T, \Gamma_1^{2n} - 2T) \n= \Gamma_1^{2n} - 4T\Gamma_1^{2n} + 4T^2 \n= \mathbf{a}(\Gamma_1^{2(n+1)}, \Gamma_1^{2(n-1)}) - 4T\Gamma_1^{2n} + 4T^2 \n= \mathbf{a}(4T\Gamma_1, \Gamma_1^{2(n-1)}) - 4T\Gamma_1^{2n} + 4T^2 \n= 4T^2.
$$
\n(58)

In total,

$$
\mathbf{D}(\Gamma_i, \Gamma_j) = \begin{cases} \Gamma_{i+j}, & i+j < n, \\ 2T + \Gamma_n, & i+j = n, \\ 2T\Gamma_{i+j-n}, & n < i+j < 2n, \\ 4T^2, & i = j = n. \end{cases}
$$

Similarly to the odd dimensional case, we note that $\{\Gamma_1^{\alpha}$ $\{a_j\}_{j=0}^n$ is a basis of $QH^*(X,L)$ as a module. To compute the ring structure we need to spell out the higher powers of Γ_1 . Specifically,

$$
\Gamma_1^{\mathbf{D}(n+1)} = \mathbf{D}(\Gamma_1, 2T + \Gamma_n) = 4T\Gamma_1.
$$

Taking the variables q, x, to correspond to T, Γ_1 , respectively, we get the desired result. \Box Remark 7.1. It follows from the computation [\(58\)](#page-31-1) that $\text{GW}_2(\Delta_n^{\otimes 3}) = 2[T^2](\textbf{D}(\Gamma_n, \Gamma_n)) = 8$.

7.2. Absolute quantum ring. In this section, we work with the basis $\{\Delta_j\}_{j=0}^n\cup\{\Delta_L\}$ of $H^*(X;\mathbb{R})$, where $\Delta_j = h^j$ as in Section [7.1](#page-31-2) and $\Delta_L = PD([L])$ as in [\(7\)](#page-7-0). For $i, j \in$ $\{0,\ldots,n,L\}$, we have

$$
\Delta_i * \Delta_j = \sum_{\substack{0 \le m \le n \\ d \ge 0}} \frac{1}{2} T^d \, \mathrm{GW}_d(\Delta_i, \Delta_j, \Delta_m) \Delta_{n-m} + \sum_{d \ge 0} \frac{(-1)^{n/2}}{2} T^d \, \mathrm{GW}_d(\Delta_i, \Delta_j, \Delta_L) \Delta_L.
$$

Proposition 7.2. The absolute quantum cohomology is given by

$$
QH^*(X) \simeq \mathbb{R}[[q]][x,y]\bigg/\!\!f,
$$

with

$$
I = (x^{n+1} - 4qx, xy, y^2 - (-1)^{n/2+1}4q - (-1)^{n/2}x^n).
$$

Proof. By the degree axiom [\(8\)](#page-8-1), $GW_d(\Delta_i, \Delta_j, \Delta_L) \neq 0$ implies $d \leq 1$. For $d = 0$, the zero axiom [\(10\)](#page-8-3) gives

$$
GW_0(\Delta_L, \Delta_L, 1) = \int_X \Delta_L^{\wedge 2} = (-1)^{n/2} 2.
$$

Recall the relation $xy = 0$ from Proposition [5.8.](#page-22-0) This implies that for $i, j \in \{0, ..., n\}$, $GW_0(\Delta_L, \Delta_i, \Delta_j) = 0$. Finally, by the degree axiom [\(8\)](#page-8-1) $GW_0(\Delta_L^{\otimes 3}) = 0$.

For $d = 1$, nontrivial values only appear when $|\Delta_i| + |\Delta_j| + |\Delta_L| = 2n$. In particular, $(i, j) \neq (L, L)$. By equation [\(41\)](#page-21-2), we have

$$
\int_L i^* \Delta_L = \int_X \Delta_L^{\wedge 2} = (-1)^{n/2} 2.
$$

Then $\eta := \frac{(-1)^{n/2}}{2} \Delta_L$ satisfies $\int_L \eta = 1$. By the open-closed relation [\(24\)](#page-11-4), for any ξ with $\int_L \xi = 0$ we have

$$
OGW_{\beta,1}(\gamma_1,\ldots,\gamma_l) = - GW_{\beta}(\eta + \xi, \Delta_L, \gamma_1,\ldots,\gamma_l).
$$

Thus, for $i, j \in \{0, ..., n\},\$

$$
GW_1(\Delta_i, \Delta_L, \Delta_j) = GW_1(\eta + \Delta_i, \Delta_L, \Delta_j) - GW_1(\eta, \Delta_L, \Delta_j) =
$$

=
$$
OGW_{1,1}(\Gamma_j) - OGW_{1,1}(\Gamma_j) = 0,
$$

while by Lemma [5.18,](#page-25-3)

 $\text{GW}_1(\Delta_L, \Delta_L, \Delta_n) = (-1)^{n/2} 2 \cdot \text{GW}_1(\eta, \Delta_L, \Delta_n) = (-1)^{n/2+1} 2 \cdot \text{OGW}_{1,1}(\Gamma_n) = (-1)^{n/2+1} 4.$

In total, we get

$$
\Delta_i * \Delta_j = \begin{cases}\n\rho(\mathbf{D}(\Gamma_i, \Gamma_j)), & i, j \in \{0, \dots, n\}, \\
\Delta_L, & i = L, j = 0, \\
-2T\Delta_L, & i = L, j = n, \\
(-1)^{n/2+1}2T + (-1)^{n/2}\Delta_n, & i = j = L, \\
0, & i = L, j \neq L, 0, n.\n\end{cases}
$$

Recall that $\Delta_n = \Delta^{*n} - 2T$. Taking the variables q, x, y, to correspond to T, Δ_1, Δ_L , respectively, we get the desired result.

7.3. Full computation of invariants. In this section we prove Theorem [8.](#page-4-2) First observe that for $I \subset \{0, 1, \ldots, n, L\}^l$ we have:

- OGW_{β,0}(Γ_I) = 0 by definition, as $H_2(X, L) \cong H_2(X)$.
- OGW_{$\beta,1$} $(\Gamma_I) = \frac{(-1)^{\frac{3n}{2}+1}}{2}$ GW_d $(PD(L), PD(L), \Gamma_I)$ by Theorem [15.](#page-11-1)
- $\mathrm{OGW}_{\beta,k}(\Gamma_I) = 0$ for $k \geq 2$, also by Theorem [15.](#page-11-1)

It will thus suffice to compute all closed Gromov-Witten invariants. We will provide a method to compute these. We first observe:

Lemma 7.3. $GW_{\beta}(PD(L)^{\otimes 2s+1}, \Delta_I) = 0$ for all β , s, and $I \subset \{0, 1, \ldots, n\}^l$.

Proof. Consider the symplectomorphism $\phi : \mathbb{CP}^{n+1} \to \mathbb{CP}^{n+1}$ given by $\phi(X) = -X$. ϕ restricts to a symplectomorphism of V, mapping L to itself. $\phi|_L$ is the antipodal map. Thus, as L is an even-dimensional sphere, $\phi^*(PD(L)) = -PD(L)$. However $\phi^*(\Delta_I) = \Delta_I$ as $\phi^* h = h$. As ϕ is a symplectomorphism we find

$$
GW_{\beta}(PD(L)^{\otimes 2s+1}, \Delta_I) = GW_{\phi*\beta}(PD(L)^{\otimes 2s+1}, \Delta_I)
$$
\n(59)

$$
= GW_{\beta}(\phi^*(PD(L)^{\otimes 2s+1}), \phi^*\Delta_I)
$$
\n(60)

$$
= (-1)^{2s+1} GW_{\beta}(PD(L)^{\otimes 2s+1}, \Delta_I). \tag{61}
$$

 \Box

The next step is to rule out any invariants with constraints only along $PD(L)$.

Lemma 7.4. $GW_{\beta}(PD(L)^{\otimes s}) = 0$ for all β and s.

Proof. For s odd this follows from the previous lemma. So assume $s = 2k$. The degree axiom of GW then implies:

$$
4r + 4r\beta + 4k - 6 = 4kr,
$$

Here $n = 2r$. Reducing this equation modulo 4 shows this is impossible.

Finally, we can obtain all closed Gromov-Witten invariants.

Lemma 7.5. All Gromov-Witten invariants are determined by the two point invariant $GW_1(\Delta_{n-1}, \Delta_n) = 4.$

Proof. First observe that by the degree equation there are no non-zero invariants $GW_{\beta}(\Delta_I)$ with $|I| < 2$, and if $|I| = 2$, the only possibility is $GW_1(\Delta_{n-1}, \Delta_n)$.

Denote $PD(L)$ by Δ_L and introduce the total order $0 < 1 < \cdots < n < L$ on $\{0, 1, \ldots, n, L\}$: J. Define the following total order on the set of possible β, I . Here we always order I to be arranged as $i_1 \leq i_2 \leq \cdots \leq i_{|I|}$.

- $(\beta_1, I_1) < (\beta_2, I_2)$ when $\beta_1 < \beta_2$.
- $(\beta, I_1) < (\beta, I_2)$ when $|I_1| < |I_2|$.
- $(\beta, I_1) < (\beta, I_2)$ when $|I_1| = |I_2|$ and $I_1 < I_2$ in the lexicographic ordering on $J^{|I_1|}$.

We will now perform induction with respect to this total order. To this end, let $GW_{\beta}(\Delta_I)$ be an invariant we want to compute, with $|I| \geq 3$. Write $I = (i_1, i_2, i_3, \overline{I})$. By the divisor and unit axioms, we may assume $i_1 > 1$ (and hence $i_j > 1$ for all j). By lemma [7.4:](#page-33-0) $i_1 \neq L$. We apply the WDVV equation [\(11\)](#page-9-2) with $u = i_1 - 1$, $v = 1$, $w = i_2$, $y = i_3$ in degree β , with extra derivatives on I . This yields:

$$
\mathrm{GW}_0(\Delta_1, \Delta_{i_1-1}, \Delta_{n-i_1}) \mathrm{GW}_\beta(\Delta_I) g^{i_1, n-i_1} \tag{62}
$$

$$
+\sum_{\mu,\nu\in J} \text{GW}_{\beta}(\Delta_1,\Delta_{i_1-1},\Delta_{\bar{I}},\Delta_{\mu}) \text{GW}_0(\Delta_{\nu},\Delta_{i_2},\Delta_{i_3}) g^{\mu,\nu} \tag{63}
$$

+
$$
\sum_{\substack{\beta_1+\beta_2=\beta\\ \beta_1,\beta_2<\beta\\ I_1\sqcup I_2=\overline{I}}} \text{GW}_{\beta_1}(\Delta_1,\Delta_{i_1-1},\Delta_\mu,\Delta_{I_1}) \text{GW}_{\beta_2}(\Delta_\nu,\Delta_{i_2},\Delta_{i_3},\Delta_{I_2})
$$
(64)

$$
\mu, \nu \in J
$$

= GW₀($\Delta_1, \Delta_{i_2}, \Delta_{n-i_2-1}$) GW_{\beta}($\Delta_{i_2+1}, \Delta_{i_1-1}, \Delta_{i_3}, \Delta_{\bar{I}}$) $g^{i_2+1, n-i_2-1}$ (65)

$$
+\sum_{\mu,\nu\in J} \text{GW}_{\beta}(\Delta_1,\Delta_{i_2},\Delta_{\bar{I}},\Delta_{\mu}) \text{GW}_0(\Delta_{\nu},\Delta_{i_1-1},\Delta_{i_3}) g^{\mu,\nu} \tag{66}
$$

+
$$
\sum_{\substack{\beta_1+\beta_2=\beta\\ \beta_1,\beta_2<\beta\\ I_1\sqcup I_2=\bar{I}\\ \mu,\nu\in J}} \text{GW}_{\beta_1}(\Delta_1,\Delta_{i_2},\Delta_{\mu},\Delta_{I_1}) \text{GW}_{\beta_2}(\Delta_{\nu},\Delta_{i_1-1},\Delta_{i_3},\Delta_{I_2})
$$
(67)

$$
(68)
$$

Now [\(62\)](#page-34-1) will determine the invariant we want to compute, as $GW_0(\Delta_1, \Delta_{i_1}, \Delta_{n-i_1}) = 4$ and $g^{i_1,n-i_1} = 1/2$. The first term in [\(63\)](#page-34-2) is smaller in the total ordering on indices, as by the divisor axiom we can reduce the number of marked points to $|I| - 1$. All terms in line [\(64\)](#page-34-3) are lower down in the ordering, as the degrees are smaller than β . The second term of [\(65\)](#page-34-4) is lower down in the ordering because the the lowest index i_1 has been lowered by 1. Also observe here that we abuse notation by writing $i_2 + 1$ and $n - i_2 - 1$. If $i_2 = L$, this first term in [\(65\)](#page-34-4) vanishes as $\Delta_1 \cup \Delta_L = 0$. The terms in line [\(66\)](#page-34-5) again have fewer interior marked points by the divisor equation, and those in line [\(67\)](#page-34-6) have smaller degrees. We have thus determined $\text{GW}_\beta(\Delta_I)$ in terms of invariants strictly lower down in the ordering. This finishes the proof by induction.

8. The quadric – dimension 2

let X be the quadric hypersurface in \mathbb{CP}^3 given by $\sum_{j=0}^2 z_j^2 - z_3^2 = 0$. Let $L \subset X$ be the real locus. In this section we will compute the relative quantum cohomology of (X, L)

8.1. Topological preliminaries. Consider the map

$$
\phi: \mathbb{CP}^1 \times \mathbb{CP}^1 \to \mathbb{CP}^3 \tag{69}
$$

$$
([x_0:x_1],[y_0:y_1]) \mapsto [x_0y_0-x_1y_1:-i(x_0y_0+x_1y_1):x_1y_0+x_0y_1:x_0y_1-x_1y_0]
$$
(70)

One then checks that ϕ is injective and $im(\phi) = X$. Thus $X \simeq \mathbb{CP}^1 \times \mathbb{CP}^1$. Let $A_1, A_2 \in$ $H_2(X;\mathbb{Z})$ be defined by $A_1 = \phi_*[\mathbb{CP}^1 \times \{[0:1]\}]$ and $A_2 = \phi_*[\{[0:1]\} \times \mathbb{CP}^1]$. Let $l = PD(A_1), l^* = PD(A_2) \in H^2(X; \mathbb{R})$. Then

$$
H^*(X; \mathbb{R}) \simeq \mathbb{R}[l, l^*]/(l^2, \quad l^{*2}).
$$

Let $L \subset X$ be given by the fixed locus of complex conjugation. Thus $L \simeq S^2$. We now compute $[L] \in H_2(X;\mathbb{Z})$. An easy computation shows that $|[L] \cap A_i| = 1$, so that $[L] =$ $\pm A_1 \pm A_2$. It remains to compute the signs. In order to do this, we consider $\phi|_L^{-1}$ $L^{-1}: L \to$ $\mathbb{CP}^1 \times \mathbb{CP}^1$. To compare orientations, it is easiest to precompose this with the (orientation preserving) stereographic projection

$$
\psi: \mathbb{R}^2 \to L \subset X \tag{71}
$$

$$
(u, v) \mapsto [2u : 2v : u^2 + v^2 - 1 : 1 + u^2 + v^2]
$$
\n
$$
(72)
$$

On the image of ψ , we have $z_3 \neq z_4$ so that the map $\phi|_{im}^{-1}$ $\lim_{i m(\psi)}^{1} : L \to \mathbb{CP}^1 \times \mathbb{CP}^1$ given by

$$
[z_0:z_1:z_2:z_3] \mapsto [z_1+iz_2:z_3-z_4], [z_3-z_4:iz_2-z_1],
$$

is well defined. The composition with ψ is then given by:

$$
(u, v) \mapsto [-(u + iv) : 1], [1 : u - iv].
$$

Now consider the maps π_i : $\mathbb{CP}^1 \times \mathbb{CP}^1 \to \mathbb{CP}^1$. The above then shows that the map $\pi_1 \circ \phi|_{im}^{-1}$ $\lim_{i \to i(\psi)} \circ \psi$ is orientation preserving, whereas the map $\pi_1 \circ \phi|_{im}^{-1}$ $\overline{\psi}_{im(\psi)}^{-1} \circ \psi$ reverses orientation. Thus

$$
[L] = A_1 - A_2 \in H_2(X; \mathbb{Z}).
$$

8.2. Absolute quantum ring. As shown in [\[3,](#page-43-6) Example 7.2] the absolute quantum cohomology of X is given by

$$
QH^*(X; \Lambda_c) \simeq \mathbb{R}[[q_1, q_2]][l, l^*]/(l^2 - q_2, l^{*2} - q_1).
$$

Here

$$
\Lambda_c = \left\{ \sum_{i=0}^{\infty} a_i T^{\beta_i} \middle| a_i \in \mathbb{R}, \beta_i \in H_2(X; \mathbb{Z}), \omega(\beta_i) \ge 0, \lim_{i \to \infty} \omega(\beta_i) = \infty \right\}.
$$

Where we have taken the identification

$$
\Lambda_c \simeq \mathbb{R}[[q_1, q_2]]
$$

via the map $T^{A_1} \mapsto q_1, T^{A_2} \mapsto q_2$. However, to compare $QH^*(X)$ with $QH^*(X, L)$ we need to work over a Novikov ring with exponents in $H_2(X, L; \mathbb{Z})$. As $[L] = A_1 - A_2$, and $H_1(L; \mathbb{Z}) = 0$ this amounts to identifying $q_1 = q_2$. We additionally change basis to $u = l + l^*$, $v = l - l^*$ to obtain:

$$
QH^*(X) \simeq \mathbb{R}[[q]][u, v]/(u^2 + v^2 - 4q, uv).
$$

Note that this latter result agrees with the quantum cohomology for the general even dimensional quadric.

8.3. Relative quantum ring. As $[L] \neq 0$, we have $QH^*(X; L) \subset QH^*(X)$ given by the kernel of integration over L, i_L . As a $\mathbb{R}[[q]]$ -module, $QH^*(X, L)$ is thus spanned by 1, u and $l \cup l^*$. From the previous section, we have that $u \star u = 2q + 2l \cup l^*$. We thus find:

$$
QH^*(X,L) \simeq \mathbb{R}[[q]][u]/(u^3-4uq).
$$

We have thus proven Theorem [6.](#page-3-2)

8.4. Full computation of invariants. $(1,0) = [\mathcal{L}], (0,1) = [\mathcal{L}^{\star}].$

Lemma 8.1. The basic Gromov-Witten invariants are given by

$$
GW_{(d_1A_1+d_2A_2)}(l \cup l^*, \alpha) = \begin{cases} 1, & (d_1, d_2, \alpha) \in \{ (1, 0, l), (0, 1, l^*) \}, \\ 1, & (d_1, d_2, \alpha) = (0, 0, l \cup l^*), \\ 0, & otherwise. \end{cases}
$$

Proof of Theorem [9.](#page-4-3) As $H^2(X)$ generates $H^*(X)$ as an algebra, the result for closed invari-ants follows from [\[5\]](#page-43-3). As $[L] \neq 0$, the result for open Gromov-Witten invariants follows from the result for closed invariants and Theorem [15.](#page-11-1)

Appendix A. Initial value

Our goal here is to prove Lemma [6.2.](#page-27-0)

A.1. **Setting up the computation.** For any space, denote by pt the map from it to a point. We use the notation of [\[12,](#page-43-1) [13\]](#page-43-8) to write $\text{OGW}_{1,3}$ as

$$
\begin{split}\n(\partial_s^3 \mathfrak{q}_{-1,0}^{\beta_1;\gamma,b})|_{s=t_j=0} &= (\partial_s^3 \mathfrak{q}_{-1,0}^{\beta_1;0,b})|_{s=t_j=0} \\
&= \langle \mathfrak{q}_{1,0}^{\beta_1}(\partial_s b, \partial_s b), \partial_s b \rangle \\
&= (-1)^{n+1} p t_*((evb_0)_*(evb_1^* \partial_s b \wedge evb_2^* \partial_s b) \wedge \partial_s b) \\
&= (-1)^{1+n \cdot \text{rel dim} evb_0} p t_* (\partial_s b \wedge (evb_0)_*(evb_1^* \partial_s b \wedge evb_2^* \partial_s b)) \\
&= pt_*(evb_0^* \partial_s b \wedge evb_1^* \partial_s b \wedge evb_2^* \partial_s b),\n\end{split} \tag{73}
$$

where b is (the) point-like bounding chain guaranteed by Theorem 2 of [\[13\]](#page-43-8).

Let $Y = L \times L \times L \setminus \Delta$ and let $Z = (evb_0 \times evb_1 \times evb_2)^{-1}(Y) \subset M_{3,0}(\beta_1)$. Observe that $g = (evb_0 \times evb_1 \times evb_2)|_Z : Z \to Y$

is a covering map. Indeed, the fibre of g over a point
$$
(x_1, x_2, x_3) \in Y
$$
 can be identified with
the pair of real oriented degree-2 curves in L passing through x_1 , x_2 , and x_3 . We claim that
for the unique choice of (relative) spin structure with $w_s = 0$, the degree of g is 2. Indeed, the
two points in the preimage of a point in Y are conjugate disks of degree 1 with three marked
points. Proposition 5.1 of [11] shows that the covering transformation that interchanges
these two disks is orientation preserving:

$$
s^+(1,3,0) \equiv \frac{2n(2n+1)}{2} + n + n + 3 + 0 + (n+1) \sum_a \frac{(k_a-1)(k_a-2)}{2}
$$

$$
\equiv n(2n+1) + 1 \equiv 0 \pmod{2}.
$$

So, the degree is ± 2 . We postpone the proof of the following proposition until later in this section.

Proposition A.1. The unique spin structure on L is such that g has degree $(-1)^{\frac{(n-2)(n-3)}{2}}2$. For $i = 1, 2, 3$, let $p_i: L \times L \times L \to L$ be the projections. Since $Y \subset L \times L \times L$ and $Z \subset \mathcal{M}_{3,0}(\beta_1)$ are open dense subsets, using that b is point-like, we obtain

$$
pt_*((evb_0)^*\partial_s b \wedge (evb_1)^*\partial_s b \wedge (evb_2)^*\partial_s b) = \int_Z g^*((p_1^*\partial_s b \wedge p_2^*\partial_s b \wedge p_3^*\partial_s b)|_Y)
$$

$$
= (-1)^{\frac{(n-2)(n-3)}{2}} 2 \int_Y p_1^*\partial_s b \wedge p_2^*\partial_s b \wedge p_3^*\partial_s b
$$

$$
= (-1)^{\frac{(n-2)(n-3)}{2}} 2 \int_{L \times L \times L} p_1^*\partial_s b \wedge p_2^*\partial_s b \wedge p_3^*\partial_s b
$$

$$
= (-1)^{\frac{(n-2)(n-3)}{2}} 2 \left(\int_L \partial_s b\right)^3
$$

$$
= (-1)^{\frac{(n-2)(n-3)}{2}} 2.
$$

Combining this calculation with equation [\(73\)](#page-36-2) we get that the value of $\overline{\text{OGW}}_{1,3}$ is $(-1)^{\frac{(n-2)(n-3)}{2}}2$.

Remark A.2. The choice of just the orientation on L has no effect on the value, since a change of orientation would contribute $(-1)^4 = +1$ to the sign of $\left(\int_L \partial_s b\right)^3$.

It is left for us to prove Proposition [A.1.](#page-37-0) At a point

$$
\mathbf{u} = [u, \vec{z} = (1, i, -1), \vec{w} = ()] \in \mathcal{M}_{3,0}(\beta_1),
$$

we need to compute the sign of the evaluation map

$$
dg_{\mathbf{u}}: T_{\mathbf{u}}\mathcal{M}_{3,0}(\beta_1) = \Gamma(u^*TX, u^*TL) \longrightarrow T_{u(1)}L \oplus T_{u(i)}T \oplus T_{u(-1)}L.
$$

In section [A.2](#page-38-0) we prove:

Lemma A.3. The sign of dg_u is independent of the choice of u .

Thus, we can pick a convenient map \mathbf{u}_0 and compute the sign $dg_{\mathbf{u}_0}$. We outline this computation. First, consider the Cauchy-Riemann boundary value problem

$$
\mathcal{E} = (E, F)
$$

given by the trivial line bundle over $(D, \partial D)$ with boundary conditions

$$
F_z = z\mathbb{R}, \qquad z \in \partial D.
$$

The Cauchy-Riemann boundary problem $\mathcal E$ has a canonical orientation and spin structure, $\mathfrak{p}_{\mathcal{E}}$, given by trivialising the bundle F_z by division by z. This then equips $\Gamma(\mathcal{E})$ with a natural orientation (see e.g. [\[11\]](#page-43-13)). In section [A.4](#page-42-0) we show:

Lemma A.4. The canonical orientation and spin structure on \mathcal{E} equip $\Gamma(\mathcal{E})$ with an orientation such that

$$
\Gamma(\mathcal{E}) \xrightarrow{Ev} F_1 \oplus F_i \oplus F_{-1}
$$

is orientation preserving.

We then show that the problem we are interested in decomposes into copies of these line bundles:

Lemma A.5. There exists an isomorphism of Cauchy-Riemann boundary value problems

$$
(u_0^*TX, u^*TL) \xrightarrow{\varphi} \mathcal{E}^{\oplus n}.
$$

Moreover, the map φ does not respect spin structures.

The proof of the above lemma is given in section [A.3.](#page-39-0) Then consider the commutative diagram:

$$
\Gamma(u^*TX, u^*TL) \xrightarrow{dg_u} T_{u(1)}L \oplus T_{u(i)}L \oplus T_{u(-1)}L
$$
\n
$$
\downarrow^{\varphi_*}
$$
\n
$$
\Gamma(\mathcal{E}^{\oplus n})
$$
\n
$$
\downarrow^{\tau}
$$
\n
$$
\Gamma(\mathcal{E})^{\oplus n} \xrightarrow{Ev} (F_1 \oplus F_i \oplus F_{-1})^{\oplus n} \xrightarrow{\sigma} F_1^{\oplus n} \oplus F_i^{\oplus n} \oplus F_{-1}^{\oplus n}
$$

Here τ is the map induced by restricting φ to the fibres over 1, i and -1 . To recover the sign of the arrow on the top row, we compute the signs of all the intermediate steps. From Lemma [A.5](#page-38-1) we have that $sgn(\varphi_*) = 1 + sgn(\varphi|_{TL})$, where the 1 comes from the fact that it doesn't respect the spin structure, and $sgn(\varphi|_{TL})$ is the sign coming from whether or not the map $\varphi : u^*TL \to F^{\oplus n}$ respects orientations. Next, note that the natural map $\Gamma(\mathcal{E}^{\oplus n}) \to \Gamma(\mathcal{E})^{\oplus n}$ is orientation preserving by [\[11,](#page-43-13) Lemma 2.16]. Then, we have the map Ev which, as shown in Lemma [A.4,](#page-37-1) is orientation preserving. The map σ is just a permutation, with sign $sgn(\sigma) = \frac{n(n-1)}{2}$. Finally, we have the map τ , which is induced by φ . It has sign $sgn(\varphi|_{TL})^3 = sgn(\varphi|_{TL})$. The overall sign is thus

$$
1 + \frac{n(n-1)}{2} \equiv \frac{(n-2)(n-3)}{2},
$$

as required. Note in particular that we don't need to compute $sgn(\varphi|_{TL})$.

A.2. **Proof of Lemma [A.3.](#page-37-2)** Showing $sgn(dg_u)$ is independent of the choice of u.

Let $Q: \mathbb{C}^{n+2} \to \mathbb{C}$ be the quadratic form given by

$$
Q(z) = \sum_{j=0}^{n} z_j^2 - z_{n+1}^2.
$$

Let

$$
\widehat{X} := \{ Q(z) = 0 \} \subset \mathbb{C}^{n+2}
$$

and the real locus

$$
\widehat{L}\subset\mathbb{R}^{n+2}.
$$

Let $e_0, \ldots, e_{n+1} \in \mathbb{C}^{n+2}$ denote the standard basis and let e_0^*, \ldots, e_{n+1}^* denote the dual basis. Let

$$
\langle \cdot, \cdot \rangle_{n+1,1} = \sum_{j=0}^{n} e_j^* \otimes e_j^* - e_{n+1}^* \otimes e_{n+1}^*
$$

39

denote the standard bilinear form of signature $(n + 1, 1)$. Let $P \subset \mathbb{R}^{n+2}$ be an oriented 3-dimensional linear subspace such that $\langle \cdot, \cdot \rangle_{n+1,1}|_P$ has signature $(2, 1)$. Let $P^{\mathbb{C}} = P \otimes \mathbb{C} \subset \mathbb{C}$ \mathbb{C}^{n+2} . Observe that $\mathbb{P}(P \cap \widehat{L})$ is isomorphic as an algebraic variety to \mathbb{RP}^1 . On the other hand, the projection map $P_L := \widehat{L} \cap P \cap \{z_{n+1} = 1\} \to \mathbb{P}(P \cap \widehat{L})$ is a diffeomorphism. We equip $\mathbb{P}(P \cap \widehat{L})$ with the induced orientation, where P_L is equipped with the following orientation. A tangent vector v to P_L is positively oriented if there is a vector $w \in P$ such that the directional derivative $\partial_w Q$ is positive, and vector $x \in P$ such that $\langle x, e_{n+1} \rangle_{n+1,1}$ is positive and w, v, x is an oriented basis for P. Equip \mathbb{RP}^1 with the orientation induced by viewing it as the boundary of the hemisphere in \mathbb{CP}^1 containing $[1:i]$. Let $\tilde{u}_P^{\mathbb{R}} : \mathbb{RP}^1 \to \mathbb{P}(P \cap \hat{L})$ be an orientation preserving algebraic isomorphism. So, $\tilde{u}_P^{\mathbb{R}}$ is uniquely determined by P up to reparameterization by $PSL_2(\mathbb{R})$. Let $\tilde{u}_P : \mathbb{CP}^1 \to \mathbb{P}(P^{\mathbb{C}} \cap \widehat{X})$ be the complexification of $\tilde{u}_P^{\mathbb{R}}$. Let

$$
\psi:D\to\mathbb{CP}^1
$$

be given by

$$
\psi(z) = [iz + i : z - 1], \quad z \in \mathbb{C}, |z| \leq 1.
$$

Let $u_P : (D, \partial D) \to (X, L)$ be the holomorphic map given by $u_P = \tilde{u}_P \circ \psi$.

Let $Gr(2, n+2)^{(2,1)}$ denote the Grassmanian of oriented subspaces $P \subset \mathbb{R}^{n+2}$ as above. Let $SO(n+1,1)_+$ denote the group of linear automorphisms $l : \mathbb{R}^{n+2} \to \mathbb{R}^{n+2}$ preserving the bilinear form $\langle \cdot, \cdot \rangle_{n+1,1}$ and satisfying

$$
\det l = 1, \qquad \langle l(e_{n+1}), e_{n+1} \rangle_{n+1,1} > 0.
$$

The group $SO(n+1,1)_+$ is connected. It acts on $Gr(2, n+2)^{(2,1)}$ and this action is transitive for $n > 1$. Moreover, $SO(n+1,1)_+$ acts on (X, L) by holomorphic symplectomorphisms and thus also acts on the moduli spaces $\mathcal{M}_{k,l}(\beta)$.

Lemma A.6. The map $Gr(2, n+2)^{(2,1)} \rightarrow \mathcal{M}_{3,0}(\beta_1)$ given by $P \mapsto u_P$ is an $SO(n+1,1)_+$ equivariant diffeomorphism.

Then, as the map $\mathcal{M}_{3,0}(\beta_1) \to L^3$ is also $SO(n+1,1)_+$ equivariant, and $SO(n+1,1)_+$ is connected (and hence acts in an orientation preserving way), we find that the sign of the evaluation map is independent of the choice of map.

A.3. Proof of Lemma [A.5.](#page-38-1) Decomposing (u^*TX, u^*TL) into line bundles.

Let

$$
P_0 = \text{Span}\{e_0, e_1, e_{n+1}\}.
$$

We compute u_{P_0} explicitly. Throughout, we shorten the subscript P_0 to 0. P_0 has associated map $u_0 : (D, \partial D) \longrightarrow (X, L)$ given by:

$$
u_0(z) = [z^2 + 1 : i(1 - z^2) : 0 : \cdots : 0 : 2z].
$$

After normalisation, the restriction to ∂D is given by:

$$
u_0(\theta) = [\cos \theta : \sin \theta : 0 : \cdots : 1]
$$

As the map u_0 is non-singular, the derivative $Du(\frac{\partial}{\partial z})$ defines an nowhere vanishing holomorphic section of u_0^*TX , call this ξ_1 . We will now extend this to a basis of sections of u_0^*TX . To this end, for $j = 2, \ldots, n$, let ξ_j be the vector field given by $(0, 0, \ldots, 0, \frac{1}{z})$ $(\frac{1}{z},0,\ldots,0)=\frac{1}{z}\partial_{x_j}$ in the affine chart $z_{n+1} \neq 0$, with coordinates x_0, \ldots, x_n . We then have:

Lemma A.7. The vector fields ξ_j extend to holomorphic sections $D \to u_0^*TX$. Moreover, they are nowhere vanishing.

Proof. The unique extension can be computed by instead going to the affine patch $z_0 \neq 0$. Using the coordinates y_1, \ldots, y_{n+1} in this patch, we find that $\xi_j = \frac{2}{1+i}$ $\frac{2}{1+z^2}\partial_{y_j}$. Thus, ξ_j extends uniquely over the apparent pole at $z = 0$, and is indeed nowhere vanishing.

Let (E_j, F_j) be the line sub-bundle of (u_0^*TX, u_0^*TL) generated by ξ_j . That is, we set $E_j = span_{\mathbb{C}} \xi_j$, and the real boundary condition is given by $F_j := E_j \cap u_0^*TL$.

Lemma A.8. The map

$$
\varphi: \mathcal{E}^{\oplus n} \to (u_0^*TX, u_0^*TL)
$$

given by

$$
\varphi_z(\lambda_1,\ldots,\lambda_n)=-i\lambda_1\xi_1+\lambda_2\xi_2+\cdots+\lambda_n\xi_n
$$

is an isomorphism.

Proof. Note first that if the map is well-defined, it is an isomorphism, as the sections ξ_j are linearly independent (and thus form a basis). It thus remains to be shown that the map respects the real boundary conditions, and is thus well-defined. For the first component, we do the computation in the affine patch $\{z_{n+1} = 1\}$. The map $u_0 : D \setminus \{0\} \to Q \subset \mathbb{C}^{n+1}$ is given by

$$
u_0(z) = (\frac{z^2+1}{2z}, i\frac{1-z^2}{2z}, 0, \dots, 0).
$$

One can check that indeed, if $|z|=1$, the quantities $\frac{z^2+1}{2z}$ $\frac{2+1}{2z}$, and $i\frac{1-z^2}{2z}$ $\frac{-z^2}{2z}$ are both real numbers. In fact, if $z = e^{i\theta}$ they are $\cos \theta$ and $\sin \theta$ respectively. The derivative can be computed as

$$
Du(\frac{\partial}{\partial z}) = (\frac{z^2 - 1}{2z^2}, -i\frac{1 + z^2}{2z^2}, 0, \dots, 0)
$$

Note that this lies indeed in the tangent bundle of Q, which is given by those $(p, v) \in$ $\mathbb{C}^{n+1} \oplus \mathbb{C}^{n+1}$ with $p \cdot v = 0$, where \cdot denotes the complex linear extension of the dot product on \mathbb{R}^{n+1} . Then, to show that φ_1 respects the real boundary conditions, note that on the left, it is given by $F_z = z \mathbb{R}$. On the right it is given by $span_{\mathbb{C}} \xi_1 \cap u_0^* TL$. Thus, let $\lambda_1 = \alpha_1 z$ for $\alpha_1 \in \mathbb{R}$. We compute

$$
-i\alpha_1 z \xi_1 = \alpha_1 (i \frac{1-z^2}{2z}, \frac{1+z^2}{2z}, 0, \ldots, 0),
$$

and, indeed, as noted before, these two quantities are real numbers, so that $-i\alpha_1 z \xi_1 \in u_0^* T L$ as required. Next, for $j \geq 2$ we have

$$
\xi_j = (0, 0, \dots, \frac{1}{z}, 0, \dots, 0)
$$

Observe that if $\lambda_j = \alpha_j z$ for $\alpha_j \in \mathbb{R}$, we have $\alpha_j z \xi_j = (0, \ldots, 0, \alpha_j, 0, \ldots, 0) \in u_0^*TL$ as required. \Box

We are now ready to compute the effect on spin structures. The spin structure on u_0^*TL is the unique spin structure obtained as follows: take the loop $u_0|_{\partial D} \subset L$. As L is simply connected, one can choose a disk D in L bounding this loop. The bundle $TL|_D$ is trivial (unique up to homotopy) as D is contractible. Restricting this trivialisation to ∂D yields a trivialisation of $u_0|_{\partial D}^*TL$. Note that as $w_2(L) = 0$, so any two choices of D yield homotopic trivialisations of $u_0|_{\partial D}^*TL$. Now equip the resulting trivial bundle u_0^*TL bundle with the trivial spin structure (i.e. the one that in this trivialisation is just the trivial $spin(n)$) bundle over $S¹$. We will explicitly write down such a trivialisation.

Lemma A.9. The natural trivialisation of u_0^*TL coming from a disk bounding $u_0|_{\partial D}$ has a framing homotopic to the one given by $\{\psi_i\}_{i=1,\dots,n}$ with

$$
\psi_1 = (\sin^2 \theta, -\cos \theta \sin \theta, \cos \theta, 0, \dots, 0)
$$

$$
\psi_2 = (-\cos \theta \sin \theta, \cos^2 \theta, \sin \theta, 0, \dots, 0)
$$

$$
\psi_j = (0, 0, 0, \dots, 0, 1, 0, \dots, 0)
$$
 for $j \ge 3$

Here ψ_i has a non-zero entry only in entry $j+1$. The vector fields are expressed by viewing $L \subset \mathbb{R}^{n+1}$ as the unit sphere.

Proof. Using the same notation as in the rest of the section, the coordinates on \mathbb{R}^{n+1} are x_0, \ldots, x_n . Let the disk $\mathbb D$ be given by $\{x_0^2 + x_1^2 + x_2^2 = 1, x_2 \ge 0 | x_i \in \mathbb R\}$. Identify $\mathbb D$ with D using stereographic projection. Note that we have $u_0|_{S^1}^*TL \simeq T\mathbb{D}|_{\partial\mathbb{D}} \oplus \mathbb{R}^{n-2}$, where the \mathbb{R}^{n-2} factor is given by the vector fields ψ_j for $j ≥ 3$. Now, trivialise TD with the standard unit vector fields ∂_x , ∂_y . Pulling these back under stereographic projection and restricting to the boundary yields the vector fields ψ_1 and ψ_2 .

Next, we look at the trivialisation induced by the isomorphism $(u_0^*TX, u_0^*TL) \simeq \mathcal{E}^{\oplus n}$. The boundary condition for $\mathcal E$ is given by $F_z = z\mathbb R$, and is trivialised by division by z. The spin structure is again given by taking the trivial spin structure with respect to this trivialisation. We have:

Lemma A.10. The trivialisation of u_0^*TL coming from the isomorphism with $\mathcal{E}^{\oplus n}$ has a framing homotopic to the one given by $\{\chi_j\}_{j=1,\dots,n}$ with

$$
\chi_1 = (\sin \theta, -\cos \theta, 0, 0, \dots, 0) \n\chi_j = (0, \dots, 0, 1, 0, \dots, 0) \text{ for } j \ge 2
$$

where χ_i has a non-zero entry only in position $j + 1$.

Proof. The sections χ_j are induced by the sections ξ_j from before. Concretely, recall that

$$
\xi_1 = (\frac{z^2 - 1}{z^2}, -i\frac{1 + z^2}{z^2}, 0, \dots, 0).
$$

The morphism $\varphi_1: F_z \to u_0^*TL$ is given by $\alpha z \mapsto -i\alpha z \xi_1$, which after substituting $z = e^{i\theta}$ indeed yields χ_1 . For $j \geq 2$ the morphism $\varphi_j : F_z \to u_0^*TL$ is given by $\alpha z \mapsto \alpha \xi_j =$ $\alpha(0,\ldots,0,1,0,\ldots,0)$ thus the induced section is indeed χ_i . .

Finally, we have:

Lemma A.11. The trivialisations induced by the basis ψ_j is not homotopic to the one induced by χ_j .

Proof. The change of basis matrix is given by

$$
\begin{pmatrix}\n\sin \theta & \cos \theta & 0 \\
-\cos \theta & \sin \theta & 0 \\
0 & 0 & I_{n-2}\n\end{pmatrix}
$$

which indeed defines a non-trivial loop in $SO(n)$.

$$
\overrightarrow{42}
$$

The conclusion is that the induced spin structures are opposed.

A.4. Proof of Proposition [A.1.](#page-37-0) Computing the sign of Ev .

It is known that global holomorphic sections of $\mathcal E$ over $(D, \partial D)$ are described as follows:

$$
\Gamma(\mathcal{E}) = \{a_0 + a_1 z + \bar{a}_0 z^2 \mid a_1 \in \mathbb{R}\}.
$$

Each copy of $\mathcal{E} \subset T_u \mathcal{M}_{3,0}(1)$ gives rise to maps

$$
evb^1 : \Gamma(\mathcal{E}) \longrightarrow F_1 \simeq \mathbb{R},
$$

\n $evi^0 : \Gamma(\mathcal{E}) \longrightarrow E_0 \simeq \mathbb{C},$

given by evaluation at $1 \in \partial D$ and $0 \in int(D)$, respectively. Lemma A.2.1 of [\[10\]](#page-43-14) says that in fact this determines an orientation-preserving isomorphism

$$
\theta = evb^1 \oplus evi^0 : \Gamma(\mathcal{E}) \stackrel{\sim}{\longrightarrow} \mathbb{R} \oplus \mathbb{C}.
$$

Using θ , we find that the following sections form a positively oriented basis for $\Gamma(\mathcal{E})$, when taken in this order:

$$
\xi^1(z) = 1 - 2z + z^2,
$$

$$
\xi^2(z) = i - iz^2,
$$

$$
\xi^3(z) = z.
$$

Evaluating ξ^j at 1, *i*, and -1, we see that

$$
\begin{pmatrix} \xi^1(1) & \xi^2(1) & \xi^3(1) \\ \xi^1(i) & \xi^2(i) & \xi^3(i) \\ \xi^1(-1) & \xi^2(-1) & \xi^3(-1) \end{pmatrix} = \begin{pmatrix} 0 & 0 & 1 \\ -2i & 2i & i \\ 4 & 0 & -1 \end{pmatrix}.
$$

By definition, the fibres of the boundary condition F of $\mathcal E$ at $z = 1, i, -1$, are, respectively

$$
F_1 = \mathbb{R}, \quad F_i = i\mathbb{R}, \quad F_{-1} = -\mathbb{R}.
$$

Thus,

$$
evb:\Gamma(\mathcal{E})\overset{\sim}{\longrightarrow}\mathbb{R}\oplus i\mathbb{R}\oplus-\mathbb{R}
$$

is represented by the matrix

$$
M = \begin{pmatrix} 0 & 0 & 1 \\ -2 & 2 & 1 \\ -4 & 0 & 1 \end{pmatrix},
$$

which has determinant

$$
\det M = 1 \cdot \begin{vmatrix} -2 & 2 \\ -4 & 0 \end{vmatrix} = 8 > 0.
$$

So, Ev is represented a block-diagonal matrix where the diagonal consists of n copies of the block M and satisfies

$$
sgn(Ev) = sgn(\det M)^n = 1.
$$

REFERENCES

- [1] A. Beauville, Quantum cohomology of complete intersections, Mat. Fiz. Anal. Geom. 2 (1995), no. 3-4, 384–398.
- [2] K. Cieliebak and E. Goldstein, A note on the mean curvature, Maslov class and symplectic area of Lagrangian immersions, J. Symplectic Geom. 2 (2004), no. 2, 261–266.
- [3] B. Crauder and R. Miranda, Quantum cohomology of rational surfaces, The moduli space of curves (Texel Island, 1994), Progr. Math., vol. 129, Birkhäuser Boston, Boston, MA, 1995, pp. 33–80, [doi:10.1007/978-1-4612-4264-2_3](http://dx.doi.org/10.1007/978-1-4612-4264-2_3).
- [4] W. Fulton and R. Pandharipande, Notes on stable maps and quantum cohomology, Algebraic geometry— Santa Cruz 1995, Proc. Sympos. Pure Math., vol. 62, Amer. Math. Soc., Providence, RI, 1997, pp. 45–96, [doi:10.1090/pspum/062.2/1492534](http://dx.doi.org/10.1090/pspum/062.2/1492534).
- [5] M. Kontsevich and Y. Manin, Gromov-Witten classes, quantum cohomology, and enumerative geometry, Comm. Math. Phys. 164 (1994), no. 3, 525–562.
- [6] W. Lerche, C. Vafa, and N. P. Warner, *Chiral rings in* $N = 2$ superconformal theories, Nuclear Phys. B 324 (1989), no. 2, 427–474, [doi:10.1016/0550-3213\(89\)90474-4](http://dx.doi.org/10.1016/0550-3213(89)90474-4).
- [7] D. McDuff and D. Salamon, J-holomorphic curves and symplectic topology, second ed., American Mathematical Society Colloquium Publications, vol. 52, American Mathematical Society, Providence, RI, 2012.
- [8] Y. Ruan and G. Tian, A mathematical theory of quantum cohomology, Math. Res. Lett. 1 (1994), no. 2, 269–278.
- [9] Y. Ruan and G. Tian, A mathematical theory of quantum cohomology, J. Differential Geom. 42 (1995), no. 2, 259–367.
- [10] J. Smith, Discrete and continuous symmetries in monotone Floer theory, Selecta Math. (N.S.) 26 (2020), no. 3, Paper No. 47, 65, [doi:10.1007/s00029-020-00575-5](http://dx.doi.org/10.1007/s00029-020-00575-5).
- [11] J. P. Solomon, Intersection theory on the moduli space of holomorphic curves with Lagrangian boundary conditions, arXiv e-prints (2006) , [arXiv:math/0606429](http://arxiv.org/abs/math/0606429).
- [12] J. P. Solomon and S. B. Tukachinsky, Relative quantum cohomology, to appear in J. Eur. Math. Soc., [doi:10.4171/JEMS/1337](http://dx.doi.org/10.4171/JEMS/1337).
- [13] J. P. Solomon and S. B. Tukachinsky, Point-like bounding chains in open Gromov-Witten theory, Geom. Funct. Anal. 31 (2021), no. 5, 1245–1320, [doi:10.1007/s00039-021-00583-3](http://dx.doi.org/10.1007/s00039-021-00583-3).
- [14] E. Witten, Topological sigma models, Comm. Math. Phys. 118 (1988), no. 3, 411–449.

SCHOOL OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, BAILRIGG, LANCASTER, LA1 4YW, UK

Email address: kai.hugtenburg@gmail.com

School of Mathematical Sciences, Tel Aviv University, Tel Aviv, 6997801, Israel Email address: sarabt1@gmail.com