

# ON THE RINGEL–HALL ALGEBRA OF THE GENTLE ONE-CYCLE ALGEBRA $\Lambda(n-1, 1, 1)$

HUI CHEN, AND DONG YANG

ABSTRACT. It is shown that the gentle one-cycle algebra  $\Lambda(n-1, 1, 1)$  has Hall polynomials. The Hall polynomials are explicitly given for all triples of indecomposable modules, and as a consequence, the Ringel–Hall Lie algebra of  $\Lambda(n-1, 1, 1)$  is shown to be isomorphic to its Riedtmann Lie algebra.

Key words: Ringel–Hall algebra, Hall polynomial, gentle one-cycle algebra  
MSC2020: 16G20, 17B37.

## 1. INTRODUCTION

Let  $A$  be a finite-dimensional algebra over a finite field  $\mathbb{F}_q$ . The *Ringel–Hall algebra*  $\mathcal{H}(A)$  of  $A$  is the  $\mathbb{Z}$ -algebra with basis the set of isomorphism classes of finite-dimensional left  $A$ -modules and the corresponding structure constants given by filtration numbers. Assume that  $A$  is representation-finite, i.e., there are only finitely many isomorphism classes of finite-dimensional  $A$ -modules. Then  $A$  is said to have Hall polynomials if, roughly speaking, the filtration numbers are the evaluation at  $q$  of some integral polynomials. In this case, we can view  $q$  as an indeterminate and evaluate  $\mathcal{H}(A)$  at  $q = 1$ . Then the  $\mathbb{Z}$ -submodule  $\mathfrak{g}(A)$  of  $\mathcal{H}(A)$  spanned by the isoclasses of indecomposable  $A$ -modules is a Lie subalgebra. In this way, the positive parts of finite-dimensional simple complex Lie algebras can be constructed from the representation theory of Dynkin quivers. See [8, 9, 10].

It is proved in [8] that Hall polynomials exist for representation-finite algebras with direct Auslander–Reiten quiver and in general this is an open problem. In this paper we consider the gentle one-cycle algebra  $A$  given by the quiver with relation

$$1 \longrightarrow 2 \longrightarrow \cdots \longrightarrow n \circlearrowleft \alpha, \quad \alpha^2,$$

whose Auslander–Reiten quiver is not directed. Over an algebraically closed field this is the derived-discrete algebra  $\Lambda(n-1, 1, 1)$  in the notation of [1]. Our first main result is

**Theorem 1.1** (Theorem 3.1).  *$A$  has Hall polynomials.*

Thus there is the Lie subalgebra  $\mathfrak{g}(A)$  of  $\mathcal{H}(A)$  (evaluated at  $q = 1$ ) of indecomposable  $A$ -modules. On the other hand, following Riedtmann’s approach [7] to consider the representation theory of  $A$  over  $\mathbb{C}$ , we can obtain another Lie algebra  $L(A_{\mathbb{C}})$ , see [4]. Our second main result is

**Theorem 1.2** (Theorem 3.4).  *$\mathfrak{g}(A)$  and  $L(A_{\mathbb{C}})$  are isomorphic Lie algebras.*

This is done by brutal force, namely, we compute the Hall polynomials for all triples of indecomposable  $A$ -modules (Proposition 3.2), and as a consequence we obtain the structure constants of  $\mathfrak{g}(A)$  (Corollary 3.3), which are the same as those of  $L(A_{\mathbb{C}})$  given in [4].

**Acknowledgement.** The authors are very grateful to Bangming Deng and Haicheng Zhang for answering their questions. The second author acknowledges support by the National Natural Science Foundation of China No.12031007.

## 2. PRELIMINARIES

In this section we recall some basic results on Ringel–Hall algebras and on the classification of indecomposable modules over the algebra  $\Lambda(n-1, 1, 1)$ .

**2.1. Ringel–Hall algebras.** Let  $K = \mathbb{F}_q$  be a finite field, and  $A$  be an associative unital finite-dimensional representation-finite  $K$ -algebra. By an  $A$ -module we will mean a finite-dimensional left  $A$ -module. For an  $A$ -module  $M$  we denote by  $[M]$  its isomorphism class. We denote by  $\mathcal{P}(A)$  the set of all isomorphism classes of  $A$ -modules.

For three  $A$ -modules  $N_1, N_2$  and  $M$ , let  $F_{N_1 N_2}^M$  be the *Hall number*, i.e., the number of submodule  $U$  of  $M$  such that  $U \cong N_2$  and  $M/U \cong N_1$ . Let  $\mathcal{H}(A)$  be the free  $\mathbb{Z}$ -module with basis  $\mathcal{P}(A)$ , and for any  $[N_1], [N_2], [M] \in \mathcal{P}(A)$  define

$$[N_1] \diamond [N_2] = \sum_{[M] \in \mathcal{P}(A)} F_{N_1 N_2}^M [M].$$

**Proposition 2.1** ([9]).  $\mathcal{H}(A)$  is an associative  $\mathbb{Z}$ -algebra with unit  $[0]$ , the isoclass of the zero module.

The algebra  $\mathcal{H}(A)$  is called the *Ringel–Hall algebra* of  $A$ . The following lemma is an immediate consequence of [5, Lemma 3.5] (see also [6, Lemma 2.1]).

**Lemma 2.2.** Let  $X, X_1, X_2, X_3, X_4, Y, M$  be  $A$ -modules (possibly 0),  $a, b \in \mathbb{C}$  and assume that  $[X] = a[X_1] \diamond [X_2] + b[X_3] \diamond [X_4]$  in  $\mathcal{H}(A)$ , then

$$F_{XY}^M = a \sum_{Z \in \mathcal{P}(A)} F_{X_1 Z}^M F_{X_2 Y}^Z + b \sum_{Z \in \mathcal{P}(A)} F_{X_3 Z}^M F_{X_4 Y}^Z.$$

Given an extension field  $E$  of  $K$  and a  $K$ -space  $V$ , put  $V^E := V \otimes_K E$ . According to [10], a field extension  $E$  of  $K$  is said to be *conservative* for the  $K$ -algebra  $A$  if the algebra  $(\text{End} M / \text{rad End } M)^E$  is a field for any indecomposable  $A$ -module  $M$ .

**Definition 2.3** ([10, Section 3]). We say that  $A$  has *Hall polynomials* provided that for all  $A$ -modules  $X, Y, M$ , there exists a polynomial  $\varphi_{XY}^M \in \mathbb{Z}[T]$  such that for any conservative field extension  $E$  of  $K$ , we have

$$\varphi_{XY}^M(|E|) = F_{X^E Y^E}^{M^E}.$$

Such a  $\varphi_{XY}^M$  is called a *Hall polynomial*.

For the existence of Hall polynomials, there are the following two general results:

**Lemma 2.4.** *Let  $S$  be a simple  $A$ -module. Then the Hall polynomials  $\varphi_{SN}^M$  and  $\varphi_{NS}^M$  exist for any  $A$ -modules  $M$  and  $N$ .*

**Lemma 2.5.** *Let  $P$  be a projective  $A$ -module. Then the Hall polynomial  $\varphi_{PN}^M$  exists for any  $A$ -modules  $M$  and  $N$ .*

*Proof.* Since  $P$  is a projective  $A$ -module, the short exact sequence

$$0 \rightarrow N \rightarrow M \rightarrow P \rightarrow 0$$

is always split. By [6, Lemma 2.7], the Hall polynomial  $\varphi_{PN}^M$  exists.  $\square$

**Theorem 2.6** ([6, Theorem 2.9]). *Assume that  $A$  is representation-finite. Then  $A$  has Hall polynomials if and only if the Hall polynomial  $\varphi_{XY}^M$  exists for any indecomposable  $A$ -module  $X$  and any  $A$ -modules  $Y$  and  $M$ .*

If  $A$  has Hall polynomials, then the *degenerate Ringel–Hall algebra*  $\mathcal{H}(A)_1$  is defined as the  $\mathbb{Z}$ -algebra with basis  $\mathcal{P}(A)$  and with multiplication

$$[N_1][N_2] = \sum_{[M]} \varphi_{N_1 N_2}^M(1)[M].$$

We denote by  $\mathfrak{g}(A)$  the  $\mathbb{Z}$ -submodule of  $\mathcal{H}(A)_1$  with basis  $[M]$ , where  $M$  runs over all indecomposable  $A$ -modules.

The following theorem is part of [10, Theorem],

**Theorem 2.7.**  *$\mathfrak{g}(A)$  is a Lie subalgebra of  $\mathcal{H}(A)_1$ , and  $\mathcal{H}(A)_1 \otimes_{\mathbb{Z}} \mathbb{Q}$  is isomorphic as a  $\mathbb{Q}$ -algebra to the universal enveloping algebra of  $\mathfrak{g}(A) \otimes_{\mathbb{Z}} \mathbb{Q}$ .*

**2.2. Representations of the algebra  $\Lambda(n-1, 1, 1)$ .** Let  $K$  be an arbitrary field. Consider the bound quiver  $(Q, I)$

$$1 \longrightarrow 2 \longrightarrow \cdots \longrightarrow n \curvearrowright_{\alpha} \alpha, \quad \alpha^2,$$

and denote the path algebra of this bound quiver by  $A_K$ . When  $K$  is an algebraically closed field, this is the derived-discrete algebra  $\Lambda(n-1, 1, 1)$  in the notation of [1]. We will identify an  $A_K$ -module with a representation of  $(Q, I)$ , i.e., a representation of  $Q$  satisfying the relation  $\alpha^2 = 0$ . The following theorem was established as [2, Theorem 3.3] for  $K = \mathbb{C}$ , but it holds true for any field  $K$  by for example [3, Section 3, Theorem].

**Theorem 2.8.** *The following representations  $U_{i,j}$ ,  $V_i$  and  $W_{i,j}$  form a complete set of representatives of the isoclasses of indecomposable  $A_K$ -modules:*

- $U_{i,j}$  for  $1 \leq j \leq i \leq n$ :

$$0 \xrightarrow{0} \cdots \xrightarrow{0} 0 \xrightarrow{0} K \xrightarrow{1} \cdots \xrightarrow{1} K \xrightarrow{e_1} K^2 \xrightarrow{1} \cdots \xrightarrow{1} K^2 \curvearrowright_{M_\alpha} M_\alpha$$

$$\begin{array}{ccccccc}
 \bullet & \longrightarrow & \cdots & \longrightarrow & \bullet & \longrightarrow & \cdots & \longrightarrow & \bullet \\
 j & & & & i & & & & n \\
 & & & & \bullet & \longrightarrow & \cdots & \longrightarrow & \bullet \\
 & & & & & & & & \bullet
 \end{array}$$

- $U_{i,j}$  for  $1 \leq i < j \leq n$ :

$$\begin{array}{cccccccccccc}
 0 & \xrightarrow{0} & \cdots & \xrightarrow{0} & 0 & \xrightarrow{0} & K & \xrightarrow{1} & \cdots & \xrightarrow{1} & K & \xrightarrow{e_2} & K^2 & \xrightarrow{I} & \cdots & \xrightarrow{I} & K^2 & \Big)_{M_\alpha} \\
 & & & & & & & & & & & & \bullet & \xrightarrow{\quad} & \cdots & \xrightarrow{\quad} & \bullet & & \\
 & & & & & & i & & & & & & j & & & & n & & \\
 & & & & & & \bullet & \xrightarrow{\quad} & \cdots & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \cdots & \xrightarrow{\quad} & \bullet & \Big)_{M_\alpha}
 \end{array}$$

- $V_i$  for  $1 \leq i \leq n$ :

$$\begin{array}{cccccccc}
 0 & \xrightarrow{0} & \cdots & \xrightarrow{0} & 0 & \xrightarrow{0} & K & \xrightarrow{1} & \cdots & \xrightarrow{1} & K & \Big)_{0} \\
 & & & & & & i & & & & n & \\
 & & & & & & \bullet & \xrightarrow{\quad} & \cdots & \xrightarrow{\quad} & \bullet & 
 \end{array}$$

- $W_{i,j}$  for  $1 \leq i \leq j < n$ :

$$\begin{array}{cccccccccccc}
 0 & \xrightarrow{0} & \cdots & \xrightarrow{0} & 0 & \xrightarrow{0} & K & \xrightarrow{1} & \cdots & \xrightarrow{1} & K & \xrightarrow{0} & 0 & \xrightarrow{0} & \cdots & \xrightarrow{0} & 0 & \Big)_{0} \\
 & & & & & & i & & & & j & & & & & & & \\
 & & & & & & \bullet & \xrightarrow{\quad} & \cdots & \xrightarrow{\quad} & \bullet & & & & & & & 
 \end{array}$$

Here,  $e_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ,  $e_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ,  $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ , and  $M_\alpha = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ .

In particular,  $A_K$  is representation-finite. The simple  $A_K$ -modules are  $S_1 = W_{1,1}, \dots, S_{n-1} = W_{n-1,n-1}$  and  $S_n = V_n$ . Moreover,  $P_1 = U_{n,1}, \dots, P_n = U_{n,n}$  (respectively,  $I_1 = W_{1,1}, \dots, I_{n-1} = W_{1,n-1}, I_n = U_{1,n}$ ) form a complete set of pairwise non-isomorphic indecomposable projective (respectively, injective)  $A_K$ -modules.

### 3. THE RINGEL–HALL ALGEBRA OF $\Lambda(n-1, 1, 1)$

In this section, we study the Ringel–Hall algebra of the algebra  $\Lambda(n-1, 1, 1)$ . More precisely, we show the existence of Hall polynomials, compute the Hall polynomials  $\varphi_{XY}^M$  for all triples  $(X, Y, M)$  of indecomposable modules and describe the structure constants of the Ringel–Hall Lie algebra.

**3.1. The existence of Hall polynomials.** Fix a finite field  $\mathbb{F}_q$  and put  $A = A_{\mathbb{F}_q}$  (see Section 2.2). The main result of this subsection is

**Theorem 3.1.** *The algebra  $A$  has Hall polynomials.*

*Proof.* By Theorem 2.6, we only need to prove that for any indecomposable  $A$ -module  $X$  the Hall polynomial  $\varphi_{XY}^M$  exists for any  $A$ -modules  $Y$  and  $M$ .

Case 1:  $X = W_{i,j}$ ,  $1 \leq i \leq j \leq n-1$ . For  $j = i$ , this follows by Lemma 2.4, since  $W_{i,i} = S_i$  is a simple module. If  $j > i$ , we have

$$[W_{i,i}] \diamond [W_{i+1,j}] = [W_{i,i} \oplus W_{i+1,j}] + [W_{i,j}], \text{ and } [W_{i+1,j}] \diamond [W_{i,i}] = [W_{i,i} \oplus W_{i+1,j}],$$

so

$$[W_{i,j}] = [W_{i,i}] \diamond [W_{i+1,j}] - [W_{i+1,j}] \diamond [W_{i,i}].$$

Therefore it follows by Lemma 2.2 that for any  $A$ -modules  $Y$  and  $M$ ,

$$F_{W_{i,j}Y}^M = \sum_{Z \in \mathcal{P}(A)} F_{W_{i,i}Z}^M F_{W_{i+1,j}Y}^Z - \sum_{Z \in \mathcal{P}(A)} F_{W_{i+1,j}Z}^M F_{W_{i,i}Y}^Z.$$

Hence by induction the Hall polynomial  $\varphi_{XY}^M$  exists for  $X = W_{i,j}$ ,  $1 \leq i \leq j \leq n-1$ .

Case 2:  $X = V_i$ ,  $1 \leq i \leq n$ . For  $i = n$  this follows by Lemma 2.4, since  $V_n = S_n$  is a simple module. For  $1 \leq i \leq n-1$ , we have

$$[W_{i,n-1}] \diamond [S_n] = [W_{i,n-1} \oplus S_n] + [V_i], \text{ and } [S_n] \diamond [W_{i,n-1}] = [W_{i,n-1} \oplus S_n],$$

so

$$[V_i] = [W_{i,n-1}] \diamond [S_n] - [S_n] \diamond [W_{i,n-1}].$$

Therefore it follows by Lemma 2.2 that for any  $A$ -modules  $Y$  and  $M$ ,

$$F_{V_iY}^M = \sum_{Z \in \mathcal{P}(A)} F_{W_{i,n-1}Z}^M F_{S_nY}^Z - \sum_{Z \in \mathcal{P}(A)} F_{S_nZ}^M F_{W_{i,n-1}Y}^Z.$$

Hence by Case 1 the Hall polynomial  $\varphi_{XY}^M$  exists for  $X = V_i$ ,  $1 \leq i \leq n$ .

Case 3:  $X = U_{i,j}$ ,  $1 \leq j \leq i \leq n$ . For  $i = n$  this follows by Lemma 2.5, since  $U_{n,j} = P_j$  is a projective module. For  $1 \leq j \leq i \leq n-1$ , we have

$$[W_{i,n-1}] \diamond [P_j] = q[W_{i,n-1} \oplus P_j] + q[U_{i,j}], \text{ and } [P_j] \diamond [W_{i,n-1}] = [W_{i,n-1} \oplus P_j],$$

so

$$[U_{i,j}] = q^{-1}[W_{i,n-1}] \diamond [P_j] - [P_j] \diamond [W_{i,n-1}].$$

Therefore it follows by Lemma 2.2 that for any  $A$ -modules  $Y$  and  $M$ ,

$$F_{U_{i,j}Y}^M = q^{-1} \sum_{Z \in \mathcal{P}(A)} F_{W_{i,n-1}Z}^M F_{P_jY}^Z - \sum_{Z \in \mathcal{P}(A)} F_{P_jZ}^M F_{W_{i,n-1}Y}^Z.$$

Hence by Case 2, there exists a rational function  $\varphi_{XY}^M$  such that  $F_{U_{i,j}Y}^M = \varphi_{XY}^M(q)$ . Since this is an integer for any  $q$ , it follows by [8, Section 2, Lemma] that  $\varphi_{XY}^M$  is a polynomial. Namely, the Hall polynomial  $\varphi_{XY}^M$  exists for  $X = U_{i,j}$ ,  $1 \leq j \leq i \leq n$ .

Case 4:  $X = U_{i,j}$ ,  $1 \leq i < j \leq n$ . We have

$$[V_i] \diamond [V_j] = q[V_i \oplus V_j] + [U_{j,i}], \text{ and } [V_j] \diamond [V_i] = [V_i \oplus V_j] + [U_{i,j}],$$

so

$$[U_{i,j}] = [V_j] \diamond [V_i] + q^{-1}[U_{j,i}] - q^{-1}[V_i] \diamond [V_j].$$

Therefore it follows by Lemma 2.2 that for any  $A$ -modules  $Y$  and  $M$ ,

$$F_{U_{i,j}Y}^M = \sum_{Z \in \mathcal{P}(A)} F_{V_jZ}^M F_{V_iY}^Z + q^{-1}F_{U_{j,i}Y}^M - q^{-1} \sum_{Z \in \mathcal{P}(A)} F_{V_iZ}^M F_{V_jY}^Z.$$

Hence by Case 2 and Case 3, there exists a rational function  $\varphi_{XY}^M$  such that  $F_{U_{i,j}Y}^M = \varphi_{XY}^M(q)$ . Then we can show that the Hall polynomial  $\varphi_{XY}^M$  exists as in Case 3.  $\square$

**3.2. Hall polynomials for indecomposable modules.** In this subsection we explicitly give the Hall polynomial  $\varphi_{XY}^M$  for indecomposable  $A$ -modules  $X, Y, M$ .

**Proposition 3.2.** *Let  $X, Y, M$  be indecomposable  $A$ -modules. Then the Hall polynomial  $\varphi_{XY}^M$  is 0 except in the following cases:*

- $\varphi_{XY}^M = T$  if the triple  $(X, Y, Z)$  is one of the following:
  - (1)  $(W_{i,j}, U_{j+1,l}, U_{i,l})$  for  $i \leq l \leq j < n$ ,
  - (2)  $(W_{i,j}, U_{j+1,j}, U_{i,j})$  for  $i < j < n$ .
- $\varphi_{XY}^M = 1$  if the triple  $(X, Y, Z)$  is one of the following:
  - (3)  $(W_{i,j}, W_{j+1,l}, W_{i,l})$  for  $i \leq j \leq l < n$ ,
  - (4)  $(W_{i,j}, V_{j+1}, V_i)$  for  $i \leq j < n$ ,
  - (5)  $(W_{i,j}, U_{j+1,l}, U_{i,l})$  for  $i \leq j < j+1 < l \leq n$ ,
  - (6)  $(W_{i,j}, U_{l,j+1}, U_{l,i})$  for  $i \leq j < j+1 \leq l \leq n$ ,
  - (7)  $(W_{i,j}, U_{l,j+1,l}, U_{l,i})$  for  $l < i \leq j < n$ ,
  - (8)  $(W_{i,j}, U_{l,j+1}, U_{l,i})$  for  $i < l \leq j < n$ ,
  - (9)  $(W_{i,j}, U_{j+1,l}, U_{i,l})$  for  $i \leq j < n$ ,
  - (10)  $(W_{i,j}, U_{i,j+1}, U_{i,i})$  for  $i \leq j < n$ ,
  - (11)  $(V_i, V_j, U_{i,j})$  for  $1 \leq i, j \leq n$ .

*Proof.* We only prove (1),(5),(11), and the proof of the other cases is similar.

Case (1): For  $i \leq l \leq j < n$ , it is clear that  $\underline{\dim} W_{i,j} + \underline{\dim} U_{j+1,l} = \underline{\dim} U_{i,l}$ , and from the structure of  $W_{i,j}$ ,  $U_{j+1,l}$  and  $U_{i,l}$  in Section 2.2,  $U_{i,l}$  has  $q$  submodules isomorphic to  $U_{j+1,l}$  with quotient isomorphic to  $W_{i,j}$ , which are:

$$0 \xrightarrow{0} \cdots \xrightarrow{0} 0 \xrightarrow{0} K\{e_1 + ce_2\} \xrightarrow{1} \cdots \xrightarrow{1} K\{e_1 + ce_2\} \hookrightarrow K^2 \xrightarrow{1} \cdots \xrightarrow{1} K^2 \bigg)_{M_\alpha}$$

$$\qquad \qquad \qquad l \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad j+1 \qquad \qquad \qquad n$$

where  $c \in K = \mathbb{F}_q$ .

Case (5): For  $i \leq j < j+1 < l \leq n$ , it is clear that  $\underline{\dim} W_{i,j} + \underline{\dim} U_{j+1,l} = \underline{\dim} U_{i,l}$ , and from the structure of  $W_{i,j}$ ,  $U_{j+1,l}$  and  $U_{i,l}$  in Section 2.2,  $U_{i,l}$  has a unique submodule isomorphic to  $U_{j+1,l}$  with quotient isomorphic to  $W_{i,j}$ .

Case (11): For  $1 \leq i, j \leq n$ , it is clear that  $\underline{\dim} V_i + \underline{\dim} V_j = \underline{\dim} U_{i,j}$ , and from the structure of  $V_i$ ,  $V_j$  and  $U_{i,j}$  in Section 2.2,  $U_{i,j}$  has a unique submodule isomorphic to  $V_j$  with quotient isomorphic to  $V_i$ .  $\square$

**3.3. The Ringel–Hall Lie algebra.** We have shown in Theorem 3.1 that the algebra  $A$  has Hall polynomials. As in the end of Section 2.1, let  $\mathfrak{g}(A)$  be the  $\mathbb{Z}$ -submodule of the degenerate Ringel–Hall algebra  $\mathcal{H}(A)_1$  with basis  $[M]$ , where  $M$  runs over all indecomposable  $A$ -modules. The following is a corollary of Theorem 2.7 and Proposition 3.2:

**Corollary 3.3.**  *$\mathfrak{g}(A)$  is a Lie subalgebra of  $\mathcal{H}(A)_1$  with the following Lie bracket:*

- (a)  $[W_{i,j}, W_{l,m}] = \delta_{j+1,l}W_{i,m} - \delta_{m+1,i}W_{l,j}$  for  $1 \leq i \leq j \leq n-1$  and  $1 \leq l \leq m \leq n-1$ ,
- (b)  $[W_{i,j}, V_l] = \delta_{j+1,l}V_i$  for  $1 \leq i \leq j \leq n-1$  and  $1 \leq l \leq n$ ,
- (c)  $[W_{i,j}, U_{l,m}] = \delta_{j+1,m}U_{l,i} + \delta_{j+1,l}U_{i,m}$  for  $1 \leq i \leq j \leq n$  and  $1 \leq l, m \leq n$ ,
- (d)  $[V_i, V_j] = U_{j,i} - U_{i,j}$  for  $1 \leq i, j \leq n$ ,

where  $\delta_{i,j}$  is the Kronecker symbol. For other pairs of indecomposable modules, the Lie bracket of them equals 0.

Using Riedtmann's construction [7], we have obtained in [4] a Lie algebra  $L(A_{\mathbb{C}})$  from the representation theory of  $A_{\mathbb{C}}$ , which is isomorphic to a Lie algebra of type  $\mathbf{BC}^+$ . By Corollary 3.3 and [4, Proposition 4.2], we have

**Theorem 3.4.** *There is an isomorphism  $\mathfrak{g}(A) \cong L(A_{\mathbb{C}})$  of Lie algebras.*

#### REFERENCES

- [1] Grzegorz Bobiński, Christof Geiß, and Andrzej Skowroński, *Classification of discrete derived categories*, Cent. Eur. J. Math. **2** (2004), no. 1, 19–49 (electronic).
- [2] Magdalena Boos and Markus Reineke, *B-orbits of 2-nilpotent matrices and generalizations*, Highlights in Lie algebraic methods, Progr. Math., vol. 295, Birkhäuser/Springer, New York, 2012, pp. 147–166.
- [3] M. C. R. Butler and Claus Michael Ringel, *Auslander-Reiten sequences with few middle terms and applications to string algebras*, Comm. Algebra **15** (1987), no. 1-2, 145–179. MR 876976 (88a:16055)
- [4] Hui Chen and Dong Yang, *On Riedtmann's Lie algebra of the gentle one-cycle algebra  $\Lambda(n-1, 1, 1)$* , Communications in Algebra **52** (2024), no. 1, 345–358.
- [5] Stanislaw Kasjan and Justyna Kosakowska, *The existence of Hall polynomials for  $x^2$ -bounded invariant subspaces of nilpotent linear operators*, J. Pure Appl. Algebra **226** (2022), no. 5, Paper No. 106921, 12.
- [6] A. R. Nasr-Isfahani, *Hall polynomials for Nakayama algebras*, Algebr. Represent. Theory **15** (2012), no. 3, 483–490.
- [7] Christine Riedtmann, *Lie algebras generated by indecomposables*, J. Algebra **170** (1994), no. 2, 526–546.
- [8] Claus Michael Ringel, *Hall algebras*, Topics in algebra, Part 1 (Warsaw, 1988), Banach Center Publ., vol. 26, PWN, Warsaw, 1990, pp. 433–447.
- [9] ———, *Hall algebras and quantum groups*, Invent. Math. **101** (1990), no. 3, 583–591.
- [10] Claus Michael Ringel, *Lie algebras arising in representation theory*, Representations of algebras and related topics (Kyoto, 1990), London Math. Soc. Lecture Note Ser., vol. 168, Cambridge Univ. Press, Cambridge, 1992, pp. 284–291.

HUI CHEN, SCHOOL OF BIOMEDICAL ENGINEERING AND INFORMATICS, NANJING MEDICAL UNIVERSITY, NANJING 211166, P. R. CHINA.

*Email address:* huichen@njmu.edu.cn

DONG YANG, DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, P. R. CHINA

*Email address:* yangdong@nju.edu.cn