A CONCISE PROOF OF THE STABLE MODEL STRUCTURE ON SYMMETRIC SPECTRA

CARY MALKIEWICH AND MARU SARAZOLA

ABSTRACT. It is well-known that the stable model structure on symmetric spectra cannot be transferred from the one on sequential spectra through the forgetful functor. We use the fibrant transfer theorem of [GMSV23] to show it can be transferred between fibrant objects, providing a new, short and conceptual proof of its existence.

1. Introduction

Symmetric spectra provide one of the most general and flexible frameworks for stable homotopy theory, as a symmetric monoidal model category. There are other model categories with good smash products, including orthogonal spectra from [MMSS01] and the S-modules of [EKMM97], but symmetric spectra are more general than these, because they only use symmetric groups in their definition. This allows us to define them in settings where a topological enrichment is not naturally present, such as the motivic setting [Hov01].

However, the theory of symmetric spectra has a famous drawback: the stable equivalences cannot be defined as the maps that induce isomorphisms on the homotopy groups (the π_* -isomorphisms), at least if we give the homotopy groups their most obvious definition. As a result, the proof of the stable model structure for symmetric spectra in [HSS00] and [MMSS01] requires additional concepts and intermediate results beyond those needed to establish the same model structure for sequential spectra (prespectra) or orthogonal spectra. This makes the theory somewhat less "user friendly" and presents a significant tradeoff compared to the other models.

A natural attempt to avoid these complications is to use the transfer theorem (see for instance [Cra95, Section 3] or the dual of [HKRS17, Theorem 2.2.1]) to produce the model structure from the adjunction relating symmetric spectra to sequential spectra,

$$\operatorname{Sp}^{\Sigma} \xrightarrow{I} \operatorname{Sp}^{\mathbb{N}}_{\operatorname{st}}.$$

If successful, this would construct a model structure on $\operatorname{Sp}^{\Sigma}$ in which a map f is a weak equivalence if and only if Uf is a π_* -isomorphism in $\operatorname{Sp}^{\mathbb{N}}_{\operatorname{st}}$. Sadly, this approach cannot succeed, because it produces the wrong class of weak equivalences: we would get the maps inducing isomorphisms on homotopy groups, rather than the maps inducing isomorphisms on generalized cohomology. These weak equivalences are "wrong" because with them the adjunction (F,U) is not a Quillen equivalence, and so we get the wrong homotopy category on the left-hand side.

Upon inspection, we see that the discrepancy between the classes of weak equivalences vanishes once we restrict to maps between fibrant objects. This suggests that one could use a more flexible version of the transfer theorem introduced recently in [GMSV23], where weak equivalences are transferred through the right adjoint only between fibrant objects, leaving room for equivalences between arbitrary objects to admit a more complicated description.

The goal of this note is to prove that this approach works, providing a short proof of the existence of the stable model structure on $\mathrm{Sp}_{\mathrm{st}}^\Sigma$. Notably, it allows us to completely avoid a treatment—and even a definition of—stable equivalences between non-fibrant objects, and only rely on well-known facts about π_* -isomorphisms in $\mathrm{Sp}_{\mathrm{st}}^\mathbb{N}$.

2. Review of model structures on spectra

In what follows, we let $\mathrm{Sp}^{\mathbb{N}}$ denote the category of sequential spectra based on simplicial sets. These are sequences of pointed simplicial sets X_n with bonding maps $X_n \wedge S^1 \to X_n$, where S^1 denotes the simplicial circle. We similarly let Sp^{Σ} denote the corresponding category of symmetric spectra, in which the levels X_n have actions by the symmetric group, and the bonding maps respect these actions in the sense of [HSS00, Def 1.2.2].

We briefly recall some of the model structures that can be defined on these categories. The first two are "level" model structures, in which the weak equivalences are maps that are equivalences at each spectrum level separately.

Theorem 2.1. [BF78, Proposition 2.2] There is a level model structure on sequential spectra $\operatorname{Sp}_l^{\mathbb{N}}$, in which weak equivalences, fibrations, and trivial fibrations are defined as the levelwise corresponding maps on $\operatorname{sSet}_{*,\operatorname{Quillen}}$.

Theorem 2.2. [HSS00, Theorem 5.1.2] There is a level model structure on symmetric spectra $\operatorname{Sp}_l^{\Sigma}$, in which weak equivalences, fibrations, and trivial fibrations are defined as the levelwise corresponding maps on $\operatorname{sSet}_{*,\operatorname{Quillen}}$.

Of course, the levelwise equivalences are not the equivalences we want in stable homotopy theory—we want a broader class of maps called stable equivalences. For sequential spectra, these are the same thing as π_* -isomorphisms.

Theorem 2.3. [BF78, Proposition 2.3] There is a stable model structure on sequential spectra $\operatorname{Sp}_{\operatorname{st}}^{\mathbb{N}}$, in which:

- weak equivalences are the π_* -isomorphisms; i.e. the maps inducing isomorphisms on all homotopy groups,
- fibrant objects are the Ω -spectra $(X_n \to \Omega X_{n+1} \text{ is an equivalence})$ which are levelwise Kan complexes, and
- trivial fibrations are the level trivial fibrations.

Theorem 2.4. [HSS00, Theorem 3.4.4] There is a stable model structure on symmetric spectra $\operatorname{Sp}_{\operatorname{st}}^{\Sigma}$, in which:

- weak equivalences are the stable equivalences; i.e. the maps $f: X \to Y$ such that $f^*: \operatorname{Map}(X, E) \to \operatorname{Map}(Y, E)$ is a weak equivalence of simplicial sets for every injective Ω -spectrum E, and
- ullet fibrant objects and trivial fibrations agree with those of $\operatorname{Sp}_{\operatorname{st}}^{\mathbb{N}}$.

All of these model structures are cofibrantly generated.

Remark 2.5. The definition of the stable equivalences for symmetric spectra requires the notion of an *injective* spectrum ([HSS00, Definition 3.1.1]), which is not recalled here. This is an intentional omission: the injectivity condition can be dropped when working between fibrant objects, which is all that is needed in our proof.

Remark 2.6. Once the stable model structures are constructed, one can use the adjunction between spectra of topological spaces and spectra of simplicial sets

$$\operatorname{Sp}_{\operatorname{Top}} \xrightarrow{\longleftarrow \square} \operatorname{Sp}.$$

By the classical transfer theorem, we get stable model structures on the categories of spectra based on topological spaces, $\mathrm{Sp}^{\mathbb{N}}_{\mathrm{Top}}$ and $\mathrm{Sp}^{\Sigma}_{\mathrm{Top}}$. As expected, each of these Topbased model structures is Quillen equivalent to its simplicial counterpart.

Sequential and symmetric spectra are related by a free-forgetful adjunction [HSS00, Thm 4.3.2]

$$\operatorname{Sp}^{\Sigma} \xrightarrow{\longleftarrow F} \operatorname{Sp}^{\mathbb{N}}$$

which can be used to transfer the level model structure on sequential spectra to obtain the level model structure on symmetric spectra. A natural question is then: is it possible to transfer the stable model structure through the free-forgetful adjunction as well? Unfortunately this strategy is not fruitful.

Remark 2.7. As observed by Hovey–Shipley–Smith in [HSS00, Section 3.1], the model structure on $\mathrm{Sp}_{\mathrm{st}}^{\Sigma}$ cannot be transferred through the free-forgetful adjunction

$$\operatorname{Sp}^{\Sigma} \xrightarrow{I} \operatorname{Sp}^{\mathbb{N}}_{\operatorname{st}}$$

Indeed, the functor U does not preserve weak equivalences, as illustrated in [HSS00, Example 3.1.10]. However, if we work only between fibrant objects, we find that a map f is a stable equivalence if and only if Uf is [HSS00, Corollary 4.3.4].

3. Existence of the model structure

Remark 2.7 suggests that the model structure on $\mathrm{Sp}_{\mathrm{st}}^{\Sigma}$ can be constructed using the notion of fibrant transfer introduced in [GMSV23]. In this section we recall the statement of this fibrant transfer theorem, then use it to construct the model structure on $\mathrm{Sp}_{\mathrm{st}}^{\Sigma}$.

Theorem 3.1. [GMSV23, Theorem 3.5] Let $(\mathcal{M}, \text{Cof}, \text{Fib}, \mathcal{W})$ be a combinatorial model category with generating set of trivial cofibrations \mathcal{K} , and let \mathcal{C} be a locally presentable category. Suppose that we have an adjunction

$$\mathcal{C} \xrightarrow{F} \mathcal{M}$$

and that the following properties are satisfied:

(1) for every map $f: X \to Y$ such that $Uf \in \text{Fib} \cap \mathcal{W}$, there is a commutative square

$$X \xrightarrow{f} Y$$

$$\iota_X \downarrow \qquad \qquad \downarrow \iota_Y$$

$$X' \xrightarrow{f'} Y'$$

with UX', UY' fibrant, $\iota_X, \iota_Y \in \operatorname{cof}(FK)$, and such that $Uf' \in \mathcal{W}$;

(2) for every object X such that UX is fibrant, there is a factorization of the diagonal morphism

$$X \xrightarrow{w} \operatorname{Path} X \xrightarrow{p} X \times X$$

with $Uw \in \mathcal{W}$ and $Up \in \text{Fib}$.

Then, there exists a combinatorial model structure on C in which a morphism f is a trivial fibration if and only if Uf is so in \mathcal{M} , and an object X is fibrant if and only if UX is fibrant in \mathcal{M} . Moreover, a morphism f between fibrant objects in C is a weak equivalence (resp. fibration) if and only if Uf is so in \mathcal{M} .

Theorem 3.2. The stable model structure on Sp^Σ can be fibrantly transferred through the free-forgetful adjunction

$$\mathrm{Sp}^{\Sigma} \xrightarrow{\stackrel{F}{\longleftarrow}} \mathrm{Sp}^{\mathbb{N}}_{\mathrm{st}}.$$

Proof. The categories of sequential and symmetric spectra are locally presentable, as we defined them from the locally presentable category of pointed simplicial sets. Moreover, the stable model structure $\mathrm{Sp}^{\mathbb{N}}_{\mathrm{st}}$ is cofibrantly generated; let \mathcal{K} denote the set of generating trivial cofibrations. We verify that conditions (1) and (2) of Theorem 3.1 hold.

For condition (1), let $f: X \to Y$ be a map of symmetric spectra such that Uf is a level trivial fibration. If Fib denotes the class of fibrations in $\operatorname{Sp}_{\operatorname{st}}^{\mathbb{N}}$, then the adjunction $F \dashv U$ gives a weak factorization system

$$(\operatorname{cof}(F\mathcal{K}), U^{-1}(\operatorname{Fib})).$$

We can then consider a factorization of the map to the terminal spectrum

$$X \xrightarrow{\iota_X} X' \xrightarrow{p} 1$$

with $\iota_X \in \operatorname{cof}(F\mathcal{K})$ and $Up \in \operatorname{Fib}$; in particular, UX' is fibrant. Taking the pushout of ι_X along f

$$X \xrightarrow{f} Y$$

$$\iota_X \downarrow \qquad \qquad \downarrow j$$

$$X' \xrightarrow{g} X' \cup_X Y$$

produces a map $j \in \operatorname{cof}(F\mathcal{K})$ (as the left class of a factorization system is always closed under pushouts). Note that maps in $\operatorname{cof}(F\mathcal{K})$ are levelwise monomorphisms – it follows that the map g is at each spectrum level a pushout of a weak equivalence along a cofibration of simplicial sets, and is therefore a weak equivalence. Since X' is an Ω -spectrum, then so is $X' \cup_X Y$ as these are preserved by level equivalences. However, $X' \cup_X Y$ may not be levelwise Kan.

To remedy this, we post-compose the diagram above with a fibrant replacement

$$i \colon X' \cup_X Y \stackrel{\sim}{\longrightarrow} \overline{X' \cup_X Y}$$

in the level model structure on $\operatorname{Sp}^{\Sigma}$. Letting $\mathcal{J} \subseteq \mathcal{K}$ denote the set of generating trivial cofibrations in $\operatorname{Sp}_l^{\Sigma}$ and $F\mathcal{J}$ the corresponding generating trivial cofibrations in $\operatorname{Sp}_l^{\Sigma}$, we have $F\mathcal{J} \subseteq F\mathcal{K}$, so the map $i \in \operatorname{cof}(F\mathcal{J})$ above is in fact also in $\operatorname{cof}(F\mathcal{K})$; hence, so is the composite ij. The spectrum $\overline{X' \cup_X Y}$, which is levelwise Kan by construction, is also an Ω -spectrum as i is a level weak equivalence. Finally, the map U(ig) is a level weak equivalence, and therefore a π_* -isomorphism. This proves condition (1).

To check condition (2), let X be a symmetric spectrum such that UX is fibrant. The cylinder object in sSet

$$\Delta[0] \sqcup \Delta[0] \hookrightarrow \Delta[1] \xrightarrow{\sim} \Delta[0]$$

yields the required path object in $\mathrm{Sp}^\Sigma_{\mathrm{st}}$

$$X \cong X^{\Delta[0]} \xrightarrow{\sim} X^{\Delta[1]} \twoheadrightarrow X^{\Delta[0] \sqcup \Delta[0]} \cong X \times X$$

To see this we observe that the cotensor $(-)^A$ commutes with the forgetful functor U, and that in $\mathrm{Sp}_{\mathrm{st}}^{\mathbb{N}}$, the above two maps are a weak equivalence and a fibration, respectively, because the model structure on $\mathrm{Sp}_{\mathrm{st}}^{\mathbb{N}}$ is simplicial [BF78, Theorem 2.3].

By Theorem 3.1, this produces a model structure on Sp^Σ in which a spectrum X is fibrant if and only if UX is fibrant in $\mathrm{Sp}^\mathbb{N}_{\mathrm{st}}$ (that is, X is an Ω -spectrum and levelwise Kan), and a map f is a trivial fibration if and only if Uf is a trivial fibration in $\mathrm{Sp}^\mathbb{N}_{\mathrm{st}}$ (that is, f is a level trivial fibration). Since fibrant objects and trivial fibrations uniquely determine the model structure (see [Joy08, Theorem 51.10]), we recover the stable model structure $\mathrm{Sp}^\Sigma_{\mathrm{st}}$ of Theorem 2.4 as desired. Moreover, Theorem 3.1 yields a description of the weak equivalences between fibrant objects: these are precisely the π_* -isomorphisms. \square

References

- [BF78] A. K. Bousfield and E. M. Friedlander. Homotopy theory of Γ-spaces, spectra, and bisimplicial sets. In *Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II*, volume 658 of *Lecture Notes in Math.*, pages 80–130. Springer, Berlin, 1978.
- [Cra95] Sjoerd E. Crans. Quillen closed model structures for sheaves. J. Pure Appl. Algebra, 101(1):35–57, 1995.
- [EKMM97] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May. Rings, modules, and algebras in stable homotopy theory, volume 47 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1997. With an appendix by M. Cole.
- [GMSV23] Léonard Guetta, Lyne Moser, Maru Sarazola, and Paula Verdugo. Fibrantly-induced model structures. arXiv:2301.07801, 2023.
- [HKRS17] Kathryn Hess, Magdalena Kędziorek, Emily Riehl, and Brooke Shipley. A necessary and sufficient condition for induced model structures. J. Topol., 10(2):324–369, 2017.

- [Hov01] Mark Hovey. Spectra and symmetric spectra in general model categories. *Journal of Pure and Applied Algebra*, 165(1):63–127, 2001.
- [HSS00] Mark Hovey, Brooke Shipley, and Jeff Smith. Symmetric spectra. J. Amer. Math. Soc., 13(1):149–208, 2000.
- $[{\rm Joy08}]$ André Joyal. Notes on logoi. http://www.math.uchicago.edu/ may/IMA/JOYAL/Joyal.pdf, 2008.
- [MMSS01] M. A. Mandell, J. P. May, S. Schwede, and B. Shipley. Model categories of diagram spectra. *Proc. London Math. Soc. (3)*, 82(2):441–512, 2001.

Department of Mathematics, Binghamton University, Binghamton, NY 13902, USA $\it Email~address: malkiewich@math.binghamton.edu$

School of Mathematics, University of Minnesota, Minneapolis MN, 55415, USA $\it Email~address:$ $\tt maru@umn.edu$