

A novel two-field pure K-essence for inflation, dark matter, dark energy and black holes

Changjun Gao

*National Astronomical Observatories, Chinese Academy of Sciences, 20A Datun Road, Beijing 100101, China and
School of Astronomy and Space Sciences, University of Chinese Academy of Sciences, 19A Yuquan Road, Beijing 100049, China*

K-essence theories are usually studied in the framework of one scalar field ϕ . Namely, the Lagrangian of K-essence is the function of scalar field ϕ and its covariant derivative. However, in this paper, we explore a two-field pure K-essence, i.e. the corresponding Lagrangian is the function of covariant derivatives of two scalar fields without the dependency of scalar fields themselves. That is why we call it pure K-essence. The novelty of this K-essence is that its Lagrangian contains the quotient term of the kinetic energies from the two scalar fields. This results in the presence of many interesting features, for example, the equation of state can be arbitrarily small and arbitrarily large. As a comparison, the range for equation of state of quintessence is from -1 to $+1$. Interestingly, this novel K-essence can play the role of inflation field, dark matter and dark energy. Finally, the absence of the scalar fields themselves in the equations of motion makes the study considerable simple such that even the exact black hole solutions can be found.

PACS numbers: 04.20.Jb

I. INTRODUCTION

K-essence is a certain class of scalar field models with nonstandard kinetic terms. The name was first coined in a paradigm for inflation by Armendariz-Picon et al. [1, 2]. Subsequently, it was found that k-essence can present interesting models for dark energy [3–9]. Right after these studies, Scherrer shows that K-essence can also serve as the unified model for dark matter and dark energy [10]. Afterwards, researches on the K-essence theories spring up.

Specifically, Chimonto and Lazkoz make an analysis on the atypical K-essence cosmologies [11]. Abramo et al. conduct the research on the stability problem of phantom k-essence theories [12]. Bonvin et al. [13] demonstrate that if one use K-essence to solve the coincidence problem and play the role of dark energy in the Universe, the fluctuations of the field have to propagate superluminally at some stage. This implies that successful K-essence models violate causality. Therefore, k-essence cannot arise as a low energy effective field theory of a causal, consistent high energy theory. Bamba [14] et al carry out a thorough investigation on the cosmological perturbations of K-essence. The causality and superluminal behavior K-essence theories are investigated by Bruneton [15]. The slow-roll conditions for thawing K-essence are derived by Chiba et al. with a separable Lagrangian K-essence [16]. Considering the unification between the swampland criteria and the attractor, Herrera studies the reconstruction of the K-essence inflationary universe [17]. Furthermore, K-essence can yield a complete picture of the evolution of the Universe, starting from the early inflation, the subsequent dark matter domination, and to the late time cos-

mic acceleration [18, 19]. Interestingly, Tian and Zhu's research [20] reveals that k-essence framework is also conducive to the Hubble tension problem [21, 22].

Very recently, Hung and Miao [23] investigate whether a quantum-induced k-essence can provide efficient reheating without affecting the observational constraints on primordial inflation. Kehayias and Scherrer find a new generic evolution for K-essence dark energy with the equation of state (EOS) of roughly cosmological constant [24]. Sarkar et al [25] show that the accretion of k-essence dark energy can lead to a faster circularising of orbits for binary black holes. Lara et al [26] investigate in depth the relation between K-essence, well-posedness of the Cauchy problem and the UV completions. Huang [27] constructs the thawing K-essence models by generating Taylor expansion coefficients of Lagrangian functions from random matrices. Chatterjee et al. make a study on the dynamical stability problem for k-essence interacting nonminimally with a perfect fluid [28]. Applying the “ $2 + 1 + 1$ ” decomposition of spacetime, Nagy et al conduct the research of spherically symmetric, static black holes with scalar hair, and naked singularities in nonminimally coupled K-essence [29]. Barvinsky et al. find that the generalized unimodular gravity can be transformed into a new form of K-essence [30]. Finally, the K-essence model in which K-essence can be responsible for both primordial inflation and the present observed acceleration, while also admitting a nonsingular de Sitter beginning of the Universe is achieved by Ferreira et al. [31].

Up to this point, what we need to emphasize is that all the above researches on K-essence are actually one-field K-essence. In specific, the Lagrangian of K-essence is the function of single field ϕ and its derivatives $\partial_\mu \phi$. How-

ever, in this paper, we shall consider two-field K-essence among which there are two scalar fields ϕ and ψ together with their derivatives. The novelty of this research is that the Lagrangian functions are significantly different from the previous ones. Exactly, they are constructed from the quotient form “potentials”, x/y with x and y the kinetic term of the fields. To our knowledge, one has never made an investigation on this sort of K-essence. On the other hand, what we consider is a pure K-essence which makes the calculation considerable simple. In fact, we can get the energy density, the pressure and the equation of state (EOS) straightforward because the equations of motion of K-essence can be exactly solved without the help of dynamical analysis. Although it seems our two-field pure K-essence can be transformed to the multi-scalar tensor theories, the equivalence is merely mathematically, not physically at least from the standpoint of speed of perturbations. The reason is that the resulting metric of field-space is not the function of dynamical field, but the auxiliary scalar fields. It is not the case for conventional multi-scalar tensor theories [32].

The paper is organized as follows. In Section II, we derive the Einstein equations and the equations of motion for the two-field pure K-essence. In Section III, we study the cosmic evolution of the K-essence in the background of Friedmann-Robertson-Walker Universe. In the first place, the Lagrangian functions of K-essence for some conventional EOS are reconstructed. Then motivated by the Lagrangian of K-essence form for the conventional EOS, we explore the so-called novel K-essence Lagrangian which has the quotient “potentials”. In this way, we obtain several unconventional EOS of K-essence. In Section IV, we transform the two-field pure K-essence to the multi-field form with resort to auxiliary scalar fields. We point that the two frameworks are merely equivalent to each other mathematically, not physically, at least from the point of view of speed of perturbations. In Section V, as the demonstration again for the simplicity of novel two-field pure K-essence, we give a black hole solution. Finally, Section VI gives the conclusion and discussion. Throughout this paper, we adopt the system of units in which $G = c = \hbar = 1$ and the metric signature $(-, +, +, +)$.

II. EQUATIONS OF MOTION

We consider the action

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{16\pi} + F + L_m \right], \quad (1)$$

where

$$F = F(x, y), \quad (2)$$

is the Lagrangian of two-field pure K-essence. Following Ref. [10] we call it pure K-essence because it does not depend explicitly on the two scalar fields, ϕ and ψ . The

Lagrangian is only the function of terms, x and y . Here x and y are the kinetic energy of scalar fields ϕ and ψ , respectively. They are defined by

$$x \equiv -\frac{1}{2} \nabla_\mu \phi \nabla^\mu \phi, \quad y \equiv -\frac{1}{2} \nabla_\mu \psi \nabla^\mu \psi. \quad (3)$$

R is the Ricci scalar and L_m is the Lagrangian for matters. Make variation of the action with respect to the metric tensor, we obtain the Einstein equations

$$G_{\mu\nu} = 8\pi (T_{\mu\nu}^k + T_{\mu\nu}^m), \quad (4)$$

where $T_{\mu\nu}^m$ and $T_{\mu\nu}^k$ represent the energy-momentum of matters and K-essence, respectively. The expression of $T_{\mu\nu}^k$ is found to be

$$T_{\mu\nu}^k = F_{,x} \nabla_\mu \phi \nabla_\nu \phi + F_{,y} \nabla_\mu \psi \nabla_\nu \psi + g_{\mu\nu} F, \quad (5)$$

with the comma denotes the derivative with respect to x and y , respectively. When $F = \text{constant}$, we obtain the energy-momentum tensor for the Einstein cosmological constant. When $F = -2x$ or $F = -2y$, we obtain the energy-momentum tensor for stiff matter. Finally, the variation of the action with respect to ϕ and ψ give their equations of motion as follows

$$\nabla_\mu [F_{,x} \nabla^\mu \phi] = 0, \quad \nabla_\mu [F_{,y} \nabla^\mu \psi] = 0. \quad (6)$$

III. COSMIC EVOLUTION

In this section, we explore the cosmic evolution of the two-field pure K-essence in the background of Friedmann-Robertson-Walker Universe. To this end, we should first derive the cosmological equations of motion.

A. cosmological equations of motion

In the background of Friedmann-Robertson-Walker universe with the line element

$$ds^2 = -dt^2 + a(t)^2 (dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2), \quad (7)$$

where $a(t)$ is the scale factor of the Universe. The Einstein equations appears as the Friedmann equation and the acceleration equation

$$3H^2 = 8\pi \left(\frac{\rho_{m0}}{a^3} + \frac{\rho_{r0}}{a^4} + \rho_K \right), \quad (8)$$

$$2\dot{H} + 3H^2 = -8\pi \left(\frac{\rho_{r0}}{3a^4} + p_K \right). \quad (9)$$

Here the energy density and pressure of K-essence are

$$\rho_K = -2xF_{,x} - 2yF_{,y} + F, \quad p_K = -F. \quad (10)$$

H is the Hubble parameter and the dot denotes the derivative with respect to the cosmic time t . ρ_{m0} and ρ_{r0} are the present-day energy density for dark matter and radiation. The equations of motion for the two scalar fields are

$$\left(a^3 F_{,x} \dot{\phi}\right)' = 0, \quad \left(a^3 F_{,y} \dot{\psi}\right)' = 0. \quad (11)$$

Eqs. (8,9,10,11) constitute the cosmological equations of motion. Observing the expressions of density and pressure, we find they are the functions of Lagrangian F . So once the equation of state $p_K = p_K(\rho_K)$ for K-essence is given, we can obtain the expression of F by solving the partial differential equations. In the next subsection, we shall make an investigation on three conventional EOSs by this way.

B. Three conventional equation-of-states

The so-called three conventional EOSs (equation-of-states) are for the Einstein cosmological constant, the constant EOS and the generalized Chaplygin gas.

1. dynamical cosmological constant

If we assume the EOS for K-essence is that for cosmological constant, taking into account Eqs. (10) and solving the differential equation

$$\omega \equiv \frac{p_K}{\rho_K} = -1, \quad (12)$$

we find

$$F = f\left(\frac{x}{y}\right). \quad (13)$$

Here and thereafter $f\left(\frac{x}{y}\right)$ stands for the arbitrary function of x/y . It is interesting that in this case, although the energy density and pressure of K-essence are constants, the scalar fields ϕ and ψ are dynamic:

$$\dot{\phi} \sim a^3, \quad \dot{\psi} \sim a^3, \quad (14)$$

for example, when

$$F = s \frac{x}{y}, \quad (15)$$

with s a constant. This has been pointed by us in Ref. [33]. It is also straightforward to obtain the Schwarzschild-de Sitter solution

$$ds^2 = -\left(1 - \frac{2M}{r} - \frac{sc_2^2 r^2}{6c_1^2}\right) dt^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2 + \left(1 - \frac{2M}{r} - \frac{sc_2^2 r^2}{6c_1^2}\right)^{-1} dr^2, \quad (16)$$

in this case. Here c_1 and c_2 are two integration constants and M is the mass of black hole. The expressions of two scalar fields are

$$\phi = \frac{2sc_2^2 r^3}{3c_1^3}, \quad \psi = -\frac{2sc_2 r^3}{3c_1^2}. \quad (17)$$

Then the energy density and pressure in this static spacetime which are the same as in the Friedmann-Robertson-Walker spacetime, remain constants. We emphasize that this is a coincidence. In general, for example in the background of anisotropic and inhomogeneous spacetime, the model of Eq. (13) or Eq. (15) would significantly differ from the Einstein cosmological constant because their energy-momentum tensors are obviously different.

2. constant equation of state

If we assume the equation of state for K-essence is a constant ω , taking into account Eqs. (10) and solving the equation

$$\omega = \frac{p_K}{\rho_K}, \quad (18)$$

we find

$$F = f\left(\frac{x}{y}\right) x^{\frac{1+\omega}{2\omega}}. \quad (19)$$

In this case, the energy density and pressure of K-essence are

$$\rho_K = \frac{\rho_{K0}}{a^{3(\omega+1)}}, \quad p_K = \omega \rho_K. \quad (20)$$

It seems the Lagrangian Eq. (19) is divergent in the limit of $\omega = 0$ (for dark matter). This is an illusion; if we parameterize the Lagrangian as below,

$$F = \frac{x^{\frac{1}{2\omega}}}{e^{\frac{1}{\omega^2}}} \sqrt{y}. \quad (21)$$

there is no divergence in the expression when $\omega = 0$.

3. Generalized Chaplygin gas

If we assume the equation of state for K-essence is that for generalized Chaplygin gas [34], taking into account Eqs. (10) and solving the equation

$$\rho_K = \frac{A}{(-p_K)^{n-1}}, \quad (22)$$

we obtain

$$F = \left[A + x^{\frac{n}{2}} f\left(\frac{x}{y}\right) \right]^{\frac{1}{n}}, \quad (23)$$

where A and n are positive constants. Observations constrains $n > 1$ in order to ensure the condition of stability and the condition of causality.

Up to this point, we conclude that the two-field pure K-essence is very powerful since it can play the role of Einstein cosmological constant, the fluid with a constant EOS and the generalized Chaplygin gas in the background of Friedmann-Robertson-Walker Universe. Of course, in the background of other spacetime, the behavior of two-field pure K-essence is certainly different from the others because they have different energy-momentum tensors. Observing the Lagrangians Eq. (13), Eq. (19) and Eq. (23), we find they are all the functions of quotient x/y . This motivates us, in the next section, to make an investigation on the cosmic evolution of K-essence by starting from a given expression of F and several unconventional EOSs are found.

C. several unconventional EOSs

In this section, we shall first assume the expression of Lagrangian function F and then explore its cosmic evolution. By this way, some novel EOSs are found.

1. The case of $F = 2y + s\frac{\sqrt{x}}{y^{2n}}$

In the first place, we consider the Lagrangian function

$$F = 2y + s\frac{\sqrt{x}}{y^{2n}}, \quad (24)$$

where s and n are constants. We note that the sign before the kinetic term is positive. This indicates ϕ is a phantom scalar field. Then the cosmological equations of motion, Eqs. (10,11) give the density and pressure of K-essence as follows

$$\rho_K = b_0 a^{\frac{3(1-4n)}{4n}} + b_1 a^{\frac{3}{2n}}, \quad (25)$$

$$p_K = -\frac{b_0}{4n} a^{\frac{3(1-4n)}{4n}} - \frac{b_1(2n+1)}{2n} a^{\frac{3}{2n}}, \quad (26)$$

where b_0 and b_1 are positive integration constants. The equation of state is

$$\omega = -\frac{\frac{b_0}{4n} a^{\frac{3(1-4n)}{4n}} + \frac{b_1(2n+1)}{2n} a^{\frac{3}{2n}}}{b_0 a^{\frac{3(1-4n)}{4n}} + b_1 a^{\frac{3}{2n}}}. \quad (27)$$

It is interesting that for large n , we have the energy density

$$\rho_K = b_0 a^{-3} + b_1, \quad (28)$$

which is the sum of dark matter and cosmological constant. Thus, when n is sufficiently large, the K-essence

unifies the cold dark matter and the Einstein cosmological constant. When $n = 1/4$, we have

$$\rho_K = b_0 + b_1 a^6. \quad (29)$$

It consists of two terms among them one is the Einstein cosmological constant and the other is a phantom. In this case, the K-essence can play the role of dark energy for present-day cosmic acceleration.

In Fig. (1) we plot the equation of state w with respect to the scale factor. In order that the energy density is always positive, we assume the two integration constants $b_0 > 0$ and $b_1 > 0$. The figure shows that when $n = -1/4$, the K-essence behaves as the stiff matter which has the equation of state $+1$. However, it is amazing that its Lagrangian function $F = 2y + s\sqrt{xy}$ is obviously different from that for conventional stiff matter $F = -2y$. If $n = 1/4$, the K-essence can play the role of cosmic dark energy as mentioned earlier. On the other hand, if $|n| \rightarrow \infty$, the K-essence unifies the Einstein cosmological constant and the cold dark matter model. Finally, when $n \approx 0$, the equation of state can be very large such that $w \rightarrow +\infty$. Now the K-essence becomes the stiffest matter in nature.




FIG. 1: The evolution of EOS for K-essence with respect to the scale factor when $n = 0.17, 0.25, 0.5, 2$ from down to top, respectively for the lower half of the graph. For the upper half of the graph, we put $n = -0.14, -0.18, -0.25, -0.5, -2$ from up to down. We assume $b_0 = b_1 = 1$.

2. The case of $F = s_1\sqrt{x} + s_2\sqrt{y} + s\left(\frac{x}{y}\right)^{\frac{n}{2}}$

Secondly, we consider the Lagrangian function

$$F = s_1\sqrt{x} + s_2\sqrt{y} + s\left(\frac{x}{y}\right)^{\frac{n}{2}}, \quad (30)$$

where s_1, s_2, s and n are constants. We recognize that s_1 and s_2 terms are actually the kinetic terms of Cuscu-

ton action [35]. The cosmological equations of motion, Eqs. (10,11) give the energy density

$$\rho_K = -\frac{s(c_2 - s_2 a^3)^n}{(c_1 - s_1 a^3)^n}, \quad (31)$$

where c_1 and c_2 are integration constants. It is straightforward to obtain the pressure p_K and equation of state ω for K-Essence by substituting the energy density into the energy conservation equation

$$\frac{d\rho_K}{dt} + 3H(\rho_K + p_K) = 0. \quad (32)$$

In order that the energy density is always positive, we let

$$s < 0, \quad s_1 < 0, \quad s_2 < 0, \quad c_1 > 0, \quad c_2 > 0. \quad (33)$$

Of course, there are other options available dependant on the values of n .




FIG. 2: The evolution of EOS for K-essence with respect to the scale factor when $n = -5, -4, -3, -2, -1, 1, 2, 3, 4, 5$ from down to top, respectively. We have put $s_1 = s_2 = -1$, $c_1 = 1$, $c_2 = 8$.

In Fig. (2) we plot the EOS w with respect to the scale factor. It shows that the EOS starts from -1 at very early universe and then decreases to negative. Subsequently it approaches -1 in the distant future. This shows the energy density emerges as nearly a constant at very early universe. Then it gets a remarkable growth. Eventually, the energy density climbs to a very huge constant again by the order of 120 which is the order Planck energy density. To show this point, we plot the evolution for the Log-ratio

$$\eta = \log \frac{\rho_K}{\rho_{K0}}. \quad (34)$$

of K-essence density ρ_K to its present-day value ρ_{K0} in Fig. (3). Thus we conclude that the fate of our Universe is to undergo an inflation a third time.




FIG. 3: The evolution of for the Log-ratio η of K-essence density to its present-day value with respect to the scale factor. We have put $s_1 = -2 \cdot 10^{-3}$, $s_2 = -10^{-3}$, $c_1 = 10$, $c_2 = 1$, $n = 170$.

In Fig. (4) and Fig. (5) we plot the evolution of EOS for K-essence with respect to the natural logarithm of scale factor. The present-day EOS is slightly less than -1 which is consistent with the astronomical observations.




FIG. 4: The evolution of EOS for K-essence with respect to the natural logarithm of scale factor. We have put $s_1 = -2 \cdot 10^{-3}$, $s_2 = -10^{-3}$, $c_1 = 10$, $c_2 = 1$, $n = 170$.

On the other hand, Fig. (2) shows that the EOS can also start from -1 at very early universe and then increases to positive, its curves bulging and distended. In the end, it arrives at -1 . This shows the energy density is nearly a constant at very early universe and then it decreases significantly. Finally, the energy density dwells on a very small constant. This indicates that the K-essence can unify the early inflation field and the later dark energy. In order to make this clear, we plot the evolution for the Log-ratio η of K-essence density to its present-day




FIG. 5: The evolution of EOS for K-essence being a part of Fig. (4), with respect to the natural logarithm of scale factor.

value, the equation of state of K-essence with respect to the natural logarithm of scale factor in Fig. (6), Fig. (7) and Fig. (8).

The Hubble radius of the present-day Universe is the order of 10^{25} meter. Inflation is generally assumed starting from the Planck length, 10^{-35} meter. So on the whole, the scale factor expands by the order of 10^{60} or 140 e-folds. In order to solve the horizon problem, 60 e-folds more or less from inflation is needed. Without the loss of generality, we assume inflation produces 70 e-folds. Therefore, inflation starts from $a \sim e^{-140}$ to $a \sim e^{-70}$. Then from $a \sim e^{-70}$ to $a \sim e^0$, the universe goes through three stages, namely, the radiation-dominated, matter-dominated and matter plus dark-energy dominated stages, respectively,

Fig. (6) tells us the early density is the order of 120 compared to the present-day cosmological constant. So the cosmological constant problem disappears. Fig. (7) shows the EOS can be very large which indicates the K-essence energy density undergoes a precipitous decline at this epoch. Fig. (8) clearly shows the EOS has -1 as its starting point and finally consequently result.

In the next subsection, we shall present the other example which can unify the early inflation field and the latter cosmic dark energy.

3. The case of $F = s_1\sqrt{x} + s_2\sqrt{y} + se^{n(\frac{x}{y})^{\frac{m}{2}}}$

Thirdly, we consider the Lagrangian

$$F = s_1\sqrt{x} + s_2\sqrt{y} + se^{n(\frac{x}{y})^{\frac{m}{2}}}, \quad (35)$$

where s_1 , s_2 , s , n and m are constants. The cosmological equations of motion, Eqs. (10,11) give the energy density

$$\rho_K = se^{n\left(\frac{c_2 - a^3 s_2}{a^3 s_1 - c_1}\right)^{\frac{m}{2}}}, \quad (36)$$




FIG. 6: The evolution of for the Log-ratio η of K-essence density to its present-day value with respect to the natural logarithm of scale factor. We have put $s_1 = -4$, $s_2 = -1$, $c_1 = 10^{-93}$, $c_2 = 10^{-93}$, $n = 200$.




FIG. 7: The evolution of EOS for K-essence with respect to the natural logarithm of scale factor. We have put $s_1 = -4$, $s_2 = -1$, $c_1 = 10^{-93}$, $c_2 = 10^{-93}$, $n = 200$.

with c_1 and c_2 two integration constants. When the scale factor a is sufficiently small, i.e. $a \rightarrow 0$, we obtain a constant energy density

$$\rho_{Ke} = se^{n\frac{c_2^{\frac{m}{2}}}{(-c_1)^{\frac{m}{2}}}}. \quad (37)$$

On the other hand, when the scale factor is sufficient large, we also have a constant energy density

$$\rho_{Kl} = se^{n\frac{(-s_2)^{\frac{m}{2}}}{s_1^{\frac{m}{2}}}}. \quad (38)$$

The Log-ratio of above constant densities is




FIG. 8: The evolution of EOS for K-essence being a part of Fig. (7), with respect to the natural logarithm of scale factor.

$$\eta \equiv \log \frac{\rho_{Ke}}{\rho_{Kl}} = \frac{1}{\ln 10} \left[n \frac{c_2^m}{(-c_1)^m} - n \frac{(-s_2)^m}{s_1^m} \right]. \quad (39)$$

If we require

$$n \frac{c_2^m}{(-c_1)^m} \sim 277, \quad n \frac{(-s_2)^m}{s_1^m} \sim 1, \quad (40)$$

we would find

$$\eta \sim 120. \quad (41)$$

Namely, the constant energy density for inflation is 120 order of magnitude larger than the present dark energy density. To make this clear and for simplicity, we consider the parameter

$$m = 1. \quad (42)$$

Then we obtain the energy density

$$\rho_K = s e^{-n \frac{a^3 s_2 - c_2}{a^3 s_1 - c_1}}. \quad (43)$$

Given that the energy conservation equation Eq. (32), the EOS can be obtained. In Fig. (9), Fig. (10) and Fig. (11), we plot the evolution for the Log-ratio η of K-essence density to its present-day value, the EOS of K-essence with respect to the natural logarithm of scale factor. To conclude, we find the Lagrangian function Eq. (35) can be used as a model of unification of early inflation field and later cosmic dark energy.

4. The case of $F = s_1 \sqrt{x} + s_2 \sqrt{y} + s \frac{\sqrt{x}}{y^{\frac{n}{2}}}$

Finally, we consider the Lagrangian




FIG. 9: The evolution of for the Log-ratio η of K-essence density to its present-day value with respect to the natural logarithm of scale factor. We have put $s_1 = -277$, $s_2 = -1$, $c_1 = 10^{-93}$, $c_2 = 10^{-93}$, $n = -277$.




FIG. 10: The evolution of EOS for K-essence with respect to the natural logarithm of scale factor. We have put $s_1 = -277$, $s_2 = -1$, $c_1 = 10^{-93}$, $c_2 = 10^{-93}$, $n = -277$.

$$F = s_1 \sqrt{x} + s_2 \sqrt{y} + s \frac{\sqrt{x}}{y^{\frac{n}{2}}}, \quad (44)$$

where s_1 , s_2 , s and n are constants. The cosmological equations of motion, Eqs. (10,11) give the energy density

$$\rho_K = \frac{1}{n} s^{\frac{1}{n}} (s_2 a^3 - c_2) a^{-3 + \frac{3}{n}} (c_1 - s_1 a^3)^{-\frac{1}{n}}, \quad (45)$$

where c_1 and c_2 are integration constants. It is straightforward to obtain the pressure p_K and thus equation of state ω




FIG. 11: The evolution of EOS for K-essence being a part of Fig. (10), with respect to the natural logarithm of scale factor.

$$\omega = -\frac{ns_1s_2a^6 - ns_2c_1a^3 - s_2c_1a^3 + c_1c_2}{n(s_2a^3 - c_2)(s_1a^3 - c_1)}, \quad (46)$$

for K-Essence by substituting the energy density into the energy conservation equation. In Fig. (12), Fig. (13), Fig. (14) and Fig. (15), we plot the EOS with respect to the scale factor. Fig. (12) shows the EOS can be very positive or negative at the early universe. The large positive and negative EOS means the energy density decreases and increases, respectively, very fast. When $a \rightarrow 0$, we have the EOS

$$\omega = -\frac{1}{n}. \quad (47)$$

Then when $n \rightarrow \pm\infty$, we have $\omega = 0$ as shown in Fig. (12). When $a \rightarrow \infty$, we have

$$\omega = -1, \quad (48)$$

which is also shown in Fig. (12). Fig. (13), Fig. (14) and Fig. (15) reveal that the EOS can cross the phantom divide either from large to small as shown in Fig. (13) and Fig. (14) or from small to large as shown in Fig. (14). They have one thing in common. Namely, they all approaches -1 in distant future.

It is of particular interest when $n = 1$. We find that in this case, the K-essence can unify inflation field, dark matter and dark energy. Eq. (47) and Eq. (48) tell us the EOS is -1 both at very early universe and later universe which is the desired feature to be the candidate of inflation field and dark energy. In Fig. (16) we plot the EOS of K-essence with respect to the natural logarithm of scale factor when $n = 1$. The figure shows inflation starts from $a \sim e^{-140}$ to $a \sim e^{-80}$, i.e., 60 e-folds are generated by inflation. Then from $a \sim e^{-80}$ to $a \sim e^{-0}$, K-essence plays the role of dark matter with EOS of 0. Finally,




FIG. 12: The evolution of EOS for K-essence with respect to the scale factor when $n = -\frac{1}{2}, -\frac{2}{3}, -1, -2, -5$ from top to down for the upper half of the graph. For the lower half of the graph, we put $n = \frac{1}{2}, \frac{2}{3}, 1, 5, 100$ from down to top, respectively. We set $s_1 = -1, s_2 = 1, c_2 = -1, c_1 = 10$ for both upper and lower graphs.




FIG. 13: The evolution of EOS for K-essence with respect to the scale factor when $n = -2$ and $n = 2$ for the upper and lower half of the graph, respectively. We set $s_2 = 100, 5, 1, 0.5, 0.25$ from down to top, respectively, for both halves. We also set $s_1 = -1, c_2 = -1, c_1 = 10$ for both upper and lower graphs.

at the vicinity of a^0 , namely, the present-day Universe, K-essence plays the role of both dark matter and dark energy. To illustrate this point, we plot the evolution of EOS with respect to cosmic redshift in Fig. (17). The circled line is for Λ CDM paradigm. As we can see, the two curves are almost overlap with each other, suggesting that the unification model holds here. In Fig. (18), we plot the evolution of for the Log-ratio η of K-essence density to its present-day value with respect to the nat-




FIG. 14: The evolution of EOS for K-essence with respect to the scale factor when $c_2 = -0.5, -20, -60, -200, -800$ from down to top. We set $n = \frac{1}{2}, s_1 = -1, s_2 = 1, c_1 = 10$.




FIG. 15: The evolution of EOS for K-essence with respect to the scale factor when $c_2 = -0.25, -8, -60, -400, -1800$ from down to top. We set $n = 2, s_1 = -1, s_2 = 1, c_1 = 10$.

ural logarithm of scale factor. There is a slope between two platforms. The slope denotes the decreasing of dark matter with the expansion of the Universe.

To summarize at this stage, the two-field pure K-essence is a potential candidate for the unification of inflation field, dark matter and dark energy. In the next section, we turn to the issue of speed of perturbations for K-essence.




FIG. 16: The evolution of EOS for K-essence with respect to the natural logarithm of scale factor when $c_2 = -1, s_1 = -10^{-14}, s_2 = 2, c_1 = 10^{-125}$.




FIG. 17: The evolution of EOS for K-essence with respect to the cosmological redshift when $c_2 = -1, s_1 = -10^{-14}, s_2 = 2, c_1 = 10^{-125}$. The circled line is for the Λ CDM paradigm.

IV. ON THE SPEED OF PERTURBATIONS

All the Lagrangian functions studied in the previous section are covered by

$$F = 2y + s_1\sqrt{x} + s_2\sqrt{y} + s\frac{x^\alpha}{y^\beta} + se^{n(\frac{x}{y})^{\frac{m}{2}}}. \quad (49)$$

The first three terms are the kinetic terms and the last two can be understood as “potential” terms. The “potential” terms are novel in that they consist of quotient expression of kinetic energies.

The above unified Lagrangian can be written as

$$F = (\phi_1 + s\alpha\phi_3 + \phi_5)x + (2 + \phi_2 - s\beta\phi_4 - \phi_6)y + s(1 - \alpha + \beta)\phi_3^{-\frac{\alpha}{1-\alpha+\beta}}\phi_4^{\frac{\beta}{1-\alpha+\beta}}$$




FIG. 18: The evolution of for the Log-ratio η of K-essence density to its present-day value with respect to the natural logarithm of scale factor when $c_2 = -1$, $s_1 = -10^{-14}$, $s_2 = 2$, $c_1 = 10^{-125}$.

$$+ \frac{1}{4} \frac{s_1^2}{\phi_1} + \frac{1}{4} \frac{s_2^2}{\phi_2} + s e^{n \left(\frac{\phi_6}{\phi_5} \right)^{\frac{m}{2}}}, \quad (50)$$

by the aid of six auxiliary scalar fields, ϕ_i with $i = 1 - 6$. Now the Lagrangian presents itself as a scalar theory with eight scalar fields. Namely, two dynamical scalars ϕ , ψ which are equipped with the kinetic terms plus six auxiliary scalars ϕ_i . The auxiliary scalars make a contribution of a scalar potential, $V(\phi_i)$.

$$V = s (1 - \alpha + \beta) \phi_3^{-\frac{\alpha}{1-\alpha+\beta}} \phi_4^{\frac{\beta}{1-\alpha+\beta}} + \frac{1}{4} \frac{s_1^2}{\phi_1} + \frac{1}{4} \frac{s_2^2}{\phi_2} + s e^{n \left(\frac{\phi_6}{\phi_5} \right)^{\frac{m}{2}}}, \quad (51)$$

We recognize immediately that it belongs to the generalized multi-field theories [32] where the linear perturbations are already completed. Then we conclude from Ref. [32] that both the entropy modes and adiabatic mode propagate with the speed of light in the framework of Lagrangian function Eq. (50). Now there is an important issue that we need to direct our attention to. The issue is that the speed of perturbations depend on the framework or gauge for the theory. Namely, the speed of perturbations in the framework of Eq. (49) is different from that in the framework of Eq. (50). We can understand this point from a simple example. For simplicity, we consider one-field pure K-essence with the Lagrangian

$$F = x^2, \quad (52)$$

It is well-known that the speed of perturbations in this framework is [2]

$$c_s^2 \equiv \frac{F_{,x}}{F_{,x} + 2xF_{,xx}} = \frac{1}{2}. \quad (53)$$

However, the Lagrangian function Eq. (54) can be transformed to be

$$F = \phi_1 x - \frac{\phi_1^2}{4}, \quad (54)$$

by introducing the auxiliary scalar field ϕ_1 . Then the speed of perturbations in this framework is [32]

$$c_s^2 \equiv \frac{F_{,x}}{F_{,x} + 2xF_{,xx}} = 1. \quad (55)$$

Therefore, from the perspective of speed of perturbations, the two frameworks are merely equivalent to each other mathematically, not physically.

V. BLACK HOLE SOLUTIONS

In the last section, we turn our attention to seek for black hole solutions with two-field pure K-essence. We find it is not a hard work because of the absence of scalar fields in the Lagrangian except for their derivatives.

A. solutions

The line element of static and spherically symmetric solution for black holes can always be written as

$$ds^2 = -U(r) dt^2 + \frac{N(r)^2}{U(r)} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2. \quad (56)$$

Since the spacetime is static and spherically symmetric, the two scalar fields ϕ and ψ are the functions of radial coordinate r , namely, $\phi = \phi(r)$ and $\psi = \psi(r)$.

At this moment in time, we are interested in the expression of F as follows

$$F = 2y + s \sqrt{\frac{x}{y}}, \quad (57)$$

with s a positive constant. Then the Einstein equations are found to be

$$\frac{U'}{rN^2} - \frac{2UN'}{rN^3} - \frac{1}{r^2} + \frac{U}{r^2N^2} = 8\pi \left(\frac{U\psi'^2}{N^2} + s \frac{\phi'}{\psi'} \right), \quad (58)$$

$$\frac{U'}{rN^2} - \frac{1}{r^2} + \frac{U}{r^2N^2} = 8\pi \left(-\frac{U\psi'^2}{N^2} + s \frac{\phi'}{\psi'} \right), \quad (59)$$

$$\frac{U'}{rN^2} - \frac{UN'}{rN^3} + \frac{U''}{2N^2} - \frac{U'N'}{2N^3} = 8\pi \left(\frac{U\psi'^2}{N^2} + s \frac{\phi'}{\psi'} \right), \quad (60)$$

and the equations of motion for ϕ and ψ are

$$\left(\frac{Nr^2}{\psi'}\right)' = 0, \quad \left(-\frac{2Ur^2\psi'}{N} + \frac{sr^2N\phi'}{\psi'^2}\right)' = 0. \quad (61)$$

Here the apostrophe ' stands for the derivative with respect to the coordinate variable r . Solving these equations, we find the solutions

$$\psi = \frac{\sqrt{6}}{12\sqrt{\pi}} \operatorname{arcsinh}\left(\frac{r^6}{a_3^6}\right), \quad (62)$$

$$\phi' = -r^5 \int \frac{3}{32\pi^{\frac{3}{2}}} \cdot \frac{\sqrt{6}\sqrt{r^6+a_3^6}(2a_1r^4+a_2a_3^6)}{sa_1r^4(r^2+a_3^2)^2(r^4-r^2a_3^2+a_3^4)^2} dr + a_4r^5, \quad (63)$$

$$N = \frac{a_1}{\sqrt{r^6+a_3^6}}, \quad (64)$$

$$U = \frac{2a_1^2\sqrt{r^6+a_3^6}}{3a_3^9} \operatorname{hypergoem}\left[\left(\frac{1}{2}, \frac{1}{6}\right), \left(\frac{7}{6}\right), -\frac{r^6}{a_3^6}\right] + \frac{a_1^2}{3a_3^6} + \frac{a_1a_2r^2}{3a_3^6} + \frac{8\pi^{\frac{3}{2}}}{9r}\sqrt{6}sa_1^2a_4\sqrt{r^6+a_3^6}. \quad (65)$$

Here the parameters a_i with $i = 1, 2, 3, 4$ are integration constants. In the limit of $r \rightarrow 0$, we have

$$U = \frac{a_1^2}{a_3^6} + \frac{8\sqrt{6}sa_4a_1^2a_3^{\frac{3}{2}}}{9r}. \quad (66)$$

On the other side, in the limit of $r \rightarrow +\infty$, we know

$$\frac{2a_1^2\sqrt{r^6+a_3^6}}{3a_3^9} \operatorname{hypergoem}\left[\left(\frac{1}{2}, \frac{1}{6}\right), \left(\frac{7}{6}\right), -\frac{r^6}{a_3^6}\right] \propto r^2, \quad (67)$$

thus

$$U \propto r^2, \quad (68)$$

when $r \rightarrow +\infty$.

B. physical significance of the parameters

By expanding the expression of U into a power series of r , we obtain

$$U = \frac{a_1^2}{a_3^6} + \frac{8\sqrt{6}sa_4a_1^2a_3^{\frac{3}{2}}}{9r} + \frac{a_1a_2}{3a_3^6}r^2 + O(r^5). \quad (69)$$

Taking account of Eq. (66) and Eq. (67), we are aware that we should compare the metric function with the

Schwarzschild-de Sitter solution. Then the parameters are required to satisfy

$$\frac{a_1^2}{a_3^6} = 1, \quad \frac{8\sqrt{6}sa_4a_1^2a_3^{\frac{3}{2}}}{9} = -2M. \quad (70)$$

So we obtain

$$a_1 = a_3^3, \quad a_4 = -\frac{3\sqrt{6}M}{8sa_3^9\pi^{\frac{3}{2}}}. \quad (71)$$

Then the metric functions turn out to be

$$U = \frac{2}{3} \frac{\sqrt{r^6+a_3^6}}{a_3^2} \operatorname{hypergoem}\left[\left(\frac{1}{2}, \frac{1}{6}\right), \left(\frac{7}{6}\right), -\frac{r^6}{a_3^6}\right] + \frac{1}{3} - \frac{2M\sqrt{r^6+a_3^6}}{a_3^3r} + \frac{a_2r^2}{3a_3^3}, \quad (72)$$

$$N = \frac{a_3^3}{\sqrt{r^6+a_3^6}}. \quad (73)$$

We observe that in the limit of

$$r \ll a_3, \quad (74)$$

we have

$$\frac{\sqrt{r^6+a_3^6}}{a_3^2} \operatorname{hypergoem}\left[\left(\frac{1}{2}, \frac{1}{6}\right), \left(\frac{7}{6}\right), -\frac{r^6}{a_3^6}\right] = 1.$$

Then the solution goes back to the Schwarzschild-de Sitter solution for $a_2 < 0$ and Schwarzschild-anti-de Sitter solution for $a_2 > 0$. Here M represents the mass of the black hole. It is obvious a_2 and a_3 have the dimension of mass and they can be understood as the scalar charges of field ϕ and ψ , respectively. Similar to the Schwarzschild-de Sitter solution, the spacetime described by Eq. (72) and Eq. (73) have generally two horizons, i.e., the black hole horizon and the cosmic horizon.

VI. CONCLUSION AND DISCUSSION

In conclusion, we have studied a novel two-field pure K-essence whose Lagrangian is the function of kinetic terms of two scalar fields while independent on the scalar fields. The absence of the scalar fields variables in the Lagrangian makes the equations of motion for scalar fields considerable simple and their analytic solutions can be found. The novelty of this k-essence lies in the 'potential' which is made up with quotient of kinetic terms of the scalar fields.

We find that this novel K-essence have many interesting features. They are listed as follows.

1. The equation of state covers a very large parameter-space, i.e. $-\infty < \omega < +\infty$ which is remarkably different from that of quintessence, i.e. $-1 < \omega < +1$.
2. The EOS starts from -1 , then increases or decreases, and ends in -1 . Then two fates of the Universe are laid before us. One is a de Sitter universe governed by a very small constant density and the other is an inflationary universe governed by a very large constant density.
3. The EOS crosses the phantom divide either from positive to negative or from negative to positive.
4. Since the EOS begins from -1 and ends in -1 , K-essence unifies the early inflation field, the later dark energy and the dark matter in between.
5. The simplicity of the equations of motion enables

us to obtain the exact solution of black holes sourced by the K-essence.

6. The equivalence of this novel K-essence to the multi-field theory is merely from the perspective of mathematics, not physics. The reason is that the metric of field space is not the function of dynamical scalar fields, but the function of auxiliary scalar fields.

ACKNOWLEDGMENTS

The work is supported by the National Key R&D Program of China grants No. 2022YFF0503404 and No. 2022SKA0110100.

-
- [1] C. Armendariz-Picon, T. Damour, and V. Mukhanov, *Phys. Lett. B* **458**, 209 (1999).
 - [2] J. Garriga and V. F. Mukhanov, *Phys. Lett. B* **458**, 219 (1999). [arXiv:hep-th/9904176].
 - [3] T. Chiba, T. Okabe, and M. Yamaguchi, *Phys. Rev. D* **62**, 023511 (2000).
 - [4] C. Armendariz-Picon, V. Mukhanov, and P. J. Steinhardt, *Phys. Rev. Lett.* **85**, 4438 (2000).
 - [5] C. Armendariz-Picon, V. Mukhanov, and P. J. Steinhardt, *Phys. Rev. D* **63**, 103510 (2001).
 - [6] T. Chiba, *Phys. Rev. D* **66**, 063514 (2002).
 - [7] L. P. Chimento and A. Feinstein, *Mod. Phys. Lett. A* **19**, 761 (2004).
 - [8] L. P. Chimento, *Phys. Rev. D* **69**, 123517 (2004), astro-ph/0311613
 - [9] M. Malquarti, E. J. Copeland and Andrew R. Liddle, *Phys. Rev. D* **68**, 023512 (2003).
 - [10] R. J. Scherrer, *Phys. Rev. Lett.* **93**, 2004 (011301).
 - [11] L. P. Chimento and R. Lazkoz, *Phys. Rev. D* **71**, 023505 (2005).
 - [12] L. R. Abramo, N. Pinto-Neto, *Phys. Rev. D* **73**, 063522 (2006).
 - [13] C. Bonvin, C. Caprini and R. Durrer, *Phys. Rev. Lett.* **97**, 081303 (2006).
 - [14] K. Bamba, J. Matsumoto and S. Nojiri, *Phys. Rev. D* **85**, 084026 (2012).
 - [15] J. P. Bruneton, *Phys. Rev. D* **75**, 085013 (2007).
 - [16] T. Chiba, S. Dutta, and R. J. Scherrer, *Phys. Rev. D* **80**, 043517 (2009).
 - [17] R. Herrera, *Phys. Rev. D* **102**, 123508 (2020).
 - [18] N. Bose and A. S. Majumdar, *Phys. Rev. D* **79**, 103517 (2009).
 - [19] N. Bose and A. S. Majumdar, *Phys. Rev. D* **80**, 103508 (2009).
 - [20] S. X. Tian and Z.-H. Zhu, *Phys. Rev. D* **103**, 043518 (2021).
 - [21] A. G. Riess, S. Casertano, W. Yuan, J. Bradley Bowers, L. Macri, J. C. Zinn, and D. Scolnic, *Astrophys. J. Lett.* **908**, L6 (2021).
 - [22] W. L. Freedman, *Astrophys. J.* **919**, 16 (2021).
 - [23] Y. S. Hung and S. P. Miao, *Phys. Rev. D* **107**, 103533 (2023).
 - [24] J. Kehayias, R. J. Scherrer, *Phys. Rev. D* **100**, 023525 (2019).
 - [25] A. Sarkar, A. Ali, K. R. Nayak, A. S. Majumdar, *Phys. Rev. D* **107**, 084038 (2023).
 - [26] G. Lara, M. Bezares, E. Barausse, *Phys. Rev. D* **105**, 064058 (2022).
 - [27] Z. Huang, *Phys. Rev. D* **104**, 103533 (2021).
 - [28] A. Chatterjee, S. Hussain, K. Bhattacharya, *Phys. Rev. D* **104**, 103505 (2021).
 - [29] C. Nagy, Z. Keresztes, L. Á. Gergely, *Phys. Rev. D* **103**, 124056 (2021).
 - [30] A. O. Barvinsky, N. Kolganov, and A. Vikman, *Phys. Rev. D* **103**, 064035 (2021).
 - [31] A. L. Ferreira, J. N. Pinto-Neto and J. Zanelli, *Phys. Rev. D* **109**, 023515 (2024).
 - [32] D. Langlois and S. Renaux-Petel, *JCAP* **0804**, 017 (2008).
 - [33] C. Gao, arXiv: gr-qc/2203.05957.
 - [34] T. Padmanabhan, *Phys. Rep.* **380**, 235 (2003).
 - [35] N. Afshordi, D. J. H. Chung, and G. Geshnizjani, *Phys. Rev. D* **75**, 083513 (2007), arXiv:hep-th/0609150.