

Conservation laws for a generalized seventh order KdV equation

M.S. Bruzón, A.P. Márquez, T.M. Garrido, E. Recio, R. de la Rosa
Universidad de Cádiz, Spain

e-mail: m.bruzon@uca.es, almuneda.marquez@uca.es, tamara.garrido@uca.es,
elena.recio@uca.es, rafael.delarosa@uca.es

Abstract

In this paper, by applying the multiplier method we obtain a complete classification of low-order local conservation laws for a generalized seventh-order KdV equation depending on seven arbitrary nonzero parameters. We apply the Lie method in order to classify all point symmetries admitted by the equation in terms of the arbitrary parameters. We find that there are no special cases of the parameters for which the equation admits extra symmetries, other than those that can be found by inspection (scaling symmetry and space and time translation symmetries). We consider the reduced ordinary differential equations and we determined all integrating factors of the reduced equation from the combined x - and t -translation symmetries. Finally, we observe that all integrating factors arise by reduction of the low-order multipliers of the generalized seventh-order KdV equation.

Keywords: Partial differential equations, multiplier method, conservation laws, first integrals.

1 Introduction

In this paper we study the following generalized seventh-order KdV equation:

$$u_t + a_1 u^3 u_x + a_2 u_x^3 + a_3 uu_x u_{2x} + a_4 u^2 u_{3x} + a_5 u_{2x} u_{3x} + a_6 u_x u_{4x} + a_7 uu_{5x} + u_{7x} = 0, \quad (1)$$

where a_i ($i = 1, \dots, 7$) are nonzero parameters and $u_{nx} = \frac{\partial^n u}{\partial x^n}$ with $n = 2, \dots, 7$.

The Korteweg-de Vries equation (KdV) was proposed to describe the steady translation water waves in a channel. The main property of this equation is the equal occurrence of non-linearity and dispersion. Some of its most important solutions, the solitary waves, are single,

localised and non-dispersive entities, having also localised finite energy density. Several methods have been applied to solve the KdV equation such as the inverse scattering method or theoretical group methods.

Many papers have been published recently studying KdV equations of different order [1, 28, 27, 29]. The integrability of this equation is important because this nonlinear evolution equation describes the behaviour of physical phenomena such as shallow water waves and plasmas. Therefore, its conservation laws are determined to predict its complete integrability.

For instance, in [30] the author proved that the parametrized seventh-order KdV equation passes the Painlevé test only in three distinct cases. Firstly, he made use of the truncated Painlevé expansion to derive their bilinear forms and then he constructed their multi-soliton solutions. Secondly, the pseudopotentials and Lax pairs were also derived. Finally, the infinite conservation laws were found using its Lax pair, and the author claimed that “all conserved densities and fluxes” are obtained with explicit recursion formulas. Moreover, in [1] the author obtained a soliton solution of the fifth-order KdV equation applying the homotopy analysis method (HAM). Besides, in [29] multiple-soliton solutions were obtained for the ninth-order KdV equation applying the Hirota bilinear method.

It is straightforward to see that equation (1) is invariant under the scaling point symmetry

$$x \rightarrow \lambda x, \quad t \rightarrow \lambda^7 t, \quad u \rightarrow \lambda^{-2} u.$$

Note that equation (1) contains all of the possible monomial terms with positive powers of u and derivatives of u that preserve the scaling symmetry. From [17] one can often find fluxes of conservation laws by an algebraic formula in terms of the corresponding sets of multipliers without integration.

Conservation laws play a significant role in the solution process of an equation. They are very useful in the study of partial differential equations (PDEs) in which certain physical properties do not change in the course of time, especially for determining conserved quantities and constants of motion. Furthermore, they are good predicting linearizations, as well as checking the accuracy of numerical solution methods. Anco and Bluman proposed the multiplier method [8, 10, 11] that gives a general treatment of a direct conservation law for PDEs. Some examples of the multiplier method can be found in [12, 13, 14, 15, 18, 19, 22, 24, 25]. In the case of evolution equations, there exists a one-to-one relationship between non-zero multipliers and non-trivial conserved currents (up to local equivalence). Moreover, conservation laws which possess a physical importance usually come from low-order multipliers. In addition, higher-order multipliers are related to the integrability of the equation.

In this work we consider the generalized seventh-order KdV equation (1), we apply the multiplier method and we determine the parameters $a_i \neq 0$ with $i = 1, \dots, 7$ for which equation (1) admits multipliers. For each multiplier, we construct the conservation laws of equation (1). Next, we perform an analysis of Lie symmetries and we use the symmetries to reduce the generalized seventh-order KdV (1) to an ODE. Furthermore, we find the integrating factors of the reduced ODE.

2 Multipliers and conservation laws of equation (1)

Conservation laws are of basic importance because they provide physical, conserved quantities for all solutions $u(x, t)$, and they can be used to check the accuracy of numerical solution methods. A general discussion of conservation laws and their applications to differential equations can be found in [7, 8, 10, 17, 18].

A local conservation law for the generalized seventh-order KdV equation (1) is a continuity equation

$$D_t T + D_x X = 0, \quad (2)$$

which holds on all solutions of equation (1), whereas the conserved density T and the spatial flux X are functions of t, x, u and derivatives of u . D_t and D_x denote the total derivative operators with respect to t and x respectively. In the case that, there is a function $\Theta(t, x, u, u_t, u_x, \dots)$ so that the conserved current vector $(T, X) = (D_x \Theta, -D_t \Theta)$ holds for every solution $u(t, x)$, equation (2) becomes an identity. Thus, this conservation law is called locally trivial. Thereby, if two conservation laws differ by a locally trivial conservation law, they are considered to be locally equivalent. A non-trivial conservation law can be written in a general form as follows

$$\frac{d}{dt} \int_{\Omega} T dx = -X \Big|_{\partial \Omega}, \quad (3)$$

where $\Omega \subseteq \mathbb{R}$ is any fixed spatial domain. Any local conservation law can be stated by using the characteristic form which arises from a divergence identity

$$\begin{aligned} D_t \tilde{T} + D_x \tilde{X} &= (u_t + a_1 u^3 u_x + a_2 u_x^3 + a_3 u u_x u_{2x} + a_4 u^2 u_{3x} \\ &+ a_5 u_{2x} u_{3x} + a_6 u_x u_{4x} + a_7 u u_{5x} + u_{7x}) Q, \end{aligned} \quad (4)$$

where $\tilde{T} = T + D_x \Theta$ and $\tilde{X} = X - D_t \Theta$ are locally equivalent to T and X , where the function $Q(t, x, u, u_t, u_x, \dots)$ which holds

$$Q = E_u(\tilde{T}), \quad (5)$$

is called a multiplier, and where E_u represents the Euler operator with respect to u [23],

$$E_u = \partial_u - D_x \partial_{u_x} - D_t \partial_{u_t} + D_x D_t \partial_{u_{xt}} + D_x^2 \partial_{u_{2x}} + \dots \quad (6)$$

For evolution equations, there is a one-to-one correspondence between non-zero multipliers and non-trivial conserved current vectors up to local equivalence [10, 23]. This correspondence comes from the characteristic equation for multipliers of the given equation. Moreover, in the case of dispersive nonlinear water waves equations, conservation laws of physical importance typically come from low-order multipliers [7]. The general form of low-order multipliers Q in terms of u and derivatives of u is given by those variables that can be differentiated to obtain a leading derivative of the KdV equation (1). Therefore, the general form for a low-order multiplier for equation (1) is

$$Q(t, x, u, u_x, u_{2x}, u_{3x}, u_{4x}, u_{5x}, u_{6x}). \quad (7)$$

Given a multiplier Q , we can obtain the conserved density using a standard method [31]

$$T = \int_0^1 uQ(x, t, \lambda u, \lambda u_x, \lambda u_{2x}, \dots) d\lambda. \quad (8)$$

All low-order multipliers can be found by solving the determining equation $E_u((u_t + a_1 u^3 u_x + a_2 u_x^3 + a_3 u u_x u_{2x} + a_4 u^2 u_{3x} + a_5 u_{2x} u_{3x} + a_6 u_x u_{4x} + a_7 u u_{5x} + u_{7x})Q) = 0$. In [10] from (5) and (8) it is defined the order of a conservation law for a generalized KdV equation as the order of the highest x derivative of u in its multiplier.

In order to obtain a general classification of low-order multipliers (7), we write and split the determining equation (4) yielding an overdetermined system for Q and the parameters a_i with $i = 1, \dots, 7$. The solutions with $a_i \neq 0$ are given by

1. For $a_2 = -a_4 + \frac{1}{2}a_3$ and a_3, a_4, a_5, a_6, a_7 nonzero constants,

$$Q_1 = 1 \quad (9)$$

2. For $a_1 = \frac{5}{49}a_6^3 - \frac{1}{49}a_6^2 a_5 - \frac{30}{49}a_6^2 a_7 + \frac{3}{49}a_6 a_5 a_7 + \frac{225}{196}a_6 a_7^2 - \frac{9}{196}a_5 a_7^2 - \frac{135}{196}a_7^3 + \frac{3}{7}a_6 a_4 + \frac{2}{21}a_6 a_2 - \frac{9}{14}a_4 a_7 - \frac{1}{7}a_2 a_7$, $a_3 = -\frac{5}{7}a_6^2 + \frac{1}{7}a_5 a_6 + \frac{45}{14}a_6 a_7 - \frac{3}{14}a_5 a_7 - \frac{45}{14}a_7^2 + a_4$,

$$Q_2 = \frac{1}{14}(2a_6 - 3a_7)u^2 + u_{2x} \quad (10)$$

3. For $a_2 = -3a_4 + a_3$, $a_5 = 5a_6 - 10a_7$,

$$Q_3 = u \quad (11)$$

4. For $a_1 = -\frac{1}{147}(a_6 - 4a_7)(4a_6^2 - 18a_6 a_7 + 8a_7^2 + 49a_4)$, $a_2 = \frac{1}{7}a_6^2 - \frac{9}{14}a_6 a_7 + \frac{2}{7}a_7^2 + 2a_4$, $a_3 = \frac{2}{7}a_6^2 - \frac{9}{7}a_6 a_7 + \frac{4}{7}a_7^2 + 6a_4$, $a_5 = 2a_6 - a_7$,

$$Q_4 = \left(\frac{4}{147}a_6^2 + \frac{8}{147}a_7^2 + \frac{1}{3}a_4 - \frac{6}{49}a_6 a_7 \right) u^3 + \left(\frac{2}{7}a_6 - \frac{a_7}{7} \right) u_{2x} u + \left(\frac{a_6}{7} - \frac{a_7}{14} \right) u_x^2 + u_{4x}, \quad (12)$$

$$Q_5 = \left(\frac{(4a_6^2 - 18a_6 a_7 + 8a_7^2 + 49a_4)(a_6 - 4a_7)u^3}{147} + \frac{1}{7}(2a_6 - a_7)(a_6 - 4a_7)u_{2x} u + \frac{1}{14}(2a_6 - a_7)(a_6 - 4a_7)u_x^2 + (a_6 - 4a_7)u_{4x} \right) t + x \quad (13)$$

5. For $a_2 = a_4, a_3 = 4a_4, a_5 = 5a_7, a_6 = 3a_7$,

$$Q_6 = \frac{1}{4}a_1 u^4 + a_4 u^2 u_{2x} + (a_4 u^2 + a_7 u_{4x})u + u_{6x} + 2u_x a_7 u_{3x} + 3/2 u_{2x}^2 a_7 \quad (14)$$

6. For $a_1 = \frac{5a_7^3}{98}, a_2 = \frac{5a_7^2}{14}, a_3 = \frac{10a_7^2}{7}, a_4 = \frac{5a_7^2}{14}, a_5 = 5a_7, a_6 = 3a_7,$

$$Q_7 = \frac{5a_7^2 u^4}{392} + \frac{5a_7 u^2 u_{2x}}{14} + \left(u_{4x} + \frac{5a_7 u_x^2}{14} \right) u + \frac{u_{6x}}{a_7} + 2u_x u_{3x} + \frac{3}{2} u_{2x}^2 \quad (15)$$

Thus, we obtain the following result:

Theorem 1 All nontrivial low-order local conservation laws admitted by the seventh-order KdV equation (1) with $a_i \neq 0, i = 1, \dots, 7$ are given by:

1. For $a_2 = -a_4 + \frac{1}{2}a_3$ and a_3, a_4, a_5, a_6, a_7 nonzero constants,

$$\begin{aligned} T_1 &= u, \\ X_1 &= u_{6x} + a_7 u u_{4x} + (a_6 - a_7) u_x u_{3x} + \left(\frac{1}{2}a_5 - \frac{1}{2}a_6 + \frac{1}{2}a_7 \right) u_{2x}^2 \\ &\quad + a_4 u^2 u_{2x} + \left(-a_4 + \frac{1}{2}a_3 \right) u u_x^2 + \frac{1}{4}a_1 u^4 \end{aligned} \quad (16)$$

2. For $a_1 = \frac{5}{49}a_6^3 - \frac{1}{49}a_6^2 a_5 - \frac{30}{49}a_6^2 a_7 + \frac{3}{49}a_6 a_5 a_7 + \frac{225}{196}a_6 a_7^2 - \frac{9}{196}a_5 a_7^2 - \frac{135}{196}a_7^3 + \frac{3}{7}a_6 a_4 + \frac{2}{21}a_6 a_2 - \frac{9}{14}a_4 a_7 - \frac{1}{7}a_2 a_7, a_3 = -\frac{5}{7}a_6^2 + \frac{1}{7}a_5 a_6 + \frac{45}{14}a_6 a_7 - \frac{3}{14}a_5 a_7 - \frac{45}{14}a_7^2 + a_4,$

$$\begin{aligned} T_2 &= \frac{1}{3}A u^3 - 7u_x^2, \\ X_2 &= (A u^2 + 14u_{2x}) u_{6x} + 7u_{4x}^2 + (-14u_{3x} + (-4a_6 + 6a_7) u u_x) u_{5x} \\ &\quad + ((4a_6 + 8a_7) u u_{2x} + (4a_6 - 6a_7) u_x^2 + a_7 A u^3) u_{4x} \\ &\quad + ((2a_6 + 4a_7) u_x u_{2x} + A(a_6 - 3a_7) u^2 u_x) u_{3x} \\ &\quad + \left(\frac{14}{3}a_5 - \frac{2}{3}a_6 - \frac{4}{3}a_7 \right) u_{2x}^3 + 14u_x u_x + (-2a_6 - 4a_7) u u_{3x}^2 \\ &\quad + \left(-\frac{3}{2}a_5 a_7 + \frac{9}{2}a_6 a_7 - a_6^2 - \frac{9}{2}a_7^2 + 7a_4 + a_5 a_6 \right) u^2 u_{2x}^2 \\ &\quad + (-2A(a_6 - 3a_7) u u_x^2 + a_4(2a_6 - 3a_7) u^4) u_{2x} \\ &\quad + (a_6^2 - \frac{9}{2}a_6 a_7 + \frac{9}{2}a_7^2 + \frac{7}{2}a_2) u_x^4 + \frac{1}{3}a_2(2a_6 - 3a_7) u^3 u_x^2 \\ &\quad + \frac{1}{3528}A^2(-6a_5 a_6 + 9a_5 a_7 + 30a_6^2 - 135a_6 a_7 + 135a_7^2 + 28a_2 + 126a_4) u^6 \end{aligned} \quad (17)$$

where $A = 2a_6 - 3a_7$

3. For $a_2 = -3a_4 + a_3, a_5 = 5a_6 - 10a_7,$

$$\begin{aligned} T_3 &= \frac{1}{2}u^2, \\ X_3 &= u u_{6x} - u_{5x} u_x + (a_7 u^2 + u_{2x}) u_{4x} - \frac{1}{2}u_{3x}^2 \\ &\quad + (a_6 - 2a_7) u u_x u_{3x} + (2a_6 - 4a_7) u u_{2x}^2 \\ &\quad + ((-a_6 + 2a_7) u_x^2 + a_4 u^3) u_{2x} + \left(\frac{1}{2}a_3 - \frac{3}{2}a_4 \right) u^2 u_x^2 \\ &\quad + \frac{1}{5}a_1 u^5 \end{aligned} \quad (18)$$

4. For $a_1 = -\frac{1}{147}(a_6 - 4a_7)(4a_6^2 - 18a_6a_7 + 8a_7^2 + 49a_4)$, $a_2 = \frac{1}{7}a_6^2 - \frac{9}{14}a_6a_7 + \frac{2}{7}a_7^2 + 2a_4$, $a_3 = \frac{2}{7}a_6^2 - \frac{9}{7}a_6a_7 + \frac{4}{7}a_7^2 + 6a_4$, $a_5 = 2a_6 - a_7$

$$T_4 = \frac{1}{2}u_{2x}^2 + \frac{1}{14}(a_7 - 2a_6)uu_x^2 + \left(\frac{1}{147}a_6^2 - \frac{3a_7a_6}{98} + \frac{2}{147}a_7^2 + \frac{1}{12}a_4\right)u^4, \quad (19)$$

5. For $a_1 = -\frac{1}{147}(a_6 - 4a_7)(4a_6^2 - 18a_6a_7 + 8a_7^2 + 49a_4)$, $a_2 = \frac{1}{7}a_6^2 - \frac{9}{14}a_6a_7 + \frac{2}{7}a_7^2 + 2a_4$, $a_3 = \frac{2}{7}a_6^2 - \frac{9}{7}a_6a_7 + \frac{4}{7}a_7^2 + 6a_4$, $a_5 = 2a_6 - a_7$

$$T_5 = \left(\frac{a_6}{2} - 2a_7\right)tu_{2x}^2 - \frac{1}{14}(-a_7 + 2a_6)(a_6 - 4a_7)tuu_x^2 + \frac{(4a_6^2 - 18a_7a_6 + 8a_7^2 + 49a_4)(a_6 - 4a_7)tu^4}{588} + xu, \quad (20)$$

6. For $a_2 = a_4, a_3 = 4a_4, a_5 = 5a_7, a_6 = 3a_7$

$$T_6 = -\frac{1}{2}u_{3x}^2 + \frac{1}{2}a_7uu_{2x}^2 - \frac{1}{2}a_4u^2u_x^2 + \frac{1}{20}a_1u^5, \quad (21)$$

7. For $a_1 = \frac{5a_7^3}{98}, a_2 = \frac{5a_7^2}{14}, a_3 = \frac{10a_7^2}{7}, a_4 = \frac{5a_7^2}{14}, a_5 = 5a_7, a_6 = 3a_7$

$$T_7 = -\frac{1}{2}a_7tu_{2x}^2 + \frac{5a_7^2tuu_x^2}{14} - \frac{5a_7^3tu^4}{392} + xu, \quad (22)$$

X_4, X_5, X_6 and X_7 are omitted here in order to save space.

3 Lie symmetries

In the Lie method, we required that the one-parameter Lie group of infinitesimal point transformations in (x, t, u) given by

$$\begin{aligned} x^* &= x + \varepsilon\xi(x, t, u) + O(\varepsilon^2), \\ t^* &= t + \varepsilon\tau(x, t, u) + O(\varepsilon^2), \\ u^* &= u + \varepsilon\eta(x, t, u) + O(\varepsilon^2), \end{aligned}$$

leaves invariant the set of solutions of equation (1), where ε is the group parameter.

Hence, an infinitesimal point symmetry for the generalized seventh-order KdV equation (1) is a generator

$$X = \xi(x, t, u)\frac{\partial}{\partial x} + \tau(x, t, u)\frac{\partial}{\partial t} + \eta(x, t, u)\frac{\partial}{\partial u}, \quad (23)$$

whose prolongation leaves invariant the equation (1). Every point symmetry (23) can be expressed by its characteristic form

$$\hat{X} = P\partial_u, \quad P = \eta - \tau u_t - \xi u_x.$$

The condition for the invariance of the equation (1) under a point symmetry (1) is given by

$$\begin{aligned}
0 = & (3a_1u^2u_x + a_3u_xu_{2x} + 2a_4uu_{3x} + a_7u_{5x})P + D_tP \\
& + (a_1u^3 + 3a_2u_x^2 + a_3uu_{2x} + a_6u_{4x})D_xP \\
& + (a_3uu_x + a_5u_{3x})D_x^2P + (a_4u^2 + a_5u_{2x})D_x^3P \\
& + a_6u_xD_x^4P + a_7uD_x^5P + D_x^7P
\end{aligned} \tag{24}$$

and holds for all solutions $u(x, t)$ of equation (1). Equation (24) is known as the symmetry determining equation. It splits with respect to x derivatives of u . This leads to a linear overdetermined system of equations on $\xi(x, t, u)$, $\tau(x, t, u)$, $\eta(x, t, u)$ along with $a_1, a_2, a_3, a_4, a_5, a_6, a_7$.

Solving the system, the only point symmetries admitted by (1) with $a_i \neq 0$, $i = 1, \dots, 7$, which are defined by the infinitesimal, are generated

$$\mathbf{X}_1 = \frac{\partial}{\partial x}, \quad \mathbf{X}_2 = \frac{\partial}{\partial t}, \quad \mathbf{X}_3 = x\frac{\partial}{\partial x} + 7t\frac{\partial}{\partial t} - 2u\frac{\partial}{\partial u}.$$

From $\mathbf{X}_1 + \mathbf{X}_2$, we obtain travelling wave reductions

$$z = x - \lambda t, \quad u = h(z),$$

where $h(z)$ satisfies

$$\begin{aligned}
& h_{7z} + a_7hh_{5z} + a_6h_zh_{4z} + a_5h_{2z}h_{3z} + a_4h^2h_{3z} + a_3hh_zh_{2z} \\
& + a_2(h_z)^3 + a_1h^3h_z - \lambda h_z = 0
\end{aligned} \tag{25}$$

with $h_{nz} = h^{(n)}(z)$ for $n = 2, \dots, 7$ (the n -th order derivative of $h(z)$).

From \mathbf{X}_3 we obtain the invariant solution

$$z = \frac{x^7}{t}, \quad u = \frac{h}{t^{\frac{1}{4n+4}}}x^{-2}h(z), \tag{26}$$

where $h(z)$ must satisfy a non-autonomous equation.

In [20, 21] used the complex singularity structure of solutions to identify an integrable case of the equations of motion for a rotating top. Ablowitz and Segur [2], and Ablowitz, Ramani, and Segur [3, 4, 5] observed this connection in the context of integrable PDEs. Their observations led to the following conjecture:

Any ODE which arises as a reduction of an integrable PDE possesses the Painlevé property, possibly after a transformation of variables.

In [30] the author showed that the parameterized seventh-order KdV equation (1) passes the Painlevé test only in three distinct cases:

- $a_1 = \frac{4}{147}, a_2 = \frac{1}{7}, a_3 = \frac{6}{7}, a_4 = \frac{2}{7}, a_5 = 3, a_6 = 2, a_7 = 1$
- $a_1 = \frac{5}{98}, a_2 = \frac{5}{14}, a_3 = \frac{10}{7}, a_4 = \frac{5}{14}, a_5 = 5, a_6 = 3, a_7 = 1$
- $a_1 = \frac{4}{147}, a_2 = \frac{5}{14}, a_3 = \frac{9}{7}, a_4 = \frac{2}{7}, a_5 = 6, a_6 = \frac{7}{2}, a_7 = 1$

Consequently, by applying the conjecture, the reduced equations for these cases possess the Painlevé property.

4 Integrating factors and first integrals for equation (25)

In [26], Sjöberg proposed the so-called double reduction method which took into consideration the relationship between symmetries and conservation laws. This method can be used to reduce a q th-order PDE into a $(q-1)$ th-order ODE. In [6], by using the direct method of the multipliers [10, 11], Anco developed symmetry properties of conservation laws of PDEs. In particular, the author provided a theoretical framework to comprehend and generalize the method proposed by Sjöberg. In this way, the author proved that conservation laws that are symmetry invariant or symmetry homogeneous have at least one important application: any symmetry-invariant conservation law will reduce to a first integral for the ODE obtained by symmetry reduction of the given PDE when symmetry-invariant solutions $u(t, x)$ are investigated. This provides a direct reduction of order of the ODE.

It is well known that for any n th-order ODE finding an integrating factor is equivalent to finding a first integral (see e.g. [9, 16]). In this section, we focus our attention on obtaining integrating factors of ODE (25) and, hence, first integrals. Furthermore, a first integral of equation (25) yields a quadrature which reduces (25) to a 6th-order ODE in terms of the original variables z, h, h_z, \dots, h_{6z} .

Following reference [16] an integrating factor of equation (25) is a function $Q(z, h, h_z, \dots, h_{6z}) \neq 0$, with $0 \leq l \leq 6$, which verifies the characteristic equation

$$D_z \tilde{X} = \left(h_{7z} + a_7 h h_{5z} + a_6 h_z h_{4z} + a_5 h_{2z} h_{3z} + a_4 h^2 h_{3z} + a_3 h h_z h_{2z} + a_2 (h_z)^3 + a_1 h^3 h_z - \lambda h_z \right) Q. \quad (27)$$

Moreover, all first integrals of ODE (25) derive from integrating factors verifying the characteristic equation (27) for any function $h(z)$. The determining equation to obtain all integrating factors is

$$\frac{\delta}{\delta h} \left((h_{7z} + a_7 h h_{5z} + a_6 h_z h_{4z} + a_5 h_{2z} h_{3z} + a_4 h^2 h_{3z} + a_3 h h_z h_{2z} + a_2 (h_z)^3 + a_1 h^3 h_z - \lambda h_z) Q \right) = 0, \quad (28)$$

where $\frac{\delta}{\delta h}$ represents the variational derivative. This equation must hold off of the set of solutions of equation (25). Once the integrating factors are found, the corresponding non-trivial first integrals are obtained by using a line integral [16].

The determining equation (28) splits with respect to the variables h_{7z}, \dots, h_{10z} . This yields a linear determining system for Q . Then, we determine all integrating factors:

1. For $a_2 = -a_4 + \frac{1}{2}a_3$ and a_3, a_4, a_5, a_6, a_7 nonzero constants

$$Q_1 = 1 \quad (29)$$

2. For $a_1 = \frac{5}{49}a_6^3 - \frac{1}{49}a_6^2a_5 - \frac{30}{49}a_6^2a_7 + \frac{3}{49}a_6a_5a_7 + \frac{225}{196}a_6a_7^2 - \frac{9}{196}a_5a_7^2 - \frac{135}{196}a_7^3 + \frac{3}{7}a_6a_4 + \frac{2}{21}a_6a_2 - \frac{9}{14}a_4a_7 - \frac{1}{7}a_2a_7$, $a_3 = -\frac{5}{7}a_6^2 + \frac{1}{7}a_5a_6 + \frac{45}{14}a_6a_7 - \frac{3}{14}a_5a_7 - \frac{45}{14}a_7^2 + a_4$, we obtain the multiplier

$$Q_2 = \frac{1}{14}(2a_6 - 3a_7)h^2 + h_{2z} \quad (30)$$

3. For $a_2 = -3a_4 + a_3$, $a_5 = 5a_6 - 10a_7$,

$$Q_3 = h \quad (31)$$

4. For $a_1 = -\frac{1}{147}(a_6 - 4a_7)(4a_6^2 - 18a_6a_7 + 8a_7^2 + 49a_4)$, $a_2 = \frac{1}{7}a_6^2 - \frac{9}{14}a_6a_7 + \frac{2}{7}a_7^2 + 2a_4$, $a_3 = \frac{2}{7}a_6^2 - \frac{9}{7}a_6a_7 + \frac{4}{7}a_7^2 + 6a_4$, $a_5 = 2a_6 - a_7$

$$Q_4 = \left(\frac{4}{147}a_6^2 + \frac{8}{147}a_7^2 + \frac{1}{3}a_4 - \frac{6}{49}a_6a_7\right)h^3 + \left(\frac{2}{7}a_6 - \frac{a_7}{7}\right)h_{2z}h + \left(\frac{a_6}{7} - \frac{a_7}{14}\right)h_z^2 + h_{4z}, \quad (32)$$

5. For $a_2 = a_4$, $a_3 = 4a_4$, $a_5 = -10a_7 + 5a_6$

$$Q_5 = \frac{1}{4}a_1h^4 + a_4h^2h_{2z} + (a_4h_z^2 + a_7h_{4z})h + h_{6z} + 2a_6h_{2z}^2 \quad (33)$$

$$-\frac{9}{2}a_7h_{2z}^2 + a_6h_zh_{3z} - a_7h_zh_{3z} \quad (34)$$

We observe that the integrating factors arise by reduction of the multipliers for equation (1).

5 Conclusions

For a generalized seventh-order KdV equation we have derived the low-order conservation laws by using the multiplier method. Moreover we have studied equation (1) from Lie symmetries viewpoint and we consider the reduced ordinary differential equation under the combined space and time translation symmetries. We have determined all integrating factors of an ordinary differential equations and we observe that the integrating factors arise by reduction of a conservation law for equation (1).

Acknowledgements

We gratefully acknowledge fruitful discussions and contributions received by Dr. Stephen Anco (Brock University). The authors express their sincere gratitude to the Plan Propio de Investigación de la Universidad de Cádiz.

References

- [1] S. Abbasbandy and F. Samadian Zakaria, Soliton solutions for the fifth-order KdV equation with the homotopy analysis method, *Nonlinear Dynamics*, **51** (2008), 83-87.
- [2] M. J. Ablowitz and H. Segur, Exact Linearization of a Painlevé Transcendent, *Physical Review Letters*, **38** (1977), 1103–1106.
- [3] M.J. Ablowitz, A. Ramani and H. Segur, Nonlinear Evolution Equations and Ordinary Differential Equations of Painlevé Type, *Lettere al Nuovo Cimento*, **23** (1978), 333–338.
- [4] M. J. Ablowitz, A. Ramani and H. Segur, A Connection between Nonlinear Evolution Equations and Ordinary Differential Equations of P-type. I, *Journal of Mathematical Physics*, **21** (1980), 715–721.
- [5] M. J. Ablowitz, A. Ramani and H. Segur, A Connection between Nonlinear Evolution Equations and Ordinary Differential Equations of P-type. II, *Journal of Mathematical Physics*, **21** (1980), 1006–1015.
- [6] S. C. Anco, Symmetry Properties of Conservation Laws, *International Journal of Modern Physics B*, **30** (2015), 1640003-1640014.
- [7] S. C. Anco, Generalization of Noether’s Theorem in Modern Form to Non-variational Partial Differential Equations, in *Recent Progress and Modern Challenges in Applied Mathematics, Modeling and Computational Science* **79**, Springer, (2017), 119–182.
- [8] S. C. Anco and G. W. Bluman, Direct construction of conservation laws from field equations, *Physical Review Letters*, **78** (1997), 2869–2873.
- [9] S. C. Anco and G. W. Bluman, Integrating factors and first integrals for ordinary differential equations, *European Journal of Applied Mathematics*, **9** (1998), 245–259.
- [10] S. C. Anco and G. W. Bluman, Direct construction method for conservation laws of partial differential equations part 1: Examples of conservation law classifications, *European Journal of Applied Mathematics*, **5** (2002), 545–566.
- [11] S. C. Anco and G. W. Bluman, Direct construction method for conservation laws of partial differential equations part 2: General treatment, *European Journal of Applied Mathematics*, **5** (2002), 567–585.
- [12] S. C. Anco and W. Feng, Group-invariant solutions of semilinear Schrodinger equations in multi-dimension, *Journal of Mathematical Physics*, **54** (2013), 12150.
- [13] S. C. Anco, M. Mohiuddin and T. Wolf, Travelling Waves and conservation laws for complex MKDV-type Equations, *Applied Mathematics and Computation*, **219** (2012), 679-698.

- [14] S. C. Anco, E. D. Avdonina, A. Gainetdinova, L. R. Galiakberova, N. H. Ibragimov and T. Wolf, Symmetries and conservation laws of the generalized Krichever-Novikov equation, *Journal of Physics A: Mathematical and Theoretical*, **49** (2016), 105201.
- [15] S. C. Anco, P. L. da Silva and I. L. Freire, A family of wave-breaking equations generalizing the Camassa-Holm and Novikov equations, *Journal of Mathematical Physics*, **56** (2015), 091506.
- [16] G. W. Bluman, S. Anco, *Symmetry and Integration Methods for Differential Equations*, Applied Mathematical Sciences series **154**, Springer, 2002.
- [17] G. W. Bluman, A. F. Cheviakov and S.C. Anco, *Applications of symmetry methods to partial differential equations*, Springer-New York, 2010.
- [18] M. S. Hashemi, A. Haji-Badali and P. Vafadar, Group Invariant Solutions and Conservation Laws of the Fornberg Whitham Equation, *Zeitschrift für Naturforschung A*, **69** (2014), 489-496.
- [19] C. M. Khalique, Exact solutions and conservation laws of a coupled integrable dispersionless system, *Filomat*, **26** (2012), 957-964.
- [20] S. Kowalevski, Sur le Problème de la Rotation d'un Corps Solide Autour d'un Point Fixé, *Acta Mathematica*, **12** (1889), 177–232.
- [21] S. Kowalevski, Sur une Propriété d'un Système d'Equations Différentielles qui Definit la Rotation d'un Corps Solide Autour d'un Point Fixé, *Acta Mathematica*, **14** (1889), 81–93.
- [22] B. Muatjetjeja and C. M. Khalique, Conservation laws for a variable coefficient variant Boussinesq system, *Abstract and Applied Analysis*, **2014** (2014), 169694.
- [23] P. J. Olver, *Applications of Lie groups to differential equations*, Springer-Verlag, 1986.
- [24] E. Recio and S. C. Anco, Conservation laws and symmetries of radial generalized nonlinear p-Laplacian evolution equations, *Journal of Mathematical Analysis and Applications*, **452** (2017), 1229–1261.
- [25] R. Tracinà, M. S. Bruzón, M. L. Gandarias and M. Torrisi, Nonlinear self-adjointness, conservation laws, exact solutions of a system of dispersive evolution equations, *Communications in Nonlinear Science and Numerical Simulations*, **19** (2014), 3036-3043.
- [26] A. Sjöberg, Double reduction of PDEs from the association of symmetries with conservation laws with applications, *Applied Mathematics and Computation*, **2** (2007), 608-616.
- [27] A. A. Soliman, A numerical simulation and explicit solutions of KdV-Burgers' and Lax's seventh-order KdV equations, *Chaos, Solitons and Fractals*, **29** (2006), 294-302.

- [28] A. Wazwaz, The extended tanh method for new solitons solutions for many forms of the fifth-order KdV equations, *Applied Mathematics and Computation*, **184** (2007), 1002-1014.
- [29] A. Wazwaz, Multiple-soliton solutions for the ninth-order KdV equation and sixth-order Boussinesq equation, *Applied Mathematics and Computation*, **203** (2008), 277-283.
- [30] G. Xu, The integrability for a generalized seventh-order KdV equation: Painlevé property, soliton solutions, Lax pairs and conservation laws, *Physica Scripta*, **89** (2014), 125201.
- [31] T. Wolf, An efficiency improved program LIEPDE for determining Lie-symmetries of PDEs, *Proceedings of Modern Group Analysis: advances analytical and computational methods in mathematical physics*, (1993), 377-385.