Towards the Mathematical Foundation of the Minimum Enclosing Ball and Related Problems

Michael N. Vrahatis

Abstract—Theoretical background is provided towards the mathematical foundation of the minimum enclosing ball problem. This problem concerns the determination of the unique spherical surface of smallest radius enclosing a given bounded set in the *d*dimensional Euclidean space. The study of several problems that are similar or related to the minimum enclosing ball problem has received a considerable impetus from the large amount of applications of these problems in various fields of science and technology. The proposed theoretical framework is based on several enclosing (covering) and partitioning (clustering) theorems and provides among others bounds and relations between the circumradius, inradius, diameter and width of a set. These enclosing and partitioning theorems are considered as cornerstones in the field that strongly influencing developments and generalizations to other spaces and non-Euclidean geometries.

Index Terms—Euclidean 1-center, Chebyshev radius and center, radii, diameter, width, property testing, theorems of Jung, Steinhagen, Perel'man, Carathéodory, Helly and Tverberg.

I. INTRODUCTION

HEORETICAL background is provided that is related L to the Minimum Enclosing Ball (MEB) problem, which refers to the determination of the unique spherical surface of smallest radius enclosing a given bounded subset of the d-dimensional Euclidean space \mathbb{R}^d (also denoted as \mathbb{E}^d). This problem is also referred in the literature using, among others, the following nomenclatures (appearing in alphabetical order): (1) Chebyshev radius and Chebyshev center, (2) douter radius, (3) Euclidean 1-center, (4) minimax problem in facility locations, (5) minimum bounding sphere, (6) minimum covering sphere, (7) minimum enclosing sphere, (8) minimum spanning ball, (9) smallest bounding ball, (10) smallest circle problem, (11) smallest covering sphere, (12) smallest enclosing ball, and (13) smallest enclosing sphere. It is worth noting that, these nomenclatures, sometimes, appear slightly different in several fields of science and technology. For example, they appear as: (a) 1-center problem in computational geometry, (b) 1-point estimator in computational statistics, (c) 1-class classification in machine learning, and (d) minimax optimization problem in operations research and optimization. We note in passing that, according to our experience, the nomenclature used most frequently worldwide is the minimum enclosing ball problem, MEB problem in short.

It is worth mentioning here that, for studying and analysing MEB and related to it problems four main characteristics of a set P in \mathbb{R}^d can be used, among others. Specifically: (a) The *circumradius* of P, *i.e.* the radius of the smallest spherical

E-mail: vrahatis@math.upatras.gr & vrahatis@upatras.gr

surface enclosing P, (b) The *inradius* of P, *i.e.* the radius of the greatest spherical surface which is contained in P, (c) The *diameter* of P, *i.e.* the maximal distance of any two points of P, and (d) The *width* of P, *i.e.* the minimal distance between a pair of parallel supporting hyperplanes (tac-hyperplanes) of P.

It is believed that the earliest known statement of the MEB problem was first posed in *circa* 300 B.C. by the considered "father of geometry" Euclid¹ in his seminal work "*Elements*" and specifically in Book IV, Proposition 5 which refers to circumscribe a circle about a given triangle [1]. The general case of finding the smallest circle enclosing a given finite set of n points in \mathbb{R}^2 was first appeared in 1857 by Sylvester² [2]. In addition, the earliest known method for tackling the above case was also proposed in 1860 by Sylvester [3]. The same method was independently proposed in 1885 by Chrystal³ [4]. Thereafter, various methods for determining the MEB of a set of n points in the plane have been proposed.

In 1901 Jung⁴ in his dissertation established seminal results regarding the MEB problem [5]. Thus, Jung was first answered to the question of best possible estimate of the radius of a sphere in \mathbb{R}^d enclosing a bounded subset of \mathbb{R}^d of a given diameter (*cf.* Jung's Theorem 7). Thereafter, generalizations of MEB problem to *d*-dimensions have been of interest for many years and this problem remains an important issue of study and further research in the field. In general, *Jung's enclosing theorem* provides an upper bound of the circumradius of a set in \mathbb{R}^d in terms of its diameter. A similar important theorem that it has been proposed by Steinhagen⁵ in 1921 [6] provides lower bounds (for *d* odd and *d* even) of the inradius of a set in terms of its width (*cf.* Steinhagen's Theorem 8).

The MEB problem can be tackled by applying *enclosing* (*covering*) theorems and it is, among others, one of the most fundamental problems in clustering. Specifically, *clustering* with respect to the *diameter* and the *radius* costs, is the task of *partitioning* a set of points in \mathbb{R}^d to subsets, where items in the same subset, named *cluster*, are similar to each other, compared to items in other clusters [7]. Usually, clustering problems arise in the analysis of large data sets. Thus, the *approximate clustering via core-sets* is used for clustering of a set of points in \mathbb{R}^d (for large *d*) by extracting properly a small set of points named *core-set* that "represents" the given set of points [8]. These issues lie within the domain of property testing. The *property testing* is motivated for designing and development of "super-fast" algorithms for analysing the global

M. N. Vrahatis is with the Department of Mathematics, University of Patras, GR-26110 Patras, Greece.

¹Euclid (c. 323 – c. 285 B.C.), Greek mathematician.

²James Joseph Sylvester (1814 – 1897), English mathematician.

³George Chrystal (1851 – 1911), Scottish mathematician.

⁴Heinrich Wilhelm Ewald Jung (1876 – 1953), German mathematician.

⁵Paul Steinhagen, German mathematician, Dr. phil. Univ. Hamburg 1920.

structure of datasets that are too large to read in their entirety in a reasonable time. These algorithms have direct access to items of a "huge" input data set and determine whether this data set satisfies a desired (predetermined global) property, or is "far" from satisfying it. They inspect relatively small portions of the data set and their complexity is measured in terms of the number of accesses to the input. The algorithms that are used are necessarily *randomized* (otherwise they may be drawing a conclusion from an atypical portion of the input) and approximate (since it is not expected the algorithm to produce an exact answer having examined only a portion of the input) (cf. [9], [10]). In these directions and trends a significant role is played by partitioning theorems, i.e. Helly-type theorems [11], that constitute fundamental results describing the ways in which convex sets intersect with each other. For an illustrative interesting application of partitioning theorems in property testing for solving the clustering problem we refer the interested reader to [10].

The paper at hand aims to contribute towards the mathematical foundation of a framework to analyze and study rigorously the MEB problem and the similar and related to it issues and aspects. Therefore, theoretical results are summarized in a number of significant enclosing (covering) and partitioning (clustering) theorems that can be utilized for tackling the MEB and the related to it problems. To this end, the rest of the paper, after Section II where a preliminary background material is provided, focuses on the following issues and aspects. In Section III several problems that are related to the MEB problem and various of their applications are shorty presented. Then, the following issues are given that forms the proposed framework. Specifically, in Section IV all possible inequalities between any two of circumradius, inradius, diameter and width of a set in \mathbb{R}^d are given. Also, various interesting properties of the regular simplex that are useful for the MEB problem are highlighted. In addition, bounds for the quotient of inner and outer radii of a compact (*i.e.* closed and bounded) convex body are presented and the determination of the inner and outer radii of three regular polytopes namely, the regular simplex, hypercube, and regular cross-polytope are provided. In Section V several enclosing and partitioning theorems are provided including, among others, the theorems of Jung, Steinhagen, Perel'man, Carathéodory, Helly and Tverberg. Also, various generalizations to other spaces and non-Euclidean geometries, mainly of Jung's theorem are highlighted. The paper ends in Section VI with concluding remarks.

II. PRELIMINARY BACKGROUND MATERIAL

The following notions are basic to the analysis and study of the MEB problem and the similar and related to it problems.

The straightforward generalization of the concept of a triangle to the *d*-dimensional Euclidean space \mathbb{R}^d is the *d*-simplex (see, *e.g.* [12]). In general, the *d*-simplex is considered to be as the smallest possible *d*-dimensional polytope, *i.e.* the convex hull of a finite number of points in \mathbb{R}^d (*cf.* [13, p. 8]).

Formally, for a set of points $\{v^k\}_{k=0}^d$ in \mathbb{R}^d that are affinely independent (*i.e.* the vectors $\{v^k - v^0\}_{k=1}^d$ are linearly independent) the convex hull $\operatorname{conv}\{v^0, v^1, \ldots, v^d\} \equiv [v^0, v^0, \ldots, v^d]$

A d-simplex is called oriented if an order has been assigned to its vertices. If $\langle v^0, v^1, \ldots, v^d \rangle$ is an orientation of $\{v^0, v^1, \ldots, v^d\}$ this is considered as being the same as any orientation that can be attained from it by an even permutation of the vertices and as the opposite of any orientation attained by an odd permutation of the vertices. The oriented d-simplex is denoted by $\sigma^d = \langle v^0, v^1, \ldots, v^d \rangle$, and it can be denoted, for example, $\langle v^0, v^1, v^2, \ldots, v^d \rangle = -\langle v^1, v^0, v^2, \ldots, v^d \rangle = \langle v^2, v^0, v^1, \ldots, v^d \rangle$. The boundary $\vartheta \sigma^d$ of an oriented d-simplex $\sigma^d = \langle v^0, v^1, \ldots, v^d \rangle$ is given by $\vartheta \sigma^d = \sum_{i=0}^d (-1)^i \langle v^0, v^1, \ldots, v^{i-1}, v^{i+1}, \ldots, v^d \rangle$. A d-dimensional polyhedron Π^d is a union of a finite number k of oriented d-simplices $\sigma^d_{(t)}$, $t = 1, 2, \ldots, k$ such that the $\sigma^d_{(t)}$ have pairwise-disjoint interiors. Thus, it can be written $\Pi^d = \sum_{t=1}^k \sigma^d_{(t)}$ and $\vartheta \Pi^d = \sum_{i=1}^k \vartheta \sigma^d_{(t)}$. The convex hull of every nonempty subset of the d + 1

since the convex hull is independent of this order the vertices

can be permuted in a list.

points that determine a d-simplex is called a face of the simplex. The faces are simplices themselves and they are also called *facets*. Specifically, for each subset of m + 1 elements $\{u^0, u^1, ..., u^m\} \subset \{v^0, v^1, ..., v^d\}$, the *m*-simplex $[u^0, u^1, \ldots, u^m]$ is called an *m*-face of $[v^0, v^1, \ldots, v^d]$. In particular, 0-faces are vertices and 1-faces are edges. If all the edges have the same length, the simplex is called *regular* or equilateral (cf. [14]). The (m-1)-simplex that defines the *i*-th (m-1)-face opposite to the vertex v^i of σ^m is denoted by $\sigma_{\neg i}^{m} = [v^{0}, v^{1}, ..., v^{i-1}, v^{i+1}, ..., v^{m}]$. Similarly, the (m-2)-simplex that determines the *j*-th (m-2)-face opposite to the vertex v^j of $\sigma_{\neg i}^m$, for $j \neq i$, is denoted by $\sigma_{\neg ij}^m = [v^0, v^1, \dots, v^{i-1}, v^{i+1}, \dots, v^{j-1}, v^{j+1}, \dots, v^m]$. The boundary of σ^m in terms of its (m-1)-faces is given by $\vartheta \sigma^m = \bigcup_{i=0}^m \sigma^m_{\neg i}$. The number of the k-faces of σ^m is equal to the binomial coefficient $\binom{m+1}{k+1}$ (cf. [15, p. 120]). For instance, for k = 0 the number of 0-faces (vertices) is $\binom{m+1}{1} = m+1$, in addition for k = 1 the number of 1-faces (edges) is $\binom{m+1}{2} = m(m+1)/2$, while for k = m-1 the number of (m-1)-faces is $\binom{m+1}{m} = m+1$.

The diameter of an m-simplex σ^m in \mathbb{R}^d $(d \ge m)$ denoted by diam (σ^m) , is the length of the longest edge (1-face) of σ^m (cf. [16, p. 812], [17, p. 607]), where the Euclidean norm, $\|\cdot\|$, is used here to measure distances. The length of the shortest edge of σ^m will be denoted by $\operatorname{shor}(\sigma^m)$. The width or breadth of σ^m denoted by $\operatorname{wid}(\sigma^m)$, is the minimum distance between a pair of parallel supporting hyperplanes (tac-hyperplanes) of σ^m .

The barycenter or centroid of an m-simplex in \mathbb{R}^d $(d \ge m) \sigma^m = [v^0, v^1, \ldots, v^m]$ and the barycenter or centroid of the *i*-th (m-1)-face $\sigma^m_{\neg i}$ for $i = 0, 1, \ldots, m$ of σ^m are respectively denoted by κ^m and κ^m_i and are given by $\kappa^m = (m+1)^{-1} \sum_{j=0}^m v^j$ and $\kappa^m_i = m^{-1} \sum_{j=0, j \ne i}^m v^j$. By convexity the barycenter of σ^m is a point in the relative interior of σ^m . The 1-simplex $\mu^m_i = [v^i, \kappa^m_i]$ for $i = 0, 1, \ldots, m$ is

called the *i*-th median of σ^m that corresponds to the vertex v^i . The m + 1 medians of σ^m concur in the barycenter κ^m of σ^m that divides each of them in the ratio 1 : m, where the longer segment being on the side of the vertex of σ^m (cf. Theorem 13). The σ^m is called orthocentric if its m + 1 altitudes intersect in a common point o^m , called its orthocenter. It should be noted that the m + 1 altitudes of an m-simplex are not necessarily concurrent if $m \ge 3$.

The (d-1)-dimensional spherical surface is denoted by S^{d-1} for which $S^{d-1} = \vartheta B^d_{r,c}$ where $B^d_{r,c} \subset \mathbb{R}^d$ denotes the standard closed Euclidean ball of radius r > 0 centered at a point c in \mathbb{R}^d , *i.e.* $B^d_{r,c} = \{x \in \mathbb{R}^d : ||x - c|| \leq r\}$. The smallest radius of the spherical surfaces S^{d-1} that enclose an *m*-simplex σ^m in \mathbb{R}^d $(d \ge m)$ is called *circumradius* and it is denoted by $\rho_{\rm cir}^m(\sigma^m)$ and the corresponding center denoted by c_{cir}^m is called *circumcenter*. If $c_{cir}^m = \kappa^m$, where κ^m is the barycenter of σ^m then the corresponding circumradius is called *barycentric circumradius* and is denoted by $\beta_{cir}^m(\sigma^m)$. The largest radius of the spherical surfaces S^{d-1} that are enclosed within σ^m is called *inradius* of σ^m , it is denoted by $\rho_{\rm inr}^m(\sigma^m)$ and the corresponding center denoted by $c_{\rm inr}^m$ is called *incenter*. If $c_{inr}^m = \kappa^m$ then the corresponding inradius denoted by $\beta_{inr}^m(\sigma^m)$ is called *barycentric inradius* and it is given by $\beta_{\text{inr}}^m(\sigma^m) = \min_{x \in \vartheta \sigma^m} \|\kappa^m - x\|$. In addition, the quantity $\theta(\sigma^m) = \beta_{\text{inr}}^m(\sigma^m)/\text{diam}(\sigma^m)$, is called *thickness* of σ^m that provides a measure for the quality or how well shaped a simplex is (cf. [16], [18]).

III. UNIQUENESS, RELATED PROBLEMS, THEORY AND Applications of the MEB Problem

The solution of the MEB problem has been proved that is *unique* in the following cases:

- (1) The Euclidean geometry (see, e.g. Welzl 1991 [19]).
- (2) The *hyperbolic geometry* (see, *e.g.* Nielsen and Hadjeres 2015 [20]).
- (3) The *Riemannian positive- definite matrix manifold* (see, *e.g.* Lang 1999 [21], Nielsen and Bhatia 2013 [22]).
- (4) In any Cartan-Hadamard manifold (see, e.g. Arnaudon and Nielsen 2013 [23]), i.e. Riemannian manifold that is complete and simply connected with non-positive sectional curvatures (see, e.g. Nielsen 2020 [24]).
- (5) In any *Bruhat-Tits space* (see, *e.g.* Lang 1999 [21]), that is complete metric space with a semi-parallelogram property, which includes the Riemannian manifold of symmetric positive definite matrices (see, *e.g.* Nielsen 2020 [24]).

On the other hand, the solution of the MEB problem may not be unique in a metric space, as for instance, in the case of *discrete Hamming metric space* (see, *e.g.* Mazumdar *et al.* 2013 [25]) which, in this case, results to NP-hard computation (see, *e.g.* Nielsen 2020 [24]).

Several problems are related or similar to the MEB problem including, among others, the following ones:

(1) The computation of the *minimum number of unit disks* on a plane required to cover the n points of a set in \mathbb{R}^2 , along with a placement of the disks, (see, *e.g.* Friederich *et al.* 2023 [26]).

- (2) The determination of the *smallest enclosing sphere of a finite number of nonempty closed subsets of* ℝ^d, as well as the closely related problem of determining a sphere with the smallest radius that intersects all of the sets (see, *e.g.* Mordukhovich *et al.* 2013 [27]).
- (3) The determination of the smallest sphere that covers at least (1 − γ)n points of a set with n points in ℝ^d, where γ ∈ (0, 1) is a parameter that it could be named outliers exclusion parameter (see, e.g. Ding 2021 [28]).
- (4) The determination of the *minimum k-enclosing ball problem* that consists of finding the ball with smallest radius that contains at least k of n given points of a set in ℝ^d (this case is similar to the previous one) (see, *e.g.* Cavaleiro and Alizadeh 2022 [29]).
- (5) The determination of the *complexity of the minimum k-enclosing ball problem* (see, *e.g.* Shenmaier 2013 [30]).
- (6) The identification and elimination of the *interior points* of the minimum radius ball enclosing a set in \mathbb{R}^d (see, *e.g.* Ahipaşaoğlu and Yıldırım 2008 [31]).
- (7) The determination of the *smallest enclosing ball of a given set of n balls* in ℝ^d (see, *e.g.* Fischer and Gärtner 2004 [32]).
- (8) The treatment of the *Euclidean k-median* problem that aims at the localization of k medians (facilities) among the given n points of a set in ℝ^d such that the sum of the distances from each point of the set to its closest median to be minimized (see, *e.g.* Kolliopoulos and Rao 2007 [33], Bhattacharya *et al.* 2021 [34]).
- (9) The treatment of the *uncapacitated facility location* problem that requires locating an undetermined number of facilities to minimize the sum of the fixed setup costs and the variable costs of serving the market demand from these facilities. In several cases of real life applications capacity limitations are incorporated on the facilities to be established. This problem is known as *capacitated facility location* problem (see, *e.g.* Verter 2011 [35], Kolliopoulos and Moysoglou 2016 [36]).
- (10) The determination of a given point in \mathbb{R}^d its *k*-nearest neighbor points of a set in \mathbb{R}^d (see, e.g. Zhang, 2022 [37], Syriopoulos et al. 2023 [38]).
- (11) The treatment of the *nearest-neighbor-preserving embeddings* that are randomized embeddings between two metric spaces which preserve the approximate nearest-neighbors (see, *e.g.* Indyk and Naor 2007 [39], Emiris *et al.* 2023 [40]).
- (12) The treatment of the *k*-center problem that aims at the determination of *k* balls with the smallest radius to cover a finite number of given points in \mathbb{R}^d . A generalized version of the *k*-center problem that aims at finding *k* balls with the smallest radius such that their union intersects all the elements of a given finite collection of nonempty closed convex sets in \mathbb{R}^d (see, *e.g.* An *et al.* 2020 [41]).
- (13) The determination of the smallest circle that encloses a set of n static points and a mobile point p in \mathbb{R}^2 (see, e.g. Banik et al. 2014 [42]).
- (14) The localization of the *center of the smallest enclosing circle of a set* of n points on a plane, where the center is

constrained to lie on a query line segment (see, *e.g.* Roy *et al.* 2009 [43]).

- (15) The computation of the *diameter*, also known as *furthest* pair, of a set of n points in \mathbb{R}^d that is the maximum pair-wise Euclidean distance between two points in the set (see, e.g. Imanparast et al. 2019 [44]).
- (16) The approximation of the *width* of a set of *n* points in \mathbb{R}^d that is the smallest distance of over all the regions between two parallel hyperplanes that enclose all points of the set (see, *e.g.* Chan 2002 [45]).
- (17) The determination of the *flatness* of a set of *n* points in \mathbb{R}^d . The mathematical notion of flatness is used in the computational metrology scientific community and it corresponds exactly to the width problem (see, *e.g.* Duncan *et al.* 1997 [46]).
- (18) The measuring of the *roundness* of a set P of n points in \mathbb{R}^d that can be determined by approximating P with a sphere S so that the maximum distance between a point of P and S is minimized. This problem is equivalent to computing the *smallest closed region lying between two concentric spheres* of radii ϱ_1 and ϱ_2 for $\varrho_1, \varrho_2 \in \mathbb{R}$ with $0 \leq \varrho_1 \leq \varrho_2$ centered at $c \in \mathbb{R}^d$ that contain P (see, *e.g.* Agarwal *et al.* 2000 [47]).
- (19) The *covering* of a set P of n points in \mathbb{R}^2 with a balanced V-shape of minimum width, where a balanced V-shape is a polygonal region in the plane contained in the union of two crossing equal-width strips (see, *e.g.* Aronov and Dulieu 2013 [48]).
- (20) The computation of the minimum (perimeter or area) axis-aligned rectangle enclosing k points of a given set of n points in the plane (see, e.g. Chan and Har-Peled 2021 [49]).
- (21) The treatment of the *unsupervised k-windows clustering* for improving the well-known and widely used *k-means clustering* aiming at a better time complexity and partitioning accuracy, and the determination of the number of clusters (see, *e.g.* Selim and Ismail 1984 [50], Vattani 2011 [51], Vrahatis *et al.* 2002 [52]).
- (22) The computation of a *minimum-width square or rectan*gular annulus that contains at least n-k points out of ngiven points in the plane, where a square or rectangular annulus is the closed region between a square or rectangle and its offset. The k excluded points are considered as outliers of the n input points (see, *e.g.* Bae 2019 [53]).
- (23) The determination of the *smallest enclosing cylinder* that is the computation of the smallest radius over all cylinders that enclose a set of n points in \mathbb{R}^d , where a cylinder of radius ρ refers to the region of all points of distance ρ from a line (see, *e.g.* Chan 2002 [45]).
- (24) The computation of the *width of simplices generated by the convex hull* of their integer vertices (see, *e.g.* Veselov *et al.* 2019 [54]).
- (25) The determination whether a given simplex in \mathbb{R}^d is covered by *d*-dimensional spheres centered at its vertices (see, *e.g.* Casado *et al.* 2011 [55]).
- (26) The determination of the *largest k-dimensional ball in a d-dimensional box* (see, *e.g.* Everett *et al.* 1998 [56]).
- (27) The determination of the minimal volume of simplices

containing a convex body M in \mathbb{R}^d , where M is a compact convex set in \mathbb{R}^d with non-empty interior (see, *e.g.* Kanazawa 2014 [57], Galicer *et al.* 2019 [58]).

- (28) The determination of the *inscribing spheres in a body* M in \mathbb{R}^d and *inscribing* M *in cylinders*, where M is a compact convex set in \mathbb{R}^d with non-empty interior having C^2 smooth boundary (see, *e.g.* Perel'man 1987 [59]).
- (29) The determination and improvement of *bounds for outer* (external) and inner (internal) radii quotient of a compact convex body in \mathbb{R}^d (see, e.g. Pukhov 1979 [60], Perel'man 1987 [59], González Merino 2017 [61]).

Theoretical issues for the analysis and study of the MEB problem and the similar and related to it issues can be found, among others, in the following fields (appearing in alphabetical order): (1) Algorithmic Information Theory, (2) Applied Mathematics, (3) Approximation Algorithms Analysis, (4) Combinatorial Convexity, (5) Combinatorial Geometry, (6) Complex Projective Geometry, (7) Computational Biology, (8) Computational Complexity, (9) Computational Geometry, (10) Computational Information Geometry, (11) Computational Intelligence, (12) Computational Mathematics, (13) Convex Analysis, (14) Convex Geometry, (15) Convexity Theory, (16) Cryptography and Cryptanalysis, (17) Data Science, (18) Discrete Geometry, (19) Discrete Mathematics, (20) Geometric Computing, (21) Geometric Optimization, (22) Geometry and Topology, (23) Global Optimization, (24) Machine Intelligence and Learning, (25) Mathematical Programming, (26) Metric Geometry, (27) Non-Euclidean Geometry, (28) Operations Research, (29) Pattern Analysis, (30) Pure Mathematics, (31) Statistics, (32) Theoretical Computer Science, (33) Theory of Complexity, (34) Theory of Computation.

It is worth mentioning that, several theoretical approaches for the MEB and the related to it problems have received a considerable impetus from their applications, that include, among others, the following ones:

- (1) Base station locations in facility locations (see, *e.g.* Plastria 2002 [62]).
- (2) Detection of differential expression genes for RNA-seq data (see, *e.g.* Zhou and Zhu 2022 [63]).
- (3) Roundness measurements in metrology (see, *e.g.* Elerian *et al.* 2021 [64]).
- (4) Gap tolerant classifiers in machine learning (see, *e.g.* Burges 1998 [65]).
- (5) k-center clustering (see, e.g. Bădoiu et al. 2002 [8]).
- (6) Solving the approximate 1-cylinder problem (see, *e.g.* Bădoiu *et al.* 2002 [8]).
- (7) Support vector clustering (see, *e.g.* Ben-Hur *et al.* 2001 [66], Bulatov *et al.* 2004 [67]).
- (8) Testing of clustering (see, e.g. Alon et al. 2000 [68], Alon et al. 2003 [69], Alon et al. 2004 [7]).
- (9) Support vector machine classification (see, *e.g.* Rizwan *et al.* 2021 [70]).
- (10) Tuning support vector machine parameters (see, *e.g.* Chapelle *et al.* 2002 [71]).
- (11) Hand gesture recognition (see, *e.g.* Ren and Zhang 2009 [72]).
- (12) Preprocessing for fast farthest neighbor query approxi-

mation (see, e.g. Goel et al. 2001 [73]).

- (13) Computation of spatial hierarchies, (*e.g.* sphere trees) (see, *e.g.* Hubbard 1996 [74]).
- (14) Collision detection (see, e.g. Hubbard 1996 [74]).
- (15) Computer graphics (see, e.g. Larsson et al. 2016 [75]).
- (16) The smallest covering cone (see, e.g. Lawson 1965 [76]).
- (17) Performance of lightweight convolutional neural network models (see, *e.g.* Tzelepi and Tefas 2019 [77]).
- (18) Similarity search in feature space (see, *e.g.* Kurniawati *et al.* 1997 [78]).
- (19) Fast fuzzy inference system training (see, *e.g.* Chung *et al.* 2009 [79]).
- (20) Classification in non-stationary environments (see, *e.g.* Heusinger and Schleif 2021 [80]).
- (21) Curve fitting for the case where a curve takes a sharp turn (see, *e.g.* Aronov and Dulieu 2013 [48]).
- (22) Minimum volume ball estimator in robust regression in statistics (see, *e.g.* Rousseeuw and Leroy 1987 [81]).
- (23) Differential privacy (see, *e.g.* Abowd 2018 [82], Ghazi *et al.* 2020 [83], Mahpud and Sheffet 2022 [84]).
- (24) Machine learning (see, *e.g.* Tsang *et al.* 2005 [85], Tsang *et al.* 2006 [86], Nielsen and Nock 2009 [87], Bauckhage *et al.* 2019 [88]).

IV. UPPER BOUNDS OF RADII OF CONVEX SETS AND COMPACT CONVEX BODIES

A. Relations and Upper Bounds of Circumradius, Inradius, Diameter and Width of Convex Sets

In 1958 Eggleston [89] considered all the twelve possible inequalities between any two of the four characteristics of a convex set P in \mathbb{R}^d : (a) *circumradius* denoted by $\rho_{cir}^d(P)$, (b) *inradius* denoted by $\rho_{inr}^d(P)$, (c) *diameter* denoted by diam(P) and (d) *width* denoted by wid(P), and categorized these inequalities as follows: The first four obvious inequalities are the following:

$$\rho_{\rm inr}^d(P) \leqslant \rho_{\rm cir}^d(P), \qquad \rho_{\rm inr}^d(P) \leqslant \frac{1}{2} {\rm wid}(P), \qquad (1a)$$

$$\operatorname{diam}(P) \leqslant 2\rho_{\operatorname{cir}}^d(P), \qquad \operatorname{wid}(P) \leqslant \operatorname{diam}(P). \tag{1b}$$

The above inequalities imply the following two ones:

$$\rho_{\rm inr}^d(P) \leqslant \frac{1}{2} {\rm diam}(P), \qquad {\rm wid}(P) \leqslant 2\rho_{\rm cir}^d(P).$$
(2)

In addition, Eggleston pointed out that since in the case where the positive real number $\lambda \in \mathbb{R}^+$ is not infinite there are no inequalities of the following four expressions:

$$\rho_{\rm cir}^d(P) \leqslant \lambda \, \rho_{\rm inr}^d(P), \qquad \rho_{\rm cir}^d(P) \leqslant \lambda \, {\rm wid}(P), \qquad (3a)$$

diam
$$(P) \leq \lambda \rho_{inr}^d(P), \quad diam(P) \leq \lambda \operatorname{wid}(P), \quad (3b)$$

and the remaining two inequalities are very interesting and have the expressions:

$$\rho_{\operatorname{cir}}^{d}(P) \leqslant \lambda \operatorname{diam}(P), \quad \operatorname{wid}(P) \leqslant \lambda \rho_{\operatorname{inr}}^{d}(P). \quad (4)$$

Particularly, the first of the above inequalities (4) is related to *Jung's theorem* [5] (*cf.* Theorem 7) and it is given by

$$\rho_{\rm cir}^d(P) \leqslant \sqrt{\frac{d}{2(d+1)}} \operatorname{diam}(P),$$
(5)

while the second inequality is related to *Steinhagen's theo*rem [6] (cf. Theorem 8) and it is given by

$$\operatorname{wid}(P) \leqslant \begin{cases} 2\sqrt{d} \,\rho_{\operatorname{inr}}^{d}(P), & \text{if } d \text{ is odd,} \\ \frac{2(d+1)}{\sqrt{d+2}} \,\rho_{\operatorname{inr}}^{d}(P), & \text{if } d \text{ is even.} \end{cases}$$
(6)

B. Explicit Upper Bounds of Radii, Width and Median of Regular Simplices

For a regular d-simplex τ^d in \mathbb{R}^d , if its diameter diam (τ^d) (*i.e.* the length of one of its equal edges) is given then the circumradius $\rho_{cir}^d(\tau^d)$, the inradius $\rho_{inr}^d(\tau^d)$ and the width wid (τ^d) of τ^d can be explicitly determined in terms of diam (τ^d) (cf. Theorems 1, 2 and 3). Also, a regular simplex is orthocentric, its medians coincide with its corresponding altitudes and the circumcenter c_{cir}^d , the incenter c_{inr}^d and orthocenter σ^m coincide with the barycenter κ^d . In addition, the circumradius coincides with the barycentric circumradius $\beta_{cir}^d(\tau^d)$, while the inradius coincides with the barycentric inradius $\beta_{inr}^m(\tau^d)$.

Theorem 1 (Circumradius of regular simplices [18]): Let τ^d be a regular *d*-simplex in \mathbb{R}^d with diameter diam (τ^d) , then the circumradius $\rho_{cir}^d(\tau^d)$ of σ^d coincides with the barycentric circumradius $\beta_{cir}^d(\tau^d)$ and

$$\rho_{\rm cir}^d(\tau^d) \equiv \beta_{\rm cir}^d(\tau^d) = \sqrt{\frac{d}{2(d+1)}} \,\,\mathrm{diam}(\tau^d). \tag{7}$$

Relation (7) provides the radius of the smallest spherical surface centered at the barycenter κ^d enclosing the regular simplex τ^d . Note that the quantity in Relation (7) constitutes the *sharp upper bound* of the inequality of the significant *Jung's enclosing theorem* (*cf.* Theorem 7).

Theorem 2 (Inradius of regular simplices [18]): Let τ^d be a regular *d*-simplex in \mathbb{R}^d with diameter diam (τ^d) , then the inradius $\rho_{inr}^d(\tau^d)$ of τ^d coincides with the barycentric inradius $\beta_{inr}^d(\tau^d)$ and

$$\rho_{\rm inr}^d(\tau^d) \equiv \beta_{\rm inr}^d(\tau^d) = \sqrt{\frac{1}{2d(d+1)}} \,\, {\rm diam}(\tau^d). \tag{8}$$

Theorem 3 (Width of regular simplices [18], [90]): Let τ^d be a regular *d*-simplex in \mathbb{R}^d with diameter diam (τ^d) , then the width wid (τ^d) of τ^d is given by:

$$\operatorname{wid}(\tau^{d}) = \begin{cases} \sqrt{\frac{2}{d+1}} \operatorname{diam}(\tau^{d}), & \text{if } d \text{ is odd,} \\ \\ \sqrt{\frac{2(d+1)}{d(d+2)}} \operatorname{diam}(\tau^{d}), & \text{if } d \text{ is even.} \end{cases}$$
(9)

Next, we provide results related to the medians of regular simplices.

Theorem 4 (Medians of regular simplices [18]): Assume that $\tau^m = [v^0, v^1, \ldots, v^m]$ is a regular *m*-simplex in \mathbb{R}^d ($d \ge m$) with edge length diam(τ^m) and barycenter κ^m . Let κ_i^m be the barycenter of the *i*-th (m-1)-face $\tau_{\neg i}^m$ of τ^m and let $\mu_i^m = [v^i, \kappa_i^m]$ be the *i*-th median that corresponds to the vertex v^i for $i = 0, 1, \ldots, m$. Then a) μ_i^m coincides with its corresponding *i*-th altitude, b) τ^m is orthocentric, c) the orthocenter o^m of τ^m coincides with its barycenter κ^m , and d) it holds that:

$$\|\mu_i^m\| = \sqrt{\frac{m+1}{2m}} \operatorname{diam}(\tau^m), \quad i = 0, 1, \dots, m.$$
 (10)

Theorem 5 (Generalization of Pythagoras' theorem [18]): Assume that $\tau^m = [v^0, v^1, \dots, v^m]$ is a regular *m*-simplex in \mathbb{R}^d $(d \ge m)$ and let κ_i^m be the barycenter of the *i*-th (m-1)-face $\tau_{\neg i}^m$ of τ^m for $i = 0, 1, \dots, m$. Then, for $j \ne i$ and $i, j = 0, 1, \dots, m$, it holds that:

$$\|v^{i} - \kappa_{i}^{m}\|^{2} + \|v^{j} - \kappa_{i}^{m}\|^{2} = \|v^{i} - v^{j}\|^{2}.$$
 (11)

In general, the regular simplex exhibits interesting properties including, among others, the maximizing properties of the regular simplex that have been given by Tanner in 1974 [91]. Particularly, Tanner has proved that the regular simplex maximizes the sum of the squared contents of all *m*-faces, for all *m*, when the sum of squared edge lengths is fixed. Thus, the regular simplex has the largest total length of all edges, total area of all 2-faces, total volume of all 3-faces, *etc.*, for a fixed sum of squared edge lengths [91]. Furthermore, in 1977 Alexander [92] proved the conjecture of Sallee⁶, which states that "Only the regular simplex has maximal width among all simplices inscribed in a sphere of \mathbb{R}^{d} ", (cf. Theorem 3, also see, *e.g.* [18], [92], [93] and [90]).

C. Inner and Outer Radii of Compact Convex Bodies and Regular Polytopes

Perel'man⁷ in 1987 in his publication "k radii of a compact convex body" [59] considered the internal and external k-radii of a compact (*i.e.* closed and bounded) convex body M in \mathbb{R}^d that are particular cases of the diameters used in Bernšteĭn and Kolmogorov approximation theory, respectively. Specifically, the internal (inner) k-radius $r_k(M)$ for k = 1, 2, ..., d of Mis defined as the radius of the greatest k-dimensional sphere contained in M, while the external (outer) k-radius $R_k(M)$ for k = 1, 2, ..., d of M is defined as the smallest radius of the solid cylinder containing M with a (d+1-k)-dimensional spherical cross section and a (k - 1)-dimensional generator. Perel'man in [59] and independently Pukhov in [60] (cf. [61]) studied the relation between these internal and external k-radii measures, and proved the following theorem:

Theorem 6 (Perel'man-Pukhov theorem [59]): Let M be a compact convex body in \mathbb{R}^d , then it holds that:

$$\frac{R_k(M)}{r_k(M)} \leqslant k+1, \quad 1 \leqslant k \leqslant d, \tag{12}$$

where $R_k(M)$ and $r_k(M)$ are the external and internal k-radii of M, respectively.

Also, Perel'man in [59] pointed out that:

(a) The problem of an upper bound for the radii quotient R_ℓ(M)/r_k(M) for k ≤ ℓ which does not depend on M is interesting. This quotient can be arbitrarily large for the case k > ℓ.

- (b) Bounds which are, in fact, least upper bounds for R_k(M)/r_k(M) is the most interesting, since the bound for R_ℓ(M)/r_k(M) for k ≤ ℓ follows from it.
- (c) Least upper bounds for the cases $R_1(M)/r_1(M)$ and $R_d(M)/r_d(M)$ can be obtained by Jung's theorem [5] and the generalization of Blaschke's theorem [94] (due to Steinhagen [6]), respectively.
- (d) It is obvious that $r_1(M) \ge r_2(M) \ge \cdots \ge r_d(M)$ and $R_1(M) \ge R_2(M) \ge \cdots \ge R_d(M)$. Also, it holds that $r_k(M) \le R_k(M)$ for any $1 \le k \le d$.
- (e) The equality $r_k(M) = R_k(M)$ holds for certain bodies, *e.g.* spheres.
- (f) Inequality (12) is also valid for finite $r_k(M)$ without the assumption that the convex body M is bounded.
- (g) The upper bound of Theorem 6 is not a least upper bound for any $1 \leq k \leq d$ and the estimate given in Inequality (12) is extremely "rough" (non-sharp). Also, precise estimates of $R_k(M)/r_k(M)$ for the cases $k = 2, 3, \ldots, d-1$ are, evidently, unknown. On the other hand, for the cases k = 1 and k = d, the maximum of $R_k(M)/r_k(M)$ can be reached on a regular d-simplex.

Similarly, in 2017 González Merino [61] studied upper bounds for the quotient of successive inner and outer radii of a convex body M in \mathbb{R}^d . In his approach the largest radius of an *i*-dimensional Euclidean disc contained in M is denoted by $r_i(M)$, while the smallest radius of a solid cylinder with *i*dimensional spherical cross-section containing M is denoted by $R_i(M)$, for any $1 \leq i \leq d$. Using these notations the *Perel'man-Pukhov inequality* (12) can be written as follows:

$$\frac{R_{d+1-i}(M)}{r_i(M)} \leqslant i+1, \quad 1 \leqslant i \leqslant d, \tag{13}$$

Inequality (13) extends two important results in convex geometry that are particular cases of it. Specifically, for i = 1Jung's inequality [5] is obtained, *i.e.*

$$\frac{R_d(M)}{r_1(M)} \leqslant \sqrt{\frac{2d}{d+1}},\tag{14}$$

while, for i = d Steinhagen's inequality [6] is attained, *i.e.*

$$\frac{R_1(M)}{r_d(M)} \leqslant \begin{cases} \sqrt{d}, & \text{if } d \text{ is odd,} \\ \frac{d+1}{\sqrt{d+2}}, & \text{if } d \text{ is even.} \end{cases}$$
(15)

Both Inequalities (14) and (15) are best possible, since the regular *d*-simplex attains equality [61].

For the first time, in 2005 Brandenber [95] completely determined the inner and outer *j*-radii for three regular polytopes which exist in every dimension *d*, namely, the *regular simplex*, (hyper-) cube, and *regular cross-polytope*. Brandenber defined the *inner j-radii* r_j as the radii of the largest *j*balls contained in *j*-dimensional slices of the polytope and the *outer j-radii* R_j as the radii of the smallest *j*-balls containing the projections of the polytope onto *j*-dimensional subspaces. In general, the inner radii r_j are also known as *Bernšteĭn diameters* and the outer radii R_j as *Kolmogorov diameters* or *Kolmogorov width*. Among the results that have been presented in [95], the results of the radii of the three types of regular

⁶George Thomas Sallee (1940 – 2019), American mathematician.

⁷Grigorii Yakovlevich Perel'man (b. 1966), Russian mathematician.

polytopes are exhibited below, where the polytopes are scaled such that their circumradius to be 1. Thus, (a) The inner radii r_j of regular simplex are given as follows:

$$r_j = \sqrt{\frac{d+1}{j(j+1)d}}.$$
(16)

(b) The inner radii r_j of both cube and regular cross-polytope are given by:

$$r_j = \sqrt{\frac{1}{j(d+1)}}.$$
(17)

(c) The outer radii R_j of regular simplex are given as follows:

$$R_{j} = \begin{cases} \sqrt{\frac{j}{d}}, & \text{if } j \notin \{1, d-1\} \text{ or } d \text{ odd,} \\ \frac{d+1}{d} \sqrt{\frac{1}{d+2}}, & \text{if } j = 1 \text{ and } d \text{ even,} \\ \frac{2d-1}{2d}, & \text{if } j = d-1 \text{ and } d \text{ even.} \end{cases}$$
(18)

(d) The outer radii R_j of both cube and regular cross-polytope are given by:

$$R_j = \sqrt{\frac{j}{d}}.$$
 (19)

Gritzmann and Klee in [93] showed that the computation of R_j is NP-hard (nondeterministic polynomial-time hard) for general simplices and many values of j.

V. MATHEMATICAL APPROACHES FOR ENCLOSING AND PARTITIONING A SET

A. Theorems of Jung, Steinhagen and Perel'man

In 1901 Jung gave seminal results regarding the MEB problem [5]. Specifically, he gave results for the case of *finite point sets* and indicated their extension to *infinite sets*, while in 1910 Jung [96] proposed necessary conditions on the smallest circle enclosing a finite set of n points in a plane. It is believed that the earliest known interesting application of Jung's theorem was first given in 1905 by Landau⁸ [97] who applied Jung's theorem in a plane in order to sharpen an inequality related to theory of analytic functions.

Theorem 7 (Jung's enclosing theorem (1901) [5]): Assume that diam(P) is the diameter of a bounded subset P of \mathbb{R}^d (containing more than a single point). Then, there exists a unique spherical surface of circumradius $\rho_{cir}^d(P)$ enclosing P, and it holds that

$$\rho_{\rm cir}^d(P) \leqslant \sqrt{\frac{d}{2(d+1)}} \,\operatorname{diam}(P).$$
(20)

This significant Jung's enclosing theorem received quite a few additional proofs due to Süss 1936 [98], Blumenthal and Wahlin 1941 [14], Eggleston 1958 [89, Th. 49, p. 111] and Guggenheimer 1977 [13, Pr. 13-6, p. 140], among others. Note that Jung's Inequality (20) is *sharp* (best possible) since the regular *d*-simplex attains equality (*cf.* Theorem 1).

⁸Edmund Georg Hermann Landau (1877 - 1938), German mathematician.

Jung's enclosing theorem is considered as a cornerstone in the field of MEB and related problems that strongly influencing later developments and generalizations to other spaces and non-Euclidean geometries. For example, to mention a few (appearing in chronological order):

- 1938 A generalization of Jung's theorem to *Minkowski spaces* has been proved by Bohnenblust [99].
- 1955 A simpler proof of Bohnenblust's result [99] has been proposed by Leichtweiss [100].
- 1959 The *expansion constant* and *Jung's constant* determined by the geometric properties of a Minkowski space have been given by Grünbaum [101].
- 1984 An *infinity dimensional Hilbert space* extension of Jung's theorem and its application to *measures of non-compactness* have been proposed by Daneš [102].
- 1985 The determination of the *Jung diameter* of the *complex* projective plane has be given and lower bounds for all *complex projective spaces* have been obtained by Katz [103].
- 1985 A generalization of Jung's theorem has been proved by Dekster [104] for which estimates are given for the sidelengths of certain inscribed simplices.
- 1995 The obtained result by Dekster [104] has been extended to the *spherical and hyperbolic spaces* by the same author [105].
- 1997 The result obtained in [104] extended by Dekster to a class of *metric spaces of curvature bounded above* that includes *Riemannian spaces* [106].
- 1997 An analogue of Jung's theorem for *Alexandrov spaces* of curvature bounded above has been presented in 1997 by Lang and Schroeder [107].
- 2006 A complete characterization regarding the *extremal sub*sets of Hilbert spaces, which constitutes an infinitedimensional generalization of the Jung theorem, has been proposed by Nguen-Khac and Nguen-Van [108].
- 2006 Generalizations of the Jung theorem for a *pair of Minkowski spaces* have been given by Boltyanski and Martini. Particularly, the authors gave generalizations of Jung's theorem for the case where there are two Minkowski metrics in \mathbb{R}^d , one for the diameter of a set $P \in \mathbb{R}^d$, and the other for the radius of the Minkowski ball containing P [109].
- 2013 Combinatorial generalizations of Jung's theorem have been presented and the "fractional" as well as the "colorful" versions of this theorem have been proved by Akopyan [110].
- 2017 An one-to-one connection between the *Jung constant* determined by Jung's theorem in an arbitrary Minkowski space and the maximal *Minkowski asymmetry* of the complete bodies within that space has been stated by Brandenberg and González Merino [111].

It is worth reminding that, Jung's theorem provides an upper bound of the circumradius of a set in terms of its diameter. A similar important theorem due to Steinhagen provides lower bounds of the inradius of a set in terms of its width.

Theorem 8 (Steinhagen's theorem (1921) [6]): Assume that P is a bounded convex subset of \mathbb{R}^d of minimal width wid(P).

Then, the inradius $\rho_{inr}^d(P)$ satisfies:

$$\rho^{d}_{\mathrm{inr}}(P) \geqslant \begin{cases} \left(2\sqrt{d}\right)^{-1} \operatorname{wid}(P), & \text{if } d \text{ is odd}, \\ \\ \left[\sqrt{d+2}/(2d+2)\right] \operatorname{wid}(P), & \text{if } d \text{ is even}. \end{cases}$$

The Steinhagen theorem has been proved in 1914 for d = 2by Blaschke [94]. An additional proof of Steinhagen's theorem has been given in 1936 by Gericke [112]. Also, Santaló in 1946 [113] gave a generalization of Jung's and Steinhagen's theorems to convex regions on the d-dimensional spherical surface. A geometrical proof of the above results for d = 2has been given in 1944, also by Santaló [114]. Santaló's result about the generalization of the Steinhagen's theorem, also generalizes the well-known Robinson's theorem that it has been proved by Robinson in 1938 [115]. In 1992 Henk [116] by studying the relation between circumradius and diameter of a convex body which has been proposed by Jung's theorem, gave a natural generalization of Jung's theorem. Similarly, in 1993 Betke and Henk [117] by analyzing the relation between inradius and width of a convex set that has been established by Steinhagen's theorem proposed respectively a natural generalization of Steinhagen's theorem.

Perel'man in 1987 [59] applied the concept of k radii (cf. Sec. IV) for the solution of two problems for compact convex bodies with C^2 smooth boundaries that have been asked by Ionin at the geometric seminar of the Novosibirsk Mathematics Institute in 1984 and are referred as Problems 8 and 9. The problems are related to inscribing spheres in a body M in \mathbb{R}^d and inscribing M in cylinders, where M is a compact convex body having C^2 smooth boundary. Specifically, Perel'man proved the following theorem:

Theorem 9 (Perel'man's enclosing theorem (1987) [59]): Let M be a compact convex body in \mathbb{R}^d with a C^2 smooth boundary. If there are k principal curvatures not greater than 1 at each point of the boundary of M (where the number k is the same for all points of the boundary), then it is possible to place in M a (k + 1)-dimensional ball of radius not less than 1/(k-2). Correspondingly, if the k principal curvatures are not less than 1 at any point of the boundary, then M can be enclosed in a solid cylinder with a (k + 1)-dimensional spherical cross section of radius not greater than 1 and a (d - k - 1)-dimensional generator for any $1 \le k \le d - 1$.

The following result permits the reduction of the MEB problem to a finite one concerning d + 1 points.

Theorem 10 (Blumenthal-Wahlin enclosing theorem (1941) [14]): If each set of d + 1 points of a subset P of \mathbb{R}^d is enclosable by a spherical surface of a given radius, then P is itself enclosable by this spherical surface.

In addition, in 1953 Gale [118] gave the following equivalent formulation of Jung's enclosing theorem.

Theorem 11 (Gale's enclosing theorem (1953) [118]): The circumscribed *d*-sphere of a regular *d*-simplex of diameter 1 will cover any *d*-dimensional set of diameter 1.

Next, we provide useful results for simplices. To this end, the following notation and assumptions are used.

Notation 1: The sum of the squares of the lengths of the m edges of an m-simplex in \mathbb{R}^d $(d \ge m)$ $\sigma^m = [v^0, v^1, \dots, v^m]$

that concur in the vertex v^i for i = 0, 1, ..., m is denoted by:

$$E(v^{i}) = \sum_{\substack{j=0\\ j\neq i}}^{m} \|v^{i} - v^{j}\|^{2}.$$
 (21)

Furthermore, the sum of the squares of the lengths of the $\binom{m}{2} = (m-1)m/2$ edges of the *i*-th (m-1)-face $\sigma_{\neg i}^m$ of σ^m is denoted as follows:

$$E(\sigma_{\neg i}^{m}) = \sum_{\substack{p=0\\p\neq i}}^{m-1} \sum_{\substack{q=p+1\\q\neq i}}^{m} \|v^{p} - v^{q}\|^{2}.$$
 (22)

In addition, the sum of the squares of the lengths of all the $\binom{m+1}{2} = m(m+1)/2$ edges of σ^m is denoted by:

$$E(\sigma^m) = \sum_{p=0}^{m-1} \sum_{q=p+1}^m \|v^p - v^q\|^2.$$
 (23)

Assumptions 1: Suppose that $\sigma^m = [v^0, v^1, \dots, v^m]$ is an m-simplex in \mathbb{R}^d $(d \ge m)$ with barycenter κ^m and diameter $\dim(\sigma^m)$. Let κ_i^m be the barycenter of the *i*-th (m-1)-face $\sigma_{\neg i}^m$ of σ^m and let $\mu_i^m = [v^i, \kappa_i^m]$ be the *i*-th median that corresponds to the vertex v^i for $i = 0, 1, \dots, m$.

Theorem 12 (Generalization of Apollonius' theorem [18]): Let Assumptions 1 hold. Then,

$$\|\mu_i^m\|^2 = \frac{1}{m^2} \left(m E(\upsilon^i) - E(\sigma_{\neg i}^m) \right), \quad i = 0, 1, \dots, m.$$
 (24)

Theorem 13 (Generalization of Commandino's theorem [18]): Let Assumptions 1 hold. Then, the m + 1 medians of σ^m concur in the barycenter κ^m of σ^m that divides each of them in the ratio 1:m, where the longer segment being on the side of the vertex of σ^m , that is to say,

$$\|\kappa^m - v^i\| = m\|\kappa^m - \kappa^m_i\|, \quad i = 0, 1, \dots, m.$$
 (25)

Theorem 14 (Barycentric distances of the vertices [18], [119]): Let Assumptions 1 hold. Then for each i = 0, 1, ..., m it holds that:

$$\|\kappa^m - v^i\| = \frac{1}{m+1} \sqrt{mE(v^i) - E(\sigma^m_{\neg i})}.$$
 (26)

Theorem 15 (Barycentric sum property [18]): Let Assumptions 1 hold. Then,

$$\sum_{i=0}^{m} \|\kappa^m - v^i\|^2 = \frac{1}{m+1} E(\sigma^m).$$
(27)

Theorem 16 (Medians sum property [18]): Let Assumptions 1 hold. Then,

$$\sum_{i=0}^{m} \|\mu_i^m\|^2 = \frac{m+1}{m^2} E(\sigma^m).$$
(28)

Theorem 17 (Simplex barycentric enclosing [18]): Let Assumptions 1 hold. Then the barycentric circumradius β_{cir}^m is unique, it is given as follows:

$$\beta_{\rm cir}^m = \frac{1}{m+1} \max_{0 \le i \le m} \left\{ m E(v^i) - E(\sigma_{\neg i}^m) \right\}^{1/2}, \tag{29}$$

and the spherical surface with center at κ^m and radius β^m_{cir} encloses σ^m .

Theorem 18 (Simplex enclosing [18]): Let Assumptions 1 hold. Suppose further that β_{cir}^m is the barycentric circumradius. Then, there exists a unique spherical surface of circumradius ρ_{cir}^m enclosing σ^m , and

$$\rho_{\rm cir}^m \leqslant \min \left\{ \beta_{\rm cir}^m, \, [m/(2m+2)]^{1/2} \operatorname{diam}(\sigma^m) \right\}.$$
(30)

The computation of ρ_{cir}^m does not require any additional computational burden since it can be easily obtained during the computation of the longest edge (diameter) of the simplex.

Note that, the barycentric circumradius $\beta_{cir}^d = \beta_{cir}^d(P)$ of a set P in \mathbb{R}^d is the supremum of the barycentric circumradii of simplices with vertices in P. Next, it has been shown in Theorem 19, that a bounded set P can be covered by a spherical surface of circumradius $\beta_{cir}^d(P)$, which in many cases gives a better result than Jung's theorem (*cf.* [120]). For instance, consider a regular 2-simplex $\tau^2 = [v^0, v^1, v^2]$ with diam $(\tau^2) = 1$ and barycenter κ^2 . By Theorem 1 we obtain for the circumradius of τ^2 , $\rho_{cir}^2(\tau^2) = 3^{-1/2}$ that is the upper bound given by Jung's Theorem 7. Consider now the 2-simplex $\sigma^2 = [\kappa^2, v^1, v^2]$. The lengths of the edges of σ^2 are $3^{-1/2}$, $3^{-1/2}$ and 1. Thus, diam $(\sigma^2) = 1$ and consequently $\rho_{cir}^2(\sigma^2) = 3^{-1/2}$. By Theorem 17 we obtain for the barycentric circumradius of σ^2 , $\beta_{cir}^d(\sigma^2) = [7/27]^{1/2}$ which is smaller than the upper bound $3^{-1/2}$ for the circumradius ρ_{cir}^2 that is given by Jung's covering Theorem 7.

Theorem 19 (Variant of Jung's theorem (1988) [120]): Assume that diam(P) is the diameter of a bounded subset P of \mathbb{R}^d (containing more than a single point) and let $\beta_{cir}^d = \beta_{cir}^d(P)$ be its barycentric circumradius. Then, there exists a unique spherical surface of circumradius ρ_{cir}^d enclosing P, and

$$\rho_{\rm cir}^d \le \min\{\beta_{\rm cir}^d(P), \, [d/(2d+2)]^{1/2} \operatorname{diam}(P)\}.$$
(31)

B. Theorems of Carathéodory, Helly, Radon and Tverberg

Carathéodory⁹ in 1907 [121] gave an important theorem in convex geometry. In 1970 Rockafellar¹⁰ [122] has pointed out that "*Carathéodory's theorem is the fundamental dimensionality result in convexity theory, and it is the source of many other results in which dimensionality is prominent.*" In general, the convex hull of a subset P of \mathbb{R}^d can be obtained by carrying out all convex combinations of elements of P. On the other hand, due to Carathéodory's theorem it is not necessary to perform combinations involving more than d + 1 elements at a time. Thus, convex combinations of the form $\sum_{i=1}^{m} \lambda_i x_i$ can be performed where $m \leq d+1$ or m = d+1 in the case of non-distinct vectors x_i (*cf.* [122]). *Carathéodory's theorem in convexity theory* states that:

Theorem 20 (Carathéodory's theorem (1907) [121]): Any point in the convex hull of a finite point set in \mathbb{R}^d is a convex combination of some at most d + 1 of these points.

Carathéodory's theorem is related to the following partitioning theorems due to Helly¹¹ and Tverberg¹².

Theorem 21 (Helly's partitioning theorem (1913) [123]): Let C_1, C_2, \ldots, C_k be a finite family of convex subsets of \mathbb{R}^d , with $k \ge d+1$. If the intersection of every d+1 of these sets is nonempty, then the whole family has a nonempty intersection, *i.e.* $\bigcap_{i=1}^k C_i \ne \emptyset$.

Theorem 22 (Tverberg's partitioning theorem (1966) [124]): Every set with at least (p-1)(d+1) + 1 points in \mathbb{R}^d can be partitioned into p subsets whose convex hulls all have at least one point in common.

Theorems 20, 21 and 22 are equivalent in the sense that each one can be deduced from another [125]. Theorem 21 proposed by Helly in 1913 [126] but not published by him until 1923. Tverberg gave the first proof of Theorem 22 in 1966 [124], while in 1981 he gave a simpler proof [127]. Tverberg's partitioning theorem constitutes a generalization of a theorem due to Radon¹³ that it has been proposed in 1921:

Theorem 23 (Radon's partitioning theorem (1921) [128]): Every set with d+2 points in \mathbb{R}^d can be partitioned into two sets whose convex hulls intersect.

Theorems 20, 21, 22 and 23 are known as *Helly-type theorems* [11]. In addition, a classical partitioning theorem for points in the plane that is also based on Carathéodory's theorem has been proved by Birch¹⁴ in 1959 [129]. Also, Adiprasito *et al.* in 2020 [130] initiated the study of the dimensionless versions of classical theorems in convexity theory. Specifically, they considered the dimensionless versions of the theorems of Carathéodory, Helly, and Tverberg. The obtained results have several "colorful" and "fractional" consequences and are particulary interesting and motivating, among others, for those who are interested in classical convexity parameters. The authors named these theorems as "*no-dimension theorems*" and proved the following theorems, among others:

Theorem 24 (No-dimension Carathéodory theorem [130]): Let P be a set of n points in \mathbb{R}^d , $r \in [n]$ (*i.e.* $r \leq n$), and $a \in \operatorname{conv} P$. Then there exists a subset Q of P with r elements (*i.e.* |Q| = r) such that for the distance between a and the convex hull of Q holds that:

$$\operatorname{dist}(a, \operatorname{conv} Q) \leq \frac{\operatorname{diam}(P)}{\sqrt{2r}}.$$

Theorem 25 (No-dimension Helly theorem [130]): Assume that K_1, K_2, \ldots, K_n are convex sets in \mathbb{R}^d and $k \in [n]$. For $J \subset [n]$ define $K(J) = \bigcap_{j \in J} K_j$. If the Euclidean unit ball B(b, 1) centered at $b \in \mathbb{R}^d$ intersects K(J) for every $J \subset [n]$ with |J| = k, then there is a point $q \in \mathbb{R}^d$ such that

$$\operatorname{dist}(q, K_i) < \frac{1}{\sqrt{k}}, \quad \forall i \in [n].$$

Theorem 26 (No-dimension Tverberg theorem [130]): Let P be a set of n points in \mathbb{R}^d , then for a given integer $1 \le k \le n$, there exists a point $q \in \mathbb{R}^d$ and a partition of P into k sets P_1, P_2, \ldots, P_k such that:

dist
$$(q, \operatorname{conv} P_i) \leq (2 + \sqrt{2}) \sqrt{\frac{k}{n}} \operatorname{diam}(P), \quad \forall i \in [k].$$

Carathéodory's theorem has obtained numerous interesting reformulations, variations, generalizations and applications (see, *e.g.* Danzer *et al.* 1963 [126], De Loera *et al.* 2019

⁹Constantin Carathéodory (1873 - 1950), Greek mathematician.

¹⁰Ralph Tyrrell Rockafellar (b. 1935), American mathematician.

¹¹Eduard Helly (1884 – 1943), Austrian mathematician.

¹²Helge Arnulf Tverberg (1935 – 2020), Norwegian mathematician.

¹³Johann Karl August Radon (1887 – 1956), Austrian mathematician.

¹⁴Bryan John Birch (b. 1931), English mathematician.

[125]). Rockafellar in 1970 [122] used Carathéodory's theorem to prove Helly's theorem and results related to *infinite systems of linear inequalities*. In addition, various generalizations, refinements, applications and conjectures of some of the classical combinatorial theorems about convex sets (Carathéodory, Helly, Radon and Tverberg theorems) have been published in 2022 by Bárány and Kalai [11]. Also, in [11] the authors point out the connection between important results from combinatorial convexity and some theorems from topology. For instance, Helly's theorem is a manifestation of the *nerve theorem* from algebraic topology, and Radon's theorem can be regarded as an early *linear* version of the *Borsuk-Ulam theorem*.

In 1982 Bárány [131] proposed the following sharp generalization of Carathéodory's theorem in the sense that the number of partition of the sets C_i 's cannot be decreased.

Theorem 27 (Bárány's partitioning theorem (1982) [131]): Let C_0, C_1, \ldots, C_d be subsets of \mathbb{R}^d and let α be a point in conv C_i (the convex hull of C_i) for $i = 0, 1, \ldots, d$ (*i.e.* $\alpha \in \bigcap_{i=0}^d \operatorname{conv} C_i$). Then there exist vectors $v^i \in C_i$ for $i = 0, 1, \ldots, d$ such that $\alpha \in \operatorname{conv} \{v^0, v^1, \ldots, v^d\}$.

In addition, in a paper published in 1986 by Polikanova and Perel'man, titled "*A remark on Helly's theorem*", the following partitioning theorem has been proved [132]:

Theorem 28 (Polikanova-Perel'man partitioning theorem (1986) [132]): Let \mathfrak{M} be a bounded family of compact convex sets in \mathbb{R}^d such that $\mu_d(\bigcap \mathfrak{M}) = 0$, where μ_d denotes the *d*-dimensional volume. Then for every $\varepsilon > 0$ there are d + 1 sets $M_1, M_2, \ldots, M_{d+1}$ in the family \mathfrak{M} such that $\mu_d(\bigcap_{i=1}^{d+1} M_i) < \varepsilon$. Also, if dim $\bigcap \mathfrak{M} = \ell$, $0 \leq \ell < d$, then there are $d - \ell + 1$ sets in the family \mathfrak{M} such that $\mu_d(\bigcap_{i=1}^{d-\ell+1} F_i) \leq \varepsilon$.

VI. CONCLUDING REMARKS

A characteristic of the MEB and related to it problems is that the theory and applications of this field of study and research are very broad and divers. Also, a large variety of methods and algorithms for approximating the MEB, diameter and width of a set have been proposed in the literature.

Our contribution is focused mainly on essentials enclosing and partitioning theorems that are considered as cornerstones in the field that strongly influencing developments and generalizations to other spaces and non-Euclidean geometries. Specifically, a contribution is provided towards the mathematical foundation of a framework to study, analyze and apply rigorously the MEB and relared to it problems.

We hopefully think that the paper at hand might contribute to a deeper understanding and rigorous tackling of the MEB problem and the similar and related to it issues and aspects.

REFERENCES

- Euclid (author), Hypsicles of Alexandria (contributor), Giovanni Mocenigo (dedicatee), *Euclid's "Elements"*. Venice: Erhard Ratdolt, May 25, 1482. [Pdf] Retrieved from the Library of Congress, [Website] https://www.loc.gov/item/2021667076/.
- [2] J. J. Sylvester, "A question in the geometry of situation," Quart. J. Pure Appl. Math., vol. 1, p. 79, 1857.
- [3] J. J. Sylvester, "On Poncelet's approximate valuation of surd forms," *Philosophical Magazine*, vol. 120, pp. 203–222, 1860.

- [4] G. Chrystal, "On the problem to construct the minimum circle enclosing n given points in a plane," *Proc. Edinburgh Math. Soc.*, vol. 3, pp. 30–33, Third Meeting, Jan. 1885.
- [5] H. W. E. Jung, "Über die kleinste Kugel, die eine räumliche Figur einschliesst," J. Reine Angew. Math., vol. 123, no. 3, pp. 241–257, 1901.
- [6] P. Steinhagen, "Über die größte Kugel in einer konvexen Punktmenge," Abh. Math. Semin. Univ. Hambg., vol. 1, no. 1, pp. 15–26, 1921.
- [7] N. Alon, S. Dar, M. Parnas, and D. Ron, "Testing of clustering," SIAM Review, vol. 46, no. 2, pp. 285–308, 2004.
- [8] M. Bădoiu, S. Har-Peled, and P. Indyk, "Approximate clustering via core-sets," in *Thirty-fourth Annual ACM Symposium on Theory of Computing (STOC 2002)*, pp. 250–257, Montreal, Quebec, Canada: ACM, 2002.
- [9] O. Goldreich, Introduction to Property Testing, Cambridge, UK: Cambridge University Press, 2017.
- [10] S. Chakraborty, R. Pratap, S. Roy, and S. Saraf, "Helly-type theorems in property testing," *Internat. J. Comput. Geom. Appl.*, vol. 28, no. 4, pp. 365–379, 2018.
- [11] I. Bárány and J. Kalai, "Helly-type problems," Bull. Amer. Math. Soc. (N.S.), vol. 59, no. 4, pp. 471–502, 2022.
- [12] M. J. Greenberg, Lectures on Algebraic Topology, N.Y.: W. A. Benjamin, 1967.
- [13] H. W. Guggenheimer, Applicable Geometry: Global and Local Convexity, Applied Mathematics Series, Huntington, N.Y.: Robert Krieger Publishing Co., Inc., 1977.
- [14] L. M. Blumenthal and G. E. Wahlin, "On the spherical surface of smallest radius enclosing a bounded subset of *n*-dimensional Euclidean space," *Bull. Amer. Math. Soc.*, vol. 47, pp. 771–777, Feb. 1941.
- [15] H. S. M. Coxeter, *Regular Polytopes*, London: Methuen & Co., Ltd., New York: Pitman Publishing Corp., 1948.
- [16] J. H. C. Whitehead, "On C¹-complexes," Ann. of Math. (2), vol. 41, no. 4, pp. 809–824, 1940.
- [17] P. S. Alexandroff and H. Hopf, *Topologie I*, Die Grundlehren der mathematischen Wissenschaften, Band 45, Berichtigter Reprint, Berlin-New York: Springer-Verlag, 1974.
- [18] M. N. Vrahatis, "Generalization of the Apollonius theorem for simplices and related problems," 2024, arXiv:2401.03232.
- [19] E. Welzl, "Smallest enclosing disks (balls and ellipsoids)," in New Results and New Trends in Computer Science, H. Maurer (ed), Lect. Notes Comput. Sci., vol. 555, pp. 359–370, Berlin: Springer, 1991.
- [20] F. Nielsen and G. Hadjeres, "Approximating covering and minimum enclosing balls in hyperbolic geometry," in *Geometric Science of Information*, (GSI 2015), F. Nielsen and F. Barbaresco (eds), *Lect. Notes Comput. Sci.*, vol. 9389, pp. 586–594, Cham, Switzerland: Springer 2015.
- [21] S. Lang, Math Talks for Undergraduates, New York, New York, USA: Springer-Verlag Inc., 1999.
- [22] F. Nielsen and R. Bhatia, *Matrix Information Geometry*, Berlin, Heidelberg, Germany: Springer-Verlag Inc., 2013.
- [23] M. Arnaudon and F. Nielsen, "On approximating the Riemannian 1center," *Comput. Geom. Theory Appl.*, vol. 46, no. 1, pp. 93–104, Jan. 2013.
- [24] F. Nielsen, "The Siegel-Klein disk: Hilbert geometry of the Siegel disk domain," *Entropy*, vol. 22, no. 9, art. no. 1019, pp. 1–41, Sep. 2020.
- [25] A. Mazumdar, Y. Polyanskiy, and B. Saha, "On Chebyshev radius of a set in Hamming space and the closest string problem," in *Proc. International Symposium on Information Theory (ISIT 2013)*, pp. 1401– 1405, Istanbul, Turkey: IEEE 2013.
- [26] R. Friederich, A. Ghosh, M. Graham, B. Hicks, and R. Shevchenko, "Experiments with unit disk cover algorithms for covering massive pointsets," *Comput. Geom. Theory Appl.*, vol. 109, 101925, pp. 1–24, 2023.
- [27] B. Mordukhovich, N. M. Nam, and C. Villalobos, "The smallest enclosing ball problem and the smallest intersecting ball problem: Existence and uniqueness of solutions," *Optim. Lett.*, vol. 7, pp. 839– 853, Jun. 2013.
- [28] H. Ding, "Stability yields sublinear time algorithms for geometric optimization in machine learning," in 29th Annual European Symposium on Algorithms (ESA 2021), P. Mutzel, R. Pagh, and G. Herman (eds), vol. 204 of LIPIcs, Article No. 38, pp. 38:1–38:19, Leibniz-Zentrum für Informatik, Wadern, Germany: Dagstuhl Publishing, 2021.
- [29] M. Cavaleiro and F. Alizadeh, "A branch-and-bound method for the minimum k-enclosing ball problem," *Oper. Res. Lett.*, vol. 50, no. 3, pp. 274–280, 2022.
- [30] V. V. Shenmaier, "The problem of a minimal ball enclosing k points," J. Appl. Ind. Math., vol. 7, no. 3, pp. 444–448, 2013.

- [31] S. D. Ahipaşaoğlu and E. A. Yıldırım "Identification and elimination of interior points for the minimum enclosing ball problem," *SIAM J. Optim.*, vol. 19, no. 3, pp. 1392–1396, 2008.
- [32] K. Fischer and B. Gärtner, "The smallest enclosing ball of balls: Combinatorial structure and algorithms," *Int. J. Comput. Geom. Appl.*, vol. 14, no. 4-5, pp. 341–378, 2004.
- [33] S. G. Kolliopoulos and S. Rao, "A nearly linear-time approximation scheme for the Euclidean k-median problem," SIAM J. Comput., vol. 37, no. 3, pp. 757–782, 2007.
- [34] A. Bhattacharya, D. Goyal, and R. Jaiswal, "Hardness of approximation for Euclidean k-median," in Approximation, Randomization, and Combinatorial Optimization Algorithms and Techniques (AP-PROX/RANDOM 2021), Leibniz International Proceedings in Informatics (LIPIcs), vol. 207, pp. 4:1-4:23, Leibniz-Zentrum für Informatik, Wadern, Germany: Dagstuhl Publishing, 2021.
- [35] V. Verter, "Uncapacitated and capacitated facility location problems," in *Foundations of Location Analysis*, H. A. Eiselt and V. Marianov (eds.), International Series in Operations Research & Management Science, vol. 155, ch. 2, pp. 25-37, Springer, 2011.
- [36] S. G. Kolliopoulos and Y. Moysoglou, "Integrality gaps for strengthened linear relaxations of capacitated facility location," *Math. Program.*, vol. 158, no. 1-2, pp. 99–141, 2016.
- [37] S. Zhang, "Challenges in KNN classification," *IEEE Trans. Knowl. Data Eng.*, vol. 34, no. 10, pp. 4663–4675, Oct. 2022.
- [38] P. K. Syriopoulos, N. G. Kalampalikis, S. B. Kotsiantis, and M. N. Vrahatis, "kNN classification: a review," *Ann. Math. Artif. Intell.*, in press, 2023.
- [39] P. Indyk and A. Naor, "Nearest-neighbor-preserving embeddings," ACM Trans. Algorithms, vol. 3, no. 3, Art. 31, pp. 1–12, 2007.
- [40] I. Z. Emiris, V. Margonis, and I. Psarros, "Near-neighbor preserving dimension reduction via coverings for doubling subsets of l₁," *Theor. Comput. Sci.*, vol. 942, pp. 169–179, 2023.
- [41] N. T. An, N. M. Nam, and X. Qin, "Solving k-center problems involving sets based on optimization techniques," J. Glob. Optim., vol. 76, pp. 189–209, 2020.
- [42] A. Banik, B. B. Bhattacharya, and S. Das, "Minimum enclosing circle of a set of fixed points and a mobile point," *Comput. Geom. Theory Appl.*, vol. 47, no. 9, pp. 891–898, Oct. 2014.
- [43] S. Roy, A. Karmakar, S. Das, and S. C. Nandy, "Constrained minimum enclosing circle with center on a query line segment," *Comput. Geom. Theory Appl.*, vol. 42, no. 6-7, pp. 632–638, Aug. 2009.
- [44] M. Imanparast, S. N. Hashemi, and A. Mohades, "Efficient approximation algorithms for point-set diameter in higher dimensions," J. Algorithms Comput., vol. 51, no. 2 pp. 47–61, Dec. 2019.
- [45] T. M. Chan, "Approximating the diameter, width, smallest enclosing cylinder, and minimum-width annulus," *Int. J. Comput. Geom. Appl.*, vol. 12, no. 1-2, pp. 67–85, 2002.
- [46] C. A. Duncan, M. T. Goodrich, and E. A. Ramos, "Efficient approximation and optimization algorithms for computational metrology," in *Proc. Eighteth Annual Symposium on Discrete Algorithms (SODA 1997)*, pp. 121–130, New Orleans, LA: ACM/SIAM, Jan. 1997.
- [47] P. K. Agarwal, B. Aronov, S. Har-Peled, and M. Sharir, "Approximation algorithms for minimum-width annuli and shells," *Discrete Comput. Geom.*, vol. 24, no. 4, pp. 687–705, 2000.
- [48] B. Aronov and M. Dulieu, "How to cover a point set with a V-shape of minimum width," *Comput. Geom. Theory Appl.*, vol. 46, no. 3, pp. 298–309, 2013.
- [49] T. M. Chan and S. Har-Peled, "Smallest k-enclosing rectangle revisited," Discrete Comput. Geom., vol. 66, no. 2, pp. 769–791, 2021.
- [50] S. Z. Selim and M. A. Ismail, "K-means-type algorithms: A generalized convergence theorem and characterization of local optimality," *IEEE Trans. Pattern Anal. Mach. Intell.*, vol. PAMI-6, no. 1, pp. 81–87, Jan. 1984.
- [51] A. Vattani, "k-means requires exponentially many iterations even in the plane," Discrete Comput. Geom., vol. 45, no. 4, pp. 596–616, 2011.
- [52] M. N. Vrahatis, B. Boutsinas, P. Alevizos, and G. Pavlides, "The new k-windows algorithm for improving the k-means clustering algorithm," J. Complexity, vol. 18, no. 1, pp. 375–391, 2002.
- [53] S. W. Bae, "Computing a minimum-width square or rectangular annulus with outliers," *Comput. Geom. Theory Appl.*, vol. 76, no. 3, pp. 33–45, 2019.
- [54] S. I. Veselov, D. V. Gribanov, and D. S. Malyshev, "FPT-algorithm for computing the width of a simplex given by a convex hull," *Moscow Univ. Comput. Math. Cybern.*, vol. 43, no. 1, pp. 1–11, 2019.
- [55] L. G. Casado, I. García, B. G. Tóth, and E. M. T. Hendrix, "On determining the cover of a simplex by spheres centered at its vertices," *J. Glob. Optim.*, vol. 50, pp. 645–655, 2011.

- [56] H. Everett, I. Stojmenovic, P. Valtr, and S. Whitesides, "The largest k-ball in a d-dimensional box," *Comput. Geom. Theory Appl.*, vol. 11, no. 2, pp. 59–67, 1998.
- [57] A. Kanazawa, "On the minimal volume of simplices enclosing a convex body," Arch. Math. (Basel), vol. 102, no. 5, pp. 489–492, 2014.
- [58] D. Galicer, M. Merzbacher, and D. Pinasco, "The minimal volume of simplices containing a convex body," *J. Geom. Anal.*, vol. 29, no. 1, pp. 717–32, 2019.
- [59] G. Ya. Perel'man, "k-radii of a convex body," Sibirsk. Mat. Zh., vol. 28, no. 4, pp. 185–186, 1987.
- [60] S. V. Pukhov "Inequalities for the Kolmogorov and Bernštein widths in Hilbert space," *Mat. Zametki*, vol. 25, no. 4, pp. 619–628, 637, 1979.
- [61] B. González Merino, "Improving bounds for the Perel'man-Pukhov quotient for inner and outer radii," J. Convex Anal., vol. 24, no. 4, pp. 1099–1116, 2017.
- [62] F. Plastria, "Continuous covering location problems," in *Facility Location: Application and Theory*, Z. Drezner and H. W. Hamacher (eds), Chapter 2, pp. 37–79, New York, USA: Springer, 2002.
- [63] Y. Zhou and J. Zhu, "FocusPaka/MEB: A Minimum Enclosing Ball (MEB) method to detect differential expression genes for RNA-seq data," GitHub packages, License: GPL-2, Version: 0.99.4, built on October 22, 2022, [Website] https://rdrr.io/github/FocusPaka/MEB/
- [64] F. A. Elerian, W. M. K. Helal, and M. A. AbouEleaz, "Methods of roundness measurement: An experimental comparative study," *J. Mech. Eng. Res. Dev.*, vol. 44, no. 9, pp. 173–183, 2021.
- [65] C. J. C. Burges, "A tutorial on support vector machines for pattern recognition," *Data Min. Knowl. Discov.*, vol. 2, no. 2, pp. 121–167, 1998.
- [66] A. Ben-Hur, D. Horn, H. T. Siegelmann, and V. Vapnik, "Support vector clustering," J. Mach. Learn. Res., vol. 2, pp. 125–137, 2001.
- [67] Y. Bulatov, S. Jambawalikar, P. Kumar, and S. Sethia, "Hand recognition using geometric classifiers," in *International Conference on Biometric Authentication (ICBA 2004)*, D. Zhang and A. K. Jain (eds), *Lect. Notes Comput. Sci.*, vol. 3072, pp. 753–759, Berlin, Heidelberg: Springer, 2004.
- [68] N. Alon, S. Dar, M. Parnas, and D. Ron, "Testing of clustering," in Forty-first Annual Symposium on Foundations of Computer Science (FOCS 2000), pp. 240–250, Los Alamitos, CA: IEEE, 2000.
- [69] N. Alon, S. Dar, M. Parnas, and D. Ron, "Testing of clustering," SIAM J. Discrete Math., vol. 16, no. 3, pp. 393–417, 2003.
- [70] A. Rizwan, N. Iqbal, R. Ahmad, and D.-H. Kim, "WR-SVM model based on the margin radius approach for solving the minimum enclosing ball problem in support vector machine classification," *Appl. Sci.*, vol. 11, no. 10, 4657, pp. 1–21, Jun. 2021.
- [71] O. Chapelle, V. Vapnik, O. Bousquet, and S. Mukherjee, "Choosing multiple parameters for support vector machines," *Mach. Learn.*, vol. 46, no. 1/3, pp. 131–159, 2002.
- [72] Y. Ren and F. Zhang, "Hand gesture recognition based on MEB-SVM," in *Proc. International Conference on Embedded Software and Systems*, Hangzhou, China, 2009, pp. 344–349.
- [73] A. Goel, P. Indyk, and K. R. Varadarajan, "Reductions among high dimensional proximity problems," in *Proc. Twelfth Annual Symposium* on Discrete Algorithms (SODA 2001), S. Rao Kosaraju (ed), pp. 769– 778, Washington, DC, USA: ACM/SIAM, 2001.
- [74] P. M. Hubbard, "Approximating polyhedra with spheres for timecritical collision detection," ACM Trans. Graph., vol. 15, no. 3, pp. 179–210, 1996.
- [75] T. Larsson, G. Capannini, and L. Källberg, "Parallel computation of optimal enclosing balls by iterative orthant scan," *Comput. Graph.*, vol. 56, pp. 1–10, May 2016.
- [76] C. L. Lawson, "The smallest covering cone or sphere (Problem 63-5* by C. Groenewoud and L. Eusanio)," *SIAM Review*, vol. 7, no. 3, pp. 415–416, 1965.
- [77] M. Tzelepi and A. Tefas, "Improving the performance of light-weight CNN models using minimum enclosing ball regularization," in *Proc.* 27th European Signal Processing Conference (EUSIPCO 2019), pp. 1–5, A Coruña, Spain: IEEE 2019.
- [78] R. Kurniawati, J. S. Jin, and J. A. Shepherd, "SS⁺-tree: An improved index structure for similarity searches in a high-dimensional feature space," in *5th Storage and Retrieval for Image and Video Databases*, I. K. Sethi and R. C. Jain (eds), vol. 3022, pp. 110–120, San Jose, CA, USA: SPIE, 1997.
- [79] F.-L. Chung, Z. H. Deng, and S. T. Wang, "From minimum enclosing ball to fast fuzzy inference system training on large datasets," *IEEE Trans. Fuzzy Syst.*, vol. 17, no. 1, pp. 173–184, Feb. 2009.

- [80] M. Heusinger and F.-M. Schleif, "Classification in non-stationary environments using coresets over sliding windows," in Advances in Computational Intelligence (IWANN 2021), I. Rojas et al. (eds), Lect. Notes Comput. Sci., vol. 12861, pp. 126–137, Cham, Switzerland: Springer Nature, 2021.
- [81] P. J. Rousseeuw and A. M. Leroy, *Robust Regression and Outlier Detection*, New York, NY: John Wiley & Sons, Inc., 1987.
- [82] J. M. Abowd, "The US Census Bureau adopts differential privacy," in Proc. 24th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining (KDD 2018), 2018, pp. 2867–2867.
- [83] B. Ghazi, R. Kumar, and P. Manurangsi, "Differentially private clustering: Tight approximation ratios," in *Proc. Thirty-Fourth Conference on Neural Information Processing Systems (NeurIPS 2020)*, art. no. 340, pp. 4040–4054, Vancouver, Canada, December 2020.
- [84] B. Mahpud and O. Sheffet, "A differentially private linear-time fPTAS for the minimum enclosing ball problem," in *Proc. Thirty-Sixth Conference on Neural Information Processing Systems (NeurIPS 2022)*, A. H. Oh *et al.* (eds), paper 9140, pp. 1–13, New Orleans, Louisiana, USA, Nov. 2022.
- [85] I. W.-H. Tsang, J. T.-Y. Kwok, and P.-M. Cheung, "Core vector machines: Fast SVM training on very large data sets," *J. Mach. Learn. Res.*, vol. 6, no. 13, pp. 363–392, 2005.
- [86] I. W.-H. Tsang, J. T.-Y. Kwok, and J. M. Zurada, "Generalized core vector machines," *IEEE Trans. Neural Netw.*, vol. 17, no. 5, pp. 1126– 1140, Sep. 2006.
- [87] F. Nielsen and R. Nock, "Approximating smallest enclosing balls with applications to machine learning," *Int. J. Comput. Geom. Appl.*, vol. 19, no. 5, pp. 389–414, 2009.
- [88] C. Bauckhage, R. Sifa, and T. Dong, "Prototypes within minimum enclosing balls," in *Artificial Neural Networks and Machine Learning* (*ICANN 2019*), I. V. Tetko *et al.* (eds), *Lect. Notes Comput. Sci.*, vol. 11731, pp. 365–376, Cham, Switzerland: Springer Nature, 2019.
- [89] H. G. Eggleston, *Convexity*, Cambridge Tracts in Mathematics and Mathematical Physics, no. 47, P. Hall and F. Smithies (gen. eds), Cambridge, England: Cambridge University Press, 1958.
- [90] S. Har-Peled and E. Robson, "On the width of the regular n-simplex," 2023, arXiv:2301.02616.
- [91] R. M. Tanner, "Some content maximizing properties of the regular simplex," *Pacific J. Math.*, vol. 52, no. 2, pp. 611–616, 1974.
- [92] R. Alexander, "The width and diameter of a simplex," *Geometriae Dedicata*, vol. 6, no. 1, pp. 87–94, 1977.
- [93] P. Gritzmann and V. Klee, "Inner and outer *j*-radii of convex bodies in finite-dimensional normed spaces," *Discrete Comput. Geom.*, vol. 7, no. 3, pp. 255–280, 1992.
- [94] W. Blaschke, "Über den grössten Kreis in einer konvexen Punktmenge," Jahresber. Dtsch. Math.-Ver., vol. 23, pp. 369–374, 1914.
- [95] R. Brandenberg, "Radii of regular polytopes," *Discrete Comput. Geom.*, vol. 33, no. 1, pp. 43–55, 2005.
- [96] H. W. E. Jung, "Über den kleinsten Kreis, der eine ebene Figur einschliesst," J. Reine Angew. Math., vol. 137, no. 4, pp. 310–313, 1910.
- [97] E. Landau, "Über einige Ungleichheitsbeziehungen in der Theorie der analytischen Funktionen," Arch. Math. Phys., vol. 11, no. 3, pp. 31–36, 1905.
- [98] W. Süss, "Durchmesser und Umkugel bei mehrdimensionalen Punktmengen," *Math. Z.*, vol. 40, no. 1, pp. 315–316, 1936.
- [99] F. Bohnenblust, "Convex regions and projections in Minkowski spaces," Ann. Math., vol. 39, no. 2, pp. 301–308, 1938.
- [100] K. Leichtweiss, "Zwei Extremalprobleme der Minkowski Geometrie," Math. Zeitschr., vol. 62, pp. 37–49, 1955.
- [101] B. Grünbaum, 'On some covering and intersection properties in Minkowski spaces," *Pacific J. Math.*, vol. 9, no. 2, pp. 487–494, 1959.
- [102] J. Daneš, "On the radius of a set in a Hilbert space," Commentat. Math. Univ. Carolin., vol. 25, no. 2, pp. 355–362, 1984.
- [103] M. Katz, "Jung's theorem in complex projective geometry," Quart. J. Math. Oxford Ser. (2), vol. 36, no. 144, pp. 451–466, 1985.
- [104] B. V. Dekster, "An extension of Jung's theorem," *Israel J. Math.*, vol. 50, no. 3, pp. 169–180, Sep. 1985.
- [105] B. V. Dekster, "The Jung theorem for the spherical and the hyperbolic spaces," Acta Math. Hungar, vol. 67, no. 4, pp. 315–331, Dec. 1995.
- [106] B. V. Dekster, "The Jung theorem in metric spaces of curvature bounded above," *Proc. Amer. Math. Soc.*, vol. 125, no. 8, pp. 2425– 2433, Aug. 1997.
- [107] U. Lang and V. Schroeder, "Jung's theorem for Alexandrov spaces of curvature bounded above," Ann. Glob. Anal. Geom., vol. 15, no. 3, pp. 263–275, 1997.

- [108] V. Nguen-Khac and K. Nguen-Van, "An infinite-dimensional generalization of the Jung theorem," *Math. Notes*, vol. 80, no. 2, pp. 224–232, 2006.
- [109] V. Boltyanski and H. Martini, "Jung's theorem for a pair of Minkowski spaces," Adv. Geom., vol. 6, no. 4, pp. 645–650, Oct. 2006.
- [110] A. V. Akopyan, "Combinatorial generalizations of Jung's theorem," Discrete Comput. Geom., vol. 49, no. 3, pp. 478–484, Mar. 2013.
- [111] R. Brandenberg and B. González Merino, "The asymmetry of complete and constant width bodies in general normed spaces and the Jung constant," *Israel J. Math.*, vol. 218, no. 1, pp. 489–510, Mar. 2017.
- [112] H. Gericke, "Über die größte Kugel in einer konvexen Punktmenge," Math. Z., vol. 40, pp. 317–320, 1936.
- [113] L. A. Santaló, "Convex regions on the n-dimensional spherical surface," Ann. Math., vol. 47, no. 3, pp. 448–459, Jul. 1946.
- [114] L. A. Santaló, "Propiedades de las figuras convexas sobre la esfera," *Mathematicae Notae*, vol. 4, pp. 11–40, 1944.
- [115] R. M. Robinson, "Note on convex regions on the sphere," Bull. Amer. Math. Soc., vol. 44, pp. 115–116, Feb. 1938.
- [116] M. Henk, "A generalization of Jung's theorem," *Geom. Dedicata*, vol. 42, pp. 235–240, 1992.
- [117] U. Betke and M. Henk, "A generalization of Steinhagen's theorem," *Abh. Math. Semin. Univ. Hambg.*, vol. 63, no. 1, pp. 165–176, 1993.
- [118] D. Gale, "On inscribing n-dimensional sets in a regular n-simplex," Proc. Amer. Math. Soc., vol. 4, no. 2, pp. 222–225, 1953.
- [119] M. N. Vrahatis, "An error estimation for the method of bisection in \mathbb{R}^d ," *Bull. Greek Math. Soc.*, vol. 27, pp. 161–174, 1986.
- [120] M. N. Vrahatis, "A variant of Jung's theorem," Bull. Greek Math. Soc., vol. 29, pp. 1–6, 1988.
- [121] C. Carathéodory, "Über den Variabilitätsbereich der Koeffizienten von Potenzreihen, die gegebene Werte nicht annehmen," *Math. Ann.*, vol. 64, no. 1, pp. 95–115, 1907.
- [122] R. T. Rockafellar, *Convex Analysis*. Princeton Mathematical Series, no. 28, Princeton, NJ: Princeton University Press, 1970.
- [123] E. Helly, "Über Mengen konvexer Körper mit gemeinschaftlichen Punkten," Jahresber. Disch. Math.-Ver., vol. 32, pp. 175–176, 1923.
- [124] H. A. Tverberg, "A generalization of Radon's theorem," J. London Math. Soc., vol. 41, pp. 123–128, 1966.
- [125] J. A. De Loera, X. Goaoc, F. Meunier, and N. Mustafa, "The discrete yet ubiquitous theorems of Carathéodory, Helly, Sperner, Tucker, and Tverberg," *Bull. Amer. Math. Soc.* (*N.S.*), vol. 56, no. 3, pp. 415–511, 2019.
- [126] L. Danzer, B. Grünbaum, and V. Klee, "Helly's theorem and its relatives," in *Convexity, Proc. Symp. Pure Math.*, vol. 7, Amer. Math. Soc., Providence, RI, 1963, pp. 101–180.
- [127] H. A. Tverberg, "A generalization of Radon's theorem II," Bull. Austral. Math. Soc., vol. 24, no. 3, pp. 321–325, 1981.
- [128] J. Radon, "Mengen konvexer Körper, die einen gemeinsamen Punkt enthalten," *Math. Ann.*, vol. 83, no. 1-2, pp. 113–115, 1921.
- [129] B. J. Birch, "On 3N points in a plane," Proc. Cambridge Philos. Soc., vol. 55, pp. 289–293, 1959.
- [130] K. Adiprasito, I. Bárány, N. Mustafa, and T. Terpai, "Theorems of Carathéodory, Helly, and Tverberg without dimension," *Discrete Comput. Geom.*, vol. 64, no. 2, pp. 233–258, 2020.
- [131] I. Bárány, "A generalization of Carathéodory's theorem," Discrete Math., vol. 40, no. 2-3, pp. 141–152, 1982.
- [132] I. V. Polikanova and G. Ya. Perel'man, "A remark on Helly's theorem," *Sibirsk. Mat. Zh.*, vol. 27, no. 5, pp. 191–194, 207, 1986.



Michael N. Vrahatis is a professor emeritus of the University of Patras, Greece. His main scientific objectives are: (a) the investigation and mathematical foundation of methods in artificial intelligence, (b) the development of innovative methods in this field, (c) the application of the acquired expertise to address challenging real-life problems in different branches of science and the search of directions of theoretical mathematical results towards applications, and (d) the dissemination of knowledge to young researchers and scholars. His current interests

focus on the thematic areas: (a) mathematics, (b) natural computing and computational intelligence, (c) global optimization, (d) reliable computing and imprecise data, and (e) neural networks and machine learning & intelligence.