THE HARD LEFSCHETZ DUALITY FOR LOCALLY CONFORMALLY ALMOST KÄHLER MANIFOLDS

SHUHO KANDA

ABSTRACT. We prove the hard Lefschetz duality for locally conformally almost Kähler manifolds. This is a generalization of that for almost Kähler manifolds studied by Cirici and Wilson. We generalize the Kähler identities to prove the duality. Based on the result, we introduce the hard Lefschetz condition for locally conformally symplectic manifolds. As examples, we give solvmanifolds which do not satisfy the hard Lefschetz condition.

1. INTRODUCTION

1.1. Background and motivation. Kähler manifolds are a special class of symplectic manifolds. The first example of a symplectic manifold which does not admit a Kähler structure was constructed by William Thurston in 1976 [\[Thu76\]](#page-13-0). Since then, methods for determining whether a symplectic manifold admits a Kähler structure have been actively studied.

One of the most basic methods is to see the ring structure of the de Rham cohomology. We focus here on the hard Lefschetz condition (HLC, for short). Let (M, ω) be a 2n-dimensional symplectic manifold. We say that (M, ω) satisfies the HLC if the $(n - j)$ -th exterior power of the symplectic form ω^{n-j} induces an isomorphism between the cohomologies $H^j(M)$ and $H^{2n-j}(M)$ for all $1 \leq j \leq n$. It is known that all compact Kähler manifolds satisfy the HLC (see [\[Huy05\]](#page-12-0), [\[GH78\]](#page-12-1)). This means that a compact symplectic manifold (M, ω) which does not satisfy the HLC never admits a Kähler structure (M, J, g) such that the associated fundamental form $g(J \cdot, \cdot)$ is ω .

We can see how a compact symplectic manifold fails to satisfy the HLC by using an additional structure of it. Every compact symplectic manifold (M, ω) has at least one almost Kähler structure (M, J, g) such that the associated fundamental form is ω (we call the structure Kähler if J is integrable). Define the space of harmonic forms of bidegree (p, q) by letting

$$
\mathcal{H}^{p,q} \coloneqq \ker(\Delta_d) \cap \mathcal{A}^{p,q},
$$

where $\mathcal{A}^{p,q}$ denotes the space of (p,q) -forms. In [\[CW20\]](#page-12-2), Cirici and Wilson studied the properties of the space $\mathcal{H}^{p,q}$. According to the generalized hard Lefschetz duality, one of the results in [\[CW20\]](#page-12-2), it is shown that the maps

$$
(1.1) \tL^{n-(p+q)}: \mathcal{H}^{p,q} \xrightarrow{\sim} \mathcal{H}^{n-q,n-p}
$$

are isomorphisms for all p, q , where L denotes an operator defined by wedging a form with ω . In general, the inclusion

$$
(1.2)\qquad \bigoplus_{p+q=j} \mathcal{H}^{p,q} \subset \mathcal{H}^j
$$

is proper, where \mathcal{H}^j denotes a space of harmonic j-forms. If (M, J, g) is Kähler, i.e. J is integrable, it turns out that the inclusion (1.2) is equal by the Kähler identities. Hence all Kähler manifolds satisfy the HLC. In this sense, we can say that the gap of the inclusion [\(1.2\)](#page-0-0) measures to what extent is the HLC holds.

We consider a class of *locally conformally symplectic* (LCS) manifolds, which is a generalization of symplectic manifolds. Let M be a manifold and ω be a nondegenerate 2-form on M. We call the pair (M, ω) LCS if, for all points of M, there exists a neighborhood U of the point and real-valued function f on U such that $(U, e^{-f}\omega)$ is symplectic. This condition is equivalent to saying that there exists a closed 1-form θ on M, which is called the Lee form, such that $d\omega = \theta \wedge \omega$. As the similar generalization of a Kähler manifold, a *locally conformally Kähler* (LCK) manifold is defined. Typical examples of LCK manifolds are Hopf manifolds, Inoue surfaces, and Thurston's manifolds,

²⁰²⁰ Mathematics Subject Classification. 32Q60; 53C55; 53D35.

which are basic examples of complex manifolds which do not admit any Kähler structures (see [\[DO98\]](#page-12-3)). The class of LCS and LCK manifolds has attracted much interests among geometers in recent years.

As in the case of symplectic manifolds, it is natural to ask whether an LCS manifold admits an LCK structure. This existence problem is explicitly proposed by Ornea and Verbitsky in [\[OV11\]](#page-13-1). A few examples are discovered only in [\[BM17\]](#page-12-4) and [\[CY16\]](#page-12-5). One of the difficulties of the problem is that, in contrast to the case of Kähler structures, there are no known topological obstructions for the existence of LCK structures. See [\[OV23\]](#page-13-2) for details.

One of the clues to the problem is the HLC. In recent years, generalizations of the HLC to LCS manifolds have been studied. In [\[AOT18\]](#page-12-6), a generalization of the HLC for LCS is proposed by using the twisted de Rham cohomology (see also [\[LV15\]](#page-12-7), where the symplectic cohomology on LCS manifolds is studied by using the twisted differential).

There are two main purpose of this paper. First, to reformulate the definition of the HLC in [\[AOT18\]](#page-12-6). Second, as a generalization of the results in [\[CW20\]](#page-12-2), to develop the hard Lefschetz duality for locally conformally almost Kähler (LCaK) manifolds, which are non-integral version of LCK manifolds.

1.2. The results of this paper. We formulate the HLC for LCS manifolds as follows:

Definition 1.1 (The hard Lefschetz condition for locally conformally symplectic manifolds, Defi-nition [4.7\)](#page-8-0). Let (M, ω) be a 2n-dimensional compact locally conformally symplectic manifold with the Lee form θ . (M, ω) satisfies the *hard Lefschetz condition* if the maps

(1.3)
$$
[L]^{n-j}: H^j_{-(n-j)/2}(M) \to H^{2n-j}_{(n-j)/2}(M)
$$

are isomorphisms for all $1 \leq j \leq n$.

In the above definition, we write $H_k^j(M)$ for the cohomology group of a complex composed of differential forms and the *twisted differential* d_k defined by $d_k \alpha := d \alpha - k \theta \wedge \alpha$ for $k \in \mathbb{R}$. Using this cohomology, which is called the twisted de Rham cohomology, we can define the Lefschetz map $[L]^{n-j}$ as follows:

(1.4)
$$
[L]^{n-j}: H_k^j(M) \to H_{k+n-j}^{2n-j}(M), \quad [\alpha] \mapsto [\omega^{n-j} \wedge \alpha].
$$

According to the results in [\[Isl21\]](#page-12-8), it turns out that the space $H_i^j(M)$ is isomorphic to the space $H^{2n-j}_{-t}(M)$ for all $t \in \mathbb{R}$ by the Hodge theory. This is why we set the parameter k in [\(1.4\)](#page-1-0) to $-(n-j)/2$ so that the relation $k = -(k + n - j)$ holds, and adopt the above definition.

We write Λ for the adjoint operator of L. Denote by $\mathcal{H}_k^{p,q}$ the space of d_k -harmonic forms of bidegree (p, q) . The space $\mathcal{H}_{k, \mathcal{J}}^j$ is a space of d_k -harmonic forms α such that $\mathcal{J}\alpha$ is also d_k -harmonic, where $\mathcal J$ is an operator induced by J .

Using the above spaces and the formulation of the HLC, we generalize the hard Lefschetz duality (1.1) as follows:

Theorem 1.2 (The hard Lefschetz duality for locally conformally almost Kähler manifolds, The-orem [4.4\)](#page-7-0). Let (M, J, g) be a 2n-dimensional compact locally conformally almost Kähler manifold with the Lee form θ . Then the operators $\{L, \Lambda, H \coloneqq [L, \Lambda]\}\$ define finite dimensional representations of $\mathfrak{sl}(2)$ on

(1.5)
$$
\bigoplus_{p,q\geq 0} \mathcal{H}^{p,q}_{-(n-(p+q))/2} \quad and \quad \bigoplus_{j\geq 0} \mathcal{H}^{j}_{-(n-j)/2,\mathcal{J}}.
$$

Furthermore, for all $0 \leq p, q$ such that $j = p + q \leq n$,

$$
L^{n-j} \colon \mathcal{H}^{p,q}_{-(n-j)/2} \, \xrightarrow{\sim} \, \mathcal{H}^{n-q,n-p}_{(n-j)/2}
$$

and

$$
L^{n-j} \colon \mathcal{H}^{j}_{-(n-j)/2,\mathcal{J}} \xrightarrow{\sim} \mathcal{H}^{2n-j}_{(n-j)/2,\mathcal{J}}
$$

are isomorphisms.

To prove this theorem, we establish the Kähler identities for LCaK manifolds (Proposition [3.2\)](#page-5-0). Furthermore, we can see the importance of the relation $k = -(k + n - j)$ in the proof.

Using this theorem, we can see how the ranks of the Lefschetz maps in [\(1.3\)](#page-1-1) drop. As examples, we give a family of LCaK manifolds which do not satisfy the HLC in Section [5.](#page-9-0) For each integer $m \geq 2$, we construct a $(2m + 2)$ -dimensional compact solvable Lie group M_m and a left-invariant LCaK structure of it. The manifolds are the same as those constructed in Section 4 of [\[AO18\]](#page-12-9). On manifolds M_2 , M_3 , and M_4 , we give the basis of the spaces in [\(1.3\)](#page-1-1) and [\(1.5\)](#page-1-2) and compute the Lefschetz maps.

Finally, we propose a conjecture on the HLC for LCK manifolds. Recall that the HLC for symplectic manifolds is always satisfied on Kähler manifolds. According to this fact, the following conjecture is naturally suggested:

Conjecture 1.3 (Conjecture [4.11\)](#page-8-1). Let (M, J, g) be a 2n-dimensional compact LCK manifold with the Lee form θ . Then, (M, J, g) satisfies the HLC.

The above conjecture is still open.

1.3. Organization of this paper. This paper is organized as follows:

- In Section [2](#page-2-0) we collect preliminaries on LCaK structures, the twisted cohomology, operators on almost complex manifolds, and the Lefschetz decomposition.
- In Section [3](#page-5-1) we write down Kähler identities for LCaK manifolds and prove them (Proposition [3.2\)](#page-5-0).
- In Section [4](#page-7-1) we define some subspaces of twisted harmonic forms. Then we prove the main theorem (Theorem [4.4\)](#page-7-0) and define the HLC for LCS manifolds (Definition [4.7\)](#page-8-0).
- In Section [5](#page-9-0) we give examples of LCaK manifolds which do not satisfy the HLC and compute the spaces defined in the previous sections with their basises.

Acknowledgements

I am deeply grateful to my supervisor Prof. Kengo Hirachi for his enormous support and helpful advice. I am also grateful to Prof. Hisashi Kasuya for helpful discussions. Lastly, I also would like to thank Jin Miyazama for several helpful comments. This research was supported by Forefront Physics and Mathematics Program to Drive Transformation (FoPM), a World-leading Innovative Graduate Study (WINGS) Program, the University of Tokyo.

2. Preliminaries

In this section we review the basics of locally conformally almost Kähler geometry, the twisted cohomology, and the Lefschetz decomposition. Throughout this paper, we only consider smooth connected manifolds without boundary which have even real dimension $2n \geq 4$. We always assume that functions and differential forms are smooth. Furthermore, the vector spaces defined in this paper, such as the space of harmonic forms, the de Rham cohomology group, etc, are always assumed to be complex vector spaces, not real vector spaces.

2.1. Locally conformally almost Kähler manifolds. Let M be a smooth manifold of dimension $2n \geq 4$ and ω be a nondegenerate 2-form on M, i.e. $\omega^{n} \neq 0$ at all points of M. The form ω is said to be *conformally symplectic* (CS, for short) on M if there exists a real-valued function f such that $e^{-f}\omega$ is symplectic on M, i.e. $d(e^{-f}\omega) = 0$. The form ω is said to be locally conformally symplectic (LCS) on M if there exists an open covering ${U_i}_{i \in I}$ of M such that $\omega|_{U_i}$ are CS on U_i for all $i \in I$. Let f_i be a real-valued function on U_i such that $e^{-f_i}\omega$ is symplectic on U_i . Then the condition that the form $e^{-f_i}\omega$ is symplectic is equivalent to $d\omega = df_i \wedge \omega$. Since ω is nondegenerate, $df_i \wedge \omega = df_j \wedge \omega$ implies $df_i = df_j$ on $U_i \cap U_j$. So the 1-forms $\{df_i\}_{i \in I}$ are glued together. Thus we obtain the following proposition which makes the above definition simpler:

Proposition 2.1. A nondegenerate 2-form ω on M is LCS if and only if there exists a closed 1-form θ on M such that $d\omega = \theta \wedge \omega$. Furthermore, ω is CS if and only if θ is exact.

The 1-form θ is determined by ω , and we call it the Lee form of ω . In these situations we call the couple (M, ω) an LCS manifold (CS manifold) with the Lee form θ .

If M admits an almost complex structure J and a hermitian metric g such that the associated fundamental form $\omega(\cdot, \cdot) := q(J \cdot, \cdot)$ is LCS (CS), we call the triple (M, J, q) a locally conformally (conformally) almost Kähler (LCaK (CaK)). Furthermore, we call the triple (M, J, g) a locally conformally (conformally) Kähler (LCK (CK)) if the almost complex structure J is integrable. The relation among the classes of manifolds is illustrated in Figure [1.](#page-3-0)

Lastly, we note that for every nondegenerate 2-forms ω on M there exists an almost complex structure J and a hermitian metric g such that the fundamental form of (M, J, g) coincides with ω.

2.2. The twisted de Rham cohomology. The twisted de Rham cohomology is also called the Morse-Novikov cohomology, the Lichnerowicz cohomology, or the adapted cohomology. Its study was initiated by Novikov [\[Nov82\]](#page-12-10), and developed by Guedira and Lichnerowicz [\[GL84\]](#page-12-11). For an introduction to the twisted de Rham cohomology, see [\[OV09\]](#page-12-12). Here we give the definitions and some basic properties of them.

Let M be a 2n-dimensional manifold. Denote by \mathcal{A}^j the space of complex-valued differential j-forms on M, and $H^j(M)$ the j-th cohomology group of the complex (A^{\bullet}, d) , where d is the exterior derivative. Let θ be a closed 1-form. We define the twisted differential d_{θ} as

$$
d_{\theta} \coloneqq d - \epsilon_{\theta} \colon \mathcal{A}^{\bullet} \to \mathcal{A}^{\bullet+1},
$$

where ϵ_{α} denotes an operator defined by $\cdot \mapsto \alpha \wedge \cdot$ for a form α . Similar to the usual differential, we have $d_{\theta}d_{\theta} = 0$. Denote by $H_{\theta}^{j}(M)$ the j-th cohomology group of the complex $(\mathcal{A}^{\bullet}, d_{\theta})$, and we call it the twisted de Rham cohomology or simply the twisted cohomology. The cohomology group $H^j_{\theta}(M)$ depends only on the cohomology class of $[\theta]$ in $H^1(M)$. Indeed, if $\theta' = \theta + df$, where f is a function on M , there exists an isomorphism of complexes:

$$
e^f: (\mathcal{A}^\bullet, d_\theta) \stackrel{\sim}{\longrightarrow} (\mathcal{A}^\bullet, d_{\theta'}).
$$

In particular, the twisted cohomology $H^j_{\theta}(M)$ is isomorphic to the de Rham cohomology $H^j(M)$ if θ is exact.

Let $k \in \mathbb{R}$ be a real number. We write $d_{k\theta}$ and $H_{k\theta}^{j}(M)$ as d_{k} and $H_{k}^{j}(M)$ respectively when θ is clear. Using these notation, we have the following Leibniz rule:

(2.1)
$$
d_{k+l}(\alpha \wedge \beta) = d_k \alpha \wedge \beta + (-1)^{\deg \alpha} \alpha \wedge d_l \beta
$$

for all $k, l \in \mathbb{R}$. From this formula, we have the following map induced by the wedge product:

(2.2) ∧ : H j1 k1 (M) × H j2 k2 (M) → H j1+j2 k1+k2 (M).

We consider the case where M is compact, orientable, and endowed with a Riemannian metric g. Denote by d_k^* the adjoint operator of d_k with respect to the metric g, and we define the *twisted* laplacian $\Delta_k := d_k^* d_k + d_k d_k^*$. Since the twisted laplacian Δ_k is elliptic, the space of differential forms A^{\bullet} has the following Hodge decomposition:

$$
\mathcal{A}^{\bullet}=\mathcal{H}^{\bullet}_k\oplus d_k\mathcal{A}^{\bullet-1}\oplus d_k^*\mathcal{A}^{\bullet+1},
$$

where $\mathcal{H}^{\bullet}_{k}$ denotes the kernel of Δ_{k} , which is called the space of *twisted harmonic forms*. As a consequence, the natural map $\mathcal{H}^{\bullet}_{k} \to H^{\bullet}_{k}(M)$ defined by $\alpha \mapsto [\alpha]$ is an isomorphism. We also have $\dim H_k^{\bullet}(M) < \infty$. The Hodge duality for the twisted cohomologies is stated as follows:

Proposition 2.2 ([\[Isl21\]](#page-12-8)). In the above situation, we have

$$
\Delta_k=\Delta_{-k}
$$

for all $k \in \mathbb{R}$. In particular, the maps

$$
* \colon \mathcal{H}^{j}_{k} \, \stackrel{\sim}{\longrightarrow} \, \mathcal{H}^{2n-j}_{-k}
$$

are defined and isomorphisms for all $0 \leq j \leq 2n$.

2.3. Operators associated with an almost complex structure. Let (M, J) be an almost complex manifold, and

$$
\mathcal{A}^j=\bigoplus_{p+q=j}\mathcal{A}^{p,q}
$$

be the bigraded decomposition of the space of j-forms A^j with respect to J. We fix a closed 1-form θ on M and $k \in \mathbb{R}$. The twisted differential d_k splits into four components:

$$
d_k = \partial_k + \partial_k + \mu + \overline{\mu},
$$

where the bidegrees of the operators ∂_k , ∂_k , μ and $\overline{\mu}$ are $(1,0)$, $(0,1)$, $(2,-1)$, and $(-1,2)$ respectively. Then we have $\partial_k = \partial - k\epsilon_{\theta^{1,0}}$ and $\overline{\partial}_k = \overline{\partial} - k\epsilon_{\theta^{0,1}}$, where $\theta^{1,0}$ and $\theta^{0,1}$ denote the $(1,0)$ -part and (0, 1)-part of θ respectively. Since the operator μ acts as zero on 0-forms, μ and $\overline{\mu}$ are linear over functions. It is well known that J is integrable if and only if $\mu = 0$. Splitting the equation $d_k d_k = 0$ into bidegrees, we obtain the following equations:

$$
\partial_k \partial_k = -\mu \partial_k - \partial_k \mu,
$$

\n
$$
\partial_k \overline{\partial}_k + \overline{\partial}_k \partial_k = -\mu \overline{\mu} - \overline{\mu} \mu,
$$

\n
$$
\mu \partial_k = -\partial_k \mu,
$$

\n
$$
\mu \mu = 0.
$$

The almost complex structure J induces the real isomorphism

$$
\mathcal{J} \coloneqq \sum_{p+q=j} \mathbf{i}^{p-q} \, \Pi^{p,q} \colon \mathcal{A}^j \to \mathcal{A}^j,
$$

where $\Pi^{p,q}$ denotes the projection $A^j \to A^{p,q}$, and i denotes the imaginary unit. Using this operator, we define a real operator d_k^c as

$$
d_k^c \coloneqq \mathcal{J}^{-1} d_k \mathcal{J}.
$$

By straightforward calculations, we have

$$
d_k^c = -\mathrm{i}(\partial_k - \overline{\partial}_k) + \mathrm{i}(\mu - \overline{\mu}).
$$

The anti-Lee form θ^c is defined by

$$
\theta^c\coloneqq -\mathcal{J}\theta = -\theta\circ J = -\mathrm{i}(\theta^{1,0}-\theta^{0,1}).
$$

Using this, we can write

$$
\mathcal{J}^{-1} \epsilon_\theta \mathcal{J} = \epsilon_{\theta^c}.
$$

So we have

$$
d_k^c = d^c - k\epsilon_{\theta^c}.
$$

2.4. The Lefschetz decomposition on the space of forms. We review some basic facts about the Lefschetz decomposition on the space of forms. See e.g. [\[Huy05\]](#page-12-0), [\[GH78\]](#page-12-1) for details.

We consider an almost hermitian manifold (M, J, g) with the associated fundamental form ω . The Lefschetz map is an operator defined by wedging a form with ω :

$$
L\coloneqq \epsilon_\omega\colon \mathcal{A}^{p,q}\to \mathcal{A}^{p+1,q+1},
$$

and its formal adjoint operator Λ is

$$
\Lambda\coloneqq (-1)^{p+q}\ast L\ast\colon \mathcal{A}^{p,q}\rightarrow \mathcal{A}^{p-1,q-1}.
$$

It is well known that the operators $\{L, \Lambda, H := [L, \Lambda]\}\$ define a representation of $\mathfrak{sl}(2)$ on

$$
\mathcal{A}^{\bullet} = \bigoplus_{j} \mathcal{A}^{j}.
$$

The $\mathfrak{sl}(2)$ -representation induces the *Lefschetz decomposition*

$$
\mathcal{A}^{p,q} = \bigoplus_{j\geq 0} L^j \mathcal{P}^{p-j,q-j},
$$

where $\mathcal{P}^{p-j,q-j}$ denotes the space ker(Λ) $\cap \mathcal{A}^{p-j,q-j}$. From the decomposition, the maps

$$
L^j\colon \mathcal{A}^{p,q} \to \mathcal{A}^{p+j,q+j}
$$

turn out to be injective for all $0 \leq j \leq n - (p + q)$. In this paper, we call the composite L^{j} also the Lefschetz map.

 $3.$ The KÄHLER IDENTITIES FOR LOCALLY CONFORMALLY ALMOST KÄHLER MANIFOLDS

Let (M, J, g) be an almost hermitian manifold and ω be the associated fundamental form. In the case where (M, J, g) is almost Kähler, i.e. ω is closed, there are remarkable relations among operators which are the so-called Kähler identities (the term "Kähler identities" is often used for the identities in the case where J is integrable, however, here we cite Weil's classical result [\[Wei58\]](#page-13-3) for the non-integrable cases. See also [\[CW20\]](#page-12-2). The references for the integrable cases, see e.g. [\[Huy05\]](#page-12-0), [\[GH78\]](#page-12-1)).

In what follows, we write δ^* for the formal adjoint operator of an operator δ . It is well known that $d^* = - * d^*$. Splitting the formula into bidegree, we have

$$
\overline{\mu}^* = - * \mu * \text{ and } \overline{\partial}^* = - * \partial *.
$$

Proposition 3.1 (The Kähler identities for almost Kähler manifolds, [\[Wei58\]](#page-13-3)). Let (M, J, g) be an almost Kähler manifold with the associated fundamental form ω . Then the following identities hold:

We generalize these identities to LCaK manifolds.

Proposition 3.2 (The Kähler identities for LCaK manifolds). Let (M, J, q) be a 2n-dimensional $LCaK$ manifold with the associated fundamental form ω and the Lee form θ . Then the following *identities hold as operators acting on* A^j :

(i)
$$
d_{k+1}L - Ld_k = 0
$$
,
\n $\partial_{k+1}L - L\partial_k = 0$,
\n $\overline{\partial}_{k+1}L - L\overline{\partial}_k = 0$,
\n(ii) $d_{k+1}^*L - Ld_k^* = -d_{-n-k+j}^c$,
\n(iii) $d_{k+1}^*L - Ld_k^* = -d_{-n-k+j}^c$,
\n(iii) $\partial_{k+1}^*L - Ld_k^* = -d_{-n-k+j}^c$,
\n(ii) $\partial_{k+1}^*L - L\overline{\partial}_k = -\overline{\partial}_{-n-k+j}$,
\n(iii) $[\mu, L] = [\overline{\mu}, L] = 0$,
\n(iii) $\overline{\partial}_k \Lambda - \Lambda \overline{\partial}_{k+1} = 0$,
\n(ii) $d_{k+1}^*L - L\overline{\partial}_k^* = -\overline{\partial}_{-n-k+j}$,
\n(iii) $[\mu, L] = [\overline{\mu}, L] = 0$,
\n(iiii) $[\mu, L] = [\overline{\mu}, L] = 0$,
\n(iv) $[\mu^*, L] = i\overline{\mu}$,
\n(iv) $[\mu^*, L] = -i\mu$,
\n(v) $[\overline{\mu^*}, L] = -i\mu$,
\n(v) $[\overline{\mu^*}, L] = -i\mu$,
\n(v) $[\overline{\mu^*}, L] = -i\mu$,

The Kähler identities for LCK manifolds, which are the special case of the above proposition, are obtained in [\[AU16\]](#page-12-13). Here we give a proof using conformal changes of the operators. Before the proof, we give some definitions and basic facts. Let f be a real-valued function. We consider the conformal change $\tilde{g} := e^{-f}g$ and the fundamental form $\tilde{\omega} := e^{-f}\omega$. Denote by * the Hodge star operator with respect to the metric \tilde{g} . It is straightforward to check that the equality

$$
(3.1) \qquad \qquad \tilde{\ast} = e^{-(n-j)f} \ast
$$

holds as an operator acting on \mathcal{A}^j . We also define the operator $d^* := -\tilde{*}d\tilde{*}$ which is the formal adjoint of d with respect to \tilde{g} . For a 1-form α we define the operator $\epsilon_{\alpha}^* := * \epsilon_{\alpha} *$ which is the formal adjoint operator of ϵ_{α} with respect to g. Taking the adjoints of the equality

$$
[d, e^{-f}] = -e^{-f} \epsilon_{df},
$$

we have

$$
[d^*, e^{-f}] = e^{-f} \epsilon_{df}^*.
$$

The operator ϵ_θ^* can be written as

$$
\epsilon_\theta^* = \iota_{\theta^\sharp},
$$

where \sharp denotes the musical isomorphism, i.e. θ^{\sharp} is the vector field determined by the relation $g(\theta^{\sharp}, \cdot) = \theta(\cdot)$ (see [\[Lee18\]](#page-12-14)). Using this, we have

$$
[\,L, \epsilon_\theta^* \,]=[\,L, \iota_{\theta^\sharp}\,]=-\epsilon_{(\iota_{\theta^\sharp}\omega)}.
$$

Combining this with the equation

$$
\iota_{\theta^{\sharp}}\omega=\omega(\theta^{\sharp},\cdot)=-g(\theta^{\sharp},J\cdot)=-\theta\circ J=\theta^c,
$$

we have the relation

(3.3) $[L, \epsilon^*_{\theta}] = -\epsilon_{\theta^c}$.

Proof of Proposition [3.2.](#page-5-0) The identities $(i)^*, (ii)^*, (iii)^*$ and $(iv)^*$ are obtained immediately by taking adjoints of the identities (i),(ii),(iii) and (iv) respectively. The first identity in (i) follows from $d_1\omega = 0$ and the Leibniz rule [\(2.1\)](#page-3-1). By splitting this identity into bidegrees, we obtain (i) and (iii). Similarly, we obtain (ii) and (iv) by splitting the first identity in (ii). Therefore, it is sufficient to prove the first identity in (ii).

Since (M, J, g) is an LCaK manifold, we can locally take a real-valued function f such that the metric $\tilde{g} = e^{-f}g$ endows (M, J) with the Kähler structure. The associated fundamental form of (M, J, \tilde{g}) is $\tilde{\omega} = e^{-f}\omega$, and we have $\theta = df$ by the uniqueness of the Lee form. From Proposition [3.1,](#page-5-2) we obtain

$$
[d^*, \tilde{L}] = -d^c,
$$

where $\tilde{L} := \epsilon_{\tilde{\omega}}$. From the equation [\(3.1\)](#page-5-3), d^* can be calculated as follows:

$$
d^* = -\tilde{*}d\tilde{*}
$$

= $-\tilde{*}d(e^{-(n-j)f})$
= $-\tilde{*}e^{-(n-j)f}(-(n-j)\epsilon_{\theta} * +d*)$
= $-e^{(n-(2n-j+1))} * e^{-(n-j)f}(-(n-j)\epsilon_{\theta} * +d*)$
= $e^f(d^* + (n-j)\epsilon_{\theta}^*).$

Using this and the equation [\(3.2\)](#page-6-0), we obtain

 \ddot{x}

$$
d^{\tilde{*}}\tilde{L} = e^f(d^*(e^{-f}L) + (n - (j + 2))\epsilon_{\theta}^*e^{-f}L)
$$

= $e^f[d^*, e^{-f}]L + d^*L + (n - j - 2)\epsilon_{\theta}^*L$
= $d^*L + (n - j - 1)\epsilon_{\theta}^*L$

and

$$
\tilde{L}d^* = Ld^* + (n-j)L\epsilon_{\theta}^*.
$$

Hence we have

$$
-d^{c} = [d^{\tilde{*}}, \tilde{L}]
$$

= [d^*, L] + (n - j - 1)\epsilon_{\theta}^*L - (n - j)L\epsilon_{\theta}^*

Therefore, we obtain

$$
d_{k+1}^* L - L d_k^* = (d^* - (k+1)\epsilon_{\theta}^*) L - L(d^* - k\epsilon_{\theta}^*)
$$

=
$$
[d^*, L] - (k+1)\epsilon_{\theta}^* L + kL\epsilon_{\theta}^*
$$

=
$$
(-d^c - (n-j-1)\epsilon_{\theta}^* L + (n-j)L\epsilon_{\theta}^*) - (k+1)\epsilon_{\theta}^* L + kL\epsilon_{\theta}^*
$$

=
$$
-(d^c + (-n-k+j)[L, \epsilon_{\theta}^*)].
$$

From the equations [\(2.3\)](#page-4-0) and [\(3.3\)](#page-6-1), we have

$$
d^{c} + (-n - k + j)[L, \epsilon_{\theta}^{*}] = d^{c} - (-n - k + j)\epsilon_{\theta^{c}}
$$

$$
= d^{c}_{-n-k+j}.
$$

This proves the proposition.

4. Twisted harmonic forms and the Lefschetz condition for LCS manifolds

4.1. The generalized Lefschetz duality for LCaK manifolds. Let (M, J, g) be a compact LCaK manifold with the Lee form θ . We define the following subspaces of harmonic forms:

$$
\mathcal{H}_k^{p,q} \coloneqq \mathcal{H}_k^{p+q} \cap \mathcal{A}^{p,q} \quad \text{and} \quad \mathcal{H}_{k,\mathcal{J}}^j \coloneqq \{ \alpha \in \mathcal{H}_k^j \mid \mathcal{J} \alpha \in \mathcal{H}_k^j \}.
$$

We have the following inclusion:

$$
\bigoplus_{p+q=j}\mathcal{H}_k^{p,q}\subset \mathcal{H}_{k,\mathcal{J}}^j
$$

.

These spaces have the following duality relations:

Proposition 4.1. Let (M, J, g) be a 2n-dimensional compact LCaK manifold with the Lee form θ. For all $0 \le j, p, q$, the following duality relations hold:

(i) (The complex conjugation). Taking the complex conjugation induces isomorphisms

$$
\mathcal{H}_k^{p,q} \stackrel{\sim}{\longrightarrow} \mathcal{H}_k^{q,p} \quad \text{and} \quad \mathcal{H}_{k,\mathcal{J}}^j \stackrel{\sim}{\longrightarrow} \mathcal{H}_{k,\mathcal{J}}^j.
$$

(ii) (The Hodge duality). The Hodge star operator * induces isomorphisms

$$
\mathcal{H}^{p,q}_k\ \stackrel{\sim}{\longrightarrow}\ \mathcal{H}^{n-q,n-p}_{-k} \quad and \quad \mathcal{H}^{j}_{k,\mathcal{J}}\ \stackrel{\sim}{\longrightarrow}\ \mathcal{H}^{2n-j}_{-k,\mathcal{J}}.
$$

Proof. The duality relations (i) hold since Δ_k and J are real operators. The duality relations (ii) are obtained by Proposition [2.2](#page-3-2) and the commutativity between the operators $*$ and \mathcal{J} .

We next prove the generalized Lefschetz duality for LCaK manifolds. Firstly, we prove the following

Proposition 4.2. Let (M, J, g) be a 2n-dimensional LCaK manifold with the Lee form θ . If $n+2k-j=0$, we have $L\alpha \in \mathcal{H}^{j+2}_{k+1,\mathcal{J}}$ and $\Lambda \alpha \in \mathcal{H}^{j-2}_{k-1,\mathcal{J}}$ for all $\alpha \in \mathcal{H}^{j}_{k,\mathcal{J}}$.

Proof. By Proposition [3.2,](#page-5-0) we obtain

$$
d_{k+1}L\alpha = Ld_k\alpha = 0
$$

and

$$
d_{k+1}^* L \alpha = -d_{-n-k+j}^c \alpha + L d_k^* \alpha
$$

= $-\mathcal{J}^{-1} d_{k-(n+2k-j)} \mathcal{J} \alpha + 0$
= $-\mathcal{J}^{-1} d_k(\mathcal{J} \alpha)$
= 0.

Therefore we have $L\alpha \in \mathcal{H}^{j+2}_{k+1,\mathcal{J}}$. By taking adjoints of all operators, we have $\Lambda\alpha \in \mathcal{H}^{j-2}_{k-1,\mathcal{J}}$ \Box

Since $\mathcal{H}_k^{p,q} \subset \mathcal{H}_{k,\mathcal{J}}^{p+q}$, we have the following

Corollary 4.3. If $n + 2k - (p + q) = 0$, we have $L\alpha \in \mathcal{H}_{k+1}^{p+1,q+1}$ and $\Lambda\alpha \in \mathcal{H}_{k-1}^{p-1,q-1}$ for all $\alpha \in \mathcal{H}_k^{p,q}$.

Combining these arguments with the Lefschetz decomposition on the space of forms, we obtain the following

Theorem 4.4 (The hard Lefschetz duality for LCaK manifolds). Let (M, J, q) be a 2n-dimensional compact LCaK manifold with the Lee form θ . Then the operators $\{L, \Lambda, H \coloneqq [L, \Lambda]\}\$ define finite dimensional representations of sl(2) on

(4.1)
$$
\bigoplus_{p,q\geq 0} \mathcal{H}^{p,q}_{-(n-(p+q))/2} \quad and \quad \bigoplus_{j\geq 0} \mathcal{H}^{j}_{-(n-j)/2,\mathcal{J}}.
$$

Furthermore, for all $0 \leq p, q$ such that $j = p + q \leq n$,

$$
L^{n-j} \colon \mathcal{H}^{p,q}_{-(n-j)/2} \, \xrightarrow{\sim} \, \mathcal{H}^{n-q,n-p}_{(n-j)/2}
$$

and

$$
L^{n-j} \colon \mathcal{H}^{j}_{-(n-j)/2,\mathcal{J}} \xrightarrow{\sim} \mathcal{H}^{2n-j}_{(n-j)/2,\mathcal{J}}
$$

are isomorphisms.

Proof. As explained in Section [2.4,](#page-4-1) $\{L, \Lambda, H\}$ defines a representation of $\mathfrak{sl}(2)$ on forms. Since $n + 2k - j = 0$, where $k = -(n - j)/2$, $\{L, \Lambda, H\}$ defines a subrepresentation on the spaces [\(4.1\)](#page-7-2) by Proposition [4.2.](#page-7-3) This proves the first statement of the theorem.

We shall prove the second statement. From the Lefschetz decomposition, the operator L^{n-j} is injective as an operator acting on \mathcal{A}^{p+q} . Moreover, by Proposition [4.1](#page-7-4) we have dim $\mathcal{H}^{p,q}_{-(n-j)/2}$ = $\dim \mathcal{H}^{n-q,n-p}_{(n-j)/2} < \infty$ and $\dim \mathcal{H}^{j}_{-(n-j)/2,\mathcal{J}} = \dim \mathcal{H}^{2n-j}_{(n-j)/2,\mathcal{J}} < \infty$. So the maps are isomorphisms.

Remark 4.5. The space of harmonic forms of bidegree (p, q) , denoted by $\mathcal{H}^{p,q}$, is defined on almost Kähler manifolds in [\[CW20\]](#page-12-2). Thus the space $\mathcal{H}_k^{p,q}$ can be regarded as a generalization of the space $\mathcal{H}^{p,q}$ to LCaK manifolds. On almost Kähler manifolds, the relations

$$
\mathcal{H}^{p,q}=\mathcal{H}^{p,q}_{\overline{\partial}}\cap\mathcal{H}^{p,q}_\mu=\mathcal{H}^{p,q}_\partial\cap\mathcal{H}^{p,q}_{\overline{\mu}},
$$

where $\mathcal{H}^{p,q}_{\delta}$ denotes a space $\mathcal{A}^{p,q} \cap \text{ker } \delta \cap \text{ker } \delta^*$ for an operator δ , hold [\[CW20\]](#page-12-2). However, on LCaK manifolds, the similar relations do not hold in general.

4.2. Relations with the twisted cohomology and the hard Lefschetz condition for LCS **manifolds.** In this section, we observe the relations of harmonic forms $\mathcal{H}_k^{p,q}$ and $\mathcal{H}_{k,\mathcal{J}}^j$ with the twisted cohomology of LCaK manifolds. We define the Lefschetz map on the twisted cohomology as follows:

$$
[L]: H_k^j(M) \to H_{k+1}^{j+2}(M), \quad [\alpha] \mapsto [\omega \wedge \alpha].
$$

The map is well-defined since the Leibniz rule [\(2.1\)](#page-3-1) holds. We call the maps $[L]$ and $[L]$ ^j also the Lefschetz maps. The maps $[L]$ and L are compatible with the following natural inclusions:

$$
\bigoplus_{p+q=j} \mathcal{H}_k^{p,q} \subset \mathcal{H}_{k,\mathcal{J}}^j \hookrightarrow H_k^j(M),
$$

Remark 4.6. On compact Kähler manifolds, the above three spaces are isomorphic for each j. In general, however, these inclusions are proper even on compact CaK manifolds [\[CW20\]](#page-12-2). We give examples of compact LCaK manifolds with the same property in Section [5.](#page-9-0)

Considering the above, we define the following condition on compact LCS manifolds:

Definition 4.7 (The hard Lefschetz condition for LCS manifolds). Let (M, ω) be a 2n-dimensional compact LCS manifold with the Lee form θ . (M, ω) satisfies the hard Lefschetz condition (HLC) if the maps

$$
[L]^{n-j} \colon H^j_{-(n-j)/2}(M) \to H^{2n-j}_{(n-j)/2}(M)
$$

are isomorphisms for all $1 \leq j \leq n$.

Remark 4.8. If (M, ω) is a CS manifold, we can take a global real-valued function f on M such that $e^{-f}\omega$ is symplectic. Then the above condition is equivalent to saying that a symplectic manifold $(M, e^{-f}\omega)$ satisfies the HLC in the classical sense. In this sense, the above condition is a generalization of the HLC for symplectic manifolds to that for LCS manifolds.

Remark 4.9. The HLC for LCS manifolds is suggested also in [\[AOT18\]](#page-12-6). However, an LCS manifold (M, ω) satisfies the HLC defined in [\[AOT18\]](#page-12-6) if and only if (M, ω) is CS and a symplectic manifold $(M, e^{-f}\omega)$ satisfies the HLC in the classical sense. So the HLC defined here is weaker than the one defined in [\[AOT18\]](#page-12-6).

Remark 4.10. The hard Lefschetz theorem for Vaisman manifolds, which are a special class of LCK manifolds, is formulated in [\[CMDNMY19\]](#page-12-15), however, the formulation has nothing to do with the hard Lefschetz theorem formulated here at least in the context of this paper.

Since the HLC for symplectic manifolds is always satisfied on compact Kähler manifolds, the following conjecture is naturally suggested:

Conjecture 4.11. Let (M, J, g) be a 2n-dimensional compact LCK manifold with the Lee form θ . Then (M, J, g) satisfies the HLC.

Remark 4.12. On some LCS manifolds, the cohomology class of ω vanishes, i.e. $[\omega] = 0 \in H_1^2(M)$. For instance, LCS manifolds constructed from contact manifolds canonically [\[Vai76\]](#page-13-4), Vaisman manifolds, and LCK manifolds with potential $[OV23]$. In these cases, the Lefschetz map $[L]$ is a zero map. So the HLC for LCS is equivalent to saying that the vanishings

$$
H^j_{-(n-j)/2}(M) = 0
$$

hold for all $1 \leq j \leq n-1$.

5. Examples

As examples, we give LCaK manifolds defined in Section 4 of [\[AO18\]](#page-12-9), and compute the spaces defined in the previous sections with their basises. Furthermore, we check that these manifolds do not satisfy the HLC. The examples we give here are solvmanifolds. So before giving the examples, we collect some facts about solvmanifolds and methods to compute their twisted cohomologies.

Let G be a simply connected Lie group and $\mathfrak g$ be the associated Lie algebra. The space $\bigwedge^{\bullet} \mathfrak g^*$ is identified with left-invariant forms on G. We take a left-invariant closed 1-form $\theta \in \mathfrak{g}^*$. Since the twisted differential d_θ is left-invariant, we obtain a complex $(\bigwedge^{\bullet} \mathfrak{g}^*, d_\theta)$ called the *twisted Chevalley*-*Eilenberg complex.* Denote by $H^{\bullet}_{\theta}(\mathfrak{g}^*)$ its cohomology groups. A discrete subgroup Γ of G is called *lattice* if the quotient $\Gamma \backslash G$ is compact. The following theorem gives a method to compute the twisted cohomology of the manifold $\Gamma\backslash G$ from the twisted Chevalley-Eilenberg complex when G is completely solvable, i.e. the endomorphisms ad_X of $\mathfrak g$ have only real eigenvalues for all $X \in \mathfrak g$.

Theorem 5.1 ([\[Hat60\]](#page-12-16)). Let G be a simply connected and completely solvable Lie group with a lattice Γ and a left-invariant closed 1-form θ . Then the natural inclusion $(\bigwedge^{\bullet} \mathfrak{g}^*, d_{\theta}) \hookrightarrow (\mathcal{A}^{\bullet}(\Gamma \backslash G), d_{\theta})$ is a quasi-isomorphism, i.e. the inclusion induces an isomorphism in cohomology.

From now on, we only consider left-invariant forms. So we write $\bigwedge^j \mathfrak{g}^*$ as \mathcal{A}^j by abuse of notation. Let $m \geq 2$ be an integer. We shall define a compact LCaK solvmanifold (M_m, J, g) of dimension $2n = 2m + 2$.

We consider a Lie algebra $\mathfrak{g}_m := \mathbb{R} \ltimes_A \mathbb{R}^{2m+1}$, where A is given by the following diagonal matrix:

$$
A = \text{diag}\left(0, \frac{1}{m}, \frac{2}{m}, \dots, 1, -\frac{1}{m}, -\frac{2}{m}, \dots, -1\right).
$$

Let G_m be the associated simply connected Lie group. It is proved in [\[AO18\]](#page-12-9) that the manifolds G_m are completely solvable, and each of the manifolds admits at least one lattice Γ_m . Thus we can compute the cohomologies of the compact manifolds $M_m \coloneqq \Gamma_m \backslash G_m$ from the associated Lie algebras. We note that the dimensions of $H^j(M_m)$ and $H^j_\theta(M_m)$ are computed in [\[AO18\]](#page-12-9).

We take $(x^1, \dots, x^{m+1}, y^1, \dots, y^{m+1})$ as the standard basis of $\mathfrak{g}_{m}^* = (\mathbb{R} \times_A \mathbb{R}^{2m+1})^*$. We use the notation $x^{\alpha_1 \cdots \alpha_p} y^{\beta_1 \cdots \beta_q}$ for the form $x^{\alpha_1} \wedge \cdots \wedge x^{\alpha_p} \wedge y^{\beta_1} \wedge \cdots \wedge y^{\beta_q}$. The structure of $(\bigwedge^{\bullet} \mathfrak{g}_{m}^*, d)$ is determined by the Leibniz rule and the following equations:

$$
dx^{1} = 0,
$$

\n
$$
dx^{2} = 0,
$$

\n
$$
dx^{3} = -(1/m)x^{13},
$$

\n
$$
dx^{4} = -(2/m)x^{14},
$$

\n
$$
dx^{m+1} = -((m-1)/m)x^{1,m+1},
$$

\n
$$
dy^{n} = (2/m)x^{1}y^{2},
$$

\n
$$
dy^{n} = (3/m)x^{1}y^{4},
$$

\n
$$
dy^{m+1} = x^{1}y^{m+1}.
$$

We consider the following forms:

$$
\omega := \sum_{\alpha=1}^{m+1} x^{\alpha} y^{\alpha}, \quad \theta := \frac{1}{m} x^1.
$$

It is straightforward to check that the form ω endows M_m with a left-invariant LCS structure with the Lee form θ . Furthermore, we give a left-invariant almost complex structure J to M_m by considering $z^{\alpha} := x^{\alpha} + iy^{\alpha}$ as (1,0)-forms, or equivalently, by defining \mathcal{J} as

$$
\mathcal{J}x^{\alpha} = -y^{\alpha}, \quad \mathcal{J}y^{\alpha} = x^{\alpha}
$$

for all $1 \le \alpha \le m+1$. Then $(x^1, \dots, x^{m+1}, y^1, \dots, y^{m+1})$ is an orthogonal basis of \mathfrak{g}^* with respect to the Riemannian metric $g(\cdot, \cdot) := \omega(\cdot, J \cdot)$. Given these structures, (M_m, J, g) is a compact LCaK manifold with the fundamental form ω and the Lee form θ . We note that the manifolds (M_m, J, g) are not LCK, i.e. *J* is not integrable, as the exterior derivative of a $(1,0)$ -form $z^2 = x^2 + iy^2$,

$$
dz^{2} = d(x^{2} + iy^{2}) = \frac{1}{m}x^{1}y^{2}
$$

= $\frac{1}{4m}(z^{1}z^{2} - z^{1}\overline{z}^{2} + \overline{z}^{1}z^{2} - \overline{z}^{1}\overline{z}^{2}),$

is not contained in $\mathcal{A}^{2,0} \oplus \mathcal{A}^{1,1}$.

We shall compute the spaces defined in the previous sections on these LCaK manifolds. We fix the orthogonal basis of $\bigwedge^j \mathfrak{g}_m^*$ as follows:

$$
\mathcal{B}^j := \{ x^{\alpha_1 \cdots \alpha_p} y^{\beta_1 \cdots \beta_q} \in \bigwedge^j \mathfrak{g}_m^* \mid 1 \le \alpha_1 < \cdots < \alpha_p \le m+1, 1 \le \beta_1 < \cdots < \beta_q \le m+1 \}.
$$

We define the weight of each element of the basis $W: \mathcal{B}^{\bullet} \to \mathbb{R}$ as follows:

$$
W(x1) = 0,
$$

\n
$$
W(x2) = 0,
$$

\n
$$
W(x3) = -1,
$$

\n
$$
W(x3) = -1,
$$

\n
$$
W(y2) = 1,
$$

\n
$$
W(y3) = 2,
$$

\n
$$
W(y3) = 2,
$$

\n
$$
W(y4) = 3,
$$

\n
$$
W(ym+1) = m,
$$

 $W(x^{\alpha_1 \cdots \alpha_p} y^{\beta_1 \cdots \beta_q}) = W(x^{\alpha_1}) + \cdots + W(x^{\alpha_p}) + W(y^{\beta_1}) + \cdots + W(y^{\beta_q}), \quad W(1) = 0.$ Using this, we can calculate the twisted differential of $\gamma \in \mathcal{B}^{\bullet}$ as follows:

(5.1)
$$
d_k(\gamma) = \frac{W(\gamma) - k}{m} x^1 \wedge \gamma.
$$

We define the subspaces of \mathcal{A}^j as follows:

$$
\mathcal{A}_{k}^{j} := \left\langle \left\{ \gamma \in \mathcal{B}^{j} \mid W(\gamma) = k \right\} \right\rangle_{\mathbb{C}},
$$

$$
\hat{\mathcal{A}}_{k}^{j} := \left\langle \left\{ \gamma \in \mathcal{B}^{j} \mid W(\gamma) \neq k, x^{1} \wedge \gamma \neq 0 \right\} \right\rangle_{\mathbb{C}},
$$

where $\langle S \rangle_{\mathbb{C}}$ denotes a C-vector space generated by the elements of the set S. Then we obtain the Hodge decomposition of the complex $(\mathcal{A}^{\bullet}, d_k)$:

Proposition 5.2. The Hodge decomposition of the complex (A^{\bullet}, d_k) with respect to g is

$$
\mathcal{A}^j = \mathcal{A}^j_k \oplus x^1 \hat{\mathcal{A}}^{j-1}_k \oplus \hat{\mathcal{A}}^j_k
$$

.

,

More precisely,

$$
\mathcal{H}^j_k = \mathcal{A}^j_k, \quad d_k \mathcal{A}^{j-1} = x^1 \hat{\mathcal{A}}^{j-1}_k \quad and \quad d_k^* \mathcal{A}^{j+1} = \hat{\mathcal{A}}^j_k
$$

where x^1V denotes a vector space $\{x^1 \wedge v \mid v \in V\}$, and \mathcal{H}_k^j denotes the space ker $\Delta_k \cap \mathcal{A}^j$.

Proof. It is obvious that the decomposition is orthogonal. From (5.1) , it follows that the space $\mathcal{A}_k^j \oplus x^1 \hat{\mathcal{A}}_k^{j-1}$ is contained in ker d_k . Moreover, the map

$$
d_k \colon \hat{\mathcal{A}}_k^{j-1} \to x^1 \hat{\mathcal{A}}_k^{j-1}
$$

is an isomorphism. So we have $d_k(\mathcal{A}^{j-1}) = x^1 \hat{\mathcal{A}}_k^{j-1}$. Therefore, the space $\mathcal{A}_k^j \oplus \hat{\mathcal{A}}_k^j$ is contained in $\ker d_k^*$. So we have $d_k^*(A^{j+1}) = \hat{A}_k^j$ and $\mathcal{H}_k^j = A_k^j$.

From this proposition, we obtain

Proposition 5.3. The LCS manifolds (M_m, ω) do not satisfy the HLC for all $m \geq 2$.

Proof. If $m = 2m'$ is even, a form $x^{m'+2} \in \mathcal{H}^1_{-m'}$ is in the kernel of the Lefschetz operator, i.e. $[L]^m([x^{m'+2}]) = 0 \in H^{2m+1}_{m'}(M_m)$. If $m = 2m' + 1$ is odd, a form $x^{2,m'+2} \in \mathcal{H}^2_{m'}$, is in the kernel of the Lefschetz operator, i.e. $[L]^{m-1}([x^{2,m'+2}]) = 0 \in H^{2m}_{m'}(M_m)$. As a consequence, the LCS manifolds (M_m, ω) do not satisfy the HLC for all $m \geq 2$.

Now we shall compute the basises of the spaces \mathcal{H}_k^j , $\mathcal{H}_{k,\mathcal{J}}^j$ and $\mathcal{H}_k^{p,q}$ where, $n+2k-j=$ $n + 2k - (p + q) = 0$ for $m = 2, 3, 4$. We only consider the case $j \leq n$ as the basises in higher degrees are obtained by taking the Hodge dual. We note that \mathcal{H}^j_k is always zero if k is not integer or $j = 0$.

$$
\bullet \ \ m=2.
$$

The basis of each of the spaces are shown in Table [1.](#page-11-0) We also have $\mathcal{H}_k^{p,q} = \{0\}$ for all p, q. The Lefschetz map $[L]^2$: $H^1_{-1}(M_2) \to H^5_1(M_2)$ is as follows:

 $[x^3] \mapsto 0.$

 \bullet $m=3$.

The basis of each of the spaces are shown in Table [2.](#page-12-17) We also have

$$
\mathcal{H}_{-1}^{1,1} = \mathcal{H}_{\mathcal{J}}^{2} \cap \ker(\mathcal{J} - id)
$$

= $\langle x^{13} + y^{13} \rangle_{\mathbb{C}}$,

$$
\mathcal{H}_{0}^{2,2} = \mathcal{H}_{\mathcal{J}}^{4} \cap \ker(\mathcal{J} - id)
$$

= $\langle x^{134}y^{4} + x^{4}y^{134}, x^{124}y^{3} + x^{3}y^{124}, x^{123}y^{2} + x^{2}y^{123} \rangle_{\mathbb{C}}$,

and $\mathcal{H}_k^{p,q} = \{0\}$ for all p, q other than those above. The Lefschetz map $[L]^2: H^2_{-1}(M_3) \to H_1^6(M_3)$ on the elements which are not represented by the elements in $\mathcal{H}^2_{-1,\mathcal{J}}$ is as follows:

$$
[x^4y^2] \mapsto 0,
$$

$$
[x^{23}] \mapsto 0.
$$

 \bullet $m=4$.

The basis of each of the spaces are shown in Table [3.](#page-12-18) We also have $\mathcal{H}_k^{p,q} = \{0\}$ for all p, q. The Lefschetz map $[L]^4$: $H_{-2}^1(M_4) \rightarrow H_2^9(M_4)$ is as follows:

 $[x^4] \mapsto 0.$

The Lefschetz map $[L]^2: H^3_{-1}(M_4) \to H^7_1(M_4)$ on the elements which are not represented by the elements in $\mathcal{H}_{-1,\mathcal{J}}^3$ is as follows:

$$
[x^1y^{14}] \mapsto 2[*(x^{45}y^5 - x^{34}y^3 - x^{24}y^2)],
$$

\n
$$
[x^{45}y^5] \mapsto -2[*(x^1y^{14})],
$$

\n
$$
[x^{34}y^3] \mapsto 2[*(x^1y^{14})],
$$

\n
$$
[x^{24}y^2] \mapsto 2[*(x^1y^{14})],
$$

\n
$$
[x^{35}y^4] \mapsto 0,
$$

\n
$$
[x^{25}y^3] \mapsto 0.
$$

Therefore, the image of this map is

$$
\{[\gamma] \in H_1^7(M_4) \mid \gamma \in \langle \ast(x^{45}y^5 - x^{34}y^3 - x^{24}y^2), \ast(x^1y^{14}) \rangle_{\mathbb{C}} \oplus \mathcal{H}_{1,\mathcal{J}}^7 \}.
$$

TABLE 1. $m = 2$

space		\mathcal{H}_{-1}^2	\mathcal{H}^4_0
dim			10
basis			$\left[x^4y^2, x^{23}\right]$ $x^1y^{123}, x^{12}y^{14}, x^{234}y^4, x^{34}y^{23}$
			$\left\ \ln \mathcal{H}_{\bullet, \mathcal{J}}^{\bullet} \right\ x^{13} \pm y^{13} \left\ x^{134} y^4 \pm x^4 \overline{y^{134}, x^{124} y^3 \pm x^3 y^{124}} \right\ $ $x^{123}y^2 \pm x^2y^{123}$

TABLE 2. $m = 3$

TABLE 3. $m = 4$

REFERENCES

[Lee18] J. M. Lee, Introduction to Riemannian manifolds, second ed., Graduate Texts in Mathematics, vol. 176, Springer, Cham, 2018.

[LV15] H. V. Lê and J. Vanžura, Cohomology theories on locally conformal symplectic manifolds, Asian J. Math. 19 (2015), no. 1, 45–82.

[Nov82] S. P. Novikov, The Hamiltonian formalism and a many-valued analogue of Morse theory, Russian Mathematical Surveys 37 (1982), no. 5, 1–56.

[OV09] L. Ornea and M. Verbitsky, Morse-Novikov cohomology of locally conformally Kähler manifolds, J. Geom. Phys. 59 (2009), no. 3, 295–305.

graduate school of mathematical sciences, the university of tokyo, 3-8-1 komaba, meguro, tokyo 153-8914, japan.

Email address: shuho@ms.u-tokyo.ac.jp