On the Π_2^1 consequences of Π_1^1 -CA₀

YUDAI SUZUKI¹ and KEITA YOKOYAMA²

^{1,2}Mathematical Institute, Tohoku University, Sendai, Japan ¹yudai.suzuki.q1@dc.tohoku.ac.jp ²keita.yokoyama.c2@tohoku.ac.jp

February 13, 2024

Abstract

The system Π_1^1 -CA₀ is known as the strongest system of the *big five* in reverse mathematics. It is known that some theorems represented by a Π_2^1 sentence, for example Kruskal's theorem, are provable from Π_1^1 -CA₀ but not provable from the second strongest system ATR₀ of the big five. However, since any Π_2^1 sentence is not equivalent to Π_1^1 -CA₀, Π_1^1 -CA₀ is too strong to prove such theorems. In this paper, we introduce a hierarchy dividing the set { $\sigma \in \Pi_2^1 : \Pi_1^1$ -CA₀ $\vdash \sigma$ }. Then, we give some characterizations of this hierarchy using some principles equivalent to Π_1^1 -CA₀: leftmost path principle, Ramsey's theorem for Σ_n^0 classes of [N]^N and determinacy for $(\Sigma_1^0)_n$ classes of N^N. As an application, our hierarchy explicitly shows that the number of application of the hyperjump operator needed to prove Σ_n^0 Ramsey's theorem or $(\Sigma_1^0)_n$ determinacy increases when the subscript *n* increases.

1 Introduction

Reverse mathematics is a program to classify theorems of mathematics according to their logical strength. In the most typical study of this area, one uses the so-called *big five* of axiomatic systems of second-order arithmetic, RCA_0 , WKL_0 , ACA_0 , ATR_0 and Π_1^1 - CA_0 . That is, for a given theorem, one shows the equivalent between the theorem and one of the big five.

In recent studies of reverse mathematics, it is also interested to find theorems that are not equivalent to any of the big five. For example, it is known that Kruskal's theorem on the embeddings of finite trees is not provable from ATR_0 and provable from Π_1^1 - CA_0 , and it is strictly weaker than Π_1^1 - CA_0 .

In [8], H. Towsner introduced relative leftmost path principle to give a new upper bound for theorems located between ATR_0 and Π_1^1 - CA_0 . He focused on an equivalent of Π_1^1 - CA_0 called the leftmost path principle which states that any ill-founded tree has a leftmost path. His argument is based on the following idea. Since the leftmost path principle is a Π_3^1 statement, the actual leftmost path is not needed to prove a Π_2^1 sentence from the leftmost path principle. Instead of the leftmost path, it is enough to use a path which behaves as a leftmost path in a certain range. Such a path is called relative leftmost path.

In this paper, we extend the idea of relativization of Π_3^1 statements to the level of *n*-th hyperjump. We define $\beta_0^1 \operatorname{\mathsf{RFN}}(n)$ as the assertion that for each set X, there is a coded ω -model \mathcal{M} of ACA₀ such that $X \in \mathcal{M}$ and $\mathcal{M} \models \exists Y(Y = \operatorname{HJ}^n(X))$. Then a witness Y works as a good approximation of the actual *n*-th hyperjump. We show that a Π_2^1 sentence provable from Π_1^1 -CA₀ is already provable from $\beta_0^1 \operatorname{\mathsf{RFN}}(n)$ for some *n*.

Intuitively, a sentence σ is provable from $\beta_0^1 \operatorname{RFN}(n)$ means that there is a proof of σ from Π_1^1 -CA₀ such that the use of the hyperjump operator in it is up to *n*-times. In this sense, Towsner's transfinite leftmost path principle TLPP is in the level of single use of the hyperjump operator. Specifically we show that there is a variant of relative leftmost path principle which is equivalent to $\beta_0^1 \operatorname{RFN}(1)$, and a variant of $\beta_0^1 \operatorname{RFN}(1)$ is enough to prove TLPP.

We apply the idea of relativization to Ramsey's theorem for $[\mathbb{N}]^{\mathbb{N}}$ and the determinacy for $\mathbb{N}^{\mathbb{N}}$. It is known that both of Ramsey's theorem for Σ_n^0 classes and the determinacy for $(\Sigma_1^0)_n$ classes are equivalent to Π_1^1 -CA₀ for n > 1. In this paper, we show that the Π_2^1 consequences of Π_1^1 -CA₀ is coincide its of the theory ACA₀ + {rel(Σ_n^0 -Ram) : $n \in \omega$ } or ACA₀ + {rel($(\Sigma_1^0)_n$ -Det) : $n \in \omega$ } where rel(σ) denotes the relativization of σ . These result imply that the number of use of the hyperjump operator to prove Σ_n^0 Ramsey's theorem or $(\Sigma_1^0)_n$ determinacy increases when the subscript n increases.

Acknowledgement

The first author's work is supported by JST, the establishment of university fellowships towards the creation of science technology innovation, Grant Number JPMJFS2102. The second author's work is partially supported by JSPS KAKENHI grant numbers 19K03601 and 21KK0045.

2 Preliminaries

In this section, we introduce basic notions for reverse mathematics. For the details, see also Simpson's textbook [4]. As usual, we write ω for the set of standard natural numbers and \mathbb{N} for the range of number variables in the language \mathcal{L}_2 of second-order arithmetic.

Reverse mathematics is a research program whose aim is to classify mathematical theorems according to their logical strength. For this purpose, we identify the logical strength of a theorem as an axiom which is needed and sufficient to prove it. The most typical researches of reverse mathematics is based on second-order arithmetic¹. That is, for a given formula T of second-order arithmetic which represents a mathematical theorem, we find an axiomatic system Γ and an axiom A such that

- Γ is sufficient to interpret T but may not be sufficient to prove T,
- A is equivalent to T over Γ .

We begin with introducing some axiomatic systems of second-order arithmetic.

Definition 2.1. Let $\theta(y, \vec{x}, \vec{X})^2$ be a formula with exactly displayed free variables. Comprehension for θ denotes the formula

$$\forall \vec{x}, \vec{X} \exists Y \forall y (y \in Y \leftrightarrow \theta(y, \vec{x}, \vec{X})).$$

Intuitively, comprehension for θ states that the set $\{y : \theta(y)\}$ exists.

- Some \mathcal{L}_2 -theories are characterized by comprehension schemas.
- RCA₀ consists of '(0, 1, +, ·, <) forms a discrete ordered semi-ring', induction for Σ⁰₁ formulas and comprehension for Δ⁰₁ formulas.
- ACA_0 consists of RCA_0 and comprehension for Σ_0^1 formulas.
- ATR_0 consists of RCA_0 and transfinite recursion for arithmetical formulas.
- Π_1^1 -CA₀ consists of RCA₀ and comprehension for Π_1^1 formulas.

 $^{^1 \}rm Sometimes$ the phrase 'second-order arithmetic' means a specific axiomatic system $Z_2.$ However, in this paper, we use this phrase as a general term for axiomatic systems of the language of second-order arithmetic.

²Here, \vec{x} and \vec{X} are abbreviations of x_1, \ldots, x_n and X_1, \ldots, X_m . When the numbers n and m of variables are not important, we use these notations.

In RCA_0 , the Turing jump operator and its iteration can be defined. We write $X^{(n)}$ and $X^{(\omega)}$ to mean the *n*-times and N-times iteration of Turing jump at X respectively. For the details, see section VIII.1 in [4].

Definition 2.2. We define ACA_0^{\prime} and ACA_0^{+} as follows.

- ACA'_0 consists of RCA_0 and $\forall n \forall X \exists Y(Y = X^{(n)})$.
- ACA_0^+ consists of RCA_0 and $\forall X \exists Y(Y = X^{(\omega)})$.

It is known that over RCA_0 , $\mathsf{ACA}_0 < \mathsf{ACA}_0' < \mathsf{ACA}_0^+$.

2.1 Trees

We introduce some notions on trees. Then we give a characterization of Π_1^1 -CA₀ by trees.

Definition 2.3. Let X be a set. We write $X^{<\mathbb{N}}$ for the set of finite sequences of elements of X and $X^{\mathbb{N}}$ for the set of infinite sequences of elements of X. More precisely, each $\sigma \in X^{<\mathbb{N}}$ is a function whose domain is $\{0, 1, \ldots, n-1\}$ for some n and $\sigma(i)$ is an element of X for any i < n. Similarly, each $f \in X^{\mathbb{N}}$ is a function whose domain is \mathbb{N} and whose codomain is X.

For each $\sigma \in X^{<\mathbb{N}}$, define the length $|\sigma|$ by the number n such that $\operatorname{dom}(\sigma) = \{0, 1, \ldots, n-1\}.$

Intuitively, we identify $\sigma \in X^{<\mathbb{N}}$ and the sequence $\langle \sigma(0), \ldots, \sigma(|\sigma|-1) \rangle$, $f \in X^{\mathbb{N}}$ and the sequence $\langle f(n) \rangle_{n \in \mathbb{N}}$.

Definition 2.4. Let σ, τ be finite sequences.

- Define the concatenation $\sigma * \tau$ by $\langle \sigma(0), \ldots, \sigma(|\sigma|-1), \tau(0), \ldots, \tau(|\tau|-1) \rangle$.
- We say σ is an initial segment of τ (write $\sigma \leq \tau$) if $|\sigma| \leq |\tau|$ and $\forall i < |\sigma|(\sigma(i) = \tau(i))$). We say σ is a proper initial segment of τ (write $\sigma \prec \tau$) if $\sigma \leq \tau$ and $\sigma \neq \tau$.
- Let f be an infinite sequence. We say σ is an initial segment of f (write $\sigma \prec f$) if $\forall i < |\sigma|(\sigma(i) = f(i))$.

Definition 2.5. Let T be a subset of $X^{<\mathbb{N}}$. We say T is a tree on X if T is closed under taking initial segments.

Let $T \subseteq X^{<\mathbb{N}}$ be a tree. A function $f \in X^{\mathbb{N}}$ is called a path of T if $(\forall n)(f[n] \in T)$. Here, f[n] denotes the initial segment of f with length = n. The set of all paths of T is denoted by [T]. We say T is ill-founded if $[T] \neq \emptyset$.

Lemma 2.6. ([4], III.7.2) ACA_0 proves the following Kőnig lemma. Let T be an infinite tree. If $\forall \sigma \in T \exists m \forall i (\sigma * \langle i \rangle \in T \rightarrow i < m)$, then T is ill-founded.

Lemma 2.7. [4, VI.1.1] The following are equivalent over RCA₀.

1. Π_1^1 -CA₀,

2. for any sequence $\langle T_k \rangle_k$ of trees, there is a set $X = \{k : [T_k] \neq \emptyset\}$.

Following the above lemma, we give a new characterization of Π_1^1 -CA₀ by using the notion of leftmost path.

Definition 2.8. (RCA₀) Let $f, g \in \mathbb{N}^{\mathbb{N}}$. We write $f <_l g$ if $\exists n(f[n] = g[n] \land f(n) < g(n))$ and $f \leq_l g$ if $f <_l g \lor f = g$. This $<_l$ forms a total order on $\mathbb{N}^{\mathbb{N}}$ and is called the lexicographical order.

Let T be a tree and $f \in [T]$. We say f is the leftmost path of T if $\forall g \in [T](f \leq_l g)$.

In the remaining of this section, we show that the equivalence of the existence of a leftmost path and Π_1^1 -CA₀. The key is that a sequence of trees can be coded by a tree.

Definition 2.9. (RCA₀) Let $\sigma \in \mathbb{N}^{<\mathbb{N}}$. For each $l < |\sigma|$, define n_l as the maximum n such that $(n, l) < |\sigma|$. Then we define $\sigma_l = \langle \sigma((0, l)), \ldots, \sigma((n_l, l)) \rangle$. Thus, each $\sigma \in \mathbb{N}^{<\mathbb{N}}$ is regarded as a sequence $\langle \sigma_l \rangle_{l < |\sigma|}$ of sequences.

Conversely, for each sequence $\langle \sigma_l \rangle_{l < L}$ of sequences, define $\bigoplus_{l < L} \sigma_l$ by

$$(\bigoplus_{l < L} \sigma_l)(n, l) = \sigma_l(n).$$

For each sequences $\langle f_l \rangle_{l \in \mathbb{N}}$ of functions, define $\bigoplus_l f_l$ by

$$(\bigoplus_{l < L} f_l)(n, l) = f_l(n).$$

Finally, for each $\langle T_l \rangle_{l \in \mathbb{N}}$ of trees, define $\bigoplus_l T_l$ by

$$\{\bigoplus_{l< L} \sigma_l : \langle \sigma_l \rangle_{l< L} \in T_0 \times \cdots \times T_{L-1}\}.$$

Lemma 2.10. (RCA₀) Let $\langle T_l \rangle_l$ be a sequence of trees. Then, the operator \bigoplus_l is a bijection between $\prod_l [T_l]$ and $[\bigoplus_l T_l]$. Moreover, this operator preserves the lexicographical order in the sense that for any $\langle f_i \rangle_i, \langle g_i \rangle_i \in \prod_i [T_i]$,

- 1. $\forall i (f_i \leq_l g_i) \to \bigoplus_i f_i \leq_l \bigoplus_i g_i,$
- 2. if $\bigoplus_i f_i$ is the leftmost path of $\bigoplus_i T_i$, then each f_i is the leftmost path of T_i .

Definition 2.11. Let T be a tree. We say T is pruned if $\forall \sigma \in T \exists n (\sigma * \langle n \rangle \in T)$.

Lemma 2.12. (RCA₀) Let T be a pruned tree and $\sigma \in T$. Then there is a T-computable path f such that $\sigma \prec f$. Moreover, we can take this f to be the leftmost of all paths extending σ .

Proof. It is immediate from the definition.

Theorem 2.13. [1, Theorem 6.5.] The following are equivalent over RCA_0 .

1. Π_1^1 -CA₀,

2. each ill-founded tree has the leftmost path.

Proof. $(1 \to 2)$. Let T be an ill-founded tree. Define $S = \{\sigma \in T : \exists f \in [T](\sigma \prec f)\}$. Then S is a pruned tree such that [S] = [T]. Since S is pruned, S has the leftmost path. Clearly this is also the leftmost path of [T].

 $(2 \to 1)$ It is enough to show that the second clause implies for any sequence $\langle T_k \rangle_k$ of trees, there is a set $X = \{k : [T_k] \neq \emptyset\}$.

Assume that each ill-founded tree has the leftmost path. Let $\langle T_k \rangle_k$ be a sequence of trees. For each k, define S_k by $\{\langle 0 \rangle * \sigma : \sigma \in T_k\} \cup \{1^l : l \in \mathbb{N}\}$ where 1^l is the sequence with length = l such that each element of it is 1. Then, each S_k has a path $1^{\infty} = \langle 1, 1, \ldots, \rangle$. Thus, $\bigoplus_k S_k$ is also ill-founded.

By the assumption, take the leftmost path f of $\bigoplus_k S_k$ and put $X = \{k : f(0,k) = 0\}$. It is easy to check that $X = \{k : [T_k] \neq \emptyset\}$.

2.2 Coded ω -models

We introduce the definition of coded ω -models in RCA_0 and give some basic properties.

Definition 2.14. (RCA₀) For a set A and a family $\langle X_i \rangle_i$, we say A is in $\langle X_i \rangle_i$ or $\langle X_i \rangle_i$ contains A (write $A \in \langle X_i \rangle_i$) if $\exists i(A = X_i)$. A set \mathcal{M} is identified with an \mathcal{L}_2 -structure $(\mathbb{N}, \langle \mathcal{M}_i \rangle_i, 0, 1, +, \cdot, \in)$. In this sense, we call \mathcal{M} a coded ω -model.

Remark 2.15. Let \mathcal{M} be a coded ω -model. For a sentence σ with parameters from \mathcal{M} , the satisfaction $\mathcal{M} \models \sigma$ is defined by the existence of an evaluating function for σ which represents Tarski's truth definition and ensures σ is true. If we work in ACA₀, then $\mathcal{M} \models \sigma$ iff the relativization $\sigma^{\mathcal{M}}$ is true. Especially, if σ is arithmeical, then $\mathcal{M} \models \sigma$ iff σ is true.

Definition 2.16. Let σ be an \mathcal{L}_2 -sentence. Define the assertion ω -model reflection of σ by

 $\forall X \exists \mathcal{M} : \omega \text{-model } (X \in \mathcal{M} \land \mathcal{M} \models \sigma).$

Lemma 2.17. Let σ be a Π_2^1 sentence. Then, over ACA₀, the ω -model reflection of σ implies $\sigma + \text{Con}(\sigma)$ where $\text{Con}(\sigma)$ is the Π_1^0 formula stating that ' σ is consistent'. Therefore, if $\text{Con}(\sigma)$ is not provable from ACA₀, then the ω -model reflection of σ is strictly stronger than σ .

Proof. Write $\sigma \equiv \forall X \exists Y \theta(X, Y)$ by an arithmetical formula θ . Assume ACA₀ and the ω -model reflection of σ . We show that $\sigma + \text{Con}(\sigma)$ holds. Let A be a set. By the ω -model reflection, take an ω -model \mathcal{M} such that $A \in \mathcal{M} \land \mathcal{M} \models \forall X \exists Y \theta(X, Y)$. Then, $\mathcal{M} \models \exists Y \theta(A, Y)$ and hence $\theta(A, B)$ holds for some $B \in \mathcal{M}$. In addition, since \mathcal{M} is a (weak-)model of σ , $\text{Con}(\sigma)$ holds.

The following characterization of ACA_0^+ is well-known.

Theorem 2.18. Over RCA_0 , the following are equivalent.

- 1. ACA_0^+ ,
- 2. the ω -model reflection of ACA₀; any set is contained in an ω -model of ACA₀,

2.3 Hyperjump and β -model reflection

In this section, we introduce the notion of hyperjump and β -models. Hyperjump is a Σ_1^1 analogue of Turing jump. As ACA₀ is characterized by Turing jumps, Π_1^1 -CA₀ is characterized by hyperjumps.

Definition 2.19. (Universal Σ_1^1 formula) Let $\mathsf{Sat}_{\Pi_0^0}$ be a universal Π_0^0 formula. We define the Σ_1^1 universal formula σ_1^1 by

$$\sigma_1^1(e, x, X) \equiv \exists f \forall y \mathsf{Sat}_{\Pi_0^0}(e, x, X, f[y]).$$

Definition 2.20. (ACA₀) Let \mathcal{M} be a coded ω -model. We say \mathcal{M} is a coded β -model if

$$\forall X \in \mathcal{M} \forall e, x(\sigma_1^1(e, x, X) \leftrightarrow \mathcal{M} \models \sigma_1^1(e, x, X)).$$

Intuitively, a coded β -model is a Σ_1^1 absolute ω -model. In fact, we can show the following.

Lemma 2.21. Let $\theta(x, X)$ be a Σ_1^1 formula. Then, ACA₀ proves that for any β -model $\mathcal{M}, \forall X \in \mathcal{M} \forall x(\theta(x, X) \leftrightarrow \mathcal{M} \models \theta(x, X))$.

Proof. Immediate from the definition of β -models.

. . . .

We next introduce the notion of hyperjump. Intuitively, the hyperjump operator is an operator to make a $\Sigma_1^{1,X}$ complete set from a set X.

Definition 2.22. (ACA₀) Let X be a set. Define the hyperjump HJ(X) by $HJ(X) = \{(e, x) : \sigma_1^1(e, x, X)\}$. We write $HJ^n(X)$ for the *n*-times iteration of the hyperjump.

Lemma 2.23. Let $\varphi(x, y, X)$ be a Σ_1^1 formula with exactly displayed free variables. Then, ACA₀ proves

$$\forall y, X, Y(Y = \mathrm{HJ}(X) \to \exists Z \leq_{\mathrm{T}} Y(Z = \{x : \varphi(x, y, X)\})).$$

That is, any $\Sigma_1^{1,X}$ -definable set is computable from $\mathrm{HJ}(X)$. Hence, any $\Pi_1^{1,X}$ -definable set is also computable from $\mathrm{HJ}(X)$.

Proof. Immediate from the definition of HJ.

Lemma 2.24 ([4] Lemma VII.2.9). Over ACA₀, for any set X, the existence of HJ(X) is equivalent to the existence of a coded β -model containing X.

Combining above two lemmas, we have that

Theorem 2.25. The following assertions are equivalent over ACA_0 .

- 1. Π_1^1 -CA₀,
- 2. $\forall X \exists Y(Y = HJ(X)),$
- 3. β -model reflection : $\forall X \exists \mathcal{M} : \beta$ -model ($X \in \mathcal{M}$).

We note that all of these assertions are Π_1^1 statements such that any of them is never equivalent to a Π_2^1 statement. In this paper, we introduce Π_2^1 variants of the second and third assertions to consider the structure of Π_2^1 consequences of Π_1^1 -CA₀.

At last, we see that a hyperjump in the ground model behaves a hyperjump in a coded ω -model.

Lemma 2.26. Let \mathcal{M} and \mathcal{M}' be models of ACA₀ such that $\mathbb{N}^{\mathcal{M}} = \mathbb{N}^{\mathcal{M}'}$ and $S^{\mathcal{M}'} \subseteq S^{\mathcal{M}}$. Here, $\mathbb{N}^{\mathcal{M}}$ is the set of numbers of \mathcal{M} and $S^{\mathcal{M}}$ is the set of sets of \mathcal{M} . Similarly, $\mathbb{N}^{\mathcal{M}'}$ is the set of numbers of \mathcal{M}' and $S^{\mathcal{M}'}$ is the set of sets of \mathcal{M}' .

Let $A, B \in \mathcal{M}'$ such that $\mathcal{M} \models B = \mathrm{HJ}(A)$. Then, $\mathcal{M}' \models B = \mathrm{HJ}(A)$. Moreover, if \mathcal{M}' is a coded ω -model in \mathcal{M} , then $\mathcal{M} \models (B = \mathrm{HJ}(A))^{\mathcal{M}'}$.

Proof. Take $\mathcal{M}, \mathcal{M}', A$ and B be as above. We first show the following claim.

Claim 2.26.1. Let $\theta(X)$ be a standard Σ_1^1 formula having exactly one set variable. Then, $\mathcal{M} \models \theta(A)$ iff $\mathcal{M}' \models \theta(A)$.

Proof of Claim. It is enough to show that $\mathcal{M} \models \theta(A)$ implies $\mathcal{M}' \models \theta(A)$. Write $\theta(X) \equiv \exists Z \varphi(X, Z)$ by an arithmetical formula φ . Assume $\mathcal{M} \models \exists Z \varphi(A, Z)$. We will show that $\mathcal{M}' \models \exists Z \leq_{\mathrm{T}} B \varphi(A, B)$. Now there is a Σ_0^0 formula θ such that both of $\mathcal{M}, \mathcal{M}'$ satisfies $\forall Z(\varphi(A, Z) \leftrightarrow \exists f \forall n \theta(A, Z[n], f[n]))$. Define trees $T, S \subseteq (2 \times \mathbb{N})^{<\mathbb{N}}$ by

$$T = \{(s,\sigma) : \forall (t,\tau) \preceq (s,\sigma)\theta(A,t,\tau)\}$$
$$S = \{(s,\sigma) : \exists (Z,f) \in [T]((s,\sigma) \prec (Z,f))\}$$

Then, T is Δ_1^0 in A and hence S is Σ_1^1 in X. Therefore, $S \leq_{\mathrm{T}} \mathrm{HJ}(A)$. Moreover, since $\exists Z\varphi(X,Z)$ holds, S is a nonempty pruned tree. Thus S has a S-computable path (Z, f). Now $Z \leq_{\mathrm{T}} S \leq_{\mathrm{T}} \mathrm{HJ}(X)$, so this completes the proof.

Now $B = \{(e, x) \in \mathbb{N}^{\mathcal{M}} : \mathcal{M} \models \sigma_1^1(e, x, A)\} = \{(e, x) \in \mathbb{N}^{\mathcal{M}'} : \mathcal{M}' \models \sigma_1^1(e, x, A)\}$. Therefore $\mathcal{M}' \models B = \mathrm{HJ}(A)$.

Lemma 2.27. (ACA₀) Let X, Y be sets such that Y = HJ(X). Then, for any coded ω -model \mathcal{M} of ACA₀, if $Y \in \mathcal{M}$, then $X \in \mathcal{M} \land \mathcal{M} \models Y = HJ(X)$.

Proof. We reason in ACA₀. Take a pair of sets X, Y such that Y = HJ(X). Let \mathcal{M} be a coded ω -model such that $Y \in \mathcal{M}$.

We first show that $X \in \mathcal{M}$. Take a number e such that

$$\forall Z(x \in Z \leftrightarrow \sigma_1^1(e, x, Z))$$

Then, $X = \{x : (e, x) \in Y\}$. Since $X \leq_{\mathrm{T}} Y, X \in \mathcal{M}$.

Therefore, $\mathcal{M} \models [Y = HJ(X)]$ remains but this is immediate from the above lemma.

Remark 2.28. In general, even if $(Y = HJ(X))^{\mathcal{M}}$ holds for a coded ω -model $\mathcal{M}, Y = HJ(X)$ may not hold. This is immediate from some results in the next section.

3 Approximation of Π_1^1 -CA₀

In this section, we introduce Π_2^1 variants of the β -model reflection $\beta_0^1 \operatorname{\mathsf{RFN}}(n)$. Then we prove that the theories $\operatorname{\mathsf{ACA}}_0 + \{\beta_0^1 \operatorname{\mathsf{RFN}}(n) : n \in \omega\}$ and $\{\sigma \in \Pi_2^1 : \Pi_1^1 \operatorname{\mathsf{-CA}}_0 \vdash \sigma\}$ proves the same sentences.

As we have seen in Theorem 2.25, Π_1^1 -CA₀ is characterized by the hyperjump operator or the β -model reflection. This means that in Π_1^1 -CA₀, we can take any (standard) finite iteration of hyperjump or β -models. More precisely, the following holds.

Observation 3.1. For any $n \in \omega$, n > 0, the following assertions are equivalent over ACA₀.

- Π_1^1 -CA₀,
- $\forall X \exists Y (Y = HJ^n(X)),$
- $\forall X \exists \mathcal{M}_0, \dots, \mathcal{M}_{n-1} : \beta$ -models $(X \in \mathcal{M}_0 \in \dots \in \mathcal{M}_{n-1})$.

From this observation, we can conjecture that theorems provable from Π_1^1 -CA₀ are classified by the number of the hyperjump operator used to proved those theorems. The following theorem ensures this conjecture.

Theorem 3.2. Let $\theta(X, Y, Z)$ be an arithmetical formula such that Π_1^1 -CA₀ $\vdash \forall X \exists Y \forall Z \theta(X, Y, Z)$. Then, there exists $n \in \omega$ such that

$$\mathsf{ACA}_0 \vdash \forall X, W(W = \mathrm{HJ}^n(X) \to \exists Y \leq_{\mathrm{T}} W \forall Z \theta(X, Y, Z))).$$

Proof. Assume $\theta(X, Y, Z)$ be an arithmetical formula such that Π_1^1 -CA₀ $\vdash \forall X \exists Y \forall Z \theta(X, Y, Z)$. For the sake of contradiction, suppose that for any $n \in \omega$, ACA₀ + $\exists X, W(W = HJ^n(X) \land \forall Y \leq_T W \exists Z \neg \theta(X, Y, Z))$ is consistent.

Let C and D be new constant set symbols. Let $\mathcal{L}_2(C, D)$ be the language extending \mathcal{L}_2 by adding C, D. Define an $\mathcal{L}_2(C, D)$ theory T by

$$T = \mathsf{ACA}_0 + \{ D_0 = C \land D_1 = \mathrm{HJ}(D_0) \land \dots \land D_k = \mathrm{HJ}(D_{k-1}) : k \in \omega \} + \{ \forall Y \leq_{\mathrm{T}} D_k \exists Z \neg \theta(C, Y, Z) : k \in \omega \}.$$

Then, by compactness theorem, T has a model $\mathcal{M} = (\mathbb{N}, \widetilde{S}, C, D)$. Define $S = \bigcup_{n \in \omega} \{X \in \widetilde{S} : \mathcal{M} \models X \leq_{\mathrm{T}} D_n\}.$

We show that (\mathbb{N}, S) is a model of Π_1^1 -CA₀ + $\neg \forall X \exists Y \forall Z \theta(X, Y, Z)$. We note that $C \in S$ and $D_n \in S$ for all $n \in \omega$. By Lemma 2.27, $(\mathbb{N}, S) \models$ $D_{n+1} = \operatorname{HJ}(D_n)$ for all n. Moreover, for any $X \in S$, there exists $Y \in S$ such that $\mathcal{M} \models Y = \operatorname{HJ}(X)$. Thus, $(\mathbb{N}, S) \models \forall X \exists Y(Y = \operatorname{HJ}(X))$ and (\mathbb{N}, S) is a β -submodel of $(\mathbb{N}, \widetilde{S})$. Let $Y \in S$. Then, $(\mathbb{N}, \widetilde{S}) \models Y \leq_{\mathrm{T}} D_k$ for some k and thus $(\mathbb{N}, \widetilde{S}) \models \exists Z \neg \theta(C, Y, Z)$. Hence $(\mathbb{N}, S) \models \exists Z \neg \theta(C, Y, Z)$. Therefore, $(\mathbb{N}, S) \models \neg \forall X \exists Y \forall Z \theta(X, Y, Z)$.

Remark 3.3. We note that this theorem generalizes the result by Montalban and Shore [2, Theorem 6.8.]. They proved that if

$$\Pi_1^1 \text{-} \mathsf{CA}_0 \vdash \forall X \exists Y \forall Z \theta(X, Y, Z),$$

then there exists $n \in \omega$ such that

$$\Pi_1^1 \text{-} \mathsf{CA}_0 \vdash \forall X \exists Y \leq_{\mathrm{T}} \mathrm{HJ}^n(X) \forall Z \theta(X, Y, Z))).$$

We give a Π_2^1 variant of the statement that any set is contained in a coded β -model. For this purpose, we consider β -submodels of a coded ω -model of ACA₀ instead of the acutual β -models.

Definition 3.4. (ACA₀) Let \mathcal{M}_0 and \mathcal{M}_1 be coded ω -models such that $\mathcal{M}_0 \in \mathcal{M}_1$. We say \mathcal{M}_0 is a coded β -submodel of \mathcal{M}_1 if

$$\forall X \in \mathcal{M}_0 \forall e, x(\mathcal{M}_0 \models \sigma_1^1(e, x, X) \leftrightarrow \mathcal{M}_1 \models \sigma_1^1(e, x, X))$$

holds. We write $\mathcal{M}_0 \in_{\beta} \mathcal{M}_1$ to mean \mathcal{M}_0 is a coded β -submodel of \mathcal{M}_1 .

Definition 3.5. For each $n \in \omega$, define $\beta_0^1 \operatorname{RFN}(n)$ as the following assertion:

 $\forall X \exists \mathcal{M}_0, \dots, \mathcal{M}_n : \text{coded } \omega \text{-models } (X \in \mathcal{M}_0 \in_\beta \dots \in_\beta \mathcal{M}_n \land \mathcal{M}_n \models \mathsf{ACA}_0).$

Remark 3.6. This form of ω -model reflection is first introduced in [3] as their $\psi_0(1, n)$. They mainly considered any (nonstandard) length of sequences of β -submodels, but we mainly consider standard length of those.

It is easy to see that the sequence $(\beta_0^1 \operatorname{RFN}(0), \beta_0^1 \operatorname{RFN}(1), \dots,)$ is strictly increasing. More precisely, the following holds.

Lemma 3.7. Let $n \in \omega$. Then, over ACA₀, $\beta_0^1 \operatorname{RFN}(n+1)$ proves the ω model reflection of $\beta_0^1 \operatorname{RFN}(n)$. Especially, $\beta_0^1 \operatorname{RFN}(n)$ is strictly stronger
than $\beta_0^1 \operatorname{RFN}(n+1)$.

Proof. Assume $\mathsf{ACA}_0 + \beta_0^1 \mathsf{RFN}(n+1)$. Take a sequence of ω -models such that $\mathcal{M}_0 \in_\beta \mathcal{M}_1 \cdots \in_\beta \mathcal{M}_n \land \mathcal{M}_n \models \mathsf{ACA}_0$. We show that $\mathcal{M}_0 \models \beta_0^1 \mathsf{RFN}(n)$. For any $X \in \mathcal{M}_0$, \mathcal{M}_n satisfies

$$\exists \mathcal{N}_0, \cdots, \mathcal{N}_{n-1} (X \in \mathcal{N}_0 \in_{\beta} \cdots \in_{\beta} \mathcal{N}_{n-1} \land \mathcal{N}_{n-1} \models \mathsf{ACA}_0)$$

via taking $(\mathcal{N}_0, \ldots, \mathcal{N}_{n-1}) = (\mathcal{M}_0, \ldots, \mathcal{M}_{n-1})$. Since this is a Σ_1^1 formula and \mathcal{M}_0 is a β -submodel of \mathcal{M}_n , \mathcal{M}_0 also satisfies

$$\exists \mathcal{N}_0, \cdots, \mathcal{N}_{n-1} (X \in \mathcal{N}_0 \in_{\beta} \cdots \in_{\beta} \mathcal{N}_{n-1} \land \mathcal{N}_{n-1} \models \mathsf{ACA}_0).$$

Since X is arbitrary, $\mathcal{M}_0 \models \beta_0^1 \operatorname{\mathsf{RFN}}(n)$.

Lemma 3.8. Over ACA₀, $\beta_0^1 \operatorname{RFN}(n)$ is equivalent to the following assertion:

 $\forall X \exists \mathcal{M} : \text{coded } \omega \text{-model } (X \in \mathcal{M} \land \mathcal{M} \models \mathsf{ACA}_0 + \exists Y(Y = \mathrm{HJ}^n(X))).$

Proof. Immediate from the fact the existence of HJ(X) is equivalent to the existence of a β -model containing X over ACA_0 .

Theorem 3.9. Let $\theta(X, Y)$ be an arithmetical formulas such that $\forall X \exists Y \theta(X, Y)$ is provable from Π_1^1 -CA₀. Then, there exists $n \in \omega$ such that $\mathsf{ACA}_0 + \beta_0^1 \mathsf{RFN}(n)$ proves $\forall X \exists Y \theta(X, Y)$.

Proof. Let σ_n be the following sentence.

 $\forall X \exists \mathcal{M} : \text{coded } \omega \text{-model } (X \in \mathcal{M} \land \mathcal{M} \models \mathsf{ACA}_0 + \exists Y(Y = \mathrm{HJ}^n(X))).$

Then, $\beta_0^1 \operatorname{RFN}(n)$ is equivalent to σ_n over ACA₀.

We show that $\forall X \exists Y \theta(X, Y)$ is provable from the theory $\mathsf{ACA}_0 + \{\sigma_n : n \in \omega\}$. Since $\forall X \exists Y \theta(X, Y)$ is provable from $\Pi_1^1 \text{-}\mathsf{CA}_0$, there exists $n \in \omega$ such that

$$\mathsf{ACA}_0 \vdash \forall X, W(W = \mathrm{HJ}^n(X) \to \exists Y \leq_{\mathrm{T}} W\theta(X, Y))$$

by Thereom 3.2. Take such an n. Assume $ACA_0 + \sigma_n$. Let X be an arbitrary set. Then, there is a coded ω -model \mathcal{M} such that

$$X \in \mathcal{M} \land \mathcal{M} \models \mathsf{ACA}_0 + \exists W(W = \mathrm{HJ}^n(X)).$$

Take such \mathcal{M} and W. Then $\mathcal{M} \models \exists Y \leq_{\mathrm{T}} W\theta(X,Y)$ and hence $\mathcal{M} \models \exists Y\theta(X,Y)$. Since θ is arithmetical, we have $\exists Y\theta(X,Y)$. This completes the proof.

Since each $\beta_0^1 \operatorname{\mathsf{RFN}}(n)$ is a Π_2^1 sentence provable from $\Pi_1^1\operatorname{\mathsf{-CA}}_0$, the theories $\operatorname{\mathsf{ACA}}_0 + \{\beta_0^1 \operatorname{\mathsf{RFN}}(n) : n \in \omega\}$ and $\{\sigma \in \Pi_2^1 : \Pi_1^1\operatorname{\mathsf{-CA}}_0 \vdash \sigma\}$ prove the same sentences.

Corollary 3.10. For any Π_2^1 sentence σ provable from Π_1^1 -CA₀, there exists exactly one $n \in \omega$ such that $n = \min\{m : \mathsf{ACA}_0 + \beta_0^1 \mathsf{RFN}(m) \vdash \sigma\}$.

The previous corollary means that Π_2^1 sentences provable from Π_1^1 -CA₀ is classified into the β_0^1 RFN hierarchy. In the following sections, we see the relationship between the β_0^1 RFN-hierarchy and some concrete Π_2^1 theorems provable from Π_1^1 -CA₀.

In the remaining of this section, we see some basic properties of $\beta_0^1 \operatorname{RFN}(n)$.

We note that $\beta_0^1 \operatorname{RFN}(0)$ is the same as the ω -model reflection of ACA₀. Thus, $\beta_0^1 \operatorname{RFN}(0)$ is equivalent to ACA₀⁺ over RCA₀. Therefore, $\beta_0^1 \operatorname{RFN}$ hierarchy is above ACA₀⁺. Moreover, it is easy to prove that even $\beta_0^1 \operatorname{RFN}(1)$ is stronger than ATR₀.

Lemma 3.11 ([4], Theorem VII.2.7.). Over ACA_0 , any coded β -model satisfies ATR_0 .

Theorem 3.12. Over ACA₀, $\beta_0^1 \operatorname{RFN}(1)$ implies the ω -model reflection of ATR₀.

Proof. We reason in $ACA_0 + \beta_0^1 \operatorname{RFN}(1)$. Let X be a set. By $\beta_0^1 \operatorname{RFN}(1)$, take coded ω -models \mathcal{M}_0 and \mathcal{M}_1 such that

$$X \in \mathcal{M}_0 \in_{\beta} \mathcal{M}_1 \land \mathcal{M}_1 \models \mathsf{ACA}_0.$$

Then, $\mathcal{M}_1 \models [\mathcal{M}_0 \models \mathsf{ATR}_0]$. Thus $\mathcal{M}_0 \models \mathsf{ATR}_0$ actually holds.

In the following section, we will show that there is a very large gap between ATR_0 and $\beta_0^1 RFN(1)$.

4 Relativized leftmost path principle and $\beta_0^1 RFN$

As we have noted in introduction, Towsner introduced a family of Π_2^1 statements called relative leftmost path principles to give an upper bound of some Π_2^1 theorems provable from Π_1^1 -CA₀ [8]. Indeed, he proved that Kruskal's theorem for trees, Nash-Williams' theorem in bqo theory and Menger's theorem for countable graphs are provable from one of the relative leftmost path principles.

In this section, we compare the relative leftmost path principles and the β_0^1 RFN-hierarchy. More precisely, we introduce a variant of relative leftmost

path principle which we call arithmetical relative leftmost path principle (ALPP) and show that ALPP is equivalent to ALPP, and we show that the strongest form of relative leftmost path principle, the transfinite relativized leftmost path principle TLPP, is weaker than $\beta_0^1 \text{RFN}(2)$. Moreover, we introduce the *n*-th iteration of ALPP denoted It^{*n*}(ALPP) which is equivalent to $\beta_0^1 \text{RFN}(n)$.

To define ALPP, we introduce the notion of arithmetical reducibility.

Definition 4.1 (Arithmetical reducibility). Let X, Y be sets. We say X is arithmetically reducible to Y (write $X \leq_{\mathrm{T}}^{\mathrm{a}} Y$) if

$$\exists n \exists Z (Z = Y^{(n)} \land X \leq_{\mathrm{T}} Z).$$

Let $\varphi(X)$ be a formula. We write $\forall X \leq_{\mathrm{T}}^{\mathrm{a}} Y \varphi(X)$ to mean $\forall X(X \leq_{\mathrm{T}}^{\mathrm{a}} Y \to \varphi(X))$ and write $\exists X \leq_{\mathrm{T}}^{\mathrm{a}} Y \varphi(X)$ to mean $\exists X(X \leq_{\mathrm{T}}^{\mathrm{a}} Y \wedge \varphi(X))$.

Remark 4.2. The notation $\leq_{\mathrm{T}}^{\mathrm{a}}$ follows from the notation $\leq_{\mathrm{W}}^{\mathrm{a}}$ of arithmetical Weihrauch reducibility.

Remark 4.3. We note that arithmetical reducibility is an extension of the notion of arithmetical definability. For $A, B \subset \omega$, we say A is arithmetically definable from B if there is an arithmetical formula $\theta(x, X)$ with exactly displayed free variables such that $A = \{x : \theta(x, B)\}$. Thus, A is arithmetically definable from B is equivalent to the condition that for any ω -model \mathcal{M} of ACA₀ containing A and $B, \mathcal{M} \models A \leq_{\mathrm{T}}^{\mathrm{a}} B$.

In contrast to this remark, for a non ω -model \mathcal{M} of ACA₀ and $A, B \in \mathcal{M}$, $\mathcal{M} \models A \leq_{\mathrm{T}}^{\mathrm{a}} B$ does not mean that there is an arithmetical formula θ such that $\mathcal{M} \models A = \{x : \theta(x, B)\}$. However, the connection between arithmetical reducibility and arithmetical definability rather holds in the following sense.

Lemma 4.4. (ACA₀) Let \mathcal{M} be a coded ω -model of ACA₀. Then, \mathcal{M} is closed under arithmetical reduction. Hence, for any arithmetical formula θ and $Y \in \mathcal{M}$,

$$\exists X \leq_{\mathrm{T}}^{\mathrm{a}} Y\theta(X) \to [\mathcal{M} \models \exists X \leq_{\mathrm{T}}^{\mathrm{a}} Y\theta(X)] \text{ and} \\ \mathcal{M} \models \forall X \leq_{\mathrm{T}}^{\mathrm{a}} Y\theta(X)] \to \forall X \leq_{\mathrm{T}}^{\mathrm{a}} Y\theta(X).$$

Proof. Let \mathcal{M} be an ω -model of ACA₀. Then, \mathcal{M} satisfies Σ_1^1 induction and $\forall Y \exists Z(Z = Y')$. Thus $\mathcal{M} \models \forall Y \forall n \exists Z(Z = Y^{(n)})$.

To show that \mathcal{M} is closed under arithmetical reduction, take $Y \in \mathcal{M}$ and X such that $X \leq_{\mathrm{T}}^{\mathrm{a}} Y$. Then, there exist n, Z such that $Z = Y^{(n)} \wedge X \leq_{\mathrm{T}} Z$. We note that $Z \in \mathcal{M}$ because the condition $Z = Y^{(n)}$ is absolute for \mathcal{M} . Thus, $X \in \mathcal{M}$.

The latter part immediately follows from the former part.

Lemma 4.5. Let $\varphi(X)$ be an arithmetical formula. Then, over ACA_0^+ , $\forall X \leq_{\mathrm{T}}^{\mathrm{a}} Y \varphi(X)$ is Δ_1^1 . Similarly, $\exists X \leq_{\mathrm{T}}^{\mathrm{a}} Y \varphi(X)$ is Δ_1^1 .

Proof. Within ACA_0^+ , $\forall X \leq_{\mathrm{T}}^{\mathrm{a}} Y \varphi(X)$ can be written as follows.

$$\forall X \leq^{\mathbf{a}}_{\mathrm{T}} Y\varphi(X) \leftrightarrow \forall Z(Z = Y^{(\omega)} \to \forall n \forall X \leq_{\mathrm{T}} Z_n\varphi(X)) \\ \leftrightarrow \exists Z(Z = Y^{(\omega)} \land \forall n \forall X \leq_{\mathrm{T}} Z_n\varphi(X)).$$

Therefore, $\forall X \leq_{\mathrm{T}}^{\mathrm{a}} Y \varphi(X)$ is Δ_1^1 . The case for $\exists X \leq_{\mathrm{T}}^{\mathrm{a}} Y \varphi(X)$ is the same.

Lemma 4.6. Assume ACA_0^+ . Then, for any X, the coded model $\mathcal{M} = \{Y : Y \leq_T^a X\}$ exists. Moreover, this is the smallest coded ω -model of ACA_0 containing X.

Definition 4.7 (Relative leftmost path principle, See [8]). Let $k \in \omega$. We define $\Delta_k^0 \text{LPP}$ as the assertion that for any ill-founded tree T, there is a path $g \in [T]$ such that

$$\forall f \in \Delta_k^0(T \oplus g) (f \in [T] \to g \leq_l f).$$

We define ALPP as the assertion that for any ill-founded tree T, there is a path $g \in [T]$ such that

$$\forall f \leq^{\mathbf{a}}_{\mathbf{T}} (T \oplus g) (f \in [T] \to g \leq_{l} f).$$

We call a witness g of ALPP an arithmetical leftmost path.

Finally, over ATR_0 , define TLPP as the following assertion: For any illfounded tree T and any well-order α , there is a path $g \in [T]$ such that

$$\forall f \leq_{\mathrm{T}} (T \oplus g)^{(\alpha)} (f \in [T] \to g \leq_l f).$$

We call such $g \neq \Delta^0_{\alpha+1}$ leftmost path.

Remark 4.8. In the definition of $\Delta_k^0 \text{LPP}$, ALPP and TLPP, we can replace $(T \oplus g)$ by g if we work in RCA₀.

From now on, we see the relationship between $\beta_0^1 \operatorname{RFN}(1)$, $\beta_0^1 \operatorname{RFN}(2)$ and ALPP, TLPP. For this comparison, we prefer the base theory to be ACA₀⁺ rather than ACA₀. However, the following lemma take the difference of the choice of the base theory away.

Lemma 4.9. Over ACA_0 , $\Delta_0^0 LPP$ implies ATR_0 .

Proof. See [8], Theorem 4.2.

Our strategy to prove $\beta_0^1 \operatorname{RFN}(1)$ from ALPP is essentially the same as for the proof of Theorem 2.13; to make a hyperjump, it is sufficient to make the set $\{i : [T_i] \neq \emptyset\}$ from given sequence of trees $\langle T_i \rangle_i$.

Lemma 4.10. There are a Σ_1^0 formula $\varphi(n, X)$ and a Π_1^0 formula $\psi(n, X)$ such that

- Over RCA_0 , $\forall n, X(\varphi(n, X) \leftrightarrow \psi(n, X))$.
- Over RCA_0 , for any $X, T = \{n : \varphi(n, X)\}$ is a tree such that $1^\infty \in [T]$.
- Over ACA_0 , for any X, if $\{n : \varphi(n, X)\}$ has a leftmost path, then HJ(X) is computable from it. Moreover, this reduction is uniform.

We may assume that there is a Turing functional $\Phi_T(X)$ defined by $\Phi_T(X) = \{n : \varphi(n, X)\}.$

Proof. Recall that there is a Σ_0^0 formula θ such that over ACA₀, HJ(X) = $\{(e, x) : \exists f \forall z \theta(e, x, X, f[z])\}$ for any X.

We give a construction of $\Phi_T(X)$ over RCA₀. For given X, define a sequence $\langle T_{e,x} \rangle_{e,x}$ of trees by $\sigma \in T_{e,x} \leftrightarrow \theta(e,x,X,\sigma)$. Then put $S_{e,x} = \{\langle 0 \rangle * \sigma : \sigma \in T_{e,x}\} \cup \{\sigma : \forall n < |\sigma|\sigma(n) = 1\}$. Finally, put $\Phi_T(X) = \bigoplus_{e,x} S_{e,x}$ where \bigoplus is the operator defined in Definition 2.9.

We show that this construction satisfies desired conditions. It is easy to see that $\Phi_T(X)$ is uniformly computable from X over RCA_0 and its has an infinite path 1^∞ . Thus, it is enough to show that $\mathrm{HJ}(X)$ is uniformly computable from the leftmost path of $\Phi_T(X)$ over ACA_0 .

We reason in ACA₀. Let f be the leftmost path of $\Phi_T(X)$. Then, for each e and x, f((0, (e, x))) = 0 iff $T_{e,x}$ has a path. Thus, $\{(e, x) : f((0, (e, x))) = 0\} = \{(e, x) : \exists f \forall z \theta(e, x, X, f[z], z)\} = \text{HJ}(X)$.

We note that we can modify the second condition as follows :

• Over RCA_0 , for any X, $\Phi_T(X) = \{n : \varphi(n, X)\}$ is an ill-founded tree which has a path uniformly computable from X, and any path computes X.

For this condition, consider the following transformation.

Definition 4.11. Let $\sigma \in \mathbb{N}^{<\mathbb{N}}$. Put

$$\sigma_{\text{even}} = \langle \sigma(0), \sigma(2), \dots, \sigma(l_e) \rangle,$$

$$\sigma_{\text{odd}} = \langle \sigma(1), \sigma(3), \dots, \sigma(l_o) \rangle,$$

where l_e is the maximum even number such that $l_e < |\sigma|$ and l_o is the maximum odd number such that $l_o < |\sigma|$. Similarly, for a function $f \in \mathbb{N}^{\mathbb{N}}$, define $f_{\text{even}} = \langle f(2n) \rangle_{n \in \mathbb{N}}$ and $f_{\text{odd}} = \langle f(2n+1) \rangle_{n \in \mathbb{N}}$.

Let X be a set and T be a tree. We identify X and its characteristic function. Thus, X is regarded as an infinite binary sequence. Define a tree $S_{T,X}$ by

$$\sigma \in S_{T,X} \leftrightarrow \sigma_{\text{even}} \in T \land \sigma_{\text{odd}} \prec X.$$

Lemma 4.12. Let X be a set and T be a tree. Then, over RCA_0 ,

- $S_{T,X}$ is uniformly computable from X and T.
- For any path f of $S_{T,X}$, f_{even} is a path of T and $f_{odd} = X$. Conversely, for $f \in [T]$ and X, the sequence $f \uplus X = \langle f(0), X(0), f(1), X(1), \ldots \rangle$ is a path of $S_{T,X}$.
- If f and g are paths of T such that $f \leq_l g$, then $f \uplus X \leq_l g \uplus X$. Especially, if f is the leftmost path of T, then the corresponding path $f \uplus X$ is the leftmost path of $S_{T,X}$.

Proof. This is immediate from the definition of $S_{T,X}$.

Therefore, by replacing $\Phi_T(X)$ with $S_{\Phi_T(X),X}$, we have

Lemma 4.13. There is a Turing functional $\Phi_T(\bullet)$ defined in RCA₀ such that

- Over RCA_0 , for any X, $\Phi_T(X)$ is an ill-founded tree which has a path uniformly computable from X, and any path computes X.
- Over ACA_0 , for any X, if $\Phi_T(X)$ has a leftmost path, then HJ(X) is computable from that path. Moreover, this reduction is uniform.

Theorem 4.14. Over ACA₀, ALPP is equivalent to $\beta_0^1 \operatorname{RFN}(1)$.

Proof. First, we show that ALPP implies $\beta_0^1 \operatorname{RFN}(1)$. Since $\Delta_0^0 \operatorname{LPP}$ implies ATR₀, it is enough to show that ACA₀⁺ +ALPP proves $\beta_0^1 \operatorname{RFN}(1)$.

Let X be a set. By ALPP, take an arithmetical leftmost path g of $\Phi_T(X)$. Let $\mathcal{M} = \{Y : Y \leq_T^a g\}$. Then $X, \Phi_T(X) \in \mathcal{M}, \mathcal{M} \models \mathsf{ACA}_0$ and g is the leftmost^{\mathcal{M}} path of $\Phi_T(X)$. Thus, \mathcal{M} satisfies $\mathrm{HJ}(X)$ exists.

We next show that $\beta_0^1 \operatorname{\mathsf{RFN}}(1)$ implies ALPP. Let T be an ill-founded tree. By $\beta_0^1 \operatorname{\mathsf{RFN}}(1)$, take an ω -model \mathcal{M} such that $T \in \mathcal{M}$ and $\mathcal{M} \models \operatorname{\mathsf{ACA}}_0 + \exists Y(Y = \operatorname{HJ}(T))$.

Now, \mathcal{M} satisfies [T has a leftmost path] because its hyperjump exists. Take the leftmost^{\mathcal{M}} path g of T. We claim that g is an arithmetical leftmost path of T. Let $f \leq_{\mathrm{T}}^{\mathrm{a}} g$ and $f \in [T]$. Since $g \in \mathcal{M}$ and $\mathcal{M} \models \mathsf{ACA}_0, f \in \mathcal{M}$. Since g is the leftmost^{\mathcal{M}} path, $g \leq_l f$.

It is shown in [8] that ALPP < TLPP. More precisely, a general form of relative leftmost path principle $\Sigma^0_{\alpha} LPP$ for a well order α is introduced, and it is proved that $\Sigma^0_{\alpha+5} LPP$ proves the consistency of $\Sigma^0_{\alpha+1} LPP$. Since ALPP is provable from $\Sigma^0_{\omega+1} LPP$, the consistency of ALPP is provable from TLPP. Thus we have

Theorem 4.15. $\beta_0^1 \text{RFN}(1) < \text{TLPP}.$

On the other hand, TLPP is strictly weaker than $\beta_0^1 \operatorname{RFN}(2)$. To prove this result, we introduce a variant of $\beta_0^1 \operatorname{RFN}(n)$.

Definition 4.16. Let σ be a Π_2^1 formula. We define $\beta_0^1 \operatorname{RFN}(n; \sigma)$ as the following assertion:

 $\forall X \exists \mathcal{M}_0, \dots, \mathcal{M}_n : \text{coded } \omega \text{-models } (X \in \mathcal{M}_0 \in_\beta \mathcal{M}_1 \dots \in_\beta \mathcal{M}_n \land \mathcal{M}_n \models \mathsf{ACA}_0 + \sigma).$

Lemma 4.17. Over ATR_0 , any β -model satisfies TLPP.

Proof. Let \mathcal{M} be a β -model. Let $T, \alpha \in \mathcal{M}$ such that

 $\mathcal{M} \models T \text{ is an ill-founded tree} \land$ $\alpha \text{ is a well order.}$

We show that \mathcal{M} has a Δ^0_{α} leftmost path.

Let $S = \{\sigma \in T : \exists f \in \mathcal{M}(\sigma \prec f)\}$. Since \mathcal{M} is a β -model, $S = \{\sigma \in T : \exists f(\sigma \prec f)\}$. Now, S is a nonempty pruned tree and hence its leftmost path g is computable from S. Moreover, g is the leftmost path of T.

Now we have $\exists g \forall f \in [T] (g \leq_l f)$ and hence

$$\exists g \forall f \leq_{\mathrm{T}} g^{(\alpha)} (f \in [T] \to g \leq_{l} f).$$

The above condition is Σ_1^1 . Therefore,

$$\mathcal{M} \models \exists g \forall f \leq_{\mathrm{T}} g^{(\alpha)} (f \in [T] \to g \leq_{l} f).$$

This completes the proof.

Theorem 4.18. (ACA₀) $\beta_0^1 \operatorname{RFN}(1; \operatorname{ATR}_0)$ proves the ω -model reflection of TLPP. Therefore, TLPP $< \beta_0^1 \operatorname{RFN}(1; \operatorname{ATR}_0)$.

Proof. Assume $ACA_0 + \beta_0^1 \operatorname{RFN}(1; ATR_0)$. Let X be a set. Take coded ω -models $\mathcal{M}_0, \mathcal{M}_1$ such that

$$X \in \mathcal{M}_0 \in_{\beta} \mathcal{M}_1 \land \mathcal{M}_1 \models \mathsf{ATR}_0.$$

Then, since \mathcal{M}_0 is a coded β -submodel of \mathcal{M}_1 , $\mathcal{M}_0 \models \mathsf{TLPP}$. This completes the proof.

Theorem 4.19. $\beta_0^1 \operatorname{\mathsf{RFN}}(2)$ proves the ω -model reflection of $\beta_0^1 \operatorname{\mathsf{RFN}}(1; \operatorname{\mathsf{ATR}}_0)$. Therefore, $\beta_0^1 \operatorname{\mathsf{RFN}}(1; \operatorname{\mathsf{ATR}}_0) < \beta_0^1 \operatorname{\mathsf{RFN}}(2)$.

Proof. Let X be a set. By $\beta_0^1 \operatorname{RFN}(2)$, take coded ω -models $\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}_2$ such that

 $X \in \mathcal{M}_0 \in_{\beta} \mathcal{M}_1 \in_{\beta} \mathcal{M}_2 \land \mathcal{M}_2 \models \mathsf{ACA}_0.$

Then $\mathcal{M}_2 \models [\mathcal{M}_1 \models \mathsf{ATR}_0]$. Thus, for any $Y \in \mathcal{M}_0$,

$$\mathcal{M}_2 \models \exists \mathcal{N}_0, \mathcal{N}_1 [Y \in \mathcal{N}_0 \in_\beta \mathcal{N}_1 \land \mathcal{N}_1 \models \mathsf{ATR}_0].$$

Since \mathcal{M}_0 is a β -submodel of \mathcal{M}_2 , we have

$$\mathcal{M}_0 \models \forall Y \exists \mathcal{N}_0, \mathcal{N}_1 [Y \in \mathcal{N}_0 \in_\beta \mathcal{N}_1 \land \mathcal{N}_1 \models \mathsf{ATR}_0].$$

Hence, $X \in \mathcal{M}_0$ and $\mathcal{M}_0 \models \beta_0^1 \operatorname{\mathsf{RFN}}(1; \operatorname{\mathsf{ATR}}_0)$. This completes the proof. \Box

Next, let us consider iterated versions of ALPP.

Definition 4.20. Let $n \in \omega$. We define $\Omega(n+1)$ as the following assertion: for any X there are f_0, \ldots, f_n such that

$$f_0 \in [\Phi_T(X)] \land \bigwedge_{i < n} f_{i+1} \in [\Phi_T(f_i)] \land$$
$$\forall g \leq^{\mathbf{a}}_{\mathbf{T}} f_n[(g \in [\Phi_T(X)] \to f_0 <_l g) \land \bigwedge_{i < n} (g \in [\Phi_T(f_i)] \to f_{i+1} <_l g)].$$

We also define $\mathsf{lt}^{n+1}(\mathsf{ALPP})$ as follows. Define $\Theta(T, f, g)$ be the following formula:

$$\Theta(T, f, g) \equiv T$$
 is a tree and $g \in [T]$ and $T, g \leq^{a}_{T} f$.

For each m < n, define ψ_m^n by

$$\psi_{0}^{n}(T_{0},\dots,T_{n},f_{0},\dots,f_{n}) \equiv \forall g \leq_{\mathrm{T}}^{\mathrm{a}} f_{0} \oplus \dots \oplus f_{n} \bigwedge_{i \leq n} (g \in [T_{i}] \to f_{i} \leq_{l} g)$$

$$\psi_{m+1}^{n} \begin{pmatrix} T_{0},\dots,T_{n-m-1}, \\ f_{0},\dots,f_{n-m-1} \end{pmatrix} \equiv \forall T_{n-m}, g_{n-m} \begin{pmatrix} \Theta(T_{n-m},g_{n-m},f_{n-m-1}) \\ \to \exists f_{n-m} \in [T_{n-m}] \psi_{m}^{n}(T_{0},\dots,T_{n-m},f_{0},\dots,f_{n-m}) \end{pmatrix}.$$

Then, define $\mathsf{It}^{n+1}(\mathsf{ALPP})$ as follows.

$$\mathsf{It}^{n+1}(\mathsf{ALPP}) \equiv \forall T_0 : \text{ ill-founded tree } \exists f_0 \in [T_0]\psi_n^n(T_0, f_0)$$

For example, $lt^1(ALPP)$ is equivalent to ALPP and $lt^2(ALPP)$ states that for any ill-founded tree T_0 , there exists a path $f_0 \in [T_0]$ such that

$$\forall T_1, g_1(\Theta(T_1, g_1, f_0) \to \exists f_1 \in [T_1] \forall g \leq^{\mathbf{a}}_{\mathbf{T}} f_0 \oplus f_1(\bigwedge_{i \leq 1} g \in [T_i] \to f_i \leq_l g)).$$

For simplicity, we write $(T,g) \in \Theta(f)$ instead of $\Theta(T, f, g)$. Then, $\mathsf{lt}^{n+1}(\mathsf{ALPP})$ states that for any ill-founded tree T_0 , there exists a path $f_0 \in [T_0]$ such that

$$\forall (T_1, g_1) \in \Theta(f_0) \exists f_1 \in [T_1] \cdots \forall (T_n, g_n) \in \Theta(f_{n-1}) \exists f_n \in [T_n] \\ \forall g \leq_{\mathrm{T}}^{\mathrm{a}} f_0 \oplus \cdots \oplus f_n(\bigwedge_{i \leq n} g \in [T_i] \to f_i \leq_l g).$$

Theorem 4.21. For $n \in \omega$, the following assertions are equivalent over ACA_0^+ .

- 1. $\beta_0^1 \operatorname{RFN}(n+1)$,
- 2. $\Omega(n+1)$,
- 3. $It^{n+1}(ALPP)$.

Proof. $(1 \rightarrow 3)$ Let T_0 be an ill-founded tree and h be a path of T_0 . By $\beta_0^1 \operatorname{RFN}(n+1)$ take a sequence of ω -models such that

$$T_0, h \in \mathcal{M}_0 \in_{\beta} \cdots \in_{\beta} \mathcal{M}_{n+1} \land \mathcal{M}_{n+1} \models \mathsf{ACA}_0.$$

Now \mathcal{M}_1 satisfies " T_0 has a leftmost path". Take the leftmost \mathcal{M}_1 path f_0 . Then, \mathcal{M}_{n+1} also satisfies " f_0 is the leftmost path of T_0 ". Let $T_1, g_1 \leq_{\mathrm{T}}^{\mathrm{a}} f_0$ such that T_1 is a tree and $g_1 \in [T_1]$. Then, \mathcal{M}_2 satisfies " T_1 has a leftmost path". Let f_1 be the leftmost \mathcal{M}_2 path. Then, \mathcal{M}_{n+1} satisfies " f_1 is the leftmost path of T_1 ". Continuing this argument, we have that

$$\forall (T_1, g_1) \in \Theta(f_0) \exists f_1 \in [T_1] \cdots \forall (T_n, g_n) \in \Theta(f_{n-1}) \exists f_n \in [T_n]$$
$$\bigwedge_{i \leq n} (\mathcal{M}_{n+1} \models "f_i \text{ is the leftmost path of } T_i").$$

Since \mathcal{M}_{n+1} is an ω -model of ACA₀, it is closed under arithmetical reduction. This completes the proof.

 $(3 \rightarrow 2)$ This is trivial.

 $(2 \to 1)$ Let X be a set. We show that there is an ω -model \mathcal{M} such that $\mathcal{M} \models \mathsf{ACA}_0 + \exists Y(Y = \mathrm{HJ}^{n+1}(X))$. Let f_0, \ldots, f_n be witnesses of $\Omega(n)$ for X. Then, $f_0 \leq_{\mathrm{T}} \cdots \leq_{\mathrm{T}} f_n$. By ACA_0^+ , take the smallest ω -model \mathcal{M} of ACA_0 containing f_n . Then, \mathcal{M} satisfies " f_0 is the leftmost path of $\Phi_T(X)$ " and hence f_0 computes $\mathrm{HJ}(X)$ in \mathcal{M} . Similarly, f_i is the leftmost path of $\Phi_T(f_{i-1})$ in \mathcal{M} and hence it computes $\mathrm{HJ}(f_i)$ in \mathcal{M} . Thus, $\mathcal{M} \models \exists Y(Y = \mathrm{HJ}^{n+1}(X))$.

As shown in [5], $\Delta_n^0 \text{LPPs}$ and variants of β -model reflection called $\Delta_n^0 \beta$ model reflections are deeply related. In the following, we see the relationship of variants of leftmost path principles and variants of β -models reflection.

Remark 4.22. We can show that

- for any $n \in \omega$, ACA₀ proves that any β -model satisfies $\Delta_n^0 \text{LPP}$,
- ACA'_0 proves that for any $n \in \mathbb{N}$, any β -model satisfies $\Delta^0_n \mathsf{LPP}$,
- ACA_0^+ proves that any β -model satisfies ALPP.

by the same proof of Lemma 4.17.

In contrast to this result, even ACA_0^+ does not prove that any β -model satisfies TLPP. To prove this, we can use the same proof as in the following lemma.

Lemma 4.23. TLPP is equivalent to

$$\forall \alpha : WO \ \forall X \exists \mathcal{M}(\alpha, X \in \mathcal{M}, \\ \mathcal{M} \models \mathsf{ACA}_0 + \exists Y(Y = \mathrm{HJ}(X))), \\ \mathcal{M} \text{ is closed under } \alpha\text{-jump}).$$

Proof. First, assume TLPP. Take a well order α . Then, $\alpha \cdot \omega$ is also a wellordering ³. We note that $\alpha \cdot \omega$ is closed under $+\alpha$ in the following sense: Recall that |L| denotes the field $\{i : (i,i) \in L\}$ of L for a linear order L. For each $(n,i) \in |\alpha \cdot \omega|$, there is a certain embedding from α to the interval [(n,i), (n+2,0)] defined by f(j) = (n+1,j). Since ATR₀ holds, there is an embedding g from α to an initial segment of [(n,i), (n+2,0)]. Then, $\sup g$ is regarded as $(n,i) + \alpha$.

Take an arbitrary X. We show that

$$\exists \mathcal{M}(\alpha, X \in \mathcal{M}, \\ \mathcal{M} \models \mathsf{ACA}_0 + \exists Y(Y = \mathrm{HJ}(X))), \\ \mathcal{M} \text{ is closed under } \alpha\text{-jump}).$$

Recall that there is a total Turing functional Φ_T which outputs an ill-founded tree for any input (see Lemma 4.13). By TLPP, take a $\Delta^0_{\alpha\cdot\omega}$ -leftmost path f of $\Phi_T(\alpha \oplus X)$. Define

$$\mathcal{M} = \{ A : A \leq_{\mathrm{T}} f^{(n,i)} \text{ for some } (n,i) \in |\alpha \cdot \omega| \}.$$

Since $\mathcal{M} \models [f \text{ is a leftmost path of } \Phi_T(\alpha \oplus X)], \ \mathcal{M} \models \exists Y(Y = HJ(X)).$ Moreover, by construction, $\mathcal{M} \models \mathsf{ACA}_0$ and closed under α -jump.

The converse is trivial because the leftmost path in \mathcal{M} is an Δ^0_{α} -leftmost path in the ground model.

Recall that a coded ω -model of ACA₀ satisfies ACA₀⁺ if it is closed under taking the ω -jump of a set. Therefore, by the same proof as above, we have

Lemma 4.24. Over ACA₀, $\Delta^0_{\omega^2}$ LPP proves $\beta^1_0 \operatorname{RFN}(1; \operatorname{ACA}^+_0)$.

Theorem 4.25. Over ACA₀, TLPP is strictly stronger than $\beta_0^1 \operatorname{RFN}(1; \operatorname{ACA}_0^+)$.

Proof. It is immediate from the fact that TLPP is strictly stronger than $\Delta^0_{\omega^2}$ LPP.

Corollary 4.26. ACA_0^+ does not prove that any β -model satisfies TLPP.

Proof. We show that if ACA_0^+ proves that any β -model satisfies TLPP, then $ACA_0 + \beta_0^1 \operatorname{RFN}(1; ACA_0^+)$ proves the consistency of TLPP. Since this yields TLPP proves the consistency of TLPP, ACA_0^+ does not prove that any β -model satisfies TLPP

³Here, $\alpha \cdot \omega$ denotes the order type of lexicographical ordering of $\mathbb{N} \times \alpha$.

Assume ACA_0^+ proves that any β -model satisfies TLPP. We work in $ACA_0 + \beta_0^1 \operatorname{RFN}(1; ACA_0^+)$. By $\beta_0^1 \operatorname{RFN}(1; ACA_0^+)$, take coded ω -models $\mathcal{M}_0, \mathcal{M}_1$ such that $\mathcal{M}_0 \in_\beta \mathcal{M}_1$ and $\mathcal{M}_1 \models ACA_0^+$. Then, $\mathcal{M}_1 \models [\mathcal{M}_0 \models \mathsf{TLPP}]$ and hence $\mathcal{M}_1 \models \mathsf{Con}(\mathsf{TLPP})$. Since $\mathsf{Con}(\mathsf{TLPP})$ is arithmetical, $\mathsf{Con}(\mathsf{TLPP})$ holds.

Theorem 4.27. Over ACA₀, $\beta_0^1 \operatorname{RFN}(n; \operatorname{ALPP})$ implies the ω -model reflection of $\beta_0^1 \operatorname{RFN}(n+1)$.

Proof. We reason in $ACA_0 + \beta_0^1 RFN(n; ALPP)$. Take an arbitrary set A. By $\beta_0^1 RFN(n; ALPP)$, take coded ω -models $\mathcal{M}_0, \ldots, \mathcal{M}_n$ such that

$$A \in \mathcal{M}_0 \in_{\beta} \cdots \in_{\beta} \mathcal{M}_{n-1} \in_{\beta} \mathcal{M}_n \land \mathcal{M}_n \models \mathsf{ACA}_0 + \mathsf{ALPP}.$$

Since ALPP and $\beta_0^1 \operatorname{RFN}(1)$ is equivalent over ACA₀, $\mathcal{M}_n \models \beta_0^1 \operatorname{RFN}(1)$.

We show that $\mathcal{M}_0 \models \beta_0^1 \operatorname{\mathsf{RFN}}(n+1)$. By $\beta_0^1 \operatorname{\mathsf{RFN}}(1)$ in \mathcal{M}_n , there are coded ω -model $\mathcal{N}_0, \mathcal{N}_1 \in \mathcal{M}_n$ such that

$$\mathcal{M}_n \models [\mathcal{M}_{n-1} \in \mathcal{N}_0 \in_\beta \mathcal{N}_1 \land \mathcal{N}_1 \models \mathsf{ACA}_0].$$

Since \mathcal{M}_{n-1} is a β -submodel of \mathcal{M}_n and \mathcal{N}_0 is an ω -submodel of \mathcal{M}_n , for any Π^1_1 sentence σ with parameters from \mathcal{M}_{n-1} ,

$$[\mathcal{M}_{n-1} \models \sigma] \to [\mathcal{M}_n \models \sigma] \\ \to [\mathcal{N}_0 \models \sigma].$$

Thus, \mathcal{M}_{n-1} is a β -submodel of \mathcal{N}_0 . Therefore, for any $X \in \mathcal{M}_0$,

$$\mathcal{M}_n \models [X \in \mathcal{M}_0 \in_{\beta} \dots \in_{\beta} \mathcal{M}_{n-1} \in_{\beta} \mathcal{N}_0 \in_{\beta} \mathcal{N}_1 \land \mathcal{N}_1 \models \mathsf{ACA}_0]$$

Therefore, for any $X \in \mathcal{M}_0$,

$$\mathcal{M}_n \models \exists \mathcal{H}_0, \dots, \mathcal{H}_{n+1} [X \in \mathcal{H}_0 \in_\beta \dots \in_\beta \mathcal{H}_{n-1} \in_\beta \mathcal{H}_n \in_\beta \mathcal{H}_{n+1} \land \mathcal{H}_{n+1} \models \mathsf{ACA}_0]$$

Since \mathcal{M}_0 is a β -submodel of \mathcal{M}_n , for any $X \in \mathcal{M}_0$,

$$\mathcal{M}_0 \models \exists \mathcal{H}_0, \dots, \mathcal{H}_{n+1} [X \in \mathcal{H}_0 \in_\beta \dots \in_\beta \mathcal{H}_{n-1} \in_\beta \mathcal{H}_n \in_\beta \mathcal{H}_{n+1} \land \mathcal{H}_{n+1} \models \mathsf{ACA}_0]$$

Thus, $\mathcal{M}_0 \models \beta_0^1 \mathsf{RFN}(n+1).$

Theorem 4.28. Over ACA₀, $\beta_0^1 \operatorname{RFN}(n+1)$ proves the ω -model reflection of $\beta_0^1 \operatorname{RFN}(n; \forall k. \Delta_k^0 \operatorname{LPP})$. Here, $\forall k. \Delta_k^0 \operatorname{LPP}$ is the assertion that

$$\forall k \forall T : \text{ill-founded tree } \exists g \in [T] \forall h \leq_{\mathrm{T}} g^{(k)} (h \in [T] \to g \leq_{l} h).$$

Proof. We reason in $ACA_0 + \beta_0^1 \operatorname{RFN}(n+1)$. Take an arbitrary set A. By $\beta_0^1 \operatorname{RFN}(n+1)$, take coded ω -models $\mathcal{M}_0, \ldots, \mathcal{M}_{n+1}$ such that

$$A \in \mathcal{M}_0 \in_{\beta} \cdots \in_{\beta} \mathcal{M}_n \in_{\beta} \mathcal{M}_{n+1} \land \mathcal{M}_{n+1} \models \mathsf{ACA}_0.$$

We show that $\mathcal{M}_0 \models \beta_0^1 \operatorname{\mathsf{RFN}}(n; \forall k. \Delta_k^0 \operatorname{\mathsf{LPP}})$. Since $\mathcal{M}_{n+1} \models \operatorname{\mathsf{ACA}}_0', \mathcal{M}_{n+1} \models [\mathcal{M}_n \models \forall k. \Delta_k^0 \operatorname{\mathsf{LPP}}]$. Therefore, for any $X \in \mathcal{M}_0$,

$$\mathcal{M}_{n+1} \models [X \in \mathcal{M}_0 \in_\beta \dots \in_\beta \mathcal{M}_n \land \mathcal{M}_n \models \forall k. \Delta_k^0 LPP].$$

Hence, for any $X \in \mathcal{M}_0$,

$$\mathcal{M}_{n+1} \models \exists \mathcal{H}_0 \cdots \mathcal{H}_n [X \in \mathcal{H}_0 \in_\beta \cdots \in_\beta \mathcal{H}_n \land \mathcal{H}_n \models \forall k. \Delta_k^0 \, \mathsf{LPP}].$$

Since \mathcal{M}_0 is a β -submodel of \mathcal{M}_{n+1} , for any $X \in \mathcal{M}_0$,

$$\mathcal{M}_0 \models \exists \mathcal{H}_0 \cdots \mathcal{H}_n [X \in \mathcal{H}_0 \in_\beta \cdots \in_\beta \mathcal{H}_n \land \mathcal{H}_n \models \forall k. \Delta_k^0 \mathsf{LPP}].$$

Therefore, $\mathcal{M}_0 \models \beta_0^1 \operatorname{\mathsf{RFN}}(n, \forall k. \Delta_k^0 \operatorname{\mathsf{LPP}}).$

Recall that any coded β -model satisfies ALPP over ACA₀⁺. Thus, by the same proof as above, we have

Theorem 4.29. Over ACA₀, $\beta_0^1 \operatorname{RFN}(n+1; \operatorname{ACA}_0^+)$ proves the ω -model reflection of $\beta_0^1 \operatorname{RFN}(n; \operatorname{ALPP})$.

Corollary 4.30. Over ACA_0 , the following holds.

 $\beta_0^1\operatorname{\mathsf{RFN}}(n;\forall k.\Delta_k^0\operatorname{\mathsf{LPP}}) < \beta_0^1\operatorname{\mathsf{RFN}}(n+1) < \beta_0^1\operatorname{\mathsf{RFN}}(n;\operatorname{\mathsf{ALPP}}) < \beta_0^1\operatorname{\mathsf{RFN}}(n+1;\operatorname{\mathsf{ACA}}_0^+).$

In the remaining of this section, we give another characterization of ALPP.

Definition 4.31. Let \mathcal{M} be a coded ω -model of ACA₀. We say \mathcal{M} is an A β -models if for any Π_2^0 formula $\theta(X)$,

$$\exists X \leq^{\mathbf{a}}_{\mathrm{T}} \mathcal{M}\theta(X) \leftrightarrow \mathcal{M} \models \exists X\theta(X).$$

Lemma 4.32. Let $\varphi(X)$ be an arithmetical formula with exactly displayed free variable. Then, ACA₀ proves that for any A β -model \mathcal{M} ,

$$\exists X \leq^{\mathbf{a}}_{\mathrm{T}} \mathcal{M}\theta(X) \leftrightarrow \mathcal{M} \models \exists X\theta(X).$$

Proof. Let $\varphi(X)$ be an arithmetical formula. Then there exists a number k and a Π_2^0 formula $\theta(X, f)$ such that ACA₀ proves

$$\forall X(\varphi(X) \leftrightarrow \exists f \leq_{\mathrm{T}} X^{(k)} \theta(X, f)).$$

We work in ACA₀. Assume \mathcal{M} is an A β -model and $\exists X \leq_{\mathrm{T}}^{\mathrm{a}} \mathcal{M}\varphi(X)$. Since φ is arithmetical, $\varphi(X)$ actually holds. Thus, there exists $f \leq_{\mathrm{T}} X^{(k)}$ such that $\theta(X, f)$. Since \mathcal{M} is a model of ACA₀, $\mathcal{M} \models \exists f \leq_{\mathrm{T}} X^{(k)}\theta(X, f)$. Hence $\mathcal{M} \models \varphi(X)$.

Definition 4.33. Define the $A\beta$ -model reflection as the assertion that for any set X there is an $A\beta$ -model containing X.

Theorem 4.34. Over ACA₀, β_0^1 RFN(1) is equivalent to A β -model reflection.

Proof. First, assume $\beta_0^1 \operatorname{\mathsf{RFN}}(1)$. Let X be a set. By $\beta_0^1 \operatorname{\mathsf{RFN}}(1)$, take a coded ω -models $\mathcal{M}_0, \mathcal{M}_1$ such that $X \in \mathcal{M}_0 \in_\beta \mathcal{M}_1 \land \mathcal{M}_1 \models \operatorname{\mathsf{ACA}}_0$.

We claim that \mathcal{M}_0 is an A β -model. Let $\theta(Y)$ be an arithmetical formula. Assume $\exists Z \leq^{\mathbf{a}}_{\mathrm{T}} \mathcal{M}_0 \theta(Z)$. Then, there exists $n \in \mathbb{N}$ such that $\exists Z \leq_{\mathrm{T}} (\mathcal{M}_0)^{(n)} \theta(Z)$. Since $\mathcal{M}_0 \in \mathcal{M}_1$ and $\mathcal{M}_1 \models \mathsf{ACA}'_0$, $\mathcal{M}_1 \models \exists Z \theta(Z)$. Since \mathcal{M}_0 is a β -submodel of \mathcal{M}_1 , $\mathcal{M}_0 \models \exists Z \theta(Z)$.

We next assume $A\beta$ -model reflection. Then, especially ACA_0^+ holds. Let X be a set. By $A\beta$ -model reflection, take an $A\beta$ -model \mathcal{M}_0 such that $X \in \mathcal{M}_0$. By ACA_0^+ , take $\mathcal{M}_1 = \{Y : Y \leq_{\mathrm{T}}^{\mathrm{a}} \mathcal{M}_0\}$.

Now $X \in \mathcal{M}_0 \in \mathcal{M}_1 \land \mathcal{M}_1 \models \mathsf{ACA}_0$. Thus, it is enough to show that \mathcal{M}_0 is a β -submodel of \mathcal{M}_1 . Let $\theta(Y)$ be an arithmetical formula. Assume $\mathcal{M}_1 \models \exists Y \theta(Y)$. Then, $\exists Y \leq^{\mathbf{a}}_{\mathrm{T}} \mathcal{M}_0 \theta(Y)$ and hence $\mathcal{M}_0 \models \exists Y \theta(Y)$. This completes the proof.

Corollary 4.35. (ACA₀.) ALPP is equivalent to $A\beta$ -model reflection.

Proof. Since ALPP is equivalent to $\beta_0^1 \operatorname{RFN}(1)$ and $\beta_0^1 \operatorname{RFN}(1)$ is equivalent to A β -model reflection, ALPP is equivalent to A β -model reflection.

For each n > 0, define $A\beta \operatorname{RFN}(n)$ as the assertion that

 $\forall X \exists \mathcal{M}_0 \cdots \mathcal{M}_{n-1} (X \in_{\beta} \cdots \in_{\beta} \mathcal{M}_{n-1} \land \mathcal{M}_{n-1} \text{ is an } \mathsf{A}\beta \text{-model}).$

Then, by the same argument as in Theorem 4.34, we have

Theorem 4.36. Over ACA₀, $\beta_0^1 \operatorname{RFN}(n)$ is equivalent to $A\beta \operatorname{RFN}(n-1)$ for any n > 0.

5 The arithmetical relativization

As we have noted, the $\beta_0^1 \text{RFN-hierarchy}$ reveals that how many times the hyperjump operator is used for prove a Π_2^1 sentence. In addition, we can classify Π_3^1 sentences provable from $\Pi_1^1\text{-}\mathsf{CA}_0$ by comparing their Π_2^1 approximations and the $\beta_0^1 \text{RFN-hierarchy}$. In this section, we summarize the idea of Π_2^1 approximations of Π_3^1 sentences.

Definition 5.1. Let $\sigma \equiv \forall X(\theta(X) \to \exists Y \forall Z \eta(X, Y, Z))$ be a Π_3^1 sentence. Define rel (σ) by $\forall X(\theta(X) \to \exists Y \forall Z \leq_{\mathrm{T}}^{\mathrm{a}} (X \oplus Y)\eta(X, Y, Z))$. We call rel (σ) the arithmetical relativization of σ .

For example, ALPP and A β RFN(n) are instances of rel(σ). Indeed, if we take $\theta(X)$ to be 'X is an ill-founded tree' and $\eta(X, Y, Z)$ to be ($Y \in [X]$) $\wedge (Z \in [X] \rightarrow Y \leq_l X$), then ALPP is equivalent to rel($\forall X(\theta(X) \rightarrow \exists Y \forall Z \eta(X, Y, Z))$).

Remark 5.2. In the case of ALPP, the following $\operatorname{rel}'(\sigma)$ is equivalent to $\operatorname{rel}(\sigma)$.

 $\operatorname{rel}'(\sigma) \equiv \forall X(\theta(X) \to \exists Y \forall Z \leq^{\mathrm{a}}_{\mathrm{T}} Y \eta(X, Y, Z)).$

In general, $rel(\sigma)$ and $rel'(\sigma)$ are not equivalent. The equivalence of $rel(\sigma)$ and $rel'(\sigma)$ is related to the notion of *cylinder* in Weihrauch reduction.

We note that if σ is a Π_3^1 sentence provable from Π_1^1 -CA₀, then rel(σ) is a Π_2^1 sentence provable from Π_1^1 -CA₀. Thus, rel(σ) is provable from ACA₀ + $\beta_0^1 \operatorname{RFN}(n)$ for some n. Let us write n_{σ} for the smallest n such that ACA₀ + $\beta_0^1 \operatorname{RFN}(n) \vdash \operatorname{rel}(\sigma)$ for such a σ . Then, we can classify Π_3^1 sentences provable from Π_1^1 -CA₀ according to n_{σ} .

We then give a lemma which is useful to evaluate n_{σ} .

Definition 5.3. Let $\varphi \equiv \forall X \exists Y \theta$ and $\psi \equiv \forall X \exists Y \eta$. For a theory *T*, we say *T* proves the lightface implication of φ to ψ if

$$T \vdash \forall X (\exists Y \theta \to \exists Z \eta).$$

Lemma 5.4. Let $\varphi \equiv \forall X \exists Y \forall Z \theta(X, Y, Z)$ and $\psi \equiv \forall X \exists Y \forall Z \eta(X, Y, Z)$ be Π_3^1 sentences such that ACA₀ proves the lightface implication of $\varphi \to \psi$. Then, ACA₀⁺ proves rel(φ) \to rel(ψ).

Proof. We reason in $\mathsf{ACA}_0^+ + \forall X \exists Y \forall Z \leq_{\mathrm{T}}^a X \oplus Y \theta(X, Y, Z)$. Let X, Y be such that $\forall Z \leq_{\mathrm{T}}^a X \oplus Y \theta(X, Y, Z)$. Take the smallest ω -model \mathcal{M} of ACA_0 containing $X \oplus Y$. Then, $\mathcal{M} \models \exists U \forall Z \theta(X, U, Z)$ via U = Y. Since \mathcal{M} is a model of ACA_0 , $\mathcal{M} \models \exists V \forall Z \eta(X, V, Z)$. Take such a V. Thus, $\exists W \forall Z \leq_{\mathrm{T}}^a X \oplus W \eta(X, W, Z)$ via W = V. For example, if ACA_0 proves $\forall X (\exists Y \forall Z \theta(X, Y, Z) \to \exists W(W = \mathrm{HJ}(X)))$, then ACA_0^+ proves $\operatorname{rel}(\forall X \exists Y \forall Z \theta) \to \beta_0^1 \mathsf{RFN}(1)$. Conversely, if ACA_0 proves $\forall X (\exists W(W = \mathrm{HJ}(X)) \to \exists Y \forall Z \theta(X, Y, Z))$, then ACA_0^+ (and hence even RCA_0) proves $\beta_0^1 \mathsf{RFN}(1) \to \operatorname{rel}(\forall X \exists Y \forall Z \theta)$.

6 Pseudo Ramsey's theorem and β RFN

In this section, we introduce Π_2^1 variants of Ramsey's theorem for $[\mathbb{N}]^{\mathbb{N}}$ which may be seen as an instance of the arithmetical relativization of usual Ramsey's theorem. Then we see the relationship between them and the β_0^1 RFNhierarchy. Here, $[X]^{\mathbb{N}}$ denotes the class of all infinite subsets of X. Topological Ramsey's theorem claims that certain subclasses of $[\mathbb{N}]^{\mathbb{N}}$ has Ramsey property in the sense that a class $\mathcal{A} \subseteq [\mathbb{N}]^{\mathbb{N}}$ has Ramsey's property if there is an infinite set $H \subseteq \mathbb{N}$ such that $[H]^{\mathbb{N}}$ is a subset of \mathcal{A} or its complement.

We first recall how to formalize Ramsey's theorem in second-order arithmetic and some existing results. For the details, see also [4].

Definition 6.1. Let $[\mathbb{N}]^{\mathbb{N}}$ denote the class of all strictly increasing sequence of natural numbers. Let $\varphi(f)$ be a formula for $f \in [\mathbb{N}]^{\mathbb{N}}$. We say φ has Ramsey's property if

$$\exists h \in [\mathbb{N}]^{\mathbb{N}} (\forall g \in [\mathbb{N}]^{\mathbb{N}} \varphi(h \circ g) \lor \forall g \in [\mathbb{N}]^{\mathbb{N}} \neg \varphi(h \circ g)).$$

In this definition, we identify $f \in [\mathbb{N}]^{\mathbb{N}}$ and its range, $\varphi(f)$ and the class $\{f \in [\mathbb{N}]^{\mathbb{N}} : \varphi(f)\}$. Especially, $h \in [\mathbb{N}]^{\mathbb{N}}$ corresponds to an infinite subset H of \mathbb{N} , $h \circ g$ corresponds to an infinite subset of H.

Definition 6.2. Let Γ be a class of formulas. Then, Γ -Ramsey's theorem (write Γ -Ram) is the assertion that any $\varphi(f) \in \Gamma$ has Ramsey's property.

Theorem 6.3. [4, VI.6.4] Over RCA_0 , Π_1^1 - CA_0 is equivalent to Σ_0^1 - Ram .

We give a finer analysis for this theorem by using the $\beta_0^1 \text{RFN}$ -hierarchy. We start with giving Π_2^1 variants of Ramsey's property which we call pseudo-Ramsey's property.

Definition 6.4. Let $\varphi(f, X)$ be a formula with exactly displayed free variables for $f \in [\mathbb{N}]^{\mathbb{N}}$. We say $\varphi(f)$ has pseudo-Ramsey's property if

$$\forall X \exists f (\forall g \leq^{\mathbf{a}}_{\mathsf{T}} f \oplus X\varphi(f \circ g) \lor \forall g \leq^{\mathbf{a}}_{\mathsf{T}} f \oplus X \neg \varphi(f \circ g)).$$

Define rel(Γ -Ram) as the assertion that any formula $\varphi \in \Gamma$ has pseudo-Ramsey's property. We note that each $\sigma \in \operatorname{rel}(\Gamma\operatorname{-Ram})$ is of the form $\operatorname{rel}([\varphi \text{ has Ramsey's property}])$ for some $\varphi \in \Gamma$ if $\Gamma \subseteq \Sigma_0^1$. Moreover, for each n > 0, there exists a formula $\varphi \in \Sigma_n^0$ such that $\operatorname{rel}(\Sigma_n^0\operatorname{-Ram})$ is axiomatizable by $\operatorname{rel}([\varphi \text{ has Ramsey's property}])$. Since $\Sigma_n^0\operatorname{-Ram}$ is provable from $\Pi_1^1\operatorname{-CA}_0$, $\operatorname{rel}(\Sigma_n^0\operatorname{-Ram})$ is provable from $\beta_0^1\operatorname{RFN}(p(n))$ for some p(n). We show that p(n) = n. It follows from the following lemma.

Lemma 6.5. ([4], VI.6.2.) (ACA₀) Let $\mathcal{M}_0, \ldots, \mathcal{M}_{k-1}$ be coded β models such that $\mathcal{M}_0 \in \cdots \in \mathcal{M}_{k-1}$. Then, for any Σ_k^0 formula $\varphi(f)$ with parameters from $\mathcal{M}_0, \varphi(f)$ has Ramsey's property via some $h \in \mathcal{M}_{k-1}$.

Theorem 6.6. rel($\Sigma_k^0 \text{Ram}$) is provable from $\beta_0^1 \text{RFN}(k)$.

Proof. Assume $\beta_0^1 \operatorname{\mathsf{RFN}}(k)$. Let $\varphi(f, X)$ be a Σ_k^0 formula for $f \in [\mathbb{N}]^{\mathbb{N}}$ with exactly displayed set variables. By $\beta_0^1 \operatorname{\mathsf{RFN}}(k)$, take coded ω -models $\mathcal{M}_0, \ldots, \mathcal{M}_k$ such that $X \in \mathcal{M}_0 \in_{\beta} \cdots \in_{\beta} \mathcal{M}_{k-1} \in_{\beta} \mathcal{M}_k$ and $\mathcal{M}_k \models \operatorname{\mathsf{ACA}}_0$. Now, by the previous lemma, there exists $f \in \mathcal{M}_{k-1}$ such that

$$\mathcal{M}_k \models \forall g \varphi(f \circ g) \lor \forall g \neg \varphi(f \circ g).$$

Since \mathcal{M}_k is a model of ACA₀, φ has pseudo-Ramsey's property via f. \Box

In the remaining of this section, we show that $\mathsf{ACA}_0^+ + \{\operatorname{rel}(\Sigma_n^0\mathsf{Ram}) : n \in \omega\}$ proves $\beta_0^1\mathsf{RFN}(n)$ for all n. In [6], it is shown that over ACA_0 , (lightface- $\Delta_2^{0,X}$) $\mathsf{Ram} \to \exists Y(Y = \mathrm{HJ}(X))$. Here lightface- $\Delta_2^{0,X}$ is the set of Δ_2^0 formulas having no set parameters other than X. Moreover, it is commented without proofs that this result can be extended to the transfinite level: $\mathsf{ATR}_0 \vdash (\mathsf{lightface}-\Delta_1^1\mathsf{Ram}) \to \forall \alpha$: recursive well order $\exists Y(Y = \mathrm{HJ}^{\alpha}(\emptyset))$). In this section, we give an explicit proof for finite level.

Definition 6.7. Let T be a tree. We say $f \in [\mathbb{N}]^{\mathbb{N}}$ majorizes T if $\exists g \in [T] \forall n(g(n) \leq f(n))$.

Definition 6.8. Let f be a function and $n \in \mathbb{N}$. Define f_{+n} by $f_{+n}(x) = f(x+n)$.

Lemma 6.9. (ACA₀) Let T be a tree and $f \in [\mathbb{N}]^{\mathbb{N}}$. Then, f majorizes T iff $\forall n > 0 \exists \tau \in T(|\tau| = n \land \forall m < n(\tau(m) \le f(m)))$. Especially, f majorizes T is an arithmetical condition.

Proof. See the proof of Lemma V.9.6. in [4]. \Box

Lemma 6.10. (Essentially [4] VI.6.1) Let $\theta(n, A)$ be the Σ_1^1 formula defining HJ(A) over ACA₀. Write $\theta(n, A) \equiv \exists f \in \mathbb{N}^{\mathbb{N}} \forall y R(n, f[y], A[y])$ by a Σ_0^0 formula R.

Assume ACA₀. For each $n \in \mathbb{N}$ and $A \subseteq \mathbb{N}$, define a tree $T(n, A) = \{\sigma : \forall y R(n, \sigma[y], A[y])\}$. For each $f \in [\mathbb{N}]^{\mathbb{N}}$ and $A \subseteq \mathbb{N}$, define $F(f, A) = \{n : \exists p(f_{+p} \text{ majorizes } T(n, A)).$

Define $\psi(f, A)$ by

 $\forall n < f(0)(f_{+2} \text{ majorizes } T(n, A) \rightarrow f_{+1} \text{ majorizes } T(n, A)).$

Then, if $h \in [\mathbb{N}]^{\mathbb{N}}$ satisfies $\forall g(\psi(h \circ g, A))$, then F(h, A) = HJ(A).

Proof. We first remark that for each n and A, T(n, A) has a path iff $n \in$ HJ(A).

Take $h \in [\mathbb{N}]^{\mathbb{N}}$ such that $\forall g(\psi(h \circ g, A))$. We show that $F(h, A) \subseteq \mathrm{HJ}(A)$. If $n \in F(h, A)$ then f_{+p} majorizes T(n, A) for some p, and hence $[T(n, A)] \neq \emptyset$. Thus, $n \in \mathrm{HJ}(A)$.

Conversely, assume $n \in HJ(A)$ and show that $n \in F(h, A)$. Now, T(n, A) has a path. We claim that $h_{+(n+2)}$ majorizes T(n, A). Let $g_0 \in [T(n, A)]$ and define $g(n) = g_0(0) + \cdots + g_0(n)$. Then $g \in [\mathbb{N}]^{\mathbb{N}}$ and it majorizes T(n, A). Assume $h_{+(n+2)}$ does not majorize T(n, A). Take k > 0 such that $\forall \tau \in T(n, A)(|\tau| = k \to \exists m < k(\tau(m) > h_{+(n+2)}(m))$. Define f by

$$f(x) = \begin{cases} h_{+(n+1)}(x) \text{ if } x < k+1 \\ h_{+(n+k+2)}(g(x)) \text{ otherwise.} \end{cases}$$

Claim 6.10.1. Now we have

- 1. $f_{+(k+1)}$ majorizes T(n, A) but
- 2. f_{+1} does not majorize T(n, A).

Proof of Claim. (1.) For any $x \in \mathbb{N}$, $f_{+(k+1)}(x) = f(x+k+1) = h_{+(n+1)}(g(x))$. Since $h \in [\mathbb{N}]^{\mathbb{N}}$, we have $h_{+(n+1)}(g(x)) \ge g(x) \ge g_0(x)$. Hence $g_0(x)$ ensures that $f_{+(k+1)}$ majorizes T(n, A).

(2.) For a contradiction, assume f_{+1} majorizes T(n, A). Then there exists $\tau_k \in T(n, A)$ such that

$$|\tau_k| = k \land \forall m < k(\tau_k(m) \le f_{+1}(m)).$$

Now, for any m < k,

$$f_{+1}(m) = f(m+1)$$

= $h_{+(n+1)}(m+1)$
= $h(m+n+2) = h_{+(n+2)}(m)$

Therefore, $\forall m < k(\tau_k(m) \leq h_{+(n+2)}(m))$. However, this contradicts the definition of k.

For each $k' \leq k$, define $g_{k'} \in [\mathbb{N}]^{\mathbb{N}}$ such that

$$g_{k'}(0) = 0,$$

$$g_{k'}(1) = 1,$$

$$g_{k'}(x+2) = \begin{cases} x+k'+n+2 \text{ if } x+k'+1 < k+1 \\ g(x+k'+1)+(n+k+2) \text{ otherwise} \end{cases}$$

Then, for any $k' \leq k, g_{k'} \in [\mathbb{N}]^{\mathbb{N}}$ and $f_{+(k'+1)} = (h \circ g_{k'})_{+2}$. Indeed, for any x,

$$f_{+(k'+1)}(x) = f(x+k'+1)$$

=
$$\begin{cases} h(x+k'+n+2) & \text{if } x+k'+1 < k+1 \\ h(g(x+k'+1)+(n+k+2)) & \text{otherwise} \end{cases}$$

and

$$\begin{aligned} (h \circ g_{k'})_{+2}(x) &= h(g_{k'}(x+2)) \\ &= \begin{cases} h(x+k'+n+2) \text{ if } x+k'+1 < k+1 \\ h(g(x+k'+1)+(n+k+2)) \text{ otherwise.} \end{cases} \end{aligned}$$

Especially, $(h \circ g_{k'})_{+1} = f_{+k'}$. Now we have

- 1. $f_{+(k+1)}$ majorizes T(n, A),
- 2. for any $k' \leq k$, if $f_{+(k'+1)}$ majorizes T(n, A) then $f_{+k'}$ majorizes T(n, A).

The second condition is proved as follows: now $n < n + 1 \le h(n + 1) = f(0) < f_{k'+1}(0) = (h \circ g_{k'})_{+2}(0)$. Therefore, by the assumption on h, if $f_{k'+1} = (h \circ g_{k'})_{+2}$ majorizes T(n, A), then $(h \circ g_{k'})_{+1} = f_{k'}$ also majorizes T(n, A).

By these conditions and arithmetical induction, we have f_{+1} majorizes T(n, A). However, this is a contradiction.

Definition 6.11. (ACA₀) For $f \in [\mathbb{N}]^{\mathbb{N}}$ and A, define $F^n(f, A)$ by

$$F^{0}(f, A) = A,$$

 $F^{n+1}(f, A) = F(f, F^{n}(f, A))$

Definition 6.12. Define an arithmetical formula $\Phi_n(f, A)$ by

$$\Phi_n(f,A) \equiv \bigwedge_{i < n} \psi(f,F^i(f,A)).$$

Lemma 6.13. Let $k \in \omega$. Then ACA_0 proves that for any $h \in [\mathbb{N}]^{\mathbb{N}}$ and A, if

$$(\forall g \Phi_k(h \circ g, A)) \lor (\forall g \neg \Phi_k(h \circ g, A))$$

then $(\forall g \Phi_k(h \circ g, A))$ holds.

Proof. Take h and A as above. For the sake of contradiction, assume $(\forall g \neg \Phi_k(h \circ g, A)).$

For each *n* define g_n by $g_n(0) = 0, g_n(x+1) = n + x + 1$. Then, since $\neg \Phi(h \circ g, A)$ holds, there exists m < h(g(0)) = h(0) such that

$$\bigvee_{i < k} (h \circ g_n)_{+2} \text{ majorizes } T(m, F^i((h \circ g_n)_{+2}, A)) \text{ but } (h \circ g_n)_{+1} \text{ does not.}$$

We note that $(h \circ g_n)_{+1}(x) = (h \circ g_n)(x+1) = h(g_n(x+1)) = h(n+x+1) = h_{+(n+1)}(x)$ and $(h \circ g_n)_{+2}(x) = h_{+(n+2)}(x)$. Hence, for each *n*, there exists m < h(0) such that

$$\bigvee_{i < k} h_{+(n+2)} \text{ majorizes } T(m, F^i(h_{+(n+2)}, A)) \text{ but } h_{+(n+1)} \text{ does not}$$

Claim 6.13.1. For each i < k and n, $F^i(h_{+(n+2)}, A) = F^i(h, A)$.

Proof of Claim. We will show by (meta-)induction on *i*.

The base step is immediate. By definition, $F^0(h_{+(n+2)}, A) = A = F^0(h, A)$.

For the induction step, assume $F^i(h_{+(n+2)}, A) = F^i(h, A)$. Then

$$\begin{aligned} F^{i+1}(h_{+(n+2)}, A) &= F(h_{+(n+2)}, F^{i}(h_{+(n+2)}, A)) \\ &= F(h_{+(n+2)}, F^{i}(h, A)) \\ &= \{x : \exists p((h_{+(n+2)})_{+p} \text{ majorizes } T(x, F^{i}(h, A)))\} \\ &= \{x : \exists p(h_{+p} \text{ majorizes } T(x, F^{i}(h, A)))\} = F(h, F^{i}(h, A)). \end{aligned}$$

This completes the proof.

Now, for each n, there exists m < h(0) such that

$$\bigvee_{i < k} h_{+(n+2)} \text{ majorizes } T(m, F^i(h, A)) \text{ but } h_{+(n+1)} \text{ does not.}$$

Put $\theta(i,n) \equiv h_{+(n+2)}$ majorizes $T(m, F^i(h, A))$ but $h_{+(n+1)}$ does not.

For each n, define m(n) be the smallest m < h(0) satisfying the above condition.

Claim 6.13.2. For each m, at most k-many n's such that m(n) = m.

Proof of Claim. Fix an m. For the sake of contradiction, assume let n_0, \ldots, n_k be such that $n_0 < \cdots < n_k$ and $m(n_j) = m$ for all j < k. Then, for each j < k, there exists an $i_j < k$ such that $\theta(i_j, n_j)$ holds.

We claim that if $j \neq j'$ then $i_j \neq i_{j'}$. Let $j < j' \leq k$. If $i_j = i_{j'}$, then

 $h_{+(n_j+2)}$ majorizes $T(m, F^{i_j}(h, A))$ but $h_{+(n_j+1)}$ does not \wedge

 $h_{+(n_{j^\prime}+2)}$ majorizes $T(m,F^{i_j}(h,A))$ but $h_{+(n_{j^\prime}+1)}$ does not .

However, since $n_j < n_{j'}$ and $h_{+(n_j+2)}$ majorizes $T(m, F^{i_j}(h, A))$, $h_{+(n_{j'}+1)}$ should majorize $T(m, F^{i_j}(h, A))$. This is a contradiction.

Now $\{i_j : j \le k\}$ should be a k+1-element subset of $\{0, \ldots, k-1\}$. This is a contradiction.

Now, m(n) is a total function such that $\{m(n) : n \in \mathbb{N}\}$ is bounded by h(0). However, for any $y \in \{m(n) : n \in \mathbb{N}\}$, there are at most k-many n such that m(n) = y. This is a contradiction.

Lemma 6.14. (ACA₀) For any $h \in [\mathbb{N}]^{\mathbb{N}}$ and A, if $\forall g \Phi_n(h \circ g, A)$ then $F^i(h, A) = \mathrm{HJ}^i(A)$ for any $i \leq n$.

Proof. Take h and A as above. Then, for the formula ψ in 6.10, $\forall g(\psi(h \circ g, A))$ holds. Hence F(h, A) = HJ(A). Thus, $\forall g(\psi(h \circ g, \text{HJ}(A)))$ also holds, and hence $F(h, F(h, A)) = F(h, \text{HJ}(A)) = \text{HJ}^2(A)$. Iterating this argument, we have $F^i(h, A) = \text{HJ}^i(A)$ for any $i \leq n$.

Corollary 6.15. Over ACA_0^+ , $\{rel(\Sigma_n^0 Ram) : n \in \omega\}$ and $\{\beta_0^1 RFN(n) : n \in \omega\}$ prove the same sentences.

Proof. As we have proved in Theorem 6.6, $\beta_0^1 \operatorname{RFN}(n)$ implies $\operatorname{rel}(\Sigma_n^0 \operatorname{Ram})$ for each n.

We show that $\beta_0^1 \operatorname{RFN}(n)$ is provable from $\{\operatorname{rel}(\Sigma_n^0 \operatorname{Ram}) : n \in \omega\}$. Assume $\operatorname{ACA}_0^+ + \{\operatorname{rel} \Sigma_n^0 \operatorname{Ram} : n \in \omega\}$. Take an arbitrary $n \in \omega$. Let X be a set.

We will show that there exists a coded ω -model \mathcal{M} such that $X \in \mathcal{M}$ and $\mathcal{M} \models \exists Y(Y = \mathrm{HJ}^n(X)).$

Now $\Phi_n(f, X)$ is an arithmetical formula for $f \in [\mathbb{N}]^{\mathbb{N}}$. Therefore, there exists an $h \in [\mathbb{N}]^{\mathbb{N}}$ such that

$$\forall g \leq^{\mathbf{a}}_{\mathbf{T}} h \oplus X(\Phi_n(h \circ g, X)) \lor \forall g \leq^{\mathbf{a}}_{\mathbf{T}} h \oplus X(\neg \Phi_n(h \circ g, X)).$$

By ACA_0^+ , take $\mathcal{M} = \{Y : Y \leq_{\mathrm{T}}^a h \oplus X\}$. Then $\mathcal{M} \models \mathsf{ACA}_0 + (\forall g \Phi_n(h \circ g, X)) \lor (\forall g \neg \Phi_n(h \circ g, X))$. Thus, by Lemma 6.13, $\mathcal{M} \models \mathsf{ACA}_0 + \forall g \Phi_n(h \circ g, X)$. Therefore, by Lemma 6.14, $\mathcal{M} \models \exists Y(Y = \mathrm{HJ}^n(X))$.

7 Pseudo determinacy and $\beta_0^1 \text{RFN}$

In this section, we introduce a Π_2^1 variant of determinacy which we call pseudo-determinacy. Then we consider the hierarchy of pseudo-determinacy for $(\Sigma_1^0)_k$ games. First of all, we introduce the notion of difference hierarchy and the determinacy.

Definition 7.1 (Difference hierarchy). Let $k \in \omega$. We define the class $(\Sigma_1^0)_k$ of formulas as follows:

- a formula is $(\Sigma_1^0)_1$ if it is Σ_1^0 ,
- a formula φ is $(\Sigma_1^0)_{k+1}$ if there are a Σ_1^0 formula ψ and a $(\Sigma_1^0)_k$ formula θ such that $\varphi \equiv \psi \wedge \neg \theta$.

We call the hierarchy formed by $\{(\Sigma_1^0)_k : k \in \omega\}$ the difference hierarchy of Σ_1^0 formulas.

Definition 7.2. We call a function $S : \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}$ a strategy⁴. Let S and S' be strategies. We say a function $f \in \mathbb{N}^{\mathbb{N}}$ is the play along (S, S') (write $f = S \otimes S'$) if

$$\forall n(f(2n) = S(f[2n]) \land f(2n+1) = S'(f[2n+1])).$$

Let $\varphi(\vec{X}, f)$ be a formula for $f \in \mathbb{N}^{\mathbb{N}}$. Let \vec{X} be sets. We say $\varphi(\vec{X}, \bullet)$ is determined if

$$\exists S_0 \forall S_1 \varphi(\vec{X}, S_0 \otimes S_1) \lor \exists S_1 \forall S_0 \neg \varphi(\vec{X}, S_0 \otimes S_1).$$

⁴Usually, a strategy is a function defined over $\bigcup_n \mathbb{N}^{2n}$ or $\bigcup_n \mathbb{N}^{2n+1}$.

Especially, if a strategy S_0 satisfies

$$\forall S_1 \varphi(\vec{X}, S_0 \otimes S_1),$$

then we say $\varphi(\vec{X}, \bullet)$ is determined via a strategy S_0 for player 0. If a strategy S_1 satisfies

$$\forall S_0 \neg \varphi(\vec{X}, S_0 \otimes S_1),$$

then we say $\varphi(\vec{X}, \bullet)$ is determined via a strategy S_1 for player 1.

We also say φ is determined if for any \vec{X} , $\varphi(\vec{X}, \bullet)$ is determined. Let Γ be a class of formulas for $f \in \mathbb{N}^{\mathbb{N}}$. We say Γ is determined (write Γ -Det) if any $\varphi \in \Gamma$ is determined.

It is well-known that over RCA_0 ,

- ATR_0 is equivalent to Σ_1^0 -Det, and
- Π_1^1 -CA₀ is equivalent to $(\Sigma_1^0)_k$ -Det for k > 1.

In this section, we consider Π_2^1 variants of the second clause.

Definition 7.3 (pseudo-determinacy). Let $\varphi(\vec{X}, f)$ be a formula for $f \in \mathbb{N}^{\mathbb{N}}$. For given sets \vec{X} , we say $\varphi(\vec{X}, \bullet)$ is pseudo-determined if

$$\exists S_0 \forall S_1 \leq^a_{\mathrm{T}} S_0 \oplus \vec{X} \varphi(\vec{X}, S_0 \otimes S_1) \lor \exists S_1 \forall S_0 \leq^a_{\mathrm{T}} S_1 \oplus \vec{X} \neg \varphi(\vec{X}, S_0 \otimes S_1)).$$

If $\varphi(\vec{X}, \bullet)$ is pseudo-determined via S, then we say S is a pseudo-winning strategy for the game defined by φ and \vec{X} . If \vec{X} is clear from the context, we just say S is a pseudo-winning strategy for the game defined by φ . If $\varphi(\vec{X}, \bullet)$ is pseudo-determined for any \vec{X} , then we say φ is pseudo-determined.

Let Γ be a class of formulas for $f \in \mathbb{N}^{\mathbb{N}}$. We say Γ is pseudo-determined if any $\varphi \in \Gamma$ is pseudo-determined. Let $\operatorname{rel}(\Gamma\operatorname{-Det})$ denote the assertion that every game defined by a formula in Γ is pseudo-determined.

We note that the statement $[\varphi \text{ is pseudo-determined}]$ is of the form $\operatorname{rel}(\psi)$ for some ψ . Moreover, each $\operatorname{rel}((\Sigma_1^0)_n\text{-}\mathsf{Det})$ is axiomatizable by a Π_2^1 sentence of the form $\operatorname{rel}(\psi)$.

Remark 7.4. For any $(\Sigma_1^0)_k$ formula $\varphi(X_1, \ldots, X_n, f)$, there is a $(\Sigma_1^0)_k$ formula $\psi(X, f)$ such that $\mathsf{RCA}_0 \vdash \forall X_1, \ldots, X_n, f(\varphi(X_1, \ldots, X_n, f) \leftrightarrow \psi(X_1 \oplus \cdots \oplus X_n, f))$. Therefore, when considering $\operatorname{rel}(\Sigma_1^0)_k$ -Det, we may assume that any $(\Sigma_1^0)_k$ formula has exactly one variable X and one function variable f.

Let $\varphi(X, f)$ be a $(\Sigma_1^0)_k$ formula. We may assume that for any X, all strategies for the game $\varphi(X, \bullet)$ computes X. Indeed, if S is a strategy for player 0, then S is enough to be defined for sequences with even length. Thus, we may assume that $X(n) = S(1^{2n+1})$. Similarly, if S is a strategy for player 1, then we may assume that $X(n) = S(1^{2n})$. Therefore, $\varphi(X, f)$ is pseudo-determined if

$$\forall X (\exists S_0 \forall S_1 \leq^a_{\mathrm{T}} S_0 \varphi(X, S_0 \otimes S_1) \lor \exists S_1 \forall S_0 \leq^a_{\mathrm{T}} S_1 \neg \varphi(X, S_0 \otimes S_1)).$$

We note that for any $k \in \omega, k > 1$, $\operatorname{rel}(\Sigma_1^0)_k$ -Det is a Π_2^1 statement provable from Π_1^1 -CA₀. Thus, each $\operatorname{rel}(\Sigma_1^0)_k$ -Det is provable from some $\beta_0^1 \operatorname{RFN}(n)$ over ACA₀. We show that $\beta_0^1 \operatorname{RFN}(k; \operatorname{ATR}_0)$ is enough to prove $\operatorname{rel}(\Sigma_1^0)_k$ -Det.

To prove $(\Sigma_1^0)_2$ determinacy from Π_1^1 -CA₀, Tanaka proved the following lemma.

Lemma 7.5. Let $\varphi(X, f)$ be a $(\Sigma_1^0)_2$ formula with exactly displayed free set variables. Assume ATR₀. For any X, if HJ(X) exists, then $\varphi(X, \bullet)$ is determined.

Proof. See [7].

We extend this lemma to $(\Sigma_1^0)_k$ formulas.

Lemma 7.6. Let $k \in \omega$ and $\varphi(X, f)$ be a $(\Sigma_1^0)_{k+1}$ formula with exactly displayed set variables. Over ATR_0 , for any set X, if $\mathrm{HJ}^k(X)$ exists, then $\varphi(X, \bullet)$ is determined.

Proof. We will show by induction on $k \in \omega$. The case k = 0 is trivial because ATR₀ proves Σ_1^0 determinacy.

Let $k \in \omega$. Assume that ATR_0 proves that for any X, if $\mathrm{HJ}^k(X)$ exists then any $(\Sigma_1^{0,X})_{k+1}$ game is determined.

We reason in ATR₀. Let X be a set such that $HJ^{k+1}(X)$ exists. Let $\varphi(A, f)$ be a $(\Sigma_1^0)_{k+2}$ formula. Then, there is a Σ_1^0 formula $\psi(A, f)$ and a $(\Sigma_1^0)_{k+1}$ formula $\theta(A, f)$ such that $\varphi(A, f) \equiv \psi(A, f) \wedge \neg \theta(A, f)$. Moreover, $\psi(A, f)$ is written as $\exists n\eta(A, f[n])$ for a Σ_0^0 formula η . Take a coded β model \mathcal{M} such that $X \in \mathcal{M} \models \exists Y(Y = HJ^k(X))$. Define W by

$$W = \{ \sigma \in \mathbb{N}^{<\mathbb{N}} : \eta(X, \sigma) \land \mathcal{M} \models \text{player } 0 \text{ wins for } \neg \theta^{\sigma}(X, \bullet) \}.$$

Here, $\neg \theta^{\sigma}(X, f)$ is the formula $\neg \theta(X, \sigma * f)$ and $\sigma * f$ denotes the concatenation of σ and f. Let $\varphi'(f) \equiv \exists n(f[n] \in W)$. Then, the game defined by φ' is determined.

Case 1: player 0 wins the game φ' via S_0 . Now, for any $\sigma \in W$, $\mathcal{M} \models \exists S_0^{\sigma} \forall S_1^{\sigma} \neg \theta(X, \sigma * S_0^{\sigma} \otimes S_1^{\sigma})$. Take a sequence $\langle S_0^{\sigma} \rangle_{\sigma \in W}$. Then, each S_0^{σ} satisfies $\forall S_1^{\sigma} \neg \theta(X, \sigma * S_0^{\sigma} \otimes S_1^{\sigma})$ because \mathcal{M} is a β -model. Consider a strategy for player 0 such that

- while the play is not in W, the strategy mimics S_0 ,
- once the play σ is in W, the strategy mimics S_0^{σ} .

Then this strategy is a winning strategy for player 0.

Case 2: player 1 wins the game φ' via S_1 . Now we have that for each play σ consistent with S_1 , if $\eta(X, \sigma)$ then

 $\mathcal{M} \models$ player 0 has no winning strategy for $\neg \theta^{\sigma}(X, \bullet)$.

By induction hypothesis, $\mathcal{M} \models \theta^{\sigma}(X, \bullet)$ is determined. Hence for each σ consistent with S_1 , if $\eta(X, \sigma)$ then $\mathcal{M} \models \exists S_1^{\sigma} \forall S_0^{\sigma} \theta(X, \sigma * S_0^{\sigma} \otimes S_1^{\sigma})$. Take a sequence $\langle S_1^{\sigma} : \sigma$ is consistent with $S_1 \land \eta(X, \sigma) \rangle$ such that $\mathcal{M} \models \forall S_0^{\sigma} \theta(X, \sigma * S_0^{\sigma} \otimes S_1^{\sigma})$. Then, since \mathcal{M} is a coded β -model, for each $S_0^{\sigma}, \forall S_1^{\sigma} \theta(X, \sigma * S_0^{\sigma} \otimes S_1^{\sigma})$ holds. Consider a strategy for player 1 such that

- while the play σ does not satisfy $\eta(X, \sigma)$, the strategy mimics S_1 ,
- once the play σ satisfies $\eta(X, \sigma)$, the strategy mimics S_1^{σ} .

Then this strategy is a winning strategy for player 1.

Theorem 7.7. For each $k \in \omega, k > 0, \beta_0^1 \operatorname{\mathsf{RFN}}(k, \operatorname{\mathsf{ATR}}_0)$ proves $\operatorname{rel}(\Sigma_1^0)_{k+1} \operatorname{\mathsf{Det}}$.

Proof. Let $\varphi(A, f)$ be a $(\Sigma_1^0)_{k+1}$ formula.

Assume $\beta_0^1 \operatorname{RFN}(k, \operatorname{ATR}_0)$. Let X be a set. By $\beta_0^1 \operatorname{RFN}(k, \operatorname{ATR}_0)$ take a coded ω -model \mathcal{M} such that $X \in \mathcal{M} \models \operatorname{ATR}_0 + \exists Y(Y = \operatorname{HJ}^k(X))$. Then, by the previous lemma, $\mathcal{M} \models \varphi(X, \bullet)$ is determined. Without loss of generality, we may assume that $\mathcal{M} \models$ player 0 wins for $\varphi(X, \bullet)$. Let $S_0 \in \mathcal{M}$ be a winning strategy of player 0. Then, for any $S_1 \leq_{\mathrm{T}}^{\mathrm{a}} S_0, S_1 \in \mathcal{M}$ and hence $\varphi(X, S_0 \otimes S_1)$.

We next see the implication from pseudo-determinacy to β_0^1 RFN. First we see the implication from rel $(\Sigma_1^0)_2$ -Det.

Lemma 7.8. There exists a $(\Sigma_1^0)_2$ formula $\varphi(f, X, Y)$ such that ACA₀ proves the following.

 $\forall \alpha : WO \forall X(\varphi(\bullet, X, \alpha) \text{ is determined} \rightarrow \exists Y(Y = (HJ(X))^{(\alpha)}).$

Proof. See [7].

Theorem 7.9. Over ACA_0^+ , $rel(\Sigma_1^0)_2$ -Det proves TLPP.

Proof. By 4.23, it is enough to show that

$$\forall \alpha : WO \,\forall X \exists \mathcal{M}(\alpha, X \in \mathcal{M}, \\ \mathcal{M} \models \mathsf{ACA}_0 + \exists Y(Y = \mathrm{HJ}(X)), \\ \mathcal{M} \text{ is closed under } \alpha\text{-jump}.$$

Let $\varphi(f, X, Y)$ be the formula in the Lemma 7.8. Take a well-ordering α and a set X. By rel $(\Sigma_1^0)_2$ -Det take a pseudo-winning strategy S for the game $\varphi(\bullet, X, \alpha \cdot \omega)$. Let \mathcal{M} be a coded ω -model of ACA₀ containing $X, \alpha \cdot \omega, S$. Then $\mathcal{M} \models \exists Y(Y = (\mathrm{HJ}(X))^{(\alpha \cdot \omega)})$.

We define a coded ω -model \mathcal{M}' by

$$\mathcal{M}' = \{ A \in \mathcal{M} : \mathcal{M} \models \exists n (A \leq_{\mathrm{T}} (\mathrm{HJ}(X))^{(\alpha \cdot n)}) \}.$$

Then, \mathcal{M}' is a coded ω -model of ACA₀ closed under α -jump. Moreover, since $\mathcal{M}' \in \mathcal{M}$ and $\mathcal{M} \models [\mathrm{HJ}(X) \in \mathcal{M}']$, $\mathcal{M}' \models \exists Y(Y = \mathrm{HJ}(X))$ by Lemma 2.27.

Corollary 7.10. $\beta_0^1 \operatorname{RFN}(1)$ is not enough to prove $\operatorname{rel}(\Sigma_1^0)_2$ -Det.

Proof. It follows from $\beta_0^1 \operatorname{RFN}(1) < \operatorname{TLPP} \le \operatorname{rel}(\Sigma_1^0)_2$ -Det.

Remark 7.11. In his paper, Tanaka pointed that ACA_0 is not strong enough to prove " Π_1^1 comprehension implies $(\Sigma_1^0)_2$ -Det" by using the axiomatic system lightface Π_1^1 comprehension. The previous corollary is another expression of this fact.

At last, we see the general case. The point is the following lemma.

Lemma 7.12. Let n > 0. Then there is a $(\Sigma_1^0)_{p(n)}$ formula $G_n(X, f)$ such that ACA₀ proves that

- $\forall X(G_n(X, \bullet) \text{ is determined} \to HJ^n(X) \text{ exists.})$
- Any winning strategy for G_n computes X.

Here, p(n) is a certain primitive recursive function.

Proof. See the proof of Proposition 6 in [3].

Theorem 7.13. Let n > 0. Then, over ACA_0^+ , $rel(\Sigma_1^0)_{p(n)}$ -Det implies $\beta_0^1 \operatorname{RFN}(n)$.

Proof. We reason in $\mathsf{ACA}_0^+ + \operatorname{rel}(\Sigma_1^0)_{p(n)}$ -Det. Let X be a set. By $\operatorname{rel}(\Sigma_1^0)_{p(n)}$, take a pseudo-winning strategy S for the game $G_n(X, \bullet)$. Let \mathcal{M} be an ω model of ACA_0 containing S. Then $\mathcal{M} \models [G_n(X, \bullet)$ is determined via S], and hence $\mathcal{M} \models \exists Y(Y = \operatorname{HJ}^n(X))$. This completes the proof. \Box

Corollary 7.14. rel $(\Sigma_1^0)_n$ -Det is just the Π_2^1 part of Π_1^1 -CA₀.

Remark 7.15. We have proved in Lemma 3.7, Theorem 7.7 and 7.9 that

- $\mathsf{TLPP} \leq \operatorname{rel}((\Sigma_1^0)_2 \operatorname{-Det}) \leq \beta_0^1 \mathsf{RFN}(1; \mathsf{ATR}_0),$
- TLPP $< \beta_0^1 \operatorname{RFN}(1; \operatorname{ATR}_0).$

Thus, at least one of TLPP or $\beta_0^1 \operatorname{RFN}(1; \operatorname{ATR}_0)$ is not equivalent to rel $((\Sigma_1^0)_2$ -Det).

Question 7.16. Can we separate rel($(\Sigma_1^0)_2$ -Det) and TLPP or $\beta_0^1 \operatorname{RFN}(1; \operatorname{ATR}_0)$?

References

- [1] Alberto Marcone. On the logical strength of nash-williams' theorem on transfinite sequences. In *Logic: from foundations to applications: European logic colloquium*, pages 327–351, 1996.
- [2] Antonio Montalbán and Richard A Shore. Conservativity of ultrafilters over subsystems of second order arithmetic. *The Journal of Symbolic Logic*, 83(2):740–765, 2018.
- [3] Leonardo Pacheco and Keita Yokoyama. Determinacy and reflection principles in second-order arithmetic. arXiv preprint arXiv:2209.04082, 2022.
- [4] Stephen G. Simpson. Subsystems of second order arithmetic. Perspectives in Logic. Cambridge University Press, Cambridge; Association for Symbolic Logic, Poughkeepsie, NY, second edition, 2009.
- [5] Yudai Suzuki and Keita Yokoyama. Searching problems above arithmetical transfinite recursion. arXiv preprint arXiv:2305.07321, 2023.
- [6] Kazuyuki Tanaka. The galvin-prikry theorem and set existen axioms. Annals of Pure and Applied Logic, 42(1):81–104, 1989.

- [7] Kazuyuki Tanaka. Weak axioms of determinacy and subsystems of analysis i: δ games. *Mathematical Logic Quarterly*, 36(6):481–491, 1990.
- [8] Henry Towsner. Partial impredicativity in reverse mathematics. J. Symbolic Logic, 78(2):459–488, 2013.