

EXT BRANCHING LAWS FOR THE GENERAL LINEAR GROUP

MOHAMMED SAAD QADRI

ABSTRACT. Let F be a non-archimedean local field. Let π_1 and π_2 be irreducible Arthur type representations of $\mathrm{GL}_n(F)$ and $\mathrm{GL}_{n-1}(F)$ respectively. We study Ext branching laws when π_1 and π_2 are products of discrete series representations and their Aubert-Zelevinsky duals. We obtain an Ext analogue of the local non-tempered Gan-Gross-Prasad conjecture in this case.

1. INTRODUCTION

Let F be a non-archimedean local field. Following the work [19] of Dipendra Prasad, there has been much interest in the study of Ext branching laws about the restriction of an irreducible representation of $\mathrm{GL}_n(F)$ to $\mathrm{GL}_{n-1}(F)$.

Let π_1 and π_2 be smooth irreducible representations of $\mathrm{GL}_n(F)$ and $\mathrm{GL}_{n-1}(F)$ respectively. It is a fundamental result due to Aizenbud-Gourevitch-Rallis-Schiffman [1] that the space $\mathrm{Hom}_{\mathrm{GL}_{n-1}(F)}(\pi_1, \pi_2)$ is at most one dimensional. It is also known that for all integers $i \geq 1$, the spaces $\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\pi_1, \pi_2)$ are finite dimensional (see [2], [19]). By the study of Ext branching laws we mean understanding when the spaces $\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\pi_1, \pi_2)$ are non-zero. If π_1 and π_2 are generic then it is a classical fact (see [18, Theorem 3], [13]) that $\mathrm{Hom}_{\mathrm{GL}_{n-1}(F)}(\pi_1, \pi_2) = \mathbb{C}$. For the corresponding Ext branching law, it was conjectured by Dipendra Prasad that when π_1 and π_2 are generic then the higher Ext spaces vanish, that is, $\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\pi_1, \pi_2) = 0$ for all integers $i \geq 1$. This has been recently proved by K.Y. Chan and G. Savin (see [9]).

The local Gan-Gross-Prasad conjectures [10] were formulated by Wee Teck Gan, Benedict Gross, and Dipendra Prasad giving rules for branching laws of certain pairs (G, H) of classical groups. These conjectures predict when a generic irreducible representation of H occurs in the quotient of a generic irreducible representation of G upon restriction. In a recent paper [11], they extended these conjectures beyond the generic case to the class of representations of *Arthur type*. For the p -adic general linear group, one direction of the local non-generic Gan-Gross-Prasad conjecture was proved by Maxim Gurevich [12] and the conjecture was completely settled by K.Y. Chan in the work [7]. In a recent work of K.Y. Chan (see [6]) the general smooth branching problem for the p -adic general linear group is resolved. It is now natural to consider the problem of formulating an Ext branching law, that is, classifying irreducible representations π_1 of $\mathrm{GL}_n(F)$ and π_2 of $\mathrm{GL}_{n-1}(F)$ such that $\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\pi_1, \pi_2) \neq 0$ for some integer $i \geq 0$. As pointed out earlier, the Ext branching for the general linear group is known when both the representations π_1 and π_2 are generic, in which case Ext branching reduces to just Hom branching.

In Remark 5.8 of the work [11], the authors mention that it would be an interesting question to give a precise condition predicting exactly when $\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\pi_1, \pi_2)$ is non-zero for some $i \geq 0$, when π_1 and π_2 are representations of Arthur type. This seems to be

a difficult problem in general. In this paper, we study Ext branching laws when π_1 and π_2 are products of discrete series representations and the Aubert-Zelevinsky duals of discrete series representations, as a first step towards answering this question. We obtain an Ext analogue of the local non-tempered Gan-Gross-Prasad conjecture in this case. Note that when either π_1 or π_2 is generic then some results on Ext branching law have been obtained in the work [7] of K.Y. Chan and this paper builds on the work [7].

We introduce the notion of *strong* Ext relevance for pairs of Arthur parameters. In order to motivate our definition we recall the original notion of relevance for Arthur type representations of $\mathrm{GL}_n(F)$. We first recall the definition of an Arthur parameter for $\mathrm{GL}_n(F)$. Let W_F denote the Weil group of the field F . An Arthur parameter of $\mathrm{GL}_n(F)$ is an admissible homomorphism,

$$\psi : W_F \times \mathrm{SL}_2(\mathbb{C}) \times \mathrm{SL}_2(\mathbb{C}) \rightarrow \mathrm{GL}_n(\mathbb{C})$$

such that the restriction of ψ to W_F has bounded image and the restriction to the two $\mathrm{SL}_2(\mathbb{C})$ factors are algebraic. The first $\mathrm{SL}_2(\mathbb{C})$ is called the Deligne SL_2 and the second $\mathrm{SL}_2(\mathbb{C})$ is called the Arthur SL_2 . We can obtain an L-parameter from an Arthur parameter by sending the Weil-Deligne group $W_F \times \mathrm{SL}_2(\mathbb{C})$ to $W_F \times \mathrm{SL}_2(\mathbb{C}) \times \mathrm{SL}_2(\mathbb{C})$ via the map,

$$\alpha : (w, g) \rightarrow \left(w, g, \begin{pmatrix} |w|^{1/2} & 0 \\ 0 & |w|^{-1/2} \end{pmatrix} \right)$$

where, $w \in W_F$ and $g \in \mathrm{SL}_2(\mathbb{C})$. Given an Arthur parameter ψ , we obtain the corresponding L-parameter $\psi \circ \alpha$. Thus by the local Langlands correspondence, we can associate an irreducible representation of $\mathrm{GL}_n(F)$ to any Arthur parameter, which turns out to be unitary. An irreducible representation of $\mathrm{GL}_n(F)$ obtained in this manner is called a representation of Arthur type.

Let $m, n \in \mathbb{Z}_{\geq 0}$ and let π_1 and π_2 be Arthur type representations of $\mathrm{GL}_m(F)$ and $\mathrm{GL}_n(F)$ respectively. Let $\mathcal{A}(\pi_1)$ and $\mathcal{A}(\pi_2)$ denote their respective Arthur parameters. Let V_d denote the unique irreducible representation of $\mathrm{SL}_2(\mathbb{C})$ of dimension d , that is, $V_d = \mathrm{Sym}^{d-1}(\mathbb{C}^2)$. In the work [11], the authors define a pair (π_1, π_2) to be relevant if there exist admissible homomorphisms $\{\phi_i\}_{i=1}^{i=r+s}$ of W_F with bounded image and positive integers $a_1, a_2, \dots, a_r, b_{r+1}, b_2, \dots, b_{r+s}, c_1, c_2, \dots, c_{r+s}$ such that,

$$\mathcal{A}(\pi_1) = \sum_{i=1}^r \phi_i \otimes V_{c_i} \otimes V_{a_i} \oplus \sum_{i=r+1}^{r+s} \phi_i \otimes V_{c_i} \otimes V_{b_{i-1}}$$

and,

$$\mathcal{A}(\pi_2) = \sum_{i=1}^r \phi_i \otimes V_{c_i} \otimes V_{a_{i-1}} \oplus \sum_{i=r+1}^{r+s} \phi_i \otimes V_{c_i} \otimes V_{b_i}.$$

It is now known by the work of K.Y. Chan (see [7, Theorem 1.3]) that the above relevance condition controls the branching law for representations of Arthur type.

Theorem 1.1. [7, Theorem 1.3] *Suppose that π_1 and π_2 are irreducible Arthur type representations of $\mathrm{GL}_n(F)$ and $\mathrm{GL}_{n-1}(F)$ respectively. Then $\mathrm{Hom}_{\mathrm{GL}_{n-1}(F)}(\pi_1, \pi_2) \neq 0$ if and only if the Arthur parameters of the representations π_1 and π_2 are GGP-relevant.*

It is known that the above relevance condition does not control the Ext branching law for Arthur type representations. This means that there exist Arthur type irreducible representations π_1 and π_2 of $\mathrm{GL}_n(F)$ and $\mathrm{GL}_{n-1}(F)$ respectively such that π_1 and π_2 are not relevant but $\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\pi_1, \pi_2) \neq 0$ for some $i \geq 0$.

For instance, let $\pi_1 = \mathbb{1}_3$ be the trivial representation of $\mathrm{GL}_3(F)$ and let $\pi_2 = \mathrm{St}_2$ be the Steinberg of $\mathrm{GL}_2(F)$. Then we have that

$$\mathrm{Ext}_{\mathrm{GL}_2(F)}^1(\mathbb{1}_3, \mathrm{St}_2) = \mathbb{C},$$

although in this case, $\mathbb{1}_3$ and St_2 do not have relevant Arthur parameters.

We now introduce the notion of *strong* Ext relevance which controls the Ext branching law in the cases we study in this paper.

Definition 1.1. Let $m, n \in \mathbb{Z}_{\geq 0}$ and let π_1 and π_2 be Arthur type representations of $\mathrm{GL}_m(F)$ and $\mathrm{GL}_n(F)$ respectively. Let $\mathcal{A}(\pi_1)$ and $\mathcal{A}(\pi_2)$ denote their respective Arthur parameters. We say that π_1 and π_2 are *strong* Ext relevant if there exist admissible homomorphisms $\{\phi_i\}_{i=1}^{i=r+s}$ and $\{\psi_i\}_{i=1}^{i=t+u}$ of W_F with bounded image and positive integers $a_1, a_2, \dots, a_r, b_{r+1}, b_{r+2}, \dots, b_{r+s}, c_1, c_2, \dots, c_{r+s}, d_1, d_2, \dots, d_t, e_{t+1}, e_{t+2}, \dots, e_{t+u}, f_1, f_2, \dots, f_{t+u}$ such that,

$$\mathcal{A}(\pi_1) = \sum_{i=1}^r \phi_i \otimes V_{c_i} \otimes V_{a_i} \oplus \sum_{i=r+1}^{r+s} \phi_i \otimes V_{c_i} \otimes V_{b_{i-1}} \oplus \sum_{i=1}^t \psi_i \otimes V_{f_i} \otimes V_{d_i} \oplus \sum_{i=t+1}^{t+u} \psi_i \otimes V_{e_{i-1}} \otimes V_{f_i}$$

and,

$$\mathcal{A}(\pi_2) = \sum_{i=1}^r \phi_i \otimes V_{c_i} \otimes V_{a_{i-1}} \oplus \sum_{i=r+1}^{r+s} \phi_i \otimes V_{c_i} \otimes V_{b_i} \oplus \sum_{i=1}^t \psi_i \otimes V_{d_{i-1}} \otimes V_{f_i} \oplus \sum_{i=t+1}^{t+u} \psi_i \otimes V_{f_i} \otimes V_{e_i}.$$

Remark 1.1. The definition stated above makes sense for arbitrary integers m and n . Later we shall specialise to the case $m = n + 1$.

Remark 1.2. A priori there are eight possible terms that might occur in the definition of strong Ext relevance where Ext might be non-zero based on the GGP relevance together with application of the Aubert-Zelevinsky involution. We enumerate these possibilities below where the first term is for $\mathrm{GL}_m(F)$ and the second term is for $\mathrm{GL}_n(F)$.

- (1) $\phi \otimes V_a \otimes V_b$ and $\phi \otimes V_a \otimes V_{b-1}$.
- (2) $\phi \otimes V_a \otimes V_{b-1}$ and $\phi \otimes V_a \otimes V_b$.
- (3) $\phi \otimes V_a \otimes V_b$ and $\phi \otimes V_{b-1} \otimes V_a$.
- (4) $\phi \otimes V_{b-1} \otimes V_a$ and $\phi \otimes V_a \otimes V_b$.
- (5) $\phi \otimes V_a \otimes V_b$ and $\phi \otimes V_b \otimes V_{a-1}$.
- (6) $\phi \otimes V_b \otimes V_{a-1}$ and $\phi \otimes V_a \otimes V_b$.
- (7) $\phi \otimes V_a \otimes V_b$ and $\phi \otimes V_{a-1} \otimes V_b$.
- (8) $\phi \otimes V_{a-1} \otimes V_b$ and $\phi \otimes V_a \otimes V_b$.

But only the sum of the first four is expected to give us a sufficient condition to obtain non-zero Ext modules. As an example consider the following. We know from the work of K.Y Chan and G. Savin [9] that the Steinberg St_n of $\mathrm{GL}_n(F)$ is projective upon restriction to $\mathrm{GL}_{n-1}(F)$ (and in fact isomorphic to the Gelfand-Graev representation upon restriction to $\mathrm{GL}_{n-1}(F)$) so that,

$$\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\mathrm{St}_n, \mathbb{1}_{n-1}) = 0$$

for all integers $i \geq 1$. Here, $\mathbb{1}_{n-1}$ denotes the trivial representation of $\mathrm{GL}_{n-1}(F)$. Now by the Euler-Poincare pairing formula of Dipendra Prasad (see [19]) we have that,

$$\dim \mathrm{Wh}(\mathrm{St}_n) \cdot \dim \mathrm{Wh}(\mathbb{1}_{n-1}) = \sum_{i \geq 0} (-1)^i \dim \mathrm{Ext}_{\mathrm{GL}_{n-1}}^i(\mathrm{St}_n, \mathbb{1}_{n-1}),$$

where $\mathrm{Wh}(\mathrm{St}_n)$ and $\mathrm{Wh}(\mathbb{1}_{n-1})$ are the spaces of Whittaker models for St_n and $\mathbb{1}_{n-1}$ respectively. Hence for $n \geq 3$, using the fact that $\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\mathrm{St}_n, \mathbb{1}_{n-1}) = 0$ for all integers $i \geq 1$, along with the Euler-Poincare pairing implies that $\mathrm{Hom}_{\mathrm{GL}_{n-1}}(\mathrm{St}_n, \mathbb{1}_{n-1}) = 0$. Hence when $n \geq 3$ we have that,

$$\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^*(\mathrm{St}_n, \mathbb{1}_{n-1}) = 0.$$

Notice that for $n \geq 3$ the Arthur parameters of St_n and $\mathbb{1}_{n-1}$,

$$\begin{aligned} \mathcal{A}(\mathrm{St}_n) &= \mathbb{1} \otimes V_n \otimes V_1, \\ \mathcal{A}(\mathbb{1}_{n-1}) &= \mathbb{1} \otimes V_1 \otimes V_{n-1}, \end{aligned}$$

are not strong Ext relevant.

We prove that strong Ext relevance gives a necessary and sufficient condition for predicting Ext non-vanishing when π_1 and π_2 are products of discrete series representations and the Aubert-Zelevinsky duals of discrete series representations. Let us state what this condition on π_1 and π_2 means, in terms of their Arthur parameters. In this case, the Arthur parameters of π_1 and π_2 are a sum of irreducible representations such that for each irreducible direct summand either the Deligne SL_2 or the Arthur SL_2 acts trivially. This means that the Arthur parameters of π_1 and π_2 are a direct sum of irreducible representations of the form $\phi \otimes V_a \otimes V_b$ such that either $a = 1$ or $b = 1$. Here, ϕ denotes an irreducible representation of W_F and V_d denotes the unique d -dimensional irreducible representation of $\mathrm{SL}_2(\mathbb{C})$.

The main result of this paper is the following Ext branching law.

Theorem 1.2. *Let π_1 and π_2 be Arthur type representations of $\mathrm{GL}_n(F)$ and $\mathrm{GL}_{n-1}(F)$ respectively. Suppose that the Arthur parameters of π_1 and π_2 are of the form,*

$$\mathcal{A}(\pi_1) = \sum_{i=1}^r \phi_i \otimes V_{a_i} \otimes V_1 \oplus \sum_{i=r+1}^{r+s} \phi_i \otimes V_1 \otimes V_{b_i}$$

and,

$$\mathcal{A}(\pi_2) = \sum_{i=1}^t \psi_i \otimes V_{c_i} \otimes V_1 \oplus \sum_{i=t+1}^{t+u} \psi_i \otimes V_1 \otimes V_{d_i}.$$

Then,

$$\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\pi_1, \pi_2) \neq 0$$

for some integer $i \geq 0$, if and only if π_1 and π_2 are strong Ext relevant.

The above result can be thought as an Ext analogue of the local non-tempered Gan-Gross-Prasad conjecture in this case.

Remark 1.3. The hypothesis on Arthur parameters in the above theorem amounts to saying that π_1 and π_2 are products of unitary representations of segment type (see Section 3.1.3). This means that π_1 and π_2 are products of unitary representations of the form $Z(\Delta)$ and $Q(\Delta)$ (see Section 2 for notation).

We now restate the definition of strong Ext relevance in representation theoretic terms. We refer to Sections 2 and 3 for unexplained notations. Given an irreducible representation π of $\mathrm{GL}_n(F)$, we let $\pi^{(h)}$ denote its highest derivative. For the sake of notational convenience, we set $\pi^- = \nu^{1/2}\pi^{(h)}$. We let $D(\pi)$ denote the Aubert-Zelevinsky involution of π .

Let $u_\rho(a, b)$ be a Speh representation (see Section 3.1 for notation) whose corresponding Arthur parameter is given as

$$\phi \otimes V_a \otimes V_b.$$

Here ϕ is an irreducible representation of W_F corresponding to the cuspidal representation ρ . Then by Theorem 3.1, the Aubert-Zelevinsky dual of $u_\rho(a, b)$ is equal to $u_\rho(b, a)$ and the corresponding Arthur parameter is equal to

$$\phi \otimes V_b \otimes V_a.$$

Also, by Lemma 3.1, the highest derivative of $\nu^{1/2}u_\rho(a, b)$ is equal to $u_\rho(a, b-1)$, that is,

$$u_\rho(a, b)^- = u_\rho(a, b-1).$$

This leads us to the following equivalent definition of strong Ext relevance in representation theoretic terms.

Definition 1.2. Let $m, n \in \mathbb{Z}_{\geq 0}$ and let π_1 and π_2 be Arthur type representations of $\mathrm{GL}_m(F)$ and $\mathrm{GL}_n(F)$ respectively. Then π_1 and π_2 are strong Ext relevant if there exist Speh representations,

$$\pi_{m,1}, \pi_{m,2}, \dots, \pi_{m,r}, \pi_{n,1}, \pi_{n,2}, \dots, \pi_{n,s}$$

and,

$$\pi_{p,1}, \pi_{p,2}, \dots, \pi_{p,t}, \pi_{q,1}, \pi_{q,2}, \dots, \pi_{q,u}$$

such that,

$$\pi_1 = \pi_{m,1} \times \dots \times \pi_{m,r} \times \pi_{p,1}^- \times \dots \times \pi_{p,t}^- \times \pi_{n,1} \times \dots \times \pi_{n,s} \times D(\pi_{q,1}^-) \times \dots \times D(\pi_{q,u}^-)$$

and,

$$\pi_2 = \pi_{m,1}^- \times \dots \times \pi_{m,r}^- \times \pi_{p,1} \times \dots \times \pi_{p,t} \times D(\pi_{n,1}^-) \times \dots \times D(\pi_{n,s}^-) \times \pi_{q,1} \times \dots \times \pi_{q,u}.$$

Let us now say a few words about what is expected in the general situation when π_1 and π_2 are representations of Arthur type. In general, the above notion of strong Ext relevance is expected to give a sufficient (but not necessary) condition in order to have non-trivial Ext branching, which we formulate in the following conjecture.

Conjecture 1.3. Let π_1 and π_2 be Arthur type representations of $\mathrm{GL}_n(F)$ and $\mathrm{GL}_{n-1}(F)$ respectively. If π_1 and π_2 are strong Ext relevant then,

$$\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\pi_1, \pi_2) \neq 0$$

for some integer $i \geq 0$.

We point out that in general, strong Ext relevance does not give us a necessary condition for Ext non-vanishing. This means that there exist Arthur type irreducible representations π_1 and π_2 of $\mathrm{GL}_n(F)$ and $\mathrm{GL}_{n-1}(F)$ respectively such that $\mathrm{Ext}_{\mathrm{GL}_{n-1}(F)}^i(\pi_1, \pi_2) \neq 0$ for some $i \geq 0$, but π_1 and π_2 are not strong Ext relevant.

Consider the following example. Let ρ denote the trivial representation of G_1 . Consider the representations $\pi_1 = u_\rho(2, 3)$ and $\pi_2 = u_\rho(3, 1) \times \rho \times \rho$ of $GL_6(F)$ and $GL_5(F)$ respectively. Then by Theorem 7.4 of [7] it follows that,

$$\text{Ext}_{GL_5(F)}^*(\pi_1, \pi_2) \neq 0.$$

Note that in this example π_1 and π_2 are not strong Ext relevant.

1.1. Remarks on the Proof. When dealing with the branching problem for the p -adic general linear group it is natural to invoke the Bernstein-Zelevinsky filtration. A very pleasant situation that might occur is the following. Suppose that one is able to show that all pieces other than exactly one in the Bernstein-Zelevinsky filtration do not contribute to the branching law. Moreover, suppose that we are also able to show that higher Ext modules for the other pieces vanish. Then we are left with exactly one piece to analyse which is quite easy to handle. Unfortunately this pleasant situation does not always occur.

Therefore rather than using the Bernstein-Zelevinsky filtration we use a variant of it introduced by K.Y. Chan in the work [7]. This variant (see Theorem 4.1) clumps together some pieces of the Bernstein-Zelevinsky filtration and hence is coarser than the Bernstein-Zelevinsky filtration. Due to some considerations involving cuspidal support one can show that all but one of the pieces in this coarser filtration cannot contribute to the Ext branching. Thus we are left with an easier situation to analyse. As a consequence one is able to replace some factors in the given representation by a suitable cuspidal representation. In the course of the proof we use the transfer lemma (Lemma 4.1) that allows us to transfer the problem from the pair (GL_n, GL_{n-1}) to the pair (GL_{n+1}, GL_n) . The variant of the Bernstein-Zelevinsky filtration introduced by K.Y. Chan and other ideas from [7] are key to our proofs.

1.2. Organization of the Paper. The organization of the paper is as follows. We introduce some preliminary notions and set up notations in Section 2. We then introduce various classes of representations in Section 3. In Section 4, we recall the coarse variant of the Bernstein-Zelevinsky filtration due to K.Y. Chan and collect some important lemmas from [7]. We prove some lemmas about extensions on the same group in Section 5. We prove a preparatory reduction lemma in Section 6. We prove Theorem 1.2 in Sections 7 and 8.

2. PRELIMINARIES

2.1. Basic Notations. Let F be a non-archimedean local field. For a p -adic group G , let $\text{Alg}(G)$ denote the category of smooth representations of G . In this paper we are interested in the study of smooth representations of the p -adic general linear group and all representations considered in this paper are smooth. For $n \in \mathbb{Z}_{\geq 0}$, we let $G_n = GL_n(F)$. Let $\text{Irr}(G_n)$ denote the collection of irreducible representations of G_n and set $\text{Irr} = \cup_{n \geq 0} \text{Irr}(G_n)$. For $\pi \in \text{Alg}(G_n)$ we set $n(\pi) = n$ and let π^\vee denote the smooth dual of π .

Given a p -adic group G and a closed subgroup H and a representation ρ of H , we let $\text{Ind}_H^G(\rho)$ (resp. $\text{c-ind}_H^G(\rho)$) denote the representation of G obtained by normalized induction (resp. normalized compact induction).

We let ν_n denote the character of G_n defined as $\nu_n(g) = |\det(g)|_F$, where $g \in G_n$. We shall usually suppress the subscript and simply denote the character as ν .

Given an irreducible representation π , we let χ_π denote its central character. Suppose that $|\chi_\pi|_F = \nu^s$. We set $e(\pi) = s$. That is, $e(\pi)$ denotes the exponent of the unramified part of χ_π .

Let \mathfrak{R}_n denote the Grothendieck group of the category of smooth representations of G_n . Set $\mathfrak{R} = \bigoplus_{n \geq 0} \mathfrak{R}_n$. The group \mathfrak{R} has the structure of a graded Hopf algebra where the product (denoted by \times) is given by normalized parabolic induction and the co-product (denoted by ∂) is defined using the Jacquet functor.

Suppose $\pi_1 \in \text{Alg}(G_{n_1})$ and $\pi_2 \in \text{Alg}(G_{n_2})$ for some $n_1, n_2 \in \mathbb{Z}_{\geq 0}$. Then $\pi_1 \times \pi_2$ denotes the representation of $G_{n_1+n_2}$ obtained by parabolic induction and is defined as,

$$\pi_1 \times \pi_2 = \text{Ind}_{P_{n_1, n_2}}^{G_{n_1+n_2}}(\pi_1 \otimes \pi_2).$$

Here P_{n_1, n_2} denotes the standard parabolic in $G_{n_1+n_2}$ with Levi $G_{n_1} \times G_{n_2}$ corresponding to the partition (n_1, n_2) . We consider $\pi_1 \otimes \pi_2$ as a representation of P_{n_1, n_2} by extending it trivially across the unipotent subgroup.

Given an irreducible representation π of G_n there exists a unique multiset $\{\rho_1, \rho_2, \dots, \rho_r\}$ consisting of cuspidal representations such that π is a subquotient of $\rho_1 \times \rho_2 \times \dots \times \rho_r$. We call this multiset the cuspidal support of π and denote it as $\text{csupp}(\pi)$. We define $\text{csupp}_{\mathbb{Z}}(\pi) = \{\nu^m \rho \mid \rho \in \text{csupp}(\pi), m \in \mathbb{Z}\}$ and we say that elements of $\text{csupp}_{\mathbb{Z}}(\pi)$ lie in the cuspidal line of π .

Consider $n_1, n_2, \dots, n_r, n \in \mathbb{Z}_{\geq 0}$ such that $n_1 + n_2 + \dots + n_r = n$. Given $\pi \in \text{Alg}(G_n)$, we let $r_{(n_1, \dots, n_r)}(\pi)$ denote the Jacquet module of π , which is a representation of $G_{n_1} \times G_{n_2} \times \dots \times G_{n_r}$. Now, given $\pi \in \text{Alg}(G_n)$, the co-product $\partial : \mathfrak{R} \rightarrow \mathfrak{R} \times \mathfrak{R}$ is defined as,

$$\partial(\pi) = \sum_{i=0}^n r_{(i, n-i)}(\pi).$$

Given $\pi \in \text{Alg}(G_n)$, we shall denote the opposite Jacquet functor corresponding to the partition (n_1, n_2) of n as $\bar{r}_{(n_1, n_2)}(\pi)$ which as a consequence of the second adjointness theorem is,

$$\bar{r}_{(n_1, n_2)}(\pi) \simeq (r_{(n_1, n_2)}(\pi^\vee))^\vee.$$

2.2. Zelevinsky and Langlands Classification. Both the Zelevinsky and Langlands classification parameterize any irreducible representation of G_n in terms of a collection of segments as explained below.

Consider $a, b \in \mathbb{C}$ such that $b - a \in \mathbb{Z}_{\geq 0}$. Given a cuspidal representation ρ of G_n , a segment Δ associated to the datum (ρ, a, b) is an ordered set of irreducible representations of G_n of the form,

$$\Delta = [a, b]_\rho = \{\nu^a \rho, \nu^{a+1} \rho, \dots, \nu^b \rho\}.$$

We set $a(\Delta) = a$ and let $b(\Delta) = b$.

Given the segment $\Delta = [a, b]_\rho$, the associated principal series,

$$\nu^a \rho \times \nu^{a+1} \rho \times \dots \times \nu^b \rho$$

has a unique irreducible submodule and unique irreducible quotient which we denote as $Z(\Delta)$ and $Q(\Delta)$ respectively.

Given cuspidal representations ρ_1 and ρ_2 of G_n , we say ρ_1 precedes ρ_2 if there exists $c \in \mathbb{Z}_{>0}$ such that $\rho_2 = \nu^c \rho_1$. In this case we write $\rho_1 \leq \rho_2$. We say that two segments

Δ_1 and Δ_2 are linked if $\Delta_1 \not\subset \Delta_2$, $\Delta_2 \not\subset \Delta_1$ and $\Delta_1 \cup \Delta_2$ is a segment. We say that Δ_1 precedes Δ_2 if Δ_1 and Δ_2 are linked and $b(\Delta_1) < b(\Delta_2)$.

Given a multiset of segments (which we call a multisegment) $\mathfrak{m} = \{\Delta_1, \Delta_2, \dots, \Delta_r\}$ such that Δ_i does not precede Δ_j for all $i < j$. The representation

$$Z(\Delta_1) \times Z(\Delta_2) \times \dots \times Z(\Delta_r)$$

has a unique irreducible submodule which we denote as $Z(\Delta_1, \Delta_2, \dots, \Delta_r)$. By the Zelevinsky classification any irreducible representation of G_n can be uniquely realized in this manner. Similarly by the Langlands classification, the representation

$$Q(\Delta_1) \times Q(\Delta_2) \times \dots \times Q(\Delta_r)$$

has a unique irreducible quotient which we denote as $Q(\Delta_1, \Delta_2, \dots, \Delta_r)$.

2.3. The Aubert-Zelevinsky Involution. Given a irreducible representation π of $GL_n(F)$ the Aubert-Zelevinsky Involution (denoted by the symbol D) takes π to another irreducible representation of $GL_n(F)$. This involution can be also defined in the more general context of any reductive p -adic group. Since we are only concerned with the general linear group in this paper we only recall the definition for GL_n .

Recall that \mathfrak{R}_n denotes the Grothendieck group of the category $\text{Alg}(G_n)$. The ring \mathfrak{R}_n is a polynomial ring in the indeterminates of the form $Z(\Delta)$ (see [21, Corollary 7.5]). Therefore it is enough to define the involution D on elements of the form $Z(\Delta)$. We define,

$$D : \mathfrak{R}_n \rightarrow \mathfrak{R}_n$$

by sending $Z(\Delta)$ to $Q(\Delta)$. It is known that the involution D takes irreducible representations to irreducible representations.

We have the following cohomological duality theorem (see [16, Theorem 2]) due to Nori and Prasad.

Theorem 2.1. *Let π_1 be a smooth irreducible representations of G_n and π_2 be any smooth representation of G_n . Then for any integer $i \geq 0$,*

$$\text{Ext}_{G_n}^i(\pi_1, \pi_2)^\vee \simeq \text{Ext}_{G_n}^{d(\pi_1)-i}(\pi_2, D(\pi_1)).$$

Here $d(\pi_1)$ is the split rank of the Levi subgroup of G_n which carries the cuspidal support of π_1 .

2.4. On Derivatives and the Bernstein-Zelevinsky Filtration. Given $\pi \in \text{Alg}(G_n)$, we want to define the i -th Bernstein-Zelevinsky derivative $\pi^{(i)}$, which is a smooth representation of G_{n-i} . We first recall the definitions of the Bernstein-Zelevinsky derivative functors that we need. Let $M_n \subset G_n$ denote the mirabolic subgroup inside G_n . We have the decomposition,

$$M_n = G_{n-1} \cdot V_n$$

where, V_n denotes the unipotent subgroup of M_n . We define functors,

$$\Phi^- : \text{Alg}(M_n) \rightarrow \text{Alg}(M_{n-1})$$

and,

$$\Psi^- : \text{Alg}(M_n) \rightarrow \text{Alg}(G_{n-1})$$

as follows.

Given $\tau \in \text{Alg}(M_n)$ we define,

$$\Phi^-(\tau) = \delta^{-1/2} \tau / \langle \tau(u).x - \psi_i(u).x : x \in \tau, u \in V_n \rangle.$$

Here, $\delta^{-1/2}$ is the modulus character of M_n and ψ_n is a non-degenerate character of V_n . Note that M_{n-1} normalizes V_n and ψ_n and therefore we may consider $\Phi^-(\tau)$ to be a smooth representation of M_{n-1} .

Similarly we define,

$$\Psi^-(\tau) = \delta^{-1/2} \tau / \langle \tau(u).x - x : x \in \tau, u \in V_n \rangle.$$

Once again, $\delta^{-1/2}$ denotes the modulus character of M_n . Since G_{n-1} normalizes V_n we have that $\Psi^-(\tau) \in \text{Alg}(G_{n-1})$.

Now given $\pi \in \text{Alg}(G_n)$, we define the i -th Bernstein-Zelevinsky derivative $\pi^{(i)} \in \text{Alg}(G_{n-i})$ as,

$$\pi^{(i)} = \Psi^-(\Phi^-)^{i-1}(\pi|_{M_n}).$$

Given, $\pi \in \text{Alg}(G_n)$, let k be the largest integer such that $\pi^{(k)} \neq 0$. Then we say that the level of π is k and write $\text{lev}(\pi) = k$.

2.4.1. Gelfand-Graev Representation. Let $U_n \subset G_n$ be the subgroup of upper triangular unipotent matrices. Fix a non-degenerate character ψ_n of U_n . The Gelfand-Graev representation of M_n (denoted by Π_n) is defined as,

$$\Pi_n = \text{c-ind}_{U_n}^{M_n} \psi_n.$$

2.4.2. Product Rule for Derivatives. We shall require the following product rule for derivatives (see [4, Corollary 4.14(c)]).

Lemma 2.1. Let $\pi_j \in \text{Alg}(G_{n_j})$ for $j = 1, 2, \dots, k$. Then the representation $(\pi_1 \times \pi_2 \times \dots \times \pi_k)^{(i)}$ has a filtration whose successive quotients are $\pi_1^{(i_1)} \times \pi_2^{(i_2)} \times \dots \times \pi_k^{(i_k)}$, where, $0 \leq i_t \leq n_t$ for $t = 1, 2, \dots, k$ and $i_1 + i_2 + \dots + i_k = i$.

2.4.3. Bernstein-Zelevinsky Filtration. Let $\pi \in \text{Alg}(G_n)$. By the Bernstein-Zelevinsky filtration (see [3]), we have a filtration of submodules $0 = V_n \subsetneq V_{n-1} \subsetneq \dots \subsetneq V_0 = \pi|_{G_{n-1}}$ such that,

$$V_i/V_{i+1} \cong \nu^{1/2} \pi^{(i+1)} \times \Pi_i.$$

Here $U_i \subset G_i$ is the subgroup of upper triangular unipotent matrices and ψ_i is a fixed non-degenerate character of U_i .

2.5. Restriction of a Principal Series to the Mirabolic Subgroup. In this subsection, we want to recall a key lemma (see [4, Proposition 4.13 (a)]) about the restriction of a principal series representation of GL_n to the mirabolic subgroup. We first recall the definitions of two types of mirabolic induction that we shall need from [4, Section 4.12].

Let M_m denote the mirabolic subgroup sitting inside G_m . Let $\text{Mat}_{m,p}$ denote the set of $m \times p$ matrices. Let $0_{m,p}$ denote the zero matrix of order $m \times p$.

Suppose r and s be integers and let $n = r + s$. Let $\rho \in \text{Alg}(G_r)$ and $\tau \in \text{Alg}(M_s)$. We define $\rho \bar{\times} \tau \in \text{Alg}(M_n)$ and $\tau \bar{\times} \rho \in \text{Alg}(M_n)$ as follows.

2.5.1. *Definition of $\rho \bar{\times} \tau$.* Consider the subgroup of M_n defined as follows.

$$H = \left\{ \begin{pmatrix} g & u \\ & m \end{pmatrix} : g \in G_r, m \in M_s, u \in \text{Mat}_{r, n-r} \right\}.$$

We extend the representation $\rho \otimes \tau$ of $G_r \times M_s$ trivially to H . We define,

$$\rho \bar{\times} \tau = \text{Ind}_H^{M_n} \rho \otimes \tau.$$

2.5.2. *Definition of $\tau \bar{\times} \rho$.* Consider the subgroup of M_n defined as follows.

$$H' = \left\{ \begin{pmatrix} g' & u & m' \\ & g & 0_{r,1} \\ & & 1 \end{pmatrix} : g \in G_r, g' \in G_{s-1}, m' \in \text{Mat}_{s-1,1}, u \in \text{Mat}_{s-1,r} \right\}.$$

Set $\tilde{\rho} = \nu^{-1/2} \rho$. We define a representation π of H' acting on the underlying space of $\tau \otimes \rho$ as,

$$\pi \begin{pmatrix} g' & u & m' \\ & g & 0_{r,1} \\ & & 1 \end{pmatrix} = \tau \begin{pmatrix} g' & m' \\ 0_{1,s-1} & 1 \end{pmatrix} \otimes \tilde{\rho}(g).$$

Then $\tau \bar{\times} \rho$ is defined as,

$$\tau \bar{\times} \rho = \text{Ind}_{H'}^{M_n} \pi.$$

We are now in a position to state the required lemma. The lemma below tells us how a principal series representation decomposes upon restriction to the mirabolic subgroup.

Lemma 2.2. [4, Proposition 4.13(a)] Let π_1 and π_2 be smooth representations of G_{n_1} and G_{n_2} respectively. Then

$$0 \rightarrow \pi_1|_{M_{n_1}} \bar{\times} \pi_2 \rightarrow (\pi_1 \times \pi_2)|_{M_{n_1+n_2}} \rightarrow \pi_1 \bar{\times} \pi_2|_{M_{n_2}} \rightarrow 0.$$

Here M_{n_1} and M_{n_2} are the mirabolic subgroups of G_{n_1} and G_{n_2} respectively.

2.6. Second Adjointness and the Identity orbit in the Geometric Lemma. Let $\pi_1 \in \text{Alg}(G_{n_1})$ and $\pi_2 \in \text{Alg}(G_{n_2})$ for some $n_1, n_2 \in \mathbb{Z}_{\geq 0}$. Set $n = n_1 + n_2$ and consider the representation $\pi_1 \times \pi_2$ of G_n . The geometric lemma (see [4]) describes the factors in a composition series of the Jacquet module $r_{(n_1, n_2)}(\pi_1 \times \pi_2)$. Each factor in the composition series of $r_{(n_1, n_2)}(\pi_1 \times \pi_2)$ corresponds to a double coset representative of $P_{n_1, n_2} \backslash G_n / P_{n_1, n_2}$ where, P_{n_1, n_2} denotes the parabolic inside G_n corresponding to the partition (n_1, n_2) . Here, the term $\pi_1 \otimes \pi_2$ in the composition series of $r_{(n_1, n_2)}(\pi_1 \times \pi_2)$ corresponds to the identity double coset. By the geometric lemma the term $\pi_1 \otimes \pi_2$ occurs as the quotient of $r_{(n_1, n_2)}(\pi_1 \times \pi_2)$.

We want to understand the structure of the opposite Jacquet module of $\bar{r}_{(n_1, n_2)}(\pi_1 \times \pi_2)$. Since we know that,

$$\bar{r}_{(n_1, n_2)}(\pi_1 \times \pi_2) \simeq (r_{(n_1, n_2)}((\pi_1 \times \pi_2)^\vee))^\vee$$

the above discussion leads to the following corollary.

Corollary 2.1. Let $\pi_1 \in \text{Alg}(G_{n_1})$ and $\pi_2 \in \text{Alg}(G_{n_2})$ for some $n_1, n_2 \in \mathbb{Z}_{\geq 0}$. Then $\pi_1 \otimes \pi_2$ occurs as a submodule of $\bar{r}_{(n_1, n_2)}(\pi_1 \times \pi_2)$.

2.7. Kunneth Formula and a lemma on Ext Vanishing. We shall require the following Kunneth formula in the course of our calculations of Ext modules.

Theorem 2.2. ([19, Theorem 3.1]) *Let H_1 and H_2 be p -adic groups and suppose that H_1 is reductive. Let E_1 and F_1 (resp. E_2 and F_2) be smooth representations of H_1 (resp. H_2). If E_1 has finite length then,*

$$\mathrm{Ext}_{H_1 \times H_2}^i(E_1 \otimes E_2, F_1 \times F_2) = \sum_{k=0}^i \mathrm{Ext}_{H_1}^k(E_1, F_1) \otimes \mathrm{Ext}_{H_2}^{i-k}(E_2, F_2).$$

The above Kunneth formula when combined with cuspidal support considerations allows us to conclude Ext vanishing in some cases as in the lemma below. The lemma below is a slight reformulation of [7, Lemma 2.4].

Lemma 2.3. Let π_1 be an irreducible representation of G_{n-i} and π_2 be a smooth representation of G_i not necessarily of finite length. Let π be an irreducible representation of G_n . Let σ be a cuspidal representation such that $\sigma \in \mathrm{csupp}(\pi_1)$ but $\sigma \notin \mathrm{csupp}(\pi)$. Then for all integers $i \geq 0$,

$$\mathrm{Ext}_{G_n}^i(\pi_1 \times \pi_2, \pi) = 0.$$

Remark 2.1. The above lemma codifies a standard argument that we shall use. In the above lemma, since π_2 is not necessarily of finite length we cannot directly conclude Ext vanishing by cuspidal support considerations. Instead we first apply second adjointness and the Kunneth formula for each simple composition factor of the opposite Jacquet module of π . Now by comparing cuspidal supports at $\sigma \in \mathrm{csupp}(\pi_1)$ and the corresponding factor in the opposite Jacquet module of π we conclude Ext vanishing.

2.8. Generic Branching Law. We recall the Hom and Ext branching law for generic representations. This will form the base case for the inductive proof of our theorems.

Theorem 2.3. ([9, Theorem 4.1]) *Let $n \geq 0$ be an integer. Let π_1 and π_2 be irreducible generic representations of G_n and G_{n-1} respectively. Then*

$$\mathrm{Hom}_{G_{n-1}}(\pi_1, \pi_2) = \mathbb{C},$$

and for all integers $i \geq 1$,

$$\mathrm{Ext}_{G_{n-1}}^i(\pi_1, \pi_2) = 0.$$

The above theorem about higher Ext vanishing was later extended to the case when π_1 and the dual of π_2 are standard representations by K.Y. Chan (see [8, Theorem 1.1]).

3. ON CLASSES OF REPRESENTATIONS OF $\mathrm{GL}_n(F)$

In this Section, we introduce various classes of representations of the p -adic general linear group. Let $\mathrm{Irr}^c(G_n)$ (resp. $\mathrm{Irr}^u(G_n)$) denote the collection of irreducible cuspidal (resp. unitary) representations of G_n . Let $\mathrm{Irr}^{c,u}(G_n) = \mathrm{Irr}^c(G_n) \cap \mathrm{Irr}^u(G_n)$ denote the class of cuspidal unitary representations. We set $\mathrm{Irr}^c = \cup_{n \geq 0} \mathrm{Irr}^c(G_n)$, $\mathrm{Irr}^u = \cup_{n \geq 0} \mathrm{Irr}^u(G_n)$ and $\mathrm{Irr}^{c,u} = \cup_{n \geq 0} \mathrm{Irr}^{c,u}(G_n)$.

3.1. On Speh and Arthur Representations. The Speh representations are the building blocks of unitary representations and are defined as follows. Let $\rho \in \text{Irr}^{c,u}$. For $a \in \mathbb{Z}_{\geq 0}$ let $\Delta(\rho, a)$ be the segment $\Delta(\rho, a) = [-\frac{(a-1)}{2}, \frac{(a-1)}{2}]_{\rho}$. We let $\delta_{\rho}(a)$ denote the square integrable representation $\delta_{\rho}(a) = Q(\Delta(\rho, a))$.

Let b be a non-negative integer. The product,

$$\nu^{\frac{(b-1)}{2}} \delta_{\rho}(a) \times \nu^{\frac{(b-1)}{2}-1} \delta_{\rho}(a) \times \dots \times \nu^{-\frac{(b-1)}{2}} \delta_{\rho}(a)$$

has a unique irreducible quotient which we denote as $u_{\rho}(a, b)$. The irreducible representations of the form $u_{\rho}(a, b)$ are known as Speh representations.

Definition 3.1. We say that an irreducible representation π of G_n is of Arthur type if there exist Speh representations $\pi_1, \pi_2, \dots, \pi_k$ such that,

$$\pi = \pi_1 \times \pi_2 \times \dots \times \pi_k.$$

3.1.1. Derivatives of Speh Representations. We shall gather some information about the derivatives of Speh representations. The knowledge of cuspidal support of the derivatives of Speh representations will be helpful in allowing us to conclude Ext vanishing.

The lemma below follows from [15, Theorem 14] when it is reformulated in terms of the Langlands classification. We refer to the discussion after Theorem 14 in loc. cit. Given $\pi \in \text{Alg}(G_n)$, recall that by the level of π (denoted $\text{lev}(\pi)$) we mean the largest integer k such that $\pi^{(k)} \neq 0$.

Lemma 3.1. Let $\pi = u_{\rho}(a, b)$ be a Speh representation. Then the level of π is equal to $n(\rho)a$, that is, $\text{lev}(\pi) = n(\rho)a$.

- (1) If $k < \text{lev}(\pi)$ then $\nu^{(m+n-2)/2} \rho \in \text{csupp}(\pi^{(k)})$.
- (2) If $k = \text{lev}(\pi)$ then $\pi^{(k)} = \nu^{-1/2} u_{\rho}(a, b-1)$.

3.1.2. Aubert-Zelevinsky Dual of a Speh Representation. We shall need the following calculation of the Aubert-Zelevinsky dual of a Speh representation. The following result is due to Tadic (see [20, Theorem B]).

Theorem 3.1. Let $a, b \in \mathbb{Z}_{\geq 0}$ and $\rho \in \text{Irr}^c$. Let $\pi = u_{\rho}(a, b)$ be a Speh representation of G_n . Then,

$$D(u_{\rho}(a, b)) = u_{\rho}(b, a).$$

The above result can be interpreted in the language of Arthur parameters as follows. Let $\pi = u_{\rho}(a, b)$ be a Speh representation of G_n that arises from an Arthur parameter of the form,

$$\psi \otimes \text{Sym}^a(\mathbb{C}^2) \otimes \text{Sym}^b(\mathbb{C}^2).$$

Then the Aubert-Zelevinsky dual of $u_{\rho}(a, b)$ is once again a Speh representation whose corresponding Arthur parameter is given as,

$$\psi \otimes \text{Sym}^b(\mathbb{C}^2) \otimes \text{Sym}^a(\mathbb{C}^2).$$

3.1.3. Definition of Speh Representations of Segment Type. We introduce some terminology for the sake of convenience. Suppose that we are given a Speh representation $u_{\rho}(a, b)$ of G_n . If $a = 1$ then the Speh representation $u_{\rho}(a, b)$ is of the form $Z(\Delta)$ for the segment $\Delta = [-\frac{(b-1)}{2}, \frac{(b-1)}{2}]_{\rho}$. If $b = 1$ then the Speh representation $u_{\rho}(a, b)$ is of the form $Q(\Delta)$ for the segment $\Delta = [-\frac{(a-1)}{2}, \frac{(a-1)}{2}]_{\rho}$.

If a Speh representation is of the form $Q(\Delta)$ or $Z(\Delta)$ for some segment Δ then we shall say that the Speh representation is of *segment type*. If $\pi \in \text{Irr}(G_n)$ is of segment type then its Arthur parameter $A(\pi)$ is such that either the Weil-Deligne SL_2 factor of $A(\pi)$ or the Arthur SL_2 factor of $A(\pi)$ acts trivially.

We shall require the following computations of Jacquet modules (see [21, Proposition 3.4] and [21, Proposition 9.5]).

Lemma 3.2. Let ρ be a cuspidal representation of G_m and consider the segment $\Delta = [\rho, \rho^{k-1}]$ where, $n = km$. If l is not divisible by m then $r_{(n-l,l)}(Q(\Delta)) = \bar{r}_{(n-l,l)}(Q(\Delta)) = r_{(n-l,l)}(Z(\Delta)) = \bar{r}_{(n-l,l)}(Z(\Delta)) = 0$. If $l = mp$ then,

$$\begin{aligned} r_{(n-l,l)}(Q(\Delta)) &= Q([\nu^p \rho, \nu^{k-1} \rho]) \otimes Q([\rho, \nu^{p-1} \rho]). \\ \bar{r}_{(n-l,l)}(Q(\Delta)) &= Q([\rho, \nu^{k-p-1} \rho]) \otimes Q([\nu^{k-p} \rho, \nu^{k-1} \rho]). \\ r_{(n-l,l)}(Z(\Delta)) &= Z([\rho, \nu^{k-p-1} \rho]) \otimes Z([\nu^{k-p} \rho, \nu^{k-1} \rho]). \\ \bar{r}_{(n-l,l)}(Z(\Delta)) &= Z([\nu^p \rho, \nu^{k-1} \rho]) \otimes Z([\rho, \nu^{p-1} \rho]). \end{aligned}$$

Remark 3.1. Suppose $Q(\Delta)$ is a unitary representation of G_n . Suppose $r_{(n-l,l)}(Q(\Delta)) = \omega_1 \otimes \omega_2$ where, $r_{(n-l,l)}(Q(\Delta))$ is the Jacquet module of $Q(\Delta)$ with respect to a proper parabolic of G_n corresponding to the partition $(n-l, l)$ of n . Then by the first computation in the above lemma it is evident that $e(\omega_1) > 0$ whereas, $e(\omega_2) < 0$. We recall that given an irreducible representation π , $e(\pi)$ denotes the exponent of the unramified part of the central character of π . We can make analogous assertions for the remaining three cases of the above lemma.

4. A COARSE VARIANT OF THE BERNSTEIN-ZELEVINSKY FILTRATION

In this Section, we recall the filtration for parabolically induced representations introduced by K.Y. Chan in the work [7]. It will be the main tool in our underlying our proofs. We also state two lemmas from [7] that follow by an application of this filtration.

4.1. Filtration for Parabolically Induced Modules. Since the following filtration on parabolically induced modules is going to play a key role in proving our results, we recall the main idea behind the proof for the sake of completeness.

Theorem 4.1. [7, Proposition 5.13] *Let π_1 and π_2 be smooth representations of G_{n_1} and G_{n_2} respectively. Then $(\pi_1 \times \pi_2)|_{G_{n_1+n_2-1}}$ has a filtration,*

$$0 = V_d \subsetneq V_{d-1} \subsetneq \dots \subsetneq V_0 = (\pi_1 \times \pi_2)|_{G_{n_1+n_2-1}}$$

such that,

$$V_0/V_1 \cong \nu^{1/2} \pi_1 \times \pi_2|_{G_{n_2-1}},$$

and for $1 \leq k \leq d-1$,

$$V_k/V_{k+1} \cong \nu^{1/2} \pi_1^{(k)} \times (\Pi_k \bar{\times} \pi_2)|_{G_{n_2+k-1}}.$$

Here, Π_k denotes the Gelfand-Graev representation of the mirabolic group M_k and $\bar{\times}$ denotes the mirabolic induction as defined in Section 2.5.

Sketch of the Proof: By Lemma 2.2, we obtain the following exact sequence,

$$0 \rightarrow \pi_1|_{M_{n_1}} \bar{\times} \pi_2 \rightarrow (\pi_1 \times \pi_2)|_{M_{n_1+n_2}} \rightarrow \pi_1 \bar{\times} \pi_2|_{M_{n_2}} \rightarrow 0.$$

On restricting $\pi_1|_{M_{n_1}} \bar{\times} \pi_2$ to $G_{n_1+n_2-1}$ we get $\nu^{1/2}\pi_1 \times \pi_2|_{G_{n_2-1}}$ which gives us the first term in our filtration.

For the submodule $\pi_1|_{M_{n_1}} \bar{\times} \pi_2$ we apply the Bernstein-Zelevinsky filtration to decompose $\pi_1|_{M_{n_1}}$ and then restrict to $G_{n_1+n_2-1}$ in order to obtain the rest of the terms our lemma.

Remark 4.1. As pointed out earlier the above filtration can be thought of as a coarse version of the Bernstein-Zelevinsky filtration where we club together some of the pieces in the Bernstein-Zelevinsky filtration. As we shall see in Lemma 6.1, an advantage of having this coarser filtration is that we can establish Ext vanishing for all but one of the pieces of this filtration making our situation easier to analyse.

4.2. Transfer Lemma. The following lemma shall allow us to transfer a branching problem for the pair (G_n, G_{n-1}) to a branching problem for the pair (G_{n-1}, G_{n-2}) and vice-versa.

Lemma 4.1. [7, Proposition 4.1] Let π_1 and π_2 be irreducible representations of G_n and G_{n-1} respectively. Then for any cuspidal representation σ of G_2 such that $\sigma \notin \text{csupp}_{\mathbb{Z}}(\nu^{-1/2}\pi_1^\vee) \cup \text{csupp}_{\mathbb{Z}}(\pi_2)$ we have that,

$$\text{Ext}_{G_{n-1}}^i(\pi_1, \pi_2^\vee) = \text{Ext}_{G_n}^i(\pi_2 \times \sigma, \pi_1^\vee)$$

for all integers $i \geq 0$.

4.3. A Replacement Lemma. We shall also require the following replacement lemma. It will be useful in the proof of Lemma 6.1 by allowing us to replace some factors in a restriction of an Arthur type representation by some cuspidal representation.

Lemma 4.2. [7, Lemma 3.6] Let $\pi_1 \in \text{Alg}(G_k)$ and $\pi_2 \in \text{Alg}(G_l)$. Let $\pi_3 \in \text{Alg}(G_n)$ with $n \geq l + k$. Let $a = (n + 1) - (k + l)$. Then, for any σ in $\text{Irr}^c(G_a)$ such that $\sigma \notin \text{csupp}_{\mathbb{Z}}(\nu^{-1/2}\pi_3)$, and for any i ,

$$\text{Ext}_{G_n}^i(\pi_1 \times (\sigma \times \pi_2)|_{G_{n-k}}, \pi_3) = \text{Ext}_{G_n}^i(\pi_1 \times (\Pi_a \bar{\times} \pi_2)|_{G_{n-k}}, \pi_3)$$

5. SOME LEMMAS ON EXTENSIONS ON THE SAME GROUP

In this Section, we prove some lemmas about extensions between representations of Arthur type on the same group. These lemmas about extensions on the same group are needed because we shall ultimately reduce the Ext branching problem to a study of extensions between representations of the same group.

5.1. Extensions between Speh Representations. We have the following lemma about Ext modules of Speh representations on the same group.

Lemma 5.1. Let π_1 and π_2 be two Speh representations of G_n such that,

$$\text{Ext}_{G_n}^i(\pi_1, \pi_2) \neq 0$$

for some integer $i \geq 0$. Then either $\pi_2 = \pi_1$ or $\pi_2 = D(\pi_1)$.

Proof. The proof follows from looking at the cuspidal supports. \square

Remark 5.1. The above lemma (proved by looking at cuspidal supports) is equivalent to the observation that if V_1, V_2, V_3 and V_4 are finite dimensional irreducible representations of $\mathrm{SL}_2(\mathbb{C})$ such that,

$$V_1 \otimes V_2 = V_3 \otimes V_4$$

then, $(V_3, V_4) = (V_1, V_2)$ or $(V_3, V_4) = (V_2, V_1)$.

5.2. Lemmas about extensions between products of unitary segment type representations. Recall that by a unitary representation of segment type (see Section 3.1.3), we mean a unitary representation of the form $Z(\Delta)$ or $Q(\Delta)$ for some segment Δ . In the proof of the lemma below we shall freely invoke Lemma 3.2 and the observations in Remark 3.1 without citing them explicitly each time.

Lemma 5.2. Let π_1 be a unitary segment type representation of G_n and let $\tau \in \mathrm{Alg}(G_k)$ (not necessarily of finite length). Let $\pi_2 \in \mathrm{Alg}(G_{n+k})$ be an irreducible unitary representation of Arthur type. Suppose that $\pi_2 = \pi_{2,1} \times \pi_{2,2} \times \dots \times \pi_{2,s}$ where, $\pi_{2,1}, \pi_{2,2}, \dots, \pi_{2,s}$ are unitary representations of segment type. Suppose that,

$$\mathrm{Ext}_{G_{n+k}}^i(\pi_1 \times \tau, \pi_2) \neq 0$$

for some integer $i \geq 0$. Then one of the following holds:

- (1) $\pi_1 = \pi_{2,m}$ for some $m \in \{1, 2, \dots, s\}$. And, $\pi_2 = \pi_1 \times \pi'_2$ where $\pi'_2 \in \mathrm{Alg}(G_k)$ is the product of the factors $\pi_{2,1}, \dots, \pi_{2,m-1}, \pi_{2,m+1}, \dots, \pi_{2,s}$. Moreover,

$$\mathrm{Ext}_{G_k}^j(\tau, \pi'_2) \neq 0$$

for some $j \leq i$.

- (2) $D(\pi_1) = \pi_{2,m}$ for some $m \in \{1, 2, \dots, s\}$. And, $\pi_2 = D(\pi_1) \times \pi''_2$ where $\pi''_2 \in \mathrm{Alg}(G_k)$ is the product of the factors $\pi_{2,1}, \dots, \pi_{2,m-1}, \pi_{2,m+1}, \dots, \pi_{2,s}$. Moreover,

$$\mathrm{Ext}_{G_k}^j(\tau, \pi''_2) \neq 0$$

for some $j < i$.

Proof. We are given that,

$$\mathrm{Ext}_{G_{n+k}}^i(\pi_1 \times \tau, \pi_2) \neq 0.$$

By second adjointness we have that,

$$\mathrm{Ext}_{G_n \times G_k}^i(\pi_1 \otimes \tau, \Omega) \neq 0 \tag{5.1}$$

for some irreducible subquotient Ω of $\bar{r}_{(n,k)}(\pi_2)$. Let us suppose that π_1 is of the form $Z(\Delta)$. The case when π_1 is of the form $Q(\Delta)$ can be handled similarly. Since π_2 is irreducible we arrange the factors of π_2 so that factors of the form $Z(\Delta)$ occur first followed by factors of the form $Q(\Delta)$. Therefore, let us suppose that $\pi_1 = Z(\Delta_0)$, $\pi_{2,i} = Z(\Delta_{2,i})$ ($i = 1, 2, \dots, k$) and $\pi_{2,j} = Q(\Delta_{2,j})$ ($j = k + 1, \dots, s$). Here, $0 \leq k \leq s$ is an integer and $\Delta_0, \Delta_{2,i}$ ($i = 1, 2, \dots, k$) and $\Delta_{2,j}$ ($j = k + 1, \dots, s$) are some segments. Note that all the representations $\pi_{2,1}, \pi_{2,2}, \dots, \pi_{2,s}$ and π_1 are unitary.

By the geometric lemma we obtain a filtration on $\bar{r}_{(n,k)}(\pi_2)$ where the subquotients are of the form,

$$(\omega_{1,1} \times \dots \times \omega_{1,k} \times \xi_{1,k+1} \times \dots \times \xi_{1,s}) \otimes (\omega_{2,1} \times \dots \times \omega_{2,k} \times \xi_{2,k+1} \times \dots \times \xi_{2,s}),$$

where $\omega_{1,i} \otimes \omega_{2,i}$ is the opposite Jacquet module of $\pi_{2,i}$ (here $i = 1, 2, \dots, k$) with respect to a suitable opposite parabolic. Similarly, $\xi_{1,j} \otimes \xi_{2,j}$ is the opposite Jacquet module of $\pi_{2,j}$ (here $j = k + 1, \dots, s$) with respect to a suitable opposite parabolic.

We wish to analyse which of the above subquotients can make a non-zero contribution in Equation 5.1. We carry out this analysis by considering the kinds of subquotients as in the following two cases.

Case 1: Suppose each of the factors $\omega_{1,1}, \omega_{2,1}, \dots, \omega_{1,k}$ is equal to the trivial representation of $G_0 = \langle e \rangle$. Let us suppose that a subquotient of this kind makes a non-zero contribution in Equation 5.1. In this case by Equation 5.1 and the Kunneth formula we conclude that,

$$\text{Ext}_{G_n}^p(\pi_1, \xi_{1,k+1} \times \dots \times \xi_{1,s}) \neq 0 \quad (5.2)$$

for some integer $p \leq i$.

Note that $e(\xi_{1,j}) \leq 0$ for all $j = k + 1, \dots, s$ whereas, $e(\pi_1) = 0$. Therefore, by comparing central characters we conclude that for all $j = k + 1, \dots, s$ either $\xi_{1,j} = Q(\Delta_{2j})$ or $\xi_{1,j}$ is the trivial representation of G_0 . We now claim that $\xi_{1,j} = Q(\Delta_{2j})$ for exactly one $j \in \{k + 1, \dots, s\}$.

Suppose for the sake of contradiction that $\xi_{1,j} = Q(\Delta_{2j})$ for two distinct integers j_1 and j_2 . Without loss of generality we may take $j_1 = k + 1$ and $j_2 = k + 2$. By Equation 5.2 we know that,

$$\text{Ext}_{G_n}^p(\pi_1, Q(\Delta_{2,k+1}) \times Q(\Delta_{2,k+2}) \times \zeta) \neq 0,$$

where, $\zeta = \xi_{1,k+3} \times \dots \times \xi_{1,s}$. Now by Frobenius reciprocity,

$$\text{Ext}_{G_l \times G_{n-l}}^p(\eta_1 \otimes \eta_2, Q(\Delta_{2,k+1}) \otimes (Q(\Delta_{2,k+2}) \times \zeta)) \neq 0,$$

where, $l = n(Q(\Delta_{2,k+1}))$ and $\eta_1 \otimes \eta_2$ is the Jacquet module of π_1 with respect to a suitable parabolic. Hence we obtain that,

$$\text{Ext}_{G_l}^q(\eta_1, Q(\Delta_{2,k+1})) \neq 0,$$

for some integer $q \leq p$. Now since $e(\eta_1) > 0$ and $e(Q(\Delta_{2,k+1})) = 0$, by comparing central characters we obtain a contradiction.

Hence we conclude that in Case 1, the only subquotient of $\bar{r}_{(n,k)}(\pi_2)$ that can make a non-zero contribution in Equation 5.1 is of the form,

$$\Omega = Q(\Delta_{2j}) \otimes \pi''$$

where, π'' is the product of segment type representations $\pi_{2,1}, \pi_{2,2}, \dots, \pi_{2,s}$ except the term $Q(\Delta_{2j})$.

Now by Equation 5.1, applying the Kunneth formula and Lemma 5.1 proves part (2) of our lemma.

Case 2: Now suppose that atleast one of the factors $\omega_{1,1}, \omega_{2,1}, \dots, \omega_{1,k}$ is a representation of G_M , where $M \geq 1$. Without loss of generality we may assume that $\omega_{1,1}$ is a representation of G_M , where $M \geq 1$. Let us suppose that a subquotient of this kind makes a non-zero contribution in Equation 5.1. Now by Equation 5.1 and the Kunneth formula we have that,

$$\text{Ext}_{G_n}^p(\pi_1, \omega_{1,1} \times \omega_{2,1} \times \dots \times \omega_{1,k} \times \xi_{1,k+1} \times \dots \times \xi_{1,s}) \neq 0$$

for some integer $p \leq i$.

Hence, by Frobenius reciprocity we have that,

$$\text{Ext}_{G_l \times G_{n-l}}^p(\eta_1 \otimes \eta_2, \omega_{1,1} \otimes (\omega_{2,1} \times \dots \times \omega_{1,k} \times \xi_{1,k+1} \times \dots \times \xi_{1,s})) \neq 0,$$

where, $l = n(\omega_{1,1})$ and $\eta_1 \otimes \eta_2$ is the Jacquet module of π_1 with respect to a suitable parabolic. Hence we have that,

$$\text{Ext}_{G_l}^q(\eta_1, \omega_{1,1}) \neq 0, \quad (5.3)$$

for some integer $q \leq 0$. Let us suppose that $\pi_1 = Z(\Delta_0) = Z([-\frac{(x-1)}{2}, \frac{(x-1)}{2}]_{\rho_1})$ and $\pi_{2,1} = Z(\Delta_{2,1}) = Z([-\frac{(y-1)}{2}, \frac{(y-1)}{2}]_{\rho_2})$, where x, y are integers such that $x, y \geq 1$, and ρ_1 and ρ_2 are some unitary cuspidal representations.

Then $\eta_1 = Z([-\frac{(x-1)}{2}, w]_{\rho_1})$ and $\omega_{1,1} = Z([z, \frac{(y-1)}{2}]_{\rho_2})$, where, w, z are integers such that $w \leq \frac{(x-1)}{2}$ and $z \geq -\frac{(y-1)}{2}$. Now by comparing cuspidal supports in Equation 5.3 we conclude that $\rho_1 = \rho_2$, $w = \frac{(y-1)}{2}$ and $z = -\frac{(x-1)}{2}$. Substituting these in the above inequalities we conclude that $x = y$. Hence $\eta_1 = \pi_1$ and $\omega_{1,1} = \pi_{2,1} \cong \pi_1$.

Hence we conclude that in Case 2, the only subquotient of $\bar{r}_{(n,k)}(\pi_2)$ that can make a non-zero contribution in Equation 5.1 is of the form,

$$\Omega = Z(\Delta_{2i}) \otimes \pi',$$

where, π'' is the product of segment type representations $\pi_{2,1}, \pi_{2,2}, \dots, \pi_{2,s}$ except the term $Z(\Delta_{2i})$. Now using Equation 5.1 and applying the Kunneth formula proves part (1) of our lemma. \square

Remark 5.2. In the above lemma, the hypothesis that π_1 and $\pi_{2,1}, \pi_{2,2}, \dots, \pi_{2,s}$ are unitary segment type representations cannot be dropped. For instance, the above lemma does not hold if we assume π_1 to be an arbitrary Speth representation.

Consider the following example. Let ρ denote the trivial representation of G_1 . Let $\pi_1 = Q([0, 1]_{\rho}, [-1, 0]_{\rho})$ and $\pi_2 = \rho \times Q([-1, 0, 1]_{\rho})$. We claim that,

$$\text{Ext}_{G_4}^1(\pi_1, \pi_2) \neq 0.$$

By Frobenius reciprocity we have that,

$$\text{Ext}_{G_4}^i(\pi_1, \pi_2) = \text{Ext}_{G_3 \times G_1}^i(\rho \otimes Q([0, 1]_{\rho}, [-1]_{\rho}), \rho \otimes Q([-1, 0, 1]_{\rho})) \neq 0.$$

Here, we have used the description of Jacquet modules of ladder representations in [14]. By comparing cuspidal supports $\rho \otimes Q([0, 1]_{\rho}, [-1]_{\rho})$ is the only subquotient of the $r_{(1,3)}(\pi_1)$ that can contribute to the above Ext module. By [17] we know that,

$$\text{Ext}_{G_3}^1(Q([0, 1]_{\rho}, [-1]_{\rho}), Q([-1, 0, 1]_{\rho})) \neq 0.$$

Applying the Kunneth formula we conclude that,

$$\text{Ext}_{G_4}^1(\pi_1, \pi_2) \neq 0.$$

Remark 5.3. The above lemma can be thought of as an Ext analogue of [7, Proposition 4.2] albeit with much more restrictive hypothesis. The Proposition 4.2 in [7] follows from [7, Theorem 9.1] (see [5] for a generalization of [7, Theorem 9.1]).

The following lemma is like a converse to Lemma 5.2. The statement of the lemma below is obvious if one takes Hom instead of Ext modules.

Lemma 5.3. Let π_1 be a unitary segment type representation of G_n and let $\tau \in \text{Alg}(G_k)$ (not necessarily of finite length). Let $\sigma \in \text{Alg}(G_k)$ be an irreducible unitary representation of Arthur type. Suppose that $\sigma = \sigma_1 \times \sigma_2 \times \dots \times \sigma_s$ where, $\sigma_1, \sigma_2, \dots, \sigma_s$ are unitary representations of segment type. Suppose that,

$$\text{Ext}_{G_k}^j(\tau, \sigma) \neq 0$$

for some integer $j \geq 0$. Then

$$\text{Ext}_{G_{n+k}}^i(\pi_1 \times \tau, \pi_1 \times \sigma) \neq 0$$

for some integer $i \geq 0$.

Proof. By second adjointness we have that,

$$\text{Ext}_{G_{n+k}}^i(\pi_1 \times \tau, \pi_1 \times \sigma) = \text{Ext}_{G_n \times G_k}^i(\pi_1 \otimes \tau, \bar{r}_P(\pi_1 \times \sigma)). \quad (5.4)$$

Here \bar{r}_P denotes the opposite Jacquet module with respect to the opposite parabolic corresponding to the partition (n, k) . By the analysis carried out in the proof of Lemma 5.2, the only subquotients of $\bar{r}_P(\pi_1 \times \sigma)$ that can make a non-zero contribution to the Ext module in the right hand side of Equation 5.4 are of the following two kinds:

- (A) $\Omega_1 = \pi_1 \otimes \sigma$. Note that by the discussion in Section 2.6, Ω_1 occurs as a *submodule* of $\bar{r}_P(\pi_1 \times \sigma)$.
- (B) $\Omega_2 = D(\pi_1) \otimes \sigma'$. Note that if Ω_2 occurs as a subquotient of $\bar{r}_P(\pi_1 \times \sigma)$ then there exist $m \in \{1, 2, \dots, s\}$ such that $\sigma_m = D(\pi_1)$. Here, $\sigma' = \pi_1 \times \sigma_1 \times \dots \times \sigma_{m-1} \times \sigma_{m+1} \times \dots \times \sigma_s$. In this case we have that $\pi_1 \times \sigma = D(\pi_1) \times \sigma'$ and once again by the discussion in Section 2.6, we can ensure that Ω_2 occurs as a *submodule* of $\bar{r}_P(\pi_1 \times \sigma)$.

We are given that

$$\text{Ext}_{G_k}^j(\tau, \sigma) \neq 0 \quad (5.5)$$

for some integer $j \geq 0$. Also, we know that

$$\text{Hom}_{G_n}(\pi_1, \pi_1) \neq 0. \quad (5.6)$$

Hence by Equation 5.5, Equation 5.6 and the Kunneth formula we conclude that,

$$\text{Ext}_{G_n \times G_k}^j(\pi_1 \otimes \tau, \pi_1 \otimes \sigma) \neq 0. \quad (5.7)$$

We define the following sets. For $t = 1, 2$ we define,

$$S_t = \{h \in \mathbb{Z} \mid \text{Ext}_{G_n \times G_k}^h(\pi_1 \otimes \tau, \Omega_t) \neq 0\}.$$

Let us denote $S = S_1 \cup S_2$. Then by Equation 5.7 it is evident that the set S is non-empty. Let i^* denote the smallest integer in the set S . Then $\text{Ext}_{G_n \times G_k}^{i^*}(\pi_1 \otimes \tau, \Omega_{t^*}) \neq 0$ where $t^* \in \{1, 2\}$.

Since Ω_{t^*} occurs as a *submodule* of $\bar{r}_P(\pi_1 \times \sigma)$ by a long exact sequence argument we get that,

$$\text{Ext}_{G_n \times G_k}^{i^*}(\pi_1 \otimes \tau, \bar{r}_P(\pi_1 \times \sigma)) \neq 0.$$

Hence by Frobenius reciprocity we conclude that,

$$\text{Ext}_{G_{n+k}}^{i^*}(\pi_1 \times \tau, \pi_1 \times \sigma) \neq 0.$$

□

We also have the following similar lemma.

Lemma 5.4. Let π_1 be a unitary segment type representation of G_n and let $\tau \in \text{Alg}(G_k)$ (not necessarily of finite length). Let $\sigma \in \text{Alg}(G_k)$ be an irreducible unitary representation of Arthur type. Suppose that $\sigma = \sigma_1 \times \sigma_2 \times \dots \times \sigma_s$ where, $\sigma_1, \sigma_2, \dots, \sigma_s$ are unitary representations of segment type. Suppose that,

$$\text{Ext}_{G_k}^j(\tau, \sigma) \neq 0$$

for some integer $j \geq 0$. Then

$$\text{Ext}_{G_{n+k}}^i(\pi_1 \times \tau, D(\pi_1) \times \sigma) \neq 0$$

for some integer $i \geq 0$.

The proof of the above lemma is along similar lines as Lemma 5.3 except that we use the fact that $\text{Ext}_{G_n}^*(\pi_1, D(\pi_1)) \neq 0$ instead of Equation 5.6 in the proof of the Lemma 5.3. The fact that $\text{Ext}_{G_n}^*(\pi_1, D(\pi_1)) \neq 0$ is a simple consequence of the duality theorem (Theorem 2.1) of Nori and Prasad.

5.3. A reformulation. We can reformulate Lemmas 5.2, 5.3 and 5.4 into a single statement in the language of Arthur parameters as follows. Let ΔSL_2 denote the diagonally embedded SL_2 sitting inside $\text{SL}_2 \times \text{SL}_2$. Note that if π_1 and π_2 are irreducible Arthur type representations of G_n then, $\mathcal{A}(\pi_1)|_{W_F \times \Delta \text{SL}_2} \cong \mathcal{A}(\pi_2)|_{W_F \times \Delta \text{SL}_2}$ if and only if π_1 and π_2 have the same cuspidal support.

Lemma 5.5. Let π_1 be a unitary segment type representation of G_n and let $\tau \in \text{Alg}(G_k)$ (not necessarily of finite length). Let $\pi_2 \in \text{Alg}(G_{n+k})$ be an irreducible unitary representation of Arthur type. Suppose that $\pi_2 = \pi_{2,1} \times \pi_{2,2} \times \dots \times \pi_{2,s}$ where, $\pi_{2,1}, \pi_{2,2}, \dots, \pi_{2,s}$ are unitary representations of segment type. Then,

$$\text{Ext}_{G_{n+k}}^*(\pi_1 \times \tau, \pi_2) \neq 0$$

if and only if π_2 decomposes as, $\pi_2 = \pi_2' \times \pi_2''$ where, $\pi_2' \in \text{Alg}(G_n)$ and $\pi_2'' \in \text{Alg}(G_k)$ such that,

$$\mathcal{A}(\pi_1)|_{W_F \times \Delta \text{SL}_2} \cong \mathcal{A}(\pi_2')|_{W_F \times \Delta \text{SL}_2}$$

and,

$$\text{Ext}_{G_k}^*(\tau, \pi_2'') \neq 0.$$

5.4. A conjecture about extensions on the same group. Since we are dealing with questions about extensions on the same group in this section, it is natural to ask when non-trivial extensions exist between Arthur type representations on the same group. The following conjecture arose in an email correspondence between Professor Dipendra Prasad and Professor K.Y. Chan.

Conjecture 5.1. Suppose π_1 and π_2 are irreducible Arthur type representations of $\text{GL}_n(F)$. Then $\text{Ext}_{\text{GL}_n(F)}^*(\pi_1, \pi_2) \neq 0$ if and only if

$$\mathcal{A}(\pi_1)|_{W_F \times \Delta \text{SL}_2} \cong \mathcal{A}(\pi_2)|_{W_F \times \Delta \text{SL}_2}.$$

Remark 5.4. Note that as observed earlier, saying that

$$\mathcal{A}(\pi_1)|_{W_F \times \Delta \text{SL}_2} \cong \mathcal{A}(\pi_2)|_{W_F \times \Delta \text{SL}_2}$$

is the same as saying that π_1 and π_2 have the same cuspidal support.

By repeated applications of Lemma 5.5 it follows that the above conjecture is true in the following special case.

Proposition 5.1. *Suppose π_1 and π_2 are irreducible Arthur type representations of $\mathrm{GL}_n(F)$. Moreover, suppose that π_1 and π_2 are products of unitary representations of segment type. Then $\mathrm{Ext}_{\mathrm{GL}_n(F)}^*(\pi_1, \pi_2) \neq 0$ if and only if*

$$\mathcal{A}(\pi_1)|_{W_F \times \Delta \mathrm{SL}_2} \cong \mathcal{A}(\pi_2)|_{W_F \times \Delta \mathrm{SL}_2}.$$

6. MAIN REDUCTION LEMMA

The following lemma is Lemma 4.3 from [7] where it is stated for Hom spaces but which works as well for Ext modules as we assert below. This lemma follows from an application of the filtration stated in Theorem 4.1. It allows us to reduce the study of Ext branching to the bottom piece of the aforementioned filtration. Since this lemma plays an important role in our arguments we recall its proof for the sake of completeness.

Lemma 6.1. Let π_1 and π_2 be Arthur type representations of G_n and G_{n-1} respectively. Suppose that

$$\pi_1 = u_{\rho_1}(a_1, b_1) \times u_{\rho_2}(a_2, b_2) \times \dots \times u_{\rho_r}(a_r, b_r)$$

and

$$\pi_2 = u_{\tau_1}(c_1, d_1) \times u_{\tau_2}(c_2, d_2) \times \dots \times u_{\tau_s}(c_s, d_s)$$

where ρ_i ($i = 1, 2, \dots, r$) and τ_j ($j = 1, 2, \dots, s$) are some unitary cuspidal representations. Moreover, let us suppose that $a_1 + b_1 \geq a_i + b_i$ and $a_1 + b_1 \geq c_j + d_j$ for all $i = 1, 2, \dots, r$ and $j = 1, 2, \dots, s$. Let σ be a unitary cuspidal representation of $G_{a_1 n(\rho_1)}$ such that $\nu^{1/2}\sigma \notin \mathrm{csupp}_{\mathbb{Z}}(\pi_2)$ and $\sigma \notin \mathrm{csupp}_{\mathbb{Z}}(\pi_1)$. Then for all integers $i \geq 0$ we have that,

$$\mathrm{Ext}_{G_{n-1}}^i(\pi_1, \pi_2) = \mathrm{Ext}_{G_{n-1}}^i(u_{\rho_1}(a_1, b_1 - 1) \times (\sigma \times \pi'_1)|_{G_a}, \pi_2)$$

where, $a = n - n(\rho_1)a_1(b_1 - 1) - 1$ and $\pi'_1 = u_{\rho_2}(a_2, b_2) \times \dots \times u_{\rho_r}(a_r, b_r)$.

Proof. By Theorem 4.1 we have that $\pi_1|_{G_{n-1}}$ is glued together from the pieces,

$$V_0/V_1 \cong \nu^{1/2}u_{\rho_1}(a_1, b_1) \times \pi'_1|_{G_{n(\pi'_1)-1}},$$

for $1 \leq k \leq d$,

$$V_k/V_{k+1} \cong \nu^{1/2}u_{\rho_1}(a_1, b_1)^{(k)} \times (\Pi_k \bar{\times} \pi'_1)|_{G_{n(\pi'_1)+k-1}}.$$

Here $d = \mathrm{lev}(u_{\rho_1}(a_1, b_1))$ is the largest integer ℓ such that $u_{\rho_1}(a_1, b_1)^{(\ell)} \neq 0$. By Lemma 3.1 we know that $d = n(\rho_1)a_1$.

Claim: We claim that among all the pieces in the above filtration the pieces other than the last piece do not contribute to the desired Ext module, that is for all integers $i \geq 0$,

$$\mathrm{Ext}_{G_{n-1}}^i(V_k/V_{k+1}, \pi_2) = 0$$

for all $0 \leq k < d$.

Proof of the Claim: The proof of the claim is based on a simple observation about the cuspidal support of derivatives of Speh representations. If $S = u_{\rho}(m, n)$ is a Speh representation then $\nu^{(m+n-2)/2}\rho \in \mathrm{csupp}(S^{(k)})$ provided $k < \mathrm{lev}(S)$, that is if $S^{(k)}$ is not

the highest derivative of S (Lemma 3.1). Suppose that $k < d = \text{lev}(u_{\rho_1}(a_1, b_1))$. Now by definition

$$\text{Ext}_{G_{n-1}}^i(V_k/V_{k+1}, \pi_2) = \text{Ext}_{G_{n-1}}^i(\nu^{1/2}u_{\rho_1}(a_1, b_1)^{(k)} \times N_k, \pi_2)$$

where $N_k = (\Pi_k \bar{\times} \pi'_1)|_{G_{n(\pi'_1)+k-1}}$. The inequalities $a_1 + b_1 \geq a_i + b_i$ and $a_1 + b_1 \geq c_j + d_j$ imply that $1/2 + (a_1 + b_1 - 2)/2$ is *strictly* greater than any element of the set,

$$S = \{(a_i + b_i - 2)/2, (c_j + d_j - 2)/2 | i = 1, 2, \dots, r; j = 1, 2, \dots, s\}.$$

So if $k < \text{lev}(u_{\rho_1}(a_1, b_1))$ we conclude that $\nu^{1/2+(a_1+b_1-2)/2}\rho_1$ lies in $\text{csupp}(u_{\rho_1}(a_1, b_1)^{(k)})$ but does not lie in the cuspidal support of π_2 . Therefore by second adjointness and comparing cuspidal supports at $\nu^{1/2+(a_1+b_1-2)/2}\rho_1$ (see Lemma 2.3) we conclude that,

$$\text{Ext}_{G_{n-1}}^i(V_k/V_{k+1}, \pi_2) = 0$$

for all $0 \leq k < d$. This completes the proof of the claim.

Since the claim is true, a long exact sequence argument implies that for all integers $i \geq 0$,

$$\begin{aligned} \text{Ext}_{G_{n-1}}^i(\pi_1, \pi_2) &= \text{Ext}_{G_{n-1}}^i(V_d/V_{d+1}, \pi_2) \\ &= \text{Ext}_{G_{n-1}}^i(u_{\rho_1}(a_1, b_1)^- \times (\Pi_d \bar{\times} \pi'_1)|_{G_a}, \pi_2). \end{aligned}$$

Here, $u_{\rho_1}(a_1, b_1)^- = \nu^{1/2}u_{\rho_1}(a_1, b_1)^{(d)} = u_{\rho_1}(a_1, b_1 - 1)$. Since $\nu^{1/2}\sigma \notin \text{csupp}_{\mathbb{Z}}(\pi_2)$ and $\sigma \notin \text{csupp}_{\mathbb{Z}}(\pi_1)$ by Lemma 4.2 it follows that,

$$\text{Ext}_{G_{n-1}}^i(\pi_1, \pi_2) = \text{Ext}_{G_{n-1}}^i(u_{\rho_1}(a_1, b_1) \times (\sigma \times \pi'_1)|_{G_a}, \pi_2).$$

□

7. PROOF OF ONE DIRECTION OF THEOREM 1.2

In this Section we shall prove one direction of Theorem 1.2. More precisely, we prove the following.

Theorem 7.1. *Let $n \geq 1$ be an integer. Let π_1 and π_2 be irreducible Arthur type representations of $\text{GL}_n(F)$ and $\text{GL}_{n-1}(F)$ respectively. Suppose that π_1 and π_2 are products of discrete series representations and the Aubert-Zelevinsky duals of discrete series representations. If $\text{Ext}_{\text{GL}_{n-1}(F)}^i(\pi_1, \pi_2) \neq 0$ for some integer $i \geq 0$, then π_1 and π_2 are strong Ext relevant.*

Proof. Suppose $\pi_1 = \pi_{1,1} \times \pi_{1,2} \times \dots \times \pi_{1,r}$ and $\pi_2 = \pi_{2,1} \times \pi_{2,2} \times \dots \times \pi_{2,s}$. Let us denote $\pi_{1,i} = u_{\rho_i}(a_i, b_i)$ and $\pi_{2,j} = u_{\tau_j}(c_j, d_j)$ where the ρ_i ($i = 1, 2, \dots, r$) and τ_j ($j = 1, 2, \dots, s$) are some cuspidal representations. We are given that $\pi_{1,1}, \pi_{1,2}, \dots, \pi_{1,r}$ and $\pi_{2,1}, \pi_{2,2}, \dots, \pi_{2,s}$ are all of segment type. This means that for all $i = 1, 2, \dots, r$ either $a_i = 1$ or $b_i = 1$. Similarly, for all $j = 1, 2, \dots, s$ either $c_j = 1$ or $d_j = 1$. We know that $\text{Ext}_{G_{n-1}}^i(\pi_1, \pi_2) \neq 0$ for some integer $i \geq 0$. We want to show that the pair (π_1, π_2) is strong Ext relevant.

Let $m(\pi_1, \pi_2)$ denote the number of non-cuspidal factors of π_1 and π_2 . The proof of the Theorem is via induction on $m(\pi_1, \pi_2)$. When $m(\pi_1, \pi_2) = 0$ then both π_1 and π_2 are generic and hence automatically strong Ext relevant. We now assume that Theorem 7.1 holds for all integers less than $m(\pi_1, \pi_2)$.

Since π_1 and π_2 are irreducible we can permute their factors and ensure that one of the following cases occurs. We can ensure that either:

- (1) $a_1 + b_1 \geq a_i + b_i$ and $a_1 + b_1 \geq c_j + d_j$ for all i and j .
- (2) $c_1 + d_1 \geq c_j + d_j$ and $c_1 + d_1 \geq a_i + b_i$ for all i and j .

We deal with the above two cases one by one.

Case 1: Let us suppose that $a_1 + b_1 \geq a_i + b_i$ and $a_1 + b_1 \geq c_j + d_j$ for all i and j . Choose a unitary cuspidal representation σ of G_2 such that $\nu^{1/2}\sigma \notin \text{csupp}_{\mathbb{Z}}(\pi_2)$ and $\sigma \notin \text{csupp}_{\mathbb{Z}}(\pi_1)$.

Since $\text{Ext}_{G_{n-1}}^i(\pi_1, \pi_2) \neq 0$ by Lemma 6.1 we have that,

$$\text{Ext}_{G_{n-1}}^i(u_{\rho_1}(a_1, b_1 - 1) \times (\sigma \times \pi'_1)|_{G_a}, \pi_2) \neq 0$$

where, $a = n - n(\rho_1)a_1(b_1 - 1) - 1$ and $\pi'_1 = u_{\rho_2}(a_2, b_2) \times \dots \times u_{\rho_r}(a_r, b_r)$. Note that $\pi_{1,1}^- = u_{\rho_1}(a_1, b_1)^- = u_{\rho_1}(a_1, b_1 - 1)$ is also of segment type.

We now invoke Lemma 5.2 and suitably rearrange the factors of π_2 to conclude the following.

- (1) $\pi_{2,1}$ is equal to either $\pi_{1,1}^-$ or $D(\pi_{1,1}^-)$. Therefore, the pair $(\pi_{1,1}, \pi_{2,1})$ is strong Ext relevant.
- (2) We have that $\text{Ext}_{G_a}^i(\sigma \times \pi'_1, \pi_{2,2} \times \dots \times \pi_{2,s}) \neq 0$. So by the induction hypothesis the pair $(\sigma \times \pi'_1, \pi_{2,2} \times \dots \times \pi_{2,s})$ is strong Ext relevant. Since $\nu^{1/2}\sigma \notin \text{csupp}_{\mathbb{Z}}(\pi_2)$, the pair $(\pi'_1, \pi_{2,2} \times \dots \times \pi_{2,s})$ is strong Ext relevant.

From the above two observations we conclude that the pair π_1 and π_2 are strong Ext relevant.

Case 2: Let us suppose that $c_1 + d_1 \geq c_j + d_j$ and $a_1 + b_1 \geq c_i + d_i$ for all i and j . Then we can apply the transfer lemma (Lemma 4.1) and deal with the pair $(\pi_2^\vee \times \tilde{\sigma}, \pi_1^\vee)$ rather than the pair (π_1, π_2) where, $\tilde{\sigma}$ is some cuspidal representation of G_2 chosen such that $\nu^{1/2}\tilde{\sigma} \notin \text{csupp}_{\mathbb{Z}}(\pi_1^\vee)$ and $\tilde{\sigma} \notin \text{csupp}_{\mathbb{Z}}(\pi_2^\vee)$. Since, $\tilde{\sigma} \notin \text{csupp}_{\mathbb{Z}}(\nu^{-1/2}\pi_1^\vee)$ observe that the pair (π_1, π_2) is strong Ext relevant if and only if the pair $(\pi_2^\vee \times \tilde{\sigma}, \pi_1^\vee)$ is strong Ext relevant. When dealing with the pair $(\pi_2^\vee \times \tilde{\sigma}, \pi_1^\vee)$ the role of $\pi_{1,1}$ is replaced by $\pi_{2,1}$. Hence it is enough to deal with Case 1. \square

8. PROOF OF OTHER DIRECTION OF THEOREM 1.2

We now prove the other direction of Theorem 1.2.

Theorem 8.1. *Let π_1 and π_2 be irreducible Arthur type representations of $\text{GL}_n(F)$ and $\text{GL}_{n-1}(F)$ respectively. Suppose that π_1 and π_2 are products of discrete series representations and the Aubert-Zelevinsky duals of discrete series representations. If π_1 and π_2 are strong Ext relevant then, $\text{Ext}_{\text{GL}_{n-1}(F)}^i(\pi_1, \pi_2) \neq 0$ for some integer $i \geq 0$.*

Proof. We set up notations as in the previous section. Suppose $\pi_1 = \pi_{1,1} \times \pi_{1,2} \times \dots \times \pi_{1,r}$ and $\pi_2 = \pi_{2,1} \times \pi_{2,2} \times \dots \times \pi_{2,s}$, where $\pi_{1,1}, \pi_{1,2}, \dots, \pi_{1,r}$ and $\pi_{2,1}, \pi_{2,2}, \dots, \pi_{2,s}$ are unitary representations of segment type. Let us denote $\pi_{1,i} = u_{\rho_i}(a_i, b_i)$ and $\pi_{2,j} = u_{\tau_j}(c_j, d_j)$ where the ρ_i ($i = 1, 2, \dots, r$) and τ_j ($j = 1, 2, \dots, s$) are some cuspidal representations.

We are given that π_1 and π_2 are strong Ext relevant. Let $m(\pi_1, \pi_2)$ denote the number of non-cuspidal factors of π_1 and π_2 . As before, the proof of the Theorem is via induction on $m(\pi_1, \pi_2)$. When $m(\pi_1, \pi_2) = 0$ then both π_1 and π_2 are generic and by Theorem 2.3

we conclude that $\text{Hom}_{G_{n-1}}(\pi_1, \pi_2) \neq 0$. We now assume that the Theorem 8.1 holds for all integers less than $m(\pi_1, \pi_2)$.

Arguing similarly as in Case 2 of the proof of Theorem 7.1, it is enough to deal with the case that $a_1 + b_1 \geq a_i + b_i$ and $a_1 + b_1 \geq c_j + d_j$ for all i and j . Choose a unitary cuspidal representation σ of G_2 such that $\nu^{1/2}\sigma \notin \text{csupp}_{\mathbb{Z}}(\pi_2)$ and $\sigma \notin \text{csupp}_{\mathbb{Z}}(\pi_1)$. Then by Lemma 6.1 we have that for all integers $i \geq 0$,

$$\text{Ext}_{G_{n-1}}^i(\pi_1, \pi_2) = \text{Ext}_{G_{n-1}}^i(\pi_{1,1}^- \times (\sigma \times \pi_1')|_{G_a}, \pi_2) \quad (8.1)$$

where, $a = n - n(\rho_1)a_1(b_1 - 1) - 1$ and $\pi_1' = u_{\rho_2}(a_2, b_2) \times \dots \times u_{\rho_r}(a_r, b_r)$. Here $\pi_{1,1}^- = u_{\rho_1}(a_1, b_1)^- = u_{\rho_1}(a_1, b_1 - 1)$.

Now since the representations π_1 and π_2 are strong Ext relevant one of the following two cases must occur.

- (A) $\pi_{2,m} = \pi_{1,1}^-$ for some $m = 1, 2, \dots, s$. In this case $\pi_2 = \pi_{1,1}^- \times \pi_2'$ where, $\pi_2' = \pi_{2,1} \times \dots \times \pi_{2,m-1} \times \pi_{2,m+1} \times \dots \times \pi_{2,s}$. In this case, the representations π_1' and π_2' are strong Ext relevant.
- (B) $\pi_{2,m} = D(\pi_{1,1}^-)$ for some $m = 1, 2, \dots, s$. In this case $\pi_2 = \pi_{1,1}^- \times \pi_2''$ where, $\pi_2'' = \pi_{2,1} \times \dots \times \pi_{2,m-1} \times \pi_{2,m+1} \times \dots \times \pi_{2,s}$. In this case, the representations π_1' and π_2'' are strong Ext relevant.

Let us suppose that Case (A) above occurs, that is, $\pi_2 = \pi_{1,1}^- \times \pi_2'$, and the representations π_1' and π_2' are strong Ext relevant. Since π_1' and π_2' are strong Ext relevant we have that the representations $\sigma \times \pi_1'$ and π_2' are strong Ext relevant. By the induction hypothesis we conclude that,

$$\text{Ext}_{G_a}^*(\sigma \times \pi_1', \pi_2') \neq 0.$$

By Lemma 5.3 and Equation 8.1 we conclude that

$$\text{Ext}_{G_{n-1}}^i(\pi_1, \pi_2) \neq 0$$

for some integer $i \geq 0$.

On the other hand suppose that Case (B) occurs. In this case, $\pi_2 = D(\pi_{1,1}^-) \times \pi_2''$, and the representations π_1' and π_2'' are strong Ext relevant. The proof proceeds exactly as in Case (A) but we invoke Lemma 5.4 instead of Lemma 5.3. □

ACKNOWLEDGEMENTS

The author would like to thank Professor Dipendra Prasad for his comments on the paper and for helpful discussions especially on Definition 1.1 of strong Ext relevance. The author would also like to thank Professor K.Y. Chan for helpful communications through Professor Dipendra Prasad.

REFERENCES

- [1] Avraham Aizenbud, Dmitry Gourevitch, Stephen Rallis, and Gérard Schiffmann. Multiplicity one theorems. *Annals of Mathematics*, pages 1407–1434, 2010.
- [2] Avraham Aizenbud and Eitan Sayag. Homological multiplicities in representation theory of p-adic groups. *Mathematische Zeitschrift*, 294:451–469, 2020.
- [3] I. Bernshtein and A. Zelevinskiy. Representations of the group $\text{GL}(n, F)$ where F is a non-archimedean local field. *Russian Mathematical Surveys*, 31(3):1, 1976.

- [4] I. Bernstein and A. Zelevinsky. Induced representations of reductive p -adic groups. I. In *Annales scientifiques de l'École Normale Supérieure*, volume 10, pages 441–472, 1977.
- [5] Kei Yuen Chan. On the product functor on inner forms of general linear group over a non-archimedean local field. *arXiv preprint arXiv:2206.10928*, 2022.
- [6] Kei Yuen Chan. Quotient branching law for p -adic $(\mathrm{GL}_{n+1}, \mathrm{GL}_n)$ I: generalized Gan-Gross-Prasad relevant pairs. *arXiv preprint arXiv:2212.05919*, 2022.
- [7] Kei Yuen Chan. Restriction for general linear groups: the local non-tempered Gan–Gross–Prasad conjecture (non-Archimedean case). *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2022(783):49–94, 2022.
- [8] Kei Yuen Chan. Ext-multiplicity theorem for standard representations of $(\mathrm{GL}_{n+1}, \mathrm{GL}_n)$. *Mathematische Zeitschrift*, 303(2):45, 2023.
- [9] Chan, Kei Yuen and Savin, Gordan. A vanishing Ext-branching theorem for $(\mathrm{GL}_{n+1}(\mathbb{F}), \mathrm{GL}_n(\mathbb{F}))$. *Duke Mathematical Journal*, 170(10):2237–2261, 2021.
- [10] Wee Teck Gan, Benedict H Gross, and Dipendra Prasad. Symplectic local root numbers, central critical L -values, and restriction problems in the representation theory of classical groups. *Astérisque*, pages 1–109, 2011.
- [11] Wee Teck Gan, Benedict H Gross, and Dipendra Prasad. Branching laws for classical groups: the non-tempered case. *Compositio Mathematica*, 156(11):2298–2367, 2020.
- [12] Maxim Gurevich. On restriction of unitarizable representations of general linear groups and the non-generic local Gan–Gross–Prasad conjecture. *Journal of the European Mathematical Society*, 24(1):265–302, 2021.
- [13] Hervé Jacquet, Ilya I Piatetskii-Shapiro, and Joseph A Shalika. Rankin-selberg convolutions. *American journal of mathematics*, 105(2):367–464, 1983.
- [14] Arno Kret and Erez Lapid. Jacquet modules of ladder representations. *Comptes Rendus Mathématique*, 350(21-22):937–940, 2012.
- [15] Erez Lapid and Alberto Mínguez. On a determinantal formula of Tadić. *American Journal of Mathematics*, 136(1):111–142, 2014.
- [16] Madhav Nori and Dipendra Prasad. On a duality theorem of Schneider–Stuhler. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2020(762):261–280, 2020.
- [17] Sascha Orlik. On extensions of generalized Steinberg representations. *Journal of Algebra*, 293(2):611–630, 2005.
- [18] Dipendra Prasad. On the decomposition of a representation of $\mathrm{GL}(3)$ restricted to $\mathrm{GL}(2)$ over a P -adic field. *Duke Math. J.*, 72(1):167–177, 1993.
- [19] Dipendra Prasad. Ext-analogues of branching laws. In *Proceedings of the International Congress of Mathematicians: Rio de Janeiro 2018*, pages 1367–1392. World Scientific, 2018.
- [20] Marko Tadić. Classification of unitary representations in irreducible representations of general linear group (non-archimedean case). In *Annales scientifiques de l'École normale supérieure*, volume 19, pages 335–382, 1986.
- [21] A. Zelevinsky. Induced representations of reductive p -adic groups. II. on irreducible representations of $\mathrm{GL}(n)$. In *Annales scientifiques de l'École Normale Supérieure*, volume 13, pages 165–210. Elsevier, 1980.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, MUMBAI 400076
 Email address: 185090012@iitb.ac.in