

Growth Rate of the Number of Empty Triangles in the Plane

Bhaswar B. Bhattacharya¹, Sandip Das², Sk Samim Islam², and Saumya Sen²

¹ University of Pennsylvania, USA

² Indian Statistical Institute, Kolkata, India

Abstract. Given a set P of n points in the plane, in general position, denote by $N_{\Delta}(P)$ the number of empty triangles with vertices in P . In this paper we investigate by how much $N_{\Delta}(P)$ changes if a point x is removed from P . By constructing a graph $G_P(x)$ based on the arrangement of the empty triangles incident on x , we transform this geometric problem to the problem of counting triangles in the graph $G_P(x)$. We study properties of the graph $G_P(x)$ and, in particular, show that it is kite-free. This relates the growth rate of the number of empty triangles to the famous Ruzsa-Szemerédi problem.

Keywords: Discrete geometry · Empty triangles · Kite-free graph.

1 Introduction

Let P be a set of n points in the plane in general position, that is, no three are on a line. We define $N_{\Delta}(P)$ as the number of empty triangles in P , that is, the number of triangles with vertices in P with no other point of P in the interior. Counting the number of empty triangles in planar point sets is a classical problem in discrete geometry (see [1–3, 5, 6, 8, 10, 11, 14] and the references therein). Specifically, Bárány and Füredi [1] showed that $N_{\Delta}(P) \geq n^2 - O(\log n)$, for any set of points P , with $|P| = n$, in general position. On the other hand, a set of n points chosen uniformly and independently at random from a convex set of area 1 contains $2n^2 + o(n^2)$ empty triangles on expectation [11, 14].


In this paper we study the growth rate of $N_{\Delta}(P)$ when a point x is removed from P . For this, let $N_{\Delta}(P \setminus \{x\})$ denote the number of empty triangles in the set $P \setminus \{x\}$ and consider the difference:

$$\Delta(x, P) = |N_{\Delta}(P) - N_{\Delta}(P \setminus \{x\})|.$$

In the following theorem we bound the above difference in terms of number of triangles in P with x as a vertex, which we denote by $V_P(x)$. To this end, denote by $K_4 \setminus \{e\}$ the *kite graph*, that is, the complete graph K_4 with one of its diagonals removed (see Figure 1).

Theorem 1. *For any set P , with $|P| = n$,*

$$\Delta(x, P) \leq V_P(x) + H(V_P(x), K_3, K_4 \setminus \{e\}), \quad (1)$$

Fig. 1: The kite graph $K_4 \setminus \{e\}$.

where $H(V_P(x), K_3, K_4 \setminus \{e\})$ is the maximum number of triangles in a $K_4 \setminus \{e\}$ -free graph on $V_P(x)$ vertices. Moreover, there exists a set P , with $|P| = n$, and a point $x \in P$ such that $\Delta(x, P) \geq CV_P(x)^{\frac{3}{2}}$, for some constant $C > 0$.

The proof of Theorem 1 is given in Section 2. To establish the upper bound in (1) we construct a graph $G_P(x)$ based on the arrangement of empty triangles in P which have x as a vertex and relate the problem of estimating $\Delta(x, P)$ to the problem of counting the number of triangles in the graph $G_P(x)$. The graph $G_P(x)$ has many interesting properties, specifically, we show that it is kite-free, which gives the upper bound in (1) (see Section 2.1). The lower bound construction is given in Section 2.2.

The extremal number $H(V_P(x), K_3, K_4 \setminus \{e\})$ that appear in the bound (1) is closely connected to the celebrated Ruzsa-Szemerédi problem, which asks for the maximum number of edges in a graph with n vertices such that every edge belongs to a unique triangle [12]. Note that if a graph G is kite-free, then every edge in G has at most one triangle passing through it. Then removing the edges which have no triangles passing through them (which does not change the number of triangles), one can relate the problem of counting triangles in G to the Ruzsa-Szemerédi problem. An application of the Szemerédi regularity lemma [13] shows that any solution to the Ruzsa-Szemerédi problem has at most $o(n^2)$ edges [12] (which is often also referred to as the diamond-free lemma). This can be improved to $n^2/e^{\Omega(\log^* n)}$ by a stronger form of the graph removal lemma [7]. This, in particular, implies that $H(V_P(x), K_3, K_4 \setminus \{e\}) = o(V_P(x)^2)$. Hence, from Theorem 1 we get the upper bound: $\Delta(x, P) = o(V_P(x)^2)$. On the other hand, the lower bound of $\Delta(x, P)$ from Theorem 1 is $\Omega(V_P(x)^{\frac{3}{2}})$. We believe the correct order of magnitude of $\Delta(x, P)$ is closer to the lower bound, because the graph $G_P(x)$ has additional geometric structure. We collect some geometric properties of the graph $G_P(x)$ in Section 3, which we believe can be of independent interest.

2 Proof of Theorem 1

We prove the upper bound in (1) in Section 2.1. The lower bound construction is given in Section 2.2.

2.1 Proof of the Upper Bound

We begin with the following simple observation:

Observation 1 $\Delta(x, P) \leq V_P(x) + I_P(x)$, where $I_P(x)$ is the number of triangles in P that contain only the point x in the interior.

Proof. Let $U_P(x)$ denote the number of empty triangles in P such that x is not a vertex of the empty triangles. Note that $N_\Delta(P) = V_P(x) + U_P(x)$ and $N_\Delta(P \setminus \{x\}) = U_P(x) + I_P(x)$. This implies, $|N_\Delta(P) - N_\Delta(P \setminus \{x\})| = |V_P(x) - I_P(x)| \leq V_P(x) + I_P(x)$. \square

Given a set P , with $|P| = n$, and a point $x \in P$, define the graph $G_P(x)$ as follows: The vertex set of $G_P(x)$ is $V(G_P(x))$, the set of triangles in P with x as one of their vertices, and there should be edge between 2 vertices in $G_P(x)$ if the corresponding triangles, say T_1 and T_2 , satisfy the following conditions:

- T_1 and T_2 share an edge,
- T_1 and T_2 are area disjoint,
- the sum of angles of T_1 and T_2 incident at x is greater than 180° .

We call the graph $G_P(x)$ the *empty triangle graph incident at x* . Figure 2 shows the graph $G_P(x)$ for a set of 4 points $P = \{x, a, b, c\}$. (It is worth noting that we use Δ in notations that count empty triangles in the point set P and K_3 to denote a triangle in the graph $G_P(x)$.)

The following lemma shows that $I_P(x)$, as defined in Observation 1, is equal to the number of triangles in $G_P(x)$.

Lemma 1. *Suppose P be a set of points in the plane, with $|P| = n$, in general position and $x \in P$. Then*

$$I_P(x) = N_{K_3}(G_P(x)),$$

where $N_{K_3}(G_P(x))$ is the number of triangles in the graph $G_P(x)$.

Proof. Let (a, b, c) be a triangle in $I_P(x)$, that is, (a, b, c) only has the point x in the interior. This corresponds to a triangle in $G_P(x)$ as shown in Figure 2.

Now, consider a triangle with vertices labeled $(1, 2, 3)$ in $G_P(x)$. The vertices 1, 2, and 3 in $G_P(x)$ correspond to 3 empty triangles in P that have the point x as one of their vertices, which we denote by T_1, T_2 , and T_3 , respectively. Since there is an edge between 1 and 2 in $G_P(x)$, the triangles T_1 and T_2 in P share a common edge, are area disjoint, and the sum of angles of T_1 and T_2 incident at x is greater than 180° . Hence, we can assume, without loss of generality, T_1




Fig. 2: A set of points $P = \{x, a, b, c\}$ and the empty triangle graph incident at x .

and T_2 are arranged as in Figure 3(a). Similarly, the pairs of triangles (T_2, T_3) and (T_1, T_3) share a common edge, are area disjoint, and the sum of their angles incident at x is greater than 180° . This implies, $T_1, T_2,$ and T_3 are mutually area disjoint, and they cannot all share the same edge. Therefore, the triangles $T_1, T_2,$ and T_3 have to be arranged as in Figure 3(b). Note that the triangle formed by the union of the 3 triangles T_1, T_2 and T_3 only contains the point x in P . Hence, for every triangle $(1, 2, 3)$ in $G_P(x)$ one gets a triangle in $I_P(x)$, formed by the union of the 3 triangles T_1, T_2 and T_3 . \square




Fig. 3: Illustration for the proof of Lemma 1.

Applying Observation 1 and Lemma 1 it follows that

$$\Delta(x, P) \leq V_P(x) + I_P(x) = V_P(x) + N_{K_3}(G_P(x)). \tag{2}$$

Therefore, to prove the upper bound in (1), it remains to show that

$$N_{K_3}(G_P(x)) \leq H(V_P(x), K_3, K_4 \setminus \{e\}) \tag{3}$$

This follows from the lemma below which shows that the graph $G_P(x)$ is kite-free.

Lemma 2. *The graph $G_P(x)$ does not contain $K_4 \setminus \{e\}$ as a subgraph, that is, $G_P(x)$ is kite-free.*

Proof. Suppose $G_P(x)$ contains a kite $K_4 \setminus \{e\}$, with vertices labeled a, b, c, d as in Figure 1. This corresponds to empty triangles T_a, T_b, T_c, T_d with x as a vertex such that (T_a, T_b, T_c) and (T_a, T_b, T_d) are mutually interior disjoint, and the pairs of triangles (T_a, T_b) , (T_b, T_c) , (T_a, T_c) , (T_a, T_d) , and (T_b, T_d) share a common edge. This means the triangles T_a, T_b, T_c must be arranged as in Figure 2(a). Hence, it is impossible to place T_d which share an edge with T_a and T_b and is interior disjoint from T_a, T_b , unless T_d coincides with T_c . This shows $G_P(x)$ cannot contain a $K_4 \setminus \{e\}$ as a subgraph. \square




Fig. 4: Example showing the lower bound in Theorem 1.

Since $G_P(x)$ is kite-free by Lemma 2, the bound in (3) follows. This together with (2) gives the upper bound in (1).

2.2 Lower Bound Construction

To prove the lower bound in Theorem 1 consider the set of points P , with $|P| = n = 3L + 1$, as shown in Figure 4. Specifically, P consists of three sets points A , B , and C , with $|A| = |B| = |C| = L$, arranged along 3 disjoint convex chains and a point x at the middle. Note that $N_\Delta(P \setminus \{x\}) = \binom{3L}{3} \sim \frac{9}{2}L^3$. Also,

$$V_P(x) = 3 \binom{L}{2} + 3L^2 = \Theta(L^2). \quad (4)$$

To compute $N_\Delta(P)$ recall that $N_\Delta(P) = V_P(x) + U_P(x)$, where $U_P(x)$ is as defined in Observation 1. Now, note that

$$U_P(x) = 3 \binom{L}{3} + 6L \binom{L}{2} \sim 3.5L^3.$$

Hence, $N_\Delta(P) \sim 3.5L^3 + \Theta(L^2)$ and

$$\Delta(x, P) = |N_\Delta(P) - N_\Delta(P \setminus \{x\})| = \Theta(L^3) = \Theta(V_P(x)^{\frac{3}{2}}),$$

from (4). This completes the proof of the lower bound in Theorem 1.

3 Properties of the Graph $G_P(x)$

In this section we collect some geometric properties of the graph $G_P(x)$. First, we show that $G_P(x)$ can contain arbitrarily large bipartite graphs.

Lemma 3. *Fix $r, s \geq 1$. Then there exists a set of points P and $x \in P$ such that the graph $G_P(x)$ contains the complete bipartite graph $K_{r,s}$.*




Fig. 5: Illustration for the proof of Lemma 3.

Proof. Consider the set of $r + s + 2$ points

$$P = \{x, x', a_1, a_2, \dots, a_r, b_1, b_2, \dots, b_s\},$$

as shown in Figure 5(a) (with $r = 2$ and $s = 3$). Specifically, the points $\{a_1, a_2, \dots, a_r\}$ and $\{b_1, b_2, \dots, b_s\}$ lie on 2 disjoint convex chains and the points x and x' are in the middle. Note that, for all $1 \leq i \leq r$ and $1 \leq j \leq s$, the triangles $xx'a_i$ and $xx'b_j$ share a common edge, are interior disjoint, and the sum of the angles incident at x is greater than 180° . This implies that the graph $G_P(x)$ contains the complete bipartite graph $K_{r,s}$. \square

The reason it is worthwhile to know whether or not $\Delta(x, P)$ contains complete bipartite graphs as subgraphs, is because of a possible approach to improve the $o(n^2)$ upper bound on $\Delta(x, P)$ through the Kővári-Sós-Turán theorem [9]. Recall that the Kővári-Sós-Turán theorem states that any graph which is $K_{r,s}$ -free, where $r \leq s$, has at most $O(n^{2-\frac{1}{r}})$ edges. Therefore, if $\Delta(x, P)$ did not contain some complete bipartite graph as a subgraph, then it would have led to a polynomial improvement over the $o(n^2)$ upper bound on $\Delta(x, P)$. Lemma 3 shows that this is not the case, hence, one cannot directly apply the Kővári-Sós-Turán theorem to improve the upper bound on $\Delta(x, P)$.

Although the Kővári-Sós-Turán result cannot be directly applied to the graph $G_P(x)$, we believe a polynomial improvement over the Ruzsa-Szemerédi upper bound on $\Delta_P(x)$ is possible, because the graph $G_P(x)$ has additional geometric structure. To illustrate this we show that the well-known Behrend’s construction [4], which gives a nearly quadratic lower bound on Ruzsa-Szemerédi problem, is not geometric realizable. We begin recalling Behrend’s construction:

Definition 1 (Behrend’s graph). *Suppose p is an odd prime and $A \subseteq \mathbb{Z}/p\mathbb{Z}$ is a set with no 3-term arithmetic progression. The Behrend’s graph $G(p, A)$ is a tripartite graph with vertices on each side of the tripartition numbered $\{0, 1, \dots, p - 1\}$ and triangles of the form $(z, z + a, z + 2a)$ modulo p , for $z \in \{0, 1, \dots, p - 1\}$ and $a \in A$.*

It is easy to check that the graph $G(p, A)$ has $3p$ vertices $3|A|p$ edges and each edge belongs to a unique triangle. For example, when $p = 3$ and $A = \{1, 2\}$ one gets the 9 vertex Paley graph shown in Figure 6. Behrend [4] constructed a set A of size $p/e^{O(\sqrt{\log p})}$ with no 3-term arithmetic progression. Using this set in the Behrend’s graph in Definition 1 one gets a lower bound of $\Omega(p^2/e^{O(\sqrt{\log p})})$ for the Ruzsa-Szemerédi problem and, hence, for $H(n, K_3, K_4 \setminus \{e\})$.

The following result shows that the graph in Figure 6 cannot be geometrically realized, that is, it is not possible to find a set of points P and $x \in P$ such that $G_P(x)$ is isomorphic to the graph in Figure 6.

Proposition 1. *The graph in Figure 6 is not geometrically realizable.*

Proposition 1 shows that Behrend’s graphs are not geometrically realizable. This, in particular, illustrates that the graph $G_P(x)$ has a richer geometric structure than the collection of kite-free graphs.




Fig. 6: The Paley graph with 9 vertices, 18 edges, and 6 triangles.

3.1 Proof of Proposition 1

We proceed by contradiction. Suppose there exists a point set P and $x \in P$ such that $G_P(x)$ is isomorphic to the graph in Figure 6. This implies $V_P(x) = 9$ and $I_P(x) = N_{K_3}(G_P(x)) = 6$ (by Lemma 1). This, in particular, means that there are 6 triangles in P which only contains the point x in the interior. Denote these triangles by $\mathcal{T} = \{T'_1, T'_2, \dots, T'_6\}$.




Fig. 7: Illustration for the proofs of Lemma 4.

Lemma 4. *There cannot be 3 triangles in the set \mathcal{T} which share a common edge.*

Proof. Let, if possible, there exists 3 triangles T'_1, T'_2, T'_3 in \mathcal{T} which share one edge (see Figure 7(a)). As all the triangles contain x as the only interior point they must have a common shareable area that contains x . Therefore, it is not possible to create a new triangle here by joining vertices of existing triangles which will contain x . Therefore, to create a new triangle which will contain x as the only interior point, the following two cases may occur.

- *Case 1:* There is a triangle $T'_4 \in \mathcal{T}$ that does not share any edge and vertex with T'_1, T'_2, T'_3 (see Figure 8 (a)) or there is a triangle $T'_4 \in \mathcal{T}$ that does not share any edge with T'_1, T'_2, T'_3 but shares one vertex with T'_1, T'_2, T'_3 . In both case $V_P(x) \geq 10$, which is a contradiction.
- *Case 2:* There is a triangle $T'_4 \in \mathcal{T}$ which shares an edge with the triangles T'_1, T'_2, T'_3 . This means T'_4 has a vertex that is not common with the vertices of T'_1, T'_2, T'_3 (see Figure 8 (b)) In this case $V_P(x) = 9$ and $I_P(x) = N_{K_3}(G_P(x)) = 4$. Consider the triangle T'_5 in $\mathcal{T} \setminus \{T'_1, T'_2, T'_3, T'_4\}$. Note that, there is at least one edge in T'_5 which does not belong to the edge set of the triangles T'_1, T'_2, T'_3, T'_4 . This edge forms an empty triangle whose one vertex is x . This implies $V_P(x) \geq 10$, which is a contradiction.

Thus, if there are 3 triangles in the set \mathcal{T} which share an edge, then it is not possible to place another 3 triangles satisfying the required geometric constraints for the graph in Figure 6. Hence, there cannot be 3 triangle in \mathcal{T} which share an edge. \square




Fig. 8: Illustrations for the proofs of Lemma 4: (a) Case 1 and (b) Case 2.

Observe that each edge of a triangle in \mathcal{T} generates an empty triangle with vertex x . Since there are 9 empty triangles with vertex x in $G_P(x)$, the number of distinct edges generated by the triangles in \mathcal{T} will be 9. Moreover, by Lemma 4 no edge of a triangle in \mathcal{T} can be shared by three triangles of \mathcal{T} . Hence, each of the 9 edges generated by the triangles in \mathcal{T} must belong to exactly 2 triangles in \mathcal{T} (since, counting with repetitions, there are a total of $6 \times 3 = 18$ edges in the triangles in \mathcal{T}). The following lemma shows that this is not geometrically realizable.

Lemma 5. *All 3 edges of any triangle in \mathcal{T} cannot be shared by other triangles in \mathcal{T} .*

Proof. Let, if possible, there exists a triangle $T'_1 \in \mathcal{T}$ whose all 3 edges are shared by the triangles T'_2, T'_3, T'_4 and T'_2, T'_3, T'_4 do not share edges in between. (see Figure 9 (a)). In this case $V_P(x) = 9$ and $I_P(x) = N_{K_3}(G_P(x)) = 4$, which is impossible by arguments similar to Case 2 of Lemma 4.

Alternatively, suppose there exists a triangle $T'_1 = (a, b, c) \in \mathcal{T}$, such that the edge (a, b) is shared by the triangle T'_2 , the edge (b, c) is shared by the triangle T'_3 , the edge (c, a) is shared by the edge T'_4 , and T'_2 and T'_3 share a common edge. This implies, there is a vertex v such that $T'_2 = (a, b, v)$ and $T'_3 = (b, c, v)$ (see Figure 9 (b)). Note that v cannot be inside the triangle (a, b, c) , since (a, b, c) has only x as the interior point. Also, v cannot be in region A , region C , and region E , because then either the triangle T'_2 or the triangle T'_3 will have more than one point in the interior. Hence, v has to be in region B , region D , or region F . In this case, the triangle T'_2 and the triangle T'_3 are area disjoint, hence only one of them can contain x in the interior (see Figure 9 (b)). This gives a contradiction and completes the proof of the lemma. \square




Fig. 9: Illustrations for the proofs of Lemma 5.

Combining Lemma 4 and Lemma 5, the result in Proposition 1 follows.

4 Conclusions

In this paper, we initiate the study of the growth rate of the number of empty triangles in the plane, by proving upper and lower bounds on the difference $\Delta(x, P)$. We relate the upper bound to the well-known Ruzsa-Szemerédi problem and study geometric properties of the triangle incidence graph $G_P(x)$. Our results show that $\Delta(x, P)$ can range from $O(V_P(x)^{\frac{3}{2}})$ and $o(V_P(x)^2)$. Understanding additional properties of the graph $G_P(x)$ is an interesting future direction, which can be useful in improving the bounds on $\Delta(x, P)$.

References

1. Imre Bárány and Zoltán Füredi. Empty simplices in euclidean space. *Canadian Mathematical Bulletin*, 30(4):436–445, 1987.
2. Imre Bárány, Jean-François Marckert, and Matthias Reitzner. Many empty triangles have a common edge. *Discrete & Computational Geometry*, 50:244–252, 2013.
3. Imre Bárány and Pavel Valtr. Planar point sets with a small number of empty convex polygons. *Studia Scientiarum Mathematicarum Hungarica*, 41(2):243–269, 2004.
4. Felix A Behrend. On sets of integers which contain no three terms in arithmetical progression. *Proceedings of the National Academy of Sciences*, 32(12):331–332, 1946.
5. Peter Brass, William OJ Moser, and János Pach. *Research problems in discrete geometry*, volume 18. Springer, 2005.
6. Paul Erdos. On some unsolved problems in elementary geometry. *Mat. Lapok*, 2(2):1–10, 1992.
7. Jacob Fox. A new proof of the graph removal lemma. *Annals of Mathematics*, pages 561–579, 2011.
8. Alfredo García. A note on the number of empty triangles. In *Spanish Meeting on Computational Geometry*, pages 249–257. Springer, 2011.
9. Tamás Kővári, Vera T Sós, and Pál Turán. On a problem of Zarankiewicz. In *Colloquium Mathematicum*, volume 3, pages 50–57. Polska Akademia Nauk, 1954.
10. Rom Pinchasi, Radoš Radoičić, and Micha Sharir. On empty convex polygons in a planar point set. *Journal of Combinatorial Theory, Series A*, 3(113):385–419, 2006.
11. Matthias Reitzner and Daniel Temesvari. Stars of empty simplices. *arXiv preprint arXiv:1808.08734*, 2018.
12. Imre Z Ruzsa and Endre Szemerédi. Triple systems with no six points carrying three triangles. *Combinatorics (Keszthely, 1976), Coll. Math. Soc. J. Bolyai*, 18(939-945):2, 1978.
13. Endre Szemerédi. *Regular partitions of graphs*. Computer Science Department, Stanford University, 1975.
14. Pavel Valtr. On the minimum number of empty polygons in planar point sets. *Studia Scientiarum Mathematicarum Hungarica*, 30:155–163, 1995.