On co-dimension 2 defect anomalies in $\mathcal{N} = 4$ SYM and (2,0) theory via brane probes in AdS/CFT

Hongliang Jiang and Arkady A. Tseytlin¹

Blackett Laboratory, Imperial College, London SW7 2AZ, U.K.

E-mail: h.jiang@imperial.ac.uk, tseytlin@imperial.ac.uk

ABSTRACT: We consider a $\frac{1}{2}$ -BPS solution for a D3 brane probe in $\operatorname{AdS}_5 \times S^5$ that has worldvolume geometry of $\operatorname{AdS}_3 \times S^1$. It intersects the boundary over a surface that represents a dimension 2 defect in the boundary $\mathcal{N} = 4$ SYM theory. The effective action of the probe brane is proportional to the logarithmically divergent volume of AdS_3 and may thus be interpreted as computing conformal anomaly of supersymmetric S^2 defect. The classical action scales as N. We compute the 1-loop correction to it due to quantum fluctuations of the D3 brane world-volume fields and compare the result to an earlier suggested expression for the defect anomaly. We also perform a similar analysis of a $\frac{1}{2}$ -BPS M5 brane probe solution in $\operatorname{AdS}_7 \times S^4$ with the world-volume geometry of $\operatorname{AdS}_5 \times S^1$ that represents a dimension 4 defect in the boundary (2,0) 6d theory. Here the classical M5 brane action computes the leading order N^2 term in *a*-anomaly of the supersymmetric S^4 defect. We perform a detailed computation of the 1-loop correction to the M5 brane effective action and thus provide a prediction for the subleading constant in the S^4 defect *a*-anomaly coefficient.

¹Also at the Institute for Theoretical and Mathematical Physics (ITMP) of Moscow U. and Lebedev Institute.

Contents

 2 S² defect anomaly in N = 4 SYM from D3 probe in AdS₅ × S⁵ 2.1 Scalar fluctuations 2.2 Fermionic fluctuations 2.3 Vector field contribution 2.4 1-loop free energy 3 S⁴ defect anomaly in (2,0) theory from M5 probe in AdS₇ × S⁴ 3.1 Scalar fluctuations 3.2 Fermionic fluctuations 3.3 Antisymmetric tensor field contribution 3.4 1-loop free energy A Scalar fluctuations of a <i>p</i>-brane in AdS_{<i>p</i>+2} × S¹ B Spin connection and projected spinor covariant derivative C Fermion mass matrix for general u₀ D Supersymmetry of M5 embedding into AdS₇ × S⁴ 	1	Introduction	1
 3 S⁴ defect anomaly in (2,0) theory from M5 probe in AdS₇ × S⁴ 3.1 Scalar fluctuations 3.2 Fermionic fluctuations 3.3 Antisymmetric tensor field contribution 3.4 1-loop free energy A Scalar fluctuations of a p-brane in AdS_{p+2} × S¹ B Spin connection and projected spinor covariant derivative C Fermion mass matrix for general u₀ D Supersymmetry of M5 embedding into AdS₇ × S⁴ 	2	S^2 defect anomaly in $\mathcal{N} = 4$ SYM from D3 probe in $AdS_5 \times S^5$ 2.1Scalar fluctuations2.2Fermionic fluctuations2.3Vector field contribution2.41-loop free energy	3 5 6 9 10
 A Scalar fluctuations of a <i>p</i>-brane in AdS_{p+2} × S¹ B Spin connection and projected spinor covariant derivative C Fermion mass matrix for general u₀ D Supersymmetry of M5 embedding into AdS₇ × S⁴ 	3	 S⁴ defect anomaly in (2,0) theory from M5 probe in AdS₇ × S⁴ 3.1 Scalar fluctuations 3.2 Fermionic fluctuations 3.3 Antisymmetric tensor field contribution 3.4 1-loop free energy 	12 14 15 17 19
 B Spin connection and projected spinor covariant derivative C Fermion mass matrix for general u₀ D Supersymmetry of M5 embedding into AdS₇ × S⁴ 	\mathbf{A}	Scalar fluctuations of a p -brane in $\mathbf{AdS}_{p+2} \times S^1$	22
C Fermion mass matrix for general u_0 D Supersymmetry of M5 embedding into $AdS_7 \times S^4$	в	Spin connection and projected spinor covariant derivative	24
D Supersymmetry of M5 embedding into ${ m AdS}_7 imes S^4$	С	Fermion mass matrix for general u_0	25
	D	Supersymmetry of M5 embedding into $\mathbf{AdS}_7 imes S^4$	27

1 Introduction

Study of defects plays an important role in the investigation of properties of QFT's in various dimensions. In particular, it reveals more information about aspects of CFT's. In general, conformal defects are characterized by a set anomaly coefficients and OPE data determined by interplay of a defect with an ambient CFT (see, e.g., [1–8] and refs. there).

In the context of the AdS/CFT duality the properties of defects may be described in terms of brane probes intersecting the boundary of AdS on which the dual CFT lives. In particular, in [5] the subleading contribution to S^2 (co-dimension 4) defect anomaly in 6d (2,0) theory was computed by quantising M2 brane probe in AdS₇ × S^4 background. The induced geometry was AdS₃ and the classical and 1-loop correction were proportional to the log divergent volume of AdS₃ with S^2 as its boundary and thus determined the leading and subleading contributions to the defect *a*-anomaly.

Our aim below will be to perform similar computations in the case of co-dimension 2 spherical defects in 4d $\mathcal{N} = 4$ SYM and 6d (2,0) theories using the dual brane probe setup.

In the SYM case we will consider a $\frac{1}{2}$ -BPS supersymmetric D3 brane probe in AdS₅ × S⁵ wrapped on AdS₃ and S¹ in AdS₅ and also on S¹ in S⁵ with the resulting induced geometry being AdS₃ × S¹ [9–11].¹ Its effective action is proportional to vol(AdS₃) = $-2\pi \log(r\Lambda)$ and thus should capture the anomaly coefficient of an S² defect in $\mathcal{N} = 4$ SYM theory.

In the (2,0) theory case we will consider a $\frac{1}{2}$ -supersymmetric M5 brane probe in $\operatorname{AdS}_7 \times S^4$ which is wrapped on $S^1 \subset \operatorname{AdS}_5$ and $S^1 \subset S^4$ with the induced geometry being $\operatorname{AdS}_5 \times S^1$ [11] (cf. also [16]). The M5 brane effective action is then proportional to $\operatorname{vol}(\operatorname{AdS}_5) = \pi^2 \log(r\Lambda)$ and should capture the *a*-anomaly of an S^4 defect in A_{N-1} (2,0) theory (cf. [6]).

Let us note that a similar M2 brane probe solution in $\operatorname{AdS}_4 \times S^7/\mathbb{Z}_k$ determines the vortex defect expectation value in the ABJM theory [15] (where the induced geometry is $\operatorname{AdS}_2 \times S^1$ which has finite volume). Other similar 1-loop computations for M-branes in AdS backgrounds were recently discussed in [17–20]. One of our motivations here is to provide more examples when semiclassical quantization of supersymmetric branes in curved spaces leads to consistent results.

In all of these cases the spectrum of fluctuations on a *p*-brane brane embedded into $\operatorname{AdS}_{p+2} \times S^q$ (q = 8 - p or 9 - p) so that the world-volume metric is $\operatorname{AdS}_p \times S^1$ will contain 2 scalar modes corresponding to fluctuations in the transverse directions of AdS_{p+2} and also q-1 scalar fluctuations from S^q . All scalars propagating on $\operatorname{AdS}_p \times S^1$ will be conformally coupled.² In addition, the first two scalars will be mixed, or equivalently, coupled to an effective abelian constant gauge potential in S^1 direction. This will result in a shift of their S^1 mode number: $n \to n \pm \frac{1}{2}(p-1)$. The fermions will be massless in $\operatorname{AdS}_p \times S^1$ but also coupled to the same constant gauge potential (with half the charge) and thus having S^1 mode number shifted as $n \to n \pm \frac{1}{4}(p-1)$.³ In addition, there will fluctuations of the world-volume gauge fields propagating in $\operatorname{AdS}_p \times S^1$ geometry: vector in the D3 brane case and self-dual tensor in the M5 brane case.

Having found the fluctuation spectra we compute the 1-loop contribution to the corresponding effective action using the standard expressions for the determinants of the scalars, fermions and world-volume vector and antisymmetric tensor fields propagating on $\operatorname{AdS}_p \times S^1$ where p is odd in the present case (see [22–28]). The coefficient of the IR log divergent vol(AdS_p) factor determines the 1-loop contribution to the defect conformal anomaly. It can

³Similar spectrum for an M2 probe in $AdS_4 \times S^7 / \mathbb{Z}_k$ was found in [21] and used in [17, 15] (direct analogy with the present case is for k = 2 when the radii of AdS_2 and S^1 are equal).

¹In general, the AdS factor of the bulk geometry may be parametrized as $ds_{AdS_{p+2}}^2 = z^{-2}(dz^2 + dx^2 + x^2ds_{S^{p-1}}^2 + dy^2) = du^2 + \cosh^2 u \, ds_{AdS_p}^2 + \sinh^2 u \, d\psi^2$ where $ds_{AdS_p}^2 = d\rho^2 + \sinh^2 \rho \, ds_{S^{p-1}}^2$. The two metrics are related by $z = r/(\cosh u \cosh \rho - \sinh u \cos \psi)$, $x = z \cosh u \sinh \rho$, $y = z \sinh u \sin \psi$. When u or ρ goes to infinity, we have $z \to 0$, i.e. reach the \mathbb{R}^{p+1} boundary containing a co-dimension 2 defect S^{p-1} with radius r. It may be described in terms of a probe brane in $\mathrm{AdS}_{p+2} \times S^q$. For closely related discussions see [10, 12–15].

² Explicitly, their kinetic operator will be $-\nabla^2 + \frac{d-2}{4(d-1)}R$ with d = p + 1 and R of unit-radius $\operatorname{AdS}_p \times S^1$, i.e. $R = R(AdS_p) = -p(p-1)$. Expanding in Fourier modes in S^1 coordinate gives a tower of scalars on AdS_p with operators $-\nabla^2_{AdS_p} + m^2$, where $m^2 = n^2 + m_0^2 = n^2 - \frac{1}{4}(p-1)^2$. Assuming the Dirichlet boundary conditions, the dimension of the corresponding dual operators at the boundary of AdS_p is then $\Delta[\Delta - (p-1)] = m^2 = n^2 - \frac{1}{4}(p-1)^2$ or $\Delta - \frac{1}{2}(p-1) = |n|$.

be represented as an infinite sum over the S^1 mode number n. This sum happens to be finite in D3 case and quadratically divergent in the M5 case. In the latter case we use the standard Riemann ζ -function regularization to define it (like in similar examples of M-brane computations with AdS_p with even p discussed in [17, 18, 20]).

This paper is organized as follows. In section 2 we consider a D3 brane probe solution in $AdS_5 \times S^5$ with $AdS_3 \times S^1$ world volume metric that should be representing the supersymmetric S^2 defect in the boundary $\mathcal{N} = 4$ SYM. We first compute its classical action that gives the leading order N term in the defect anomaly. We then find the quadratic fluctuations of the probe brane fields near this brane configuration and find the value of the 1-loop correction to its effective action that contributes a finite constant term to the defect anomaly. The value of this constant that we find does not appear to match the expression suggested earlier in [2, 3].

In section 3 we perform a similar computation in the case of a supersymmetric M5 brane probe in $\operatorname{AdS}_7 \times S^4$ that should be related to an S^4 defect in 6d (2,0) theory. Here the worldvolume metric is $\operatorname{AdS}_5 \times S^1$ and the classical value of the action scales as N^2 . We compute the 1-loop correction to the effective action and regularizing it using the ζ -function get a prediction for the subleading constant term in the S^4 defect anomaly. It is not clear at the moment how to compare the result we found in the probe brane setup with the holographic computation of the S^4 defect anomaly in [8] which considered only the supergravity bubbling geometry approach where the numbers of all M5 branes are of the same order and thus the leading anomaly term superficially scales as N^3 . A special choice of the parameters of the solution which corresponds to a particular probe limit with leading scaling being N^2 remains to be understood.

Appendix A contains a general derivation of the scalar quadratic fluctuation Lagrangian for similar BPS brane configurations. The explicit form of the spin connection and the fermion covariant derivative are given in Appendix B. In Appendix C we show that like the scalar fluctuation action, the fermion mass matrix does not have a non-trivial dependence on the value of the "radial" position u_0 of the probe brane in AdS. The supersymmetry of the probe M5 brane solution studied in section 3 is demonstrated in Appendix D.

2 S² defect anomaly in $\mathcal{N} = 4$ SYM from D3 probe in AdS₅ × S⁵

Let us first consider the conformal anomaly associated with a surface defect in $\mathcal{N} = 4$ SYM from the dual AdS₅ × S⁵ perspective (see [12, 13]). We shall parametrize the AdS₅ × S⁵ background as (cf. footnote 1)

$$ds_{10}^2 = L^2 \left(du^2 + \cosh^2 u \, ds_{AdS_3}^2 + \sinh^2 u \, d\psi^2 \right) + L^2 \left(d\theta^2 + \sin^2 \theta \, d\phi^2 + \cos^2 \theta \, ds_{S^3}^2 \right) \,, \quad (2.1)$$

$$F_5 = dC_4 , \qquad C_4 = L^4 (\cosh^4 u - 1) \operatorname{vol}_{AdS_3} \wedge d\psi + \cdots , \qquad L^4 = 4\pi g_s N \alpha'^2 , \quad (2.2)$$

where in (2.2) dots stand for "magnetic" terms that make $F_5 = dC_4$ (anti-)self-dual. The bosonic part of the action for a D3 brane probe in this background is

$$S = -T_3 \left[\int d^4 \xi \, \sqrt{-\det(G_{\alpha\beta} + 2\pi\alpha' \mathcal{F}_{\alpha\beta})} - \int C_4 \right] \,, \qquad T_3 = \frac{1}{(2\pi)^3 g_s \alpha'^2} = \frac{N}{2\pi^2 L^4} \,, \ (2.3)$$

where $G_{\alpha\beta} = G_{mn}(X)\partial_{\alpha}X^{m}\partial_{\beta}X^{n}$. Here we are considering the case of Minkowski signature (-+...+) but will rotate to the Euclidean one when discussing the value of the classical and 1-loop free energy as we will be interested in the case when the boundary of AdS₃ representing the defect is S^{2} .⁴

We shall consider the solution for which the probe D3 brane is wrapped on AdS_3 and also on the circle parametrized by the 2π periodic angles ψ and ϕ so that

$$u = u_0 , \qquad \theta = \frac{\pi}{2} , \qquad \phi = \psi .$$
 (2.4)

The world-volume gauge field $\mathcal{F}_{\alpha\beta}$ will have vanishing background. We will identify the AdS₃ coordinates with the first 3 world-volume coordinates ξ^{α} ($\alpha = \hat{0}, \hat{1}, \hat{2}$) and $\phi = \psi$ with $\xi^{\hat{3}}$.⁵

Such configuration preserves half of supersymmetry [29, 12, 14] (this may be shown along the same lines as for the similar M5 brane case in Appendix D). The induced metric on the brane is then that of $AdS_3 \times S^1$ with equal radii

$$ds_{D3}^2 = L^2 \cosh^2 u_0 (ds_{AdS_3}^2 + d\psi^2) = L^2 \cosh^2 u_0 \ \mathsf{g}_{\alpha\beta}(\xi) d\xi^{\alpha} d\xi^{\beta} , \qquad (2.5)$$

where $\mathbf{g}_{\alpha\beta}$ is the metric of the unit-radius $\mathrm{AdS}_3 \times S^1$.

The defect is represented by the boundary of AdS_3 that we shall assume to be S^2 (see footnote 1). The leading large N contribution to the corresponding free energy is given by the classical value of the Euclidean action S_E of the D3 brane probe that is found to be⁶

$$F^{(0)} = S_E = T_3 L^4 2\pi \operatorname{vol}(AdS_3) = -\frac{N}{2\pi^2} (2\pi)^2 \log(r\Lambda) = -2N \log(r\Lambda) .$$
 (2.6)

Here we used that the regularized volume of unit-radius AdS_3 whose boundary is S^2

$$\operatorname{vol}(AdS_3) = -2\pi \log(r\Lambda) , \qquad (2.7)$$

where $\Lambda \to \infty$ is an IR cutoff (or UV cutoff in the dual gauge theory side) and r is the radius of S^2 . Note that the dependence on the position u_0 cancelled out in (2.6). The dependence on this modulus parameter will be absent also in the quantum corrections discussed below.

On the gauge theory side, the corresponding S^2 defect free energy $F_{\text{def}} \equiv F$ defined in terms of the partition function as $Z_{\text{def}} = Z e^{-F}$ may be represented as

$$F = -\frac{b}{3}\log(r\Lambda) , \qquad (2.8)$$

⁴Note that the formal structure of the spectrum of fluctuations of the brane near the classical configuration will not depend on the signature choice.

⁵We shall use the following notation: m, n, ... will stand for target space indices (0, 1, ...9); $\alpha, \beta, ...$ will be the brane world-volume indices; i, j, k will denote the AdS₅ indices. Explicit values of the world-volume indices will be indicated with hats, e.g., $\hat{1}$, etc. Indices along the tangent space directions will be underlined, e.g., $\underline{9}$ or $\hat{\underline{1}}$, etc.

⁶Note that for the choice of C_4 in (2.3) (different from one in [10] but the same as in [13]) we have this potential regular at u = 0: for small u the metric in (2.1) contains $du^2 + u^2 d\psi^2$ so that $C_4 \sim u^2 d\psi \wedge \operatorname{vol}_{AdS_3}$ is analytic when expressed in terms of Cartesian coordinates (cf. also footnote 20). As a result, the classical value of the action is non-zero.

where b is the corresponding defect conformal anomaly coefficient. The general expression for the coefficient b was found in [2, 3] and was shown to depend only on the Levi group of the defect. For a single probe D3 brane, the resulting defect on the boundary corresponds to the Levi group $S(U(1) \times U(N-1))$, as discussed in [10]. In this case we get

$$b = 3[N^2 - 1 - (N - 1)^2] = 6N - 6.$$
(2.9)

The defect free energy in (2.8) is expected to be matched to the corresponding value of the effective action of the D3 brane probe in $AdS_5 \times S^5$ background. Indeed, the leading order N term in (2.8), (2.9) is in agreement with the one following from the classical value of the D3 brane action in (2.6), i.e.

$$b^{(0)} = 6N (2.10)$$

The subleading $b^{(1)} = -6$ term in (2.9) should come from the quantum 1-loop contribution of the D3 brane fluctuations around the above background. According to (2.9) all higher (2-loop, etc.) D3 brane corrections should be absent which should be a consequence of the supersymmetry of this problem.

Our aim will be to compute the 1-loop correction to (2.6). We will need to add together the fluctuations of the "transverse" scalars, fermions and world-volume gauge vector. The structure of the 1-loop computation is similar to the one described in [30, 31] in the case of the solution of [32] where the induced metric on D3 brane in $AdS_5 \times S^5$ was $AdS_2 \times S^2$.

2.1 Scalar fluctuations

Choosing the static gauge

$$AdS_3 = \{\xi^{\hat{0}}, \xi^{\hat{1}}, \xi^{\hat{2}}\}, \qquad \psi = \xi^{\hat{3}} \in [0, 2\pi], \qquad (2.11)$$

and considering fluctuations of the two "transverse" AdS₅ coordinates $u = u_0 + \delta u$, $\phi = \xi^{\hat{3}} + \delta \phi$ one finds for the quadratic fluctuation part of the D3 brane action in (2.3) (see Appendix A for details of the derivation of the scalar fluctuation action)

$$\int (\sqrt{-G} - C_4) \to c \int d^4 \xi \sqrt{-\mathbf{g}} \Big(\mathbf{g}^{\alpha\beta} \partial_\alpha \delta u \partial_\beta \delta u + \mathbf{g}^{\alpha\beta} \tanh^2 u_0 \partial_\alpha \delta \phi \partial_\beta \delta \phi + 4 \tanh u_0 \delta u \partial_3 \delta \phi \Big)$$
$$= c \int d^4 \xi \sqrt{-\mathbf{g}} \Big[\mathbf{g}^{\alpha\beta} \partial_\alpha \bar{\chi} \partial_\beta \chi + i(\bar{\chi} \partial_\beta \chi - \chi \partial_\beta \bar{\chi}) \Big] \,. \tag{2.12}$$

$$= c \int d^{4}\xi \sqrt{-\mathbf{g}} \Big[\mathbf{g}^{\alpha\beta} \partial_{\alpha} \bar{\chi} \partial_{\beta} \chi + i(\bar{\chi} \partial_{\hat{3}} \chi - \chi \partial_{\hat{3}} \bar{\chi}) \Big] , \qquad (2.12)$$

$$\chi \equiv \delta u + i \tanh u_0 \,\delta \psi \,, \qquad c \equiv \frac{1}{2} \cosh^2 u_0 \,L^4 \,. \tag{2.13}$$

Note that in the second line u_0 enters only via the overall factor c that can be rescaled away. Since $\mathbf{g}^{\alpha\beta}\partial_{\alpha}\bar{\chi}\partial_{\beta}\chi + i(\bar{\chi}\partial_{3}\chi - \chi\partial_{3}\bar{\chi}) = \mathbf{g}^{ij}\partial_{i}\bar{\chi}\partial_{j}\chi + (\partial_{3}\chi - i\chi)(\partial_{3}\bar{\chi} + i\bar{\chi}) - \bar{\chi}\chi$ (where i, j label AdS₃ directions) we get a conformally coupled complex scalar in AdS₃ × S¹⁷ also coupled to an effective constant U(1) gauge field with $A_{3} = 1$ with charge q = 1.

⁷To recall, the conformally coupled 4d scalar operator $-\nabla^2 + \frac{1}{6}R$ for $R(AdS_3 \times S^1) = -6$ becomes $-\nabla^2 - 1$ (see also footnote 2).

Expanding in Fourier modes in $\xi^{\hat{3}}$ we get 2 towers of scalar modes in AdS₃ with masses⁸

$$m^2 = n^2 \pm 2n = (n \pm 1)^2 - 1$$
, $n = 0, \pm 1, \pm 2, \dots$. (2.14)

The corresponding values of the AdS_3 boundary field dimensions $\Delta(\Delta - 2) = m^2$ are thus given by

$$\Delta - 1 = |n \pm 1| . \tag{2.15}$$

Here we assume the Dirichlet boundary conditions for the scalars (as appropriate for a defect interpretation) so that $\Delta - 1 \ge 0$.

To find the quadratic action of the 4 transverse fluctuations in S^5 let us set $\theta = \frac{\pi}{2} + v$ and $ds_{S^3}^2 = d\varphi_1^2 + \cos^2 \varphi_1 (d\varphi_2^2 + \sin^2 \varphi_2 d\varphi_3^2)$ and introduce 4 cartesian coordinates $X^a = \{x, y, z, w\}$ such that

$$v = \sqrt{x^2 + y^2 + z^2 + w^2}$$
, $\tan \varphi_1 = \frac{x}{\sqrt{y^2 + z^2 + w^2}}$, $\tan \varphi_2 = \frac{\sqrt{w^2 + z^2}}{y}$, $\tan \varphi_3 = \frac{w}{z}$.
(2.16)

Then from (2.3) we get $(C_4$ does not contribute at this order, cf. (2.13))

$$\int \sqrt{-G} \rightarrow c \int d^4 \xi \sqrt{-\mathsf{g}} \sum_{a=1}^4 \left(\mathsf{g}^{\alpha\beta} \partial_\alpha \mathsf{X}^a \partial_\beta \mathsf{X}^a - \mathsf{X}^a \mathsf{X}^a \right) \,. \tag{2.17}$$

Thus we get 4 conformally coupled scalars in $AdS_3 \times S^1$. Expanding in S^1 modes we get 4 towers of scalar operators with masses

$$m^2 = n^2 - 1$$
, $n = 0, \pm 1, \pm 2, \dots$. (2.18)

Again assuming the Dirichlet boundary conditions the corresponding 2d scaling dimensions are

$$\Delta - 1 = |n| . \tag{2.19}$$

2.2 Fermionic fluctuations

The quadratic fermionic part of the D3 brane action in a target space with a non-trivial F_5 background and no world-volume gauge field may be written as (see, e.g., [33])⁹

$$S_f = \int d^4 \xi \, \sqrt{-g} \, g^{\alpha\beta} \, \bar{\Theta} (1 - \tilde{\Gamma}_{D3}) \, \tilde{\Gamma}_{\alpha} \widehat{D}_{\beta} \Theta \,, \qquad (2.20)$$

where $g_{\alpha\beta}$ is the induced metric in the static gauge (2.5) (i.e. $AdS_3 \times S^1$) and

$$\nabla_{m} = \partial_{m} + \frac{1}{4} \Omega_{\overline{m}}^{\underline{n}\underline{k}} \Gamma_{\underline{n}\underline{k}} , \qquad \Gamma_{\alpha} = \partial_{\alpha} X^{m} \Gamma_{m} , \qquad \Gamma_{m} = E_{\overline{m}}^{\underline{m}} \Gamma_{\underline{m}} , \qquad E_{\overline{m}}^{\underline{m}} E_{\overline{n}}^{\underline{n}} \eta_{\underline{m}\underline{n}} = G_{mn} ,$$

$$\{\Gamma_{\underline{m}}, \Gamma_{\underline{n}}\} = 2\eta_{\underline{m}\underline{n}} , \qquad \{\Gamma_{\alpha}, \Gamma_{\beta}\} = 2g_{\alpha\beta} . \qquad (2.22)$$

⁸Note that our D3 brane embedding and fluctuation spectrum is different from the one in [9] where the brane was not wrapping ψ of AdS₅ or effectively u_0 was set to zero from the start.

⁹Here we ignore the overall constant factor of $\frac{1}{2}T_3$. The complete form of the action for a D3 brane in $AdS_5 \times S^5$ background was given in [34].

 $\Omega_m^{\underline{nk}}$ is the spin connection in the 10d target space. For index notation see footnote 5.

The fermion field is $\Theta = \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}$ where θ_I are 10d positive chirality MW spinors satisfying $\theta^* = iC\Gamma^{\underline{0}}\theta$ and (we use $\epsilon^{\underline{012\cdots9}} = 1$)

$$\Gamma_{10}\theta_I = \theta_I , \qquad \Gamma_{10} = -\frac{1}{10!} \epsilon^{\underline{m}_0 \dots \underline{m}_9} \Gamma_{\underline{m}_0} \dots \Gamma_{\underline{m}_9} = \Gamma^{\underline{0}} \Gamma^{\underline{1}} \dots \Gamma^{\underline{9}} . \qquad (2.23)$$

Also, we use $\bar{\Theta} = i\Theta^{\dagger}\tilde{\Gamma}^{\underline{0}} = \Theta^{T}C$. In a proper basis, both θ_{I} have 32 real components. $\tilde{\Gamma}_{\alpha}$ stands for $\Gamma_{\alpha} \otimes 1_{2}$ where $1_{2} = \delta_{IJ}$. In (2.21) we suppressed 1_{2} factors and σ_{2} acts on the index I = 1, 2. Finally, $\tilde{\Gamma}_{D3}$ in (2.20) is defined as (here $\epsilon^{\underline{0}\hat{1}\hat{2}\hat{3}} = 1$)

$$\tilde{\Gamma}_{D3} = \Gamma_{D3} \otimes (-i\sigma_2) , \qquad \Gamma_{D3} = \frac{\epsilon^{\alpha_1 \cdots \alpha_4}}{4!\sqrt{-g}} \Gamma_{\alpha_1 \cdots \alpha_4} = \Gamma_{\underline{\hat{0}\hat{1}\hat{2}\hat{3}}} , \qquad (\tilde{\Gamma}_{D3})^2 = 1 .$$
(2.24)

Following [33] we will fix the κ -symmetry gauge by imposing

$$\tilde{\Gamma}_{10}\Theta = \Theta , \qquad \tilde{\Gamma}_{10} = \Gamma_{10} \otimes \sigma_3 , \qquad \Gamma_{10} = \Gamma^{\underline{01}\dots\underline{9}} .$$
(2.25)

This is equivalent to $\theta_2 = 0$ so from now on we set $\theta_1 \equiv \vartheta$. Then we get for (2.20)

where we used that $\{\Gamma_{D3}, \Gamma_{\alpha}\} = 0$.

We shall set L = 1 and label the 10d coordinates as

$$X^{0,1,2} = AdS_3 , \quad X^3 = u , \qquad X^4 = \psi , \quad X^5 = \theta , \quad X^6 = \phi , \quad X^{7,8,9} = S^3 , \quad (2.27)$$

where in the static gauge $X^{0,1,2} = \xi^{\hat{0},\hat{1},\hat{2}}$, $X^3 = u = u_0$, $X^4 = \psi = \xi^{\hat{3}}$, $X^5 = \theta = \frac{\pi}{2}$, $X^6 = \phi = \xi^{\hat{3}}$, $X^{7,8,9}$ =const. The corresponding spin connection components are given in Appendix B. In the present case of $AdS_5 \times S^5$ we find that the induced covariant derivative in (2.21) is given by (see (B.15))

$$\Gamma^{\alpha}\partial_{\alpha}X^{m}\nabla_{m} = \nabla + 2\tanh u_{0}\Gamma_{\underline{3}} + \frac{1}{2\cosh u_{0}}\Gamma_{\underline{643}} , \qquad (2.28)$$

where ∇ denotes the Dirac operator on $\operatorname{AdS}_3 \times S^1$. As a result, we get from (2.26) the action for ϑ of the standard Dirac form

$$S_f = \int d^4 \xi \, \sqrt{-g} \, \bar{\vartheta} (\nabla + \mathbf{M}) \vartheta \,, \qquad (2.29)$$

$$M = 2 \tanh u_0 \Gamma_{\underline{3}} + \frac{1}{2 \cosh u_0} \Gamma_{\underline{643}} - \frac{1}{16} \Gamma_{D3} \Gamma^{\alpha} \not\!\!\!/ _5 \Gamma_{\alpha} .$$
(2.30)

To compute the contribution of the F_5 term in (2.21) to M we note that the self-dual F_5 corresponding to C_4 in (2.2) is given by

$$F_5 = -4L^4(1+*)\operatorname{vol}_{AdS_5} = -4L^{-1} \left(E^{\underline{0}} \wedge E^{\underline{1}} \wedge E^{\underline{2}} \wedge E^{\underline{3}} \wedge E^{\underline{4}} + E^{\underline{5}} \wedge E^{\underline{6}} \wedge E^{\underline{7}} \wedge E^{\underline{8}} \wedge E^{\underline{9}} \right) .$$
(2.31)

$$\mathbf{M} = 2 \tanh u_0 \,\Gamma_{\underline{3}} + \frac{1}{2 \cosh u_0} \Gamma_{\underline{643}} + \frac{1}{4} \Gamma_{D3} \Gamma^{\alpha} \left(\Gamma^{\underline{0}\cdots\underline{4}} + \Gamma^{\underline{5}\cdots\underline{9}} \right) \Gamma_{\alpha} \,, \tag{2.32}$$

where Γ_{D3} is given in (2.24). We have checked explicitly that the result for the fermionic spectrum does not depend on u_0 : the dependence on u_0 can be absorbed into a rotation of ϑ by a u_0 dependent phase in the (<u>46</u>) plane.¹⁰ Thus we may thus take the limit $u_0 \to 0$ and keep only the leading terms (as is easy to see, they are non-singular).

Using that $\Gamma^i(\Gamma^{\underline{0}\cdots\underline{4}} + \Gamma^{\underline{5}\cdots\underline{9}})\Gamma_i = 3(\Gamma^{\underline{0}\cdots\underline{4}} - \Gamma^{\underline{5}\cdots\underline{9}})$ for $\alpha = i = \hat{0}, \hat{1}, \hat{2}$ and $\Gamma^6(\Gamma^{\underline{0}\cdots\underline{4}} + \Gamma^{\underline{5}\cdots\underline{9}})\Gamma_6 = -(\Gamma^{\underline{0}\cdots\underline{4}} - \Gamma^{\underline{5}\cdots\underline{9}})$ for $\alpha = \hat{3}$, together with $\Gamma_{D3} = \Gamma_{\underline{0}\hat{1}\hat{2}\hat{3}} = \Gamma_{\underline{0}\underline{1}\underline{2}6}$, we get

$$\mathbf{M} = \frac{1}{2}\Gamma_{\underline{643}} + \frac{1}{4}(3-1)\Gamma_{D3}\left(\Gamma^{\underline{0}\cdots\underline{4}} - \Gamma^{\underline{5}\cdots\underline{9}}\right) = \frac{1}{2}\Gamma_{\underline{643}} + \frac{1}{2}\Gamma_{\underline{0126}}\left(\Gamma^{\underline{0}\cdots\underline{4}} - \Gamma^{\underline{5}\cdots\underline{9}}\right) = \frac{1}{2}\Gamma_{\underline{0125789}} \ . \ (2.33)$$

Due to the chirality constraint in (2.23), i.e. $\Gamma^{\underline{01}\cdots\underline{9}}\vartheta = \vartheta$, we conclude that acting on ϑ the mass operator takes the following simple form

$$\mathbf{M} = \frac{1}{2} \Gamma_{\underline{0125789}} \Gamma^{\underline{01}\dots\underline{9}} = -\frac{1}{2} \Gamma_{\underline{643}} = \frac{1}{2} \Gamma_{\underline{346}} .$$
 (2.34)

Thus the contribution of the F_5 term in (2.30) is twice opposite that of the normal component of the spin connection in (2.28), i.e. it effectively reverses the sign of the former.¹¹

Expanding ϑ in modes in $\xi^{\hat{3}}$ the Dirac operator in (2.29) on $\operatorname{AdS}_3 \times S^1$ reduces to that on AdS_3 (we use that $\Gamma^{\hat{3}} = \Gamma^{\hat{6}}$)

$$i(\nabla + \mathbf{M}) = i \nabla_{AdS_3} + i \Gamma^{\hat{3}} \partial_{\hat{3}} + i \mathbf{M} \quad \rightarrow \quad i \nabla_{AdS_3} - \hat{M} , \qquad \qquad \hat{M} = n \Gamma_{\underline{6}} + \frac{1}{2} i \Gamma_{\underline{643}} . \tag{2.35}$$

Equivalently, we may write the operator in (2.35) as $i \nabla AdS_3 + i \Gamma^{\hat{3}} (\partial_{\hat{3}} - \frac{1}{2} \Gamma_{\underline{43}})$, i.e. we get a set of 4 massless fermions in $AdS_3 \times S^1$ coupled to a constant U(1) gauge potential in $\hat{3}$ direction.¹²

Since $\Gamma_{\underline{6}}^2 = 1$, $(i\Gamma_{\underline{643}})^2 = 1$ and $[\Gamma_{\underline{6}}, \Gamma_{\underline{643}}] = 0$ we conclude that \hat{M} has eigenvalues $m_f = \pm n \pm \frac{1}{2}$ $(n = 0, \pm 1, ...)$. Thus we find 4 towers of 3d fermions with such masses. The corresponding dimensions of the boundary operators are then (assuming again the standard, i.e. the Dirichlet, boundary conditions)

$$\Delta - 1 = |m_f| = |n \pm \frac{1}{2}|, \qquad n = 0, \pm 1, \pm 2, \dots .$$
(2.36)

¹⁰This rotation reflects the fact that on the classical solution both $X^4 = \psi$ and $X^6 = \phi$ are equal to $\xi^{\hat{3}}$ and thus have the same projections on the world volume, leading to an effective mixing of the Γ -matrices in the (<u>46</u>) directions.

¹¹The same mass operator is found also for generic u_0 , see Appendix C.

¹²Since $\Gamma_{\underline{43}}$ commutes with $\Gamma^{\underline{3}} = \Gamma^{\underline{6}}$ it can be diagonalized with $\pm i$ as eigenvalues, i.e. the U(1) gauge field is $A_{\underline{3}} = 1$ with the fermion charges being $q = \pm \frac{1}{2}$.

2.3 Vector field contribution

As the world-volume vector gauge field in (2.3) has no background value its contribution to 1loop partition function is the same as of a Maxwell field propagating on $\operatorname{AdS}_3 \times S^1$ background with the standard action $\int d^4 \xi \sqrt{-g} \, \mathcal{F}^{\alpha\beta} \mathcal{F}_{\alpha\beta}$. The partition function of a Maxwell vector on a general curved 4d background may be written as

$$Z_1 = \det(-\nabla^2) \left[\det(-g_{\alpha\beta}\nabla^2 + R_{\alpha\beta}) \right]^{-1/2} .$$
(2.37)

In the unit-radius $\operatorname{AdS}_3 \times S^1$ case we have $R_{\alpha\beta} = (-2g_{ij}, 0)$ where g_{ij} is the AdS_3 metric. Then $\det(-g_{\alpha\beta}\nabla^2 + R_{\alpha\beta}) = \det(-\nabla^2) \det(-\nabla^2 - 2)_{ij}$. Splitting A_{μ} into the longitudinal and transverse parts we get $\det(-\nabla^2 - 2)_{\mu\nu} = \det(-\nabla^2) \det(-\nabla^2 - 2)_{ij,\perp}$ where $(-\nabla^2 - 2)_{ij,\perp}$ is defined on a transverse A_{μ} depending on $\operatorname{AdS}_3 \times S^1$ coordinates. Thus

$$Z_1 = \left[\det(-\nabla^2 - 2)_{ij,\perp}\right]^{-1/2}, \qquad \nabla^2 = \nabla^2_{AdS_3} + \partial^2_{\hat{3}}. \qquad (2.38)$$

Expanding the transverse vector A_{μ} in modes in $\xi^{\hat{3}}$ we thus get a tower of transverse 3d vectors with masses

$$m_1^2 = n^2 - 2 . (2.39)$$

The corresponding boundary dimension is found from $\Delta(\Delta - 2) = m_1^2 + 1 = n^2 - 1$ and thus with the Dirichlet boundary condition choice

$$\Delta - 1 = |n| , \qquad n = 0, \pm 1, \pm 2, \dots .$$
(2.40)

One can check directly that for n = 0 the same result is found by first dimensionally reducing the Maxwell action to AdS₃ (i.e. getting a 3d Maxwell field plus a massless scalar) and then quantizing in the $\nabla_i A^i = 0$ gauge.

The resulting fluctuation spectrum in (2.15), (2.19), (2.36), (2.40)) is that of the supersymmetric $\mathcal{N} = 4$ vector multiplet defined on $\operatorname{AdS}_3 \times S^1$. It can be indeed organized into $\mathcal{N} = 2$ supermultiplets on AdS_3 as described in [35]. Recall that in flat 4 dimensions the $\mathcal{N} = 4$ vector multiplet is a superposition of one $\mathcal{N} = 2$ vector multiplet (vector, 2 real scalars, 2 Weyl fermions) and one $\mathcal{N} = 2$ hypermultiplet (4 real scalars and 2 Weyl fermions). In the present $\operatorname{AdS}_3 \times S^1$ case we also get a collection of $\mathcal{N} = 2$ vector multiplet and a hypermultiplet with masses of fields as given above. In terms of AdS_3 towers of fields we have:¹³

(i) vector multiplet containing 1 vector with $\Delta - 1 = |n|$, 2 scalars with $\Delta - 1 = |n|$, 2 fermions with $\Delta - 1 = |n \pm \frac{1}{2}|$;

(ii) hypermultiplet containing 2 scalars with $\Delta - 1 = |n|$, 2 scalars with $\Delta - 1 = |n \pm 1|$, 2 fermions $\Delta - 1 = |n \pm \frac{1}{2}|$.

 $^{^{13}}$ In the notation of [35] adapted to the 2d boundary theory these are (2,2) vector multiplet and (2,2) hypermultiplet.

2.4 1-loop free energy

The expressions for 1-loop determinants of the relevant fields in AdS_3 can be found, e.g., in [22, 24, 26]. We shall always assume Dirichlet boundary conditions so that $\Delta - 1 \ge 0.14$ In particular, for a real scalar we get¹⁵

$$F_0 = \frac{1}{2}\log\det(-\nabla_{AdS_3}^2 + m^2) = -\frac{1}{12\pi}(\Delta - 1)^3\operatorname{vol}(AdS_3) , \qquad (\Delta - 1)^2 = m^2 + 1 . \quad (2.41)$$

For the vector contribution we find (see (2.38))

$$F_1 = \frac{1}{2} \log \det(-\nabla_{AdS_3}^2 + m^2)_{ij,\perp} = -\frac{2}{12\pi} (\Delta - 1) \left[(\Delta - 1)^2 - 3 \right] \operatorname{vol}(AdS_3), \qquad (\Delta - 1)^2 = m^2 + 2.$$
(2.42)

$$F_{1/2} = \frac{1}{2} \log \det(-\nabla_{AdS_3}^2 + \frac{1}{4}R + m_f^2) = -\frac{2}{12\pi} (\Delta - 1) \left[(\Delta - 1)^2 - \frac{3}{4} \right] \operatorname{vol}(AdS_3) , \quad (\Delta - 1)^2 = m_f^2.$$
(2.43)

Here vol (AdS_3) is given by (2.7), i.e. each contribution scales as $\log(r\Lambda)$.

Let us first consider the 1-loop result found for the collection of fields of the standard $\mathcal{N} = 4$ vector multiplet on $\mathrm{AdS}_3 \times S^1$, i.e. for 6 conformally coupled scalars, a gauge vector and 4 Weyl fermions.¹⁶ Upon expansion in S^1 modes that gives a set of AdS_3 fields for each value of $n = 0, \pm 1, \ldots$ 6 scalars with $m^2 = n^2 + \frac{1}{6}R = n^2 - 1$, a vector with $m^2 = n^2 - 2$ and 4 fermions with $m^2 = n^2$. Then using (2.41),(2.42),(2.43) the total 1-loop free energy is found to vanish

$$F^{(1)} = 6F_0 + F_1 - 4F_{1/2} = -\frac{1}{12\pi} P \operatorname{vol}(AdS_3) , \qquad P = \sum_{n=-\infty}^{\infty} P_n , \qquad (2.44)$$

$$P_n = 6|n|^3 + 2|n|(n^2 - 3) - 8|n|(n^2 - \frac{3}{4}) = 0.$$
(2.45)

¹⁴In general, if, e.g., for a scalar in AdS_{p+1} we have $(\Delta - \frac{p}{2})^2 = m^2$, then for the standard Dirichlet case $\Delta = \Delta_+$ where $\Delta_+ - \frac{p}{2} = |m| \ge 0$.

¹⁵Let us recall that for an operator defined on symmetric traceless transverse spin s field in AdS₃ [26]

$$F_{s} = \frac{1}{2} \log \det(-\nabla_{AdS_{3}}^{2} + m^{2})_{\mathrm{T},\perp} = -g_{s} \frac{\operatorname{vol}(AdS_{3})}{\operatorname{vol}(S^{2}) [2\Gamma(\frac{3}{2})]^{2}} \lim_{z \to 0} \frac{\partial}{\partial z} \int_{0}^{\infty} d\lambda \frac{\lambda^{2} + s^{2}}{[\lambda^{2} + (\Delta - 1)^{2}]^{z}}$$
$$= -\frac{1}{12\pi} g_{s} (\Delta - 1) \Big[(\Delta - 1)^{2} - 3s^{2} \Big] \operatorname{vol}(AdS_{3}) , \qquad (\Delta - 1)^{2} = m^{2} + s + 1 .$$

where $g_0 = 1$ and $g_s = 2$ for s > 0. To get the free energy for a massless gauge field in AdS₃ one is to add the contribution of the ghost operator. As discussed above, here for n = 0 the vector contribution F_1 is different from the one for a 3d s = 1 gauge field as it also contains an extra massless scalar part that cancels the ghost determinant contribution. Note also that if one dualises the massless 3d vector in AdS₃ to a massless scalar the latter will be subject to the Neumann boundary condition so will have the opposite sign of the free energy contribution compared to the standard Dirichlet massless scalar. Then the total contribution of a vector in AdS₃ × S¹ dimensionally reduced to AdS₃ (i.e. a combination of a 3d vector and a massless scalar) will be zero, in agreement with the vanishing of the vector contribution in (2.45) or (2.47) for n = 0.

¹⁶Here do not introduce a coupling to an extra U(1) gauge field so the global supersymmetry on $AdS_3 \times S^1$ is not preserved.

The cancellation of the n^3 terms is due to the balance of the numbers of bosonic and fermionic degrees of freedom. The cancellation of the linear in n terms (which would produce a quadratic divergence in the sum in (2.44)) is related to the general fact that for the $\mathcal{N} = 4$ vector multiplet defined on a curved 4-space the coefficient of quadratic UV divergence (determined for log det $(-\nabla^2 + X)$ by the Seeley coefficient $b_2 = \text{tr}(\frac{1}{6}R - X)$) can be shown to vanish: the conformally coupled scalars have $b_2 = 0$ while the vector contribution cancels against the fermionic one.¹⁷

This observation should also apply to the case of the fluctuation spectrum we have found above: it corresponds to the fields of the $\mathcal{N} = 4$ vector multiplet on $\mathrm{AdS}_3 \times S^1$ coupled also in a specific way to a constant U(1) gauge potential in $\hat{3}$ direction (which in the present case originates from a non-trivial embedding of the D3 brane into the target space $\mathrm{AdS}_5 \times S^5$ background). Its presence shifts the values of n for 2 scalars and the fermion modes. It cannot alter the cancellation of UV divergences but may contribute a non-trivial constant term to the analog of the sum in (2.44).

Combining the contributions of the AdS₃ modes in (2.15),(2.19),(2.36),(2.40), i.e. 2 conformally coupled scalars with shifts ± 1 , 4 scalars with shift 0, a vector and 2 sets of fermions with shifts $\pm \frac{1}{2}$ we get the following counterpart of (2.44),(2.45)

$$F^{(1)} = \frac{1}{6} P \log(r\Lambda) , \qquad P = \sum_{n=-\infty}^{\infty} P_n , \qquad (2.46)$$
$$P_n = |n+1|^3 + |n-1|^3 + 4|n|^3 + 2|n|(n^2 - 3)$$
$$-4|n+\frac{1}{2}|\left[(n+\frac{1}{2})^2 - \frac{3}{4}\right] - 4|n-\frac{1}{2}|\left[(n-\frac{1}{2})^2 - \frac{3}{4}\right] . \qquad (2.47)$$

For $n \neq 0$ we learn that $P_n = 0$, i.e. all order n^3 and n terms cancel which should be a consequence of underlying supersymmetry. The non-trivial contribution thus come just from the n = 0 level states: from 2 "shifted" scalars and the fermions

$$P = P_0 = 2 - 4\left(-\frac{1}{4} - \frac{1}{4}\right) = 4 , \qquad b^{(1)} = -\frac{1}{2}P = -2 . \qquad (2.48)$$

Adding $F^{(1)}$ in (2.46),(2.48) to the classical contribution in (2.6) we finish with

$$F = F^{(0)} + F^{(1)} = -2\left(N - \frac{1}{3}\right)\log(r\Lambda) .$$
(2.49)

This appears to disagree with the prediction in [2, 3], i.e. $F = -2(N-1)\log(r\Lambda)$ as given in (2.8),(2.9). The reason for this disagreement remains to be understood. One issue might be the choice of boundary conditions of some low-lying modes. Although the Dirichlet boundary conditions are the simplest and most natural ones here, the choice of the Neumann boundary conditions might also be possible and help to resolve the discrepancy.

¹⁷Note also that in the special case of $\operatorname{AdS}_3 \times S^1$ the b_4 Seeley coefficient also vanishes (in agreement with no log UV divergence in (2.45)): this space is conformally flat, i.e. Weyl tensor is zero and also the 4d Euler density R^*R^* vanishes for any $M^3 \times S^1$ space.

3 S^4 defect anomaly in (2,0) theory from M5 probe in AdS₇ × S^4

Let us now consider a similar computation in the case of $\frac{1}{2}$ BPS configuration of M5 brane in AdS₇ × S⁴ with induced metric AdS₅ × S¹. Following [13] we parametrize AdS₇ × S⁴ as $(u \in (0, \infty); \psi, \phi \in [0, 2\pi])^{18}$

$$ds_{11}^2 = L_A^2 \left(du^2 + \cosh^2 u \, ds_{AdS_5}^2 + \sinh^2 u \, d\psi^2 \right) + L^2 \left(d\theta^2 + \sin^2 \theta \, d\phi^2 + \cos^2 \theta \, ds_{S^2}^2 \right) \,, \quad (3.1)$$

$$F_4 = dC_3 = 3L^3 \operatorname{vol}_{S^4} = 3L^3 \sin \theta \cos^2 \theta \, d\theta \wedge d\phi \wedge \operatorname{vol}_{S^2} \,, \tag{3.2}$$

$$L_A = 2L$$
, $L_A^3 = 8\pi N \ell_P^3$. (3.3)

We will assume the Minkowski signature of this 11d background but at the end will rotate to the Euclidean one as will be interested in the case when the boundary of AdS_5 is S^4 .

The bosonic part of the action of an M5 brane in a 11d supergravity background may be written as [36-38]

$$S = -T_5 \left\{ \int d^6 \xi \left[\sqrt{-\det(G_{\alpha\beta} + i\widetilde{\mathcal{H}}_{\alpha\beta})} - \frac{1}{4}\sqrt{-G} \widetilde{\mathcal{H}}^{\star \alpha\beta} \widetilde{\mathcal{H}}_{\alpha\beta} \right] - \int \left(C_6 + \frac{1}{2}\mathcal{H} \wedge C_3 \right) \right\}, \quad (3.4)$$

$$H_{\mu\nu\lambda} = 3\partial_{[\mu}A_{\nu\lambda]} , \qquad H_{\mu\nu\lambda} = H_{\mu\nu\lambda} - C_{\mu\nu\lambda} , \qquad H^{\star\mu\nu\lambda} = \frac{1}{6\sqrt{-G}} \epsilon^{\mu\nu\lambda\alpha\beta\gamma} H_{\alpha\beta\gamma} , \quad (3.5)$$
$$\widetilde{H}_{\mu\nu} = H^{\star}_{\mu\nu\lambda} U^{\lambda} , \qquad \widetilde{H}^{\star}_{\mu\nu} = H_{\mu\nu\lambda}U^{\lambda} , \qquad U_{\lambda}(\xi) \equiv \frac{\partial_{\lambda}a(\xi)}{\sqrt{(\partial_{\mu}a)^{2}}} ,$$
$$T_{5} = \frac{1}{(2\pi)^{5}\ell_{P}^{6}} = \frac{2N^{2}}{\pi^{3}L_{A}^{6}} . \qquad (3.6)$$

Here $G_{\alpha\beta} = \partial_{\alpha} X^m \partial_{\beta} X^n G_{mn}(X(\xi))$ (X^m are 11d coordinates), $C_{\mu\nu\lambda} = C_{mnk} \partial_{\mu} X^m \partial_{\nu} X^n \partial_{\lambda} X^k$ and $H_{\alpha\beta\gamma}$ (which is self-dual on shell) is the field strength of the world-volume antisymmetric gauge field $A_{\alpha\beta}(\xi)$. The auxiliary scalar $a(\xi)$ may be fixed by a gauge choice $a(\xi) = \xi^{\hat{5}}$ [39] and will play no role below. The 6-form potential C_6 is defined by¹⁹

$$dC_6 = F_4^{\star} - \frac{1}{2}C_3 \wedge F_4 , \qquad (3.7)$$

Then from (3.2) we get²⁰

$$C_6 = L_A^6(\cosh^6 u - 1) \mathrm{vol}_{AdS_5} \wedge d\psi .$$
 (3.8)

¹⁸Note that the 2π periodicity of ψ guarantees that there is no singularity for $u \to 0$. Here $vol(S^4) = \int vol_{S^4} = \frac{8}{3}\pi^2$.

 $^{^{19}}F_4^*$ is the 11d dual of F_4 . Note also that $d(dC_6) = 0$ on the equations of motion for C_3 (assuming there is no 11d gravitino background).

²⁰ In general, the WZ term in (3.4) should be defined in terms of an integral of $F_7 + \frac{1}{2}H_3 \wedge F_4$ over 7-space with 6d boundary (cf. also [32]). Then it is invariant under "large" gauge transformations that change C_6 and may in principle change its integral. The result does not depend on a choice of 7-space as long as the charge quantization condition is satisfied. Note that the shift by -1 in (3.8) is required for the potential C_6 not to be singular near the origin $u \to 0$ (cf. footnote 6).

The BPS solution for the M5 brane wrapped on $AdS_5 \subset AdS_7$ and also on two circles ψ and ϕ has vanishing 3-form $H_{\mu\nu\lambda}$ and is a direct analog of the D3 brane solution in (2.4)

$$u = u_0 , \qquad \theta = \frac{\pi}{2} , \qquad \phi = 2\psi .$$
 (3.9)

Here the factor of 2 in the relation between ϕ and ψ is related to the factor of 2 ratio of the AdS₇ and S^4 radii in (3.3). u_0 is an arbitrary modulus. As both ψ and ϕ are 2π periodic the brane wraps twice around the ϕ circle of S^4 . This M5 embedding preserves half of supersymmetry of the AdS₄ × S^7 background (see Appendix D).

The induced metric on M5 brane is then

$$ds_{M5}^{2} = L_{A}^{2} \left(\cosh^{2} u_{0} \ ds_{AdS_{5}}^{2} + \sinh^{2} u_{0} \ d\psi^{2}\right) + 4L^{2} d\psi^{2} = L_{A}^{2} \cosh^{2} u_{0} \left(ds_{AdS_{5}}^{2} + d\psi^{2}\right)$$
$$= g_{\alpha\beta} d\xi^{\alpha} d\xi^{\beta} = L_{A}^{2} \cosh^{2} u_{0} \ \mathsf{g}_{\alpha\beta} d\xi^{\alpha} d\xi^{\beta} , \qquad (3.10)$$

where $\mathbf{g}_{\alpha\beta}$ is the metric of unit-radius $\mathrm{AdS}_5 \times S^1$.

The classical value of the Euclidean M5 brane action S_E corresponding to the Minkowski one $S = -T_5 \left(\int d^6 \xi \sqrt{-\det G} - \int C_6 \right)$ found from (3.4) is

$$F^{(0)} = S_E = 2\pi L_A^6 T_5 \operatorname{vol}(AdS_5) = 4N^2 \log(r\Lambda) .$$
(3.11)

Here we assumed that AdS_5 has S^4 boundary and used that for global odd-dimensional AdS space one has (cf. (2.7))

$$\operatorname{vol}(AdS_{2n+1}) = \frac{2(-1)^n \pi^n}{\Gamma(n+1)} \log(r\Lambda) \xrightarrow{n=2} \operatorname{vol}(AdS_5) = \pi^2 \log(r\Lambda) , \qquad (3.12)$$

where Λ is an IR cutoff and r is the radius of S^4 . The value of the classical action does not depend on u_0 and the same will be true also for the contribution of the quantum fluctuations.

As usual, we express the free energy on S^4 in terms of the *a*-anomaly coefficient as

$$F = 4a \log(r\Lambda) . \tag{3.13}$$

Then the expression in (3.11) corresponds to the leading large N value of the *a*-anomaly for the S^4 defect in (2,0) theory being²¹

$$a^{(0)} = N^2 . (3.14)$$

The leading N^2 scaling is consistent with the expectation that it should be effectively determined by gauge theory degrees of freedom (cf. [8]).²²

²¹Incidentally, this is 4 times the (large N part of) conformal anomaly of $SU(N) \mathcal{N} = 4$ SYM theory on S^4 . There should not be any direct connection to the $\mathcal{N} = 4$ SYM anomaly which has dual description in terms of the 10d supergravity (string theory) on $AdS_5 \times S^5$.

²²This is also consistent with the discussion in Appendix B of [40] although there the defect had shape $S^1 \times S^3$ and thus the coefficient of conformal anomaly was zero.

Our aim will be to compute the subleading correction to (3.11) or (3.14) coming from the quantum M5 brane fluctuations near the above classical solution. There appears to be no alternative (2,0) theory result for this subleading coefficient known at the moment (cf. [8]) so our 1-loop M5 brane computation will provide a prediction for it.

The set of fluctuations will be the same as for a (2,0) 6d multiplet (5 scalars, 4 fermions and self-dual 2-form) on $\operatorname{AdS}_5 \times S^1$ background with all scalars conformally coupled to the metric and 2 scalars and fermions coupled also and to a constant U(1) potential in S^1 direction reflecting the presence of a non-trivial target space geometry and the F_4 flux. As in the D3 brane case, the latter should be also responsible for preservation of the global supersymmetry of the world-volume theory defined on $\operatorname{AdS}_5 \times S^1$ at the level of the quadratic fluctuation action.

3.1 Scalar fluctuations

The derivation of the scalar fluctuation action is directly analogous to the one in section 2.1 in D3 brane case. Let us choose a static gauge where AdS_5 and ψ coordinates are not fluctuating, i.e.

$$AdS_5 = \{\xi^{\hat{0}}, ..., \xi^{\hat{4}}\}, \qquad u = u_0 + \delta u, \qquad \psi = \xi^{\hat{5}}, \qquad \phi = 2\xi^{\hat{5}} + \delta\phi, \qquad (3.15)$$

and also S^2 coordinates are fluctuating near trivial vacuum values. Here $\xi^{\hat{5}}$ is 2π periodic.

Specializing the discussion in Appendix A to the case of *p*-brane with p = 5 we get the following action for the fluctuations of u and ϕ (ignoring overall constant factor $\sim \cosh^4 u_0$)

$$S \to \int d^6 \xi \sqrt{-\mathsf{g}} \,\mathcal{L}(\chi) \,, \qquad \mathcal{L} = \mathsf{g}^{\alpha\beta} \partial_\alpha \chi \partial_\beta \bar{\chi} + 2i(\bar{\chi} \partial_{\hat{5}} \chi - \chi \partial_{\hat{5}} \bar{\chi}) \,, \qquad (3.16)$$

$$\chi = \delta u + \frac{i}{2} \tanh u_0 \delta \phi , \qquad \qquad ds^2 = \mathsf{g}_{\alpha\beta} d\xi^\alpha d\xi^\beta = \mathsf{g}_{ij} d\xi^i d\xi^j + (d\xi^5)^2 , \qquad (3.17)$$

where \mathbf{g}_{ij} is the unit-radius metric on AdS₅. Eq. (3.16) is a direct analog of (2.12) and thus we conclude that we get a complex scalar which is conformally coupled²³ to curvature of AdS₅ × S¹ and is also coupled to a constant U(1) gauge potential $A_{\hat{5}} = 1$ with charge 2

$$\mathcal{L} = \mathsf{g}^{ij}\partial_i\chi\partial_j\bar{\chi} + (\partial_{\hat{5}}\chi - 2i\chi)(\partial_{\hat{5}}\bar{\chi} + 2i\bar{\chi}) - 4\bar{\chi}\chi \ . \tag{3.18}$$

Expanding in modes in S^1 coordinate $\xi^{\hat{5}}$ we get 2 towers of real scalars on AdS₅ with masses

$$m^2 = (n \pm 2)^2 - 4$$
, $n = 0, \pm 1, \pm 2, \dots$, (3.19)

and thus with 4d boundary dimensions $\Delta(\Delta - 4) = m^2$. Assuming as in the D3 brane case (cf. (2.15)) the Dirichlet boundary conditions, i.e. $\Delta = \Delta_+$ we get

$$\Delta - 2 = |n \pm 2| . \tag{3.20}$$

²³In 6d a conformally coupled scalar has kinetic operator $-\nabla^2 + \frac{1}{5}R$ and $R(AdS_5 \times S^1) = -20$ (cf. footnote 2).

The analysis of the remaining fluctuations in θ and S^2 directions is again analogous to the D3 brane case in (2.16),(2.17): setting $\theta = \frac{\pi}{2} + v$, $ds_{S^2}^2 = d\varphi_1^2 + \cos^2 \varphi_1 \ d\varphi_2^2$ and introducing 3 Cartesian coordinates $X^a = \{x, y, z\}$ as $v = \sqrt{x^2 + y^2 + z^2}$, $\tan \varphi_1 = \frac{x}{\sqrt{y^2 + z^2}}$, $\tan \varphi_2 = \frac{z}{y}$, we find that the quadratic fluctuation action for X^a is

$$\int d^4 \xi \sqrt{-\mathsf{g}} \sum_{a=1}^3 \left(\mathsf{g}^{\alpha\beta} \partial_\alpha \mathsf{X}^a \partial_\beta \mathsf{X}^a - 4 \mathsf{X}^a \mathsf{X}^a \right) \,, \tag{3.21}$$

which describes 3 real conformally coupled scalars on $AdS_5 \times S^1$. Expanding in modes in $\xi^{\hat{5}}$ we get 3 towers of scalar operators with masses and scaling dimensions given by (assuming Dirichlet boundary conditions)

 $m^2 = n^2 - 4$, $\Delta - 2 = |n|$, $n = 0, \pm 1, \dots$. (3.22)

3.2 Fermionic fluctuations

The quadratic fermionic part of the κ -symmetric M5 brane action in a general background which is solution of 11d supergravity may be written as $[36-38, 41, 42]^{24}$

$$S_{f} = \int d^{6}\xi \,\sqrt{-G} \left[G^{\alpha\beta} \,\partial_{\alpha} X^{m} \,\bar{\vartheta} \,\Gamma_{m} \widehat{D}_{\beta} \vartheta \right. \\ \left. - \frac{1}{5!} \epsilon^{\mu\nu\kappa\lambda\sigma\rho} \partial_{\mu} X^{m} \partial_{\nu} X^{n} \partial_{\kappa} X^{k} \partial_{\lambda} X^{l} \partial_{\sigma} X^{r} \,\bar{\vartheta} \,\Gamma_{mnklr} \widehat{D}_{\rho} \vartheta \right] \,, \tag{3.23}$$

$$G_{\alpha\beta} = \partial_{\alpha} X^{m} \partial_{\beta} X^{n} G_{mn}(X) , \quad G_{mn} = E^{\underline{a}}_{\overline{m}} E^{\underline{b}}_{\overline{n}} \eta_{\underline{a}\underline{b}} , \quad \Gamma_{m} = E^{\underline{a}}_{\overline{m}}(X) \Gamma_{\underline{a}} , \quad \Gamma_{\alpha} = \partial_{\alpha} X^{m} \Gamma_{m} ,$$

$$\widehat{D}_{\alpha} = \partial_{\alpha} X^{m} D_{m}, \qquad D_{m} = \nabla_{m} - \frac{1}{288} (\Gamma^{pnkl}_{\ m} + 8\Gamma^{pnk} \delta^{l}_{m}) F_{pnkl} .$$
(3.24)

Here we ignored dependence on $H_{\mu\nu\lambda}$ that is not relevant in the present case. As in (2.20) we omitted the overall factor of brane tension that can be absorbed into a rescaling of ϑ . We use Minkowski notation with ϑ being a 32 component 11d Majorana spinor.²⁵ D_m is the generalized 11d spinor covariant derivative [46] and $\nabla_m = \partial_m + \frac{1}{4}\Gamma_{\underline{ab}}\Omega_m^{\underline{ab}}$.

Specifying to the bosonic background (3.9) we may use the background value $g_{\alpha\beta}$ of the induced metric action (3.23) may be rewritten as (cf. (2.20),(2.24))

$$S_f = \int d^6 \xi \sqrt{-g} \, g^{\alpha\beta} \, \bar{\vartheta} (1 - \Gamma_*) \Gamma_\alpha \widehat{D}_\beta \vartheta \, , \qquad (3.25)$$

$$\Gamma_* \equiv \frac{\epsilon^{\alpha\beta\gamma\mu\nu\sigma}}{6!\sqrt{-g}} \partial_{\alpha} X^m \partial_{\beta} X^n \partial_{\gamma} X^k \partial_{\mu} X^l \partial_{\nu} X^p \partial_{\sigma} X^q \Gamma_{mnklpq} = \frac{\epsilon^{\alpha\beta\gamma\mu\nu\sigma}}{6!\sqrt{-g}} \Gamma_{\alpha\beta\gamma\mu\nu\sigma} = \Gamma_{\underline{\hat{0}}\underline{\hat{1}}\underline{\hat{2}}\underline{\hat{3}}\underline{\hat{1}}\underline{\hat{5}}} , \quad (3.26)$$

²⁴Note that in the particular cases of the maximally supersymmetric $AdS_4 \times S^7$ or $AdS_7 \times S^4$ backgrounds the fermionic part of the M5 brane action may be written in an explicit form including also higher orders in ϑ [43, 44, 42, 45].

²⁵To recall (cf. footnote 5), we use Latin letters m, n, \cdots to label the spacetime coordinates, and Greek letters α, β, \ldots to label the world-volume coordinates. When numbering the coordinates, $0, 1, \cdots, 9, 10$ will be used for the spacetime, while hatted numbers $\hat{0}, \hat{1}, \cdots$ – for the world-volume indices. For both spacetime and world-volume indices we use underlined letters to denote indices along the tangent directions.

where $\Gamma_* \equiv \Gamma_{M5}$ is the analog of $\tilde{\Gamma}_{D3}$ in (2.20) with $\Gamma^2_* = 1$. We shall label the coordinates as follows:

$$AdS_{5} = \{X^{0}, ..., X^{4}\} = \{\xi^{\hat{0}}, ..., \xi^{\hat{4}}\}, \qquad X^{5} = u = u_{0}, \qquad X^{6} = \psi = \xi^{\hat{5}}, X^{7} = \theta = \frac{\pi}{2}, \qquad X^{8} = \phi = 2\xi^{\hat{5}}, \qquad S^{2} = \{X^{9}, X^{10}\}.$$
(3.27)

Then F_4 in (3.2) may be written as

$$F_4 = 3L^3 \sin\theta \cos^2\theta \, d\theta \wedge d\phi \wedge \operatorname{vol}_{S^2} = 3L^{-1} \, E^{\underline{7}} \wedge E^{\underline{8}} \wedge E^{\underline{9}} \wedge E^{\underline{10}} \,. \tag{3.28}$$

For m < 7, the second term in brackets in D_m in (3.24) does not contribute and we get

$$D_m = \nabla_m - \frac{1}{288} \Gamma^{pnkl}{}_m F_{pnkl} = \nabla_m - \frac{1}{12} \Gamma^{789\,10}{}_m F_{789\,10} , \qquad m < 7 , \qquad (3.29)$$

$$\sum_{\alpha=\hat{0}}^{4} \Gamma^{\alpha} D_{\alpha} = \sum_{m=0}^{4} \Gamma^{m} D_{m} = \sum_{m=0}^{4} \Gamma^{m} \nabla_{m} - \frac{5}{4L} \Gamma^{\underline{78910}} , \qquad (3.30)$$

$$D_6 = \nabla_6 - \frac{1}{4L} \,\Gamma^{\underline{78910}}_{6} \,. \tag{3.31}$$

For m = 8 the first term in brackets in D_m in (3.24) does not contribute and we get

$$D_8 = \nabla_8 - \frac{1}{36} \Gamma^{pnk} F_{pnk8} = \nabla_8 - \frac{1}{6} \Gamma^{79\ 10} F_{789\ 10} = \nabla_8 - \frac{1}{2L} E_8^{\underline{8}} \Gamma^{\underline{79\ 10}} , \qquad (3.32)$$

$$\partial_{\hat{5}} X^m D_m = D_6 + 2D_8 = \nabla_6 + 2\nabla_8 - \frac{1}{4L} \Gamma^{\underline{78910}}_{6} - \frac{1}{L} \Gamma^{\underline{78910}}_{-8} , \qquad (3.33)$$

$$\Gamma^{\hat{5}}\partial_{\hat{5}}X^{m}D_{m} = \Gamma^{\hat{5}}(D_{6} + 2D_{8}) , \qquad \Gamma^{\hat{5}} = \frac{\sinh u_{0}\Gamma_{\underline{6}} + \Gamma_{\underline{8}}}{L_{A}\cosh^{2}u_{0}} .$$
(3.34)

Also, computing the spin connection gives (see (B.15) in Appendix B)

$$\Gamma^{\alpha}\partial_{\alpha}X^{m}\nabla_{m} = \nabla + \frac{3}{2L}\tanh u_{0}\Gamma_{\underline{5}} + \frac{1}{4L\cosh u_{0}}\Gamma_{\underline{865}} , \qquad (3.35)$$

where ∇ corresponds to $\mathrm{AdS}_5 \times S^1$.

Like in the D3 case one expects that the resulting fermionic action should have no nontrivial dependence on u_0 . Indeed, in Appendix C we will show that the fermion spectrum is the same for all values of u_0 . Thus, to simplify the presentation, here we may just consider the limiting case of $u_0 \rightarrow 0$ ignoring the subleading terms. Then in total

$$\Gamma^{\alpha}\partial_{\alpha}X^{m}D_{m} = \nabla + \frac{1}{2L}\Gamma_{\underline{8}}\left(\frac{1}{2}\Gamma_{\underline{65}} + \frac{3}{2}\Gamma_{\underline{79}10}\right).$$
(3.36)

The fermionic action (3.25) then may be written as²⁶

$$S_f = \int d^6 \xi \sqrt{-\mathsf{g}} \,\bar{\vartheta} (1 - \Gamma_*) \big(\nabla \!\!\!/ + \mathrm{M} \big) \vartheta \,\,, \tag{3.37}$$

$$\Gamma_* = \Gamma_{\underline{\hat{0}}\cdots\underline{\hat{4}}\underline{\hat{5}}} = \Gamma_{\underline{0}\cdots\underline{48}} , \qquad M = \frac{1}{2}\Gamma_{\underline{8}} \Big(\Gamma_{\underline{65}} + 3\Gamma_{\underline{79}\,\underline{10}} \Big) . \qquad (3.38)$$

 $^{^{26}}$ We scale out $L_A = 2L$ so that $g_{\alpha\beta} \rightarrow \mathbf{g}_{\alpha\beta}$ is the unit-radius $\mathrm{AdS}_5 \times S^1$ metric and ignore the overall constant factor.

Note that Γ_* in (3.25) anticommutes with ∇ and M. We shall fix the κ -symmetry gauge by

$$(1 - \Gamma_*)\vartheta = 0. (3.39)$$

Then acting on ϑ we have $\Gamma_{\underline{0}\dots\underline{48}} = +1$. We may also use that in the conventions assumed in (3.23) one has $\Gamma_{\underline{0}\dots\underline{456789}\,\underline{10}} = +1$ (see (D.17),(D.22)). Then $\Gamma_{\underline{5679}\,\underline{10}} = -1$ or $\Gamma_{\underline{79}\,\underline{10}} = \Gamma_{\underline{56}}$ (see (D.24)). As a result, we may simplify M in (3.38) to

$$M = \frac{1}{2}(1-3)\Gamma_{\underline{865}} = -\Gamma_{\underline{865}} , \qquad (3.40)$$

which is the same result as found in (C.18).

Like in the D3 case (cf. (2.34)) the final expression for the mass operator is given by a combination of the contributions of the "transverse" part of the spin connection and of F_4 in (3.24). After diagonalization of M the resulting action (3.37) or $\int d^6\xi \sqrt{-g} \,\bar{\vartheta}(\nabla + M)\vartheta$ thus describes 2 sets of 6d fermions in $\mathrm{AdS}_5 \times S^1$ with masses ± 1 .

Expanding ϑ in modes in $\xi^{\hat{5}}$ (assuming periodic boundary condition as required by preservation of supersymmetry) the Dirac operator on $\operatorname{AdS}_5 \times S^1$ reduces to that on AdS_5 as (recall that $\Gamma^{\hat{5}} = \Gamma^{\underline{8}}$ in the case of $u_0 \to 0$)

$$i(\nabla + \mathbf{M}) = i \nabla_{AdS_5} + i \Gamma^{\underline{5}} \partial_{\underline{5}} + i \mathbf{M} \quad \rightarrow \quad i \nabla_{AdS_5} - \hat{M} , \qquad \qquad \hat{M} = n \Gamma_{\underline{8}} - i \Gamma_{\underline{568}} . \tag{3.41}$$

Since $\Gamma_8^2 = 1$, $(i\Gamma_{568})^2 = 1$ and $[\Gamma_8, \Gamma_{568}] = 0$ we conclude that \hat{M} has eigenvalues $m_f = \pm n \pm 1.^{27}$ Thus we find 2 towers of 6d fermions with such masses. The corresponding scaling dimensions of the boundary operators are then (cf. (2.36))

$$\Delta - 2 = |m_f| = |n \pm 1| , \qquad n = 0, \pm 1, \pm 2, \dots .$$
(3.42)

3.3 Antisymmetric tensor field contribution

Since the self-dual $H_{\mu\nu\lambda}$ field in the M5 brane action (3.4) has no background value its contribution to the 1-loop free energy is the same as half of that of rank 2 antisymmetric tensor $A_{\mu\nu}$ propagating on $\mathrm{AdS}_5 \times S^1$.

The partition function for $A_{\mu\nu}$ with the standard action $\int d^6 \xi \sqrt{-g} H_{\mu\nu\lambda} H^{\mu\nu\lambda}$ in a general 6d curved background is given by (see, e.g., [47–49] and refs. there)

$$Z_2 = (\det \hat{\Delta}_2)^{-1/2} \det \Delta_1 \ (\det \hat{\Delta}_0)^{-3/2} , \qquad (3.43)$$

Here the Hodge-DeRham operators $\hat{\Delta}_p$ are

$$(\hat{\Delta}_2)^{\alpha\beta}_{\mu\nu} = -\nabla^2 \delta^{\alpha\beta}_{\mu\nu} + 2R^{[\alpha}_{[\mu}\delta^{\beta]}_{\nu]} - R_{\mu\nu}{}^{\alpha\beta} , \qquad (\hat{\Delta}_1)^{\nu}_{\mu} = -\nabla^2 \delta^{\nu}_{\mu} + R^{\nu}_{\mu} , \qquad \hat{\Delta}_0 = -\nabla^2 . \quad (3.44)$$

²⁷Since $[\Gamma_{\underline{8}}, \Gamma_{\underline{568}}] = 0$ the action corresponding to (3.41) may be interpreted as that of massless fermions in AdS₅ × S¹ coupled to U(1) gauge potential in 5 direction with charge 1. This is similar to the D3 brane case in (2.35) where the charge was $\frac{1}{2}$.

Let us specify (3.43) to $AdS_5 \times S^1$ with equal radii =1. For AdS_5 we have $R_{ijkl} = -(g_{ik}g_{jl} - g_{il}g_{jk})$, $R_{ij} = -4g_{ij}$, $R = R_i^i = -20$. Thus from (3.44) splitting $A_{\mu\nu} = (A_{ij}, A_i = A_{i5})$ and also the vector ghost $C_{\mu} = (C_i, C = C_5)$ we get (i = 0, 1, 2, 3, 4)

$$A^{\mu\nu}\hat{\Delta}_2 A_{\mu\nu} = A^{ij}(-\nabla^2 - 6)A_{ij} + A^i(-\nabla^2 - 4)A_i , \qquad (3.45)$$

$$C^{\mu}\hat{\Delta}_{1}C_{\mu} = C^{i}(-\nabla^{2} - 4)C_{i} + C(-\nabla^{2})C , \qquad \nabla^{2} = \nabla^{2}_{AdS_{5}} + \partial^{2}_{5} . \qquad (3.46)$$

We may express det $\hat{\Delta}_1$ as (cf. (2.37) and discussion below it)²⁸

$$\det \hat{\Delta}_1 = \det(-\nabla^2 - 4)_{\perp} \ [\det(-\nabla^2)]^2 \ . \tag{3.47}$$

Similarly, applying the redefinition $A_{ij} = A_{ij}^{\perp} + \nabla_i V_j^{\perp} - \nabla_j V_i^{\perp}$ and accounting for the Jacobian factor we get

$$Z_2(AdS_5 \times S^1) = \left[\frac{(\det \hat{\Delta}_1(-4))^2 (\det \hat{\Delta}_0(0))^2}{\det \hat{\Delta}_2(-6) \det \hat{\Delta}_1(-4) (\det \hat{\Delta}_0(0))^3}\right]^{1/2} = \left[\frac{1}{\det \hat{\Delta}_{2\perp}(-6)}\right]^{1/2}.$$
 (3.48)

Here $\hat{\Delta}_p(X) \equiv -\nabla^2 + X$ are defined on *p*-forms with AdS₅ indices and $\hat{\Delta}_{2\perp}(-6)A_{ij}^{\perp} = (-\nabla^2_{AdS_5} - \partial_5^2 - 6)A_{ij}^{\perp}$.

One can give an alternative derivation of (3.48) as follows. Let us split $H^2_{\mu\nu\lambda} = H^2_{ijk} + 3H^2_{ij5}$ and fix the gauge as $A_{i5} = 0$ so that $H^2_{ij5} = (\partial_5 A_{ij})^2$. Then for $A_{ij} = A^{\perp}_{ij} + \partial_i C^{\perp}_j - \partial_j C^{\perp}_i$ we get $H^2_{ijk} = A^{\perp}_{ij} \hat{\Delta}_2(-6)A^{\perp}_{ij}$. From $(\partial_5 A_{ij})^2$ we find that determinant of ∂_5^2 cancels against the ghost determinant. Including also the contribution of the Jacobian we end up with (3.48). The same expression (3.48) was given also in [50, 51].

Expanding $A_{ij}^{\perp}(\xi^k, \xi^5)$ in S^1 modes we get a tower of transverse antisymmetric tensor fields in AdS₅ with masses

$$m^2 = n^2 - 6 (3.49)$$

Let us recall how this case fits into the general discussion of fields in AdS₅ corresponding to representations of SO(2, 4). Let ϕ be a massive $(\Delta > 2 + j_1 + j_2 \text{ for } j_1 j_2 \neq 0 \text{ or } \Delta > 1 + j_1 + j_2$ for $j_1 j_2 = 0$) or massless $(\Delta = 2 + j_1 + j_2, \ j_1 j_2 \neq 0)$ field in AdS₅ corresponding the SO(2, 4)representation $(\Delta; j_1, j_2)$. $h_1 = j_1 + j_2 = s$ and $h_2 = j_1 - j_2$ are integer for bosons and half-integer for fermions (in the bosonic case, h_1 and $|h_2|$ are the lengths of a two-row Young tableau). According to [52, 53], the covariant AdS₅ equation of motion for a bosonic transverse field ϕ is (for $j_1 \geq j_2$)²⁹

$$\mathcal{O}\phi = 0$$
, $\mathcal{O} = -\nabla_{AdS_5}^2 + m^2$, $m^2 = (\Delta - 2)^2 - 4 - 2j_1$. (3.50)

 $\frac{1}{2^8 \text{Set } A_i = A_i^{\perp} + \partial_i \varphi. \text{ We use that } \int dA e^{-\int A^2} \to \int dA_{\perp} \, d\varphi \, \sqrt{\det(-\nabla^2)} \, e^{-\int A_{\perp}^2} \text{ and } A^i(-\nabla^2 - 4)A_i = A_i^{\perp}(-\nabla^2 - 4)A_i^{\perp} + \varphi(-\nabla^2)^2 \varphi. \text{ Note also that for } \nabla_i \text{ derivatives } -\nabla^i(-\nabla^2)\nabla_i = \nabla^4 + \nabla^i[\nabla^2, \nabla_i] = \nabla^4 - 4\nabla^2.$

²⁹This equation is also for the fermionic fields after squaring the 5d Dirac operator. For a generic fermion spinor-tensor field Ψ one has $(\nabla + \Delta - 2) \Psi = 0$ [54]. After squaring, this turns out to be $\left[-\nabla_{AdS_5}^2 + \frac{1}{4}R - 2j_1 + 1 + (\Delta - 2)^2 \right] \Psi = 0$, where $R = R(AdS_5) = -20$. This gives the same m^2 as in (3.50).

The partition function for such massive field is then given by 30

$$Z(\Delta; j_1, j_2) = \left[\det(-\nabla_{AdS_5}^2 + m^2)_{\perp}\right]^{-1/2}.$$
(3.51)

The antisymmetric tensor case in (3.48),(3.49) corresponds to the sum of two (self-dual and anti self-dual) representations (cf. (3.50))

$$(\Delta; 1, 0) + (\Delta; 0, 1)$$
, $\Delta - 2 = |n|$. (3.52)

The collection of 2 scalars with masses/dimensions in (3.19),(3.20), 3 scalars in (3.22), the fermions in (3.42) and the self-dual rank 2 tensor (3.52) form 5d supermultiplets that represent the (2,0) multiplet defined on $AdS_5 \times S^1$ in a way consistent with preservation of supersymmetry, i.e. requiring conformal coupling of all scalars to the curvature and a particular coupling of 2 scalars and all fermions to a constant U(1) gauge potential (cf. [55, 35, 56]).³¹

3.4 1-loop free energy

The free energy corresponding to (3.51) can be computed explicitly like in the AdS₃ case in (2.41)-(2.43). In the case of the Euclidean AdS₅ with boundary S^4 it is proportional to $vol(AdS_5) = \pi^2 log(r\Lambda)$ and thus is proportional to the 4d conformal *a*-anomaly coefficient (see [22, 59, 25, 27] and, in particular, [28])

$$F(\Delta; j_1, j_2) = \frac{1}{2} \log \det(-\nabla_{AdS_5}^2 + m^2)_{\perp} = -\frac{1}{2} \zeta'_{(\Delta; j_1, j_2)}(0) = 4a(\Delta; j_1, j_2) \log(r\Lambda)$$
(3.53)

$$a(\Delta; j_1, j_2) = -\frac{1}{96\pi} (-1)^{2(j_1+j_2)} (2j_1+1)(2j_2+1) \lim_{z \to 0} J'(z) , \qquad (3.54)$$

$$J(z) = \int_0^\infty d\lambda \, \frac{\left[\lambda^2 + (j_1 - j_2)^2\right] \left[\lambda^2 + (j_1 + j_2 + 1)^2\right]}{\left[\lambda^2 + (\Delta - 2)^2\right]^z} \,, \qquad (\Delta - 2)^2 = m^2 + 4 + 2j_1 \,.$$

This gives, assuming Dirichlet boundary conditions so that $\Delta - 2 \ge 0$,

$$a(\Delta; j_1, j_2) = \frac{1}{1440} (-1)^{2(j_1+j_2)} (2j_1+1)(2j_2+1) (\Delta - 2) \\ \times \left[3(\Delta - 2)^4 - 10(j_1^2 + j_2^2 + j_1 + j_2 + \frac{1}{2})(\Delta - 2)^2 + 15(j_1 - j_2)^2(j_1 + j_2 + 1)^2 \right].$$
(3.55)

As a result, we get for a real AdS_5 scalar contribution

$$a(\Delta; 0, 0) = \frac{1}{1440} (\Delta - 2) \left[3 (\Delta - 2)^4 - 5(\Delta - 2)^2 \right] .$$
 (3.56)

³⁰In the massless case of $\Delta = 2 + s$ one needs to take into account the contribution of the corresponding ghosts that belong to the representation $(\Delta + 1; j_1 - \frac{1}{2}, j_2 - \frac{1}{2})$ (see, e.g., [25]).

³¹It is interesting to note that by applying an analytic continuation to $S^5 \times S^1$ with the angle of S^1 identified with period β one gets a similar system of fields whose supersymmetric partition function computes the Schur index [57] of the (2,0) tensor multiplet (see section 5 in [20]). Similar relation is true also in the D3 brane case in section 2 upon analytic continuation to $S^3 \times S^1$ world volume theory (cf. [58]).

The contribution of the self-dual antisymmetric tensor is³²

$$a(\Delta; 1, 0) = \frac{3}{1440} (\Delta - 2) \left[3 (\Delta - 2)^4 - 25(\Delta - 2)^2 + 60 \right].$$
 (3.57)

The fermion contribution is

$$a(\Delta; \frac{1}{2}, 0) = a(\Delta; 0, \frac{1}{2}) = -\frac{2}{1440} \left(\Delta - 2\right) \left[3\left(\Delta - 2\right)^4 - \frac{25}{2}\left(\Delta - 2\right)^2 + \frac{135}{16}\right].$$
 (3.58)

Let us first consider the case of (2,0) multiplet of 5 conformally coupled scalars, self-dual tensor and 4 fermions propagating on $\operatorname{AdS}_5 \times S^1$ with all scalars and fermions not coupled to a U(1) potential in 5 direction, i.e. without shift of S^1 mode number n. Expanding in Fourier modes in this case we get system of AdS_5 fields with all the fields having $\Delta - 2 = |n|$. Thus we get for the free energy in $\operatorname{AdS}_5 \times S^1$ (cf. (2.44),(3.14))

$$F = \frac{2}{\pi^3} a \operatorname{vol}(AdS_5 \times S^1) = 4a \log(r\Lambda) , \qquad (3.59)$$

$$a_{(2,0)} = \sum_{n=-\infty}^{\infty} \left[5a(\Delta;0,0) + a(\Delta;1,0) + 4a(\Delta;\frac{1}{2},0) \right] = \frac{1}{1440}P , \qquad P = \sum_{n=-\infty}^{\infty} P_n , \quad (3.60)$$

$$P_n = 5|n|(3n^4 - 5n^2) + 3|n|(3n^4 - 25n^2 + 60) - 8|n|(3n^4 - \frac{25}{2}n^2 + \frac{135}{16}) = \frac{225}{2}|n|, \quad (3.61)$$

$$P = 225 \sum_{n=1}^{\infty} n = 225 \zeta_R(-1) = -\frac{225}{12} , \qquad a_{(2,0)} = -\frac{5}{384} . \qquad (3.62)$$

In contrast to the case of the $\mathcal{N} = 4$ multiplet in $\mathrm{AdS}_3 \times S^1$ in (2.44),(2.45) where the 1-loop free energy was UV finite and vanishing here $F^{(1)}$ is quadratically divergent. As in other similar examples (see, e.g., [17–19]) we used the Riemann ζ -function regularization to compute the resulting sum.

The presence of the quadratic UV divergence was, in fact, expected. In 6d the free energy $F^{(1)} = \frac{1}{2} \log \det(-\nabla^2 + X)$ has a UV divergent part given by (in heat kernel regularization)

$$F_{\infty}^{(1)} = \frac{1}{(4\pi)^3} \int d^6 \xi \sqrt{g} \left(\frac{1}{6} b_0 \Lambda^6 + \frac{1}{4} b_2 \Lambda^4 + \frac{1}{2} b_4 \Lambda^2 + b_6 \log \Lambda \right) , \qquad (3.63)$$

where b_k are the Seeley's coefficients. $b_0 = \text{tr1}$ counts total number of degrees of freedom and thus vanishes for a supersymmetric model. One can check that $b_2 = \text{tr}(\frac{1}{6}R - X)$ also vanishes in a combination of 5 conformally coupled scalars ($\hat{\Delta}_0 = -\nabla^2 + \frac{1}{5}R$), 4 massless

³²The contribution of the self-dual antisymmetric tensor to the 1-loop free energy is by definition half that of the standard antisymmetric tensor. If one dimensionally reduces the antisymmetric tensor action $\int H^2_{\mu\nu\lambda}$ to AdS₅ (i.e. considers only the n = 0 mode of the S^1 expansion) one gets a collection of a rank 2 tensor $\int H^2_{ijk}$ and massless vector $\int F^2_{ij} (A_i \equiv A_{i5})$ 5d actions. Dualizing the former to a vector A'_i (which can be done by a path integral transformation and thus preserves the expression for the partition function modulo zero mode contribution absent in the present case) one thus gets a collection of two 5d vectors. Their total contribution to *a*-anomaly, is however, zero as their boundary conditions are opposite (cf. footnote 15 for a similar remark in the AdS₃ context). Thus the total contribution to *a*-anomaly of the n = 0 mode of the antisymmetric tensor is zero, in agreement with the general expressions in (3.61),(3.65).

fermions $(\hat{\Delta}_{1/2} = -\nabla^2 + \frac{1}{4}R)$ and the self-dual antisymmetric tensor in (3.43),(3.44). This is consistent with the cancellation of n^3 terms in (3.61).

At the same time, one finds that b_4 and b_6 coefficients do not vanish in general. The b_6 coefficient for the (2,0) multiplet (that determines its conformal anomaly) was explicitly computed in [49] and is given by a combination of the 6d Euler density and 3 cubic invariants built out of the 6d Weyl tensor. The Euler density vanishes for a space like $M^5 \times S^1$ and the Weyl tensor vanishes in the case of the conformally flat $AdS_5 \times S^1$ space. This is consistent with the absence of the log UV divergence in (3.60). Complementing the discussion in [49] and computing b_4 for the (2,0) multiplet on a general curved 6d space we get³³

$$b_4 = \frac{1}{4}R_{\mu\nu\lambda\rho}^2 - R_{\mu\nu}^2 + \frac{1}{10}R^2 . \qquad (3.64)$$

Computing this for $AdS_5 \times S^1$ we get non-vanishing result for b_4 , in agreement with noncancellation of order n terms in (3.61) leading to quadratic UV divergence.³⁴

Let us now turn to the case of our interest when the (2,0) multiplet on $AdS_5 \times S^1$ originates from the supersymmetric M5 brane embedded into $AdS_7 \times S^4$ and is thus coupled also to an effective constant U(1) gauge potential in 5 direction (with 2 scalars having charge ± 2 and the fermions charge ± 1 which results in the shifts of n in (3.20), (3.42)). In this case P_n in (3.60) can be written as

$$P_{n} = |n-2| [3(n-2)^{4} - 5(n-2)^{2}] + |n+2| [3(n+2)^{4} - 5(n+2)^{2}] + 3|n|(3n^{4} - 5n^{2}) + 3|n|(3n^{4} - 25n^{2} + 60) - 4|n+1| [3(n+1)^{4} - \frac{25}{2}(n+1)^{2} + \frac{135}{16}] - 4|n-1| [3(n-1)^{4} - \frac{25}{2}(n-1)^{2} + \frac{135}{16}].$$
(3.65)

As a result,

$$P_0 = \frac{241}{2}, \qquad P_1 = \frac{1297}{2}, \qquad P_{|n|>1} = \frac{1305}{2}|n|.$$
 (3.66)

The large n asymptotics of P_n is again $\sim n$ as in (3.61) consistent with expected presence of a quadratic UV divergence.³⁵ Then (cf. (3.62))

$$P = \frac{241}{2} + 1297 + 1305 \sum_{n=2}^{\infty} n = \frac{15}{4} , \qquad a^{(1)} = \frac{1}{1440} P = \frac{1}{384} . \qquad (3.67)$$

³³In a general number of dimensions for an operator $\hat{\Delta} = -\nabla^2(A) + X$ defined on a vector bundle with connection A_{μ} one has (up to a total derivative term) $b_4 = \text{tr} \left[\frac{1}{12}F_{\mu\nu}^2 + \frac{1}{180}(R_{\mu\nu\lambda\rho}^2 - R_{\mu\nu}^2) + \frac{1}{2}(\frac{1}{6}R - X)^2\right]$. ³⁴The fact that b_4 is non-vanishing for 6d (2,0) multiplet is analogous to non-vanishing of b_4 for the 4d

 $[\]mathcal{N} = 4$ multiplet.

 $^{^{35}}$ Note that coupling to a constant UV gauge field should not a priori change the values of the coefficients b_p of the UV divergent terms (3.63). This is true, however, if one uses a covariant 6d regularization which is not the case here. Here we first expand in S^1 modes, then define the resulting 5d determinants using spectral ζ -function and at the end sum over n. A cut off on n is obviously not covariant in 6d and thus the structure of (subleading) power divergences here is a priori sensitive to shifts of n.

Here like in (3.62) we used the ζ_R -function to define the sum. Combining this 1-loop value of *a* with the classical one in (3.14) we thus get the following prediction for the S^4 defect anomaly coefficient

$$a = N^2 + \frac{1}{384} + \mathcal{O}(N^{-2}) . aga{3.68}$$

This result can not be directly compared with the defect anomaly coefficient computed in [8] using the bubbling solution in supergravity where it has N^3 scaling at leading order. The precise limit of the parameters of the solution in [8] that corresponds to the probe limit in which the anomaly coefficient should scale as N^2 remains to be understood.³⁶

Acknowledgements

We are grateful to M. Beccaria, N. Drukker, S. Giombi, C. Herzog and R. Metsaev for useful discussions and comments, and also thank O. Aharony, A. Chalabi and D. Sorokin for related correspondence. This work was supported in part by the STFC Consolidated Grants ST/T000791/1 and ST/X000575/1.

A Scalar fluctuations of a *p*-brane in $AdS_{p+2} \times S^1$

The scalar fluctuations of a *p*-brane embedded in a supersymmetric way in $AdS_{p+2} \times S^1$ can be computed in a universal way for any *p*, thus covering the cases of p = 3 and 5 discussed in the main text. The relevant part of the background is

$$ds^{2} = L_{A}^{2} \left(du^{2} + \cosh^{2} u \, ds_{AdS_{p}}^{2} + \sinh^{2} u \, d\psi^{2} \right) + L^{2} d\phi^{2} , \qquad (A.1)$$

$$C_{p+1} = L_A^{p+1}(\cosh^{p+1} u - 1) \operatorname{vol}_{AdS_p} \wedge d\psi \equiv \mathsf{C}_{p+1} \operatorname{vol}_{AdS_p} \wedge d\psi .$$
(A.2)

The configuration of the probe *p*-brane related to co-dimension 2 half-supersymmetric defect in the boundary theory is such that it wrapps AdS_p and also (see [13] and refs. there)

$$u = u_0 = \text{const}$$
, $L_A \psi = L \phi$. (A.3)

The induced metric on the probe p-brane is then that of the equal-radii $AdS_p \times S^1$ space

$$ds_{p+1}^2 = L_A^2 \cosh^2 u_0 \, ds_{AdS_p}^2 + L_A^2 \sinh^2 u_0 d\psi^2 + L_A^2 d\psi^2 = L_A^2 \cosh^2 u_0 \, \left(ds_{AdS_p}^2 + d\psi^2 \right) \,. \tag{A.4}$$

Starting with the standard action $S \sim -\int d^{p+1}\xi \sqrt{-G} + \int C_{p+1}$ of a probe *p*-brane in the background (A.1),(A.2) let us find the resulting quadratic fluctuation action in the static gauge

$$AdS_p = \{\xi^{\alpha}\}, \qquad u = u_0 + \delta u, \qquad \psi = \xi^{\hat{p}}, \qquad \phi = \frac{L_A}{L}\xi^{\hat{p}} + \delta\phi.$$
 (A.5)

Then the induced metric $G_{\alpha\beta}$ has the following quadratic fluctuation part

$$ds_G^2 = L_A^2 \cosh^2(u_0 + \delta u)(ds_{AdS_p}^2 + d\psi^2) + 2L_A L d\delta\phi d\psi + (L_A^2 d\delta u^2 + L^2 d\delta\phi^2) + \dots$$
(A.6)

³⁶We thank J. Estes, B. Suzzoni and P. Capuozzo for a correspondence on this issue.

This leads to^{37}

$$\sqrt{-G} = \sqrt{-g} l_0^{p+1} \left[1 + \left(\frac{L}{L_A \cosh^2 u_0} \partial_{\psi} \delta \phi + (p+1) \tanh u_0 \, \delta u \right) + \frac{1}{2 \cosh^2 u_0} \left(\mathsf{g}^{\alpha\beta} \partial_{\alpha} \delta u \, \partial_{\beta} \delta u + \frac{L^2 \tanh^2 u_0}{L_A^2} \mathsf{g}^{\alpha\beta} \partial_{\alpha} \delta \phi \, \partial_{\beta} \delta \phi + (p+1) (\cosh^2 u_0 + p \sinh^2 u_0) \, \delta u^2 + \frac{2L(p-1) \tanh u_0}{L_A} \delta u \, \partial_{\psi} \delta \phi \right) + \cdots \right],$$
(A.7)

where $l_0 = L_A \cosh u_0$ and $g_{\alpha\beta}$ is the metric of the unit-radius $\operatorname{AdS}_p \times S^1$. For the variation of the potential C_{p+1} in (A.2) we get

$$\mathsf{C}_{p+1} = -L_A^{p+1} + l_0^{p+1} \Big[1 + (p+1) \tanh u_0 \,\delta u + \frac{1}{2} (p+1)(1+p \tanh^2 u_0) \,\delta u^2 + \cdots \Big] \,. \tag{A.8}$$

Combined with (A.7) this gives

$$\sqrt{-G} - \mathsf{C}_{\mathsf{p}+1} = L_A^{p+1} + l_0^{p+1} \Big[\frac{L}{L_A \cosh^2 u_0} \partial_{\psi} \delta\phi + \frac{1}{2 \cosh^2 u_0} \Big(\mathsf{g}^{\alpha\beta} \partial_{\alpha} \delta u \, \partial_{\beta} \delta u \\ + \frac{L^2 \tanh^2 u_0}{L_A^2} \mathsf{g}^{\alpha\beta} \partial_{\alpha} \delta\phi \, \partial_{\beta} \delta\phi + 2(p-1) \frac{L \tanh u_0}{L_A} \delta u \, \partial_{\psi} \, \delta\phi \Big) + \cdots \Big] .$$
(A.9)

As expected, the linear fluctuation part is a total derivative and the quadratic fluctuation part of the action is given by

$$S - S_0 = -\frac{l_0^{p+1}}{2\cosh^2 u_0} \int d^{p+1}\xi \,\sqrt{-\mathsf{g}} \Big[\mathsf{g}^{\alpha\beta}\partial_\alpha\chi\partial_\beta\bar{\chi} + \frac{i}{2}(p-1)(\bar{\chi}\partial_\psi\chi - \chi\partial_\psi\bar{\chi}) \Big] \,, \qquad (A.10)$$

$$\chi \equiv \delta u + i \frac{L}{L_A} \tanh u_0 \,\delta\phi \;. \tag{A.11}$$

Rescaling χ we find that a canonically normalized scalar fluctuation action is

$$\frac{1}{2}\int d^{p+1}\xi\sqrt{-\mathsf{g}}\left(\mathsf{g}^{ij}\partial_i\chi\partial_j\bar{\chi} + \left[\partial_\psi\chi - \frac{i}{2}(p-1)\chi\right]\left[\partial_\psi\bar{\chi} + \frac{i}{2}(p-1)\bar{\chi}\right] - \frac{1}{4}(p-1)^2\chi\bar{\chi}\right), \quad (A.12)$$

where i, j are indices of the AdS_p part of the brane metric.

This action describes a conformally coupled complex scalar on $\operatorname{AdS}_p \times S^1$ coupled also to a constant U(1) potential $A = d\xi^{\hat{p}}$ with charge $q = \frac{1}{2}(p-1)$. Fourier expanding in $\psi = \xi^{\hat{p}}$, i.e. $\chi = \sum_n e^{in\xi^{\hat{p}}} \chi_n$ we get a tower of scalars on AdS_p with masses

$$m^2 = \left(n - \frac{p-1}{2}\right)^2 - \left(\frac{p-1}{2}\right)^2, \qquad n = 0, \pm 1, \pm 2, \dots.$$
 (A.13)

The corresponding boundary dimensions are defined by $\Delta[\Delta - (p-1)] = m^2$ or for the Dirichlet boundary condition choice $(\Delta = \Delta_+)$ we get

$$\Delta - \frac{p-1}{2} = |n - \frac{p-1}{2}| . \tag{A.14}$$

Since χ is complex, we actually have two towers of states with opposite shift, namely $\Delta - \frac{p-1}{2} = |n \pm \frac{p-1}{2}|$.

 $[\]overline{{}^{37}\text{We use that for } \tilde{g}_{\mu\nu} = g_{\mu\nu} + h_{\mu\nu} \text{ one has } \sqrt{-\tilde{g}} = \sqrt{-g} \left(1 + \frac{1}{2}h^{\mu}{}_{\mu} + \frac{1}{8}(h^{\mu}{}_{\mu})^2 - \frac{1}{4}h_{\mu\nu}h^{\mu\nu} + \cdots\right). \text{ Let us note also that given a metric } ds_g^2 = g_{\mu\nu}dx^{\mu}dx^{\nu} = g_{ab}dx^adx^b + g_{\theta\theta}d\theta^2 \ (a \neq \theta) \text{ and } ds_{\tilde{g}}^2 = ds_g^2 + dFd\theta \text{ where } F = F(x) \text{ is a function of } x^{\mu} \text{ then } \sqrt{-\tilde{g}} = \sqrt{-g} \left(1 + \frac{1}{2}g^{\theta\theta}\partial_{\theta}F - \frac{1}{8}g^{\theta\theta}g^{\mu\nu}\partial_{\mu}F\partial_{\nu}F\right).$

B Spin connection and projected spinor covariant derivative

Here we will compute the spin connection contribution to the induced fermionic covariant derivative. Let us consider the following metric (cf. (A.1))

$$ds^{2} = L_{A}^{2} \left(du^{2} + \cosh^{2} u \, ds_{AdS_{p}}^{2} + \sinh^{2} u \, d\psi^{2} \right) + L^{2} \left(d\theta^{2} + \sin^{2} \theta d\phi^{2} \right) \,. \tag{B.1}$$

We will label the target space coordinates as follows:

$$X^{0,1,\dots,p-1} = AdS_p , \quad X^p = u , \quad X^{p+1} = \psi , \quad X^{p+2} = \theta , \quad X^{p+3} = \phi .$$
 (B.2)

We shall assume that a *p*-brane wraps $AdS_p \times S^1$ as in (A.3). Since $\theta = \frac{\pi}{2}$ the resulting induced metric on the brane is the same as in (A.4). For this classical brane configuration we have (we will ignore bosonic fluctuations here, cf. (A.5))

$$X^{0,1,\dots,p-1} = AdS_p = X^{\hat{0},\hat{1},\dots,\hat{p-1}}, \quad X^p = u = u_0, \quad X^{\widehat{p+1}} = \psi = \xi^{\widehat{p}}, \quad (B.3)$$

$$X^{p+2} = \theta = \frac{\pi}{2}$$
, $X^{p+3} = \phi = \frac{L_A}{L}\xi^{\hat{p}}$. (B.4)

The vielbein components for (B.1) can be chosen as

$$E_m^{\underline{m}} = L_A \cosh u \, \hat{E}_m^{\underline{m}} \,, \qquad m = 0, ..., p - 1$$
 (B.5)

$$E_{p}^{p} = L_{A} , \quad E_{p+1}^{p+1} = L_{A} \sinh u , \quad E_{p+2}^{p+2} = L , \quad E_{p+3}^{p+3} = L \sin \theta ,$$
 (B.6)

where the underlined indices correspond to tangent space directions and \hat{E}_m^m is the vielbein for the unit-radius AdS_p . The corresponding spin connection components along the normal directions to the brane world volume are found to be

$$\Omega_m^{\underline{m}\,\underline{p}} = \frac{\tanh u}{L_A} E_m^{\underline{m}} = \sinh u \,\hat{E}_m^{\underline{m}} \,, \qquad m = 0, ..., p - 1 \,, \tag{B.7}$$

$$\Omega_{p+1}^{\underline{p+1}\underline{p}} = \frac{1}{L_A \tanh u} E_{p+1}^{\underline{p+1}} = \cosh u , \qquad \Omega_{p+3}^{\underline{p+3}\underline{p+2}} = \cos \theta . \tag{B.8}$$

The projected covariant derivative and Γ -matrix are given by

$$\partial_{\alpha} X^m \nabla_m = \partial_{\alpha} X^m \left(\partial_m + \frac{1}{4} \Omega_m^{\underline{ab}} \Gamma_{\underline{ab}} \right) , \qquad \Gamma_{\alpha} = \partial_{\alpha} X^m \Gamma_m . \tag{B.9}$$

When $\alpha = \hat{0}, \cdots, \widehat{p-1}$, we have $\partial_{\alpha} X^m = \delta^m_{\alpha}$, so

$$\partial_{\alpha} X^m \nabla_m = \partial_{\alpha} + \frac{1}{4} \Omega^{\underline{a}\underline{b}}_{\alpha} \Gamma_{\underline{a}\underline{b}} = \partial_{\alpha} + \frac{1}{4} \Omega^{\underline{\beta}\underline{\gamma}}_{\alpha} \Gamma_{\underline{\beta}\underline{\gamma}} + \frac{1}{2} \sinh u_0 \hat{E}^{\underline{\alpha}}_{\alpha} \Gamma_{\underline{\alpha}\underline{p}} .$$
(B.10)

Since $\Gamma^{\alpha} = E^{\alpha}_{\underline{\alpha}}\Gamma^{\underline{\alpha}} = \frac{1}{L_A \cosh u_0} \hat{E}^{\alpha}_{\underline{\alpha}}\Gamma^{\underline{\alpha}}$, we have

$$\Gamma^{\alpha}\partial_{\alpha}X^{m}\nabla_{m} = \bigvee_{AdS_{p}} + \frac{1}{2L_{A}\cosh u_{0}}\sinh u_{0}\hat{E}_{\alpha}^{\underline{\alpha}}\hat{E}_{\underline{\alpha}'}^{\underline{\alpha}'}\Gamma^{\underline{\alpha}'}\Gamma_{\underline{\alpha}\underline{p}} = \bigvee_{AdS_{p}} + \frac{p}{2L_{A}}\tanh u_{0}\Gamma_{\underline{p}} , \quad (B.11)$$

where $\nabla_{AdS_p} = \Gamma^{\alpha}\partial_{\alpha} + \frac{1}{4}\Omega^{\beta\gamma}_{\alpha}\Gamma_{\beta\gamma}$ is the Dirac operator on AdS_p . When $\alpha = \hat{p}$, we have $\partial_{\alpha}X^m = \delta^m_{p+1} + \frac{L_A}{L}\delta^m_{p+3}$ so that

$$\partial_{\hat{p}} X^m \nabla_m = \nabla_{p+1} + \frac{L_A}{L} \nabla_{p+3} + \frac{1}{2} \cosh u_0 \Gamma_{\underline{p+1p}} .$$
(B.12)

Since

$$\Gamma^{\hat{p}} = \frac{1}{L_A \cosh u_0} \hat{E}^{\hat{p}}_{\underline{\hat{p}}} \Gamma^{\underline{\hat{p}}} = \frac{\sinh u_0 \Gamma_{\underline{p+1}} + \Gamma_{\underline{p+3}}}{L_A \cosh^2 u_0} , \qquad (B.13)$$

we find

$$\Gamma^{\hat{p}}\partial_{\hat{p}}X^{m}\nabla_{m} = \Gamma^{\hat{p}}\partial_{\hat{p}} + \frac{\cosh u_{0}}{2}\Gamma^{\hat{p}}\Gamma_{\underline{p+1p}} = \Gamma^{\hat{p}}\partial_{\hat{p}} + \frac{\tanh u_{0}}{2L_{A}}\Gamma_{\underline{p}} + \frac{1}{2L_{A}\cosh u_{0}}\Gamma_{\underline{p+3}}\underline{p+1}\underline{p} .$$
(B.14)

Combining (B.11) and (B.14), we finally get

$$\Gamma^{\alpha}\partial_{\alpha}X^{m}\nabla_{m} = \bigvee_{AdS_{p}\times S^{1}} + \frac{p+1}{2L_{A}}\tanh u_{0}\Gamma_{\underline{p}} + \frac{1}{2L_{A}\cosh u_{0}}\Gamma_{\underline{p+3}\,\underline{p+1}\,\underline{p}} . \tag{B.15}$$

C Fermion mass matrix for general u_0

When studying fermionic fluctuations in sections 2.2 and 3.2 we considered the limiting case of $u_0 \rightarrow 0$. Here we show that the equivalent fermion mass matrix is obtained for the general value of u_0 , in both D3 and M5 brane cases: the u_0 dependence can be eliminated by a spinor rotation.

D3 brane case

The fermion mass matrix in the D3 brane case is given by (2.32):

$$M = 2 \tanh u_0 \Gamma_{\underline{3}} + \frac{1}{2 \cosh u_0} \Gamma_{\underline{643}} + \frac{1}{4} \Gamma_{D3} \Gamma^{\alpha} \left(\Gamma^{\underline{0}\cdots\underline{4}} + \Gamma^{\underline{5}\cdots\underline{9}} \right) \Gamma_{\alpha} .$$
(C.1)

Let us consider the rotation matrix

$$\mathcal{R} = \exp(\gamma \Gamma_{\underline{46}}) = \cos\gamma + \Gamma_{\underline{46}} \sin\gamma , \qquad \mathcal{R}^{-1} = \exp(-\gamma \Gamma_{\underline{46}}) = \cos\gamma - \Gamma_{\underline{46}} \sin\gamma , \qquad (C.2)$$

where $(\Gamma_{\underline{46}})^2 = -1$ and γ is related to u_0 by

$$\frac{1+\tan\gamma}{1-\tan\gamma} = e^{u_0} , \qquad \tan\gamma = \tanh\frac{u_0}{2} . \qquad (C.3)$$

Then $\mathcal{R}\Gamma_m \mathcal{R}^{-1} = \Gamma_m$ for $m \neq 4, 6$ and

$$\mathcal{R}\Gamma_{\underline{4}}\mathcal{R}^{-1} = \frac{1}{\cosh u_0} \left(\Gamma_{\underline{4}} - \sinh u_0 \Gamma_{\underline{6}} \right) , \qquad \mathcal{R}\Gamma_{\underline{6}}\mathcal{R}^{-1} = \frac{1}{\cosh u_0} \left(\Gamma_{\underline{6}} + \sinh u_0 \Gamma_{\underline{4}} \right) , \quad (C.4)$$

This enables us to write the world-volume components of Γ -matrices as

$$\Gamma_{\hat{3}} = \Gamma_4 + \Gamma_6 = \sinh u_0 \,\Gamma_{\underline{4}} + \Gamma_{\underline{6}} = \cosh u_0 \,\mathcal{R}\Gamma_{\underline{6}}\mathcal{R}^{-1} , \qquad \Gamma^{\underline{3}} = \Gamma_{\underline{3}} = \mathcal{R}\Gamma_{\underline{6}}\mathcal{R}^{-1} , \qquad (C.5)$$

$$\Gamma_{D3} \equiv \Gamma_{\underline{\hat{0}123}} = \Gamma_{\underline{012}} \mathcal{R} \Gamma_{\underline{6}} \mathcal{R}^{-1} = \mathcal{R} \Gamma_{\underline{0126}} \mathcal{R}^{-1} .$$
(C.6)

Then after a detailed computation, one finds

$$M \cosh u_0 = 2 \sinh u_0 \Gamma_{\underline{3}} + \frac{1}{2} \Gamma_{\underline{643}} + \frac{1}{4} \cosh u_0 \Gamma_{D3} \Gamma^\alpha \left(\Gamma^{\underline{0}\cdots\underline{4}} + \Gamma^{\underline{5}\cdots\underline{9}} \right) \Gamma_\alpha$$
$$= \mathcal{R} \left[\frac{1}{2} \Gamma_{\underline{346}} \Gamma^{\underline{01}\cdots\underline{9}} + \sinh u_0 \Gamma_{\underline{3}} \left(1 - \Gamma^{\underline{01}\cdots\underline{9}} \right) \right] \mathcal{R}^{-1} .$$
(C.7)

Rotating the fermions as $\vartheta \to \vartheta' = \mathcal{R}^{-1}\vartheta$ we get for the gauge-fixed fermionic action (cf. (2.29))

$$S_f = \int d^4 \xi \, \sqrt{-g} \, \bar{\vartheta} \Big(\nabla \!\!\!/ + \mathcal{M} \Big) \vartheta \to \int d^4 \xi \, \sqrt{-g} \, \bar{\vartheta}' \Big(\nabla \!\!\!/ + \mathcal{M}' \Big) \vartheta' \,, \tag{C.8}$$

where we ignore the overall constant factor and the rotated mass matrix is

$$M' = \mathcal{R}^{-1}(M \cosh u_0)\mathcal{R} = \frac{1}{2}\Gamma_{\underline{346}}\Gamma^{\underline{01}\dots\underline{9}} + \sinh u_0\Gamma_{\underline{3}}(1 - \Gamma^{\underline{01}\dots\underline{9}}) = \frac{1}{2}\Gamma_{\underline{346}} .$$
(C.9)

In the final equality we have used the chirality constraint $\Gamma^{\underline{01}\cdots\underline{9}}\vartheta' = \vartheta'$ as M' is acting on a MW spinor. Thus the fermionic action is independent of u_0 , up to an overall constant factor that can be absorbed into a rescaling of the fermionic field and in the present context does not change the value of the fermionic determinant.

M5 brane case

Here the fermion action is given by (cf. (3.37),(3.38))

$$S_f = \int d^6 \xi \sqrt{-g} \, g^{\alpha\beta} \, \bar{\vartheta} (1 - \Gamma_*) \Big(\nabla \!\!\!/ + \mathrm{M} \Big) \vartheta \tag{C.10}$$

where ∇ is the Dirac operator on $\mathrm{AdS}_5 \times S^1$ and

$$\mathbf{M} = \left(\frac{3}{2L} \tanh u_0 \Gamma_{\underline{5}} + \frac{1}{4L \cosh u_0} \Gamma_{\underline{865}}\right) + \left(-\frac{5}{4L} \Gamma^{\underline{78910}} - \frac{1}{4L} \Gamma^{\hat{5}} \Gamma^{\underline{78910}}_{6} - \frac{1}{L} \Gamma^{\hat{5}} \Gamma^{\underline{78910}}_{8}\right), \quad (C.11)$$

where the first bracket is the contribution of the normal part of the spin connection (3.35) while the second is the contribution of F_4 terms in D_m in (3.24) (see (3.30),(3.33)). Introducing the u_0 -dependent rotation matrix as in (C.2) so that $\mathcal{R}\Gamma_m\mathcal{R}^{-1} = \Gamma_m$ for $m \neq \underline{6}, \underline{8}$ and

$$\mathcal{R}\Gamma_{\underline{6}}\mathcal{R}^{-1} = \frac{1}{\cosh u_0} \Big(\Gamma_{\underline{6}} - \sinh u_0 \Gamma_{\underline{8}}\Big), \qquad \mathcal{R}\Gamma_{\underline{8}}\mathcal{R}^{-1} = \frac{1}{\cosh u_0} \Big(\Gamma_{\underline{8}} + \sinh u_0 \Gamma_{\underline{6}}\Big), \qquad (C.12)$$

we find after a detailed computation that the mass matrix (C.11) may be written as

$$2L \cosh u_0 M = \mathcal{R} \Big[3 \sinh u_0 \Gamma_{\underline{5}} + \frac{1}{2} \Gamma_{\underline{865}} - \frac{3}{2} \Gamma_{\underline{789} \underline{10}} + 3 \sinh u_0 \Gamma_{\underline{769} \underline{10}} \Big] \mathcal{R}^{-1} \\ = \mathcal{R} \Big[3 \sinh u_0 \Gamma_{\underline{5}} \Big(1 - \Gamma_{\underline{5679} \underline{10}} \Big) + \frac{1}{2} \Gamma_{\underline{865}} \Big(1 + 3\Gamma_{\underline{5679} \underline{10}} \Big) \Big] \mathcal{R}^{-1} . \quad (C.13)$$

Let us also rotate the fermions so that $\mathcal{R}^{-1}\vartheta = \vartheta'$ and fix the κ -symmetry gauge as in (3.39), i.e.

$$(1-\Gamma_*)\vartheta = 0 = (1-\mathcal{R}\Gamma_{\underline{0}\cdots\underline{48}}\mathcal{R}^{-1})\mathcal{R}\vartheta' = \mathcal{R}(1-\Gamma_{\underline{0}\cdots\underline{48}})\vartheta' = 0 \quad \rightarrow \quad (1-\Gamma_{\underline{0}\cdots\underline{48}})\vartheta' = 0 \;. \; (C.14)$$

We also have $\bar{\vartheta}'(1 + \Gamma_{0\dots 48}) = 0$. The fermion action (C.10) becomes

$$\int d^4 \xi \, \sqrt{-\mathsf{g}} \, \bar{\vartheta}' \Big(\nabla \!\!\!/ + \mathrm{M}' \Big) \vartheta' \,, \qquad (\mathrm{C.15})$$

where we have scaled out the overall constant factor (so that \mathbf{g} is the metric of the unit-radius $AdS_5 \times S^1$) and

$$M' = \mathcal{R}^{-1}(2L\cosh u_0 M)\mathcal{R} = 3\sinh u_0 \Gamma_{\underline{5}} \left(1 - \Gamma_{\underline{5679} 10}\right) + \frac{1}{2}\Gamma_{\underline{865}} \left(1 + 3\Gamma_{\underline{5679} 10}\right) . \quad (C.16)$$

Note that here the first term multiplied by $\Gamma_{\underline{5}} - \Gamma_{\underline{679 \ 10}}$ does not contribute to the gauge-fixed action as it commutes with $1 - \Gamma_*$.

From the analysis of supersymmetry preserved by the M5 brane embedding in Appendix D it follows that assuming that the fermionic action contains the projector $1 - \Gamma_*$ as in (3.25) then $s_1 = 1$ in (D.16) and thus preservation of supersymmetry is consistent with the choice of Γ -matrix representation were $s_2 = 1$ in (D.17). Then the gauge-fixed fermion should be subject to (D.24), i.e.

$$\Gamma_{\underline{5679\,10}}\,\vartheta' = -\vartheta' \,\,, \tag{C.17}$$

and thus finally we can replace (C.16) by

$$M' = \frac{1}{2} \Gamma_{\underline{865}} \left(1 + 3\Gamma_{\underline{5679} \ \underline{10}} \right) = -\Gamma_{\underline{865}} , \qquad (C.18)$$

which is equivalent to (3.40).

Supersymmetry of M5 embedding into $AdS_7 \times S^4$ D

Given the definition of covariant derivative D_m in (3.24) and the F_4 background in (3.28), we find that (cf. (3.31), (3.32))

$$D_m = \nabla_m - \frac{1}{4L} \Gamma_m \hat{\Gamma} , \qquad m < 7; \qquad D_m = \nabla_m + \frac{1}{2L} \Gamma_m \hat{\Gamma} , \qquad m \ge 7 ; \qquad (D.1)$$
$$\hat{\Gamma} \equiv \Gamma_{789\,10}. \qquad (D.2)$$

$$\Gamma \equiv \Gamma_{\underline{789\,10}}.\tag{D.2}$$

Note that $[\Gamma_m, \hat{\Gamma}] = 0$ when m < 7, and $\{\Gamma_m, \hat{\Gamma}\} = 0$ when $m \ge 7$. The 11d Killing spinor equation follows from the condition of the vanishing of the local supersymmetry variation of the 11d gravitino (see, e.g., [60])

$$\delta\psi_m = D_m \epsilon = 0 . \tag{D.3}$$

Since $[\Gamma_m \hat{\Gamma}, \Gamma_n \hat{\Gamma}] = 0$ when m < 7 and $n \ge 7$, the AdS₇ and S⁴ parts of ϵ factorize

$$\epsilon = \epsilon_{AdS_7} \epsilon_{S^4} = \mathcal{M}_{AdS_7} \mathcal{M}_{S^4} \epsilon_0 , \qquad [\mathcal{M}_{AdS_7}, \mathcal{M}_{S^4}] = 0 , \qquad (D.4)$$

where ϵ_0 is a constant spinor.

Let us first consider the AdS₇ part and set $L_A = 2L = 1$ for simplicity. Let us first consider the m = u = 5 (cf. (3.27),(3.1)) component of (D.3), i.e.

$$D_u \epsilon = \left(\partial_u - \frac{1}{2}\Gamma_{\underline{u}}\hat{\Gamma}\right)\epsilon = 0 \qquad \to \qquad \epsilon = e^{\frac{1}{2}u\Gamma_{\underline{u}}\hat{\Gamma}}\epsilon' , \qquad (D.5)$$

where ϵ' is independent of u. Next, for $m = \psi = 6$ we get³⁸

$$D_{\psi}\epsilon = \left[\partial_{\psi} - \frac{1}{2}\left(\cosh u\,\Gamma_{\underline{u}\underline{\psi}} + \sinh u\,\Gamma_{\underline{\psi}}\hat{\Gamma}\right)\right]\epsilon = \left[\partial_{\psi} - \frac{1}{2}e^{\frac{1}{2}u\,\Gamma_{\underline{u}}\hat{\Gamma}}\Gamma_{\underline{u}\underline{\psi}}e^{-\frac{1}{2}u\Gamma_{\underline{u}}\hat{\Gamma}}\right]\epsilon = 0 , \quad (D.6)$$

Substituting (D.5) into (D.6), we get that

$$\left(\partial_{\psi} - \frac{1}{2}\Gamma_{\underline{u}\underline{\psi}}\right)\epsilon' = 0 \qquad \rightarrow \qquad \epsilon' = e^{\frac{1}{2}\psi\Gamma_{\underline{u}\underline{\psi}}}\epsilon'' . \tag{D.7}$$

For $m = 0, \dots, 5$, we have

$$D_m = \tilde{\nabla}_m - \frac{1}{2} \Big(\cosh u \,\Gamma_m \hat{\Gamma} - \sinh u \,\Gamma_{m\underline{u}} \Big) = \tilde{\nabla}_m - \frac{1}{2} e^{\frac{1}{2} u \Gamma_{\underline{u}} \hat{\Gamma}} \Gamma_m \hat{\Gamma} e^{-\frac{1}{2} u \Gamma_{\underline{u}} \hat{\Gamma}} , \qquad (D.8)$$

where $\tilde{\nabla}_m$ has spin connection components along AdS₅ only. This means we can write the Killing spinor in AdS_7 parametrized as in (3.1) in terms of the Killing spinor on AdS_5 (independent of u and ψ) as

$$\epsilon_{AdS_7} = e^{\frac{1}{2}u\Gamma_{\underline{u}}\hat{\Gamma}}e^{\frac{1}{2}\psi\Gamma_{\underline{u}\underline{\psi}}}\epsilon_{AdS_5} , \qquad \left(\hat{\nabla}_m - \frac{1}{2}\Gamma_m\hat{\Gamma}\right)\epsilon_{AdS_5} = 0 . \tag{D.9}$$

Similarly, for S^4 components of (D.3) we get $(\theta = 7, \phi = 8)^{39}$

$$D_{\theta}\epsilon = \left(\partial_{\theta} + \frac{1}{2}\Gamma_{\underline{\theta}}\hat{\Gamma}\right)\epsilon = 0 \qquad \to \qquad \epsilon = e^{-\frac{1}{2}\theta\Gamma_{\underline{\theta}}\hat{\Gamma}}\epsilon' , \qquad (D.10)$$

$$D_{\phi}\epsilon = \left[\partial_{\phi} + \frac{1}{2}(-\cos\theta\Gamma_{\underline{\theta}\underline{\phi}} + \sin\theta\Gamma_{\underline{\phi}}\hat{\Gamma})\right]\epsilon = \left(\partial_{\phi} - \frac{1}{2}e^{-\frac{1}{2}\Gamma_{\underline{\theta}}\hat{\Gamma}}\Gamma_{\underline{\theta}\underline{\phi}}e^{\frac{1}{2}\Gamma_{\underline{\theta}}\hat{\Gamma}}\right)\epsilon = 0.$$
(D.11)

Substituting (D.10) into (D.11), we find

$$\left(\partial_{\phi} - \frac{1}{2}\Gamma_{\underline{\theta}\underline{\phi}}\right)\epsilon' = 0 \qquad \rightarrow \qquad \epsilon' = e^{-\frac{1}{2}\phi\Gamma_{\underline{\phi}\underline{\theta}}}\epsilon'' .$$
 (D.12)

For the remaining m = 9,10 components of D_m (corresponding to the $S^2 \subset S^4$ angles $\varphi_1 = 9, \ \varphi_2 = 10$) we get

$$D_m = \hat{\nabla}_m + \frac{1}{2} e^{-\frac{1}{2}\Gamma_{\underline{\theta}}\hat{\Gamma}} \Gamma_m \hat{\Gamma} e^{\frac{1}{2}\Gamma_{\underline{\theta}}\hat{\Gamma}} , \qquad (D.13)$$

where $\hat{\nabla}_m$ contains only the S^2 spin connection. As a result,⁴⁰

$$\epsilon_{_{S^4}} = e^{-\frac{1}{2}\theta\Gamma_{\underline{\theta}}\hat{\Gamma}}e^{-\frac{1}{2}\phi\Gamma_{\underline{\phi}\underline{\theta}}}\epsilon_{_{S^2}} , \qquad \left(\hat{\nabla}_m + \frac{1}{2}\Gamma_m\hat{\Gamma}\right)\epsilon_{_{S^2}} = 0 . \tag{D.14}$$

³⁸We use the following relations which are valid for i, j < 7: $e^{\frac{1}{2}\alpha\Gamma_i\hat{\Gamma}}\Gamma_{ij}e^{-\frac{1}{2}\alpha\Gamma_i\hat{\Gamma}} = \cosh \alpha\Gamma_{ij} + \sinh \alpha\Gamma_j\hat{\Gamma}$ and $e^{\frac{1}{2}\alpha\Gamma_i\hat{\Gamma}}\Gamma_j\hat{\Gamma}e^{-\frac{1}{2}\alpha\Gamma_i\hat{\Gamma}} = \sinh \alpha\Gamma_{ij} + \cosh \alpha\Gamma_j\hat{\Gamma}.$ ³⁹We use that for $i, j \ge 7$: $e^{\frac{1}{2}\alpha\Gamma_i\hat{\Gamma}}\Gamma_{ij}e^{-\frac{1}{2}\alpha\Gamma_i\hat{\Gamma}} = \cos \alpha\Gamma_{ij} + \sin \alpha\Gamma_j\hat{\Gamma}$ and $e^{\frac{1}{2}\alpha\Gamma_i\hat{\Gamma}}\Gamma_j\hat{\Gamma}e^{-\frac{1}{2}\alpha\Gamma_i\hat{\Gamma}} = \cos \alpha\Gamma_j\hat{\Gamma} - \cos \alpha\Gamma_j\hat{\Gamma}$

⁴⁰The explicit solution for ϵ_{g2} is $e^{-\frac{1}{2}\varphi_1\Gamma_{\underline{\varphi_1}}\hat{\Gamma}}e^{-\frac{1}{2}\varphi_2\Gamma_{\underline{\varphi_2}\varphi_1}}$.

 $[\]sin \alpha \Gamma_{ij}$.

Combining (D.9) and (D.14), we find the Killing spinor on $AdS_7 \times S^4$ can be written as

$$\epsilon = \mathcal{M}\epsilon_0 , \qquad \mathcal{M} = \mathcal{M}_{AdS_7}\mathcal{M}_{S^4} = e^{\frac{1}{2}u\Gamma_{\underline{u}}\hat{\Gamma}}e^{\frac{1}{2}\psi\Gamma_{\underline{u}\underline{\psi}}}\mathcal{M}_{AdS_5}e^{-\frac{1}{2}\theta\Gamma_{\underline{\theta}}\hat{\Gamma}}e^{-\frac{1}{2}\phi\Gamma_{\underline{\phi}\underline{\theta}}}\mathcal{M}_{S^2}$$
$$= e^{\frac{1}{2}u\Gamma_{\underline{u}}\hat{\Gamma}}e^{\frac{1}{2}\psi\Gamma_{\underline{u}\underline{\psi}}}e^{-\frac{1}{2}\theta\Gamma_{\underline{\theta}}\hat{\Gamma}}e^{-\frac{1}{2}\phi\Gamma_{\underline{\phi}\underline{\theta}}}\mathcal{M}_{AdS_5\times S^2} , \qquad (D.15)$$

where $\mathcal{M}_{AdS_5 \times S^2}$ depends only on the AdS₅ and S^2 coordinates.

Let us now consider the M5 brane configuration in (3.27) and find the amount of global supersymmetry it preserves (for a general discussion see, e.g., [61, 16, 62]). The supersymmetry condition for the brane embedding is determined by the projector $\frac{1}{2}(1+\Gamma_*)$ orthogonal to the one that enters the κ -symmetry transformation of ϑ in the M5 brane action (3.23),(3.25),(3.26) and thus the κ -symmetry gauge on θ in (3.39).⁴¹

To account for possible orientation choice ambiguity let us introduce the parameter $s_1 = \pm 1$ and assuming the fermionic action (and thus also the gauge fixing condition (3.39)) contains the projector $1 - s_1\Gamma_*$ ($s_1 = 1$ in (3.25)) consider in general the condition $(1 + s_1\Gamma_*)\epsilon = 0$, i.e. (cf. (3.34))

$$\Gamma_* \epsilon = -s_1 \epsilon , \qquad \Gamma_* = \Gamma_{\underline{\hat{0}}\underline{\hat{1}}\underline{\hat{2}}\underline{\hat{3}}\underline{\hat{4}}\underline{\hat{5}}} = \frac{1}{\cosh u_0} \Gamma_{\underline{0}\cdots\underline{4}} \Big(\sinh u_0 \Gamma_{\underline{6}} + \Gamma_{\underline{8}} \Big) , \qquad s_1 = \pm 1 . \quad (D.16)$$

Let us also set

$$\Gamma_{11} \equiv \Gamma_{\underline{0}\cdots\underline{9}\,\underline{10}} = s_2 , \qquad \Gamma_{11}^2 = 1 , \quad s_2 = \pm 1 , \qquad (D.17)$$

where s_2 is introduced to account for a freedom in choice of Γ -matrix representation. Note that under $\Gamma_m \to -\Gamma_m$ we have $\Gamma_* \to \Gamma_*$, $\Gamma_{11} \to -\Gamma_{11}$ and M5 brane action (3.23) stays the same (up to overall sign) provided one also changes $F_4 \to -F_4$ in the covariant derivative D_m in (3.24). Equivalently, the Killing spinor in this case is still given by (D.15) with $\Gamma_m \to -\Gamma_m$ (with $\hat{\Gamma}$ in (D.2) staying invariant).

In (D.16) ϵ is the Killing spinor of the AdS₇ × S⁴ background (D.15), specialised to the brane solution (3.9), i.e. $u = u_0$, $\phi = 2\psi$, $\theta = \frac{\pi}{2}$. Since the brane extends along AdS₅ and is localized at a point in S², we do not need track the dependence of the Killing spinor in (D.15) on those coordinates and may effectively set $\mathcal{M}_{AdS_5 \times S^2} = 1$. Then

$$\epsilon = \bar{\mathcal{M}} \epsilon_0 , \qquad \qquad \bar{\mathcal{M}} = e^{\frac{1}{2}u_0\Gamma_{\underline{u}}\hat{\Gamma}} e^{\frac{1}{2}\psi\Gamma_{\underline{u}\underline{\psi}}} e^{-\frac{\pi}{4}\Gamma_{\underline{\theta}}\hat{\Gamma}} e^{-\psi\Gamma_{\underline{\phi}\underline{\theta}}} . \tag{D.18}$$

Then condition (D.16) may be written as

$$K\epsilon_0 = 0$$
, $K \equiv \overline{\mathcal{M}}^{-1}(1 + s_1\Gamma_*)\overline{\mathcal{M}}$. (D.19)

Using (D.17) we get for $\hat{\Gamma}$ in (D.2)

$$\Gamma \equiv \Gamma_{\underline{789}\,\underline{10}} = s_2 \Gamma_{\underline{0}\cdots\underline{6}} \ . \tag{D.20}$$

⁴¹In general [61], the variation of ϑ in (3.25) under the κ -symmetry and target space supersymmetry is $\delta\vartheta = (1 - \Gamma_*)\kappa + \epsilon$. Upon gauge fixing $(1 - \Gamma_*)\vartheta = 0$, i.e. $\vartheta = (1 + \Gamma_*)\tilde{\vartheta}$. The preservation of the gauge condition implies $(1 - \Gamma_*)\delta\vartheta = (1 + \Gamma_*)\delta\vartheta = 0$ and thus the condition for unbroken global supersymmetry of the brane embedding is $(1 + \Gamma_*)\epsilon = 0$.

One can show that in general

$$K = 1 - s_1 s_2 + s_1 s_2 \left(1 - \tanh u_0 \cos \psi \, \Gamma_{\underline{6910}} + \tanh u_0 \sin \psi \, \Gamma_{\underline{678910}} \right) \left(1 - \Gamma_{\underline{5678}} \right) \,. \quad (D.21)$$

To have a non-trivial constant ϵ_0 solution of (D.19) we are thus to require

$$s_1 s_2 = 1$$
, $(1 - \Gamma_{\underline{5678}}) \epsilon_0 = 0$. (D.22)

Since $\Gamma_{\underline{5678}}^2 = 1$ we thus get a projector implying preservation of half of the original supersymmetry.

Note that performing the rotation $\varepsilon = \mathcal{R}^{-1} \epsilon$ discussed in Appendix C, we get Γ_* in (D.16) transformed to its $u_0 = 0$ value $\Gamma_{0...48}$ and thus the condition (D.16) becomes (cf. (D.17))

$$\Gamma_*\varepsilon = \Gamma_{\underline{0}\cdots\underline{48}}\,\varepsilon = -s_1\varepsilon\,,\qquad \Gamma_{\underline{5679}\,\underline{10}}\,\varepsilon = -\Gamma_{\underline{0}\cdots\underline{48}}\Gamma_{\underline{0}\cdots\underline{9}\,\underline{10}} = s_1s_2\varepsilon = \varepsilon\,. \tag{D.23}$$

At the same time, the κ -symmetry gauge condition (3.39) on the fermionic field ϑ that involves the projector complementary to the one in (D.16) reads

$$\Gamma_*\vartheta' = \Gamma_{\underline{0}\cdots\underline{48}}\vartheta' = s_1\vartheta' , \qquad \Gamma_{\underline{5679}\,\underline{10}}\vartheta' = -s_1s_2\vartheta' = -\vartheta' . \qquad (D.24)$$

This is the condition we used in the main text to arrive at the expression for the mass operator in (3.40).

References

- N. Andrei et al., Boundary and Defect CFT: Open Problems and Applications, J. Phys. A 53 (2020) 453002 [1810.05697].
- [2] K. Jensen, A. O'Bannon, B. Robinson and R. Rodgers, From the Weyl Anomaly to Entropy of Two-Dimensional Boundaries and Defects, Phys. Rev. Lett. 122 (2019) 241602 [1812.08745].
- [3] A. Chalabi, A. O'Bannon, B. Robinson and J. Sisti, Central charges of 2d superconformal defects, JHEP 05 (2020) 095 [2003.02857].
- [4] Y. Wang, Defect a-theorem and a-maximization, JHEP 02 (2022) 061 [2101.12648].
- [5] N. Drukker, S. Giombi, A. A. Tseytlin and X. Zhou, Defect CFT in the 6d (2,0) theory from M2 brane dynamics in AdS₇×S⁴, JHEP 07 (2020) 101 [2004.04562].
- [6] A. Chalabi, C. P. Herzog, A. O'Bannon, B. Robinson and J. Sisti, Weyl anomalies of four dimensional conformal boundaries and defects, JHEP 02 (2022) 166 [2111.14713].
- [7] N. Drukker, O. Shahpo and M. Trépanier, Quantum holographic surface anomalies, J. Phys. A 57 (2024) 085402 [2311.14797].
- [8] P. Capuozzo, J. Estes, B. Robinson and B. Suzzoni, Holographic Weyl Anomalies for 4d Defects in 6d SCFTs, 2310.17447.
- N. R. Constable, J. Erdmenger, Z. Guralnik and I. Kirsch, Intersecting D-3 branes and holography, Phys. Rev. D 68 (2003) 106007 [hep-th/0211222].

- [10] N. Drukker, J. Gomis and S. Matsuura, Probing N=4 SYM With Surface Operators, JHEP 10 (2008) 048 [0805.4199].
- [11] E. D'Hoker, J. Estes, M. Gutperle, D. Krym and P. Sorba, Half-BPS supergravity solutions and superalgebras, JHEP 12 (2008) 047 [0810.1484].
- [12] N. Drukker, J. Gomis and D. Young, Vortex Loop Operators, M2-branes and Holography, JHEP 03 (2009) 004 [0810.4344].
- [13] M. Gutperle and C. F. Uhlemann, Janus on the Brane, JHEP 07 (2020) 243 [2003.12080].
- [14] V. Gupta, Holographic M5 branes in $AdS_7 \times S^4$, JHEP 12 (2021) 032 [2109.08551].
- [15] N. Drukker and O. Shahpo, Vortex loop operators and quantum M2-branes, 2312.17091.
- [16] O. Lunin, 1/2-BPS states in M theory and defects in the dual CFTs, JHEP 10 (2007) 014 [0704.3442].
- [17] S. Giombi and A. A. Tseytlin, Wilson Loops at Large N and the Quantum M2-Brane, Phys. Rev. Lett. 130 (2023) 201601 [2303.15207].
- [18] M. Beccaria, S. Giombi and A. A. Tseytlin, Instanton contributions to the ABJM free energy from quantum M2 branes, JHEP 10 (2023) 029 [2307.14112].
- [19] M. Beccaria, S. Giombi and A. A. Tseytlin, (2,0) theory on S5 ×S¹ and quantum M2 branes, Nucl. Phys. B 998 (2024) 116400 [2309.10786].
- [20] M. Beccaria and A. A. Tseytlin, Large N expansion of superconformal index of k?=?1 ABJM theory and semiclassical M5 brane partition function, Nucl. Phys. B 1001 (2024) 116507 [2312.01917].
- [21] M. Sakaguchi, H. Shin and K. Yoshida, Semiclassical Analysis of M2-brane in AdS_4xS^7/Z_k , JHEP 12 (2010) 012 [1007.3354].
- [22] R. Camporesi and A. Higuchi, Spectral functions and zeta functions in hyperbolic spaces, J.Math.Phys. 35 (1994) 4217.
- [23] R. Camporesi and A. Higuchi, The Plancherel measure for p-forms in real hyperbolic spaces, J. Geom. Phys. 15 (1994) 57.
- [24] S. Giombi, A. Maloney and X. Yin, One-loop Partition Functions of 3D Gravity, JHEP 08 (2008) 007 [0804.1773].
- [25] S. Giombi, I. R. Klebanov, S. S. Pufu, B. R. Safdi and G. Tarnopolsky, AdS Description of Induced Higher-Spin Gauge Theory, JHEP 10 (2013) 016 [1306.5242].
- [26] S. Giombi and I. R. Klebanov, One Loop Tests of Higher Spin AdS/CFT, JHEP 12 (2013) 068 [1308.2337].
- [27] S. Giombi, I. R. Klebanov and B. R. Safdi, *Higher Spin* AdS_{d+1}/CFT_d at One Loop, *Phys. Rev.* D 89 (2014) 084004 [1401.0825].
- [28] M. Beccaria and A. A. Tseytlin, Higher spins in AdS₅ at one loop: vacuum energy, boundary conformal anomalies and AdS/CFT, JHEP 11 (2014) 114 [1410.3273].
- [29] K. Skenderis and M. Taylor, Branes in AdS and pp wave space-times, JHEP 06 (2002) 025 [hep-th/0204054].

- [30] A. Faraggi and L. A. Pando Zayas, The Spectrum of Excitations of Holographic Wilson Loops, JHEP 05 (2011) 018 [1101.5145].
- [31] E. I. Buchbinder and A. A. Tseytlin, 1/N correction in the D3-brane description of a circular Wilson loop at strong coupling, Phys. Rev. D 89 (2014) 126008 [1404.4952].
- [32] N. Drukker and B. Fiol, All-genus calculation of Wilson loops using D-branes, JHEP 02 (2005) 010 [hep-th/0501109].
- [33] L. Martucci, J. Rosseel, D. Van den Bleeken and A. Van Proeyen, Dirac actions for D-branes on backgrounds with fluxes, Class. Quant. Grav. 22 (2005) 2745 [hep-th/0504041].
- [34] R. R. Metsaev and A. A. Tseytlin, Supersymmetric D3-brane action in AdS(5) x S5, Phys. Lett. B 436 (1998) 281 [hep-th/9806095].
- [35] O. Aharony, M. Berkooz, A. Karasik and T. Vaknin, Supersymmetric field theories on $AdS_p \times S^q$, JHEP 04 (2016) 066 [1512.04698].
- [36] I. A. Bandos, K. Lechner, A. Nurmagambetov, P. Pasti, D. P. Sorokin and M. Tonin, Covariant action for the superfive-brane of M theory, Phys. Rev. Lett. 78 (1997) 4332 [hep-th/9701149].
- [37] M. Aganagic, J. Park, C. Popescu and J. H. Schwarz, World volume action of the M theory five-brane, Nucl. Phys. B 496 (1997) 191 [hep-th/9701166].
- [38] P. S. Howe, E. Sezgin and P. C. West, Covariant field equations of the M theory five-brane, Phys. Lett. B 399 (1997) 49 [hep-th/9702008].
- [39] P. Pasti, D. P. Sorokin and M. Tonin, Covariant action for a D = 11 five-brane with the chiral field, Phys. Lett. B 398 (1997) 41 [hep-th/9701037].
- [40] L. Santilli and C. F. Uhlemann, 3d defects in 5d: RG flows and defect F-maximization, JHEP 06 (2023) 136 [2305.01004].
- [41] I. A. Bandos, K. Lechner, A. Nurmagambetov, P. Pasti, D. P. Sorokin and M. Tonin, On the equivalence of different formulations of the M theory five-brane, Phys. Lett. B 408 (1997) 135 [hep-th/9703127].
- [42] P. Claus, Super M-brane actions in AdS(4) x S7 and AdS(7) x S4, Phys. Rev. D 59 (1999) 066003 [hep-th/9809045].
- [43] P. Claus, R. Kallosh and A. Van Proeyen, M five-brane and superconformal (0,2) tensor multiplet in six-dimensions, Nucl. Phys. B 518 (1998) 117 [hep-th/9711161].
- [44] B. de Wit, K. Peeters, J. Plefka and A. Sevrin, The M theory two-brane in AdS(4) x S7 and AdS(7) x S4, Phys. Lett. B 443 (1998) 153 [hep-th/9808052].
- [45] M. Sakaguchi and K. Yoshida, Open M-branes on AdS(4/7) x S7/4 revisited, Nucl. Phys. B 714 (2005) 51 [hep-th/0405109].
- [46] E. Cremmer and B. Julia, The SO(8) Supergravity, Nucl. Phys. B 159 (1979) 141.
- [47] Y. N. Obukhov, The geometric approach to antisymmetric tensor field theory, Phys. Lett. B 109 (1982) 195.
- [48] E. S. Fradkin and A. A. Tseytlin, Quantum Properties of Higher Dimensional and Dimensionally Reduced Supersymmetric Theories, Nucl. Phys. B 227 (1983) 252.

- [49] F. Bastianelli, S. Frolov and A. A. Tseytlin, Conformal anomaly of (2,0) tensor multiplet in six-dimensions and AdS/CFT correspondence, JHEP 0002 (2000) 013 [hep-th/0001041].
- [50] J. R. David and J. Mukherjee, Hyperbolic cylinders and entanglement entropy: gravitons, higher spins, p-forms, JHEP 01 (2021) 202 [2005.08402].
- [51] J. R. David and J. Mukherjee, Partition functions of p-forms from Harish-Chandra characters, JHEP 09 (2021) 094 [2105.03662].
- [52] R. R. Metsaev, Lowest eigenvalues of the energy operator for totally (anti)symmetric massless fields of the n-dimensional anti-de Sitter group, Class. Quant. Grav. 11 (1994) L141.
- [53] R. R. Metsaev, Massive totally symmetric fields in AdS(d), Phys. Lett. B 590 (2004) 95 [hep-th/0312297].
- [54] R. R. Metsaev, Fermionic fields in the d-dimensional anti-de Sitter space-time, Phys. Lett. B 419 (1998) 49 [hep-th/9802097].
- [55] S. Gukov, Comments on N=2 AdS orbifolds, Phys. Lett. B 439 (1998) 23 [hep-th/9806180].
- [56] O. Aharony, M. Berkooz and S.-J. Rey, Rigid holography and six-dimensional $\mathcal{N} = (2,0)$ theories on $AdS_5 \times \mathbb{S}^1$, JHEP **03** (2015) 121 [1501.02904].
- [57] H.-C. Kim and S. Kim, M5-branes from gauge theories on the 5-sphere, JHEP 05 (2013) 144 [1206.6339].
- [58] M. Beccaria and A. Cabo-Bizet, Large N Schur index of $\mathcal{N} = 4$ SYM from semiclassical D3 brane, 2402.12172.
- [59] D. E. Diaz and H. Dorn, Partition functions and double-trace deformations in AdS/CFT, JHEP 05 (2007) 046 [hep-th/0702163].
- [60] H. Lu, C. N. Pope and J. Rahmfeld, A Construction of Killing spinors on Sⁿ, J. Math. Phys. 40 (1999) 4518 [hep-th/9805151].
- [61] E. Bergshoeff, R. Kallosh, T. Ortin and G. Papadopoulos, Kappa symmetry, supersymmetry and intersecting branes, Nucl. Phys. B 502 (1997) 149 [hep-th/9705040].
- [62] J. Simon, Brane Effective Actions, Kappa-Symmetry and Applications, Living Rev. Rel. 15 (2012) 3 [1110.2422].