# Computational complexity of isometric tensor network states

Daniel Malz

Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen, Denmark

Rahul Trivedi

Electrical and Computer Engineering, University of Washington, Seattle, Washington 98195, USA

We determine the computational power of isometric tensor network states (isoTNS), a variational ansatz originally developed to numerically find and compute properties of gapped ground states and topological states in two dimensions. By mapping 2D isoTNS to 1+1D unitary quantum circuits, we find that computing local expectation values in isoTNS is BQP-complete. We then introduce *injective* isoTNS, which are those isoTNS that are the unique ground states of frustration-free Hamiltonians, and which are characterized by an injectivity parameter  $\delta \in (0, 1/D]$ , where D is the bond dimension of the isoTNS. We show that injective isoTNS (small  $\delta$ ) are still BQP-complete, but that there exists an efficient classical algorithm to compute local expectation values in strongly injective isoTNS ( $\eta \geq 0.41$ ). Sampling from isoTNS corresponds to monitored quantum dynamics and we exhibit a family of isoTNS that undergo a phase transition from a hard regime to an easy phase where the monitored circuit can be sampled efficiently. Our results can be used to design provable algorithms to contract isoTNS. Our mapping between ground states of certain frustration-free Hamiltonians to open circuit dynamics in one dimension fewer may be of independent interest.

### I. INTRODUCTION

One of the central goals in condensed matter theory and many-body physics is to describe and analyze ground states of local Hamiltonians. Since the dimension of the Hilbert space of a lattice system grows exponentially with the number of sites, computing properties of even modestly sized systems requires approximations. One particular successful strategy has been to variationally search for ground states in restricted families of states. For example, the celebrated density matrix renormalization group method [1] for one dimensional systems is effectively a variational method over the class of matrix product states (MPS) [2, 3]. MPS are so successful, because they efficiently capture ground states of gapped Hamiltonians [4], and simultaneously are easy to use on classical computers (in terms of computational complexity) [5]. In recent years, it has also become increasingly relevant that MPS can also efficiently be prepared on quantum computers using a linear depth circuit [6], or approximately even in logarithmic depth [7].

In two and higher dimensions, the MPS ansatz has been extended to projected entangled pair states (PEPS) [8–10], which are also highly expressive [11] and have been used successfully to gain insight into longstanding problems in condensed-matter physics [12–19]. However, the usefulness of PEPS for numerical calculations limited by the fact that contracting them in general is computationally hard (#P-complete [20]), and thus even using a quantum computer, they are hard to prepare [20] or contract [21].

While approximate contraction strategies for PEPS have brought considerable success [12–19], another strategy is to define subsets of PEPS that admit efficient algorithms to optimize them. An early example of such

a simplification were sequentially generated states [22]. Leveraging MPS techniques, these states allow for efficient computation of expectation values everywhere, but unlike PEPS, increasing bond dimension seems to improve their expressibility only modestly [22]. More recently, isometric tensor network states (isoTNS) have been introduced [23], which generalize MPS in a less restrictive way. For instance, they provably capture exactly the renormalization-group fixed point of all string-net liquids [24], a constant-depth quantum circuits [24], some thermal states [25] and certain fine-tuned critical states [26] and to some degree also quantum dynamics [27, 28].

IsoTNS may inhabit a sweet spot, being computationally strictly easier than PEPS, yet with some evidence towards their success. To understand better what certain ansätze can and cannot capture, it is important to understand the classical and computational complexity associated to using them. This has motivated work to characterize the (quantum and classical) computational complexity of PEPS [20] and injective PEPS [29]. Here, we therefore explore the computational complexity of isoTNS. Our results hold equally well for plaquette-PEPS [30].

#### **II. SUMMARY OF RESULTS**

We build on the result that isoTNS can be written in terms of a linear-depth sequential quantum circuit and thus can be regarded as being generated from a chain of ancillas that sequentially interact with the physical sites of the lattice [30]. The time evolution of the ancillas after tracing the physical sites maps the 2D isoTNS to a 1+1D circuit of quantum channels, which needs to be

TABLE I. Summary of results of this paper. We determine the classical complexity of computing local expectation values to constant additive error, producing samples according to the probability distribution obtained by measuring the state to constant additive error, and producing samples to multiplicative error. For a specific familiy of injective isoTNS we can show that sampling undergoes some sort of "measurement-induced phase transition" to an easy phase, but we do not know if what happens for general strongly injective isoTNS, hence the question mark.

	Weakly/Non injective isoTNS	Strongly injective isoTNS
EXPECTATION TO $O(1)$ ADDITIVE ERROR SAMPLING TO $O(1)$ ADDITIVE ERROR SAMPLING TO $O(1)$ MULTIPLICATIVE ERROR	Hard unless BQP=BPP Hard unless BQP=BPP Hard unless postBQP = postBPP	$\begin{array}{l} \operatorname{Easy} \\ ? \\ \operatorname{Hard\ unless\ } postBQP = postBPP \end{array}$

simulated to compute a local expectation value on a site corresponding to some later time, which provides a way to efficiently compute computing expectation values in isoTNS using a quantum computer. Our first result is then to construct isoTNS where computing local expectation values is BQP-complete [31], which means that a machine or algorithm that can compute local expectation values in isoTNS is as powerful as a quantum computer.

Mirroring injective PEPS, which are those states that are unique ground states of frustration-free parent Hamiltonians) [32], we introduce injective isoTNS. They are characterized by an injectivity parameter  $\delta \in (0, 1/D]$ which is defined as the size of the smallest singular value of the PEPS tensor. We show that injectivity necessarily adds depolarizing noise at a rate  $\eta = \delta^2 D^2$  to the 1+1D circuit of the ancillas, where D is the bond dimension of the isoTNS. Our second result is then that local expectation values in weakly injective isoTNS is still BQPcomplete, which we show by embedding a fault-tolerant circuit that is robust to small amounts of noise. Conversely, we provide an explicit algorithm to compute local expectation values that runs in  $poly(1/\epsilon)$  time (independent of system size) for strongly injective isoTNS for which  $\eta \ge 0.41$ .

Sampling from states to additive error is at least as hard as computing local observables, so the hardness of local expectation values extends to sampling. We show that sampling from isoTNS maps to monitored quantum circuits, which are known to undergo a complexity transition under strong monitoring [33]. We construct a class of isoTNS that as a function of the injectivity parameter also undergoes such a complexity transition, which provides an intriguing link between 2D states and 1D dynamics. We also consider sampling to multiplicative precision, which we find is hard even in the maximally injective case.

We summarise our results in Table I.

### **III. RELATED PRIOR WORK**

Schuch et al. [20] have used a duality between PEPS and postselected quantum circuits to show that a machine capable of preparing any PEPS is equivalent to a postselected quantum computer. In contrast, by virtue of the isometry condition, isoTNS are dual to quantum circuits, which naturally lead to the result that a machine that can prepare isoTNS is equivalent to a quantum computer.

Anshu et al. [29] use a similar (to the present work) fault-tolerant embedding of quantum circuits in injective PEPS to show that injective PEPS are BQP-hard. They also show that determining expectation values of nonlocal observables to multiplicative precision is #P-hard. In injective isoTNS, we find that local expectation values are BQP-complete and thus hard on classical computers. Note that while it is clear that isoTNS can be prepared on a quantum computer, this is not clear for injective PEPS as considered in Ref. [29]. It may well be the case that their complexity is greater than BQP.

Many results exist on the difficulty of sampling [34], even from shallow-depth circuits [35]. To show that sampling from isoTNS to multiplicative error is hard independent of the injectivity parameter, we establish a link between maximally injective isoTNS and IQP circuits (circuits of commuting gates), which are known to be hard to sample from unless postBQP=postBPP [36].

Our results establishing that computing local expectation values in sufficiently injective isoTNS is easy draw on early results showing that noisy quantum computers undergo a complexity transition [37, 38], which uses a similar percolation result to the present paper. Similarly, after mapping SAMPLING in isoTNS to monitored quantum circuits, we can draw on the large body of literature that establishes conditions under which such monitored circuits become easy to simulate [33, 39, 40].

Another result that establishes conditions under which there is an efficient classical algorithm to compute local expectation values in injective PEPS is obtained in Ref. [41]. There, it was shown that local observables in a normalized injective PEPS on d-dimensions can be computed to a precision  $\epsilon$  by just restricting the PEPS to a  $\log(\epsilon^{-1})$  length cube around the local observable. In 2D, this would also yield a  $poly(\epsilon^{-1})$  algorithm for computing the local observable. The key distinction between this and our result is that the result in Ref. [41] does not rely on isometry, but instead they require that the frustrationfree parent Hamiltonian of the PEPS is gapped and remains uniformly gapped when removing terms, which is an unrelated and potentially stronger condition.

IsoTNS can also be viewed as being prepared from a chain of ancillas interacting sequentially with arrays of qubits [30], similar to how MPS can be prepared from a single emitter [6]. This links efforts to find efficient classical numerical approaches based on tensor networks to efforts to utilize qubit-efficient "holographic" ways of simulating quantum systems [42–47]. Our work directly applies to states generated in such schemes and may thus also be useful to understand the power and limitation of such approaches.

## IV. ISOTNS, QUANTUM CIRCUITS, AND CHANNEL CIRCUITS

Most of our results are based on interpreting isoTNS contraction as physical evolution of the virtual degrees of freedom, which builds on representing them as sequential circuits [30], which we recall in the following.

#### A. Matrix product states (MPS)

Recall that MPS of bond dimension D and physical dimension d can be thought of as being generated sequentially by a D-dimensional ancilla  $\mathcal{H}_{\mathcal{A}} \simeq \mathbb{C}^{D}$  [6] interacting sequentially with a sequence of d-dimensional qudits  $\mathcal{H}_{\mathcal{B}} \simeq \mathbb{C}^{d}$  (each initially in  $|0\rangle$ ). In one step of a protocol, a new qudit is attached to the chain and a unitary is performed on the joint qudit-ancilla system  $\mathcal{H}_{\mathcal{A}} \otimes \mathcal{H}_{\mathcal{B}}$ . The whole operation can expressed as an isometry  $V : \mathcal{H}_{\mathcal{A}} \to \mathcal{H}_{\mathcal{A}} \otimes \mathcal{H}_{\mathcal{B}}$ . If  $\{|i\rangle\}, i \in \{0, \ldots, d-1\}$ and  $\{|\alpha\rangle\}, \alpha \in \{0, \ldots, D-1\}$  are respectively bases for qudit and ancilla, we can write

$$V = \sum_{i=0}^{d-1} \sum_{\alpha,\beta=0}^{D-1} V_{\alpha\beta}^{i} |i\rangle_{\mathcal{P}} |\alpha\rangle_{\mathcal{V}} \langle\beta|.$$
(1)

Here and in the rest of the article, we use Latin superscript letters for physical indices/legs, Greek subscript letters for the virtual (ancilla) indices/legs and a subscript  $\mathcal{P}$  ( $\mathcal{V}$ ) to label states in the physical (virtual) Hilbert space. The isometry condition implies that contracting the outgoing legs with the corresponding legs on the conjugate tensor yields the identity on the remaining legs, *viz*.

$$\sum_{i=0}^{d-1} V^{i\dagger} V^i = \xrightarrow{\rightarrow} = = 1, \qquad (2)$$

where we also introduced graphical notation in which tensors are represented by boxes and their indices/legs by lines. Here and in the following, we draw the physical legs in black and the ancilla/virtual legs in red, and we graphically depict the space the isometry acts on (the input) by legs with ingoing arrows, and the target space (the output) by outgoing arrows. Applying N isometries, and requiring that final state be a product state of ancilla and physical qudits, the reduced state of the qudits is an MPS in canonical form. Conversely, any MPS can be written in this way [6].

For later convenience, we allow the ancilla dimension to vary across the system (as a function of time). In particular, we truncate the ancilla dimension such that the reduced state of the ancilla is always full rank, which means that before the first step and after the last step, the ancilla dimension is 1. This allows us to write the MPS as

$$|\psi_{\rm MPS}\rangle = \sum_{i_1\dots i_N=0}^{d-1} V_{[n]}^{i_n} \cdots V_{[1]}^{i_1} |i_N \dots i_1\rangle$$
$$= \underbrace{\stackrel{i_1 \quad i_2}{\longrightarrow} \dots \quad \stackrel{i_N}{\longrightarrow} \dots \quad (3)$$

Notice that the first isometry (leftmost) is in fact just a state (the input dimension is 1) and the last isometry is a unitary (input and output dimension agree).

During the generation process, the reduced state of the ancilla evolves under the action of the channels obtained by tracing over the chain qudits. In the nth step, the channel reads

$$\Phi_{n}: \rho \mapsto \operatorname{tr}_{\mathcal{P}}(V_{[n]}\rho V_{[n]}^{\dagger}) = \sum_{i=0}^{d-1} V_{[n]}^{i}\rho V_{[n]}^{i\dagger}.$$
 (4)

This allows us to calculate the expectation value of an observable  $\mathcal{O}$  on the *n*th site as

$$\langle \psi_{\text{MPS}} | \mathcal{O}_n | \psi_{\text{MPS}} \rangle = \text{tr}(V_{[n]} \rho_{n-1} V_{[n]}^{\dagger} \mathcal{O}).$$
 (5)

Equivalently, we could say that the transfer matrix of the MPS is a CPTP map.

#### B. Isometric tensor network states

The isometry property can be generalized to define isometric tensor network states two dimensions [23]. We can think of an  $N_x \times N_y$ -dimensional isoTNS as being produced by  $N_x + N_y$  ancillas. The ancillas fly through each other in the following way:  $N_y$  are arranged as a vertical stack and fly left to right, and  $N_x$  are arranged as a horizontal stack and fly bottom to top. Whenever their worldlines cross, we apply an isometry that acts on the two ancillas and generates a physical qudit. For  $(N_x, N_y) = (4, 2)$ , the process and state can be illustrated as

$$|\psi\rangle = \prod_{m,n}^{\leftarrow} V_{[mn]} = \xrightarrow{\uparrow} \qquad \uparrow \qquad (6)$$

Here we have used the symbol  $\overline{\prod}$  to denote that the isometries need to be applied in the order of the arrows, i.e., isometries earlier in the worldline are applied first

and appear further right in the product. Note that in our convention the arrows point in the direction of time and opposite to those in Ref. [23].

As for MPS, expectation values can be computed by following the time evolution of the ancillas until the required row and column as in Eq. (4). The reduced state of the ancillas  $\rho_{\mathcal{V}}$  after tracing the physical degree of freedom evolves under the channels [*cf.* Eq. (4)]

$$\Phi_{mn}: \rho_{\mathcal{V}} \mapsto \operatorname{tr}_{\mathcal{P}}(V_{[mn]}\rho_{\mathcal{V}}V_{[mn]}^{\dagger}) = \sum_{i=0}^{d-1} V_{[mn]}^{i}\rho_{\mathcal{V}}V_{[mn]}^{i\dagger}$$

$$= \bigvee_{\mathcal{V}} \bigvee_$$

where in the illustration we have suppressed indices and the rest of the legs of  $\rho$  for clarity. Here,  $V_{[mn]}$  acts on the reduced state of two incoming ancillas (the  $n^{\text{th}}$ ancilla from the vertical stack and the  $m^{\text{th}}$  ancilla from the horizontal stack). To evaluate the expectation value of an observable  $\mathcal{O}$  at site (m, n) we need to first apply all "earlier" channels

$$\langle \psi | \mathcal{O}_{mn} | \psi \rangle = \operatorname{tr}_{\mathcal{P}} \left[ V_{mn} \left( \prod_{\substack{k,l \le m,n, \\ (k,l) \ne (m,n)}}^{\leftarrow} \Phi_{kl}(|\vec{0}\rangle\langle\vec{0}|) \right) V_{mn}^{\dagger} \mathcal{O} \right].$$
(8)

Note that this is merely a reinterpretation of tensor network contraction, but where the fact that we have isometries allows us to interpret the contraction as CPTP channels acting on the virtual degrees of freedom.

The definition of isoTNS can straightforwardly be extended to arbitrary directed acyclic graphs and with mixed bond dimensions, but for the rest of this work we focus on the physically relevant two-dimensional states.

#### C. Projected-entangled pair states and injectivity

Projected-entangled pair states (PEPS) is an alternative way to think about tensor network states.

**Definition 1 (PEPS)** Given a graph G = (W, E) with vertices W and edges E, we associate to each edge  $e \in E$ an (unnormalized) entangled pair state  $|\phi_e\rangle = \sum_i^{D_e} |ii\rangle$ , where  $D_e$  is called bond dimension. This maximally entangled pair is distributed to the two vertices connected by the edge. As a result, we obtain the state  $|\Phi\rangle =$  $\bigotimes_e |\phi_e\rangle$ , which is defined on a lattice given by the vertices, where each vertex v has the local Hilbert space dimension  $\dim \mathcal{H}_v = \prod_{e \in E_v} D_e$ . A PEPS is obtained by applying a linear map to each vertex that maps ("projects") the halves of the entangled pairs at that vertex to a physical space  $P : \mathbb{C}_{\prod_e D_e} \to \mathbb{C}_{d_v}$ , where  $d_v$  is the physical dimension at vertex v. An isoTNS written as in Eq. (6) is a PEPS with projectors

$$\begin{array}{c} \checkmark P \rightleftharpoons \\ & \downarrow \end{array} = P^{k}_{\alpha\beta\gamma\delta} = V^{k}_{\alpha\beta,\gamma\delta} = \begin{array}{c} \checkmark V \checkmark \\ & \checkmark V \end{array}, \qquad (9)$$

Thus, P and V are same tensor, but V is thought of as a map from  $D^2$  to  $d * D^2$ , whereas P is a map from  $D^4$  to d.

The most studied condition that can be imposed on PEPS is *injectivity*. Injective PEPS are the unique ground states of local frustration-free Hamiltonians [32]. They have the property that every tensor has an inverse acting on the physical index. Thus, we require that  $d \ge D^4$  and that every PEPS projector  $P_{[ij]}$  has an inverse  $P_{[ij]}^{-1}$  such that

$$\sum_{i=1}^{d} \left( P_{[ij]}^{-1} \right)_{\alpha\beta\gamma\eta}^{i} \left( P_{[ij]} \right)_{\alpha'\beta'\gamma'\eta'}^{i} = \delta_{\alpha\alpha'} \delta_{\beta\beta'} \delta_{\gamma\gamma'} \delta_{\eta\eta'}.$$
 (10)

Graphically, this condition reads

$$\xrightarrow{\longrightarrow} P^{-1} \rightarrow \longleftarrow P \xrightarrow{\longleftarrow} = \underbrace{\qquad} \dots \qquad (11)$$

To avoid pathological situations in which tensors are arbitrarily close to non-injective ones, we will demand that all singular values of P are greater or equal to some  $\delta > 0$ , or equivalently that  $||P^{-1}||_{\infty} \leq 1/\delta$ . A similar condition has been used before [29, 41, 48] and the corresponding states (without the isometry condition) have been dubbed  $\delta$ -injective PEPS [29]. (We note that defining the PEPS tensor as acting on *normalized* entangled pair states corresponds to diving each element of P by a factor D and a corresponding change in  $\delta$ .)

#### D. Injective isoTNS and depolarizing noise

Injectivity imposed on isoTNS has the interesting consequence that the ancillas are guaranteed to experience some amount of local depolarizing noise. In the following computations we take the bond dimensions to be Dacross all bonds for simplicity, but a generalization to inhomogeneous bond dimension is straightforward.

Let us first characterize what values  $\delta$  can take. Since the Frobenius norm is equal to the element-wise 2-norm, we have  $\|V\|_F = \|P\|_F = \sqrt{\sum_i^{D^4} \sigma_i^2}$ , where  $\{\sigma_i\}$  are the singular values of P. Since V is an isometry mapping a  $D^2$ -dimensional space to  $dD^2$ , we have  $\|V\|_F = D$ . Now consider the spectral norm  $\|P\|_{\infty}$ . For a fixed Frobenius norm, it is minimized when all its  $D^4$  singular values are equal, which in our case gives  $1/D \leq \|P\|_{\infty}$ . In turn it is maximized if one singular value is large and all the others are as small as possible, i.e., equal to  $\delta$ . Together with the lower bound, this yields

$$1/D \le ||P||_{\infty} \le \sqrt{D^2 - \delta^2 (D^4 - 1)}.$$
 (12)

Both inequalities can only be satisfied if

$$\delta \le 1/D. \tag{13}$$

If this bound is saturated, all singular values of P are 1/D, which means it is a (rescaled) isometry as well. (The scaling factor D stems from our definition of PEPS in terms of unnormalized entangled pairs, see Definition 1.) States obtained by acting with an isometry on maximally entangled pairs have been considered before and are called *isometric PEPS* [49]. These PEPS have strictly finite nearest neighbour correlations and can be prepared using a constant-depth circuit.

Away from this limit, for smaller  $\delta$ , we find that injectivity causes the ancillas to experience depolarizing noise (a similar result was very recently obtained for a special family of PEPS, namely ground states that encode circuits through teleportation [29]). To establish this, we insert a resolution of the identity inside the trace in Eq. (7)

$$\mathbb{1}_{d} = (\mathbb{1}_{d} - \eta M) + \eta M, \qquad M = P^{-1\dagger} P^{-1} / D^{2}.$$
(14)

Recall that P maps virtual to physical space, so M maps physical space to physical space. This allows us to decompose the channel  $\Phi_{mn}$  [Eq. (7)] as a sum of a depolarizing channel  $\mathcal{E}_{depol}$  of strength  $\eta$  and another channel  $\mathcal{E}_1$ 

$$\Phi_{mn}(\rho) = \operatorname{tr}_{\mathcal{P}}[(\mathbb{1} - \eta M)V_{[mn]}\rho V_{[mn]}^{\dagger}] + \eta \mathbb{1}/D^{2}$$
  
=  $(1 - \eta)\mathcal{E}_{1}(\rho) + \eta \mathcal{E}_{\operatorname{depol}}(\rho).$  (15)


To see that  $\mathcal{E}_1$  is a proper channel note that it can be written in terms of Kraus operators by diagonalizing M. Taking U to be the unitary that diagonalizes  $\mathbb{1} - \eta M = U^{\dagger} \Lambda U$  for some diagonal matrix  $\Lambda = \text{diag}(\lambda_1, \lambda_2, \ldots)$ , we have

$$\mathcal{E}_1(\rho) = (1-\eta)^{-1} \sum_i K^i \rho K^{i\dagger}, \qquad (16)$$

where we define the Kraus operators  $K^i = \sqrt{\lambda_i} \langle i | UV$  and require  $\mathbb{1} - \eta M \ge 0$  such that  $\lambda_i \ge 0$ . To see that  $\mathcal{E}_1$  is trace preserving, we can check

$$\sum_{i} K^{i\dagger} K^{i} = V^{\dagger} (\mathbb{1} - \eta M) V = \mathbb{1} - \frac{\eta}{D^{2}} V^{\dagger} (P^{-1})^{\dagger} P^{-1} V.$$
(17)

We can evaluate the second term in Eq. (17) as



Thus,  $\sum_{i} K^{i\dagger} K^{i} = (1-\eta)\mathbb{1}$ . By assumption,  $\|D^2 M\|_{\infty} \leq 1/\delta^2$ , so the maximum  $\eta$  we can choose is  $D^2\delta^2$ .

## V. COMPLEXITY OF COMPUTING LOCAL EXPECTATION VALUES

### A. Worst-case hardness of isoTNS

To show that computing local expectation values in isoTNS with constant bond dimension is BQP-complete, first note that following Eq. (6), isoTNS can efficiently be prepared on quantum computers, which can be used to sample from the state. For the converse statement, that any problem in BQP can be mapped to isoTNS, we show that one can embed a brickwork circuit into isoTNS. For simplicity, we consider D = 2, d = 4 and three types of tensor: (i) a gate tensor G that implements a unitary gate on the ancilla degrees of freedom (with arbitrary  $U_{\mathcal{V}}$ acting only on the ancillas),


(ii) a *swap tensor* that swaps the ancillas into the physical space

$$S = \delta^i_{\alpha} \delta^j_{\beta} |ij\rangle_{\mathcal{P} \mathcal{V}} \langle \alpha \beta | = \bigcup =$$
 (20)

(note that the red outgoing legs in the last illustration have dimension 1 and are drawn just for visual consistency), (iii) and an *identity tensor* that simply swaps the ancillas and initializes the physical qubits in  $|0\rangle$ 

$$I = |00\rangle_{\mathcal{P}} SWAP_{\mathcal{V}} = \underset{|0\rangle^{\otimes 2}}{\stackrel{\bigstar}{\longleftarrow}} = \underset{(21)}{\stackrel{\bigstar}{\longrightarrow}} I$$

Given any one-dimensional brickwork circuit with nearest-neighbour gates, we can now define a corresponding isoTNS using the following procedure: (i) replace all unitaries by the corresponding gate tensor G, (ii) at the end of the circuit place a layer of swap tensors S, (iii) fill up the remaining sites with the identity tensor to obtain a square lattice geometry. For example for a circuit of four qubits and depth five, we have (suppressing physical legs for clarity)



The size of the isoTNS is linear in both number of qubits and depth and the reduced state of the physical qudits corresponding to sites with the swap tensor (marked in yellow) is the output of the circuit. Computing local expectation values on these sites is therefore BQP-complete.

# B. Strongly injective isoTNS can be contracted efficiently

Equation (22) defines a highly artificial state and one might wonder what happens to states that might more realistically appear in the study of ground states. In the absence of topological order, PEPS are typically injective (or normal), which is equivalent to the state being the unique ground state of a nearest-neighbour frustrationfree Hamiltonian [32]. Intuitively, injectivity means that every observable on the virtual space can be accessed using a suitably defined observable on the physical space and as we have shown, this leads to depolarizing noise. Depolarizing noise reduces the computational power of the ancillas, eventually leading to a transition to a classically simulable regime, of a type first predicted more than 20 years ago [37, 38]. We will use this to show two things: (i) strongly injective isoTNS can be contracted efficiently, and (ii) weakly injective isoTNS are still BQP-complete.

To find a provably efficient classical algorithm, we adapt a method due to Aharonov [38] to map contracting the tensor network to percolation. To evaluate a given (constant-support) observable, we randomly sample from possible assignments of channels to sites. Specifically, using Eq. (15), the time evolution of the ancilla degrees of freedom can be cast as a stochastic process, where we apply a brickwall circuit of nearest-neighbour channels, and each channel is randomly chosen as either the depolarizing channel  $\mathcal{E}_{depol}$  (empty site) with probability  $\eta$  or as  $(\Phi - \eta \mathcal{E}_{depol})/(1 - \eta)$  (occupied site) with probability  $1 - \eta$ , where  $\Phi$  is the channel given in Eq. (7). To compute the observable in a given sample, we only need to contract the connected cluster of filled sites (see Fig. 1). If  $\eta \geq 0.41$ , then the equivalent site percolation is subcritical (i.e. the filled sites do not percolate) and the cluster connected to the site of the local observable will be independent of the system size [50].

To make the algorithm efficient, we only accept the sample if the cluster size lies below some cutoff  $s_{\rm th}$ . As a result, there is systematic error in the computation of the expectation value. A standard result in percolation theory is the exponential suppression of clusters of large sizes [50]: on a lattice of n sites and when the effective 2D percolation problem is subcritical, the size of the cluster connected to the local observable, S, as a random variable satisfies

$$\operatorname{Prob}(\mathcal{S} \ge s) \le O(s^2 \exp(-s/\xi_0)), \tag{23}$$

for some constant  $\xi_0$  [51]. The expectation value of a local observable  $\mathcal{O}$  is computed using only accepted samples, which gives an error that we can bound as

$$\begin{aligned} |\mathbb{E}(\langle \mathcal{O} \rangle | \mathcal{S} \leq s_{\rm th}) - \langle \mathcal{O} \rangle | \\ &= \frac{|\langle \mathcal{O} \rangle - \mathbb{E}(\langle \mathcal{O} \rangle | \mathcal{S} \geq s_{\rm th}) \operatorname{Prob}(\mathcal{S} \geq s_{\rm th})|}{1 - \operatorname{Prob}(\mathcal{S} \geq s_{\rm th})} \\ &\leq 2 \|\mathcal{O}\| O(s_{\rm th}^2 \exp(-s_{\rm th}/\xi_0)), \end{aligned}$$
(24)

FIG. 1. Illustration of one sample in the algorithm to compute local expectation values. Each site is randomly assigned the depolarizing channel (empty crossing) or the channel  $\mathcal{E}_1$  (blue tensor). In the easy phase, sites with tensors do not percolate and clusters are typically small. As a result, to compute an observable on the red site, we only need to contract the cluster connected to the sites (blue) and in fact only the part of the cluster in the past light cone (gray background).

have that where we used the fact  $|\langle \mathcal{O} \rangle|, |\mathbb{E}(\langle \mathcal{O} \rangle|\mathcal{S} \ge s_{\mathrm{th}})| \le ||\mathcal{O}||.$ Thus, to compute  $\langle \mathcal{O} \rangle$  to a precision  $\epsilon$ , the a cluster size threshold  $s_{\rm th} = \Theta(\log(\epsilon^{-1}))$  is sufficient. This yields a run-time of  $poly(\epsilon^{-1})$  for contracting each individual accepted sample, and  $\Theta(\epsilon^{-2})$  samples are needed to reduce the variance of the estimated average to  $O(\epsilon)$ . Putting these two estimates together, we conclude that, in the subcritical regime, local observables in injective isoTNS can be computed in time independent in the system size and inverse-polynomial in the desired precision.

#### C. Weakly injective isoTNS are still hard

Next, we consider the question of hardness of computing local observables in injective isoTNS for any  $\delta$ . Building on the idea in Ref. [29] and from the fact that a 1D quantum circuit with nearest-neighbour gates can be fault tolerant [52, 53], we show that for  $\delta$  below some threshold, computing local observables is BQP-hard. To encode a 1D fault tolerant quantum circuit into the ancilla dynamics, we need to construct injective and isometric tensors that approximately implement (a) a given two-qubit unitary U and (b) a restart operation on one of the ancilla qubits.

As in Section VA, we take bond dimension D = 2, but now we need physical dimension  $d = D^4 = 16$  to obtain injective tensors. First, we perturb the gate tensor Eq. (19) to injectivity. To this end, consider a slightly depolarized unitary

$$\mathcal{E}_U(\rho) = (1-p)U\rho U^{\dagger} + p\frac{\mathbb{1}\otimes\mathbb{1}}{4}\operatorname{tr}(\rho).$$
(25)

As a channel on  $\mathbb{C}^D \otimes \mathbb{C}^D$ ,  $\mathcal{E}_U$  has full Kraus rank and thus is described by  $(D^2)^2 = D^4$  linearly independent Kraus operators  $A^1, A^2 \dots A^{D^4}$ . We can then choose the isometry from the input ancillas to the output ancillas and physical qudit to be the Stinespring dilation of  $A^i$ 

$$V := \sum_{i=1}^{D^4} A^i \otimes |i\rangle, \tag{26}$$

where the states  $|1\rangle, |2\rangle \dots |D^4\rangle$  are a basis for the Hilbert space  $\mathbb{C}^d$  corresponding to the physical qudit. Furthermore, as calculated in Appendix A, if we define the PEPS tensor P in terms of V [see Eq. (9)], we have  $||P^{-1}||_{\infty} = 1/\delta = 2/\sqrt{p}$ .

Next, we perturb the restart gate to injectivity. In the ideal case, it should reset (one of the) ancilla qubit, i.e., implement the map  $\rho \to \text{tr}_1(\rho) \otimes |0\rangle\langle 0|$ . To make it injective, we again implement a slightly depolarized version of this channel

$$\mathcal{E}_{\rm res}(\rho) = (1-p)|0\rangle\!\langle 0| \otimes {\rm Tr}_1(\rho) + p \frac{\mathbb{1}^{\otimes 2}}{4} {\rm Tr}(\rho).$$
(27)

As with the channel applying the unitary gate, this channel also has full Kraus rank and be described by  $D^4$  linearly independent Kraus operators, which can then be mapped to an isometry via the Stinespring dilation, which gives again a PEPS tensor with  $||P^{-1}||_{\infty} = 1/\delta = 2/\sqrt{p}$ .

Now, we use the circuit to isoTNS embedding described in section VA and depicted schematically in Eq. (22), but instead of embedding a 1D quantum circuit into it directly, we embed its fault tolerant version (which would include both two-qubit unitaries and restart operations) together with the decoding circuit. If  $\delta$  is small enough, then the noise strength is below the fault tolerance threshold. We point out that while it may appear as if the identity tensors in the circuit Eq. (22) introduce an error, but in fact they are only relevant at the boundary of the circuit, when they propagate an ancilla that is part of the circuit. Thus, they contribute depolarizing noise at the same rate as all the other gates and are corrected automatically. We conclude that this embedding establishes that computing local observables would again become BQP-complete when p is small enough, and thus computing local observables on (weakly) injective isoTNS is as hard as on general isoTNS in the worst case.

## VI. COMPLEXITY OF SAMPLING FROM ISOTNS

SAMPLING is a computational task where we are required to return K samples that are distributed according to the probability distribution  $p(\vec{x}) = |\langle \vec{x} | \psi \rangle|^2$ , where  $|\vec{x}\rangle$  are computational basis states.

Sampling from output states of random circuits with or without depolarizing noise has been studied intensely [34] as it was perceived to be a potential avenue to demonstrate quantum computational advantage [54]. In Section VIA we show that sampling to multiplicative error is hard even from maximally injective isoTNS, which are equal to isometric PEPS. This result closely mirrors earlier results that establish hardness when sampling from low-depth circuits or cluster states [35, 55].

Another sampling problem that has been studied intensely in recent years is if instead of depolarizing noise, random unitary circuits are interspersed with measurements. In this setting, high measurement rates drive the system to an area-law state, whereas for low measurement rates, the system tends to a volume-law state, in a phenomenon that has been dubbed "measurementinduced phase transition" (MIPT) [33, 39, 40]. In the thermodynamic limit, there is a sharp transition between these regimes and this setting has elicited tremendous interest [56].

At face value, sampling from states and sampling from the measurements added to a dynamical circuits are not directly related, as the former probes properties of the state, and the latter of the dynamics. However, in line with the central theme of this paper, we can use the fact that isoTNS are closely related to physical dynamics to establish a map between sampling from the isoTNS and monitored quantum dynamics of the ancillas. In particular, this allows us to prove the existence of a MIPT when sampling from certain isoTNS as a function of the injectivity parameter. To show this, we construct a oneparameter family of isoTNS that interpolates from the provably hard regime to a provably easy regime in Section VIB.

## A. Hardness to sample to multiplicative precision

We first consider the problem of sampling an injective isoTNS to multiplicative precision. More specifically, we ask whether there can exist an efficient classical algorithm that can sample from a probability distribution  $p_{\rm cl}(\vec{x})$  such that

$$\frac{1}{1+\epsilon}p(\vec{x}) \le p_{\rm cl}(\vec{x}) \le (1+\epsilon)p(\vec{x}),\tag{28}$$

where  $\epsilon \geq 0$  controls the accuracy.

In this section, we establish that even the most injective isoTNS (i.e., isoTNS with  $\delta = 1/D$ , which saturate the bound in Eq. (13)), are hard to sample from within a multiplicative error. Our conclusion is based on the simple insight that with post-selection, even highly injective isoTNS can encode post-selected quantum computations. It has been previously established in Ref. [36] that families of quantum states that, with postselection, can encode postselected quantum computation are unlikely to be classically samplable to multiplicative precision since that would effectively imply that postBPP = postBQP(or that the polynomial hierarchy collapses to the third level), which is considered unlikely. This establishes a contrast between computation of local observables and sampling for isometric tensor networks, similar to what is known for constant depth quantum circuits.

8

We restrict ourselves to bond-dimension D = 2 and physical dimension  $d = D^4 = 16$ . We first note that a two-qubit unitary U on the ancillary qubits can be applied, on post-selection of the physical qudit, by choosing the isoTNS tensor  $P^k$  via

$$P_U^k = \frac{1}{4} U \sigma_k \text{ for } k \in \{0, 1, 2 \dots 15\},$$
(29)

where  $\sigma_0, \sigma_1 \dots \sigma_{15}$  are 2-qubit Pauli operators and  $\sigma_0 = 1$ . Clearly, on postselecting the physical qudit in  $|0\rangle$ , we effectively apply the unitary U on the ancillas. It can also be verified by explicit computation that  $||P^{-1}||_{\infty} = 2$ , thus saturating the bound in Eq. (13). Furthermore, at the end of the computation, we also need to swap an ancilla qubits with the physical qudits. This can be implemented in an isoTNS by using  $P^k$  as

$$P^{k} = \frac{1}{2} |k_0 k_1 \rangle \langle k_2 k_3 |, \qquad (30)$$

where  $k := \{k_0, k_1, k_2, k_3\} \in \{0, 1\}^4$ . This tensor also saturates the bound in Eq. (13) and thus is maximally injective. Furthermore, post-selecting the physical qudit on  $|0\rangle \cong |0, 0, 0, 0\rangle$  effectively post-selects the ancillas on  $|0, 0\rangle$  and post-selecting the physical qudit on a state of the form  $|k_0, k_2, k_0, k_3\rangle$  (or  $|k_0, k_1, k_2, k_1\rangle$ ) performs a Z-basis measurement on one of the ancillas. Using the two maximally injective tensors shown above, it is clear that any post-selected quantum computation can be encoded into a maximally injective isoTNS with postselection, and hence this isoTNS remains hard to sample from classically to multiplicative precision.

# B. Measurement induced phase transition (MIPT) in sampling

In this section we take the isoTNS into which we have embedded a quantum circuit Eq. (22) and perturb it such that it becomes injective, but we choose the perturbation in such a way that it lets us prove that sampling becomes easy once the state is injective enough. To obtain an injective version of Eq. (22), we first need to embed each physical qubit in a larger Hilbert space  $d = D^4 = 16$ . We decompose the physical Hilbert space at each site into four qubits. Note that all tensors in Eq. (22) can directly be adapted to this setting for example by taking  $V \to V \otimes |0\rangle \otimes |0\rangle$ .

Now consider the following isometry that maps  $\mathbb{C}^2 \to (\mathbb{C}^2)^{\otimes 3}$  (i.e., one ancilla qubit to the ancila qubit and two physical qubits)

$$W = \sum_{i,j} |ij\rangle K_{ij} = \boxed{W},$$
  

$$K_{00} = \sqrt{1/2 - \delta^2} \mathbb{1} + i\delta |1\rangle \langle 1| \qquad K_{01} = i\delta |1\rangle \langle 0|,$$
  

$$K_{11} = \sqrt{1/2 - \delta^2} \mathbb{1} + i\delta |0\rangle \langle 0| \qquad K_{10} = i\delta |0\rangle \langle 1|,$$
  
(31)

Measuring the physical qubits in the computational basis after applying W on a single-qubit state yields unequal outcomes (01 or 10) with probability  $p_{01} + p_{10} = \delta^2$ , independent of the input state. Observing 01 or 10 projects the qubit into  $|1\rangle$  or  $|0\rangle$ , respectively.

We now construct the one-parameter family of tensors that lead to a complexity transition in sampling, which is directly connected to what has been dubbed "measurement-induced phase transition" in monitored quantum circuits [33]. Specifically, we take the same circuit as previously, Eq. (22), but with the perturbed gate tensor

$$\tilde{V} = (W \otimes W)U = \underbrace{\begin{matrix} U \\ W \end{matrix}}_{U} \underbrace{\begin{matrix} U \\ U \end{matrix}}_{U}.$$
(32)

The other tensors we perturb to injectivity as before. Due to our construction of W, now all PEPS tensors P have minimum singular value  $\delta$ .

To simulate sampling, we evolve the ancillas as before in the direction of time, but instead of tracing over the physical sites, we sample from the sites as we go. In the general case, computing this time evolution is difficult (since otherwise we could solve problems in BQP). However, when measuring the physical qubits, in each layer and for each ancilla we have a  $\delta^2$  chance of getting the outcomes 01 or 10 and thus resetting and disentangling the corresponding ancilla qubit. This breaks the bond to the following tensor and for  $\delta^2 > 0.5$  we are above the bond percolation threshold and large clusters are exponentially suppressed.

To rigorously assess the computational effort required to get a certain error, we again introduce a cutoff size  $s_{\rm th}$ and discard samples in which the largest cluster exceeds this size. The probability to find a cluster larger than  $s > s_0$  (for some constant  $s_0$ ) anywhere in the system, and thus the fraction of samples we have to discard, is bounded by (this time with an additional factor of N)

$$\operatorname{Prob}(\mathcal{S} \ge s) \le Ns^2 \exp(-s/\xi_0), \tag{33}$$

where  $\xi_0 > 0$  is again some constant. To ensure that the fraction of rejected samples falls below  $\epsilon$ , we can take  $s_{\rm th} = \xi_0 \log(N/\epsilon) + \xi_0 \log(\log^3(N/\epsilon))$ . Clusters of this size can be contracted in  $\operatorname{poly}(N/\epsilon)$  computational effort, which makes this algorithm efficient.

#### VII. OUTLOOK AND DISCUSSIONS

We have studied the computational complexity of isoTNS, a variational class of tensor network states with the additional constraint that its tensors are isometries. Using a duality between the properties of these states in two dimensions and dynamics of systems in one dimension, we have shown that both computing the local expectation value and sampling from isoTNS have hard and easy phases. Aside from computational complexity, there are also other properties of the states that can be derived from properties of the dynamics. Very recently, isoTNS with power-law correlations have been constructed [26], where the emergence of the power law is connected to the reduced 1D dynamics of the ancillas being critical. Similarly, one can find isoTNS with volume-law states along one column.

It will interesting to explore whether it is possible to use properties of the states and relate them back to properties of the dynamics. For example, approximate Markov states have the property that if two regions Aand C are screened by an intermediate third region B, the mutual information of regions A and C decays with the width of B (typically exponentially) [57]. If an isoTNS has exponential decay of mutual information in this sense, the corresponding ancilla dynamics will be rapidly mixing. Note that the easy phase for injective isoTNS that we have identified corresponds to rapidly mixing ancilla evolution and thus the states that lie in this phase are automatically approximate Markov states.

IsoTNS can be defined on any graph, and one natural direction to explore in the future would be to study their complexity as a function of properties of the underlying graph. Another potential direction is to use the map between isoTNS and open system dynamics to define a continuous version of isoTNS.

Finally, it remains to test our results, conditions, and the algorithms we provide on isoTNS obtained from numerically optimizing for a ground state. We note here that the algorithms we provide to compute local expectation values or to sample can be improved in a number of ways. One promising way to improve the algorithm to compute local expectation values is to improve the decomposition of the channels. Instead of decomposing into depolarizing noise and the rest, one can numerically find the entanglement breaking channel with the largest rate into which the original channel can be decomposed, which would allow one to extend the easy phase.

#### ACKNOWLEDGMENTS

We acknowledge support from the Novo Nordisk Fonden under grant number NNF22OC0071934 and NNF20OC0059939. DM would like to thank Freek Witteveen for insightful discussions, the International Quantum Tensor Network (IQTN) for organizing meetings that have inspired this work, and the Flatiron Institute for hosting one of them, where the idea for this paper was born.

# Appendix A: Calculation of $\delta$ for fault tolerant construction

Perturbing gate tensor to injectivity: Consider the following channel on a qudit  $\mathbb{C}^k$ 

$$\mathcal{E}_U(\rho) = (1-p)U\rho U^{\dagger} + p\operatorname{tr} \rho \frac{\mathbb{1}}{k}.$$
 (A1)

For results concerning embedding a two-qubit unitary into an injective isoTNS, we simply set k = 4 and interpret U as the two-qubit unitary.  $\mathcal{E}_U$  has  $k^2$  Kraus operators  $A^{\alpha,\beta}$  for  $\alpha,\beta \in \{1,2...k\}$ ,

$$A^{\alpha,\beta} = \begin{cases} \gamma_0 U + \sqrt{\frac{p}{k}} |\alpha\rangle\!\langle\alpha|U, & \text{if } \alpha = \beta, \\ \sqrt{\frac{p}{k}} |\alpha\rangle\!\langle\beta|U, & \text{if } \alpha \neq \beta, \end{cases}$$
(A2)

where

$$\gamma_0 = -\sqrt{\frac{p}{k}} + \sqrt{1 - p + \frac{p}{k}}.$$
 (A3)

Note that the operators  $A^{\alpha,\beta}$  are linearly independent, and consequently the PEPS tensor A formed by the Kraus operators is injective. We can now compute the condition number of this tensor. From the bond-space to physical-space, the PEPS tensor A effectively implements the map

$$A|\alpha,\alpha\rangle := (\mathbb{1}\otimes U^{\dagger}) \left[\gamma_0 \sum_{\alpha'=1}^k |\alpha',\alpha'\rangle + \sqrt{\frac{p}{k}} |\alpha,\alpha\rangle\right], \quad (A4)$$

$$A|\alpha,\beta\rangle := \sqrt{\frac{p}{k}} (\mathbb{1} \otimes U^{\dagger})|\alpha,\beta\rangle.$$
(A5)

Note that  $\tilde{A} = (\mathbb{1} \otimes U)A$  is block-diagonal within the subspaces  $S_{\text{diag}} = \text{span}\{|\alpha, \alpha\rangle : \alpha \in \{1, 2...k\}\}$  and  $S_{\text{diag}}^{\perp} = \text{span}\{|\alpha, \beta\rangle : \alpha, \beta \in \{1, 2...k\}\}$ . Furthermore,

$$\tilde{A}\big|_{\mathcal{S}_{\text{diag}}^{\perp}} = \sqrt{\frac{p}{k}}\mathbb{1} \text{ and } \tilde{A}\big|_{\mathcal{S}_{\text{diag}}} = \sqrt{\frac{p}{k}}\mathbb{1} + k\gamma_0 |\Phi\rangle\langle\Phi|,$$
 (A6)

where  $|\Phi\rangle = \sum_{\alpha=1}^{k} |\alpha, \alpha\rangle / \sqrt{k}$ . Thus we obtain

$$\delta = \sigma_{\min}(A) = \lambda_{\min}(\tilde{A}) = \sqrt{\frac{p}{k}}.$$
 (A7)

Perturbing restart tensor to injectivity: Consider now the two qudit channel on  $\mathbb{C}^k \otimes \mathbb{C}^k$ 

$$\mathcal{E}_{\rm res}(\rho) = (1-p)|0\rangle\!\langle 0| \otimes {\rm tr}_1(\rho) + p\,{\rm tr}(\rho)\frac{\mathbb{1}^{\otimes 2}}{k^2}.$$
 (A8)

The  $k^4$  Kraus operators for this channel are given by  $A^{\alpha_1,\alpha_2;\beta}$ , where  $\alpha_1,\alpha_2 \in \{0,1...k-1\}, \beta \in \{0,1,2...k^2-1\}$ 

$$A^{\alpha_1,\alpha_2;\beta} = \begin{cases} \sqrt{1-p+\frac{p}{k^2}}|0\rangle\!\langle\alpha_2|\otimes\sigma_\beta & \text{for } \alpha_1=0,\\ \sqrt{\frac{p}{k^2}}|\alpha_1\rangle\!\langle\alpha_2|\otimes\sigma_\beta & \text{otherwise,} \end{cases}$$
(A9)

where  $\sigma_0 = 1$ ,  $\sigma_1, \sigma_2 \dots \sigma_{k^2-1}$  are traceless matrices and  $\operatorname{tr}(\sigma_i^{\dagger}\sigma_j) = \delta_{i,j}$  i.e.  $\sigma_0, \sigma_1 \dots \sigma_{k^2-1}$  form an orthonormal basis for matrices in  $\mathbb{C}^{k \times k}$ . Note also that  $A^{\alpha_1, \alpha_2;\beta}$  are orthogonal i.e.  $\operatorname{tr}(A^{\alpha_1, \alpha_2;\beta\dagger}A^{\alpha'_1, \alpha'_2;\beta'}) = 0$  unless  $\alpha_1 = \alpha'_1, \alpha_2 = \alpha'_2, \beta = \beta'$ . Therefore, viewing the Kraus operators as a map from the bond indices to the physical

- [1] S. R. White, Density matrix formulation for quantum renormalization groups, Phys. Rev. Lett. **69**, 2863 (1992).
- [2] S. Östlund and S. Rommer, Thermodynamic limit of density matrix renormalization, Physical Review Letters 75, 3537 (1995).
- [3] U. Schollwöck, The density-matrix renormalization group in the age of matrix product states, Ann. Phys. (N. Y). **326**, 96 (2011).
- [4] F. Verstraete and J. I. Cirac, Matrix product states represent ground states faithfully, Phys. Rev. B 73, 094423 (2006).
- [5] D. Perez-Garcia, F. Verstraete, M. M. Wolf, and J. I. Cirac, Matrix Product State Representations, Qu. Inf. Comp. 7, 401 (2007), arXiv:0608197 [quant-ph].
- [6] C. Schön, E. Solano, F. Verstraete, J. I. Cirac, and M. M. Wolf, Sequential Generation of Entangled Multiqubit States, Phys. Rev. Lett. 95, 110503 (2005).
- [7] D. Malz and J. I. Cirac, Few-body analog quantum simulation with rydberg-dressed atoms in optical lattices, PRX Quantum 4, 020301 (2023).
- [8] Y. Hieida, K. Okunishi, and Y. Akutsu, Numerical renormalization approach to two-dimensional quantum antiferromagnets with valence-bond-solid type ground state, New Journal of Physics 1, 7 (1999).
- [9] N. Maeshima, Y. Hieida, Y. Akutsu, T. Nishino, and K. Okunishi, Vertical density matrix algorithm: A higher-dimensional numerical renormalization scheme based on the tensor product state ansatz, Physical Review E 64, 016705 (2001).
- [10] F. Verstraete and J. I. Cirac, Renormalization algorithms for quantum-many body systems in two and higher dimensions, (2004), cond-mat/0407066.
- [11] F. Verstraete, M. M. Wolf, D. Perez-Garcia, and J. I. Cirac, Criticality, the Area Law, and the Computational Power of Projected Entangled Pair States, Phys. Rev. Lett. 96, 220601 (2006).
- [12] J. Jordan, R. Orús, G. Vidal, F. Verstraete, and J. I. Cirac, Classical simulation of infinite-size quantum lattice systems in two spatial dimensions, Physical Review Letters 101, 250602 (2008).
- [13] P. Corboz, T. Rice, and M. Troyer, Competing states in the t-j model: Uniform d-wave state versus stripe state, Physical Review Letters 113, 046402 (2014).
- [14] P. Corboz and F. Mila, Crystals of bound states in the magnetization plateaus of the shastry-sutherland model, Physical Review Letters 112, 147203 (2014).
- [15] H. Liao, Z. Xie, J. Chen, Z. Liu, H. Xie, R. Huang, B. Normand, and T. Xiang, Gapless spin-liquid ground state in the s = 1/2 kagome antiferromagnet, Physical Review Letters 118, 137202 (2017).
- [16] B.-X. Zheng, C.-M. Chung, P. Corboz, G. Ehlers, M.-P. Qin, R. M. Noack, H. Shi, S. R. White,

indices, we obtain

$$\sigma_{\min}(A) = \sqrt{\frac{p}{k^2}}.$$
 (A10)

S. Zhang, and G. K.-L. Chan, Stripe order in the underdoped region of the two-dimensional Hubbard model, Science **358**, 1155 (2017).

- [17] J.-Y. Chen, L. Vanderstraeten, S. Capponi, and D. Poilblanc, Non-abelian chiral spin liquid in a quantum antiferromagnet revealed by an ipeps study, Physical Review B 98, 184409 (2018).
- [18] J. L. Jiménez, S. P. G. Crone, E. Fogh, M. E. Zayed, R. Lortz, E. Pomjakushina, K. Conder, A. M. Läuchli, L. Weber, S. Wessel, A. Honecker, B. Normand, C. Rüegg, P. Corboz, H. M. Rønnow, and F. Mila, A quantum magnetic analogue to the critical point of water, Nature **592**, 370 (2021).
- [19] W.-Y. Liu, J. Hasik, S.-S. Gong, D. Poilblanc, W.-Q. Chen, and Z.-C. Gu, Emergence of gapless quantum spin liquid from deconfined quantum critical point, Physical Review X 12, 031039 (2022).
- [20] N. Schuch, M. M. Wolf, F. Verstraete, and J. I. Cirac, Computational complexity of projected entangled pair states, Phys. Rev. Lett. 98, 140506 (2007).
- [21] I. Arad and Z. Landau, Quantum computation and the evaluation of tensor networks, SIAM Journal on Computing 39, 3089 (2010).
- [22] M. C. Bañuls, D. Pérez-García, M. M. Wolf, F. Verstraete, and J. I. Cirac, Sequentially generated states for the study of two-dimensional systems, Phys. Rev. A 77, 052306 (2008).
- [23] M. P. Zaletel and F. Pollmann, Isometric Tensor Network States in Two Dimensions, Phys. Rev. Lett. 124, 037201 (2020).
- [24] T. Soejima, K. Siva, N. Bultinck, S. Chatterjee, F. Pollmann, and M. P. Zaletel, Isometric tensor network representation of string-net liquids, Phys. Rev. B 101, 085117 (2020).
- [25] W. Kadow, F. Pollmann, and M. Knap, Isometric tensor network representations of two-dimensional thermal states, Phys. Rev. B 107, 205106 (2023).
- [26] Y.-J. Liu, K. Shtengel, and F. Pollmann, Topological quantum phase transitions in 2d isometric tensor networks, (2023), arXiv:2312.05079 [quant-ph].
- [27] S.-H. Lin, M. P. Zaletel, and F. Pollmann, Efficient simulation of dynamics in two-dimensional quantum spin systems with isometric tensor networks, Phys. Rev. B 106, 245102 (2022).
- [28] Y. Kim, A. Eddins, S. Anand, K. X. Wei, E. van den Berg, S. Rosenblatt, H. Nayfeh, Y. Wu, M. Zaletel, K. Temme, and A. Kandala, Evidence for the utility of quantum computing before fault tolerance, Nature 618, 500 (2023).
- [29] A. Anshu, N. P. Breuckmann, and Q. T. Nguyen, Circuitto-hamiltonian from tensor networks and fault tolerance, (2023), arXiv:2309.16475 [quant-ph].
- [30] Z.-Y. Wei, D. Malz, and J. I. Cirac, Sequential generation of projected entangled-pair states,

Phys. Rev. Lett. 128, 010607 (2022).

- [31] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information*, 10th ed. (Cambridge University Press, 2011).
- [32] D. Perez-Garcia, F. Verstraete, J. Cirac, and M. Wolf, PEPS as unique ground states of local Hamiltonians, Quantum Inf. Comput. 8, 650 (2008).
- [33] B. Skinner, J. Ruhman, and A. Nahum, Measurement-Induced Phase Transitions in the Dynamics of Entanglement, Phys. Rev. X 9, 031009 (2019).
- [34] D. Hangleiter and J. Eisert, Computational advantage of quantum random sampling, Reviews of Modern Physics 95, 035001 (2023).
- [35] B. M. Terhal and D. P. DiVincenzo, Adaptive quantum computation, constant depth quantum circuits and arthur-merlin games, Quant. Inf. Comp. Vol. 4 (No. 2), pages 134–145 (2004), arXiv:quant-ph/0205133 [quant-ph].
- [36] M. J. Bremner, R. Jozsa, and D. J. Shepherd, Classical simulation of commuting quantum computations implies collapse of the polynomial hierarchy, Proc. R. Soc. A 467, 459 (2010).
- [37] D. Aharonov and M. Ben-Or, Polynomial simulations of decohered quantum computers (1996) arXiv:quant-ph/9611029v1.
- [38] D. Aharonov, Quantum to classical phase transition in noisy quantum computers, Phys. Rev. A 62, 062311 (2000).
- [39] A. Chan, R. M. Nandkishore, M. Pretko, and G. Smith, Unitary-projective entanglement dynamics, Physical Review B 99, 224307 (2019).
- [40] Y. Li, X. Chen, and M. P. Fisher, Measurement-driven entanglement transition in hybrid quantum circuits, Phys. Rev. B 100, 134306 (2019).
- [41] M. Schwarz, O. Buerschaper, and J. Eisert, Approximating local observables on projected entangled pair states, Physical Review A 95, 060102 (2017).
- [42] T. J. Osborne, J. Eisert, and F. Verstraete, Holographic Quantum States, Phys. Rev. Lett. 105, 260401 (2010).
- [43] I. H. Kim, Holographic quantum simulation, (2017), arXiv:1702.02093 [quant-ph].
- [44] J. Kim, D. Yang, S. hoon Oh, and K. An, Coherent singleatom superradiance, Science 359, 662 (2018).

- [45] J. Borregaard, M. Christandl, and D. Stilck França, Noise-robust exploration of many-body quantum states on near-term quantum devices, npj Quantum Information 7, 10.1038/s41534-021-00363-9 (2021).
- [46] D. Niu, R. Haghshenas, Y. Zhang, M. Foss-Feig, G. K.-L. Chan, and A. C. Potter, Holographic simulation of correlated electrons on a trapped-ion quantum processor, PRX Quantum 3, 030317 (2022).
- [47] S. Anand, J. Hauschild, Y. Zhang, A. C. Potter, and M. P. Zaletel, Holographic quantum simulation of entanglement renormalization circuits, PRX Quantum 4, 030334 (2023).
- [48] M. Schwarz, K. Temme, and F. Verstraete, Preparing Projected Entangled Pair States on a Quantum Computer, Phys. Rev. Lett. 108, 110502 (2012).
- [49] N. Schuch, I. Cirac, and D. Pérez-García, PEPS as ground states: Degeneracy and topology, Ann. Phys. (N. Y). **325**, 2153 (2010).
- [50] D. Stauffer and A. Aharony, *Introduction To Percolation Theory* (Taylor & Francis, 1992).
- [51] M. Z. Bazant, Largest cluster in subcritical percolation, Physical Review E 62, 1660 (2000).
- [52] D. Aharonov and M. Ben-Or, Fault-tolerant quantum computation with constant error rate, (1999), arXiv:quant-ph/9906129 [quant-ph].
- [53] D. Gottesman, Fault-tolerant quantum computation with local gates, Journal of Modern Optics 47, 333 (2000).
- [54] A. Bouland, B. Fefferman, C. Nirkhe, and U. Vazirani, On the complexity and verification of quantum random circuit sampling, Nature Physics 15, 159 (2018).
- Wang, [55] X. Gao. S.-T. and L.-M. Duan. Quantum for simulating supremacy а translation-invariant ising spin model. Physical Review Letters **118**, 040502 (2017).
- [56] M. P. Fisher, V. Khemani, A. Nahum, and S. Vijay, Random quantum circuits, Annual Review of Condensed Matter Physics 14, 335 (2023).
- [57] P. Hayden, R. Jozsa, D. Petz, and A. Winter, Structure of states which satisfy strong subadditivity of quantum entropy with equality, Communications in Mathematical Physics 246, 359 (2004).