

**AFFINE LAUMON SPACE AND CONTRAGREDIENT DUAL
VERMA MODULE OF $U_q(\widehat{\mathfrak{gl}}_n)$**

CHE SHEN

ABSTRACT. We study the action of the quantum group $U_q(\widehat{\mathfrak{gl}}_n)$ on the equivariant K-theory of affine Laumon spaces. We show that, at any highest weight away from the critical level, this can be identified with the contragredient dual Verma module of $U_q(\widehat{\mathfrak{gl}}_n)$, improving earlier results of Braverman-Finkelberg and Neguț. The proof uses a variant of stable envelopes first introduced by Maulik-Okounkov in the study of Nakajima quiver varieties.

CONTENTS

1. Introduction	2
1.1. Laumon space	2
1.2. Affine Laumon space	2
1.3. Strategy of proof	3
2. Laumon spaces and quantum affine algebra	3
2.1. Affine Laumon space	3
2.2. Quantum group $U_q(\widehat{\mathfrak{gl}}_n)$	6
2.3. Specializing to finite Laumon spaces and $U_q(\mathfrak{sl}_n)$	10
3. $K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{fin}})$ and dual Verma module of $U_q(\mathfrak{sl}_n)$	11
4. $K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}})$ and dual Verma module of $U_q(\widehat{\mathfrak{gl}}_n)$	14
4.1. Stable envelopes	15
4.2. $U_q(\widehat{\mathfrak{gl}}_n)$ action almost preserves $K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}})$	19
Appendix A. Existence of stable envelope	22
References	26

1. INTRODUCTION

1.1. **Laumon space.** Laumon spaces parametrize flags of locally-free sheaves on \mathbb{P}^1 :

$$(1) \quad \mathcal{F}_1 \subset \dots \subset \mathcal{F}_{n-1} \subset \mathcal{O}_{\mathbb{P}^1}^{\oplus n}$$

whose fibres near $\infty \in \mathbb{P}^1$ match a fixed full flag of subsheaves of \mathbb{C}^n . Laumon spaces are disconnected, with components indexed by vectors $\mathbf{d} = (d_1, \dots, d_{n-1}) \in \mathbb{Z}_{\geq 0}^{n-1}$ that keep track of the degree of the sheaves in (1). We denote these connected components by $\mathcal{M}_{\mathbf{d}}^{\text{fin}}$.¹ The torus $\mathbb{T} = (\mathbb{C}^*)^{n+1}$ acts on $\mathcal{M}_{\mathbf{d}}^{\text{fin}}$ by scaling the fibres of $\mathcal{O}_{\mathbb{P}^1}^{\oplus n}$ as well as scaling the base \mathbb{P}^1 .

The relation between Laumon spaces and representation theory of \mathfrak{sl}_n , as well as their affine analogs, was extensively studied in the literature, see e.g. [5, 6, 12, 18]. (See also [3, 9] in the context of supersymmetric gauge theories.) In particular, it was proved in [2] that the localized equivariant K-theory of Laumon space

$$K^{\text{loc, fin}} := \bigoplus_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^{n-1}} K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{fin}}) \otimes_{K_{\mathbb{T}}(\text{pt})} \text{Frac } K_{\mathbb{T}}(\text{pt})$$

has a natural action of the quantum group $U_q(\mathfrak{sl}_n)$ and can be identified with the universal Verma module of $U_q(\mathfrak{sl}_n)$ over $\text{Frac } K_{\mathbb{T}}(\text{pt})$, whose highest weight corresponds to equivariant parameters (u_1, \dots, u_n) . It is natural to ask what module we get when specializing the parameters to certain highest weight

$$(2) \quad u_i = q^{a_i}, \quad a_i \in \mathbb{C}, \quad i = 1, \dots, n$$

The answer is less obvious than it seems to be because certain coefficients might become 0 or ∞ making it no longer a Verma module. In fact, as a corollary of theorem 3.1, we prove that the (non-localized) K-theory

$$K^{\text{int, fin}} := \bigoplus_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^{n-1}} K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{fin}})$$

becomes the *contragredient dual* of the Verma module of $U_q(\mathfrak{sl}_n)$ under the specialization (2). A similar result was obtained in [5] when considering (non-quantized) $U(\mathfrak{gl}_n)$ acting on the cohomology of Laumon space.

1.2. **Affine Laumon space.** There is an affine analogue of the above result. Consider the affine Laumon space $\mathcal{M}_{\mathbf{d}}$, see section 2.1 for the definition. [14, 18] constructed a geometric action of $U_q(\widehat{\mathfrak{gl}}_n)$ on the localized equivariant K-theory

$$K^{\text{loc}} := \bigoplus_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^n} K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}) \otimes_{K_{\mathbb{T}}(\text{pt})} \text{Frac } K_{\mathbb{T}}(\text{pt})$$

and [14] further proved that this makes K^{loc} the universal Verma module of $U_q(\widehat{\mathfrak{gl}}_n)$ over $\text{Frac } K_{\mathbb{T}}(\text{pt})$. Similar to the finite case, we prove in corollary 4.2 that

$$K^{\text{int}} := \bigoplus_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^n} K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}})$$

¹The upper script “fin” stands for finite, which reflects its relation to the finite dimensional Lie algebra, whereas affine Laumon space relates to the affine Lie algebra.

becomes the contragredient dual Verma module of $U_q(\widehat{\mathfrak{gl}}_n)$ when specializing the equivariant parameters to certain highest weight. As we will explain in section 4.2, the highest weight must be away from the critical level due to a mild denominator in the $U_q(\widehat{\mathfrak{gl}}_n)$ action.

1.3. Strategy of proof. When $d_n = 0$, the affine Laumon space $\mathcal{M}_{\mathbf{d}}$ becomes the (finite) Laumon space $\mathcal{M}_{(d_1, \dots, d_{n-1})}^{\text{fin}}$, so the results discussed in section 1.1 could be a corollary of section 1.2. However, in this paper, we first present a proof for the finite Laumon space independent of the proof in the general case. The proof makes crucial use of a certain upper triangular property of PBW basis, see section 3. Some key ideas are reused in the proof for the affine case.

For affine Laumon space, the upper triangular property of PBW basis no longer holds and new methods are required. It turns out that a variant of stable envelope, first defined in [11] and further developed in [16, 17], provides a way to “restore” the upper triangular property and plays a key role in the proof. In section 4, we explain in detail how to use the properties of stable envelope and the so-called “rigidity argument” to compute their pairing with the PBW basis. For completeness, we include the proof of the existence of stable envelope in the appendix.

Acknowledgement. I am grateful to my advisor Andrei Okounkov for many helpful discussions and suggestions. I would also like to thank Shaoyun Bai, Sam DeHority, Yixuan Li, Andrei Neguț, Leonid Rybnikov, Spencer Tamagni, Tianqing Zhu for many useful discussions and comments.


2. LAUMON SPACES AND QUANTUM AFFINE ALGEBRA

This section collects some basic properties of affine Laumon space as well as the action of quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_n)$ on the localized K-theory of it. We will closely follow the notations and figures in [14].

2.1. Affine Laumon space.

2.1.1. Definition as a quiver variety. Given an integer $n \geq 2$ and a degree $\mathbf{d} \in \mathbb{Z}_{\geq 0}^n$, Affine Laumon spaces $\mathcal{M}_{\mathbf{d}}$ can be described in two equivalent ways, either as the moduli space of framed parabolic sheaves on $\mathbb{P}^1 \times \mathbb{P}^1$, or as a quiver variety. c.f. [14] section 3. We will use the latter description in this paper.

Consider the following quiver:



Fix spaces V_1, \dots, V_n of dimension d_1, \dots, d_n , consider the vector space of linear maps

$$(3) \quad M_{\mathbf{d}} = \bigoplus_{i=1}^n (\text{Hom}(V_i, V_i) \oplus \text{Hom}(V_{i-1}, V_i) \oplus \text{Hom}(W_i, V_i) \oplus \text{Hom}(V_{i-1}, W_i))$$

(By convention, $V_{n+1} = V_1, V_0 = V_n$.) Elements of the vector space $M_{\mathbf{d}}$ will be quadruples $(X_i, Y_i, A_i, B_i)_{1 \leq i \leq n}$. Consider the quadratic map

$$M_{\mathbf{d}} \xrightarrow{\nu} \bigoplus_{i=1}^{n-1} \text{Hom}(V_{i-1}, V_i)$$

$$\nu(X_i, Y_i, A_i, B_i)_{1 \leq i \leq n} = \bigoplus_{i=2}^{n-1} (X_i Y_i - Y_i X_{i-1} + A_i B_i)$$

A point in $M_{\mathbf{d}}$ is stable if V_i are generated by X and Y acting on the images of A maps. And

$$\mathcal{M}_{\mathbf{d}} = \nu^{-1}(0)^{\text{stable}} / GL_{\mathbf{d}}$$

There is a torus action by

$$(4) \quad \mathbb{T} = (\mathbb{C}^*)^{n+2} = T_W \times \mathbb{C}_q^* \times \mathbb{C}_p^*$$

Let (u_1, \dots, u_n, q, p) denote² the characters of \mathbb{T} , then \mathbb{T} scales the quadruple (X_i, Y_i, A_i, B_i) with weight

$$(q^{-2}, p^{-2\delta_i^1}, u_i^{-2}, u_i^2 q^{-2} p^{-2\delta_i^1})$$

(i.e. p only acts on Y_1 and B_1 .) Note that we are using a 2^{n+2} -fold cover of the usual torus action to avoid square roots in formulas.

Define

$$K^{\text{int}} := \bigoplus_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^n} K_{\mathbf{d}}^{\text{int}} := \bigoplus_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^n} K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}})$$

which is a module over

$$(5) \quad \mathbb{Z}_{\mathbf{u}} := \mathbb{Z}[u_1^{\pm}, \dots, u_n^{\pm}, q^{\pm}, p^{\pm}] = K_{\mathbb{T}}(\text{pt})$$

We also define localized equivariant K-theory

$$K^{\text{loc}} := K^{\text{int}} \otimes_{\mathbb{Z}_{\mathbf{u}}} \mathbb{F}_{\mathbf{u}}$$

² p was denoted \bar{q} in [14]

where

$$(6) \quad \mathbb{F}_{\mathbf{u}} := \text{Frac } \mathbb{Z}_{\mathbf{u}} = \mathbb{Q}(u_1, \dots, u_n, q, p)$$

2.1.2. *Fixed points.* The fixed points are parametrized by n -tuple of 2d partitions $\boldsymbol{\lambda} = (\lambda^1, \dots, \lambda^n)$ where each λ^i itself is a 2d partition. We will simply call $\boldsymbol{\lambda}$ a partition for brevity. (It should be clear from the context whether we are talking about a single 2d partition or an n -tuple of 2d partitions.) For a box \square in the (x, y) position of the partition λ^k , we say that \square has color $y + k \bmod n$. (Note that x, y starts from 0.) Write $|\boldsymbol{\lambda}| = \mathbf{d}$ where $\mathbf{d} = (d_1, \dots, d_n) \in \mathbb{Z}_{\geq 0}^n$ if $\boldsymbol{\lambda}$ has d_i boxes of color i , $i = 1, 2, \dots, n$. Fixed points of $\mathcal{M}_{\mathbf{d}}$ are parametrized by partitions $\boldsymbol{\lambda}$ with $|\boldsymbol{\lambda}| = \mathbf{d}$. We will abuse notation to use $\boldsymbol{\lambda}$ to denote either a fixed point or a partition.

We will use the normalized fixed point basis

$$|\boldsymbol{\lambda}\rangle := \frac{[\mathcal{O}_{\boldsymbol{\lambda}}]}{[\Lambda \bullet (T\mathcal{M}_{\mathbf{d}}|_{\boldsymbol{\lambda}})]}$$

where $[\mathcal{O}_{\boldsymbol{\lambda}}]$ is the structure sheaf supported at the point $\boldsymbol{\lambda}$. With this notation, the equivariant localization formula becomes

$$c = \sum_{|\boldsymbol{\lambda}| = \mathbf{d}} c|_{\boldsymbol{\lambda}} \cdot |\boldsymbol{\lambda}\rangle$$

where $c \in K_{\top}(\mathcal{M}_{\mathbf{d}})$.

2.1.3. *Local coordinates.* For any fixed point $\boldsymbol{\lambda}$, there is an open subset $U_{\boldsymbol{\lambda}} \ni \boldsymbol{\lambda}$ defined in the following way: for any box $\square = (i, j)$ in the k -th partition λ^k of $\boldsymbol{\lambda}$, define the vector

$$v_{\square} := Y_k^j X_k^i A_k w_k \in V_{\text{col } \square}$$

Let $U_{\boldsymbol{\lambda}} \subset \mathcal{M}_{\mathbf{d}}$ be the open subset where $\{v_{\square}\}_{\square \in \boldsymbol{\lambda}}$ form a basis of V_1, \dots, V_n . Writing matrices (X, Y, A, B) in the basis $\{v_{\square}\}_{\square \in \boldsymbol{\lambda}}$, we see that $U_{\boldsymbol{\lambda}}$ is a closed subset inside an affine space, cut out by the moment map condition $\nu = 0$. The fixed point $\boldsymbol{\lambda}$ is the origin of this affine space. We will refer to this as the standard coordinate on $U_{\boldsymbol{\lambda}}$.

2.1.4. *Tautological bundle.* By construction, affine Laumon space comes with tautological bundles $\mathcal{V}_i, i = 1, \dots, n$ defined by globalizing V_i in equation (3).

The K-theory class of the tangent bundle can be described in terms of \mathcal{V}_i 's, see formula (3.23) of [14]:

$$(7) \quad [T\mathcal{M}_{\mathbf{d}}] = \sum_{i=1}^n \left[\left(1 - \frac{1}{q^2}\right) \left(\frac{\mathcal{V}_i}{\mathcal{V}_{i-1}} - \frac{\mathcal{V}_i}{\mathcal{V}_i}\right) + \frac{\mathcal{V}_i}{u_i^2} + \frac{u_{i+1}^2}{\mathcal{V}_i q^2} \right]$$

where by convention $\mathcal{V}_{n+i} = \mathcal{V}_i p^{-2}, u_{n+i}^2 = u_i^2 p^{-2}$ for $i \in \mathbb{Z}$.

Given a partition $\boldsymbol{\lambda} = (\lambda^1, \dots, \lambda^n)$, for a box \square in the (x, y) position of λ^k , define the weight of the box to be

$$\chi_{\square} := u_k^2 q^{2x} p^{2\lfloor \frac{y-k}{n} \rfloor}$$

With this notation, we have

$$\mathcal{V}_i|\lambda = \sum_{\substack{\square \in \lambda \\ \text{col } \square = i}} \chi_{\square}$$

As an example, consider the partition

$$(8) \quad \begin{array}{c} \boxed{u_1^2 p^2} \\ \boxed{u_1^2} \\ \boxed{u_1^2} \quad \boxed{u_1^2 q^2} \\ \boxed{u_1^2} \quad \boxed{u_1^2 q^2} \end{array} \quad \begin{array}{c} \boxed{u_2^2} \\ \boxed{u_2^2} \quad \boxed{u_2^2 q^2} \end{array} \quad \begin{array}{c} \boxed{u_3^2 p^2} \\ \boxed{u_3^2} \\ \boxed{u_3^2} \\ \boxed{u_3^2} \quad \boxed{u_3^2 q^2} \end{array}$$

This is an example of $n = 3$, $\mathbf{d} = (4, 5, 5)$. Boxes of color 1,2,3 are colored as pink, green, yellow. The weight χ_{\square} of each box is also labeled.

2.2. Quantum group $U_q(\widehat{\mathfrak{gl}}_n)$. We briefly recall the definition of quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_n)$ using RTT relations and Drinfeld double, see section 2.11 of [13] for details.

For $1 \leq i \leq n$ and $j > i$, let $[i; j]$ denote the length n vector whose k -th entry is equal to the number of integers equal to $k \bmod n$ in the interval $[i, j]$. (For example, when $n = 3$, $[1; 5] = (2, 1, 1)$). Define the positive half of $U_q(\widehat{\mathfrak{gl}}_n)$ by

$$(9) \quad U_q^+(\widehat{\mathfrak{gl}}_n) := \langle e_{[i;j]} \rangle_{1 \leq i < j \leq n} / \text{relations}$$

where the relations are

$$(10) \quad q^2 e_{[i;j]} e_{[i';j']} - q^2 e_{[i';j']} e_{[i;j]} \\ = (q - q^{-1}) \left(\sum_{\substack{a \equiv i \\ i' < a \leq j'}} e_{[a;j']} e_{[i+i'-a;j]} - \sum_{\substack{a \equiv j \\ i' \leq a < j'}} e_{[i;j+j'-a]} e_{[i';a]} \right)$$

(here q^2 refers to an integer power of q that won't play a role in this paper.) And $U_q^-(\widehat{\mathfrak{gl}}_n)$ is defined similarly.

The quantum group $U_q(\widehat{\mathfrak{gl}}_n)$ is defined using Drinfeld double, i.e. as a vector space

$$(11) \quad U_q(\widehat{\mathfrak{gl}}_n) = U_q^+(\widehat{\mathfrak{gl}}_n) \otimes \mathbb{Q}(q)[\psi_1^{\pm}, \dots, \psi_n^{\pm}, c^{\pm}] \otimes U_q^-(\widehat{\mathfrak{gl}}_n)$$

See formula (2.33), (2.34), (2.46) in [13] for the relations among the tensor factors.

2.2.1. Anti-involution. [13] constructed an embedding of $U_q(\widehat{\mathfrak{gl}}_n)$ into the shuffle algebra \mathcal{A} . The latter is a $\mathbb{Q}(q, p)$ algebra whose positive and negative half both have color-symmetric polynomials as basis. The shuffle algebra \mathcal{A} has an anti-involution τ which sends a polynomial in the positive/negative half to the same polynomial in the other half.

τ preserves $U_q(\widehat{\mathfrak{gl}}_n) \otimes \mathbb{Q}(q, p)$ and thus induces an anti-involution on it (still denoted τ).³ Note that although $U_q(\widehat{\mathfrak{gl}}_n)$ has nothing to do with p , the anti-involution τ does. This has to do with the fact that

$$U_q(\widehat{\mathfrak{gl}}_n) = U_q(\widehat{\mathfrak{sl}}_n) \otimes U_q(\widehat{\mathfrak{gl}}_1)$$

and the anti-involution on $U_q(\widehat{\mathfrak{sl}}_n)$ and on $U_q(\widehat{\mathfrak{gl}}_1)$ can be scaled independently, and thus gives a family of inequivalent anti-involution of $U_q(\widehat{\mathfrak{gl}}_n)$. Setting p to a number means choosing one anti-involution in this family. The τ defined here is natural in this setting because the associated Shapovalov form can be identified with Euler character pairing of K-theory with a twist, see section 2.2.4.

2.2.2. Universal Verma module and Shapovalov form. Recall the definition of $\mathbb{Z}_{\mathbf{u}}, \mathbb{F}_{\mathbf{u}}$ in equation (5), (6). Let

$$(12) \quad V^{\text{loc}} := U_q(\widehat{\mathfrak{gl}}_n) \otimes_{U_q^{\leq 0}(\widehat{\mathfrak{gl}}_n)} \mathbb{F}_{\mathbf{u}}$$

be the universal verma module of $U_q(\widehat{\mathfrak{gl}}_n)$, where the negative half $U_q^-(\widehat{\mathfrak{gl}}_n)$ act on $\mathbb{F}_{\mathbf{u}}$ by 0 and ψ_i act by scalars u_i/q^i and c act by scalar $q^n p$. The superscript “loc” stresses the fact that it is over the fraction field $\mathbb{F}_{\mathbf{u}}$.

The Shapovalov form on V^{loc} is defined by

$$(13) \quad \begin{aligned} (|\emptyset\rangle, |\emptyset\rangle) &= 1 \\ (g|u\rangle, |v\rangle) &= (|u\rangle, \tau(g)|v\rangle) \end{aligned}$$

where \emptyset stand for the highest weight vector in V^{loc} .

2.2.3. PBW basis. Defining PBW basis amounts to choosing an ordering of the root generators. To each multi-partition λ we define an element $e_{\lambda} \in U_q^+(\widehat{\mathfrak{gl}}_n)$ in the following way: split λ into vertical strips, then multiply the root generators corresponding to each strip from left to right. For example, the partition in (8) corresponds to

$$e_{\lambda} = e_{[1;5]} e_{[1;3]} e_{[2;4]} e_{[2;3]} e_{[3;7]} e_{[3;4]}$$

For each $\mathbf{d} \in \mathbb{Z}_{\geq 0}^n$, the set $\{e_{\lambda}\}_{|\lambda|=\mathbf{d}}$ form a basis of the degree \mathbf{d} piece of $U_q^+(\widehat{\mathfrak{gl}}_n)$, and any product of $e_{[i;j]}$'s can be written as a $\mathbb{Z}[q, q^{-1}]$ combination of $\{e_{\lambda}\}$.

Let $|e_{\lambda}\rangle := e_{\lambda}|\emptyset\rangle \in V^{\text{loc}}$. We will call them PBW basis. And let $|e_{\lambda}\rangle^* \in V^{\text{loc}}$ be their dual under Shapovalov form, i.e.

$$(|e_{\lambda}\rangle, |e_{\mu}\rangle^*) = \delta_{\lambda\mu}$$

³It's more common to further multiply it by an element in the Cartan part so that τ is a coalgebra homomorphism. We omit it here since it only multiplies the dual Verma module by a constant power of q (depending on degree).

2.2.4. *Action on $K_T(\mathcal{M}_{\mathbf{d}})$.* As was shown in [2, 14, 18], there is a natural action of $U_q(\widehat{\mathfrak{gl}}_n)$ on K^{loc} defined by correspondences. We briefly recall the construction and refer to [14] section 3,4 for more details.

The geometric realization relies on certain varieties $\mathfrak{Z}_{[i;j]}, \overline{\mathfrak{Z}}_{[i;j]}$ together with K-theory classes $[\mathfrak{Z}_{[i;j]}^+] \in K_T(\mathfrak{Z}_{[i;j]})$, $[\overline{\mathfrak{Z}}_{[i;j]}^-] \in K_T(\overline{\mathfrak{Z}}_{[i;j]})$ and maps

$$\begin{array}{ccc} & \mathfrak{Z}_{[i;j]} \text{ or } \overline{\mathfrak{Z}}_{[i;j]} & \\ p^+ \text{ or } \overline{p}^+ \swarrow & & \searrow p^- \text{ or } \overline{p}^- \\ \mathcal{M}_{\mathbf{d}^+} & & \mathcal{M}_{\mathbf{d}^-} \end{array}$$

The action of the root generators are given by

$$\begin{aligned} e_{[i;j]}(\alpha) &= p^+([\mathfrak{Z}_{[i;j]}^+]) \cdot p^{-*}(\alpha) \\ f_{[i;j]}(\alpha) &= \overline{p}^-([\overline{\mathfrak{Z}}_{[i;j]}^-]) \cdot p^{+*}(\alpha) \end{aligned}$$

for any class $\alpha \in K^{\text{loc}}$. These operators form an action of $U_q(\widehat{\mathfrak{gl}}_n)$.

The action can also be written explicitly in normalized fixed point basis

$$(14) \quad \langle \lambda | e_{[i;j]} | \mu \rangle = R_{ij}^+(\lambda | \mu) \prod_{\blacksquare \in \lambda \setminus \mu} \left[(q - q^{-1}) \zeta \left(\frac{\chi_{\blacksquare}}{\chi_{\mu}} \right) \tau_+(\chi_{\blacksquare}) \right]$$

where

$$(15) \quad R_{ij}^+ = \text{Sym} \left[\frac{1}{\left(1 - \frac{z_{i+1}}{z_i q^2}\right) \cdots \left(1 - \frac{z_{j-1}}{z_{j-2} q^2}\right)} \prod_{i < a < b < j} \zeta \left(\frac{z_b}{z_a} \right) \right]$$

(See [14] for the exact definition of ζ and τ_+ and how to evaluate R_{ij} on $\lambda \setminus \mu$.) In particular, the product of ζ and τ_+ above can be simplified to

$$(16) \quad \zeta \left(\frac{z}{\chi_{\lambda}} \right) \tau_+(z) = \frac{\prod_{\text{addable } \square \text{ of } \lambda}^{\text{col } \square = \text{col } z+1} \left(\frac{\sqrt{\chi_{\square}}}{q} - \frac{zq}{\sqrt{\chi_{\square}}} \right)}{\prod_{\text{removable } \square \text{ of } \lambda}^{\text{col } \square = \text{col } z} \left(\sqrt{\chi_{\square}} - \frac{z}{\sqrt{\chi_{\square}}} \right)}$$

In the moduli-of-sheaves description of affine Laumon space, $\mathfrak{Z}_{[i;j]}$ can be defined by Hecke-type operators that twist the sheaves at $0 \in \mathbb{P}^1$. To define $\mathfrak{Z}_{[i;j]}$ in the quiver description, fix $\mathbf{d}^+, \mathbf{d}^-$ such that $\mathbf{d}^+ - \mathbf{d}^- = [i; j]$. Fix two sets of vector spaces $\mathbf{V}^+, \mathbf{V}^-$ of dimension $\mathbf{d}^+, \mathbf{d}^-$ and a flag of subspaces

$$\mathbf{V}^+ = \mathbf{V}^j \xrightarrow{\rightarrow_{j-1}} \mathbf{V}^{j-1} \xrightarrow{\rightarrow_{j-2}} \cdots \xrightarrow{\rightarrow_{i+1}} \mathbf{V}^{i+1} \xrightarrow{\rightarrow_i} \mathbf{V}^i = \mathbf{V}^-$$

(\rightarrow_k means only changing the k -th component of \mathbf{V} by a one-dimensional kernel.) View $M_{\mathbf{d}^+}$ (defined in (3)) as linear maps $(\mathbf{X}, \mathbf{Y}, \mathbf{A}, \mathbf{B})$ acting on \mathbf{V}^+ and consider the subspace $Z_{[i;j]} \subset M_{\mathbf{d}^+}$ where \mathbf{X}, \mathbf{Y} satisfy certain compatibility condition with the subspaces. Let η be the moment map on $Z_{[i;j]}$ induced from ν on $M_{\mathbf{d}^+}$. Let $P_{[i;j]}$ be the parabolic subgroup that preserve the subspaces. Define

$$\mathfrak{Z}_{[i;j]} = \eta^{-1}(0)^s / P_{[i;j]}$$

The spaces $\overline{\mathfrak{Z}}_{[i;j]}$ can be defined in a similar way.

Lemma 2.1. *The maps p^+, p^-, \bar{p}^+ are proper.*

Proof. p^+, \bar{p}^+ are closed embeddings, so we only need to prove for $p = p^-$. For each local chart $U_\lambda \subset \mathcal{M}_{d^-}$ as defined in section 2.1.3, we prove that we can factor the map $p^{-1}(U_\lambda) \rightarrow U_\lambda$ by

$$(17) \quad p^{-1}(U_\lambda) \rightarrow U_\lambda \times \mathbb{P}^{N_1} \times \dots \times \mathbb{P}^{N_{j-i}} \rightarrow U_\lambda$$

where the first map is closed embedding while the second map is projection, and \times indicates a (possibly non-trivial) projective bundle.

Recall that $\mathfrak{Z}_{[i;j]}$ is defined to be $\eta^{-1}(0)^s/P_{[i;j]}$. η takes value in $\text{Hom}(V_{k-1}^-, V_k^+)$ because all X, Y kill the 1-dimensional kernel. Instead of considering $\eta^{-1}(0)$, consider the larger space

$$Z' = \eta^{-1}(\text{Hom}(V_{k-1}^-, \ker(V_k^+ \rightarrow V_k^-))),$$

i.e. we relax the moment map condition on the 1-dimensional kernel. Note that we still keep the stability condition that the image of A under X, Y generate V^+ . Pick the standard basis $\{v_\square\}_{\square \in \lambda}$ on U_λ as a basis of V_k^- . For V_k^+ , use the same basis for V_k^- plus a vector v_{ker} in the 1-dimensional kernel. Fixing a point $(X^-, Y^-, A^-, B^-) \in U_\lambda$, then a point $(X^+, Y^+, A^+, B^+) \in Z'$ in the fibre over it still has some freedom in matrix coefficients. These are coefficients in the kernel direction, coming in Y^+ (those boxes that doesn't have another box on top), X^+ (those boxes coming from Y , as well as the right most bottom box. If $A = 0$ then the right most bottom box transfer to A). In any case there will be $\dim V_{k-1}^- + 1$ undetermined coefficients in X_k^+, Y_k^+ together. They cannot all be 0 because of stability condition. They can be simultaneously scaled by our choice of v_{ker} . This shows that $Z'/P_{[i;j]}$ is an iterated projective bundle over U_λ . Now we add back the moment map conditions, which cut out the closed subspace $p^{-1}(U_\lambda)$ inside $Z'/P_{[i;j]}$. This gives the desired maps (17). \square

Remark 2.2. *Note that the map \bar{p}^- is not proper. We will address this in section 4.2.*

Under this action, there is an isomorphism ([14], see also [2] for the finite case)

$$(18) \quad \Phi : V^{\text{loc}} \xrightarrow{\cong} K^{\text{loc}}$$

sending highest weight vector in V^{loc} to the structure sheaf in degree 0. The Shapovalov form on V^{loc} induces a bilinear form on K^{loc} given by (cf. [2] proposition 2.29)

$$(19) \quad (\mathcal{F}, \mathcal{G}) = \text{const}_d \cdot \chi(\mathcal{F} \otimes \mathcal{G} \otimes \mathcal{D}_d^{-1})$$

where

$$\mathcal{D}_d = \prod_{i=1}^n \det \mathcal{V}_i$$

is the determinant line bundle of tautological bundles and const_d is an invertible element in \mathbb{Z}_u (i.e. monomial with coefficient 1) depending on d .

2.3. Specializing to finite Laumon spaces and $U_q(\mathfrak{sl}_n)$. When $d_n = 0$, affine Laumon space becomes the finite Laumon space initially defined in [10], which we denote by $\mathcal{M}_{\mathbf{d}}^{\text{fin}}$ where $\mathbf{d} = (d_1, \dots, d_{n-1})$, see [14] remark 3.3. In this section, we spell out what the above constructions mean in this special case.

The quiver becomes (c.f. [12] section 2(i))




Figure 2.3

The fixed points are parametrized by $n-1$ tuples of 2d partitions $\lambda = (\lambda^1, \dots, \lambda^{n-1})$ where each λ^i itself is a 2d partition and such that λ^i has at most $n-i$ rows. We will call such λ a Kostant partition, following the nomenclature in [7].

Below is an example of $n = 4$ with degree $\mathbf{d} = (4, 4, 3)$. Boxes of color 1,2,3 are colored as pink, green, yellow.

$$(20) \quad \begin{array}{cccc} & \text{yellow} & & \\ & u_1^2 & & \\ & \text{green} & \text{green} & \\ & u_1^2 & u_1^2 q^2 & \\ \text{pink} & u_1^2 & u_1^2 q^2 & u_1^2 q^4 & u_1^2 q^6 \\ & \text{green} & \text{green} & & \text{yellow} \\ & u_2^2 & u_2^2 q^2 & & u_3^2 \end{array}$$

For the quantum group action, only root generators $e_{[i;j]}, f_{[i;j]}$ with $1 \leq i < j \leq n$ preserves $K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{fin}})$. Together with ψ_i they generate a subalgebra

$$U_q(\mathfrak{sl}_n) \hookrightarrow U_q(\widehat{\mathfrak{gl}}_n)$$

The generators $e_{[i;i+1]}$ correspond to the usual generators of $U_q(\mathfrak{sl}_n)$ multiplied by $q - q^{-1}$. And from relation (10) we have

$$qe_{[i+1;j]}e_{[i;i+1]} - e_{[i;i+1]}e_{[i+1;j]} = (q - q^{-1})e_{[i;j]}$$

for $1 \leq i < j \leq n$.

It is worth noting that for finite Laumon space, the relative normal bundle

$$(21) \quad [T\mathcal{M}_{\mathbf{d}}] - [T\mathfrak{Z}_{[i;j]}] = j - i + \left(1 - \frac{1}{q^2}\right) \left(\sum_{i \leq a < j} \frac{\mathcal{V}_{a+1}^+}{\mathcal{L}_a} - \sum_{i \leq a < j} \frac{\mathcal{V}_a^+}{\mathcal{L}_a} \right) - \sum_{a=i+1}^{j-1} \frac{\mathcal{L}_a}{\mathcal{L}_{a-1}q^2} + \sum_{a=i}^{j-1} \frac{u_{a+1}^2}{\mathcal{L}_a q^2}$$

Here, \mathcal{V}_a^+ (resp. \mathcal{V}_a^-) is the tautological bundle for $\mathcal{M}_{\mathbf{d}^+}$ (resp. $\mathcal{M}_{\mathbf{d}^-}$) and \mathcal{L}_a is their difference on $\mathfrak{Z}_{[i;j]}$. This is, of course, equivalent to formula (14), but is more convenient for the proof in the next section.

3. $K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{FIN}})$ AND DUAL VERMA MODULE OF $U_q(\mathfrak{sl}_n)$

In this section, we will focus on the finite Laumon space $K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{fin}})$ and all notations will be adapted to the finite case. \mathbf{d} will denote a vector of length $n - 1$, and boldface letters (e.g. $\boldsymbol{\lambda}$) will denote Kostant partitions as explained in section 2.3. $K^{\text{loc}}, K^{\text{int}}, V^{\text{loc}}, \Phi$ are understood as defined for $\mathcal{M}_{\mathbf{d}}^{\text{fin}}$ and $U_q(\mathfrak{sl}_n)$.

Theorem 3.1. (1) For any partition $\boldsymbol{\lambda}$, Φ maps the dual PBW basis $|e_{\boldsymbol{\lambda}}\rangle^*$ into $K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{fin}})$ where $\mathbf{d} = |\boldsymbol{\lambda}|$.

(2) For any $\mathbf{d} \in \mathbb{Z}_{\geq 0}^{n-1}$, $\{|e_{\boldsymbol{\lambda}}\rangle^*\}_{|\boldsymbol{\lambda}|=\mathbf{d}}$ form a $\mathbb{Z}_{\mathbf{u}}$ -basis of $K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{fin}})$.

Remark 3.2. A similar result was obtained in [5] theorem 3.5 when studying equivariant cohomology of Laumon space as (non-quantum) $U(\mathfrak{gl}_n)$ -module. The approach we use here is different from theirs.

Since the action of $U_q(\mathfrak{sl}_n)$ preserves $K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{fin}})$, we have

Corollary 3.3. Fix $q_0 \in \mathbb{C}$ a non-zero complex number and $a_1, \dots, a_n \in \mathbb{C}$. Let $\mathbb{C}_{\mathbf{a}}$ denote the evaluation module of $\mathbb{Z}_{\mathbf{u}}$, i.e. the target of the map

$$\mathbb{Z}_{\mathbf{u}} = \mathbb{Z}[u_1^{\pm}, \dots, u_n^{\pm}, q^{\pm}] \xrightarrow{u_i=q^{a_i}, q=q_0} \mathbb{C}$$

Then $K^{\text{int}} \otimes_{\mathbb{Z}_{\mathbf{u}}} \mathbb{C}_{\mathbf{a}}$ is isomorphic to the dual Verma module of $U_{q_0}(\mathfrak{sl}_n)$ with highest weight (b_1, \dots, b_{n-1}) where $b_i = a_i - a_{i+1} + 1$

Proof of Theorem 3.1. First, observe that part (1) of the theorem implies part (2). Indeed, given any $\alpha \in K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{fin}})$ and any PBW basis $|e_{\boldsymbol{\lambda}}\rangle$, using the pairing (19), we have

$$(\alpha, \Phi(|e_{\boldsymbol{\lambda}}\rangle)) = (\tau(e_{\boldsymbol{\lambda}})\alpha, \Phi(|\emptyset\rangle)).$$

Note that $\tau(e_{\boldsymbol{\lambda}})\alpha$ is an integral K-theory class due to Lemma 2.1. Thus the pairing lives in $\mathbb{Z}_{\mathbf{u}}$. By definition, the dual PBW basis satisfy

$$(\Phi(|e_{\boldsymbol{\lambda}}\rangle^*), \Phi(|e_{\boldsymbol{\mu}}\rangle)) = \delta_{\boldsymbol{\lambda}}^{\boldsymbol{\mu}}$$

This implies that α is in the $\mathbb{Z}_{\mathbf{u}}$ -linear span of $\{\Phi(|e_{\boldsymbol{\lambda}}\rangle^*)\}_{|\boldsymbol{\lambda}|=\mathbf{d}}$.

Thus, we only need to prove that $|e_{\boldsymbol{\lambda}}\rangle^* \in K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}}^{\text{fin}})$. We define a partial order on the set of fixed points of $\mathcal{M}_{\mathbf{d}}^{\text{fin}}$ for a given \mathbf{d} . If $\boldsymbol{\mu}$ can be obtained from $\boldsymbol{\lambda}$ by moving a vertical strip of boxes rightward, then we say $\boldsymbol{\mu} \prec \boldsymbol{\lambda}$. If $\boldsymbol{\mu} \prec \boldsymbol{\lambda}, \boldsymbol{\lambda} \prec \boldsymbol{\nu}$, we say $\boldsymbol{\mu} \prec \boldsymbol{\nu}$. Refine this to a total order, which we denote by $<$. In this order, the flat partition (i.e. each partition has only one row) is the smallest.

Lemma 3.4. $|e_{\boldsymbol{\lambda}}\rangle$ is only non-zero on $|\boldsymbol{\mu}\rangle$ for $\boldsymbol{\mu} \prec \boldsymbol{\lambda}$

Proof. Recall that formula (14) implies that $\langle \lambda | e_{[i;j]} | \mu \rangle$ is only non-zero if λ can be obtained from μ by adding boxes of color $i, i + 1, \dots, j - 1$.

We prove the lemma by induction. Assume the lemma is proven for partitions smaller than λ . Let λ' be λ removing the left most strip $[i; j]$. Let $\mu' \prec \lambda'$ and let μ be μ' adding vertical strip $[i; j]$ to the left. Then by definition $\mu \prec \lambda$. The induction hypothesis implies that $\langle \nu | e_{[i;j]} | \mu' \rangle \neq 0$ only if $\nu \prec \mu'$. (Otherwise there's a minimal μ'_0 which under $e_{[i;j]}$ produces a fixed point greater than μ which is impossible.) So $\nu \prec \mu \prec \lambda$ and we are done. \square

Dual PBW basis is uniquely determined by the property that

$$\langle e_\lambda \rangle^*, |e_\mu \rangle = \delta_\lambda^\mu$$

For each partition λ with $|\lambda| = \mathbf{d}$, we will construct elements $g_\lambda \in K_{\mathbf{d}}^{\text{int}}$ satisfying the following properties:

- (P1) $g_\lambda | \mu \rangle \neq 0$ only if $\lambda \prec \mu$
(P2) $g_\lambda | \lambda \cdot |e_\lambda \rangle | \lambda = \Lambda^\bullet(T\mathcal{M}_{\mathbf{d}}^{\text{fin}} | \lambda)^\vee$

Combining with Lemma 3.4, we see that for $\mu < \lambda$, $(g_\lambda, |e_\mu \rangle)$ must be 0 since at least one of them vanish on any fixed point. On the other hand, $(g_\lambda, |e_\lambda \rangle)$ only get contribution from fixed point λ , so (P2) ensures that it is equal to 1 after scaling g_λ by an invertible element in $\mathbb{Z}_{\mathbf{u}}$ which account for the const_d and twist by $\mathcal{D}_{\mathbf{d}}$ in formula (19). To sum up, the inner product between a suitable rescaling of $\{g_\lambda\}$ and $\{|e_\mu \rangle\}$ is upper-triangular with 1 on the diagonal.

$$\begin{pmatrix} 1 & & & & & \\ & 1 & & & & * \\ & & \ddots & & & \\ & 0 & & & & 1 \\ & & & & & & 1 \end{pmatrix}$$

This will allow us to construct $|e_\lambda \rangle^*$ as a $\mathbb{Z}_{\mathbf{u}}$ -linear combination of g_λ and thus $|e_\lambda \rangle^* \in K_{\mathbf{d}}^{\text{int}}$.

The desired class g_λ is constructed through attracting manifold with respect to the torus action. Recall that if a torus $\mathbb{T} = (\mathbb{C}^*)^k$ acts on a variety X , given a fixed point $p \in X^{\mathbb{T}}$ and a subtorus $\sigma : \mathbb{C}^* \rightarrow \mathbb{T}$, attracting manifold is defined to be

$$\text{Attr}_p := \{x \in X \mid \lim_{t \rightarrow 0} \sigma(t) \cdot x = p\}$$

In our situation, consider the torus \mathbb{T} action on $\mathcal{M}_{\mathbf{d}}^{\text{fin}}$ as in (4) and let u_1, \dots, u_n, q be the weights of \mathbb{T} . Let Attr_λ be the attracting locus of λ with respect to a subtorus where tangent weight q^{-k} is attracting for $k > 0$. $u_j^2/u_i^2 q^k$ is attracting for $j > i$ and any k . (For example, choose $\sigma(t) = (t^{2|\mathbf{d}|}, t^{4|\mathbf{d}|}, \dots, t^{2n|\mathbf{d}|}, t^{-1})$.) Let

$$g_\lambda := i_*[\overline{\mathcal{O}_{\text{Attr}_\lambda}}] \in K_T(\mathcal{M}_{\mathbf{d}}^{\text{fin}})$$

Here $\overline{\text{Attr}_\lambda}$ is the closure of Attr_λ , a closed subvariety of $\mathcal{M}_{\mathbf{d}}^{\text{fin}}$, and i is the inclusion map. It remains to prove that g_λ satisfies properties (P1) and (P2). Now the proof reduces to the following two propositions:

Proposition 3.5. *For fixed points $\lambda, \mu \in K_T(\mathcal{M}_d^{\text{fin}})$, if $g_\lambda|_\mu \neq 0$, then $\lambda \prec \mu$.*

Proof. For any point μ such that $\lambda \not\prec \mu$, there exists a color c and $i \leq c$ such that

$$\sum_{k=i}^c \lambda_{[c]}^k < \sum_{k=i}^c \mu_{[c]}^k$$

Recall the convention about color in section 2.1.2. (Otherwise we can always move things rightward to go from μ to λ .)

Let C be the set of color c boxes in μ^i, \dots, μ^c . Consider the open subset

$$(22) \quad U_C = \{v_\square\}_{\square \in C} \text{ are linearly independent}$$

This is an open subset in $\mathcal{M}_d^{\text{fin}}$ that contains μ .

On the other hand, consider the standard coordinates on the open subset U_λ defined in section 2.1.3. For any point in $U_\lambda \cap U_C$, the matrix X_c must have a non-zero coordinate that goes from a box in partition $\geq i$ to a partition $< i$, due to the linearly independence in (22). This coordinate has weight $u_{<i}^2/u_{\geq i}^2$ and thus this point cannot be in the attracting locus of λ . Note that U_λ is invariant under T so $\text{Attr}_\lambda \subset U_\lambda$. Thus $\text{Attr}_\lambda \cap U_C = \emptyset$. This finishes the proof. \square

Proposition 3.6. $\Lambda^\bullet T_\lambda^\vee \text{Attr}_\lambda$ is equal to $|e_\lambda\rangle|_\lambda$ up to invertible constant in \mathbb{Z}_u .

Note that this implies property (P2) because $g_\lambda|_\lambda = T_{\lambda, \text{repelling directions}}^\vee$

Proof. Prove by induction. Let μ be the partition obtained from λ by removing the left most vertical strip $[i; j)$. Assume that the proposition is proved for μ . Let's examine how the two expressions change for λ .

Lemma 3.4 implies that the coefficient $|e_\lambda\rangle|_\lambda$ comes only from $|e_\mu\rangle|_\mu$. So it changes by the exterior power of relative tangent bundle (21) up to invertible constants. So to prove the proposition, it suffices to prove that

$$(21) = T_\lambda \text{Attr}_\lambda - T_\mu \text{Attr}_\mu$$

(Note that $\Lambda^\bullet V$ and $\Lambda^\bullet V^\vee$ differ by an invertible constant, thus we get rid of the dual in the above formula.)

On the other hand, $T_\lambda^\vee \text{Attr}_\lambda$ can be computed from formula (7). To compute the change of weights from μ to λ , recall that all new boxes are added in partition μ^i while existing boxes are all in partition $\mu^{\geq i}$, so they contribute to attracting weight if either they appear in the denominator, or they appear in the numerator but the

denominator is a box in the same partition. More precisely, we have

$$(23) \quad T_\lambda^\vee \text{Attr}_\lambda - T_\mu^\vee \text{Attr}_\mu = \\ j - i + \left(1 - \frac{1}{q^2}\right) \left(\sum_{i \leq a < j} \frac{\mathcal{V}_{a+1}^+}{\mathcal{L}_a} - \sum_{i \leq a < j} \frac{\mathcal{V}_a^+}{\mathcal{L}_a} \right) + \sum_{a=i}^{j-1} \frac{u_{a+1}^2}{\mathcal{L}_a q^2} \\ - \sum_{a=i+1}^{j-1} \frac{\mathcal{L}_a}{\mathcal{L}_{a-1} q^2}$$

Here, the first line is when \mathcal{L}_a (corresponding to the added box) appear in the denominator. The $j - i$ at the beginning is to remove the constant factor appeared in $\mathcal{V}_a^+/\mathcal{L}_a$. The second line is when \mathcal{L}_a appears in the numerator with denominator being something in partition i . This only happens in the term $(1 - q^{-2})(\mathcal{V}_{i+1}/\mathcal{V}_i)$ in (7). Formula (23) exactly matches with formula (21) and this finishes the proof. \square

4. $K_\tau(\mathcal{M}_\mathbf{d})$ AND DUAL VERMA MODULE OF $U_q(\widehat{\mathfrak{gl}}_n)$

We will prove the analog of theorem 3.1 in the affine case:

Theorem 4.1. (1) For any partition λ , Φ maps the dual PBW basis $|e_\lambda\rangle^*$ into $K_\tau(\mathcal{M}_\mathbf{d})$ where $\mathbf{d} = |\lambda|$.

(2) For any $\mathbf{d} \in \mathbb{Z}_{\geq 0}^n$, $\{|e_\lambda\rangle^*\}_{|\lambda|=\mathbf{d}}$ form a $\mathbb{Z}_\mathbf{u}$ -basis of $K_\tau(\mathcal{M}_\mathbf{d})$.

The theorem will be proved in section 4.1.

The analog of corollary 3.3 requires a slight modification due to the possible denominators in $f_{[i;j]}$ action.

Corollary 4.2. Fix $q_0 \in \mathbb{C}$ a non-zero complex number, $a_1, \dots, a_n, l \in \mathbb{C}$ such that q_0^l is not a root of unity. Let \mathbb{C}_a denote the evaluation module of $\mathbb{Z}_\mathbf{u}$, i.e. the target of the map

$$\mathbb{Z}_\mathbf{u} = \mathbb{Z}[u_1^\pm, \dots, u_n^\pm, q^\pm, p^\pm] \xrightarrow{u_i=q^{a_i}, p=q^l, q=q_0} \mathbb{C}$$

Then $K^{\text{int}} \otimes_{\mathbb{Z}_\mathbf{u}} \mathbb{C}_a$ is isomorphic to the dual Verma module of $U_{q_0}(\widehat{\mathfrak{gl}}_n)$ where ψ_i and c act by q^{i-a_i} and q^{n+l} .

This will be proved in section 4.2. The main task is to control the denominators appeared in the action, see proposition 4.12.

Remark 4.3. In the above corollary, $l = 0$ corresponds to the critical level $c = q^n$. Away from the critical level, we can always find generic q_0 so that the result applies.

Remark 4.4. Using the RTT integral form of $U_q(\widehat{\mathfrak{gl}}_n)$ as defined in [8], one can define the integral form of universal Verma module similar to (12), as well as the integral form of universal dual Verma module. Then the results above says that, after a mild localization, K^{int} can be identified with the universal dual Verma module.

4.1. Stable envelopes. In this section, we use a variant of the notion of stable envelope first introduced in [11] to prove the main theorem 4.1.

The same argument as in section 3 shows that part (1) of the theorem implies part (2). To establish part (1), it suffices to find K-theory classes whose pairing with PBW basis is an upper triangular matrix. This is where stable envelopes come in handy.

4.1.1. To begin with, we collect the ingredients used in the definition of stable envelopes. Consider the subtorus $\mathbf{A} = (u_1, \dots, u_n, p) \subset \mathbb{T}$ and fix a 1-parameter subgroup

$$(24) \quad \sigma_{\mathbf{A}}(t) = (t^{k_1}, \dots, t^{k_n}, t^s)$$

with $k_1 < k_2 < \dots < k_n < k_1 - s$. From now on, we will only consider attracting manifold for \mathbf{A} action with respect to $\sigma_{\mathbf{A}}$. The connected components of \mathbf{A} fixed points of $\mathcal{M}_{\mathbf{d}}$ are affine spaces and each contains exactly one point of $(\mathcal{M}_{\mathbf{d}})^{\mathbb{T}}$.

Let

$$\begin{aligned} \mathcal{V} &= \bigoplus_{i=1}^n \mathcal{A}_i \\ \mathcal{U} &= \bigoplus_{i=1}^n \mathcal{B}_i \\ L &= \left(\prod_{i=1}^n \det \mathcal{V}_i \right)^{\epsilon} \end{aligned}$$

where $\mathcal{A}_i, \mathcal{B}_i, \mathcal{V}_i$ denote the tautological bundle from A_i, B_i, V_i and $\epsilon > 0$ is a very small real number. (L will be used in the definition of stable envelopes, and the non-integer power ϵ will be interpreted properly.)

It's straightforward to check that

Lemma 4.5. *For each fixed point $\lambda \in \mathcal{M}_{\mathbf{d}}$,*

$$\begin{aligned} (\mathcal{V}|\lambda)_{mov} &= (T\mathcal{M}_{\mathbf{d}}|\lambda)_{rep} \\ \mathcal{U}|\lambda &= (T\mathcal{M}_{\mathbf{d}}|\lambda)_{attr} \end{aligned}$$

where the subscript *mov* means the moving (non-fixed) part with respect to the torus \mathbf{A} and *rep/attr* means repelling/attracting directions.

Definition 4.6. *For an element $\alpha \in K_{\mathbb{T}}(X)$ where X is an algebraic variety with torus \mathbb{T} action. Suppose $\mathbf{A} \subset \mathbb{T}$ is a subtorus such that X is fixed under \mathbf{A} action so that $K_{\mathbb{T}}(X) = K_{\mathbb{T}/\mathbf{A}}(X) \otimes K_{\mathbf{A}}(pt)$. Then the degree of α is defined as the Newton polygon of α viewed as a Laurent polynomial with coefficients in $K_{\mathbb{T}/\mathbf{A}}(X)$. In other words, let*

$$S = \{w \in \text{char}(A) \mid \text{coefficient of } a^w \text{ in } \alpha \text{ is not } 0\}$$

Then $\text{deg}_{\mathbf{A}} \alpha$ is the convex hull of S in $\text{char}(A) \otimes \mathbb{R}$.

We write $\mu \succ \lambda$ if μ is in the full attracting locus (i.e. taking attracting locus and closure iteratively) of λ . This defines a partial ordering on the fixed points.

Combinatorially, this means that μ can be obtained from λ by moving a vertical strip of boxes upwards. We also define a total ordering on the fixed points. We say that $\mu > \lambda$ if $h(\mu) > h(\lambda)$ where h is the sum of the height of boxes in μ or λ .

Proposition 4.7. *For each λ , there exists $s_\lambda \in K_{\mathbb{T}}(\mathcal{M}_{\mathbf{d}})$ such that*

- (i) s_λ is supported on the full attracting manifold of λ .
- (ii) $s_\lambda|_\lambda = \Lambda^\bullet(\mathcal{V}_\lambda^{mov})^\vee$
- (iii) For any $\mu \succ \lambda$,

$$\deg_{\mathbb{A}} s_\lambda|_\mu + \text{weight}_{\mathbb{A}} L|_\lambda \subset \deg_{\mathbb{A}} \Lambda^\bullet(\mathcal{V}_\lambda^{mov})^\vee + \text{weight}_{\mathbb{A}} L|_\mu$$

The classes s_λ above can be viewed as a generalized version of K-theoretic stable envelope introduced in [16]. Their existence was essentially proved in [17] theorem 2. The version we use here is slightly different in that $\mathcal{M}_{\mathbf{d}}$ does not have a polarization. But as indicated in section 2.1.4 of [17], having a vector bundle \mathcal{V} satisfying lemma 4.5 is good enough. The proof can be adapted to this situation with minor changes. We include the proof in appendix A for completeness.

4.1.2. We will leverage the “rigidity argument” introduced in [16] to compute the inner product of stable envelopes and PBW basis. The logic can be summarized as follows: first, by localization formula (27), the inner product of K-theory classes can be computed by summing over fixed points. The degree constraint (property (iii) in the definition of s_λ , and lemma 4.9 below) implies that contribution from each fixed point is “bounded” in the sense defined below. Hence, the left hand side of (27) also has this property. The properness of correspondences imply that LHS is a Laurent polynomial. Together with boundedness, this implies that LHS is constant in \mathbb{A} . Thus it can be computed by sending equivariant variables in \mathbb{A} to any limit, for which we can choose a convenient one to make most fixed point contributions vanish, see proposition 4.11.

For a function $f \in \mathbb{Z}_{\mathbf{u}}$ and a 1-parameter subgroup $\xi(t) : \mathbb{C}^* \rightarrow \mathbb{A}$, denote

$$\lim_{\xi(t) \rightarrow \infty} f := \lim_{t \rightarrow \infty} f|_{(u_1, \dots, u_n, p) = \xi(t)}$$

(Note that q is left untouched in f in the above definition.) We say that f remains bounded in any limit of \mathbb{A} (or bounded for short) if for any subgroup $\xi(t)$, $\lim_{\xi(t) \rightarrow \infty} f$ is a well-defined function of q .

Recall the subgroup $\sigma_{\mathbb{A}}(t)$ defined in formula (24).

Lemma 4.8. *Given fixed point λ, μ , let*

$$f_{\lambda\mu} = \frac{s_\lambda|_\mu}{\Lambda^\bullet(T_\mu^{<0})^\vee}$$

then

- (1) If $\mu < \lambda$, then $s_\lambda|_\mu = 0$
- (2) If $\mu > \lambda$, then $f_{\lambda\mu}$ is bounded in any limit of \mathbb{A} , and $\lim_{\sigma_{\mathbb{A}}(t) \rightarrow \infty} f_{\lambda\mu} = 0$
- (3) If $\mu = \lambda$, then $f_{\lambda\mu} = 1$

Proof. This follows from the definition of s_λ . (In fact, the stronger statement with \langle (resp. \succ) replaced by $\not\prec$ (resp. \succ) is also true.) \square

Let

$$(25) \quad |\widetilde{e_\lambda}\rangle := |e_\lambda\rangle \left/ \prod_{i=1}^n u_{i+1}^{d_i} \right.$$

(By convention $u_{n+1} = u_1 p^{-2}$.)

Lemma 4.9. *Given fixed point λ, μ , let*

$$g_{\lambda\mu} = \frac{|\widetilde{e_\lambda}\rangle|_\mu}{\Lambda \bullet (T_{\mu}^{\geq 0})^\vee}$$

then $g_{\lambda\mu}$ remains bounded in any limit of A . Furthermore, if $\mu < \lambda$, then $\lim_{\sigma_\Lambda(t) \rightarrow \infty} g_{\lambda\mu} = 0$.

Proof. Recall that $|e_\lambda\rangle$ is constructed by applying $e_{[i;j]}$ coming from the columns of λ . We will prove that each $e_{[i;j]}$ produce a coefficient that is bounded in A when properly combined with other factors in $g_{\lambda\mu}$.

To this end, recall formula (16), where we can rewrite

$$(26) \quad \zeta\left(\frac{z}{\chi_\lambda}\right)\tau_+(z) = \frac{\prod_{\substack{\text{col } \square = \text{col } z+1 \\ \text{addable } \square \text{ of } \lambda \text{ not on bottom row}}} \left(\frac{\sqrt{\chi_\square}}{q} - \frac{zq}{\sqrt{\chi_\square}}\right)}{\prod_{\substack{\text{col } \square = \text{col } z \\ \text{removable } \square \text{ of } \lambda}} \left(\sqrt{\chi_\square} - \frac{z}{\sqrt{\chi_\square}}\right)} \cdot \left(1 - \frac{zq^2}{\chi_\square}\right) \cdot \frac{\sqrt{\chi_\square}}{q}$$

where \square stand for the addable box on the bottom row of the $i+1$ -th partition in λ where $i = \text{col } z$. In particular, $\chi_\square = u_{i+1}^2 q^{\lambda_{i+1}^{i+1}}$. Thus the last term in (26) cancels the normalization factor in (25). The first term in (26) is obviously bounded since removable box in color i and addable box of color $i+1$ comes in pairs. From lemma 4.5, for each \blacksquare there is a corresponding attracting direction with weight

$$w = \frac{u_{i+1}^2}{\chi_{\blacksquare} q^{\dots}}$$

where q^{\dots} stand for some power of q . Then

$$\frac{1 - \frac{\chi_{\blacksquare} q^2}{\chi_\square}}{1 - w^{-1}}$$

is bounded. On the other hand, the $1/(1 - z_{i+1}/z_i q^2)$ appeared in $R^+(\lambda \setminus \mu)$ is obviously bounded. This proves that $g_{\lambda\mu}$ is bounded for any limit of A .

When $\mu < \lambda$, at least one z_{i+1}/z_i in $R(\lambda \setminus \mu)$ goes to ∞ in the limit $\lim_{\sigma_\Lambda(t) \rightarrow \infty}$ since box in lower places goes to ∞ faster. This makes $g_{\lambda\mu} \rightarrow 0$ in the limit. \square

Recall the pairing on $K_T(\mathcal{M}_d)$ defined in (19). Let $\widetilde{s}_\lambda = s_\lambda \otimes \mathcal{D}_d$ to compensate for the twist in the pairing.

Lemma 4.10. *For any fixed point λ, μ , the pairing $(\tilde{s}_\lambda, \widetilde{|e_\mu\rangle}) \in \mathbb{Z}[q, q^{-1}]$, i.e., it is a Laurent polynomial that only depends on q .*

Proof. By lemma 2.1, $(\tilde{s}_\lambda, \widetilde{|e_\mu\rangle}) \in \mathbb{Z}_u$. By localization formula,

$$(27) \quad (\tilde{s}_\lambda, \widetilde{|e_\mu\rangle}) = \sum_{\text{fixed point } \nu} f_{\lambda\nu} g_{\mu\nu}$$

By lemma 4.8 and 4.9, each term on the right hand side is bounded in any limit of A , thus so is the left hand side. But the LHS is a Laurent polynomial in \mathbb{Z}_u , thus it must not depend on u_1, \dots, u_n, p , hence an element in $\mathbb{Z}[q, q^{-1}]$. \square

Proposition 4.11. *For $\mu > \lambda$, $(\tilde{s}_\lambda, \widetilde{|e_\mu\rangle}) = 0$. For $\mu = \lambda$, $(\tilde{s}_\lambda, \widetilde{|e_\lambda\rangle}) = q^N$ for some integer N .*

Proof. Since $(\tilde{s}_\lambda, \widetilde{|e_\mu\rangle})$ does not depend on u_1, \dots, u_n, p , it can be computed in any limit of A .

For $\mu > \lambda$, for any fixed point ν , either $\mu > \nu$ or $\nu > \lambda$ (or both), hence by lemma 4.8 and 4.9, either $\lim_{\sigma_A(t) \rightarrow \infty} g_{\mu\nu} = 0$ or $f_{\lambda\nu} = 0$. So the RHS of (27) is 0.

For $\mu = \lambda$, we compute $(\tilde{s}_\lambda, \widetilde{|e_\lambda\rangle})$ in the limit $\sigma_A(t) \rightarrow 0$. The only contribution comes from $f_{\lambda\lambda} g_{\lambda\lambda}$. Note that other fixed point may have non-trivial contribution *before* taking the limit, but they all vanish in the limit because, similar to the proof of lemma 4.9, at least one of z_{i+1}/z_i in R^+ goes to ∞ because moving boxes lower makes it larger. For the same reason, the symmetrization at the beginning of R^+ in formula (15) can be dropped.

Now consider rational functions of the form

$$\frac{1 - \frac{\lambda_{\square}}{\lambda_{\blacksquare}} q^k}{1 - \frac{\lambda_{\square}}{\lambda_{\blacksquare}}}$$

for some integer k . If \square, \blacksquare are not in the same partition λ^i of λ , then the limit is either 1 or q^k , and can be dropped for the purpose of this proof. The non-trivial factors come from the following (again we are focusing on $e_{[i,j]}$ coming from one column of λ as in the proof of lemma 4.9, and note that all boxes must land in standard position in the i -th partition, and when applying it there is no box above the $(j-i)$ -th row):

- the term in (26) is only non-trivial when the removable box is next to \blacksquare and the addable box is one row above \blacksquare , this results in a factor of

$$\frac{1 - q^{2(\lambda_m^i - \lambda_{m+1}^i + 1)}}{1 - q^2}$$

where $m = \text{col } \blacksquare - i$.

- in formula (14), each \blacksquare comes with a factor of $q - q^{-1}$, this cancels the denominator above.

- in R^+ , for adjacent boxes \square, \blacksquare with $\text{col } \square = m - i + 1, \text{col } \blacksquare = m - i,$

$$\frac{1}{1 - \frac{\chi_{\square}}{\chi_{\blacksquare} q^2}} = \frac{1}{1 - q^{2(\lambda_{m+1}^i - \lambda_m^i - 1)}}$$

This almost cancels the numerator in the first item, except for one term from $m = j$. (Note that $\lambda_{j+1}^i = 0$ because we apply columns from right to left.)

- The pure q weight in $\Lambda^\bullet(T_{\lambda}^{\geq 0})^\vee$ (i.e., $\Lambda^\bullet(T_{\lambda}^{=0})^\vee$). It can be computed from formula (7) that each $e_{[i;j]}$ will increase this by $1 - q^{2\lambda_j^i + 1}$, this cancels the remaining term from the first item.

So we see that $(\widetilde{s}_{\lambda}, |\widetilde{e}_{\lambda}\rangle)$ is indeed a power of q . □

Therefore, the pairing between \widetilde{s}_{λ} and $|\widetilde{e}_{\mu}\rangle$ when λ, μ ranging through all fixed points is a lower triangular matrix with powers of q on the diagonal and hence its inverse lies in $\mathbb{Z}_{\mathbf{u}}$. Thus, the dual PBW basis are $\mathbb{Z}_{\mathbf{u}}$ -linear combinations of \widetilde{s}_{λ} 's, and hence lives in $K_{\top}(\mathcal{M}_{\mathbf{d}})$. This finishes the proof of theorem 4.1.

4.2. $U_q(\widehat{\mathfrak{gl}}_n)$ action almost preserves $K_{\top}(\mathcal{M}_{\mathbf{d}})$. The action of $e_{[i;j]}$ and $\tau(e_{[i;j]})$ preserves K^{int} by lemma 2.1. However, $f_{[i;j]}$ does not. Nevertheless, we can control the denominator for the $f_{[i;j]}$ action and thus proves corollary 4.2.

For a given $[i;j]$, consider the element $\tau(f_{[i;j]})$ and write it as a $\mathbb{Q}(q, p)$ -linear combination of PBW basis e_{λ} , i.e.

$$(28) \quad \tau(f_{[i;j]}) = \sum_{|\lambda|=[i;j]} c_{\lambda} e_{\lambda}$$

where the coefficients c_{λ} are, a priori, elements of $\mathbb{Q}(q, p)$. We will control the denominators of c_{λ} and this in turn controls the denominator of $f_{[i;j]}$ action.

Let

$$(29) \quad \mathbb{Z}[q^{\pm}, p^{\pm}]_{1-p^{\#}} := \left\{ \frac{f}{g} \mid f \in \mathbb{Z}[q^{\pm}, p^{\pm}], g = \prod_{i=1}^m (1 - p^{2k_i}) \text{ for some integers } k_1, \dots, k_m \right\}$$

Proposition 4.12. *The coefficients c_{λ} live in $\mathbb{Z}[q^{\pm}, p^{\pm}]_{1-p^{\#}}$*

This will be proved in section 4.2.2. This implies that $f_{[i;j]}$ can be written as a sum of products of $\tau(e_{[i';j']})$ whose coefficients has denominator no worse than a product of $1 - p^{2k}$. Hence the action of $U_q(\widehat{\mathfrak{gl}}_n)$ preserves $K_{1-p^{\#}}^{\text{int}}$, where the subscript $1 - p^{\#}$ means inverting elements of the form $1 - p^{2k}, k \in \mathbb{Z}$. This combined with theorem 4.1 proves corollary 4.2.

4.2.1. Recall some basic facts about shuffle algebra and $U_q(\widehat{\mathfrak{gl}}_n)$ from [13, 15].

It was proved in [13] that

$$U_q(\widehat{\mathfrak{gl}}_n) = U_q(\widehat{\mathfrak{sl}}_n) \otimes U_q(\widehat{\mathfrak{gl}}_1)$$

$e_{[i;i+1]}, i = 1, 2, \dots, n$ generate $U_q^+(\widehat{\mathfrak{sl}}_n)$ and $P_k, k = 1, 2, \dots$ generate $U_q^+(\widehat{\mathfrak{gl}}_1)$. Similar for the negative half. Let $e_i := e_{[i;i+1]}$ in what follows.

There is a non-degenerate bilinear pairing between $U_q^+(\widehat{\mathfrak{gl}}_n)$ and $U_q^-(\widehat{\mathfrak{gl}}_n)$ given by

$$\begin{aligned} \langle e_i, f_j \rangle &= (q - q^{-1})\delta_j^i \\ \langle P_k, P_l \rangle &= \delta_{k+l} k \cdot \frac{(q^{nk} - q^{-nk})(c^k - c^{-k})}{(p^k - p^{-k})(q^{nk}p^k - q^{-nk}p^{-k})} \end{aligned}$$

on generators and satisfies

$$\langle a * a', b \rangle = \langle a \otimes a', \Delta(b) \rangle$$

From these we can deduce that the pairing

$$(30) \quad \langle e_{i_1} \dots e_{i_m} P_{j_1} \dots P_{j_l}, f_{i'_1} \dots f_{i'_m} P_{-j'_1} \dots P_{-j'_l} \rangle$$

(assuming j and j' are in increasing order since the P 's commute) is non-zero only when $m = m', l = l', j_i = j'_i$ for $i = 1, \dots, l$, in which case the pairing is a

$$\prod_{i=1}^l \langle P_{j_i}, P_{-j_i} \rangle \cdot \text{a rational function in } q$$

It is shown in [13] proposition 4.2 that

$$(31) \quad \langle F_{[i;i+nk]}, P_k \rangle = (-q^{-1})^{nk} (-1)^{k-1} \frac{q^k - q^{-k}}{p^k - p^{-k}}$$

$$(32) \quad \langle P_{-k}, E_{[i;i+nk]} \rangle = (-1)^{k-1} \frac{q^k - q^{-k}}{p^k q^{nk} - p^{-k} q^{-nk}}$$

combined with lemma 3.20 of *loc. cit.* which simplifies the pairing between root generator and a product of several terms, we see that

$$(33) \quad \langle f_{[i;j]}, e_{i_1} \dots e_{i_m} P_{j_1} \dots P_{j_l} \rangle = \prod_{i=1}^l \frac{1}{p^k - p^{-k}} \cdot \text{a rational function of } q$$

$$(34) \quad \langle f_{i_1} \dots f_{i_m} P_{-j_1} \dots P_{-j_l}, e_{[i;j]} \rangle = \prod_{i=1}^l \frac{1}{p^k q^{nk} - p^{-k} q^{-nk}} \cdot \text{a rational function of } q$$

4.2.2. To prove proposition 4.12, we first prove a weaker result

Proposition 4.13. *Let c_λ be as defined in (28). Then c_λ live in $\mathbb{Q}(q)[p^\pm]_{1-p^\#}$, where the subscript $1 - p^\#$ is defined in the same way as (29).*

Proof. Let S_+ (resp. S_-) be a subset of monomials of the form $e_{i_1} \dots e_{i_m} P_{j_1} \dots P_{j_l}$ (resp. $f_{i_1} \dots f_{i_m} P_{-j_1} \dots P_{-j_l}$) that forms a basis of $U_q^+(\widehat{\mathfrak{gl}}_n)$ (resp. $U_q^-(\widehat{\mathfrak{gl}}_n)$).

Write

$$f_{[i;j]} = \sum_{s \in S_-} b_s \cdot s$$

We claim that $b_s \in \mathbb{Q}(q)[p^\pm]$. Indeed, b_s can be computed from the pairing $\langle f_{[i;j]}, s \rangle$ for $s \in S_+$ as in (33). Compare this the pairing between S_+ and S_- in (30), we deduce that for $s = f_{i_1} \dots f_{i_m} P_{-j_1} \dots P_{-j_l}$,

$$b_s \in \prod_{i=1}^l (p^{j_i} q^{n_{j_i}} - p^{-j_i} q^{-n_{j_i}}) \cdot \mathbb{Q}(q)$$

and in particular it is an element of $\mathbb{Q}(q)[p^\pm]$.

Our next task is to write P_k in terms of $e_{[i;j]}$. Similar to the above, write

$$e_{[i;i+nk]} = \sum_{s \in S_+} a_s \cdot s$$

(S_+ must contain P_k for dimension reason.) The same argument as above shows that for $s = e_{i_1} \dots e_{i_m} P_{j_1} \dots P_{j_l}$,

$$a_s \in \prod_{i=1}^l (p^{j_i} - p^{-j_i}) \cdot \mathbb{Q}(q)$$

In particular, the coefficient for P_k is $p^k - p^{-k}$. This shows (by induction on P_i 's) that P_k can be written as a sum of products of $e_{[i;j]}$'s where the coefficients are in $\mathbb{Q}(q)[p^\pm]_{1-p^\#}$.

Combining the two paragraphs above, and note that

$$\tau(f_i) = e_i, \tau(P_{-k}) = P_k$$

we deduce that $\tau(f_{[i;j]})$ can be written as a sum of products of $e_{[i;j]}$'s where the coefficients are in $\mathbb{Q}(q)[p^\pm]_{1-p^\#}$.


Since any product of $e_{[i;j]}$'s can be written as a $\mathbb{Q}(q)$ -linear combination of PBW basis e_λ , this finishes the proof of the proposition. \square

Proof of proposition 4.12. Note that

$$c_\lambda = (\tau(f_{[i;j]}) \cdot |0\rangle, |e_\lambda\rangle^*) = (|0\rangle, f|e_\lambda\rangle^*)$$

where $|0\rangle = [\mathcal{O}_{\mathcal{M}_0}]$ is the structure sheaf in degree $\mathbf{d} = 0$. So we only need to prove that $f|e_\lambda\rangle^*$ has no factor in $\mathbb{Q}(q)$, which is equivalent to prove that the correspondence $\overline{\mathfrak{F}}_{[i;j]}$ has no tangent weight that only involves q . (Note that here we used that $|e_\lambda\rangle^*$ are integral K-theory classes, which is what we proved in the main theorem.) This in turn is equivalent to prove that the \mathbf{A} -fixed (as opposed to \mathbf{T} -fixed) points in $\overline{\mathfrak{F}}_{[i;j]}$ are isolated.

To this end, recall that $\overline{\mathfrak{F}}_{[i;j]}$ is a closed subset of $\mathcal{M}_{[i;j]}$. The \mathbf{A} -fixed points of $\mathcal{M}_{[i;j]}$ are given by the product of quivers of the form



Each X_i can be gauge fixed to have the form

$$\begin{pmatrix} 0 & & & & * \\ 1 & 0 & & & * \\ & 1 & & & * \\ & & \ddots & & \vdots \\ & & & \ddots & \vdots \\ & & & & 1 & * \end{pmatrix}$$

and Y_i are constant. Since X_i 's are nilpotent in $\overline{\mathfrak{J}}$, this forces all $*$ to be 0 and hence the fixed points are isolated. This finishes the proof of the proposition. \square

APPENDIX A. EXISTENCE OF STABLE ENVELOPE

In this section, we adapt the proof in [17] to the case of affine Laumon space $\mathcal{M}_{\mathbf{d}}$ to prove proposition 4.7.

First, we prove some basic properties that will be used later.

Proposition A.1. $\mathcal{M}_{\mathbf{d}}$ has a \mathbb{T} -equivariant cell decomposition, i.e. There is a decreasing filtration

$$\mathcal{M}_{\mathbf{d}} = X^m \supset X^{m-1} \supset \dots \supset X^1 \supset X^0 = \emptyset$$

such that each X^i is a \mathbb{T} -stable closed subvariety and $X^i \setminus X^{i-1}$ is an affine space.

Proof. For the proof of this proposition, we consider the attracting manifold with respect to the subgroup

$$(35) \quad \sigma(t) = (t^{k_1}, \dots, t^{k_n}, t^l, t^s)$$

where

$$k_1 < k_2 < \dots < k_n < k_1 - s \text{ and } l \ll k_{i+1} - k_i, l \ll k_1 - s - k_n$$

thus q^{-k} is attracting for $k > 0$. $u_j^2/u_i^2 q^k$ is attracting for $j > i$ and any k . $u_j^2/(u_i^2 p^{2m} q^k)$ is attracting for $m > 0$ and any i, j, k .

By Bialynicki-Birula theorem [1, 4] (plus an equivariant compactification as in [11]) each attracting manifold Attr_λ is an affine space. We prove that every point in $\mathcal{M}_\mathbf{d}$ is in an attracting manifold, thus they give a cell decomposition of $\mathcal{M}_\mathbf{d}$.

Given $(X, Y, A, B) \in \mathcal{M}_\mathbf{d}$, construct the partition $\lambda = (\lambda^1, \dots, \lambda^n)$ in the following way: For a given color i , let r_k denote the number of (color i) boxes in the k -th row of λ^{i-k} for $k = 0, 1, 2, \dots$ where upperscript $i - k$ is understood mod n . And we refer to these boxes as boxes in group k (of color i). Determine r_0, r_1, r_2, \dots inductively by taking r_k to be the maximum integer such that $\{v_{\square}\}_{\square \in \text{group} \leq k}$ are linearly independent. It's easy to see that the numbers determined this way form a partition λ . Writing the point (X, Y, A, B) in the standard basis of U_λ , by construction, all non-zero entries in X_i corresponds to mapping \square_1 to \square_2 where either \square_1 is right to \square_2 in the same row (hence this entry has weight q^{-N} for some N), or \square_1 is lower than \square_2 (hence by our construction the weight is attracting). A similar argument shows that non-zero entries of the Y_i 's are also attracting.

□

Apply Bialynicki-Birula theorem to the subtours inverse to (35), we also have

Lemma A.2. *For each λ , there is an affine space containing λ in $\mathcal{M}_\mathbf{d}$ where the action of \mathbb{T} are the repelling weights at λ .*

For an indivisible character w , we will denote the subspace spanned by the directions whose weight is a multiple of w by $C_{\lambda, w}$.

Lemma A.3. *Let A' be a subtorus of A . Let $F \subset \mathcal{M}_\mathbf{d}$ be a fixed component of A' . Then*

(1) *For any fixed point $\lambda, \mu \in F$,*

$$\deg_{A'} \mathcal{V}_\lambda^{mov} = \deg_{A'} \mathcal{V}_\mu^{mov}$$

(2) *For any class $\gamma \in K_{\mathbb{T}}(F)$, $\deg_{A'} \gamma$ is contained in the union of all $\deg_{A'} \gamma|_\lambda$ where λ runs over all A fixed points contained in F .*

Proof. Part (1): Decompose $\mathcal{V}|_F$ according to A' weights. For a subbundle \mathcal{W} with weight w , if $w \neq 0$, then it contributes to the Newton polygon by sweeping it with $w \cdot \text{rk} \mathcal{W}$, same for λ and μ . If $w = 0$, either it's not in the moving part or it has a non-trivial \mathbb{T}/A' weight, it won't change the Newton polygon.

Part (2): this follows from the fact that restriction to fixed points is injective for $K_{\mathbb{T}}(F)$, which in turn follows from the cell decomposition of F induced from lemma A.1. □

Proof of proposition 4.7. Consider all fixed points smaller or equal to λ : $\lambda = \lambda_1 \geq \lambda_2 \geq \dots$ and let F_i be the component in $(\mathcal{M}_\mathbf{d})^A$ containing λ_i . The attracting manifold of each F_i is an affine bundle over F_i .

Let

$$X_k = \mathcal{M}_{\mathbf{d}} \setminus \bigcup_{i>k} \text{Attr}_{F_i}, i = 0, 1, 2, \dots$$

Thus $X_0 \subset X_1 \subset X_2 \subset \dots$ with $X_k \setminus X_{k-1} = \text{Attr}_{F_k}$.⁴ We will inductively construct $s_{\lambda}|_{X_k}, k = 1, 2, \dots$

For $k = 1$, let $s_{\lambda}|_{X_1} = \mathcal{O}_{\text{Attr}_{F_1}}$.

Assume the induction is done for a given i . Consider the short exact sequence

$$0 \rightarrow K_{\mathbb{T}}(\text{Attr}(F_i)) \rightarrow K_{\mathbb{T}}(X_{i+1}) \rightarrow K_{\mathbb{T}}(X_i) \rightarrow 0$$

(The left hand side is exact because we know the pushforward is multiplication by the repelling directions.) By induction hypothesis we have constructed $\alpha := s_{\lambda}|_{X_i}$. Let $\mu = \lambda_{i+1}$.

For any repelling direction w in T_{λ} , let A' be the codimension 1 subtours

$$1 \rightarrow A' \rightarrow A \xrightarrow{a^w} \mathbb{C}^* \rightarrow 1$$

$C_{\lambda,w}$ is fixed under A' , and the fixed component of A' containing $C_{\lambda,w}$ contains at least one fixed point of A (again due to the cell decomposition). The lemma above implies that

$$\deg_{A'} \alpha|_{C_{\mu,w} \setminus \mu} + \text{weight}_{A'} L|_{\lambda} \subset \deg_{A'} \Lambda^{\bullet} (\mathcal{Y}_{\lambda}^{mov})^{\vee} + \text{weight}_{A'} L|_{\mu}$$

Let $\mathcal{R} = K_{\mathbb{T}/A}(\text{pt}), \mathcal{R}[A] := \mathcal{R}[u_1^{\pm}, \dots, u_n^{\pm}, p^{\pm}] = K_{\mathbb{T}}(\text{pt})$. Let

$$(36) \quad P = \Lambda^{\bullet} T_{\mu}^{mov} = \prod_{\text{indivisible } \nu} P_{\nu}(a^{\nu})$$

Let Δ be the Newton polygon of P . Let

$$s = \text{weight}(L|_{\mu}) - \text{weight}(L|_{\lambda}) \in \text{char}(A) \otimes \mathbb{Q}$$

(s was denoted λ in [17].) Since the choice of lifting only affects the value at fixed point μ , the induction step reduces to the following local statement:

Proposition A.4. *Let P be as above, let $\alpha \in \mathcal{R}[A]/P$ satisfy the condition that for any ν appeared in equation (36), there exists $\alpha_{\nu} \in \mathcal{R}[A]$ such that*

$$(37) \quad \deg_{A'} \alpha_{\nu} \subset \deg_{A'} P + s$$

and $\pi(\alpha_{\nu}) = \alpha$ where $\pi : \mathcal{R}[A] \rightarrow \mathcal{R}[A]/P$ is the natural projection, Then there exists $\alpha' \in \mathcal{R}[A]$ such that

$$\deg_A \alpha' \subset \deg_A P + s$$

and $\pi(\alpha') = \alpha$.

Proof of proposition A.4. As in [17], the degree constraint in proposition 4.7 is translated into being a section of a certain line bundle on a toric variety.

Let

$$\Delta_s = \Delta + s$$

⁴For affine Laumon space, $X_0 = \emptyset$. But the proof should work in more general situations.

Δ_s defines a toric variety \overline{A} together with a line bundle $\mathcal{O}(\Delta_s)$ on it by

$$(\overline{A}, \mathcal{O}(\Delta_s)) = \text{Proj} \bigoplus_{n \geq 0} \mathcal{P}_{n\Delta_s, A}$$

It follows immediately from the construction that

Lemma A.5. $H^0(\overline{A}, \Delta_s)$ can be identified with \mathcal{R} -linear combinations of lattice points in Δ_s .

Note that the toric variety defined by Δ and by Δ_s are the same (but the line bundles are different.)

Consider the closed subvariety defined by $P = 0$ inside A , defined by viewing P as an element in degree 1 in the defining graded algebra. We have

Lemma A.6. For an element α satisfying the condition of proposition A.4, there exists $\tilde{\alpha} \in H^0(\overline{A}, \mathcal{O}(\Delta_s)|_{P=0})$ such that $\tilde{\alpha}|_A = \alpha$.

Proof. Consider any affine chart $U \subset \overline{A}$ and let $\mathcal{R}[U]$ denote the \mathcal{R} -linear combination of lattice points in U . We need to prove that the element $\alpha \in \mathcal{R}[A]/P$ can be extended to $\mathcal{R}[U]/P$. (The lifts on different affine charts agree because they agree on the open subset A .)

To this end, we need to prove the statement that given a Laurent polynomial P and a vertex of its Newton polytope Δ , let α be a Laurent polynomial which satisfy condition (37) where ν runs through all edges at this vertex, then we can add a multiple of P to α so that its degree lies in the cone at this vertex. We prove this statement by induction on the dimension of the polytope.

For each wall of the cone of U , induction hypothesis (or condition (37) in the base case) implies that we can add a multiple of P to make all coefficient lie on the correct side of the wall. In addition, doing this for a wall won't affect walls that are already "cleared". \square

Now consider the short exact sequence

$$0 \rightarrow \mathcal{O}(\Delta)^{-1} \rightarrow \mathcal{O} \rightarrow \mathcal{O}|_{P=0} \rightarrow 0$$

tensoring with $\mathcal{O}(\Delta_s)$ we get

$$0 \rightarrow \mathcal{O}(\Delta)^{-1} \otimes \mathcal{O}(\Delta_s) \rightarrow \mathcal{O}(\Delta_s) \rightarrow \mathcal{O}(\Delta_s)|_{P=0} \rightarrow 0$$

As was shown in [17], the cohomology of the first term vanish, thus $\tilde{\alpha} \in H^0(\overline{A}, \mathcal{O}(\Delta_s)|_{P=0})$ lifts to a class $\tilde{\alpha}' \in H^0(\overline{A}, \mathcal{O}(\Delta_s))$. By lemma A.5, $\tilde{\alpha}'$ corresponds to a class $\alpha' \in \mathcal{R}[A]$ which lifts α . \square

The above proposition finishes the inductive step and thus the proof of theorem 4.7 \square

REFERENCES

- [1] Andrzej Białynicki-Birula, *Some theorems on actions of algebraic groups*, Annals of mathematics **98** (1973), no. 3, 480–497. ↑23
- [2] Alexander Braverman and Michael Finkelberg, *Finite difference quantum Toda lattice via equivariant k -theory*, Transformation Groups **10** (2005), 363–386. ↑2, 8, 9
- [3] Mathew Bullimore, Tudor Dimofte, Davide Gaiotto, Justin Hilburn, and Hee-Cheol Kim, *Vortices and vermas*, arXiv preprint arXiv:1609.04406 (2016). ↑2
- [4] Neil Chriss and Victor Ginzburg, *Representation theory and complex geometry*, Vol. 42, Springer, 1997. ↑23
- [5] Boris Feigin, Michael Finkelberg, Igor Frenkel, and Leonid Rybnikov, *Gelfand–tsetlin algebras and cohomology rings of laumon spaces*, Selecta Mathematica **17** (2011), 337–361. ↑2, 11
- [6] Boris Feigin, Michael Finkelberg, Andrei Negut, and Leonid Rybnikov, *Yangians and cohomology rings of laumon spaces*, Selecta Mathematica **17** (2011), no. 3, 573–607. ↑2
- [7] Michael Finkelberg and Alexander Kuznetsov, *Parabolic sheaves on surfaces and affine lie algebra* (2000). ↑10
- [8] Michael Finkelberg and Alexander Tsymbaliuk, *Shifted quantum affine algebras: integral forms in type a* , Arnold Mathematical Journal **5** (2019), no. 2-3, 197–283. ↑14
- [9] Justin Hilburn, Joel Kamnitzer, and Alex Weekes, *Bfn springer theory*, Communications in Mathematical Physics (2023), 1–68. ↑2
- [10] Gérard Laumon, *Un analogue global du cône nilpotent* (1988). ↑10
- [11] Daves Maulik and Andrei Okounkov, *Quantum groups and quantum cohomology*, arXiv preprint arXiv:1211.1287 (2012). ↑3, 15, 23
- [12] Hiraku Nakajima, *Handsaw quiver varieties and finite w -algebras*, arXiv preprint arXiv:1107.5073 (2011). ↑2, 10
- [13] Andrei Neguț, *Quantum toroidal and shuffle algebras*, arXiv preprint arXiv:1302.6202 (2013). ↑6, 20
- [14] ———, *Affine laumon spaces and a conjecture of kuznetsov*, arXiv preprint arXiv:1811.01011 (2018). ↑2, 3, 4, 5, 8, 9, 10
- [15] ———, *The pbw basis of $U_{q,p}(\mathfrak{gl}_n)$* , arXiv preprint arXiv:1905.06277 (2019). ↑20
- [16] Andrei Okounkov, *Lectures on k -theoretic computations in enumerative geometry*, arXiv preprint arXiv:1512.07363 (2015). ↑3, 16
- [17] ———, *Inductive construction of stable envelopes*, Letters in Mathematical Physics **111** (2021), 1–56. ↑3, 16, 22, 24, 25
- [18] Aleksander Tsymbaliuk, *Quantum affine gelfand–tsetlin bases and quantum toroidal algebra via k -theory of affine laumon spaces*, Selecta Mathematica **16** (2010), no. 2, 173–200. ↑2, 8