

Fixed-sparsity matrix approximation from matrix-vector products

Noah Amsel
New York University
noah.amsel@nyu.edu

Tyler Chen
New York University
tyler.chen@nyu.edu

Feyza Duman Keles
New York University
fd2135@nyu.edu

Diana Halikias
Cornell University
dh736@cornell.edu

Cameron Musco
UMass Amherst
cmusco@cs.umass.edu

Christopher Musco
New York University
cmusco@nyu.edu

Abstract

We study the problem of approximating a matrix \mathbf{A} with a matrix that has a fixed sparsity pattern (e.g., diagonal, banded, etc.), when \mathbf{A} is accessed only by matrix-vector products. We describe a simple randomized algorithm that returns an approximation with the given sparsity pattern with Frobenius-norm error at most $(1+\epsilon)$ times the best possible error. When each row of the desired sparsity pattern has at most s nonzero entries, this algorithm requires $O(s/\epsilon)$ non-adaptive matrix-vector products with \mathbf{A} . We also prove a matching lower-bound, showing that, for any sparsity pattern with $\Theta(s)$ nonzeros per row and column, any algorithm achieving $(1+\epsilon)$ approximation requires $\Omega(s/\epsilon)$ matrix-vector products in the worst case. We thus resolve the matrix-vector product query complexity of the problem up to constant factors, even for the well-studied case of diagonal approximation, for which no previous lower bounds were known.

1 Introduction

Learning about a matrix \mathbf{A} from matrix-vector product (matvec) queries¹ $\mathbf{x}_1, \dots, \mathbf{x}_m \mapsto \mathbf{A}\mathbf{x}_1, \dots, \mathbf{A}\mathbf{x}_m$ is a widespread task in numerical linear algebra, theoretical computer science, and machine learning. Algorithms that only access \mathbf{A} through matvec queries (also known as *matrix-free* algorithms) are useful when \mathbf{A} is not known explicitly, but admits efficient matvecs. Common settings include when \mathbf{A} is a solution operator for a linear partial differential equation [Kar+21; SO21; BET22], a function of another matrix that can be applied with Krylov subspace methods [GS92; Hig08], the Hessian of a neural network that can be applied via back-propagation [Pea94], or a data matrix in compressed sensing applications [KK11]. Matrix-free algorithms are often the methods of choice due to practical considerations such as memory usage and data movement. If the queries are chosen *non-adaptively* — i.e., the i -th query vector \mathbf{x}_i does not depend on the results $\mathbf{A}\mathbf{x}_1, \dots, \mathbf{A}\mathbf{x}_{i-1}$

¹In some settings, it may also be possible to do matrix-transpose-vector queries $\mathbf{y} \mapsto \mathbf{A}^\top \mathbf{y}$. It is an open question to understand when matrix-transpose-vector queries may be beneficial [BHOT24].

from previous queries — queries can be parallelized, potentially leading to substantial runtime improvements [Mur+23]. Non-adaptive queries can also be computed in a single pass over \mathbf{A} , and thus, non-adaptive matvec query algorithms are an important tool in streaming and distributed computation [CW09; MT20].

In many cases, the cost of matrix-free algorithms is dominated by the cost of the matvec queries. As such, a key goal is to understand the minimum number of queries required to solve a given problem, also known as the query complexity. Any algorithm automatically yields an upper bound on the query complexity, whereas it can be more challenging to prove lower bounds. Problems for which the matvec query complexity have been extensively studied include low-rank approximation [HMT11; SER18; TW23; BCW22; BN23; HT23], spectrum approximation [Ski89; WWAF06; SW23], trace and diagonal estimation [Gir87; Hut89; BKS07; MMMW21], the approximation of linear system solutions and the action of matrix functions [GS92; Gre97; BHSW20], and matrix property testing [SWYZ21; NSW22].

An important class of problems considers recovering \mathbf{A} from matvec queries when \mathbf{A} is structured, or relatedly, finding the nearest approximation to \mathbf{A} within a structured class of matrices. Example classes that have been studied extensively in this setting include low-rank matrices [HMT11; TW23], hierarchical low-rank matrices [LLY11; Mar16; LM22a; LM22b; SO21; HT23], diagonal matrices [BKS07; TS11; BN22; DM23], sparse matrices [CPR74; CM83; CC86; WEV13; DSBN15], and beyond [WSB11; SKO21].

1.1 Fixed-sparsity matrix approximation

In this work, we focus on the task of approximating \mathbf{A} with a matrix of a *specified* sparsity pattern, with error competitive with the best approximation of the given sparsity pattern. This is a natural task; indeed, there are many existing linear algebra algorithms for matrices of a given sparsity pattern (e.g. diagonal, tridiagonal, banded, block diagonal, etc.), so it is a common goal to obtain an approximation compatible with such algorithms. As we discuss in Section 1.3, several important special cases including diagonal approximation and exact recovery of matrices with known-sparsity have been studied extensively in prior work.

Formally, using “ \circ ” to indicate the Hadamard (entrywise) product and $\|\cdot\|_F$ to denote the Frobenius norm, we consider the following problem:

Problem 1 (Best approximation by a matrix of fixed sparsity). *Given a matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ and a binary matrix $\mathbf{S} \in \{0, 1\}^{n \times d}$, find a matrix $\tilde{\mathbf{A}}$ so that $\tilde{\mathbf{A}} = \mathbf{S} \circ \tilde{\mathbf{A}}$ and*

$$\|\mathbf{A} - \tilde{\mathbf{A}}\|_F \leq (1 + \varepsilon) \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_F.$$

Observe that $\mathbf{S} \circ \mathbf{A}$ is the matrix of sparsity \mathbf{S} nearest to \mathbf{A} in the Frobenius norm:

$$\mathbf{S} \circ \mathbf{A} = \operatorname{argmin}_{\mathbf{X}=\mathbf{S} \circ \mathbf{X}} \|\mathbf{A} - \mathbf{X}\|_F.$$

Hence, Problem 1 is asking for a *near-optimal* approximation to \mathbf{A} of the given sparsity \mathbf{S} .² In particular, if \mathbf{A} already has sparsity pattern \mathbf{S} , then $\mathbf{A} = \mathbf{S} \circ \mathbf{A}$, so solving Problem 1 will recover \mathbf{A} exactly.

²This is reminiscent of the low-rank approximation problem, in which we aim to find a rank- k approximation to \mathbf{A} competitive with the best rank- k approximation [HMT11; TW23].

In addressing [Problem 1](#), it will also be beneficial to consider the closely related problem of recovering the “sparse-part” of a matrix:

Problem 2 (Best approximation to on-sparsity-pattern entries). *Given a matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ and a binary matrix $\mathbf{S} \in \{0, 1\}^{n \times d}$, find a matrix $\tilde{\mathbf{A}}$ so that $\tilde{\mathbf{A}} = \mathbf{S} \circ \tilde{\mathbf{A}}$ and*

$$\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}}^2 \leq \varepsilon \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2.$$

Note the presence of the squared norms in [Problem 2](#). Since $\tilde{\mathbf{A}}$ has the same sparsity as \mathbf{S} ; i.e. $\tilde{\mathbf{A}} = \mathbf{S} \circ \tilde{\mathbf{A}}$,

$$\|\mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}}^2 = \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2 + \|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}}^2. \quad (1)$$

Thus, using the fact that $\sqrt{1+2\varepsilon} < 1+\varepsilon$ for all $\varepsilon > 0$, a solution to [Problem 2](#) with accuracy 2ε immediately yields a solution to [Problem 1](#) with accuracy ε . Conversely, if $\varepsilon \in (0, 1)$ so that $\sqrt{1+3\varepsilon} \geq 1+\varepsilon$, then a solution to [Problem 1](#) with accuracy ε yields a solution to [Problem 2](#) with accuracy 3ε . In this sense, the problems are equivalent.

1.2 Our Contributions and Roadmap

Our first contribution is to analyze a simple algorithm ([Algorithm 1](#)) that solves [Problems 1](#) and [2](#). When the sparsity pattern \mathbf{S} has at most s non-zero entries per row, this algorithm uses $m = O(s/\varepsilon)$ non-adaptive matrix-vector product queries. Specifically, the algorithm computes $\mathbf{Z} = \mathbf{A}\mathbf{G}$, where \mathbf{G} is a $d \times m$ matrix with independent standard normal entries, and then outputs the matrix

$$\tilde{\mathbf{A}} = \underset{\mathbf{X}=\mathbf{S} \circ \mathbf{X}}{\operatorname{argmin}} \|\mathbf{Z} - \mathbf{X}\mathbf{G}\|_{\mathbb{F}}. \quad (2)$$

We make no claims about the novelty of the algorithm, which follows immediately from ideas in compressed sensing. In [Section 2](#), using standard tools from random matrix theory and high dimensional probability, we provide an analysis of [Algorithm 1](#) and prove the following:

Theorem 1. *Consider any $\mathbf{A} \in \mathbb{R}^{n \times d}$ and any $\mathbf{S} \in \{0, 1\}^{n \times d}$ with at most s nonzero entries per row. Then, for any $m \geq s + 2$, using m randomized matrix-vector queries, [Algorithm 1](#) returns a matrix $\tilde{\mathbf{A}}$, equal to $\mathbf{S} \circ \mathbf{A}$ in expectation, satisfying*

$$\mathbb{E} \left[\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}}^2 \right] \leq \frac{s}{m-s-1} \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2.$$

The above inequality is equality if each row of \mathbf{S} has exactly s non-zero entries.

Owing to [\(1\)](#) and Jensen’s inequality, [Theorem 1](#) also gives an expectation bound for [Problem 1](#). Setting $m = O(s/\varepsilon)$ implies that [Problems 1](#) and [2](#) are solved in expectation. Using Markov’s inequality, we derive a probability bound:

Corollary 1. *In the setting of [Theorem 1](#), for any $\varepsilon > 0$ and $\delta \in (0, 1)$, if $m \geq s + 2$ then*

$$m \geq s \left(\frac{1}{2\delta\varepsilon} + 1 \right) + 1 \implies \mathbb{P} \left[\|\mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}} \geq (1 + \varepsilon) \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}} \right] \leq \delta.$$

Hence, using $O(s/\varepsilon)$ [Algorithm 1](#) solves [Problem 1](#) except with small constant probability, say $\leq 1/100$.

We proceed, in [Section 3](#), to study lower bounds for the matvec query complexity of [Problems 1](#) and [2](#). We show that up to constant factors, the upper bound in [Corollary 1](#) is optimal, even for the stronger class of adaptive matvec query algorithms:

Theorem 2. *Fix $\gamma \in (0, 1)$. Then there exist constants $c, C > 0$ (depending only on γ) such that the following holds:*

For any $\varepsilon \in (0, c)$ and integer $s \geq 1$, there is a distribution on (symmetric) matrices $\mathbf{A} \in \mathbb{R}^{d \times d}$ such that, for any sparsity pattern \mathbf{S} whose rows and columns each have between γs and s nonzero entries, and for any (possibly randomized) algorithm that uses $m < Cs/\varepsilon$ (possibly adaptive) matrix-vector queries to \mathbf{A} to output $\tilde{\mathbf{A}}$ with $\tilde{\mathbf{A}} \circ \mathbf{S} = \tilde{\mathbf{A}}$,

$$\mathbb{P}\left[\|\mathbf{A} - \tilde{\mathbf{A}}\|_{\text{F}} \leq (1 + \varepsilon) \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\text{F}}\right] \leq \frac{1}{25}.$$

Thus, $\Omega(s/\varepsilon)$ queries are required to solve [Problem 1](#) with any reasonable probability. Note that since solving [Problem 1](#) gives a solution to [Problem 2](#), the lower bound [Theorem 2](#) also implies that $\Omega(s/\varepsilon)$ queries are required to solve [Problem 2](#). Our approach uses an invariance property of Wishart matrices after a sequence of adaptive queries [[BHSW20](#)]. Note that since our hard instance is symmetric, [Theorem 2](#) also holds against algorithms which are allowed to use matvec queries with \mathbf{A}^{T} .

Our lower bound applies even to the well-studied case of best approximation by a diagonal matrix, and more broadly, to approximation by a banded matrix. To the best of our knowledge, our lower bounds are the first (adaptive or non-adaptive) for [Problems 1](#) and [2](#) even for these special cases.

In [Section 4](#), we compare our algorithm to widely-used coloring-based methods for fixed-sparsity pattern matrix approximation [[CPR74](#); [CM83](#); [CC86](#), etc.]. We show that there are situations where coloring methods perform worse than the algorithm described in this paper by a quadratic factor or more. Specifically, even for matrices with $\leq s$ non-zeros per row *and* column, they can require $O(s^2)$ instead of $O(s)$ matvec queries. We also discuss a setting in which coloring methods can outperform the algorithm from this paper.

Finally, in [Section 5](#) we present several numerical experiments on test problems to illustrate the sharpness of our upper bound, and in [Section 6](#) we discuss the potential for future work, including the potential for algorithms which combine the algorithm in this paper with coloring-algorithms to obtain more robustness to noise.

1.3 Past work

To the best of our knowledge, [Problems 1](#) and [2](#) have not been previously stated explicitly in the given generality. However, there is a range of past work that studies special cases of these problems. We categorize these works into the zero error case, where $\mathbf{A} = \mathbf{S} \circ \mathbf{A}$, and the nonzero error case, where $\mathbf{A} \neq \mathbf{S} \circ \mathbf{A}$. In addition, we discuss how [Problems 1](#) and [2](#) differ from the standard sparse-recovery problem in compressed sensing.

1.3.1 Zero error

Some of the earliest work relating to [Problems 1](#) and [2](#) seeks to recover Jacobian and Hessian matrices with a known sparsity pattern from matvecs [[CPR74](#); [CM83](#); [CC86](#)]. These methods make use of the fact that a *graph coloring* of a particular graph, induced by the sparsity pattern of \mathbf{A} , can be used to obtain a set of query vectors which are sufficient to exactly recover \mathbf{A} . In many (but not all) cases, exact recovery is possible with s queries. Coloring methods have also influenced many algorithms for the nonzero error setting. We compare our results to such coloring-based methods in [Section 4](#), arguing that [Algorithm 1](#) always performs better in the zero-error setting, since it always requires just s queries.

More generally, [[HT23](#)] studies the matvec query complexity of exact recovery of a wide range of linearly parameterized matrix families, proving matching upper and lower-bounds on the number of queries required. In particular, it is shown that recovering a diagonal matrix requires one query, recovering a block-diagonal matrix with $s \times s$ blocks requires s queries, and recovering a tridiagonal matrix requires 3 queries. Recovering a general $n \times d$ matrix is shown to require d queries. With probability one, [Algorithm 1](#) matches these lower bounds (see [Proposition 1](#)).

1.3.2 Nonzero error

The most theoretically well-studied instance of [Problem 2](#) is arguably the case $\mathbf{S} = \mathbf{I}$; i.e. the task of approximating the diagonal of a matrix. For this task, it is common to use Hutchinson’s diagonal estimator, defined as

$$\mathbf{d}_m = \left[\sum_{j=1}^m \mathbf{r}_j \circ (\mathbf{A}\mathbf{r}_j) \right] \diamond \left[\sum_{j=1}^m \mathbf{r}_j \circ \mathbf{r}_j \right], \quad (3)$$

where “ \diamond ” indicates entrywise division and the entries of the vectors \mathbf{r}_j are all independent random variables with mean zero and variance one. A number of analyses of this estimator have been given for various distributions [[BKS07](#); [TS11](#); [BN22](#); [HIS23](#); [DM23](#)]. In particular, [[BN22](#); [DM23](#)] give error bounds for [Problem 2](#), showing it suffices to set $m = O(1/\varepsilon)$, matching [Theorem 1](#) up to constant factors. In fact, when $\mathbf{S} = \mathbf{I}$ our [Algorithm 1](#) is equivalent to (3) if the query vectors \mathbf{r}_i are Gaussian. We detail this connection in [Remark 3](#).

Past work has also studied [Problem 1](#) with the goal of approximating a potentially non-sparse matrix by a sparse matrix. For instance, there is a long line of work on approximating matrix functions $\mathbf{A} = f(\mathbf{H})$ from matvecs. If \mathbf{H} is sparse, then the entries of $\mathbf{A} = f(\mathbf{H})$ decay exponentially away from the nonzero entries of \mathbf{H} under mild assumptions on $f(x)$ [[DMS84](#); [BR08](#); [BBR13](#)]. As such, it is reasonable to approximate \mathbf{A} with a sparse matrix of a sparsity similar to \mathbf{H} . This observation has been used in matrix approximation algorithms [[TS11](#); [SLO13](#); [FSS21](#); [PN23](#)]. Broadly speaking, these algorithms aim to combine the coloring methods described above with the estimator \mathbf{d}_m described in (3). For banded matrices, one can use the existing analyses of \mathbf{d}_m to analyze the performance of these methods, showing that they solve [Problem 2](#) to accuracy ε using $O(s/\varepsilon)$ matrix-vector queries. We include a note on this in [Section 4.1](#), as we were unable to find such an analysis in the literature. [Theorem 1](#) shows that [Algorithm 1](#) matches this bound for arbitrary sparsity patterns.

More recently, motivated by the field of partial differential equation (PDE) learning, there has been widespread interest in learning the solution operators of PDEs from input-output

data of forcing terms and solutions, analogous to matrix-vector products [SKO21; SO21; BHT23; Kar+21]. The method in [SO21] obtains a fixed-sparsity approximation to the sparse Cholesky factorization of the solution operator by coloring, which is provably accurate for certain problems. This makes use of the fact that in certain settings a fixed-sparsity Cholesky factorization is accurate and can be efficiently computed [SKO21]. This is broadly related to the (factorized) sparse approximate inverse problem for obtaining preconditioners [BT99]. The method in [BHT23] also derives a continuous analogue of a generalized coloring algorithm for targeting low-rank subblocks of hierarchical matrices [LM22b], and then recovering these subblocks using the randomized SVD [HMT11]. The final step of this algorithm reduces to the recovery of a block diagonal matrix.

1.3.3 The sparse recovery problem in compressed-sensing

It is important to contrast the aims and methods of this paper with the rich literature on compressed sensing and sparse recovery [EK12; FR13, etc.].

Given access to a length d vector \mathbf{a} through linear measurements of \mathbf{a} : $\mathbf{M} \mapsto \mathbf{M}\mathbf{a}$, the goal of the ℓ_2/ℓ_2 sparse recovery problem is to obtain an s -sparse vector $\tilde{\mathbf{a}}$ for which

$$\|\mathbf{a} - \tilde{\mathbf{a}}\|_2 \leq (1 + \varepsilon) \min_{\mathbf{a}' \text{ } s\text{-sparse}} \|\mathbf{a} - \mathbf{a}'\|_2.$$

Critically, the support of $\tilde{\mathbf{a}}$ is not known ahead of time. In fact, if the support were specified, this would be a trivial problem; simply take \mathbf{M} to have s rows, each a standard basis vector corresponding to an entry of the support.

A number of past works [WSB11; WEV13; DSBN15, etc.] have also studied a matrix version of this problem in which one aims to obtain an sd -sparse matrix $\tilde{\mathbf{A}}$ for which

$$\|\mathbf{A} - \tilde{\mathbf{A}}\|_F \leq (1 + \varepsilon) \min_{\mathbf{X} \text{ } sd\text{-sparse}} \|\mathbf{A} - \mathbf{X}\|_F,$$

using only bi-linear measurements of \mathbf{A} : $(\mathbf{U}, \mathbf{V}) \mapsto \mathbf{U}\mathbf{A}\mathbf{V}^T$. Note that the matrix problem is actually equivalent to a restricted version of the vector problem. Indeed, if $\mathbf{a} = \text{vec}(\mathbf{A})$ and $\mathbf{M} = \mathbf{U} \otimes \mathbf{V}$, then $\mathbf{M}\mathbf{a} = \text{vec}(\mathbf{U}\mathbf{A}\mathbf{V}^T)$. Here $\text{vec}(\cdot)$ forms a vector by stacking the columns of a matrix on top of one another and \otimes is the Kronecker product. In the zero error setting it has been shown that if \mathbf{A} is an $n \times n$ matrix with sufficiently distributed sparsity, one can use a convex program to stably recover \mathbf{A} using $O(\sqrt{\text{nnz}(\mathbf{A}) \cdot \log n})$ queries on each side [DSBN15].

Even for the general vector recovery problem, such algorithms necessarily have worse dependencies on ε and d than the bounds we prove for the algorithm described in the next section. In particular, algorithms solving the ℓ_2/ℓ_2 sparse recovery problem necessarily require $\Omega(s/\varepsilon^2 + s \log(ds)/\varepsilon)$ linear measurements [PW11]. In fact, even if the problem is relaxed, so that the output vector $\tilde{\mathbf{a}}$ is allowed to be non-sparse, $\Omega(s \log(d/s)/\varepsilon)$ queries are required [PW11]. In contrast, our upper-bound Theorem 1/Corollary 1 have no dependence on the dimension d .

1.4 Notation

For a set S , S^c indicates the complement (determined from context). For $d \geq 1$, we define $[d] = \{1, 2, \dots, d\}$. For $R \subset [n]$ and $C \subset [d]$, $[\mathbf{X}]_{R,C}$ indicates the $|R| \times |C|$ submatrix of a $n \times d$ matrix \mathbf{X} corresponding to the rows in R and columns in C . If R or C contain only one element, we will simply write this element. Likewise, when $R = [n]$ or $C = [d]$, we will use a colon; e.g. $[\mathbf{X}]_{1,:}$ is the first column of \mathbf{X} .

We denote the Frobenius norm of a matrix \mathbf{X} by $\|\mathbf{X}\|_F$, the transpose by \mathbf{X}^\top , and the pseudo-inverse by \mathbf{X}^\dagger . We use “ \circ ” to denote the Hadamard (entrywise) product. Specifically, for matrices \mathbf{X} and \mathbf{Y} , $\mathbf{X} \circ \mathbf{Y}$ is the matrix defined by $[\mathbf{X} \circ \mathbf{Y}]_{i,j} = [\mathbf{X}]_{i,j}[\mathbf{Y}]_{i,j}$. We use $\mathbf{0}$ and $\mathbf{1}$ to denote matrices of all zeros or ones, with size determined from context.

Throughout \mathbb{P} will be used to indicate probabilities, \mathbb{E} the expectation of a random variable, and \mathbb{V} the variance. We denote by $\mathcal{N}(\mu, \sigma^2)$ the Gaussian distribution with mean μ and variance σ^2 . We use $\text{Gaussian}(n, d)$ (or $\text{Gaussian}(d)$ if $n = d$) to denote the distribution on $n \times d$ matrices, where each entry of the matrix is independent and identically distributed (iid) with distribution $\mathcal{N}(0, 1)$.

2 An algorithm and upper bound

We begin by writing down an explicit algorithm ([Algorithm 1](#)) for solving [Problems 1](#) and [2](#). This algorithm proceeds row-by-row, taking a advantage of the fact that different rows of the solution to [\(2\)](#) do not depend on one another (except through the common use of $\mathbf{Z} = \mathbf{A}\mathbf{G}$). For each row, we can solve for the entries of $\tilde{\mathbf{A}}$ via an appropriate least squares problem.

Algorithm 1 Fixed-sparse-matrix recovery

```

1: procedure FIXED-SPARSE-MATRIX-RECOVERY( $\mathbf{A}, \mathbf{S}, m$ )
2:   Form  $\mathbf{G} \sim \text{Gaussian}(d, m)$  ▷  $d \times m$  iid Gaussian matrix
3:   Compute  $\mathbf{Z} = \mathbf{A}\mathbf{G}$  ▷  $m$  non-adaptive matvec queries
4:   for  $i = 1, 2, \dots, n$  do
5:     Let  $S_i = \{j : [\mathbf{S}]_{i,j} = 1\}$  ▷ nonzero entries of  $i$ th row of  $\mathbf{S}$ 
6:     Let  $\mathbf{z}_i^\top = [\mathbf{Z}]_{i,:}$  ▷  $i$ -th row of  $\mathbf{Z}$ 
7:     Let  $\mathbf{G}_i^\top = [\mathbf{G}]_{S_i,:}$  ▷ submatrix formed by taking the rows from  $S_i$ 
8:     Compute  $\tilde{\mathbf{a}}_i = \mathbf{G}_i^\dagger \mathbf{z}_i$  ▷ solve a  $m \times |S_i|$  least squares problem
9:     Set  $[\tilde{\mathbf{A}}]_{i,S_i} = \tilde{\mathbf{a}}_i^\top$  and  $[\tilde{\mathbf{A}}]_{i,S_i^c} = \mathbf{0}^\top$  ▷ construct  $i$ th row of  $\tilde{\mathbf{A}}$ 
10:  return  $\tilde{\mathbf{A}}$ 

```

Note that in the case that \mathbf{A} has nonzeros only in positions where \mathbf{S} is nonzero, [Algorithm 1](#) will exactly recover \mathbf{A} as long as the least squares problem for each row is fully determined.

Proposition 1. *If $\mathbf{S} \circ \mathbf{A} = \mathbf{A}$ and $m \geq s$, then [Algorithm 1](#) returns a matrix $\tilde{\mathbf{A}} = \mathbf{A}$ with probability one.*

Proof. Consider the i -th row, and let $\mathbf{x}_i \in \mathbb{R}^{|S_i|}$ be the set of non-zero entries in that row. The corresponding $\mathbf{G}_i \in \mathbb{R}^{m \times |S_i|}$ is full rank $|S_i|$ with probability one. Observe that $\mathbf{z}_i = \mathbf{G}_i \mathbf{x}_i$. Thus, since \mathbf{G}_i is full-rank, $\tilde{\mathbf{a}}_i = \mathbf{G}_i^\dagger \mathbf{z} = \mathbf{G}_i^\dagger \mathbf{G}_i \mathbf{x}_i = \mathbf{x}_i$. Thus, we recover the row exactly. By a union bound, with probability one, this simultaneously happens for all the rows. \square

Our main focus will be the case where \mathbf{A} may have nonzeros off of the specified sparsity pattern. We first recall a standard result from high dimensional probability.

Proposition 2. *Let $\mathbf{G} \sim \text{Gaussian}(p, q)$. Then, for compatible matrices \mathbf{X} and \mathbf{Y} ,*

$$\mathbb{E}[\|\mathbf{X}\mathbf{G}\mathbf{Y}\|_{\mathbb{F}}^2] = \|\mathbf{X}\|_{\mathbb{F}}^2 \|\mathbf{Y}\|_{\mathbb{F}}^2. \quad (4)$$

Moreover, if $p - q \geq 2$, then

$$\mathbb{E}[\|\mathbf{G}^\dagger\|_{\mathbb{F}}^2] = \frac{q}{p - q - 1}. \quad (5)$$

Proof. The expression (4) is an elementary calculation; see for instance [HMT11, Proposition A.1]. The expression (5) follows from the fact that $\mathbf{G}^\top \mathbf{G}$ is invertible with probability one, and $(\mathbf{G}^\top \mathbf{G})^{-1}$ has a inverse Wishart distribution, which, for $p - q \geq 2$ has mean $\mathbf{I}/(p - q - 1)$ [Mui82, §3.2 (12)]. Since $\|\mathbf{G}^\dagger\|_{\mathbb{F}}^2 = \text{tr}((\mathbf{G}^\dagger)^\top (\mathbf{G}^\dagger)) = \text{tr}((\mathbf{G}^\dagger)^\top (\mathbf{G}^\dagger))$, the result follows from the linearity of the expectation; see for instance [HMT11, Proposition A.5]. \square

Using Proposition 2, we establish the following theorem.

Theorem 1. *Consider any $\mathbf{A} \in \mathbb{R}^{n \times d}$ and any $\mathbf{S} \in \{0, 1\}^{n \times d}$ with at most s nonzero entries per row. Then, for any $m \geq s + 2$, using m randomized matrix-vector queries, Algorithm 1 returns a matrix $\tilde{\mathbf{A}}$, equal to $\mathbf{S} \circ \mathbf{A}$ in expectation, satisfying*

$$\mathbb{E}[\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}}^2] \leq \frac{s}{m - s - 1} \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2.$$

The above inequality is equality if each row of \mathbf{S} has exactly s non-zero entries.

We also obtain a probability bound:

Corollary 1. *In the setting of Theorem 1, for any $\varepsilon > 0$ and $\delta \in (0, 1)$, if $m \geq s + 2$ then*

$$m \geq s \left(\frac{1}{2\delta\varepsilon} + 1 \right) + 1 \implies \mathbb{P}[\|\mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}} \geq (1 + \varepsilon) \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}] \leq \delta.$$

Proof of Theorem 1. The algorithm processes $\mathbf{Z} = \mathbf{A}\mathbf{G} \in \mathbb{R}^{n \times m}$ sequentially to approximate the rows of \mathbf{A} . Fix i and let S_i be the indices of the nonzero entries of $[\mathbf{S}]_{i,:}$ and $\mathbf{z}_i^\top = [\mathbf{Z}]_{i,:}$ be the i -th row of \mathbf{Z} and $S_i^c = [d] \setminus S_i$. Let $\mathbf{G}_i^\top = [\mathbf{G}]_{S_i,:}$ and $\widehat{\mathbf{G}}_i^\top = [\mathbf{G}]_{S_i^c,:}$ be submatrices of \mathbf{G} formed by taking the rows of \mathbf{G} in S_i and S_i^c respectively. Define $\mathbf{x}_i^\top = [\mathbf{A}]_{i,S_i}$ and $\mathbf{y}_i^\top = [\mathbf{A}]_{i,S_i^c}$ and observe that

$$\mathbf{z}_i^\top := [\mathbf{A}\mathbf{G}]_{i,:} = \mathbf{x}_i^\top \mathbf{G}_i^\top + \mathbf{y}_i^\top \widehat{\mathbf{G}}_i^\top.$$

To enforce the sparsity pattern, [Algorithm 1](#) tries to recover $\mathbf{x}_i \in \mathbb{R}^s$ from $\mathbf{z}_i \in \mathbb{R}^m$ by solving the least squares problem:

$$\tilde{\mathbf{a}}_i := \mathbf{G}_i^\dagger \mathbf{z}_i = \mathbf{G}_i^\dagger (\mathbf{G}_i \mathbf{x}_i + \widehat{\mathbf{G}}_i \mathbf{y}_i) = \mathbf{x}_i + \mathbf{G}_i^\dagger \widehat{\mathbf{G}}_i \mathbf{y}_i.$$

Here we have used that \mathbf{G}_i is full-rank with probability one.

Since \mathbf{G}_i and $\widehat{\mathbf{G}}_i$ are independent, clearly $\mathbb{E}[\tilde{\mathbf{a}}_i] = \mathbf{x}_i$ as $\mathbb{E}[\widehat{\mathbf{G}}_i] = \mathbf{0}$. Thus, [Algorithm 1](#) outputs an unbiased estimator for $\mathbf{S} \circ \mathbf{A}$.

As long as $m \geq |S_i| + 2$, it follows from standard results in random matrix theory that

$$\begin{aligned} \mathbb{E}[\|\mathbf{x}_i - \tilde{\mathbf{a}}_i\|_2^2] &= \mathbb{E}[\|\mathbf{G}_i^\dagger \widehat{\mathbf{G}}_i \mathbf{y}_i\|_2^2] \\ &= \mathbb{E}[\mathbb{E}[\|\mathbf{G}_i^\dagger \widehat{\mathbf{G}}_i \mathbf{y}_i\|_2^2 \mid \mathbf{G}_i]] \\ &= \mathbb{E}[\|\mathbf{G}_i^\dagger\|_{\mathbb{F}}^2 \cdot \|\mathbf{y}_i\|_2^2] && \text{(4) in Proposition 2} \\ &= \frac{|S_i|}{m - |S_i| - 1} \cdot \|\mathbf{y}_i\|_2^2 && \text{(5) in Proposition 2} \\ &\leq \frac{s}{m - s - 1} \cdot \|\mathbf{y}_i\|_2^2, \end{aligned}$$

where we have used that $|S_i| \leq s$ in the final line (and hence we have equality if $|S_i| = s$).

Let $\tilde{\mathbf{A}}$ be the output of [Algorithm 1](#). Then, by the linearity of expectation,

$$\begin{aligned} \mathbb{E}[\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}}^2] &= \sum_{i=1}^n \mathbb{E}[\|\mathbf{x}_i - \tilde{\mathbf{a}}_i\|_2^2] \\ &\leq \frac{s}{m - s - 1} \sum_{i=1}^n \|\mathbf{y}_i\|_2^2 \\ &= \frac{s}{m - s - 1} \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2. \end{aligned}$$

Observe that we have equality if $|S_i| = s$ for each row $i \in [n]$. □

Proof of Corollary 1. Applying Markov's inequality to [Theorem 1](#), we find

$$\mathbb{P}[\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}}^2 \geq \alpha] \leq \frac{s}{m - s - 1} \frac{\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2}{\alpha}. \quad (6)$$

Set $\alpha = 2\varepsilon \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2$. Then, using that $\sqrt{1 + 2\varepsilon} \leq 1 + \varepsilon$ for all $\varepsilon > 0$ and recalling [\(1\)](#) gives that

$$\begin{aligned} \mathbb{P}[\|\mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}} \geq (1 + \varepsilon) \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}] &\leq \mathbb{P}[\|\mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}}^2 \geq (1 + 2\varepsilon) \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2] \\ &= \mathbb{P}[\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}}^2 \geq 2\varepsilon \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2] \\ &\leq s / ((m - s - 1)(2\varepsilon)). \end{aligned}$$

By assumption $m \geq s(1/(2\delta\varepsilon) + 1) + 1$, which gives the result. □

We now make several comments about [Algorithm 1](#) and our analysis.

Remark 1. Our bound in [Corollary 1](#) has an unfavorable $O(1/\delta)$ dependence on the failure probability δ . One could apply Markov’s inequality to each row and Hoeffding’s inequality to the sum to obtain a dependence $O(\log(n/\delta))$. However this has a dependence on the dimension n which we would like to avoid. In [Appendix C](#) we show that one can apply a high-dimensional analog of the “median trick” to obtain an algorithm with a $O(\log(1/\delta))$ failure probability (without any dependence on the dimensions n and d).

Remark 2. If \mathbf{A} and \mathbf{S} are symmetric, then it is better to return $(\tilde{\mathbf{A}} + \tilde{\mathbf{A}}^\top)/2$ than $\tilde{\mathbf{A}}$ since, by the triangle inequality,

$$\|\mathbf{A} - (\tilde{\mathbf{A}} + \tilde{\mathbf{A}}^\top)/2\|_F = \|(\mathbf{A} - \tilde{\mathbf{A}})/2 + (\mathbf{A} - \tilde{\mathbf{A}})^\top/2\|_F \leq \|\mathbf{A} - \tilde{\mathbf{A}}\|_F.$$

Remark 3. If the entries of the \mathbf{r}_i are Gaussian, then the diagonal estimator \mathbf{d}_m from [\(3\)](#) is equivalent to [\(2\)](#) with $\mathbf{S} = \mathbf{I}$. Let \mathbf{r}_j denote the j th column of \mathbf{G} . By definition, $\mathbf{z}_i = [[\mathbf{A}\mathbf{r}_1]_i, \dots, [\mathbf{A}\mathbf{r}_m]_i]^\top$ and in this case,

$$\mathbf{G}_i^\top := [\mathbf{G}]_{S_i, \cdot} = [\mathbf{G}]_{i, \cdot} = [[\mathbf{r}_1]_i, \dots, [\mathbf{r}_m]_i]$$

is a vector. The i -th row of \mathbf{d}_m is

$$[\mathbf{d}_m]_i := \frac{\sum_{j=1}^m [\mathbf{r}_j]_i \cdot [\mathbf{A}\mathbf{r}_j]_i}{\sum_{j=1}^m [\mathbf{r}_j]_i^2} = \frac{\mathbf{G}_i^\top \mathbf{z}_i}{\mathbf{G}_i^\top \mathbf{G}_i} = \mathbf{G}_i^\dagger \mathbf{z}_i.$$

In this sense, [Algorithm 1](#) for computing [\(2\)](#) is a generalization of [\(3\)](#) to non-diagonal sparsity patterns. Interestingly, however, we have not seen [\(3\)](#) interpreted in terms of a least-squares problem or pseudoinverse in the literature. This is perhaps because past work focused on diagonal estimation ([Problem 2](#)) rather than approximation by a diagonal ([Problem 1](#)).

Remark 4. [Algorithm 1](#) requires solving n least squares problems with a coefficient matrix of size $m \times s$. So, in addition to the application dependent cost of computing $\mathbf{Z} = \mathbf{A}\mathbf{G}$, its runtime is just $O(nms^2)$. There are a number of practical improvements which can be made upon implementation. First, for many sparsity patterns, the matrices \mathbf{G}_i and \mathbf{G}_{i+1} differ only by a permutation and low-rank update. Thus, by downdating/updating appropriate quantities, the cost of solving all n least-squares problems may be lower than n times the cost of solving a single system. In addition, a posteriori variance estimates could also be obtained through Jack-knife type techniques [\[ET23\]](#).

3 A lower-bound for adaptive algorithms

[Algorithm 1](#) solves [Problem 1](#) using $O(s/\varepsilon)$ matvec queries. In this section, we show that there are distributions of matrices and sparsity patterns for which no matvec query algorithm can reliably solve [Problem 1](#) using $\Omega(s/\varepsilon)$ matvecs. In particular, we show the following:

Theorem 2. *Fix $\gamma \in (0, 1)$. Then there exist constants $c, C > 0$ (depending only on γ) such that the following holds:*

For any $\varepsilon \in (0, c)$ and integer $s \geq 1$, there is a distribution on (symmetric) matrices $\mathbf{A} \in \mathbb{R}^{d \times d}$ such that, for any sparsity pattern \mathbf{S} whose rows and columns each have between γs and s nonzero entries, and for any (possibly randomized) algorithm that uses $m < Cs/\varepsilon$ (possibly adaptive) matrix-vector queries to \mathbf{A} to output $\tilde{\mathbf{A}}$ with $\tilde{\mathbf{A}} \circ \mathbf{S} = \tilde{\mathbf{A}}$,

$$\mathbb{P}\left[\|\mathbf{A} - \tilde{\mathbf{A}}\|_F \leq (1 + \varepsilon) \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_F\right] \leq \frac{1}{25}.$$

This implies that for certain hard instances of [Problems 1](#) and [2](#), our [Corollary 1](#) and thus [Theorem 1](#) are optimal up to constants. In particular, adaptivity can only improve constants; it will not lead to an improved dependence on s or ϵ . If \mathbf{A} is known to have a particular structure, it is possible that adaptive algorithms may perform better than non-adaptive algorithms for these problems.

We note that the condition $\epsilon < c$ is benign. In particular, if $C < c/2$, then $m \leq Cs/\epsilon$ implies $m \leq s/2$, in which case one cannot solve [Problem 1](#), even in the zero error case, due to a parameter counting argument.

3.1 Key technical tools

Before we prove [Theorem 2](#), we introduce several key results.

Our hard distribution will be $\mathbf{A} = \mathbf{G}^\top \mathbf{G}$, where $\mathbf{G} \sim \text{Gaussian}(d)$. This is a special case of a so-called Wishart matrix. Our lower bound will make use of the fact that the conditional distribution of a Wishart matrix after a sequence of adaptive matrix-vector queries still looks like a slightly smaller transformed Wishart matrix [[BHSW20](#), Lemma 3.4]. Similar hard input distributions have been used in a number of lower-bounds for matvec query tasks [[SER18](#); [BHSW20](#); [JPWZ21](#); [Che+23](#)]. We believe other simple distributions such as $\mathbf{A} = \mathbf{G}$ or $\mathbf{A} = \mathbf{G} + \mathbf{G}^\top$ would also suffice to prove something like [Theorem 2](#). We have chosen to use $\mathbf{A} = \mathbf{G}^\top \mathbf{G}$ because it is symmetric, and it allows us to use a conceptually intuitive anti-concentration result based on the Berry–Esseen theorem.

The following is essentially Lemma 3.4 from [[BHSW20](#)], restated to suit our needs:

Proposition 3. *Suppose $\mathbf{G} \sim \text{Gaussian}(d, r)$. Let $\mathbf{x}_1, \dots, \mathbf{x}_m$ and $\mathbf{y}_1 = \mathbf{G}^\top \mathbf{G} \mathbf{x}_1, \dots, \mathbf{y}_m = \mathbf{G}^\top \mathbf{G} \mathbf{x}_m$ be such that, for each $j = 1, \dots, m$, \mathbf{x}_j was chosen based only on the query vectors $\mathbf{x}_1, \dots, \mathbf{x}_{j-1}$ and the outputs $\mathbf{y}_1, \dots, \mathbf{y}_{j-1}$.*

Then, there is an $n \times n$ orthonormal matrix \mathbf{V}_m and an $n \times n$ matrix $\mathbf{\Delta}_m$, each constructed solely as functions of $\mathbf{x}_1, \dots, \mathbf{x}_m$ and $\mathbf{y}_1, \dots, \mathbf{y}_m$, and a matrix $\mathbf{G}_m \sim \text{Gaussian}(d-m, r-m)$ independent of $\mathbf{x}_1, \dots, \mathbf{x}_m$ and $\mathbf{y}_1, \dots, \mathbf{y}_m$ such that

$$\mathbf{V}_m^\top \mathbf{G}^\top \mathbf{G} \mathbf{V}_m = \mathbf{\Delta}_m + \begin{bmatrix} \mathbf{0}_{m,m} & \mathbf{0}_{m,d-m} \\ \mathbf{0}_{d-m,m} & \mathbf{G}_m^\top \mathbf{G}_m \end{bmatrix}.$$

We have included a proof in [Appendix B.1](#) for completeness.

We will also use the following bound about the anti-concentration of independent random variables, which we prove in [Appendix B.2](#). This is an immediate consequence of the Berry–Esseen Theorem and a basic anti-concentration result for Gaussians.

Proposition 4. *There exists a constant $C > 0$ such that, if X_1, \dots, X_k are independent random variables with $\mathbb{V}[X_i] \geq \sigma^2$, and $\mathbb{E}[|X_i - \mathbb{E}[X_i]|^3] \leq \rho$, and if we define*

$$X = X_1 + \dots + X_k,$$

then for any $t \in \mathbb{R}$ and $\alpha > 0$, if $k > C\rho^2/(\alpha^2\sigma^6)$,

$$\mathbb{P}\left[|X - t| < \alpha\sigma\sqrt{k}\right] < \alpha.$$

Using [Proposition 4](#), we can derive a more specific consequence which we will use directly in the proof of [Theorem 2](#). The proof of this result is also contained in [Appendix B.2](#).

Lemma 1. *There exists a constant $C > 0$ such that, if $\mathbf{x} = \mathbf{G}\mathbf{u}$ and $\mathbf{y} = \mathbf{G}\mathbf{v}$, where $\mathbf{G} \sim \text{Gaussian}(k)$ and $\mathbf{u}, \mathbf{v} \in \mathbb{R}^k$ such that $\|\mathbf{u}\|_2 \leq 1$ and $\|\mathbf{v}\|_2 \leq 1$, then for any $\alpha > 0$ and $t \in \mathbb{R}$, if $k > C/(\alpha^2 \|\mathbf{u}\|_2^6 \|\mathbf{v}\|_2^6)$ then*

$$\mathbb{P}\left[|\mathbf{x}^\top \mathbf{y} - t| < \alpha \|\mathbf{u}\|_2 \|\mathbf{v}\|_2 \sqrt{k}\right] < \alpha.$$

Finally, we will use the following observation about sparsity patterns that overlap all sufficiently large principal submatrices.

Lemma 2. *Let $\gamma < 1$ and suppose $\mathbf{S} \in \{0, 1\}^{d \times d}$ is binary matrix for which each row and column has between γs and s non-zero entries. Let $I \subset [d]$ with $|I| \geq 2d/(2 + \gamma)$. Then the principal submatrix $[\mathbf{S}]_{I,I}$ of \mathbf{S} contains at least $\gamma ds/(2 + \gamma)$ nonzero entries.*

Proof. Let $I \subset [d]$ with $|I| \geq 2d/(2 + \gamma)$. Note that

$$\|[\mathbf{S}]_{I,[d]}\|_{\text{F}}^2 = \sum_{i \in I} \sum_{j \in [d]} [\mathbf{S}]_{i,j} \geq |I| \cdot \gamma s = \frac{2\gamma}{2 + \gamma} ds.$$

Next, note that, since $|I^c| \leq d - 2d/(2 + \gamma) = \gamma d/(2 + \gamma)$,

$$\|[\mathbf{S}]_{I,I^c}\|_{\text{F}}^2 = \sum_{i \in I} \sum_{j \in I^c} [\mathbf{S}]_{i,j} \leq |I^c| \cdot s \leq \frac{\gamma}{2 + \gamma} ds.$$

Finally, since $[d]$ is partitioned into I and I^c ,

$$\|[\mathbf{S}]_{I,I}\|_{\text{F}}^2 = \|[\mathbf{S}]_{I,[d]}\|_{\text{F}}^2 - \|[\mathbf{S}]_{I,I^c}\|_{\text{F}}^2 \geq \frac{2\gamma}{2 + \gamma} ds - \frac{\gamma}{2 + \gamma} ds = \frac{\gamma}{2 + \gamma} ds. \quad \square$$

3.2 Proof of Theorem 2

We now have the tools necessary to prove [Theorem 2](#). The general strategy will be to show that the conditional distribution of a Wishart matrix after a sequence of (adaptive) queries is hard to approximate. That is, that the on-sparsity entries are anti-concentrated (conditioned on the queries) relative to the off-sparsity mass, which is $O(d^{3/2})$ with high probability. We will then use this result to prove [Theorem 2](#).

Lemma 3 (Main technical lemma). *Fix $\gamma \in (0, 1)$. Then there exist constants $c_1, c_2, c_3 > 0$ (depending only on γ) such that the following holds:*

Suppose $\mathbf{A} = \mathbf{G}^\top \mathbf{G}$ where $\mathbf{G} \sim \text{Gaussian}(r, d)$. Then, for any sparsity pattern \mathbf{S} whose rows and columns each have between γs and s nonzero entries, and for any (possibly randomized) algorithm that uses m (possibly adaptive) matrix-vector queries to \mathbf{A} to output $\tilde{\mathbf{A}}$ with $\tilde{\mathbf{A}} \circ \mathbf{S} = \tilde{\mathbf{A}}$, if $m \leq c_1 d$ and, for any $\alpha \in (0, 1)$, $r > m + c_3/\alpha^2$, then

$$\mathbb{P}\left[\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\text{F}}^2 < c_2 \alpha^2 ds(r - m)\right] < \alpha.$$

Proof. Fix $\gamma \in (0, 1)$. Let C denote the absolute constant in [Lemma 1](#), define

$$c_1 = \frac{\gamma}{4 + 2\gamma} \quad c_2 = \frac{c_1^2}{4} \quad c_3 = \frac{C}{c_1^6},$$

and suppose m , r , and d are integers such that

$$m \leq c_1 d, \quad r - m > \frac{c_3}{\alpha^2}.$$

Suppose we do m (possibly adaptive) queries to \mathbf{A} . [Proposition 3](#) implies that there exists an $d \times d$ matrix $\mathbf{\Delta}$, and $d \times (d - m)$ matrix \mathbf{V} with orthonormal columns, both constructed solely as functions of the queries and measurements, and a matrix $\widehat{\mathbf{G}} \sim \text{Gaussian}(r - m, d - m)$ independent of the queries and measurements such that

$$\mathbf{A} = \mathbf{\Delta} + \mathbf{V}\mathbf{W}\mathbf{V}^\top, \quad \mathbf{W} = \widehat{\mathbf{G}}^\top \widehat{\mathbf{G}}.$$

Let \mathbf{S} be any sparsity pattern for which each row and column has between γs and s non-zeros; i.e. for which we can apply [Lemma 2](#).

Let $\mathbf{T} \in \mathbb{R}^{d \times d}$ be any matrix with sparsity \mathbf{S} determined solely as a function of the queries and measurements, and hence independent of \mathbf{G} . Without loss of generality, we will absorb $\mathbf{S} \circ \mathbf{\Delta}$ into \mathbf{T} . Note also that it suffices to assume \mathbf{T} is deterministic, as the following argument holds for all possible draws of a random \mathbf{T} (and by extension, for the expectation over random draws of \mathbf{T}).

Define the set of indices

$$P = \left\{ (i, j) \in [d] \times [d] : |[\mathbf{S} \circ \mathbf{V}\mathbf{W}\mathbf{V}^\top]_{i,j} - [\mathbf{T}]_{i,j}|^2 > (\alpha/2)^2 c_1 (r - m) \right\},$$

and the event

$$E = \left\{ |P| \geq c_1 d s \right\}.$$

Note that if E holds, we have that

$$\begin{aligned} \|\mathbf{S} \circ \mathbf{A} - \mathbf{T}\|_F^2 &\geq \sum_{(i,j) \in P} |[\mathbf{S} \circ \mathbf{V}\mathbf{W}\mathbf{V}^\top]_{i,j} - [\mathbf{T}]_{i,j}|^2 \\ &\geq |P| \cdot (\alpha/2)^2 c_1 (r - m) \geq c_1 d s \cdot (\alpha/2)^2 c_1 (r - m) = \alpha^2 c_2 d s (r - m), \end{aligned}$$

so it remains to show $\mathbb{P}[E] \geq 1 - \alpha$.

Towards this end, let \mathbf{v}_i be the i th row of \mathbf{V} , and define

$$I = \{i \in [d] : \|\mathbf{v}_i\|_2^2 \geq c_1\}, \quad M = \{(i, j) \in I \times I : [\mathbf{S}]_{i,j} = 1\}.$$

We must have $|I| \geq 2d/(2 + \gamma)$. Otherwise, since \mathbf{V} has orthonormal columns so that $\|\mathbf{v}_i\|_2^2 \leq 1$, we would have:

$$d - m = \|\mathbf{V}\|_F^2 = \sum_{i=1}^d \|\mathbf{v}_i\|_2^2 < \sum_{i \in I^c} c_1 + \sum_{i \in I} 1 < c_1 d + \frac{2d}{2 + \gamma} = (1 - c_1)d,$$

which contradicts our assumption $m \leq c_1 d \Leftrightarrow d - m \geq (1 - c_1)d$. Then, since $|I| \geq 2d/(2 + \gamma)$, [Lemma 2](#) and our assumption on \mathbf{S} give that

$$|M| \geq \frac{\gamma ds}{2 + \gamma} = 2c_1 ds.$$

We now show that if $(i, j) \in M$ then $(i, j) \in P$ with high probability. Fix arbitrary $(i, j) \in M$ (and note that this means $i, j \in I$). Since $[\mathbf{S}]_{i,j} = 1$, note that $[\mathbf{S} \circ \mathbf{V}\mathbf{V}^\top]_{i,j} = \mathbf{x}_i^\top \mathbf{x}_j$, where $\mathbf{x}_i = \widehat{\mathbf{G}}\mathbf{v}_i$ and $\mathbf{x}_j = \widehat{\mathbf{G}}\mathbf{v}_j$ and \mathbf{v}_i and \mathbf{v}_j are the i -th and j -th rows of \mathbf{V} respectively. We also have that $\|\mathbf{v}_i\|_2^2, \|\mathbf{v}_j\|_2^2 \geq c_1$. With this in mind, our choice of constants implies

$$r - m \geq \frac{c_3}{\alpha^2} = \frac{C}{\alpha^2 c_1^6} \geq \frac{C}{\alpha^2 \|\mathbf{v}_i\|_2^6 \|\mathbf{v}_j\|_2^6}.$$

Hence, applying [Lemma 1](#),

$$\begin{aligned} \mathbb{P}[(i, j) \notin P] &= \mathbb{P}\left[|[\mathbf{S} \circ \mathbf{V}\mathbf{V}^\top]_{i,j} - [\mathbf{T}]_{i,j}| < (\alpha/2)c_1\sqrt{r - m}\right] \\ &\leq \mathbb{P}\left[|\mathbf{x}_i^\top \mathbf{x}_j - [\mathbf{T}]_{i,j}| < (\alpha/2)\|\mathbf{v}_i\|_2\|\mathbf{v}_j\|_2\sqrt{r - m}\right] < \frac{\alpha}{2}. \end{aligned}$$

Now, by Markov's inequality, we have that

$$\mathbb{P}\left[\sum_{(i,j) \in M} \mathbf{1}[(i, j) \notin P] \geq \frac{1}{\alpha} \cdot \frac{\alpha}{2}|M|\right] \leq \alpha.$$

Since $|M| \geq 2c_1 ds$, this then implies that

$$\mathbb{P}[E] = \mathbb{P}\left[|P| \geq c_1 ds\right] \geq \mathbb{P}\left[|P| \geq \frac{|M|}{2}\right] \geq 1 - \alpha.$$

This proves the result. \square

With [Lemma 3](#), the proof of [Theorem 2](#) is straightforward. The basic idea is that if $r = d$ and α is constant then $\|\mathbf{S} \circ \mathbf{A} - \widetilde{\mathbf{A}}\|_F^2$ is typically $\Omega(sd^2)$. However, by simple computation, it is not hard to see that $\|\mathbf{S} \circ \mathbf{A}\|_F^2 \leq \|\mathbf{A}\|_F^2$ is typically $O(d^3)$. Thus, if we set $d = O(s/\varepsilon)$, we typically will have that $\|\mathbf{S} \circ \mathbf{A} - \widetilde{\mathbf{A}}\|_F^2 > \varepsilon\|\mathbf{S} \circ \mathbf{A}\|$ as long as $m \leq c_1 d = O(s/\varepsilon)$.

Proof of Theorem 2. Fix $\gamma \in (0, 1)$ and let c_1, c_2, c_3 be the constants from [Lemma 3](#). We will make the following assignments:³

$$\alpha = \frac{1}{25}, \quad b_1 = 50, \quad b_2 = \frac{c_2 \alpha^2 (1 - c_1)}{6b_1}, \quad b_3 = \max\left\{2, \frac{(1 - c_1)b_2 \alpha^2}{c_3}\right\}, \quad C_1 = c_1 b_2.$$

Fix $\varepsilon > 0$. We will show that if

$$m \leq C_1 \frac{s}{\varepsilon}, \quad \frac{b_2}{\varepsilon} \in \mathbb{Z}, \quad \varepsilon < \frac{1}{b_3} \leq \frac{1}{2}, \tag{7}$$

³We have labeled the constants so that if c_i depends on c_j , then $i > j$.

then there is a distribution on matrices such that for any sparsity pattern with between γs and s entries per row,

$$\mathbb{P}\left[\|\mathbf{A} - \tilde{\mathbf{A}}\|_{\text{F}} \leq (1 + 2\varepsilon)\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\text{F}}\right] \leq \frac{1}{25}. \quad (8)$$

The assumption $b_2/\varepsilon \in \mathbb{Z}$ will subsequently be removed.

Towards this end, assume (7), and set

$$d = b_2 \frac{s}{\varepsilon}, \quad r = d, \quad \mathbf{A} = \mathbf{G}^{\text{T}} \mathbf{G}, \quad \mathbf{G} \sim \text{Gaussian}(r, d).$$

Define events,

$$E = \left\{ \|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\text{F}}^2 \geq c_2 \alpha^2 ds(r - m) \right\}, \quad F = \left\{ \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\text{F}}^2 \leq b_1 d^3 \right\}.$$

By assumption, $m \leq C_1 s/\varepsilon = c_1 d$ and

$$r - m \geq (1 - c_1)d = (1 - c_1) \cdot b_2 \frac{s}{\varepsilon} > (1 - c_1) b_2 \frac{1}{b_3} = \frac{c_3}{\alpha^2}.$$

Therefore, applying Lemma 3, we have that $\mathbb{P}[E] \geq 49/50$.

It is easy to show that $\mathbb{E}[\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\text{F}}^2] \leq d^3$ (see Fact 2 for a derivation), so since $b_1 = 50$, an application of Markov's inequality implies $\mathbb{P}[F] \geq 49/50$.

Note that

$$6\varepsilon b_1 d^3 \leq 6\varepsilon b_1 d \cdot b_2 \frac{s}{\varepsilon} \cdot \frac{r - m}{1 - c_1} \leq c_2 \alpha^2 ds(r - m).$$

By a union bound, both E and F hold with probability at least $24/25$. Then since $\mathbf{S} \circ \mathbf{A}$ and $\tilde{\mathbf{A}}$ have disjoint support and $\varepsilon < 1/2$, as noted in (1),

$$\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\text{F}}^2 \geq (1 + 6\varepsilon)\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\text{F}}^2 \geq (1 + 2\varepsilon)^2 \|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\text{F}}^2.$$

We now relax the assumption $c_5/\varepsilon \in \mathbb{Z}$. For arbitrary $\varepsilon > 0$, define $\tilde{\varepsilon} = b_2/\lceil b_2/\varepsilon \rceil$. Then $b_2/\tilde{\varepsilon} \in \mathbb{Z}$ and $\tilde{\varepsilon} \leq \varepsilon$. Moreover, if $\varepsilon \leq b_2/2$, then $\varepsilon \leq b_2 \tilde{\varepsilon}/(b_2 - \tilde{\varepsilon}) \leq 2\tilde{\varepsilon}$. Set $c = \min\{1/b_3, b_2/2\}$. Therefore, since $1/\varepsilon \leq 1/\tilde{\varepsilon}$, if

$$m \leq C_1 \frac{s}{\varepsilon}, \quad \varepsilon < c, \quad (9)$$

then the conditions in (7) hold (with $\tilde{\varepsilon}$), and so we have the result in (8) (with $\tilde{\varepsilon}$). Since $\varepsilon \leq 2\tilde{\varepsilon}$, this implies there is a distribution on matrices and sparsity patterns for which the result of the theorem holds.

Thus, the proof is complete with $C = 2C_1$. \square

4 Comparison with coloring methods

A number of methods for sparse-matrix and operator recovery based on *graph colorings* have been proposed [CPR74; CM83; SLO13; SO21; FSS21, etc.].⁴ To the best of our knowledge,

⁴These methods are sometimes called probing methods in the matrix function trace estimation literature.

such methods were first considered in the 1970s, and are based on the following observation: Partition column indices $[d]$ into sets $\{C_1, \dots, C_k\}$ such that the columns of \mathbf{A} corresponding to any given C_i have disjoint support. Then we can recover all of the columns in a given set C_i with a single matrix-vector product: a vector which is supported only on C_i . The sets C_i can be obtained by coloring a graph. In particular, form a graph on d vertices, where there is an edge between vertices i and j if and only if the i -th and j -th columns of \mathbf{A} have overlapping support. A k -coloring of this graph gives the partition $\{C_1, \dots, C_k\}$.

For matrices which cannot be colored with a small number of colors, one might still use coloring methods to partition the columns of the matrix, and then recover the relevant entries within each partition using an algorithm which can handle noise (e.g. [Algorithm 1](#) or Hutchinson’s diagonal estimator). In [Section 4.1](#) we analyze this approach for banded matrices. The remainder of this section provides some extreme cases to illustrate some potential pros and cons of coloring-based methods in comparison to our proposed method.

4.1 Analysis of coloring methods on banded matrices

We will describe how the estimator \mathbf{d}_m defined in [\(3\)](#) can be combined with coloring methods to solve [Problems 1](#) and [2](#). Note that if the entries of \mathbf{r}_i are chosen as independent Rademacher random variables (i.e. each entry is independently $+1$ with probability $1/2$ and -1 with probability $1/2$), then [\(3\)](#) simplifies, as $\mathbf{r}_i \circ \mathbf{r}_i$ is always the all-ones vector. It is then easy to show $\mathbb{E}[\mathbf{d}_m] = \text{diag}(\mathbf{A})$ and $\mathbb{E}[\|\text{diag}(\mathbf{A}) - \mathbf{d}_m\|_2^2] = \|\mathbf{A} - \mathbf{I} \circ \mathbf{A}\|_F^2/m$.

For any integer $b \geq 0$, let $\mathbf{S} \in \{0, 1\}^{d \times d}$ be the sparsity pattern of banded matrices of bandwidth $s = 2b + 1$; that is, $[\mathbf{S}]_{i,j} = \mathbf{1}(|i - j| \leq b)$. For convenience, we will assume $d = ks$, for some integer $k \geq 1$. This sparsity pattern yields a natural coloring-based partitioning of $[d]$ into the sets $C_i = \{i + js : j \in [k]\}$ for $i \in [s]$.

For each $i \in [s]$, define the $d \times k$ matrices

$$\mathbf{A}^{(i)} = [\mathbf{A}]_{:,C_i}, \quad \mathbf{S}^{(i)} = [\mathbf{S}]_{:,C_i}.$$

Observe that if $\mathbf{A} = \mathbf{A} \circ \mathbf{S}$, then we could recover all of the entries in $\mathbf{A}^{(i)}$ by multiplying with the all-ones vector, since there would be exactly one nonzero in each row of $\mathbf{A}^{(i)}$.

Let $\mathbf{v}^{(i)} \in \{-1, +1\}^k$ have independent Rademacher entries. Define $\mathbf{c}^{(i)} = \mathbf{S}^{(i)}\mathbf{v}^{(i)}$ and consider the vector

$$\mathbf{y}^{(i)} = \mathbf{c}^{(i)} \circ (\mathbf{A}^{(i)}\mathbf{v}^{(i)}).$$

Since the entries of $\mathbf{v}^{(i)}$ are independent, mean zero, and variance 1, a direct computation shows that

$$\mathbb{E}[\mathbf{y}^{(i)}] = (\mathbf{S}^{(i)} \circ \mathbf{A}^{(i)})\mathbf{1}, \quad \mathbb{E}[\|\mathbf{y}^{(i)} - (\mathbf{S}^{(i)} \circ \mathbf{A}^{(i)})\mathbf{1}\|_2^2] = \|\mathbf{A}^{(i)} - \mathbf{S}^{(i)} \circ \mathbf{A}^{(i)}\|_F^2.$$

Matrix-vector products with $\mathbf{A}^{(i)}$ can be computed with a single product to \mathbf{A} . Thus, using s matrix-vector products, we obtain an unbiased estimator for $\mathbf{S} \circ \mathbf{A}$ with expected squared error $\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_F^2$. Averaging t independent copies of this estimator will reduce the variance by a factor of t . Hence, using $m \geq s/\varepsilon$ matrix-vector products, one obtains an algorithm with expected squared error bounded by $\varepsilon\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_F^2$. While this algorithm is well-known in the literature [[SLO13](#); [FSS21](#), etc.], to the best of our knowledge, an analysis like the one described here has not been written down. As we discuss in the next section, this is marginally better than [Theorem 1](#).

4.2 Coloring algorithms can be better

The previous example shows that if \mathbf{S} is a banded matrix, the expected squared error of the coloring-based method described above after m matrix-vector products is better by a factor of $(m - s - 1)/m$ than [Algorithm 1](#). If s is large, m is not much larger than s , and the off-sparsity mass is large, this may be relevant. However, when m is large relative to s or if the off-sparsity mass is small, this difference is not so important.

Coloring based methods can also outperform [Algorithm 1](#) because the error of coloring methods decouples entirely between colors. Thus, to recover the entries within a given color, algorithms do not need to pay for large off-sparsity entries in a different color. An example matrix for which this observation leads to an arbitrarily large improvement over [Algorithm 1](#) is shown in the left panel of [Figure 1](#). One can also use extra queries to reduce the variance in only colors with large off-sparsity mass.

Such an improvement is clearly tied to how much of the off-sparsity mass can be ignored by the coloring scheme. If this mass is small, then coloring-based methods unnecessarily use extra queries for each color. For instance, the middle panel of [Figure 1](#) shows an example where coloring would not be so beneficial.

4.3 Coloring algorithms can be worse

In the zero-error case, it is clear that an s -sparse matrix requires at least s colors (and hence matvecs) to recover \mathbf{A} . As noted in [Proposition 1](#), [Algorithm 1](#) requires exactly s matvecs in the zero-error case, matching this lower bound. However, coloring-based methods can fail to match this lower bound, under-performing [Algorithm 1](#).

In particular, for any $s, d \geq 1$, there are s -sparse matrices with d columns which do not have a column partitioning into fewer than d partitions; i.e. for which every pair of columns has intersecting support and hence coloring-based approach do no better than the trivial algorithm which reads the matrix column-by-column, using d matvecs. A natural assumption, motivated by the banded case, is that the matrix \mathbf{A} is s -doubly sparse. That is, there are at most s nonzeros in any given row or column. For an s -doubly sparse matrix, the maximum vertex degree of the graph described above is trivially bounded by s^2 , so a greedy coloring will result in at most $s^2 + 1$ colors. The following example shows there are s -doubly sparse matrices for which $\Omega(s^2)$ colors are required. Thus, while coloring based methods would require $\Omega(s^2)$ matvec queries to recover such a matrix, [Algorithm 1](#) requires only s queries.

In particular, for any integer $k \geq 1$, define the $k^2 \times k^2$ matrix \mathbf{A} by

$$[\mathbf{A}]_{pk+i, qk+j} = \begin{cases} 1 & i = q \text{ or } j = p \\ 0 & \text{otherwise} \end{cases}, \quad p, q, i, j \in [k].$$

This sparsity pattern is represented in the left panel of [Figure 1](#). Any given row or column of this matrix has exactly $s = 2k - 1$ nonzeros. However, the support of every column overlaps. Indeed, for columns $x = pk + i$ and $y = qk + j$ (with $i, j \in [k]$),

$$[\mathbf{A}]_{ik+q, x} = [\mathbf{A}]_{jk+p, y} = 1, \quad [\mathbf{A}]_{jk+p, x} = [\mathbf{A}]_{ik+q, y} = 1.$$




Figure 1: Left: Visualization of a matrix described in Section 4.2 for which Algorithm 1 is not the best method for recovering the diagonal (intensity indicates magnitude of entries of \mathbf{A}). In particular, the diagonal of the matrix can be recovered using exactly 2 queries, while Algorithm 1 will require many queries to overcome the large noise in the off-diagonal blocks. Middle: Visualization of a matrix for which using the same colorings as the matrix on the left panel will not help. Right: Visualization of the hard sparsity pattern described in Section 4.3 with $k = 10$. Here black pixels correspond to one and white pixels to zero. Note that while each row and column of the matrix has only $O(k)$ nonzeros, each pair of the k^2 columns has overlapping support.

Therefore, exact coloring based approaches require $d = k^2 = O(s^2)$ colors; i.e. they do no better than the trivial upper bound of d queries. In contrast, Algorithm 1 would require only s queries.

5 Numerical Experiments

In this section we provide several numerical experiments which illustrate the performance of Algorithm 1. These problems are modeled after similar problems from the literature. Code to reproduce the figures can be found at https://github.com/tchen-research/fixed_sparsity_matrix_approximation.

5.1 Model problem

We consider the matrix $\mathbf{A} = \mathbf{M}^{-1}$, where $\mathbf{M} = \text{tridiag}(-1, 4, -1)$. This class of matrices exhibits exponential decay away from the diagonal and was used in experiments in past work [BS15; FSS21].

We take \mathbf{A} to be 1000×1000 and, for varying values of $b \geq 0$, we set \mathbf{S} to be a symmetric banded matrix of maximum total bandwidth $2b + 1$; i.e. $[\mathbf{S}]_{i,j} = \mathbf{1}(|i - j| \leq b)$. We then compute the approximation error $\|\mathbf{A} - \tilde{\mathbf{A}}\|_F$ and recovery error $\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_F$ for the output of Algorithm 1 run using a varying number of matvec queries m .

The results are illustrated in Figure 2. Here the convergence of Algorithm 1 is matched well by the upper bound in Theorem 1 for the expected squared error (note that the plot shows the error, not the expected squared error).




Figure 2: Approximation of model problem matrix $\mathbf{A} = \text{tridiag}(-1, 4, -1)^{-1}$ by a matrix of total bandwidth s for varying values of s . The solid circles indicate the root mean squared error of Algorithm 1 over 20 independent runs of the algorithm, and the shaded region indicates the 10%-90% range. The dotted lines are the $\sqrt{s/(m - s - 1)}\|\mathbf{A} - \mathbf{S} \circ \tilde{\mathbf{A}}\|_F$ (left) and $\sqrt{1 + s/(m - s - 1)}\|\mathbf{A} - \mathbf{S} \circ \tilde{\mathbf{A}}\|_F$ (right).

5.2 Trefethen Primes

We let $\mathbf{A} = \mathbf{M}^{-1}$, where \mathbf{M} be the 1000×1000 matrix whose entries are zero everywhere except for the primes $2, 3, 5, 7, \dots, 7919$ along the main diagonal and the number 1 in all the positions $[\mathbf{B}]_{i,j}$ with $|i - j| \in \{1, 2, 4, 8, \dots, 512\}$.⁵ An example (somewhat different from our example) involving this matrix was used in [PN23].




Figure 3: Left: Log-scale of the nonzero entries of \mathbf{M} , which range in magnitude from 1 to 7919. Middle: Log-scale of the nonzero entries of \mathbf{A} . Right: Sample sparsity pattern \mathbf{S} corresponding to $b = 5$.

For $b > 0$, we defined a sparsity pattern \mathbf{S} to be such that, for each $t \in \{1, 2, 4, 8, \dots, 512\}$, $[\mathbf{S}]_{i,j} = 1$ whenever $|i - j \pm t| \leq b$ and zero otherwise. In other words, the sparsity pattern consists of bandwidth $2b + 1$ bands centered nonzero entries of $\mathbf{M} = \mathbf{A}^{-1}$. This is illustrated in Figure 3.

The results are shown in Figure 4. Here, the convergence of Algorithm 1 is somewhat better than the upper bound Theorem 1. This is because many rows of the sparsity pattern have far fewer than s entries. From the proof of Theorem 1, it is clear how to obtain an exact

⁵In problem 7 of the “A Hundred-dollar, Hundred-digit Challenge” in SIAM News, readers are asked to compute the (1,1) entry of a larger but analogously defined matrix \mathbf{A} to 100 digits of accuracy [Tre02].

characterization of the expected squared error.




Figure 4: Approximation of “Trefethen primes” inverse matrix by a multi-banded matrix for varying values of s . The solid circles indicate the root mean squared error of Algorithm 1 over 100 independent runs of the algorithm, and the shaded region indicates the 10%-90% range. The dotted lines are the $\sqrt{s/(m-s-1)}\|\mathbf{A}-\mathbf{S}\circ\tilde{\mathbf{A}}\|_F$ (left) and $\sqrt{1+s/(m-s-1)}\|\mathbf{A}-\mathbf{S}\circ\tilde{\mathbf{A}}\|_F$ (right).

6 Outlook

This work raises a number of interesting practical and theoretical questions. We now comment on three.

It is clear that there is potential for combining coloring methods with algorithms such as Algorithm 1 in order to avoid paying for large off-sparsity entries in parts of a matrix while simultaneously maintaining the robustness to noise enjoyed by Algorithm 1. One approach to combining these two paradigms is to use adaptive queries to identify portions of the matrix with large-mass, and then find a coloring based on this information. We believe further study in this direction may yield algorithms which work better in many practical situations. Of course, our lower bound Theorem 2 shows that only constant factors can be improved for some families of problem instances.

The present paper focuses only on the Frobenius norm. It would be valuable to understand the analogous problem in other norms such as the matrix 2-norm. For other norms, $\mathbf{S}\circ\mathbf{A}$ is not necessarily the best approximation to \mathbf{A} with sparsity \mathbf{S} . For instance, if $\mathbf{A} = [1, 1; 1, 1]$ and $\mathbf{S} = [1, 0; 0, 0]$, then

$$\operatorname{argmin}_{\mathbf{X}=\mathbf{S}\circ\mathbf{X}}\|\mathbf{A}-\mathbf{X}\|_2 = \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix} \neq \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \mathbf{S}\circ\mathbf{A}.$$

Loosely speaking, minimizing the operator-norm approximation error requires the rows of the error matrix to be small and unaligned, whereas the Frobenius norm problem only requires them to be small.

Finally, it is of broad interest to understand algorithms and lower bounds for richer structures of matrices, such as the sum of sparse and low-rank matrices and matrices with hierarchical low-rank structure. This paper is a starting point for investigating these problems, and we hope that future work will explore them in greater depth.

Acknowledgements

Diana Halikias was supported by the Office of Naval Research (ONR) under grant N00014-23-1-2729. Cameron Musco was partially supported by NSF Grant 2046235. Christopher Musco was partially supported by NSF Grant 2045590.

We thank Robert J. Webber for informing us of the operator-norm example in [Section 6](#).

A Discussion on relative error recovery

Instead of [Problem 2](#), one may hope for a guarantee like

$$\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}} \leq \varepsilon \|\mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}. \quad (10)$$

Unfortunately, we cannot expect to be able to obtain such an approximation using fewer than d queries in general; for $\varepsilon < 1$, this would require differentiating matrices for which $\mathbf{S} \circ \mathbf{A}$ is zero from those for which it is arbitrarily small.

Even so, in many situations where one might hope to recover the sparse part of a matrix, $\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}$ can be bounded by a constant multiple of $\|\mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}$. In such cases, [Theorem 1](#) implies that [Algorithm 1](#) returns an approximation satisfying (10) using $O(s/\varepsilon^2)$ queries. One instance where this can be expected is if $\mathbf{A} = f(\mathbf{M})$, where \mathbf{M} has sparsity \mathbf{S} . It is known that matrix functions of sparse matrices have entries which decay exponentially away from the sparsity pattern [\[BR08\]](#). Another instance in which this holds is if \mathbf{S} contains the identity and \mathbf{A} is diagonally dominant; i.e. if $|\mathbf{A}_{i,i}| \geq \sum_{j \neq i} |\mathbf{A}_{i,j}|$. Indeed, in this case

$$\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2 \leq \|\mathbf{A} - \mathbf{I} \circ \mathbf{A}\|_{\mathbb{F}}^2 = \sum_{i=1}^d \sum_{\substack{j=1 \\ j \neq i}}^d |\mathbf{A}_{i,j}|^2 \leq \sum_{i=1}^d |\mathbf{A}_{i,i}|^2 = \|\mathbf{I} \circ \mathbf{A}\|_{\mathbb{F}}^2 \leq \|\mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2.$$

Using [Lemma 3](#), we can derive lower-bounds. For instance, if in the proof of [Theorem 2](#) we replace F by

$$F = \left\{ \|\mathbf{I} \circ \mathbf{A}\|_{\mathbb{F}}^2 \leq b_1 d^3 \right\}$$

and b_1 by 150, then by [Fact 2](#), the proof of [Theorem 2](#) can be repeated and results in a bound $\Omega(1/\varepsilon^2)$ queries for relative error diagonal approximation of PSD matrices.

In fact, if we set r as large constant multiple of d , then \mathbf{A} becomes diagonally dominant with high probability, and the same lower bound still holds. In light of the above upper bound, [Algorithm 1](#) is optimal for relative error diagonal approximation on diagonally dominant matrices.

B Lower Bound Lemmas

In this section, we provide the proofs of the key lemmas used to prove the lower bounds stated in [Section 3](#). These lemmas are all standard in the literature, but we include the proofs

for the benefit of the reader, as they are, for the most part, self-contained and interesting.

B.1 Adaptive queries to Wishart matrices

In this section we provide a proof of [Proposition 3](#), which is essentially [[BHSW20](#), Lemma 3.4]. This is mostly included for completeness. First, we consider what happens after a single non-adaptive query.

Lemma 4. *Suppose $\mathbf{G} \sim \text{Gaussian}(d, r)$. Let \mathbf{x} be a unit-length query chosen independently of \mathbf{G} , and define $\mathbf{y} = \mathbf{G}^\top \mathbf{G} \mathbf{x}$. Then, there is an $n \times n$ orthonormal matrix $\hat{\mathbf{X}} = [\mathbf{x} \ \mathbf{X}]$, constructed solely as a function of \mathbf{x} , and a matrix $\mathbf{H} \sim \text{Gaussian}(d-1, r-1)$ independent of \mathbf{x} and \mathbf{y} such that*

$$\hat{\mathbf{X}}^\top \mathbf{G}^\top \mathbf{G} \hat{\mathbf{X}} = \begin{bmatrix} \mathbf{x}^\top \mathbf{y} & \mathbf{y}^\top \mathbf{X} \\ \mathbf{X}^\top \mathbf{y} & (\mathbf{x}^\top \mathbf{y})^{-2} \mathbf{X}^\top \mathbf{y} \mathbf{y}^\top \mathbf{X} \end{bmatrix} + \begin{bmatrix} 0 & \mathbf{0}_{1,d-1} \\ \mathbf{0}_{d-1,1} & \mathbf{H}^\top \mathbf{H} \end{bmatrix}.$$

Proof. Solely based on \mathbf{x} , extend to an orthonormal matrix:

$$\hat{\mathbf{X}} = [\mathbf{x} \ \mathbf{X}].$$

For instance, append the identity to \mathbf{x} , delete the first column which is dependent on the previous columns, then orthonormalize sequentially using Gram–Schmidt.

Since $\mathbf{y} = \mathbf{G}^\top \mathbf{G} \mathbf{x}$,

$$\hat{\mathbf{X}}^\top \mathbf{G}^\top \mathbf{G} \hat{\mathbf{X}} = \begin{bmatrix} \mathbf{x}^\top \mathbf{y} & \mathbf{y}^\top \mathbf{X} \\ \mathbf{X}^\top \mathbf{y} & \mathbf{X}^\top \mathbf{G}^\top \mathbf{G} \mathbf{X} \end{bmatrix}.$$

The first row and column of this matrix depend only on \mathbf{x} and \mathbf{y} . We will now show $\mathbf{X}^\top \mathbf{G}^\top \mathbf{G} \mathbf{X}$ is the sum of a matrix depending on \mathbf{x} and \mathbf{y} and Gaussian matrix independent of \mathbf{x} and \mathbf{y} .

Define $\mathbf{r} = \|\mathbf{G} \mathbf{x}\|^{-1} \mathbf{G} \mathbf{x}$, and note that $\|\mathbf{r}\| = 1$. Solely based on \mathbf{r} , extend to an orthonormal matrix:

$$\hat{\mathbf{R}} = [\mathbf{r} \ \mathbf{R}].$$

Since $\hat{\mathbf{R}}$ is orthonormal, $\mathbf{r} \mathbf{r}^\top + \mathbf{R} \mathbf{R}^\top = \mathbf{I}$, and

$$\begin{aligned} \mathbf{X}^\top \mathbf{G}^\top \mathbf{G} \mathbf{X} &= \mathbf{X}^\top \mathbf{G}^\top \mathbf{r} \mathbf{r}^\top \mathbf{G} \mathbf{X} + \mathbf{X}^\top \mathbf{G}^\top \mathbf{R} \mathbf{R}^\top \mathbf{G} \mathbf{X} \\ &= \|\mathbf{G} \mathbf{x}\|^{-2} \mathbf{X}^\top \mathbf{G}^\top \mathbf{G} \mathbf{x} \mathbf{x}^\top \mathbf{G}^\top \mathbf{G} \mathbf{X} + \mathbf{X}^\top \mathbf{G}^\top \mathbf{R} \mathbf{R}^\top \mathbf{G} \mathbf{X}. \end{aligned}$$

Thus, using that $\|\mathbf{G} \mathbf{x}\|^2 = \mathbf{x}^\top \mathbf{G}^\top \mathbf{G} \mathbf{x} = \mathbf{x}^\top \mathbf{y}$,

$$\hat{\mathbf{X}}^\top \mathbf{G}^\top \mathbf{G} \hat{\mathbf{X}} = \begin{bmatrix} \mathbf{x}^\top \mathbf{y} & \mathbf{y}^\top \mathbf{X} \\ \mathbf{X}^\top \mathbf{y} & (\mathbf{x}^\top \mathbf{y})^{-1} \mathbf{X}^\top \mathbf{y} \mathbf{y}^\top \mathbf{X} \end{bmatrix} + \begin{bmatrix} 0 & \mathbf{0}_{1,n-1} \\ \mathbf{0}_{n-1,1} & \mathbf{X}^\top \mathbf{G}^\top \mathbf{R} \mathbf{R}^\top \mathbf{G} \mathbf{X} \end{bmatrix}.$$

It remains to show $\mathbf{R}^\top \mathbf{G} \mathbf{X}$ is a $(r-1) \times (d-1)$ Gaussian matrix independent of \mathbf{x} and \mathbf{y} .

First, note that since \mathbf{X} is chosen only based on \mathbf{x} (which is independent of \mathbf{G}), $\mathbf{G} \hat{\mathbf{X}}$ consists of iid Gaussians. Thus, the columns of $\mathbf{G} \hat{\mathbf{X}}$ are mutually independent of one another and \mathbf{x} , and hence $\mathbf{G} \mathbf{X}$ is mutually independent of \mathbf{x} and \mathbf{y} . Finally, since \mathbf{R} depends only on $\mathbf{r} = (\mathbf{x}^\top \mathbf{y})^{-1/2} \mathbf{G} \mathbf{x}$, \mathbf{R} is independent of $\mathbf{G} \mathbf{X}$. Thus, $\mathbf{R}^\top \mathbf{G} \mathbf{X}$ has iid Gaussian entries independent of \mathbf{x} and \mathbf{y} (and hence any matrices constructed solely from \mathbf{x} and \mathbf{y}). \square

We will now prove the general statement.

Proof of Proposition 3. We proceed by induction. Suppose that after t queries, the result of the lemma holds. Let \mathbf{x}_{t+1} be a query chosen based solely on $\mathbf{x}_1, \dots, \mathbf{x}_t$ and $\mathbf{y}_1, \dots, \mathbf{y}_t$ and hence independent of \mathbf{G}_t .

Then by the inductive hypothesis,

$$\mathbf{G}^\top \mathbf{G} \mathbf{x}_{t+1} = \mathbf{V}_t \Delta_t \mathbf{V}_t^\top \mathbf{x}_{t+1} + \mathbf{V}_t^\top \begin{bmatrix} \mathbf{0}_{t,t} & \mathbf{0}_{t,d-t} \\ \mathbf{0}_{d-t,t} & \mathbf{G}_t^\top \mathbf{G}_t \end{bmatrix} \mathbf{V}_t^\top \mathbf{x}_{t+1}.$$

Let \mathbf{x} denote the bottom $d-t$ entries of $\mathbf{V}_t^\top \mathbf{x}_{t+1}$, normalized to have length 1. By Lemma 4, querying $\mathbf{G}_t^\top \mathbf{G}_t$ results in the factorization

$$\hat{\mathbf{X}}^\top \mathbf{G}_t^\top \mathbf{G}_t \hat{\mathbf{X}} = \Delta + \begin{bmatrix} 0 & \mathbf{0}_{1,d-t-1} \\ \mathbf{0}_{d-t-1,1} & \mathbf{H}^\top \mathbf{H} \end{bmatrix},$$

where Δ and $\hat{\mathbf{X}}$ are constructed solely as functions of \mathbf{x} and $\mathbf{G}_t^\top \mathbf{G}_t \mathbf{x}$ (and hence of $\mathbf{x}_1, \dots, \mathbf{x}_{t+1}$ and $\mathbf{y}_1, \dots, \mathbf{y}_{t+1}$), $\hat{\mathbf{X}}$ is orthonormal, and $\mathbf{H} \sim \text{Gaussian}(d-t-1, r-t-1)$ is independent of \mathbf{x} (and hence of $\mathbf{x}_1, \dots, \mathbf{x}_{t+1}$ and $\mathbf{y}_1, \dots, \mathbf{y}_{t+1}$).

Define $\mathbf{G}_{t+1} = \mathbf{H}$ and the matrices

$$\tilde{\mathbf{X}} = \begin{bmatrix} 1 & \mathbf{0}_{1,t-1} \\ \mathbf{0}_{t-1,t} & \hat{\mathbf{X}} \end{bmatrix}, \quad \mathbf{V}_{t+1} = \mathbf{V}_t \tilde{\mathbf{X}}, \quad \Delta_{t+1} = \tilde{\mathbf{X}}^\top \Delta_t \tilde{\mathbf{X}} + \begin{bmatrix} \mathbf{0}_{t,t} & \mathbf{0}_{t,d-t} \\ \mathbf{0}_{d-t,t} & \Delta \end{bmatrix}.$$

Clearly \mathbf{V}_{t+1} is orthonormal and \mathbf{V}_{t+1} and Δ_{t+1} are constructed solely as functions of $\mathbf{x}_1, \dots, \mathbf{x}_{t+1}$ and $\mathbf{y}_1, \dots, \mathbf{y}_{t+1}$. We easily verify that

$$\mathbf{V}_{t+1}^\top \mathbf{G}^\top \mathbf{G} \mathbf{V}_{t+1} = \tilde{\mathbf{X}}^\top \Delta_t \tilde{\mathbf{X}} + \begin{bmatrix} \mathbf{0}_{t,t} & \mathbf{0}_{t,d-t} \\ \mathbf{0}_{d-t,t} & \hat{\mathbf{X}}^\top \mathbf{G}_t^\top \mathbf{G}_t \hat{\mathbf{X}} \end{bmatrix} = \Delta_{t+1} + \begin{bmatrix} \mathbf{0}_{t+1,t+1} & \mathbf{0}_{t+1,d-(t+1)} \\ \mathbf{0}_{d-(t+1),t+1} & \mathbf{G}_{t+1}^\top \mathbf{G}_{t+1} \end{bmatrix}$$

The result is proved as the base case $t = 0$ is trivial. \square

B.2 Anti-concentration for sums

In this section we will prove Proposition 4 and Lemma 1.

We begin recalling the Berry–Esseen Theorem for non-identically distributed summands.

Proposition 5 (Berry–Esseen; see e.g. [Ser80, §1.9]). *There exists a constant $C > 0$ such that, if X_1, \dots, X_k are independent random variables with $\mathbb{E}[X_i] = 0$, $\mathbb{E}[X_i^2] = \sigma_i^2$, and $\mathbb{E}[|X_i|^3] = \rho_i$, and if we define*

$$Y = \frac{X_1 + \dots + X_k}{(\sigma_1^2 + \dots + \sigma_k^2)^{1/2}},$$

then the CDF F_Y of Y is near to the CDF Φ of a standard Gaussian in that,

$$\forall z : |F_Y(z) - \Phi(z)| \leq C \frac{\rho_1 + \dots + \rho_k}{(\sigma_1^2 + \dots + \sigma_k^2)^{3/2}}.$$

In addition, we note a simple anti-concentration bound for Gaussians.

Lemma 5. *Let $Z \sim \mathcal{N}(0, 1)$. Then, for any $t \in \mathbb{R}$ and any $\alpha > 0$,*

$$\mathbb{P}[|Z - t| < \alpha] < \sqrt{\frac{2}{\pi}} \alpha.$$

Proof. Note that the density for Z is $f_Z(x) = (1/\sqrt{2\pi}) \exp(-x^2/2)$, and $f_Z(x) \leq 1/\sqrt{2\pi}$ for all x . Hence,

$$\mathbb{P}[|Z - t| < \alpha] = \int_{t-\alpha}^{t+\alpha} f_Z(x) dx \leq \frac{2\alpha}{\sqrt{2\pi}} = \sqrt{\frac{2}{\pi}} \alpha. \quad \square$$

Proof of Proposition 4. Without loss of generality, we can assume $\mathbb{E}[X_i] = 0$ for all i by absorbing the means into t . By assumption we have that

$$\hat{\sigma} = \sqrt{\sigma_1^2 + \dots + \sigma_k^2} \geq \sigma\sqrt{k}, \quad \rho_1 + \dots + \rho_k \leq k\rho.$$

Let $Y = X/\hat{\sigma}$ with cumulative distribution F_Y and let $t' = t/\hat{\sigma}$. Then, with C denoting the constant from Proposition 5, we apply Proposition 5, Lemma 5, and the bounds above to obtain

$$\begin{aligned} \mathbb{P}[|X - t| < \alpha\sigma\sqrt{k}] &\leq \mathbb{P}[|X - t| < \alpha\hat{\sigma}] \\ &= \mathbb{P}[|Y - t'| < \alpha] \\ &= F_Y(t' + \alpha) - F_Y(t' - \alpha) \\ &\leq \Phi(t' + \alpha) - \Phi(t' - \alpha) + \frac{2Ck\rho}{k^{3/2}\sigma^3} \\ &\leq \sqrt{\frac{2}{\pi}} \alpha + \frac{2C\rho}{\sigma^3\sqrt{k}}. \end{aligned}$$

Since $(1 - \sqrt{2/\pi}) > 0.2$, the result follows by the choice $k > 100C^2\rho^2/(\alpha^2\sigma^6)$, and relabeling C . \square

Proof of Lemma 1. Let \mathbf{g}_ℓ denote the ℓ -th row of \mathbf{G} and define $x_\ell = \mathbf{u}^\top \mathbf{g}_\ell$ and $y_\ell = \mathbf{v}^\top \mathbf{g}_\ell$. In order to apply Proposition 4 to the sum

$$\mathbf{x}^\top \mathbf{y} = \sum_{\ell=1}^k x_\ell y_\ell,$$

which has independent terms since the \mathbf{g}_ℓ are independent, we must obtain upper-and lower-bounds on the variance and an upper bound on the third centered absolute moment of each term.

We will first bound the variance. By direct computation (see Fact 1 for a derivation),

$$\mathbb{V}[x_\ell y_\ell] = \mathbb{V}[\mathbf{g}_\ell^\top \mathbf{v} \mathbf{u}^\top \mathbf{g}_\ell] = \|\mathbf{v} \mathbf{u}^\top + \mathbf{u} \mathbf{v}^\top\|_{\mathbb{F}}^2 / 2 \geq \|\mathbf{u}\|_2^2 \|\mathbf{v}\|_2^2.$$

Here we have used that

$$\|\mathbf{v} \mathbf{u}^\top + \mathbf{u} \mathbf{v}^\top\|_{\mathbb{F}}^2 = 2\|\mathbf{v} \mathbf{u}^\top\|_{\mathbb{F}}^2 + 2(\mathbf{v}^\top \mathbf{u})^2 \geq 2\|\mathbf{v}\|_2^2 \|\mathbf{u}\|_2^2.$$

We now argue the third absolute moments are bounded. Note that x_ℓ and y_ℓ are both normally distributed with mean zero and variance at most 1 (since $\|\mathbf{u}\|_2 \leq 1$ and $\|\mathbf{v}\|_2 \leq 1$). Then $x_\ell y_\ell$ is sub-exponential with constant width parameter [Ver18, Lemma 2.7.7], and hence $\mathbb{E}[|x_\ell y_\ell - \mathbb{E}[x_\ell y_\ell]|^3] \leq \rho$ for some ρ .

Hence, applying Proposition 4, for any $\alpha > 0$, and provided $k > C\rho^2/(\alpha^2\|\mathbf{u}\|_2^6\|\mathbf{v}\|_2^6)$, where C is the constant from Proposition 4,

$$\mathbb{P}\left[|\mathbf{x}^\top \mathbf{y} - t| < \alpha\|\mathbf{u}\|_2\|\mathbf{v}\|_2\sqrt{k}\right] < \alpha.$$

Relabeling C gives the result. \square

B.3 Other facts

Fact 1. For a matrix $\mathbf{A} \in \mathbb{R}^{d \times d}$, if $\mathbf{g} \sim \text{Gaussian}(d, 1)$, then $\mathbb{V}[\mathbf{g}^\top \mathbf{A} \mathbf{g}] = \|\mathbf{A} + \mathbf{A}^\top\|_F^2/2$.

Proof. Since $\mathbf{g}^\top \mathbf{A} \mathbf{g} = \mathbf{g}^\top (\mathbf{A}^\top + \mathbf{A}) \mathbf{g}/2$, without loss of generality we can assume \mathbf{A} is symmetric. Let $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^\top$ be the eigendecomposition of \mathbf{A} , then $\mathbf{h} = \mathbf{U}^\top \mathbf{g} \sim \text{Gaussian}(d, 1)$ and $\mathbf{h}^\top \mathbf{\Lambda} \mathbf{h}$ is a linear-combination of independent Chi-squared random variables with one degree of freedom (and variance 2). The result follows since $\|\mathbf{A}\|_F^2 = \|\mathbf{\Lambda}\|_F^2$. \square

Fact 2. For $r, d \geq 1$, suppose $\mathbf{G} \sim \text{Gaussian}(r, d)$. Then

$$\mathbb{E}[\|\mathbf{I} \circ \mathbf{G}^\top \mathbf{G}\|_F^2] = d(2r + r^2), \quad \mathbb{E}[\|\mathbf{G}^\top \mathbf{G} - \mathbf{I} \circ \mathbf{G}^\top \mathbf{G}\|_F^2] = (d^2 - d)r.$$

Proof. Write $\mathbf{A} = \mathbf{G}^\top \mathbf{G}$. The diagonal entries of \mathbf{A} are distributed as Chi-squared random variables with r degrees of freedom. These entries have mean r and variance $2r$. Likewise, the off-diagonal entries of \mathbf{A} are distributed as the inner product of two independent standard normal Gaussian vectors of length r . These entries therefore have mean zero and variance r . Therefore, for $i \neq j$,

$$\mathbb{E}[\mathbf{A}_{i,i}^2] = \mathbb{V}[\mathbf{A}_{i,i}] + \mathbb{E}[\mathbf{A}_{i,i}]^2 = 2r + r^2, \quad \mathbb{E}[\mathbf{A}_{i,j}^2] = \mathbb{V}[\mathbf{A}_{i,j}] = r.$$

The result follows by linearity of expectation. \square

C High probability algorithm

The bound Theorem 1 for Algorithm 1 has an unfavorable dependence $O(1/\delta)$ on the failure probability δ . We will now use a high-dimensional version of the ‘‘median trick’’ to improve the dependence on the failure probability to logarithmic.

Theorem 3. Consider any $\mathbf{A} \in \mathbb{R}^{n \times d}$ and any $\mathbf{S} \in \{0, 1\}^{n \times d}$ with at most s nonzero entries per row. For any $\varepsilon > 0$ and $\delta \in (0, 1)$, if $m \geq s + 2$ and additionally

$$m \geq s \left(\frac{90}{\varepsilon} + 1 \right) + 1, \quad r \geq 10 \log \left(\frac{1}{\delta} \right),$$

Algorithm 2 Fixed-sparse-matrix recovery (boosted)

-
- 1: **procedure** BOOSTED-FIXED-SPARSE-MATRIX RECOVERY($\mathbf{A}, \mathbf{S}, m, r$)
 - 2: Run [Algorithm 1](#) independently r times to get $\tilde{\mathbf{A}}_1, \dots, \tilde{\mathbf{A}}_r$
 - 3: $\forall i, j$: define $d_{i,j} = \|\tilde{\mathbf{A}}_i - \tilde{\mathbf{A}}_j\|_{\mathbb{F}}$
 - 4: $\forall i$: define B_i as the $\lceil r/2 \rceil$ -th smallest $d_{i,j}$ $\triangleright |\{j \in [r] : d_{i,j} \leq B_i\}| = \lceil r/2 \rceil$
 - 5: Compute $i^* = \operatorname{argmin}_i B_i$ $\triangleright \forall i : B_{i^*} \leq B_i$
 - 6: **return** $\tilde{\mathbf{A}}_{i^*}$
-

then, using $m \cdot r$ matrix-vector queries, [Algorithm 2](#) returns a matrix $\tilde{\mathbf{A}}$ satisfying:

$$\mathbb{P}[\|\mathbf{A} - \tilde{\mathbf{A}}\|_{\mathbb{F}} < (1 + \varepsilon)\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}] \geq 1 - \delta.$$

Proof. Let $\tilde{\varepsilon} = \frac{2}{9}\varepsilon$. Define the set

$$P = \{i \in [r] : \|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}_i\|_{\mathbb{F}}^2 \leq \tilde{\varepsilon}\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2\}.$$

In (6) of the proof of [Corollary 1](#), we show that

$$\mathbb{P}[\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}_i\|_{\mathbb{F}}^2 \geq \varepsilon\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2] \leq s/((m - s - 1)\tilde{\varepsilon}).$$

Hence, if $m \geq s((20/\tilde{\varepsilon}) + 1) + 1$, then $\mathbb{P}[i \in P] \geq \frac{19}{20}$. Define the event

$$E = \{|P| > \lceil r/2 \rceil\}.$$

A standard result [[AV79](#), Prop 2.4 (a)] asserts that with $q = 19/20$,

$$\sum_{k=0}^{\lceil r/2 \rceil} \binom{r}{k} q^k (1-q)^{r-k} \leq \exp\left(-\frac{rq}{2} \left(1 - \frac{1}{2q}\right)^2\right) = \exp\left(-\frac{81r}{760}\right).$$

In addition, $\delta \geq \exp(-r/10)$ by definition of r . Therefore, $\mathbb{P}[E] \geq 1 - \delta$.

We will condition on E for the remainder of the proof. By the triangle inequality, for any indices $i, j \in P$,

$$d_{i,j} = \|\tilde{\mathbf{A}}_i - \tilde{\mathbf{A}}_j\|_{\mathbb{F}} \leq \|\tilde{\mathbf{A}}_i - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}} + \|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}_j\|_{\mathbb{F}} \leq 2\sqrt{\tilde{\varepsilon}}\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}.$$

Since $|P| > \lceil r/2 \rceil$, then for each $i \in P$, there are at least $\lceil r/2 \rceil$ indices j satisfying $d_{i,j} \leq 2\sqrt{\tilde{\varepsilon}}\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}$. Thus by definition of B_i ,

$$B_i \leq 2\sqrt{\tilde{\varepsilon}}\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}.$$

By definition of B_{i^*} , there are at least $\lceil r/2 \rceil$ indices j for which

$$\|\tilde{\mathbf{A}}_{i^*} - \tilde{\mathbf{A}}_j\|_{\mathbb{F}} \leq 2\sqrt{\tilde{\varepsilon}}\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}$$

Simultaneously, $|P| > \lceil r/2 \rceil$. Since $|P| + \lceil r/2 \rceil > \lceil r/2 \rceil + \lceil r/2 \rceil = r$, the pigeonhole principle ensures there is at least one j^* for which $j^* \in P$ and

$$\|\tilde{\mathbf{A}}_{i^*} - \tilde{\mathbf{A}}_{j^*}\|_{\mathbb{F}} \leq 2\sqrt{\tilde{\varepsilon}}\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}.$$

Applying the triangle inequality, we find that

$$\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}_{i^*}\|_{\mathbb{F}} \leq \|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}_{j^*}\|_{\mathbb{F}} + \|\tilde{\mathbf{A}}_{i^*} - \tilde{\mathbf{A}}_{j^*}\|_{\mathbb{F}} \leq 3\sqrt{\tilde{\varepsilon}}\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}},$$

which implies $\|\mathbf{S} \circ \mathbf{A} - \tilde{\mathbf{A}}_{i^*}\|_{\mathbb{F}}^2 \leq 9\tilde{\varepsilon}\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2$. Adding $\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}^2$ to both sides (see (1)) and taking square roots,

$$\|\mathbf{A} - \tilde{\mathbf{A}}_{i^*}\|_{\mathbb{F}} \leq \sqrt{1 + 9\tilde{\varepsilon}}\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}} = \sqrt{1 + 2\varepsilon}\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}} \leq (1 + \varepsilon)\|\mathbf{A} - \mathbf{S} \circ \mathbf{A}\|_{\mathbb{F}}. \quad \square$$

References

- [AV79] D. Angluin and L. Valiant. “Fast Probabilistic Algorithms for Hamiltonian Circuits and Matchings”. In: *Journal of Computer and System Sciences* 18.2 (1979), pp. 155–193 (cited on page 26).
- [BBR13] M. Benzi, P. Boito, and N. Razouk. “Decay Properties of Spectral Projectors with Applications to Electronic Structure”. In: *SIAM Review* 55.1 (2013), pp. 3–64 (cited on page 5).
- [BCW22] A. Bakshi, K. L. Clarkson, and D. P. Woodruff. “Low-Rank Approximation with $1/\epsilon^{1/3}$ Matrix-Vector Products”. In: *Proceedings of the 54th Annual ACM SIGACT Symposium on Theory of Computing*. STOC 2022. Rome, Italy: Association for Computing Machinery, 2022, pp. 1130–1143 (cited on page 2).
- [BET22] N. Boullé, C. J. Earls, and A. Townsend. “Data-driven discovery of Green’s functions with human-understandable deep learning”. In: *Sci. Rep.* 12.1 (2022), p. 4824 (cited on page 1).
- [BHOT24] N. Boullé, D. Halikias, S. E. Otto, and A. Townsend. *Operator learning without the adjoint*. 2024 (cited on page 1).
- [BHSW20] M. Braverman, E. Hazan, M. Simchowitz, and B. Woodworth. “The Gradient Complexity of Linear Regression”. In: *Proceedings of Thirty Third Conference on Learning Theory*. Vol. 125. Proceedings of Machine Learning Research. PMLR, 2020, pp. 627–647 (cited on pages 2, 4, 11, 22).
- [BHT23] N. Boullé, D. Halikias, and A. Townsend. “Elliptic PDE learning is provably data-efficient”. In: *Proceedings of the National Academy of Sciences* 120.39 (2023), e2303904120 (cited on page 6).
- [BKS07] C. Bekas, E. Kokiopoulou, and Y. Saad. “An estimator for the diagonal of a matrix”. In: *Applied Numerical Mathematics* 57.11–12 (2007), pp. 1214–1229 (cited on pages 2, 5).
- [BN22] R. A. Baston and Y. Nakatsukasa. *Stochastic diagonal estimation: probabilistic bounds and an improved algorithm*. 2022 (cited on pages 2, 5).
- [BN23] A. Bakshi and S. Narayanan. “Krylov Methods are (nearly) Optimal for Low-Rank Approximation”. In: *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)*. IEEE, 2023 (cited on page 2).
- [BR08] M. Benzi and N. Razouk. “Decay bounds and $O(n)$ algorithms for approximating functions of sparse matrices”. In: *Electron. Trans. Numer. Anal.* 28 (2007-2008), pp. 16–39 (cited on pages 5, 21).

- [BS15] M. Benzi and V. Simoncini. “Decay Bounds for Functions of Hermitian Matrices with Banded or Kronecker Structure”. In: *SIAM Journal on Matrix Analysis and Applications* 36.3 (2015), pp. 1263–1282 (cited on page 18).
- [BT99] M. Benzi and M. Tuma. “A comparative study of sparse approximate inverse preconditioners”. In: *Applied Numerical Mathematics* 30.2–3 (1999), pp. 305–340 (cited on page 6).
- [CC86] T. F. Coleman and J.-Y. Cai. “The Cyclic Coloring Problem and Estimation of Sparse Hessian Matrices”. In: *SIAM Journal on Algebraic Discrete Methods* 7.2 (1986), pp. 221–235 (cited on pages 2, 4, 5).
- [Che+23] S. Chewi, J. De Dios Pont, J. Li, C. Lu, and S. Narayanan. “Query lower bounds for log-concave sampling”. In: *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)*. IEEE, 2023 (cited on page 11).
- [CM83] T. F. Coleman and J. J. Moré. “Estimation of Sparse Jacobian Matrices and Graph Coloring Blems”. In: *SIAM Journal on Numerical Analysis* 20.1 (1983), pp. 187–209 (cited on pages 2, 4, 5, 15).
- [CPR74] A. R. Curtis, M. J. D. Powell, and J. K. Reid. “On the Estimation of Sparse Jacobian Matrices”. In: *IMA Journal of Applied Mathematics* 13.1 (1974), pp. 117–119 (cited on pages 2, 4, 5, 15).
- [CW09] K. L. Clarkson and D. P. Woodruff. “Numerical linear algebra in the streaming model”. In: *Proceedings of the Forty-First Annual ACM Symposium on Theory of Computing*. 2009, pp. 205–214 (cited on page 2).
- [DM23] P. Dharangutte and C. Musco. “A Tight Analysis of Hutchinson’s Diagonal Estimator”. In: *Symposium on Simplicity in Algorithms (SOSA)*. Society for Industrial and Applied Mathematics, 2023, pp. 353–364 (cited on pages 2, 5).
- [DMS84] S. Demko, W. F. Moss, and P. W. Smith. “Decay rates for inverses of band matrices”. In: *Mathematics of Computation* 43.168 (1984), pp. 491–499 (cited on page 5).
- [DSBN15] G. Dasarathy, P. Shah, B. N. Bhaskar, and R. D. Nowak. “Sketching Sparse Matrices, Covariances, and Graphs via Tensor Products”. In: *IEEE Transactions on Information Theory* 61.3 (2015), pp. 1373–1388 (cited on pages 2, 6).
- [EK12] Y. C. Eldar and G. Kutyniok. *Compressed sensing: theory and applications*. Cambridge; New York: Cambridge University Press, 2012 (cited on page 6).
- [ET23] E. N. Epperly and J. A. Tropp. *Efficient error and variance estimation for randomized matrix computations*. 2023 (cited on page 10).
- [FR13] S. Foucart and H. Rauhut. *A Mathematical Introduction to Compressive Sensing*. Springer New York, 2013 (cited on page 6).
- [FSS21] A. Frommer, C. Schimmel, and M. Schweitzer. “Analysis of Probing Techniques for Sparse Approximation and Trace Estimation of Decaying Matrix Functions”. In: *SIAM Journal on Matrix Analysis and Applications* 42.3 (2021), pp. 1290–1318 (cited on pages 5, 15, 16, 18).
- [Gir87] D. Girard. *Un algorithme rapide pour le calcul de la trace de l’inverse d’une grande matrice*. 1987 (cited on page 2).
- [Gre97] A. Greenbaum. *Iterative Methods for Solving Linear Systems*. Philadelphia, PA, USA: Society for Industrial and Applied Mathematics, 1997 (cited on page 2).

- [GS92] E. Gallopoulos and Y. Saad. “Efficient Solution of Parabolic Equations by Krylov Approximation Methods”. In: *SIAM Journal on Scientific and Statistical Computing* 13.5 (1992), pp. 1236–1264 (cited on pages 1, 2).
- [Hig08] N. J. Higham. *Functions of Matrices*. Society for Industrial and Applied Mathematics, 2008 (cited on page 1).
- [HIS23] E. Hallman, I. C. F. Ipsen, and A. K. Saibaba. “Monte Carlo Methods for Estimating the Diagonal of a Real Symmetric Matrix”. In: *SIAM Journal on Matrix Analysis and Applications* 44.1 (2023), pp. 240–269 (cited on page 5).
- [HMT11] N. Halko, P. G. Martinsson, and J. A. Tropp. “Finding structure with randomness: probabilistic algorithms for constructing approximate matrix decompositions”. In: *SIAM Rev.* 53.2 (2011), pp. 217–288 (cited on pages 2, 6, 8).
- [HT23] D. Halikias and A. Townsend. “Structured matrix recovery from matrix-vector products”. In: *Numerical Linear Algebra with Applications* 31.1 (2023) (cited on pages 2, 5).
- [Hut89] M. Hutchinson. “A Stochastic Estimator of the Trace of the Influence Matrix for Laplacian Smoothing Splines”. In: *Communications in Statistics - Simulation and Computation* 18.3 (1989), pp. 1059–1076 (cited on page 2).
- [JPWZ21] S. Jiang, H. Pham, D. Woodruff, and R. Zhang. “Optimal Sketching for Trace Estimation”. In: *Advances in Neural Information Processing Systems*. Vol. 34. Curran Associates, Inc., 2021, pp. 23741–23753 (cited on page 11).
- [Kar+21] G. E. Karniadakis, I. G. Kevrekidis, L. Lu, P. Perdikaris, S. Wang, and L. Yang. “Physics-informed machine learning”. In: *Nat. Rev. Phys.* 3.6 (2021), pp. 422–440 (cited on pages 1, 6).
- [KK11] W. A. Kalender and W. A. Kalender. *Computed tomography: fundamentals, system technology, image quality, applications; [including 64-slice spiral CT]*. eng. 3., rev. ed. Erlangen: Publicis Publ, 2011 (cited on page 1).
- [LLY11] L. Lin, J. Lu, and L. Ying. “Fast construction of hierarchical matrix representation from matrix–vector multiplication”. In: *Journal of Computational Physics* 230.10 (2011), pp. 4071–4087 (cited on page 2).
- [LM22a] J. Levitt and P.-G. Martinsson. *Linear-Complexity Black-Box Randomized Compression of Rank-Structured Matrices*. 2022 (cited on page 2).
- [LM22b] J. Levitt and P.-G. Martinsson. *Randomized Compression of Rank-Structured Matrices Accelerated with Graph Coloring*. 2022 (cited on pages 2, 6).
- [Mar16] P.-G. Martinsson. “Compressing Rank-Structured Matrices via Randomized Sampling”. In: *SIAM Journal on Scientific Computing* 38.4 (2016), A1959–A1986 (cited on page 2).
- [MMMWW21] R. A. Meyer, C. Musco, C. Musco, and D. P. Woodruff. “Hutch++: Optimal Stochastic Trace Estimation”. In: *Symposium on Simplicity in Algorithms (SOSA)*. Society for Industrial and Applied Mathematics, 2021, pp. 142–155 (cited on page 2).
- [MT20] P.-G. Martinsson and J. A. Tropp. “Randomized numerical linear algebra: Foundations and algorithms”. In: *Acta Numerica* 29 (2020), pp. 403–572 (cited on page 2).

- [Mui82] R. J. Muirhead. *Aspects of Multivariate Statistical Theory*. Wiley, 1982 (cited on page 8).
- [Mur+23] R. Murray, J. Demmel, M. W. Mahoney, N. B. Erichson, M. Melnichenko, O. A. Malik, L. Grigori, P. Luszczek, M. Dereziński, M. E. Lopes, T. Liang, H. Luo, and J. Dongarra. *Randomized Numerical Linear Algebra : A Perspective on the Field With an Eye to Software*. 2023 (cited on page 2).
- [NSW22] D. Needell, W. Swartworth, and D. P. Woodruff. “Testing Positive Semidefiniteness Using Linear Measurements”. In: *2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS)*. IEEE, 2022 (cited on page 2).
- [Pea94] B. A. Pearlmutter. “Fast Exact Multiplication by the Hessian”. In: *Neural Computation* 6.1 (1994), pp. 147–160 (cited on page 1).
- [PN23] T. Park and Y. Nakatsukasa. *Approximating Sparse Matrices and their Functions using Matrix-vector products*. 2023 (cited on pages 5, 19).
- [PW11] E. Price and D. P. Woodruff. “(1 + eps)-Approximate Sparse Recovery”. In: *2011 IEEE 52nd Annual Symposium on Foundations of Computer Science*. IEEE, 2011 (cited on page 6).
- [SER18] M. Simchowitz, A. El Alaoui, and B. Recht. “Tight query complexity lower bounds for PCA via finite sample deformed wigner law”. In: *Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing*. STOC ’18. ACM, 2018 (cited on pages 2, 11).
- [Ser80] R. J. Serfling. *Approximation Theorems of Mathematical Statistics*. Wiley, 1980 (cited on page 23).
- [Ski89] J. Skilling. “The Eigenvalues of Mega-dimensional Matrices”. In: *Maximum Entropy and Bayesian Methods*. Springer Netherlands, 1989, pp. 455–466 (cited on page 2).
- [SKO21] F. Schäfer, M. Katzfuss, and H. Owhadi. “Sparse Cholesky Factorization by Kullback–Leibler Minimization”. In: *SIAM Journal on Scientific Computing* 43.3 (2021), A2019–A2046 (cited on pages 2, 6).
- [SLO13] A. Stathopoulos, J. Laeuchli, and K. Orginos. “Hierarchical Probing for Estimating the Trace of the Matrix Inverse on Toroidal Lattices”. In: *SIAM Journal on Scientific Computing* 35.5 (2013), S299–S322 (cited on pages 5, 15, 16).
- [SO21] F. Schäfer and H. Owhadi. *Sparse recovery of elliptic solvers from matrix-vector products*. 2021 (cited on pages 1, 2, 6, 15).
- [SW23] W. Swartworth and D. P. Woodruff. “Optimal Eigenvalue Approximation via Sketching”. In: *Proceedings of the 55th Annual ACM Symposium on Theory of Computing*. STOC ’23. ACM, 2023 (cited on page 2).
- [SWYZ21] X. Sun, D. P. Woodruff, G. Yang, and J. Zhang. “Querying a Matrix through Matrix-Vector Products”. In: *ACM Trans. Algorithms* 17.4 (2021) (cited on page 2).
- [Tre02] N. Trefethen. “A Hundred-dollar, Hundred-digit Challenge”. In: *SIAM News* 35.1 (2002) (cited on page 19).
- [TS11] J. M. Tang and Y. Saad. “A probing method for computing the diagonal of a matrix inverse”. In: *Numerical Linear Algebra with Applications* 19.3 (2011), pp. 485–501 (cited on pages 2, 5).

- [TW23] J. A. Tropp and R. J. Webber. “Randomized algorithms for low-rank matrix approximation: Design, analysis, and applications”. In: *arXiv preprint arXiv:2306.12418* (2023) (cited on page 2).
- [Ver18] R. Vershynin. *High-Dimensional Probability*. Cambridge University Press, 2018 (cited on page 25).
- [WEV13] T. Wimalajeewa, Y. C. Eldar, and P. K. Varshney. *Recovery of Sparse Matrices via Matrix Sketching*. 2013 (cited on pages 2, 6).
- [WSB11] A. Waters, A. Sankaranarayanan, and R. Baraniuk. “SpaRCS: Recovering low-rank and sparse matrices from compressive measurements”. In: *Advances in Neural Information Processing Systems*. Vol. 24. Curran Associates, Inc., 2011 (cited on pages 2, 6).
- [WWAF06] A. Weiße, G. Wellein, A. Alvermann, and H. Fehske. “The kernel polynomial method”. In: *Reviews of Modern Physics* 78.1 (2006), pp. 275–306 (cited on page 2).