
AN ACCELERATED DISTRIBUTED STOCHASTIC GRADIENT METHOD WITH MOMENTUM

A PREPRINT

Kun Huang

The Chinese University of Hong Kong, Shenzhen
School of Data Science (SDS)
Shenzhen, Guangdong, China
kunhuang@link.cuhk.edu.cn

Shi Pu

The Chinese University of Hong Kong, Shenzhen
School of Data Science (SDS)
Shenzhen, Guangdong, China
pushi@cuhk.edu.cn

Angelia Nedić

Arizona State University
School of Electrical, Computer and Energy Engineering
Tempe, AZ, United States
angelia.nedich@asu.edu

February 20, 2024

ABSTRACT

In this paper, we introduce an accelerated distributed stochastic gradient method with momentum for solving the distributed optimization problem, where a group of n agents collaboratively minimize the average of the local objective functions over a connected network. The method, termed “Distributed Stochastic Momentum Tracking (DSMT)”, is a single-loop algorithm that utilizes the momentum tracking technique as well as the Loopless Chebyshev Acceleration (LCA) method. We show that DSMT can asymptotically achieve comparable convergence rates as centralized stochastic gradient descent (SGD) method under a general variance condition regarding the stochastic gradients. Moreover, the number of iterations (transient times) required for DSMT to achieve such rates behaves as $\mathcal{O}(n^{5/3}/(1-\lambda))$ for minimizing general smooth objective functions, and $\mathcal{O}(\sqrt{n/(1-\lambda)})$ under the Polyak-Łojasiewicz (PL) condition. Here, the term $1-\lambda$ denotes the spectral gap of the mixing matrix related to the underlying network topology. Notably, the obtained results do not rely on multiple inter-node communications or stochastic gradient accumulation per iteration, and the transient times are the shortest under the setting to the best of our knowledge.

1 Introduction

We investigate how a group of networked agents $\mathcal{N} := \{1, 2, \dots, n\}$ collaborate to solve the following distributed optimization problem:

$$\min_{x \in \mathbb{R}^p} f(x) := \frac{1}{n} \sum_{i=1}^n f_i(x), \quad (1)$$

where each agent i has access only to its local objective function $f_i : \mathbb{R}^p \rightarrow \mathbb{R}$. Problem (1) arises in various fields, including signal processing, distributed estimation, and machine learning. Specifically, solving large-scale problems such as training foundation models [3, 6, 54, 30] in specialized and centralized clusters can be costly [50, 44], while decentralized training offers a promising alternative that relieves the robustness bottleneck and reduces the high latency at the central server [28]. In this paper, we consider the typical setting where each agent can query unbiased noisy gradients of $f_i(x)$ in the form of $g_i(x; \xi)$ that satisfies a certain variance condition, where ξ represents a random sample or a batch of random samples.

Decentralization may slow down the optimization process due to the partial communication over sparse networks. For instance, the number of iterations required for the distributed subgradient descent (DGD) method [29] to reach certain accuracy threshold significantly grows with the network size [36], downgrading its performance compared to the centralized subgradient method. When stochastic gradients are used, however, several distributed methods such as those in [21, 37, 11, 42, 40, 34, 1, 51, 47, 12, 17] have exhibited the so-called ‘‘asymptotic network independent (ANI)’’ property [36]. In other words, these methods achieve the same convergence rates as their centralized counterparts after a finite transient time has passed. Such a property ensures that the required number of iterations to achieve high accuracy do not increase significantly with the network size. Nevertheless, it is still critical to develop algorithms with shorter transient times that typically depend on the network size and topology. Currently, the best known transient times for minimizing general smooth objective functions are $\mathcal{O}(n^3/(1 - \lambda)^2)$ [1, 12] and $\mathcal{O}(n/(1 - \lambda))$ [11, 1, 51] with and without the strong convexity (or the Polyak-Łojasiewicz condition) on the objective functions, respectively.

Chebyshev Acceleration (CA) has been a popular technique for achieving faster consensus among networked agents during the distributed optimization process [38]. When CA is combined with stochastic gradient accumulation using a large batch size at every iteration, the optimal convergence guarantee can be achieved [24, 53]. However, CA requires inner loops of multiple communication steps which may affect the practical performance of the algorithms, and large batches are not always available. Inspired by the recently developed Loopless Chebyshev Acceleration (LCA) technique that works without inner loops [39], we consider in this paper a new method termed ‘‘Distributed Stochastic Momentum Tracking (DSMT)’’ that incorporates two key features. First, the LCA technique is utilized to accelerate the communication process and avoids multiple communication steps between two successive stochastic gradient computations. Second, each agent employs an auxiliary variable to track the average momentum parameter over the network. We show that DSMT improves the transient times over the existing works for minimizing smooth objective functions, with or without the Polyak-Łojasiewicz (PL) condition. In particular, the convergence rates of DSMT are derived under the most general condition on the stochastic gradients under the distributed setting [15, 20, 19], enhancing the applicability of the method.

It is worth noting that, the convergence result of DSMT implies the momentum parameter can benefit the algorithmic convergence of distributed stochastic gradient methods, particularly related to the network topology. Such an observation can potentially inspire future development of distributed optimization algorithms.

1.1 Related Works

There exists a rich literature on the distributed implementation of stochastic gradient methods over networks. For example, early works including [4, 27, 5, 32, 33] suggest that distributed stochastic gradient methods may achieve comparable performance with centralized SGD under specific settings. More recently, the paper [21] first demonstrates the ANI property enjoyed by the distributed stochastic gradient descent (DSGD) method, followed by the study in [35, 17, 36]. Subsequent research that considers gradient tracking based methods [47, 34, 16, 46, 49] and primal-dual like methods [11, 51, 43, 12] also demonstrate the ANI property while relieving from the data heterogeneity challenge encountered by DSGD. Both types of methods are unified in [1].

Recently, there exists a line of works that achieve the optimal iteration complexity in distributed stochastic optimization by integrating stochastic gradient accumulation and accelerated communication techniques such as Chebyshev Acceleration (CA) [53, 24]. In particular, the works in [53] and [24] obtain the optimal iteration complexity under smooth objective functions with and without the PL condition, respectively. For finite sum problems, the work in [25] achieves the optimal guarantee by further incorporating a variance reduction technique.¹ Nevertheless, CA that relies on inner loops of multiple communication steps may not be communication-efficient, and sampling large batches for computing the stochastic gradients is not always practical [46]. Therefore, we are motivated to develop an efficient distributed stochastic gradient method that samples constant batches for computing the stochastic gradients and performs one round of communication at every iteration.

The momentum method [31] is popular for accelerating the convergence of first-order methods. Recently, it has been applied to the distributed setting [7, 52, 45, 22], resulting in practical improvements. However, it remains unclear whether momentum parameters can benefit the convergence of distributed stochastic gradient methods theoretically.

1.2 Main Contribution

The main contribution of this paper is two-fold.

¹It is worth mentioning that when the objective functions take a finite sum form, distributed random reshuffling (RR) methods can further improve the iteration complexity [9, 13].

Method	\mathfrak{g}	Additional Assumption	Transient Time Nonconvex	Transient Time PL condition
DSGD [21]	$(\sigma^2, 0)$	$(\zeta^2, 0)$ -BGD	$\mathcal{O}\left(\frac{n^3}{(1-\lambda)^4}\right)$	$\mathcal{O}\left(\frac{n}{(1-\lambda)^2}\right)$
DSGD [37]	(σ^2, η^2)	$(\zeta^2, 0)$ -BGD	/	$\mathcal{O}\left(\frac{n}{(1-\lambda)^2}\right)$
ED [1]	$(\sigma^2, 0)$	/	$\mathcal{O}\left(\frac{n^3}{(1-\lambda)^2}\right)$	$\mathcal{O}\left(\frac{n}{1-\lambda}\right)$
EDAS[11]	(σ^2, η^2)	/	/	$\mathcal{O}\left(\frac{n}{1-\lambda}\right)$
DSGT [1]	$(\sigma^2, 0)$	/	$\mathcal{O}\left(\max\left\{\frac{n^3}{(1-\lambda)^2}, \frac{n}{(1-\lambda)^{8/3}}\right\}\right)$	$\mathcal{O}\left(\max\left\{\frac{n}{1-\lambda}, \frac{n^{1/3}}{(1-\lambda)^{4/3}}\right\}\right)$
DSGT-HB [7]	$(\sigma^2, 0)$	/	$\mathcal{O}\left(\frac{n^3}{(1-\lambda)^4}\right)$	/
QG-DSGDm [22]	$(\sigma^2, 0)$	$(\zeta^2, 0)$ -BGD	$\mathcal{O}\left(\frac{n^3}{(1-\lambda)^4}\right)$	/
Momentum Tracking [41]	$(\sigma^2, 0)$	/	$\mathcal{O}\left(\frac{n^3}{(1-\lambda)^8}\right)$	/
DSMT (This paper)	ABC	f_i lower bounded	$\mathcal{O}\left(\frac{n^{5/3}}{1-\lambda}\right)$	$\mathcal{O}\left(\sqrt{\frac{n}{1-\lambda}}\right)$

Table 1: Comparison of different methods regarding the assumptions and the transient times for minimizing smooth objective functions. The “ \mathfrak{g} ” column describes the conditions assumed on the stochastic gradients, where the notation (σ^2, η^2) represents the relaxed growth condition, i.e., $\mathbb{E}[\|g_i(x; \xi_i) - \nabla f_i(x)\|^2 | x] \leq \sigma^2 + \eta^2 \|\nabla f_i(x)\|^2, \forall i \in \mathcal{N}$, and ABC stands for Assumption 1.2. The notation (ζ^2, ψ^2) -BGD stands for the bounded gradient dissimilarity assumption that $\frac{1}{n} \sum_{i=1}^n \|\nabla f_i(x) - \nabla f(x)\|^2 \leq \zeta^2 + \psi^2 \|\nabla f(x)\|^2$.

Firstly, the proposed DSMT method achieves improved convergence rates and better transient times over the existing distributed stochastic gradient methods (see Table 1). Specifically, for general smooth objective functions, DSMT shortens the state-of-the-art transient time from $\mathcal{O}(n^3/(1-\lambda)^2)$ to $\mathcal{O}(n^{5/3}/(1-\lambda))$, where $1-\lambda$ denotes the spectral gap of the mixing matrix that may scale as $\mathcal{O}(1/n^2)$ for sparse networks. When the objective functions further satisfy the PL condition, DSMT shortens the state-of-the-art transient time from $\mathcal{O}(n/(1-\lambda))$ to $\mathcal{O}(\sqrt{n/(1-\lambda)})$.

Secondly, the convergence results of DSMT only assumes the so-called ABC condition [15, 20, 19] on the stochastic gradients, which is the most general variance condition under the distributed settings to our knowledge. By comparison, existing works depend on the relaxed growth condition (RGC) or the bounded variance [2] regarding the stochastic gradients or the bounded gradient dissimilarity (BGD) assumption concerning the data heterogeneity among the agents. Such a generalization expands the practical applicability of DSMT given that the ABC condition covers counterexamples that do not satisfy RGC [10, 15].

1.3 Notation and Assumptions

Throughout this paper, column vectors are considered by default unless specified otherwise. We use $x_{i,k} \in \mathbb{R}^p$ to denote the iterate of agent i at the k -th iteration. For the sake of clarity and presentation, we introduce the stacked variables as follows:

$$\begin{aligned} \mathbf{x}_k &:= (x_{1,k}, x_{2,k}, \dots, x_{n,k})^\top \in \mathbb{R}^{n \times p}, \\ \nabla F(\mathbf{x}_k) &:= (\nabla f_1(x_{1,k}), \nabla f_2(x_{2,k}), \dots, \nabla f_n(x_{n,k}))^\top \in \mathbb{R}^{n \times p}, \\ A_{\#} &:= \begin{pmatrix} A \\ A \end{pmatrix} \in \mathbb{R}^{2n \times p}. \end{aligned}$$

We use $\bar{x} \in \mathbb{R}^p$ to denote the averaged variables of x_i among the agents. For instance, the variable $\bar{x}_k := 1/n \sum_{i=1}^n x_{i,k}$ stands for the average of all the agents’ iterates at the k -th iteration. We use $\|\cdot\|$ to denote the Frobenius norm for a matrix by default and the ℓ_2 norm for a vector. The term $\langle a, b \rangle$ stands for the inner product of two vectors $a, b \in \mathbb{R}^p$. For two matrices $A, B \in \mathbb{R}^{n \times p}$, $\langle A, B \rangle$ is defined as $\langle A, B \rangle := \sum_{i=1}^n \langle A_i, B_i \rangle$, where A_i (and B_i) represents the i -row of A (and B).

We next introduce the standing assumptions. Assumption 1.1 is common that requires the objective functions to be smooth and lower bounded.

Assumption 1.1. Each $f_i(x) : \mathbb{R}^p \rightarrow \mathbb{R}$ is L -smooth, i.e.,

$$\|\nabla f_i(x) - \nabla f_i(x')\| \leq L \|x - x'\|, \forall x, x' \in \mathbb{R}^p.$$

In addition, each $f_i(x)$ is bounded from below, i.e., $f_i(x) \geq f_i^* := \inf_x f_i(x) > -\infty, \forall x \in \mathbb{R}^p$. We also denote $f^* := \inf_x f(x)$.

Let the filtration $\{\mathcal{F}_k\}_{k \geq 0}$ be generated by $\{\xi_{i,\ell} | i \in \mathcal{N}, \ell = 0, 1, \dots, k-1\}$. We consider the ABC condition in Assumption 1.2 regarding the stochastic gradients.

Assumption 1.2. Assume each agent has access to an conditionally unbiased stochastic gradient $g_i(x; \xi_i)$ of $\nabla f_i(x)$, i.e., $\mathbb{E}[g_i(x; \xi_{i,k}) | \mathcal{F}_k] = \nabla f_i(x)$, and there exist constants $C, \sigma \geq 0$ such that for any $k \in \mathbb{N}, i \in \mathcal{N}$,

$$\mathbb{E} \left[\|g_i(x; \xi_{i,k}) - \nabla f_i(x)\|^2 \middle| \mathcal{F}_k \right] \leq C [f_i(x) - f_i^*] + \sigma^2, \quad (2)$$

In addition, the stochastic gradients are independent across different agents at each $k \geq 0$.

It is noteworthy that the relaxed growth condition (RGC) implies (2) given that each f_i has Lipschitz continuous gradients and is lower bounded [15], however, there exist examples that only satisfy the ABC condition but not RGC [15, 10]. Moreover, Assumption 1.2 generally holds for the empirical risk minimization problems if the stochastic gradients are queried through uniformly sampling with replacement [10].

We assume the agents in the network are connected via a graph $\mathcal{G} = (\mathcal{N}, \mathcal{E})$ with $\mathcal{E} \subseteq \mathcal{N} \times \mathcal{N}$ representing the set of edges connecting the agents. In particular, $(i, i) \in \mathcal{E}$ for all $i \in \mathcal{N}$. The set of neighbors for agent i is denoted by $\mathcal{N}_i = \{j \in \mathcal{N} : (i, j) \in \mathcal{E}\}$. The element w_{ij} in the weight matrix $W \in \mathbb{R}^{n \times n}$ represents the weight of the edge between agents i and j . Regarding the network topology, we consider Assumption 1.3 that is standard in the distributed optimization literature. The condition guarantees that the spectral norm λ of the matrix $(W - \mathbf{1}\mathbf{1}^\top/n)$ is strictly less than one.

Assumption 1.3. The graph \mathcal{G} is undirected and connected, i.e., there exists a path between any two nodes in \mathcal{G} . There is a direct link between i and j ($i \neq j$) in \mathcal{G} if and only if $w_{ij} > 0$ and $w_{ji} > 0$; otherwise, $w_{ij} = w_{ji} = 0$. The mixing matrix W is nonnegative and doubly stochastic, i.e., $\mathbf{1}^\top W = \mathbf{1}^\top$ and $W\mathbf{1} = \mathbf{1}$. In addition, we assume $w_{ii} > 0$ for any $i \in \mathcal{N}$.

1.4 Organization

The rest of this paper is organized as follows. In Section 2, we introduce the proposed DSMT algorithm starting with its motivation. We then proceed to conduct a preliminary analysis in Section 3. The main convergence results of DSMT under smooth objective functions with or without the PL condition are presented in Section 4. We then provide numerical experiments in Section 5 and conclude the paper in Section 6.

2 A Distributed Stochastic Momentum Tracking Method

In this section, we introduce the proposed algorithm, Distributed Stochastic Momentum Tracking (DSMT). We start with the motivation and the intuition behind DSMT.

2.1 Motivation

The proposed algorithm relies heavily on the so-called Loopless Chebyshev Acceleration (LCA) technique stated in Lemma 2.1 below.

Lemma 2.1 (Lemma 11 in [39]). Suppose the mixing matrix W is symmetric and positive semidefinite. Define $\eta_w := 1/(1 + \sqrt{1 - \lambda^2})$, then $\tilde{\rho}_w := \sqrt{\eta_w} \sim \mathcal{O}(1 - \sqrt{1 - \lambda^2})$, and for any $A \in \mathbb{R}^{n \times p}$ and $k \geq 0$,

$$\left\| \tilde{\Pi} \tilde{W}^k \tilde{\Pi} A_\# \right\|^2 \leq c_0 \tilde{\rho}_w^{2k} \|\Pi A\|^2, \quad c_0 = 14,$$

where

$$\tilde{\Pi} := \begin{pmatrix} \Pi & \mathbf{0} \\ \mathbf{0} & \Pi \end{pmatrix}, \quad \Pi := I - \frac{\mathbf{1}\mathbf{1}^\top}{n}, \quad A_\# := \begin{pmatrix} A \\ A \end{pmatrix}, \quad \tilde{W} := \begin{pmatrix} (1 + \eta_w)W & -\eta_w I \\ I & \mathbf{0} \end{pmatrix}.$$

The above lemma implies that, by employing an augmented mixing matrix \tilde{W} , the consensus rate among multiple agents can be accelerated compared to applying the basic consensus procedure using the matrix W . In addition, when LCA is employed in distributed optimization algorithms, there is no need to perform inner loops of multiple communication rounds. For instance, the Optimal Gradient Tracking (OGT) method proposed in [39] achieves the optimal communication and computation complexities simultaneously without inner loops of multiple communication rounds for minimizing smooth strongly convex objective functions using full gradients.

In the following discussion, we adhere to the notations introduced in Lemma 2.1. Additionally, we use $[\tilde{\mathbf{x}}_k]_{1:n}$ to represent the first n rows of $\tilde{\mathbf{x}}_k \in \mathbb{R}^{2n \times p}$.

It is noteworthy that applying the LCA technique to distributed stochastic gradient methods while achieving improved transient times is a nontrivial task. A recent attempt in [49] realized the same transient time $\mathcal{O}(n/(1-\lambda))$ as those presented in [11, 1, 51]. We discuss the reasons of this result as follows. In [49], the LCA technique is directly combined with the Distributed Stochastic Gradient Tracking (DSGT) method described in (3), where $\mathbf{g}_k := (g_1(x_{1,k}; \xi_{1,k}), \dots, g_n(x_{n,k}; \xi_{n,k}))^\top \in \mathbb{R}^{n \times p}$ denotes the stacked stochastic gradient at the k -th iteration.

$$\mathbf{x}_{k+1} = W(\mathbf{x}_k - \alpha \mathbf{y}_k), \quad (3a)$$

$$\mathbf{y}_{k+1} = W(\mathbf{y}_k + \mathbf{g}_{k+1} - \mathbf{g}_k), \mathbf{y}_0 = \mathbf{g}_0. \quad (3b)$$

Here, the auxiliary variables \mathbf{y}_k track the averaged stochastic gradients among the agents since $\bar{y}_k = \bar{g}_k$ based on Assumption 1.3 and induction from (3b). One common approach for analyzing the convergence of DSGT involves estimating the following term (see, e.g., [34, 49] or (29)):

$$\mathbb{E} \left[\|\mathbf{x}_k - \mathbf{1}\bar{x}_k^\top\|^2 \right] + \frac{\alpha^2 \mathcal{C}_0}{(1-\lambda)^2} \mathbb{E} \left[\|\mathbf{y}_k - \mathbf{1}\bar{y}_k^\top\|^2 \right], \quad (4)$$

for some constant $\mathcal{C}_0 > 0$ that is independent of n and $(1-\lambda)$. However, this leads to an unsatisfactory performance of the algorithm, primarily due to the bias in \mathbf{y}_k , i.e., $\mathbb{E}[\mathbf{y}_k - \mathbf{g}_k | \mathcal{F}_k] \neq 0$.

To alleviate the bias in \mathbf{y}_k , we introduce alternative variables, \mathbf{z}_k , for \mathbf{y}_k to track. Specifically, the update \mathbf{y}_k is as follows:

$$\mathbf{y}_{k+1} = W(\mathbf{y}_k + \mathbf{z}_{k+1} - \mathbf{z}_k). \quad (5)$$

Simultaneously, the variables \mathbf{z}_k follow the update defined in (6):

$$\mathbf{z}_{k+1} - \mathbf{z}_k = -(1-\beta)(\mathbf{z}_k - \nabla F(\mathbf{x}_{k+1})) + (1-\beta)[\mathbf{g}_{k+1} - \nabla F(\mathbf{x}_{k+1})], \quad (6)$$

where $\beta \in (0, 1)$. With the help of the additional term $(1-\beta)$ in (6), the impact of the stochastic gradient variance is better controlled, as detailed in Lemma 3.7. Such a strategic adjustment enables the cancellation of the extra term $1/(1-\lambda)$ introduced by (4).

Remark 2.1. *Substituting the update (5) and (6) into the DSGT method (3) leads to the following relations.*

$$\begin{aligned} \mathbf{x}_{k+1} &= W(\mathbf{x}_k - \alpha \mathbf{y}_k), \\ \mathbf{z}_{k+1} &= \beta \mathbf{z}_k + (1-\beta) \mathbf{g}_{k+1}, \\ \mathbf{y}_{k+1} &= W(\mathbf{y}_k + \mathbf{z}_{k+1} - \mathbf{z}_k), \mathbf{y}_0 = \mathbf{z}_0 = (1-\beta) \mathbf{g}_0. \end{aligned} \quad (7)$$

Invoking Assumption 1.3, we can deduce that $\bar{y}_k = \bar{z}_k$ by induction. Consequently, the behavior of \bar{x}_k as described in (8) resembles the update of the Stochastic Gradient Descent with Momentum (SGDM) method [31]. It turns out that \mathbf{y}_k tracks the momentum variable \mathbf{z}_k , and this is why the proposed method is named ‘‘distributed stochastic momentum tracking’’.

$$\begin{aligned} \bar{x}_{k+1} &= \bar{x}_k - \alpha \bar{z}_k, \\ \bar{z}_{k+1} &= \beta \bar{z}_k + (1-\beta) \bar{g}_k. \end{aligned} \quad (8)$$

The above discussions motivate the proposed method in the next part.

2.2 The Proposed Method

We begin by presenting the compact form of the Distributed Stochastic Momentum Tracking (DSMT) method in (9). In essence, DSMT leverages the Loopless Chebyshev Acceleration (LCA) technique for the update (7).

$$\tilde{\mathbf{x}}_{k+1} = \tilde{W}(\tilde{\mathbf{x}}_k - \alpha(\mathbf{y}_k)_\#) \quad (9a)$$

$$\mathbf{x}_{k+1} = [\tilde{\mathbf{x}}_{k+1}]_{1:n} \quad (9b)$$

$$\mathbf{z}_{k+1} = \beta\mathbf{z}_k + (1 - \beta)\mathbf{g}_{k+1} \quad (9c)$$

$$\tilde{\mathbf{y}}_{k+1} = \tilde{W}(\tilde{\mathbf{y}}_k + (\mathbf{z}_{k+1})_\# - (\mathbf{z}_k)_\#) \quad (9d)$$

$$\mathbf{y}_{k+1} = [\tilde{\mathbf{y}}_{k+1}]_{1:n}, \quad (9e)$$

where \tilde{W} defined in Lemma 2.1 signifies the application of the LCA technique, and $(\mathbf{y}_k)_\# := (\mathbf{y}_k^\top, \mathbf{y}_k^\top)^\top \in \mathbb{R}^{2n \times p}$. Consequently, the updates in (9b) and (9e) are required to obtain \mathbf{x}_{k+1} and $(\mathbf{y}_{k+1})_\#$ in the subsequent iterations. The updates in (9c) and (9d) align with the rationale explained for (7), representing a critical step in improving the convergence rate compared to the method outlined in [49].

The formal description of the DSMT method is outlined in Algorithm 1. At each iteration, agent i first performs an approximate gradient descent step in Line 4 to obtain $x_{i,k+\frac{1}{2}}$ and $x_{i,k+\frac{1}{2}}^l$. Then, after communicating with neighboring agents the updated $x_{i,k+\frac{1}{2}}$ in Line 5, the new iterates $x_{i,k+1}$ and $x_{i,k+1}^l$ are obtained. Subsequently, agent i queries a noisy gradient based on $x_{i,k+1}$ and completes the update of $z_{i,k+1}$ in Line 7. The remaining procedures aim to track the averaged variable \bar{z}_k given the initialization $\mathbf{y}_0 = \mathbf{z}_0 = (1 - \beta)\mathbf{g}_0$.

Algorithm 1 Distributed Stochastic Momentum Tracking (DSMT)

- 1: Initialize $x_{i,0} = x_{i,0}^l \in \mathbb{R}^p$ for all agent $i \in \mathcal{N}$, determine $W = [w_{ij}] \in \mathbb{R}^{n \times n}$, stepsize α and parameter β .
Initialize $y_{i,0} = y_{i,0}^l = z_{i,0} = (1 - \beta)g_{i,0}$. Input η_w .
 - 2: **for** $k = 0, 1, 2, \dots, K - 1$ **do**
 - 3: **for** Agent $i = 1, 2, \dots, n$ in parallel **do**
 - 4: Update $x_{i,k+\frac{1}{2}} = x_{i,k} - \alpha y_{i,k}$ and $x_{i,k+\frac{1}{2}}^l = x_{i,k}^l - \alpha y_{i,k}$.
 - 5: Communicate and update $x_{i,k+1} = (1 + \eta_w) \sum_{j \in \mathcal{N}_i} w_{ij} x_{j,k+\frac{1}{2}} - \eta_w x_{i,k+\frac{1}{2}}^l$ and $x_{i,k+1}^l = x_{i,k+\frac{1}{2}}^l$.
 - 6: Acquires a stochastic gradient $g_{i,k+1} = g_i(x_{i,k+1}; \xi_{i,k+1}) \in \mathbb{R}^p$.
 - 7: Update $z_{i,k+1} = \beta z_{i,k} + (1 - \beta)g_{i,k+1}$.
 - 8: Update $y_{i,k+\frac{1}{2}} = y_{i,k} + z_{i,k+1} - z_{i,k}$ and $y_{i,k+\frac{1}{2}}^l = y_{i,k}^l + z_{i,k+1} - z_{i,k}$.
 - 9: Communicate and update $y_{i,k+1} = (1 + \eta_w) \sum_{j \in \mathcal{N}_i} w_{ij} y_{j,k+\frac{1}{2}} - \eta_w y_{i,k+\frac{1}{2}}^l$ and $y_{i,k+1}^l = y_{i,k+\frac{1}{2}}^l$.
 - 10: **end for**
 - 11: **end for**
 - 12: Output $x_{i,T}$ for all agent $i \in \mathcal{N}$.
-

Remark 2.2. We compare DSMT method with the DSGT method. It is worth noting that DSMT necessitates additional information of λ compared to DSGT. This is common for constructing accelerated or optimal methods including those in [24, 53, 38]. Regarding the communication cost per iteration, each agent in the DSMT method communicates variables $(x_{i,k}, y_{i,k})_{k \geq 0}$, which aligns with the communication load of DSGT. The additional cost primarily stems from the storage of variables $(x_{i,k}^l, y_{i,k}^l, z_{i,k})_{k \geq 0}$.

Remark 2.3. In the analysis, we introduce the definition $\mathbf{z}_{-1} := \mathbf{0}$. It is worth noting that the relation $\mathbf{z}_0 = \beta\mathbf{z}_{-1} + (1 - \beta)\mathbf{g}_0$ remains valid due to the initialization $\mathbf{z}_0 = (1 - \beta)\mathbf{g}_0$ in Algorithm 1. As a consequence, the subsequent analysis involving \mathbf{z}_{-1} is valid.

3 Preliminary Analysis

In this section, we present several preliminary results. We start with establishing the recursions involving the averaged variables \bar{x}_k , \bar{y}_k , and \bar{z}_k .

Lemma 3.1. Let Assumption 1.3 hold and $\mathbf{y}_0 = \mathbf{z}_0 = (1 - \beta)\mathbf{g}_0$, we have for any $k \geq 0$ that

$$\begin{aligned} \bar{x}_{k+1} &= \bar{x}_k - \alpha \bar{y}_k, \\ \bar{z}_{k+1} &= \beta \bar{z}_k + (1 - \beta) \bar{g}_{k+1}, \\ \bar{y}_{k+1} &= \bar{z}_{k+1}. \end{aligned}$$

Proof. See Appendix A.1. □

As highlighted in Remark 2.1, the update of \bar{x}_k can be viewed as an approximate version of the SGDM method. Such an observation prompts us to investigate the behavior of a series of auxiliary variables $\{\bar{d}_k\}$ defined as follows:

$$\bar{d}_k := \begin{cases} \bar{x}_k, & k = 0 \\ \frac{1}{1-\beta}\bar{x}_k - \frac{\beta}{1-\beta}\bar{x}_{k-1}, & k \geq 1. \end{cases} \quad (10)$$

The behavior of \bar{d}_k is more like the update of SGD compared to that of \bar{x}_k ; see Lemma 3.2 below. Similar arguments have been explored in prior works, including [48, 8, 23].

Lemma 3.2. *Let Assumption 1.3 hold. Define \bar{d}_k as in (10), we have*

$$\bar{d}_{k+1} = \bar{d}_k - \alpha \bar{g}_k, \quad k \geq 0,$$

and

$$\bar{d}_k - \bar{x}_k = -\frac{\alpha\beta}{1-\beta}\bar{z}_{k-1}, \quad k \geq 0. \quad (11)$$

Proof. See Appendix A.2. □

Next, we consider the relation between $\mathbb{E}[f(\bar{d}_k) - f^*]$ and $\mathbb{E}[f(\bar{x}_k) - f^*]$ in light of (11) in Lemma 3.2. As a consequence, the follow-up analysis needs not involve the term $\mathbb{E}[f(\bar{x}_k) - f^*]$ which may appear due to Assumption 1.2.

Lemma 3.3. *Let Assumptions 1.1-1.3 hold. Set the stepsize α to satisfy $\alpha \leq (1-\beta)/(3\beta L)$. We have*

$$\mathbb{E}[f(\bar{x}_k) - f^*] \leq \mathbb{E}[f(\bar{d}_k) - f^*] + \frac{\alpha\beta}{1-\beta}\mathbb{E}[\|\nabla f(\bar{x}_k)\|^2] + \frac{\alpha\beta}{1-\beta}\mathbb{E}[\|\bar{z}_{k-1}\|^2].$$

Proof. See Appendix A.3. □

Our primary goal in this section is to locate and analyze a Lyapunov function to facilitate the convergence analysis. We start with Lemma 3.4 that presents the recursion of $\mathbb{E}[f(\bar{d}_k) - f^*]$ in light of Assumption 1.1 and Lemma 3.2.

Lemma 3.4. *Let Assumptions 1.1-1.3 hold. Set the stepsize α to satisfy $\alpha \leq \min\{1/(4L), 1/(2C)\}$. We have*

$$\begin{aligned} \mathbb{E}[f(\bar{d}_{k+1}) - f^*] &\leq \left(1 + \frac{2\alpha^2 CL}{n}\right)\mathbb{E}[f(\bar{d}_k) - f^*] - \frac{\alpha}{2}\left(1 - \frac{4\alpha^2 CL}{n(1-\beta)}\right)\mathbb{E}[\|\nabla f(\bar{x}_k)\|^2] \\ &\quad + \frac{\alpha L^2}{n}\mathbb{E}[\|\Pi\mathbf{x}_k\|^2] + \frac{\alpha^3\beta^2 L(L+2C)}{(1-\beta)^2}\mathbb{E}[\|\bar{z}_{k-1}\|^2] + \frac{\alpha^2 L(2C\sigma_f^* + \sigma^2)}{n}, \end{aligned}$$

where $\sigma_f^* := f^* - \frac{1}{n}\sum_{i=1}^n f_i^*$.

Proof. See Appendix A.4. □

Lemma 3.4 inspires us to consider the recursion of $\mathbb{E}[\|\bar{z}_k\|^2]$ as follows.

Lemma 3.5. *Let Assumptions 1.1-1.3 hold. Set the stepsize α to satisfy $\alpha \leq \min\{(1-\beta)/(2\beta L), 1/(4C\beta)\}$. We have*

$$\begin{aligned} \mathbb{E}[\|\bar{z}_k\|^2] &\leq \frac{1+\beta}{2}\mathbb{E}[\|\bar{z}_{k-1}\|^2] + \frac{2C(1-\beta)^2}{n}\mathbb{E}[f(\bar{d}_k) - f^*] + 3(1-\beta)\mathbb{E}[\|\nabla f(\bar{x}_k)\|^2] \\ &\quad + \frac{(1-\beta)^2(2C\sigma_f^* + \sigma^2)}{n} + \frac{2(1-\beta)L}{n}\left(L + \frac{C(1-\beta)}{n}\right)\mathbb{E}[\|\Pi\mathbf{x}_k\|^2]. \end{aligned}$$

Proof. See Appendix A.5. □

We next consider the recursion of $\mathbb{E}[\|\mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top)\|^2]$.

Lemma 3.6. *Let Assumptions 1.1-1.3 hold. Set the stepsize α to satisfy $\alpha \leq (1 - \beta)/(3L)$. We have for any $k \geq 0$ that*

$$\begin{aligned} \mathbb{E} \left[\|\mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top)\|^2 \right] &\leq \beta \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right] + 2\alpha n(L + C) \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] \\ &\quad + 2(1 - \beta)L(L + C) \mathbb{E} \left[\|\Pi \mathbf{x}_k\|^2 \right] + 2n(1 - \beta)^2 C \mathbb{E} \left[f(\bar{d}_k) - f^* \right] + 2\alpha n(1 - \beta) C \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] \\ &\quad + n(1 - \beta)^2 (2C\sigma_f^* + \sigma^2), \end{aligned}$$

where $\mathbf{x}_{-1} := \mathbf{x}_0$ and $\mathbf{z}_{-1} = \mathbf{0}$.

Proof. See Appendix A.6. □

Next, we consider the consensus error terms $\mathbb{E}[\|\Pi \mathbf{x}_k\|^2]$ and $\mathbb{E}[\|\Pi \mathbf{y}_k\|^2]$ similar to those in [39]. One of the key steps in the derivation of Lemma 3.7 involves identifying the error term $\mathbb{E}[\|\mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top)\|^2]$, as discussed in Subsection 2.1. This particular step is essential for preserving the benefits of the gradient tracking technique, which reduces the impact of data heterogeneity.

Lemma 3.7. *Let Assumptions 1.1-1.3 hold. Set the stepsize α and the parameter β to satisfy*

$$\alpha \leq \min \left\{ \frac{1}{2(L + C)}, \sqrt{\frac{1 - \tilde{\rho}_w}{6c_0^2 L(L + C)}}, \sqrt{\frac{1 - \beta}{4CL}} \right\}, \quad 1 > \beta \geq \tilde{\rho}_w.$$

Then, there exist sequences $\{\mathcal{R}_k^x\}$ and $\{\mathcal{R}_k^y\}$ such that

$$\begin{aligned} \mathbb{E} \left[\|\Pi \mathbf{x}_k\|^2 \right] &\leq \mathbb{E} \left[\|\tilde{\Pi} \bar{\mathbf{x}}_k\|^2 \right] \leq \mathcal{R}_k^x, \\ \mathbb{E} \left[\|\Pi \mathbf{y}_k\|^2 \right] &\leq \mathbb{E} \left[\|\tilde{\Pi} \bar{\mathbf{y}}_k\|^2 \right] \leq \mathcal{R}_k^y, \end{aligned}$$

and

$$\mathcal{R}_{k+1}^x \leq \tilde{\rho}_w \mathcal{R}_k^x + \frac{\alpha^2 c_0 \tilde{\rho}_w}{1 - \tilde{\rho}_w} \mathcal{R}_k^y, \quad (12)$$

$$\begin{aligned} \mathcal{R}_{k+1}^y &\leq \frac{1 + \tilde{\rho}_w}{2} \mathcal{R}_k^y + \frac{15c_0(1 - \beta)^2 L(L + C)}{1 - \tilde{\rho}_w} \mathcal{R}_k^x + \frac{3c_0(1 - \beta)^2 \beta}{1 - \tilde{\rho}_w} \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right] \\ &\quad + 9\alpha n c_0(1 - \beta)(C + L) \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] + 12n c_0(1 - \beta)^2 C \mathbb{E} \left[f(\bar{d}_k) - f^* \right] \\ &\quad + 20\alpha n c_0(1 - \beta)(C + L) \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] + 6n c_0(1 - \beta)^2 (2C\sigma_f^* + \sigma^2), \end{aligned} \quad (13)$$

where $c_0 = 14$ is defined in Lemma 2.1 and

$$\begin{aligned} \mathcal{R}_0^x &\leq c_0 \|\Pi \mathbf{x}_0\|^2 + \frac{\alpha^2 c_0 \tilde{\rho}_w}{1 - \tilde{\rho}_w} \mathbb{E} \left[\|\Pi \mathbf{y}_0\|^2 \right], \\ \mathcal{R}_0^y &:= c_0 \mathbb{E} \left[\|\Pi \mathbf{y}_0\|^2 \right] + n c_0 \tilde{\rho}_w (1 - \beta)^2 (2C\sigma_f^* + \sigma^2). \end{aligned} \quad (14)$$

Proof. See Appendix A.7. □

We are now ready to introduce the Lyapunov function \mathcal{L}_k defined in (15):

$$\begin{aligned} \mathcal{L}_k &:= \mathbb{E} \left[f(\bar{d}_k) - f^* \right] + \frac{3\alpha L^2}{n(1 - \tilde{\rho}_w)} \mathcal{R}_k^x + \frac{12\alpha^3 c_0 \tilde{\rho}_w L^2}{n(1 - \tilde{\rho}_w)^3} \mathcal{R}_k^y \\ &\quad + \frac{4\alpha^3 L(L + 2C)}{(1 - \beta)^3} \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] + \frac{48\alpha^3 c_0^2 \tilde{\rho}_w L^2}{n(1 - \tilde{\rho}_w)^3} \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right]. \end{aligned} \quad (15)$$

Lemma 3.8 below presents the recursion for \mathcal{L}_k by combining the results in Lemmas 3.4-3.7.

Lemma 3.8. *Let Assumptions 1.1-1.3 hold. Set the stepsize α and the parameter β to satisfy*

$$\alpha \leq \min \left\{ \sqrt{\frac{(1 - \tilde{\rho}_w)^2}{240c_0^2L(L+C)}}, \frac{1 - \beta}{24(L+C)}, \left(\frac{(1 - \tilde{\rho}_w)^3}{4032c_0^2(1 - \beta)L^2(C+L)} \right)^{1/3} \right\},$$

$$1 > \beta \geq \tilde{\rho}_w.$$

We have

$$\begin{aligned} \mathcal{L}_{k+1} \leq & \left(1 + \frac{4\alpha^2CL}{n} + \frac{240\alpha^3c_0^2\tilde{\rho}_wCL^2(1-\beta)^2}{(1-\tilde{\rho}_w)^3} \right) \mathcal{L}_k - \frac{\alpha}{4} \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] + \frac{\alpha^2L(2C\sigma_f^* + \sigma^2)}{n} \\ & + \left[\frac{72c_0^2\tilde{\rho}_wL^2(1-\beta)^2}{(1-\tilde{\rho}_w)^3} + \frac{4L(L+2C)}{n(1-\beta)} \right] \alpha^3(2C\sigma_f^* + \sigma^2). \end{aligned}$$

Moreover, denote $\Delta_0 := f(\bar{x}_0) - f^*$, the term \mathcal{L}_0 can be upper bounded by

$$\begin{aligned} \mathcal{L}_0 &= \Delta_0 + \frac{3\alpha L^2}{n(1-\tilde{\rho}_w)} \mathcal{R}_0^x + \frac{12\alpha^3c_0\tilde{\rho}_wL^2}{n(1-\tilde{\rho}_w)^3} \mathcal{R}_0^y + \frac{48\alpha^3c_0^2\tilde{\rho}_wL^2}{n(1-\tilde{\rho}_w)^3} \|\nabla F(\mathbf{x}_0)\|^2 \\ &\leq 2\Delta_0 + \frac{9\alpha L^2 \|\Pi \mathbf{x}_0\|^2}{n(1-\tilde{\rho}_w)} + \frac{27\alpha^3c_0^2L^2(1-\beta)^2(2C\sigma_f^* + \sigma^2)}{(1-\tilde{\rho}_w)^2} + \frac{72\alpha^3c_0^2L^2 \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{n(1-\tilde{\rho}_w)^3}. \end{aligned} \quad (16)$$

Proof. See Appendix A.8. □

In summary, Lemma 3.4 inspires us to derive the recursions of the corresponding error terms, outlined in Lemmas 3.5-3.7, and to consider the Lyapunov function in (15). Moreover, due to the fact that $\|\tilde{\Pi}\tilde{W}\|_2$ may be greater than one, the consensus error terms $\mathbb{E}[\|\Pi \mathbf{x}_k\|^2]$ and $\mathbb{E}[\|\Pi \mathbf{y}_k\|^2]$ are proved to be R -linear sequences with ‘‘additional errors’’ similar as those in [39]. Notably, the analysis of DSMT differs from [39] mainly due to the need for dealing with the stochastic gradients under the general variance condition (2) and the momentum variable \mathbf{z}_k .

4 Main Results

In this section, we present the main results, which outline the convergence rates and the transient times of the DSMT method for minimizing smooth objective functions with or without the PL condition. These results are detailed in Subsections 4.1 and 4.2, respectively. The formal definitions of transient times $K_{\text{Transient}}^{(\text{NCVX})}$ and $K_{\text{Transient}}^{(\text{PL})}$ under general smooth objective functions and smooth objective functions satisfying the PL condition are defined, as follows:

$$K_{\text{Transient}}^{(\text{NCVX})} := \inf_K \left\{ \min_{t=0,1,\dots,k-1} \mathbb{E} \left[\|\nabla f(\bar{x}_t)\|^2 \right] \leq \mathcal{O} \left(\frac{1}{\sqrt{nk}} \right), \forall k \geq K \right\} \quad (17a)$$

$$K_{\text{Transient}}^{(\text{PL})} := \inf_K \left\{ \frac{1}{n} \sum_{i=1}^n \mathbb{E} [f(x_{i,k}) - f^*] \leq \mathcal{O} \left(\frac{1}{nk} \right), \forall k \geq K \right\}. \quad (17b)$$

Note that $\mathcal{O}(1/\sqrt{nk})$ and $\mathcal{O}(1/(nk))$ represent the convergence rates of the centralized stochastic gradient method under the two considered conditions, respectively.

4.1 General Nonconvex Case

We begin by presenting the convergence result of DSMT for minimizing smooth nonconvex objective functions, as demonstrated in Theorem 4.1 below.

Theorem 4.1. *Let Assumptions 1.1-1.3 hold. For a given $K \geq 1$, let the stepsize α and the parameter β to satisfy*

$$\alpha \leq \min \left\{ \sqrt{\frac{(1 - \tilde{\rho}_w)^2}{240c_0^2L(L+C)}}, \frac{1 - \beta}{24(L+C)}, \left(\frac{(1 - \tilde{\rho}_w)^3}{4032c_0^2(1 - \beta)L^2(C+L)} \right)^{1/3}, \right.$$

$$\left. \sqrt{\frac{n}{8CLK}}, \left(\frac{(1 - \tilde{\rho}_w)^3}{480c_0^2CL^2(1 - \beta)^2K} \right)^{1/3} \right\}, \quad 1 > \beta \geq \tilde{\rho}_w.$$

Then, we have

$$\begin{aligned} \min_{k=0,1,\dots,K-1} \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] &\leq \frac{24\Delta_0}{\alpha K} + \frac{108L^2 \|\Pi \mathbf{x}_0\|^2}{n(1-\tilde{\rho}_w)K} + \frac{324c_0^2\alpha^2 L^2(1-\beta)^2(2C\sigma_f^* + \sigma^2)}{(1-\tilde{\rho}_w)^2 K} \\ &+ \frac{4\alpha L(2C\sigma_f^* + \sigma^2)}{n} + \frac{864c_0^2\alpha^2 L^2 \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{n(1-\tilde{\rho}_w)^3 K} \\ &+ \left[\frac{L(L+2C)}{n(1-\beta)} + \frac{18c_0^2\tilde{\rho}_w L^2(1-\beta)^2}{(1-\tilde{\rho}_w)^3} \right] 16\alpha^2(2C\sigma_f^* + \sigma^2). \end{aligned} \quad (18)$$

In addition, if we set

$$\alpha = \frac{1}{\sqrt{\frac{L(2C\sigma_f^* + \sigma^2)K}{3n\Delta_0} + \gamma}}, \quad \beta = 1 - \frac{1}{n^{1/3}}(1-\tilde{\rho}_w), \quad (19)$$

where

$$\gamma := \left(\frac{4032c_0^2 L^2(C+L)}{n^{1/3}(1-\tilde{\rho}_w)^2} \right)^{1/3} + \frac{24c_0 n^{1/3}(L+C)}{1-\tilde{\rho}_w} + \sqrt{\frac{8CLK}{n}} + \left(\frac{480c_0^2 CL^2 K}{n^{2/3}(1-\tilde{\rho}_w)} \right)^{1/3},$$

then

$$\begin{aligned} \min_{k=0,1,\dots,K-1} \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] &= \mathcal{O} \left(\sqrt{\frac{\Delta_0 L(2C\sigma_f^* + \sigma^2)}{nK}} + \sqrt{\frac{\Delta_0^2 CL}{nK}} + \frac{\Delta_0 (CL^2)^{1/3}}{n^{2/9}(1-\tilde{\rho}_w)^{1/3} K^{2/3}} \right. \\ &\left. + \frac{n^{1/3} \Delta_0 (L+C) + \|\Pi \mathbf{x}_0\|^2 L^2/n}{(1-\tilde{\rho}_w)K} + \frac{n^{1/3} \Delta_0 L}{K^2} + \frac{\Delta_0 L \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{(1-\tilde{\rho}_w)^3 (2C\sigma_f^* + \sigma^2) K^2} \right), \end{aligned} \quad (20)$$

where $\mathcal{O}(\cdot)$ hides the numerical constants.

Proof. We first state Lemma 4.1 from [26, Lemma 6] that provides a direct link connecting Lemma 3.8 to the desired results.

Lemma 4.1. *If there exist constants $a, b, c \geq 0$ and non-negative sequences $\{s_t\}, \{q_t\}$ such that for any t , we have*

$$s_{t+1} \leq (1+a)s_t - bq_t + c,$$

then it holds that

$$\min_{t=0,1,\dots,T-1} q_t \leq \frac{(1+a)^T}{bT} s_0 + \frac{c}{b}.$$

Applying Lemma 4.1 to Lemma 3.8 leads to

$$\begin{aligned} \min_{k=0,1,\dots,K-1} \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] &\leq \frac{4 \left(1 + \frac{4\alpha^2 CL}{n} + \frac{240\alpha^3 c_0^2 \tilde{\rho}_w CL^2 (1-\beta)^2}{(1-\tilde{\rho}_w)^3} \right)^K}{\alpha K} \mathcal{L}_0 + \frac{4\alpha L(2C\sigma_f^* + \sigma^2)}{n} \\ &+ \left[\frac{L(L+2C)}{n(1-\beta)} + \frac{18c_0^2\tilde{\rho}_w L^2(1-\beta)^2}{(1-\tilde{\rho}_w)^3} \right] 16\alpha^2(2C\sigma_f^* + \sigma^2). \end{aligned} \quad (21)$$

Letting

$$\alpha \leq \min \left\{ \sqrt{\frac{n}{8CLK}}, \left(\frac{(1-\tilde{\rho}_w)^3}{480c_0^2 CL^2 (1-\beta)^2 K} \right)^{1/3} \right\}$$

yields

$$\left(1 + \frac{4\alpha^2 CL}{n} + \frac{240\alpha^3 c_0^2 \tilde{\rho}_w CL^2 (1-\beta)^2}{(1-\tilde{\rho}_w)^3} \right)^K \leq \exp \left\{ \left[\frac{4\alpha^2 CL}{n} + \frac{240\alpha^3 c_0^2 \tilde{\rho}_w CL^2 (1-\beta)^2}{(1-\tilde{\rho}_w)^3} \right] K \right\} \leq \exp(1). \quad (22)$$

Substituting (22) and the expression of \mathcal{L}_0 in (16) into (21) yields (18).

Set α as in (19), then

$$\begin{aligned} \frac{\Delta_0}{\alpha K} &= \sqrt{\frac{\Delta_0 L(2C\sigma_f^* + \sigma^2)}{3nK}} + \Delta_0 \sqrt{\frac{8CL}{nK}} + \frac{\Delta_0(480c_0^2 CL^2)^{1/3}}{n^{2/9}(1-\tilde{\rho}_w)^{1/3} K^{2/3}} \\ &\quad + \frac{\Delta_0[4032c_0^2 L^2(C+L)]^{1/3}}{n^{1/9}(1-\tilde{\rho}_w)^{2/3} K} + \frac{24c_0\Delta_0 n^{1/3}(L+C)}{(1-\tilde{\rho}_w)K}, \\ \frac{4\alpha L(2C\sigma_f^* + \sigma^2)}{n} &\leq 4\sqrt{\frac{3L(2C\sigma_f^* + \sigma^2)\Delta_0}{nK}}, \quad \alpha^2 \leq \frac{3n\Delta_0}{L(2C\sigma_f^* + \sigma^2)K}. \end{aligned} \quad (23)$$

Substituting (23) into (18) yields the desired result (20). \square

Subsequently, we compute the transient time required for DSMT to achieve comparable performance to the centralized SGD method when minimizing smooth objective functions, as detailed in Corollary 4.2 below.

Corollary 4.2. *Let Assumptions 1.1-1.3 hold. Set the stepsize α and the parameter β to satisfy (19). Then the transient time $K_{\text{Transient}}^{(\text{NCVX})}$ required for DSMT method to achieve comparable performance as the centralized SGD method behaves as*

$$K_{\text{Transient}}^{(\text{NCVX})} = \max \left\{ \mathcal{O} \left(\frac{n^{5/3}}{1-\lambda} \right), \mathcal{O} \left(\frac{\|\Pi \mathbf{x}_0\|^4}{(1-\lambda)n} \right), \mathcal{O} \left(\frac{n^{1/3} \left(\sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2 \right)^{2/3}}{1-\lambda} \right) \right\}. \quad (24)$$

In addition, if we initialize all agents at the same solution and assume that $\sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2 = \mathcal{O}(n^2)$, the transient time $K_{\text{Transient}}^{(\text{NCVX})}$ stated in (24) further reduces to

$$K_{\text{Transient}}^{(\text{NCVX})} = \mathcal{O} \left(\frac{n^{5/3}}{1-\lambda} \right).$$

Remark 4.1. *It is worth noting that initializing all the agents at the same solution and assuming $\sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2 = \mathcal{O}(n^2)$ are mild conditions. In practice, the term $\sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2$ often behaves as $\mathcal{O}(n)$ assuming \bar{x}_0 is fixed.*

Proof. We hide the constants that are independent of n and $(1-\lambda)$ in the following. From (20), we have

$$\begin{aligned} \min_{k=0,1,\dots,K-1} \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] &= \mathcal{O} \left(\frac{1}{\sqrt{nK}} \right) \mathcal{O} \left(1 + \frac{n^{5/(18)}}{(1-\tilde{\rho}_w)^{1/3} K^{1/6}} + \frac{n^{5/6}}{(1-\tilde{\rho}_w)\sqrt{K}} \right. \\ &\quad \left. + \frac{\|\Pi \mathbf{x}_0\|^2}{(1-\tilde{\rho}_w)\sqrt{nK}} + \frac{n^{5/6}}{K^{3/2}} + \frac{\sqrt{n} \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{(1-\tilde{\rho}_w)^3 K^{3/2}} \right). \end{aligned} \quad (25)$$

Noting that

$$1 - \tilde{\rho}_w = 1 - \sqrt{\frac{1}{1 + \sqrt{1 - \lambda^2}}} = \frac{\sqrt{1 - \lambda^2}}{\sqrt{1 + \sqrt{1 - \lambda^2}} \left(1 + \sqrt{1 + \sqrt{1 - \lambda^2}} \right)} \sim \mathcal{O} \left(\sqrt{1 - \lambda} \right). \quad (26)$$

According to the definition for transient times in (17a), we obtain the desired result (24) by invoking (25) and (26). \square

4.2 PL Condition

In this part, we will consider a specific nonconvex condition known as the Polyak-Łojasiewicz (PL) condition in Assumption 4.3. Overparameterized models often satisfy this condition. Notably, the strong convexity condition implies the PL condition [14].

Assumption 4.3. There exists $\mu > 0$, such that the aggregate function $f(x) = \frac{1}{n} \sum_{i=1}^n f_i(x)$ satisfies

$$2\mu (f(x) - f^*) \leq \|\nabla f(x)\|^2, \quad (27)$$

for all $x \in \mathbb{R}^p$, where $f^* := \inf_{x \in \mathbb{R}^p} f(x)$.

In light of Assumption 4.3, we construct a ‘‘contractive’’ recursion for the quantity $\mathbb{E}[f(\bar{d}_k) - f^*]$.

Lemma 4.4. Let Assumptions 1.1-1.3 and 4.3 hold, and let $K > 0$. Set \bar{d}_k as in (10) and let the stepsize α to satisfy $\alpha \leq \min\{\mu/(4LC), (1 - \beta)/(2L)\}$. We have

$$\begin{aligned} \mathbb{E}[f(\bar{d}_{k+1}) - f^*] &\leq \left(1 - \frac{\alpha\mu}{2}\right) \mathbb{E}[f(\bar{d}_k) - f^*] + \frac{\alpha L^2}{n} \mathbb{E}[\|\Pi \mathbf{x}_k\|^2] + \frac{\alpha^2 L(2C\sigma_f^* + \sigma^2)}{n} \\ &\quad + \frac{\alpha^3 \beta L(L+C)}{(1-\beta)^2} \mathbb{E}[\|\bar{z}_{k-1}\|^2]. \end{aligned}$$

Proof. See Appendix B.1. □

Thanks to Assumption 4.3, we are now able to show that \mathcal{L}_k is a Q -linear sequence with ‘‘additional errors’’ in Lemma 4.5. The derivation is similar to those in the proof of Lemma 3.8.

Lemma 4.5. Let Assumptions 1.1-1.3 and 4.3 hold, and let $k \geq 0$. Set the stepsize α and the parameter β to satisfy

$$\begin{aligned} \alpha &\leq \min \left\{ \frac{1-\beta}{3\mu}, \frac{1-\beta}{486c_0^2(L+C)}, \sqrt{\frac{(1-\tilde{\rho}_w)^2}{240c_0^2L(L+C)}}, \sqrt{\frac{1-\tilde{\rho}_w}{324c_0^2(L+C)^2}}, \sqrt{\frac{\mu(1-\beta)^2}{960c_0^2L(L+C)^2}} \right\}, \\ 1 &\geq \beta \geq 1 - \frac{11(1-\tilde{\rho}_w)}{12}. \end{aligned}$$

We have

$$\mathcal{L}_{k+1} \leq \left(1 - \frac{\alpha\mu}{4}\right) \mathcal{L}_k + \left[\frac{L}{n} + \frac{72\alpha c_0^2 \tilde{\rho}_w L^2 (1-\beta)^2}{(1-\tilde{\rho}_w)^3} + \frac{4\alpha L(L+2C)}{n(1-\beta)}\right] \alpha^2 (2C\sigma_f^* + \sigma^2). \quad (28)$$

Proof. See Appendix B.2. □

We can obtain the convergence rate of DSMT in Theorem 4.2 by noting another Lyapunov function \mathcal{H}_k :

$$\mathcal{H}_k := \mathcal{R}_k^x + \frac{3\alpha^2 c_0 \tilde{\rho}_w}{(1-\tilde{\rho}_w)^2} \mathcal{R}_k^y + \frac{405\alpha^3 n c_0^2 (C+L)}{(1-\tilde{\rho}_w)^3} \mathbb{E}[\|\bar{z}_{k-1}\|^2] + \frac{54\alpha^2 c_0^2 \tilde{\rho}_w}{(1-\tilde{\rho}_w)^2} \mathbb{E}[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2]. \quad (29)$$

Theorem 4.2. Let Assumptions 1.1-1.3 and 4.3 hold. Set the stepsize α and the parameter β to satisfy

$$\alpha \leq \min \left\{ \frac{1-\beta}{3\mu}, \frac{1-\beta}{486c_0^2(L+C)}, \sqrt{\frac{(1-\tilde{\rho}_w)^2}{240c_0^2L(L+C)}}, \sqrt{\frac{1-\tilde{\rho}_w}{324c_0^2(L+C)^2}}, \sqrt{\frac{\mu(1-\beta)^2}{960c_0^2L(L+C)^2}} \right\}, \quad (30)$$

$$1 \geq \beta \geq 1 - \frac{11(1-\tilde{\rho}_w)}{12}. \quad (31)$$

We have

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_{i,k}) - f^*] &\leq 33\mathcal{L}_0 \left(1 - \frac{\alpha\mu}{4}\right)^k + \frac{5L\mathcal{H}_0}{n} \left(\frac{5+\beta}{6}\right)^k + \frac{20\alpha L(2C\sigma_f^* + \sigma^2)}{n\mu} \\ &\quad + \left[\frac{48L\mathcal{S}_1}{\mu} + \frac{504c_0^2(1+\mathcal{S}_2)(1-\beta)}{1-\tilde{\rho}_w} + \frac{32(6L+C)(1-\tilde{\rho}_w)}{n\mu(1-\beta)}\right] \frac{4\alpha^2 L(2C\sigma_f^* + \sigma^2)}{1-\tilde{\rho}_w}, \end{aligned}$$

where

$$\mathcal{S}_1 := \frac{18c_0^2(1-\beta)^2}{(1-\tilde{\rho}_w)^2} + \frac{(L+2C)(1-\tilde{\rho}_w)}{nL(1-\beta)}, \quad \mathcal{S}_2 := \frac{L}{n\mu} + \frac{L\mathcal{S}_1}{\mu},$$

and

$$\begin{aligned} \mathcal{H}_0 &= \mathcal{R}_0^x + \frac{3\alpha^2 c_0 \tilde{\rho}_w}{(1-\tilde{\rho}_w)^2} \mathcal{R}_0^y + \frac{54\alpha^2 c_0^2 \tilde{\rho}_w}{(1-\tilde{\rho}_w)^2} \|\nabla F(\mathbf{1}\bar{x}_0^\top)\|^2 \\ &\leq 2c_0 \|\Pi \mathbf{x}_0\|^2 + \frac{12\alpha^2 n c_0^2 (1-\beta)^2 (C\Delta_0 + 2C\sigma_f^* + \sigma^2)}{(1-\tilde{\rho}_w)^2} + \frac{72\alpha^2 c_0^2 \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{(1-\tilde{\rho}_w)^2}. \end{aligned} \quad (32)$$

Proof. See Appendix B.3. □

Based on Theorem 4.2, we can first compare the non-exponentially decreasing terms in the error bounds for different distributed stochastic gradient methods under a constant stepsize α ; see Table 2. It can be seen that the DSMT method exhibits the lowest static error when compared to the state-of-art methods by choosing $\beta = \mathcal{O}\left(1 - \sqrt{(1-\lambda)/n}\right)$, $n \geq 2$.

Method	Static Error
DSGD [37]	$\mathcal{O}\left(\frac{\alpha}{n} + \frac{\alpha^{3/2}}{1-\lambda} + \frac{\alpha^2}{(1-\lambda)^2}\right)$
ED [1]	$\mathcal{O}\left(\frac{\alpha}{n} + \frac{\alpha^2}{1-\lambda} + \frac{\alpha^4}{n(1-\lambda)^3}\right)$
DSGT [1]	$\mathcal{O}\left(\frac{\alpha}{n} + \frac{\alpha^2}{1-\lambda} + \frac{\alpha^4}{n(1-\lambda)^4}\right)$
DSMT	$\mathcal{O}\left(\frac{\alpha}{n} + \left[\frac{1-\beta}{1-\lambda} + \frac{1}{n(1-\beta)}\right] \alpha^2\right)$
DSMT $\beta = \mathcal{O}(1 - \sqrt{(1-\lambda)/n})$	$\mathcal{O}\left(\frac{\alpha}{n} + \frac{\alpha^2}{\sqrt{n(1-\lambda)}}\right)$

Table 2: Comparison of different methods regarding the non-exponentially decreasing terms for minimizing smooth objective functions satisfying the PL condition.

Table 2 also shows that proper choice of β improves the final error of DSMT, demonstrating the benefits of momentum acceleration in decentralized optimization methods.

By considering specific stepsize α and parameter β , we can write down the following convergence result of DSMT based on Theorem 4.2.

Corollary 4.6. *Let the conditions in Theorem 4.2 hold. Furthermore, let the stepsize α and parameter β satisfy*

$$\alpha = \alpha_1 = \frac{4}{\mu K} \ln \left(\frac{33n\mu^2 K \mathcal{L}_0}{(2C\sigma_f^* + \sigma^2)L} \right), \quad \beta = 1 - \frac{(1 - \tilde{\rho}_w)}{\sqrt{n}}, \quad n \geq 2, \quad (33)$$

for some given $K \geq 1$. Then, the final iterates $x_{i,K}$ for all $i \in \mathcal{N}$ satisfy

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E} [f(x_{i,K}) - f^*] &\leq 33\mathcal{L}_0 \exp\left(-\frac{\alpha\mu K}{4}\right) + \frac{5L\mathcal{H}_0}{n} \exp\left(-\frac{(1 - \tilde{\rho}_w)K}{\sqrt{n}}\right) \\ &+ \tilde{\mathcal{O}}\left(\frac{(2C\sigma_f^* + \sigma^2)}{nK} + \frac{(2C\sigma_f^* + \sigma^2)}{\sqrt{n(1-\lambda)K^2}}\right), \end{aligned} \quad (34)$$

where $\tilde{\mathcal{O}}(\cdot)$ hides the term $\ln\left(\frac{33n\mu^2 K \mathcal{L}_0}{(2C\sigma_f^* + \sigma^2)L}\right)$ and some constants that are independent of n and $(1-\lambda)$. In particular, we have

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E} [f(x_{i,K}) - f^*] &= \tilde{\mathcal{O}} \left\{ \left[\|\Pi \mathbf{x}_0\|^2 + \frac{(\Delta_0 + C\sigma_f^* + \sigma^2)}{\mu^2 K^2} + \frac{\sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2 L^2}{\mu^2 (1 - \tilde{\rho}_w)^2 K^2} \right] \exp\left(-\frac{(1 - \tilde{\rho}_w)K}{6\sqrt{n}}\right) \right. \\ &+ \left[\Delta_0 + \frac{L^2 \|\Pi \mathbf{x}_0\|^2}{n\mu(1 - \tilde{\rho}_w)K} + \frac{L^2(2C\sigma_f^* + \sigma^2)}{n\mu^3 K^3} + \frac{L^2 \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{\mu^3 (1 - \tilde{\rho}_w)^3 K^3} \right] \exp\left(-\frac{(1 - \tilde{\rho}_w)\mu K}{4L}\right) \\ &\left. + \frac{L(C\sigma_f^* + \sigma^2)}{n\mu^2 K} + \frac{L(C\sigma_f^* + \sigma^2)}{\sqrt{n}(1 - \tilde{\rho}_w)\mu^2 K^2} \right\}. \end{aligned} \quad (35)$$

Proof. Since $\beta = 1 - (1 - \tilde{\rho}_w)/\sqrt{n}$, the constants \mathcal{S}_1 and \mathcal{S}_2 become

$$\mathcal{S}_1 = \frac{18c_0^2}{n} + \frac{L + 2C}{\sqrt{n}L}, \quad \mathcal{S}_2 = \frac{L}{n\mu} + \frac{18c_0^2 L}{n\mu} + \frac{L + 2C}{\sqrt{n}\mu}.$$

Denote

$$\mathcal{S}_3 := \frac{48L}{\mu} \left(\frac{18c_0^2}{\sqrt{n}} + \frac{L+2C}{L} \right) + 504c_0^2(1 + \mathcal{S}_1) + \frac{32(6L+C)}{\mu} \sim \mathcal{O}(1),$$

we have

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E} [f(x_{i,K}) - f^*] &\leq 33\mathcal{L}_0 \exp\left(-\frac{\alpha\mu K}{4}\right) + \frac{5L\mathcal{H}_0}{n} \exp\left(-\frac{(1-\tilde{\rho}_w)K}{6\sqrt{n}}\right) \\ &\quad + \frac{20\alpha L(2C\sigma_f^* + \sigma^2)}{n\mu} + \frac{4\alpha^2 \mathcal{S}_3 L(2C\sigma_f^* + \sigma^2)}{\sqrt{n}(1-\tilde{\rho}_w)}. \end{aligned} \quad (36)$$

Let $\bar{\alpha}$ satisfy (30). It suffices to discuss the following two cases:

Case I: If $\bar{\alpha} < \alpha_1$, then we substitute $\bar{\alpha}$ into (36) and obtain

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E} [f(x_{i,K}) - f^*] &\leq 33\mathcal{L}_0 \exp\left(-\frac{\bar{\alpha}\mu K}{4}\right) + \frac{5L\mathcal{H}_0}{n} \exp\left(-\frac{(1-\tilde{\rho}_w)K}{6\sqrt{n}}\right) \\ &\quad + \frac{80L(2C\sigma_f^* + \sigma^2) \ln\left(\frac{33n\mu^2 K \mathcal{L}_0}{(2C\sigma_f^* + \sigma^2)L}\right)}{n\mu^2 K} + \frac{64L\mathcal{S}_3(2C\sigma_f^* + \sigma^2) \ln^2\left(\frac{33n\mu^2 K \mathcal{L}_0}{(2C\sigma_f^* + \sigma^2)L}\right)}{\sqrt{n}(1-\tilde{\rho}_w)\mu^2 K^2}. \end{aligned} \quad (37)$$

In this case, we have $\bar{\alpha} \sim \mathcal{O}((1-\tilde{\rho}_w)/[\sqrt{n}(L+C)])$. Then,

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E} [f(x_{i,K}) - f^*] &= \tilde{\mathcal{O}} \left\{ \left[\|\Pi \mathbf{x}_0\|^2 + \frac{(C\Delta_0 + C\sigma_f^* + \sigma^2)}{\mu^2 K^2} + \frac{\sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{\mu^2(1-\tilde{\rho}_w)^2 K^2} \right] \exp\left(-\frac{(1-\tilde{\rho}_w)K}{6\sqrt{n}}\right) \right. \\ &\quad + \left[\Delta_0 + \frac{L^2 \|\Pi \mathbf{x}_0\|^2}{n\mu(1-\tilde{\rho}_w)K} + \frac{L^2(2C\sigma_f^* + \sigma^2)}{n\mu^3 K^3} + \frac{L^2 \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{\mu^3(1-\tilde{\rho}_w)^3 K^3} \right] \exp\left(-\frac{(1-\tilde{\rho}_w)\mu K}{4\sqrt{n}(L+C)}\right) \\ &\quad \left. + \frac{L(C\sigma_f^* + \sigma^2)}{n\mu^2 K} + \frac{L(C\sigma_f^* + \sigma^2)}{\sqrt{n}(1-\tilde{\rho}_w)\mu^2 K^2} \right\}. \end{aligned} \quad (38)$$

Case II: If $\bar{\alpha} \geq \alpha_1$, then we substitute α_1 into (36) and obtain

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E} [f(x_{i,K}) - f^*] &\leq \frac{(2C\sigma_f^* + \sigma^2)L}{n\mu^2 K} + \frac{5L\mathcal{H}_0}{n} \exp\left(-\frac{(1-\tilde{\rho}_w)K}{6\sqrt{n}}\right) \\ &\quad + \frac{80L(2C\sigma_f^* + \sigma^2) \ln\left(\frac{33n\mu^2 K \mathcal{L}_0}{(2C\sigma_f^* + \sigma^2)L}\right)}{n\mu^2 K} + \frac{64L\mathcal{S}_3(2C\sigma_f^* + \sigma^2) \ln^2\left(\frac{33n\mu^2 K \mathcal{L}_0}{(2C\sigma_f^* + \sigma^2)L}\right)}{\sqrt{n}(1-\tilde{\rho}_w)\mu^2 K^2}. \end{aligned} \quad (39)$$

In this case, we have from (39) that


$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E} [f(x_{i,K}) - f^*] &= \tilde{\mathcal{O}} \left\{ \left[\|\Pi \mathbf{x}_0\|^2 + \frac{(\Delta_0 + C\sigma_f^* + \sigma^2)}{\mu^2 K^2} + \frac{\sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2 L^2}{\mu^2(1-\tilde{\rho}_w)^2 K^2} \right] \exp\left(-\frac{(1-\tilde{\rho}_w)K}{6\sqrt{n}}\right) \right. \\ &\quad \left. + \frac{L(C\sigma_f^* + \sigma^2)}{n\mu^2 K} + \frac{L(C\sigma_f^* + \sigma^2)}{\sqrt{n}(1-\tilde{\rho}_w)\mu^2 K^2} \right\}. \end{aligned} \quad (40)$$

Combining (37) and (39) yields the desired result (34). Combining (38) and (40) leads to the desired result (35). \square

Finally, we are able to derive the transient time of DSMT under smooth objective functions satisfying the PL condition.

Corollary 4.7. *Let the conditions in Corollary 4.6 hold. Then the transient time $K_{\text{Transient}}^{(PL)}$ required for DSMT to achieve comparable performance as centralized SGD method when minimizing smooth objective functions satisfying the PL condition behaves as*

$$K_{\text{Transient}}^{(PL)} = \tilde{\mathcal{O}} \left(\sqrt{\frac{n}{1-\lambda}} \right), \quad (41)$$

Figure 1: Illustration of ring graph topology with $n = 16$.

where $\tilde{\mathcal{O}}(\cdot)$ hides the following logarithm terms and other constants that are independent of n and $(1 - \lambda)$:

$$\max \left\{ \ln \left(n \|\Pi \mathbf{x}_0\|^2 K \right), \ln \left(\frac{33n\mu^2 K \mathcal{L}_0}{(2C\sigma_f^* + \sigma^2)L} \right), \ln \left(\frac{n}{K} \right), \ln \left(\frac{n \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{(1 - \lambda)^{3/2} K} \right), \ln \left(\frac{\|\Pi \mathbf{x}_0\|^2}{\sqrt{1 - \lambda}} \right) \right\}.$$

Proof. We hide the constants that are independent of n and $(1 - \lambda)$ in the following. Based on (35), we have

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E} [f(x_{i,K}) - f^*] &= \mathcal{O} \left(\frac{1}{nK} \right) \tilde{\mathcal{O}} \left\{ \left[nK \|\Pi \mathbf{x}_0\|^2 + \frac{n}{K} + \frac{n \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{(1 - \tilde{\rho}_w)^2 K} \right] \exp \left(-\frac{(1 - \tilde{\rho}_w)K}{\sqrt{n}} \right) \right. \\ &\quad \left. + \left[nK + \frac{\|\Pi \mathbf{x}_0\|^2}{(1 - \tilde{\rho}_w)} + \frac{1}{K^2} + \frac{n \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{(1 - \tilde{\rho}_w)^3 K^2} \right] \exp \left(-\frac{(1 - \tilde{\rho}_w)K}{\sqrt{n}} \right) + 1 + \frac{\sqrt{n}}{(1 - \tilde{\rho}_w)K} \right\}. \end{aligned} \quad (42)$$

Invoking the definition of $K_{\text{Transient}}^{(\text{PL})}$ and the relation $(1 - \tilde{\rho}_w) \sim \mathcal{O}(\sqrt{1 - \lambda})$ in (26) yields the desired result (41). \square

5 Simulations

This section presents two numerical examples that illustrate the performance of DSMT compared with the existing methods over a ring graph (Figure 1). We evaluate these algorithms on a strongly convex problem (43) (satisfying the PL condition) and a general nonconvex problem (44) to verify the theoretical findings. Overall, the results demonstrate that the incorporation of either the momentum tracking technique or Loopless Chebyshev Acceleration (LCA) enhances the performance of distributed stochastic gradient methods. Furthermore, the DSMT method outperforms existing methods due to the effective combination of these two techniques.

5.1 Logistic Regression

We consider a binary classification problem using logistic regression (43) on the CIFAR-10 [18] dataset. Each agent possesses a distinct local dataset $\mathcal{S}_i = \{(u_j, v_j)\}$ selected from the whole dataset \mathcal{S} . Here, the variable $u_j \in \mathbb{R}^p$ denotes the image input and $v_j \in \mathbb{R}$ represents the label. In particular, we consider a *heterogeneous* data setting, where data samples are sorted based on their labels and partitioned among the agents. The classifier can then be obtained by solving the following optimization problem using all the agents' local datasets $\mathcal{S}_i, i = 1, 2, \dots, n$:

$$\begin{aligned} \min_{x \in \mathbb{R}^p} f(x) &= \frac{1}{n} \sum_{i=1}^n f_i(x), \\ f_i(x) &:= \frac{1}{|\mathcal{S}_i|} \sum_{j \in \mathcal{S}_i} \log [1 + \exp(-x^\top u_j v_j)] + \frac{\rho}{2} \|x\|^2, \end{aligned} \quad (43)$$

where ρ is set as 0.2.

We evaluate the performance of several methods, including DSMT, DSGT [37], EDAS [11], DSGD [37], DSGT-HB [7], centralized SGD (CSGD), and centralized SGDM (CSGDM), in the context of classifying airplanes and trucks on the CIFAR-10 dataset using a constant step size. The results are presented in Figure 2. Additionally, to facilitate comparison, we implement the DSMT method without the LCA technique, denoted as DSMT_noLCA in subsequent discussions.




Figure 2: Comparison among DSMT, DSGT, EDAS, DSGD, CSGD, and CSGDM for solving Problem (43) on the CIFAR-10 dataset using a constant stepsize. The stepsize is set as $\alpha = 0.01$ for all the methods. The momentum parameter is set as $\beta = \tilde{\rho}_w$ for DSMT, DSMT_noLCA, and SGDM. The results are averaged over 10 repeated experiments.

The performance of decentralized methods shown in Figures 2a and 2b reveals that, as the network size decreases (resulting in a larger value of $(1 - \lambda)$), the methods tend to exhibit more comparable performance to centralized SGD. Among these decentralized methods, DSMT achieves the most similar accuracy compared to centralized methods, especially when the network topology is not well-connected (Figure 2a).

Comparing those decentralized methods incorporating momentum acceleration, they all behave more similar to CSGDM than CSGD, aligning with the arguments in Subsection 2.1. However, a comparison between DSMT_noLCA and DSGT-HB reveals the importance of appropriately employing the momentum technique to enhance the practical performance. Straightforward combination may not yield significant improvements, as seen from the performance comparison of DSGT and DSGT-HB in Figure 2. Such a comparison implies the effectiveness of the momentum tracking technique.

Comparing the performance of DSMT and DSMT_noLCA in Figure 2, it can be concluded that the LCA technique indeed accelerates the convergence process. Such an effect becomes more pronounced when the network topology degrades, as evidenced in the transition from Figure 2b to Figure 2a.

In summary, the incorporation of either the momentum tracking technique or LCA method proves effective in enhancing the practical performance of distributed stochastic gradient methods. Notably, the DSMT method outperforms the aforementioned alternatives by combining these two techniques. Such a comparison corroborates the theoretical findings in Section 4.

5.2 Nonconvex Logistic Regression

In this part, we consider a binary classification problem (44) classifying airplanes and trucks on the CIFAR-10 dataset. The parameter ω is set as $\omega = 0.05$ and $[x]_q$ denotes the q -element of $x \in \mathbb{R}^p$. The other settings follow the same as those in Subsection 5.1. We compared the aforementioned methods over ring graphs with $n = 100$ (Figure 3a) and

$n = 50$ (Figure 3b), respectively.

$$\begin{aligned} \min_{x \in \mathbb{R}^p} f(x) &= \frac{1}{n} \sum_{i=1}^n f_i(x), \\ f_i(x) &:= \frac{1}{|S_i|} \sum_{j \in S_i} \log [1 + \exp(-x^\top u_j v_j)] + \frac{\omega}{2} \sum_{q=1}^p \frac{[x]_q^2}{1 + [x]_q^2}. \end{aligned} \quad (44)$$

It can be seen from Figure 3 that solely applying momentum technique to DSGT does not enhance its performance. In Figure 3b, DSMT_noLCA demonstrates a slight improvement over DSGT-HB. However, DSMT consistently outperforms other methods, capitalizing on the combined efficacy of momentum tracking technique and LCA method. Notably, the difference between DSMT and EDAS becomes negligible as the number of iteration increases, suggesting that the application of LCA to EDAS could potentially enhance its convergence. Such a phenomenon inspires further research of interest.




Figure 3: Comparison among DSMT, DSGT, EDAS, DSGD, CSGD, and CSGDM for solving Problem (44) on the CIFAR-10 dataset using a constant stepsize. The stepsizes are set as 0.02 for all methods. The momentum parameter is set as $\beta = 1 - (1 - \tilde{\rho}_w)/n^{1/3}$ for DSMT, DSMT_noLCA, and SGDM. The results are averaged over 10 repeated experiments.

6 Conclusion

This paper focuses on addressing the distributed stochastic optimization problem over networked agents. The proposed algorithm, Distributed Stochastic Momentum Tracking (DSMT), leverages momentum tracking technique as well as the Loopless Chebyshev Acceleration (LCA) method to enhance the performance of distributed stochastic gradient methods over networks. In particular, DSMT shortens the transient times for decentralized stochastic gradient methods without requiring multiple communication per iteration for smooth objective functions with or without the Polyak-Łojasiewicz (PL) condition under the most general variance condition in distributed settings. Such a condition for stochastic gradients enables the wide range of application of DSMT. Experimental results also corroborate to such theoretical findings. The momentum tracking technique is also of independent interest and can potentially inspire future development of distributed optimization algorithms.

A Proofs for the General Nonconvex Case

A.1 Proof of Lemma 3.1

Define $\underline{x}_k^\top := \frac{1}{n} \mathbf{1}^\top [\tilde{\mathbf{x}}_k]_{n+1:2n}$ and $\bar{x}_k^\top := \frac{1}{2n} \mathbf{1}^\top \tilde{\mathbf{x}}_k$. From (9a), we have

$$\begin{aligned}\bar{x}_{k+1} &= \frac{1}{2} [(2 + \eta_w)\bar{x}_k - \eta_w \underline{x}_k] - \alpha \bar{y}_k, \\ \mathbf{x}_{k+1} &= (1 + \eta_w)W(\mathbf{x}_k - \alpha \mathbf{y}_k) - \eta_w([\tilde{\mathbf{x}}_k]_{n+1:2n} - \alpha \mathbf{y}_k), \\ [\tilde{\mathbf{x}}_{k+1}]_{n+1:2n} &= \mathbf{x}_k - \alpha \mathbf{y}_k.\end{aligned}\tag{45}$$

Therefore, it suffices to show $\bar{x}_k = \underline{x}_k = \bar{x}_k$ for any $k \geq 0$, which we prove by induction. When $k = 0$, $\bar{x}_0 = \underline{x}_0 = \bar{x}_0$ since $\tilde{\mathbf{x}}_0 = (\mathbf{x}_0)_\#$. Suppose $\bar{x}_k = \underline{x}_k = \bar{x}_k$ for some $k \geq 0$, then (45) leads to

$$\begin{aligned}\bar{x}_{k+1} &= \bar{x}_k - \alpha \bar{y}_k, \\ \bar{x}_{k+1} &= (1 + \eta_w)\bar{x}_k - \eta_w \underline{x}_k - \alpha \bar{y}_k = \bar{x}_k - \alpha \bar{y}_k, \\ \underline{x}_{k+1} &= \bar{x}_k - \alpha \bar{y}_k.\end{aligned}$$

Hence, we have $\bar{x}_k = \underline{x}_k = \bar{x}_k$ for any $k \geq 0$. Similar line of analysis can show that $\bar{y}_{k+1} = \bar{y}_{k+1} = \underline{y}_{k+1} = \bar{y}_k + \bar{z}_{k+1} - \bar{z}_k$. The update form for \bar{y}_k and the initialization $\mathbf{y}_0 = \mathbf{z}_0$ imply $\bar{y}_k = \bar{z}_k$ for any $k \geq 0$.

A.2 Proof of Lemma 3.2

Recall from Lemma 3.1 that $\bar{x}_{k+1} = \bar{x}_k - \alpha \bar{z}_k$. We first show $\bar{d}_{k+1} = \bar{d}_k - \alpha \bar{g}_k$. When $k = 0$, we have

$$\bar{d}_1 - \bar{d}_0 = \frac{1}{1-\beta} \bar{x}_1 - \frac{\beta}{1-\beta} \bar{x}_0 - \bar{x}_0 = \frac{1}{1-\beta} (\bar{x}_0 - \alpha \bar{z}_0) - \frac{1}{1-\beta} \bar{x}_0 = -\alpha \bar{g}_0,$$

where the last equality holds by initializing $\mathbf{z}_0 = (1-\beta)\mathbf{g}_0$.

For the case $k \geq 1$, we have

$$\bar{d}_{k+1} - \bar{d}_k = \frac{1}{1-\beta} (\bar{x}_{k+1} - \beta \bar{x}_k - \bar{x}_k + \beta \bar{x}_{k-1}) = \frac{1}{1-\beta} (-\alpha \bar{z}_k + \beta \alpha \bar{z}_{k-1}) = -\alpha \bar{g}_k.$$

Next we show $\bar{d}_k - \bar{x}_k = -\frac{\alpha\beta}{1-\beta} \bar{z}_{k-1}$, which is because for $k \geq 1$,

$$\bar{d}_k - \bar{x}_k = \frac{1}{1-\beta} \bar{x}_k - \frac{\beta}{1-\beta} \bar{x}_{k-1} - \bar{x}_k = \frac{\beta}{1-\beta} (\bar{x}_k - \bar{x}_{k-1}) = -\frac{\alpha\beta}{1-\beta} \bar{z}_{k-1}.\tag{46}$$

Due to $\mathbf{z}_{-1} = \mathbf{0}$ and $\bar{d}_0 = \bar{x}_0$, the relation (46) also holds for $k = 0$.

A.3 Proof of Lemma 3.3

Given that $\bar{x}_k = \bar{d}_k + \alpha\beta\bar{z}_{k-1}/(1-\beta)$, $k \geq 0$ from Lemma 3.2, we have

$$f(\bar{x}_k) - f^* \leq f(\bar{d}_k) - f^* + \frac{\alpha\beta}{1-\beta} \langle \nabla f(\bar{d}_k) - \nabla f(\bar{x}_k) + \nabla f(\bar{x}_k), \bar{z}_{k-1} \rangle + \frac{\alpha^2\beta^2 L}{2(1-\beta)^2} \|\bar{z}_{k-1}\|^2\tag{47}$$

$$\leq f(\bar{d}_k) - f^* + \frac{\alpha\beta}{2(1-\beta)} \|\nabla f(\bar{x}_k)\|^2 + \frac{\alpha\beta[1+3\alpha\beta L/(1-\beta)]}{2(1-\beta)} \|\bar{z}_{k-1}\|^2,\tag{48}$$

where we invoke $\|\nabla f(\bar{x}_k) - \nabla f(\bar{d}_k)\| \leq \alpha\beta L \|\bar{z}_{k-1}\|/(1-\beta)$ for $k \geq 0$. Taking the full expectation on both sides yields the desired result.

A.4 Proof of Lemma 3.4

In light of Lemma 3.2 that $\bar{d}_{k+1} = \bar{d}_k - \alpha \bar{g}_k$ for $k \geq 0$, we have from the descent lemma and Assumption 1.2 that

$$\begin{aligned}
\mathbb{E} [f(\bar{d}_{k+1}) | \mathcal{F}_k] &\leq f(\bar{d}_k) - \alpha \left\langle \nabla f(\bar{d}_k) - \nabla f(\bar{x}_k), \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\rangle \\
&\quad - \alpha \left\langle \nabla f(\bar{x}_k), \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\rangle + \frac{L}{2} \mathbb{E} [\|\alpha \bar{g}_k\|^2 | \mathcal{F}_k] \\
&\leq f(\bar{d}_k) + \alpha \|\nabla f(\bar{d}_k) - \nabla f(\bar{x}_k)\|^2 + \frac{\alpha}{4} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\|^2 - \frac{\alpha(1-\alpha L)}{2} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\|^2 \\
&\quad - \frac{\alpha}{2} \|\nabla f(\bar{x}_k)\|^2 + \frac{\alpha}{2} \left\| \nabla f(\bar{x}_k) - \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\|^2 + \frac{\alpha^2 L}{2} \mathbb{E} \left[\left\| \bar{g}_k - \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\|^2 \middle| \mathcal{F}_k \right] \\
&\leq f(\bar{d}_k) + \alpha L^2 \|\bar{d}_k - \bar{x}_k\|^2 - \frac{\alpha}{2} \|\nabla f(\bar{x}_k)\|^2 - \frac{\alpha}{8} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\|^2 + \frac{\alpha L^2}{2n} \sum_{i=1}^n \|\bar{x}_k - x_{i,k}\|^2 \\
&\quad + \frac{\alpha^2 L}{n} \left[\frac{C}{n} \sum_{i=1}^n (f_i(x_{i,k}) - f_i^*) + \sigma^2 \right],
\end{aligned} \tag{49}$$

where we let $\alpha \leq 1/(4L)$ and invoke Assumption 1.2 for the last inequality.

Noting that

$$\begin{aligned}
f_i(x_{i,k}) - f_i^* &\leq f_i(\bar{x}_k) - f_i^* + \langle \nabla f_i(\bar{x}_k), x_{i,k} - \bar{x}_k \rangle + \frac{L}{2} \|x_{i,k} - \bar{x}_k\|^2 \\
&\leq f_i(\bar{x}_k) - f_i^* + \frac{1}{2L} \|\nabla f_i(\bar{x}_k)\|^2 + L \|x_{i,k} - \bar{x}_k\|^2 \\
&\leq 2(f_i(\bar{x}_k) - f_i^*) + L \|x_{i,k} - \bar{x}_k\|^2,
\end{aligned} \tag{50}$$

we have

$$\frac{1}{n} \sum_{i=1}^n (f_i(x_{i,k}) - f_i^*) \leq 2(f(\bar{x}_k) - f^*) + 2\sigma_f^* + \frac{L}{n} \|\mathbf{x}_k - \mathbf{1}\bar{x}_k\|^2. \tag{51}$$

In light of the relation $\bar{d}_k - \bar{x}_k = -\alpha\beta\bar{z}_{k-1}/(1-\beta)$ and $\mathbf{z}_{-1} = \mathbf{0}$, we have

$$\|\bar{d}_k - \bar{x}_k\|^2 = \frac{\alpha^2\beta^2}{(1-\beta)^2} \|\bar{z}_{k-1}\|^2, k \geq 0. \tag{52}$$

Combing (49)-(52), taking the full expectation, letting $\alpha \leq 1/(2C)$, and invoking Lemma 3.3 yield the desired result for $k \geq 0$.

A.5 Proof of Lemma 3.5

According to Lemma 3.1, we have

$$\bar{z}_k = \beta\bar{z}_{k-1} + (1-\beta) \left(\bar{g}_k - \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right) + (1-\beta) \left[\frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) - \nabla f(\bar{x}_k) \right] + (1-\beta) \nabla f(\bar{x}). \tag{53}$$

Then, by Assumption 1.2 and Jensen's inequality, we have

$$\begin{aligned}
\mathbb{E} \left[\|\bar{z}_k\|^2 \middle| \mathcal{F}_k \right] &\leq \beta \|\bar{z}_{k-1}\|^2 + \frac{(1-\beta)^2}{n} \left[\frac{C}{n} \sum_{i=1}^n (f_i(x_{i,k}) - f_i^*) + \sigma^2 \right] + 2(1-\beta) \|\nabla f(\bar{x}_k)\|^2 \\
&\quad + \frac{2(1-\beta)L^2}{n} \|\mathbf{x}_k - \mathbf{1}\bar{x}_k^\top\|^2 \\
&\leq \beta \|\bar{z}_{k-1}\|^2 + \frac{2C(1-\beta)^2}{n} [f(\bar{x}_k) - f^*] + 2(1-\beta) \|\nabla f(\bar{x}_k)\|^2 + \frac{(1-\beta)^2(2C\sigma_f^* + \sigma^2)}{n} \\
&\quad + \frac{2(1-\beta)L}{n} \left(L + \frac{C(1-\beta)}{n} \right) \|\mathbf{x}_k - \mathbf{1}\bar{x}_k^\top\|^2.
\end{aligned} \tag{54}$$

Taking the full expectation on both sides and invoking Lemma 3.3 yield the desired result.

A.6 Proof of Lemma 3.6

First consider

$$\begin{aligned}
\mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top) &= \beta \mathbf{z}_{k-1} + (1-\beta) \mathbf{g}_k - \nabla F(\mathbf{1}\bar{x}_k^\top) \\
&= \beta [\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)] + (1-\beta) [\mathbf{g}_k - \nabla F(\mathbf{x}_k)] + (1-\beta) [\nabla F(\mathbf{x}_k) - \nabla F(\mathbf{1}\bar{x}_k^\top)] \\
&\quad - \beta [\nabla F(\mathbf{1}\bar{x}_k^\top) - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)], \quad k \geq 1.
\end{aligned} \tag{55}$$

From (55) and Jensen's inequality, we have

$$\begin{aligned}
\mathbb{E} \left[\|\mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top)\|^2 \middle| \mathcal{F}_k \right] &\leq \beta \|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 + 2(1-\beta) \|\nabla F(\mathbf{x}_k) - \nabla F(\mathbf{1}\bar{x}_k^\top)\|^2 \\
&\quad + \frac{2\beta^2}{1-\beta} \|\nabla F(\mathbf{1}\bar{x}_k^\top) - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 + (1-\beta)^2 \mathbb{E} \left[\|\mathbf{g}_k - \nabla F(\mathbf{x}_k)\|^2 \middle| \mathcal{F}_k \right] \\
&\leq \beta \|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 + \frac{2\alpha^2\beta^2nL^2}{1-\beta} \|\bar{z}_{k-1}\|^2 + 2(1-\beta)L^2 \|\Pi\mathbf{x}_k\|^2 \\
&\quad + n(1-\beta)^2(2C\sigma_f^* + \sigma^2) + n(1-\beta)^2C \left\{ 2[f(\bar{x}_k) - f^*] + \frac{L}{n} \|\Pi\mathbf{x}_k\|^2 \right\},
\end{aligned} \tag{56}$$

where Assumption 1.2 and the upper bound (57) are invoked for the last inequality. The inequality (57) holds due to Lemma 3.1 that $\bar{x}_k = \bar{x}_{k-1} - \alpha\bar{z}_{k-1}$

$$\mathbb{E} \left[\|\nabla F(\mathbf{1}\bar{x}_k^\top) - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right] \leq nL^2 \mathbb{E} \left[\|\bar{x}_k - \bar{x}_{k-1}\|^2 \right] \leq \alpha^2 nL^2 \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right]. \tag{57}$$

Taking full expectation on both sides of (56), and substituting the result of Lemmas 3.3 yield that

$$\begin{aligned}
\mathbb{E} \left[\|\mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top)\|^2 \right] &\leq \beta \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right] + \frac{2\alpha^2\beta^2nL^2}{1-\beta} \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] + 2(1-\beta)L^2 \mathbb{E} \left[\|\Pi\mathbf{x}_k\|^2 \right] \\
&\quad + n(1-\beta)^2C \left\{ 2\mathbb{E} [f(\bar{d}_k) - f^*] + \frac{2\alpha\beta}{1-\beta} \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] + \frac{2\alpha\beta}{1-\beta} \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] + \frac{L}{n} \mathbb{E} \left[\|\Pi\mathbf{x}_k\|^2 \right] \right\} \\
&\quad + n(1-\beta)^2(2C\sigma_f^* + \sigma^2) \\
&\leq \beta \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right] + 2\alpha n \left(\frac{\alpha L^2}{1-\beta} + (1-\beta)C \right) \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] + 2(1-\beta)L(L+C) \mathbb{E} \left[\|\Pi\mathbf{x}_k\|^2 \right] \\
&\quad + n(1-\beta)^2(2C\sigma_f^* + \sigma^2) + 2n(1-\beta)^2C \mathbb{E} [f(\bar{d}_k) - f^*] + 2\alpha n(1-\beta)C \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right], \quad k \geq 1.
\end{aligned}$$

Letting $\alpha \leq (1-\beta)/L$ leads to the desired result for $k \geq 1$. For the case $k = 0$, recall that $\mathbf{x}_{-1} = \mathbf{x}_0$, $\mathbf{z}_{-1} = \mathbf{0}$ and $\mathbf{z}_0 = (1-\beta)\mathbf{g}_0$, then

$$\begin{aligned}
\mathbf{z}_0 - \nabla F(\mathbf{1}\bar{x}_0^\top) &= \beta [\mathbf{z}_{-1} - \nabla F(\mathbf{1}\bar{x}_{-1}^\top)] + (1-\beta) [\mathbf{g}_0 - \nabla F(\mathbf{x}_0)] + (1-\beta) [\nabla F(\mathbf{x}_0) - \nabla F(\mathbf{1}\bar{x}_0^\top)] \\
&\quad - \beta [\nabla F(\mathbf{1}\bar{x}_0^\top) - \nabla F(\mathbf{1}\bar{x}_{-1}^\top)].
\end{aligned} \tag{58}$$

Since $[\nabla F(\mathbf{1}\bar{x}_0^\top) - \nabla F(\mathbf{1}\bar{x}_{-1}^\top)] = \mathbf{0}$ due to $\mathbf{x}_{-1} = \mathbf{x}_0$, similar to the derivations of (56), we have

$$\begin{aligned} \mathbb{E} \left[\|\mathbf{z}_0 - \nabla F(\mathbf{1}\bar{x}_0^\top)\|^2 \right] &\leq \beta \mathbb{E} \left[\|\mathbf{z}_{-1} - \nabla F(\mathbf{1}\bar{x}_{-1}^\top)\|^2 \right] + (1 - \beta)L^2 \|\Pi \mathbf{x}_0\|^2 \\ &\quad + n(1 - \beta)^2(2C\sigma_f^* + \sigma^2) + n(1 - \beta)^2C \left\{ 2\mathbb{E}[f(\bar{x}_0) - f^*] + \frac{L}{n} \|\Pi \mathbf{x}_0\|^2 \right\} \\ &\leq \beta \mathbb{E} \left[\|\mathbf{z}_{-1} - \nabla F(\mathbf{1}\bar{x}_{-1}^\top)\|^2 \right] + (1 - \beta)L(L + C) \|\Pi \mathbf{x}_0\|^2 \\ &\quad + n(1 - \beta)^2(2C\sigma_f^* + \sigma^2) + 2n(1 - \beta)^2C\mathbb{E}[f(\bar{d}_0) - f^*], \end{aligned} \quad (59)$$

where we invoke that $\bar{x}_{-1} = \bar{x}_0 = \bar{d}_0$. As a consequence, combining the recursions for $k = 0$ and $k \geq 1$ yields the desired result.

A.7 Proof of Lemma 3.7

Note that $\tilde{\Pi}\tilde{W} = \tilde{\Pi}\tilde{W}\tilde{\Pi}$. Then,

$$\begin{aligned} \mathbb{E} \left[\left\| \tilde{\Pi}\tilde{\mathbf{x}}_{k+1} \right\|^2 \middle| \mathcal{F}_k \right] &= \mathbb{E} \left[\left\| \tilde{\Pi}\tilde{W}\tilde{\Pi}\tilde{\mathbf{x}}_k - \alpha\tilde{\Pi}\tilde{W}\tilde{\Pi}(\mathbf{y}_k)_\# \right\|^2 \middle| \mathcal{F}_k \right] \\ &\leq \frac{1}{q} \left\| \tilde{\Pi}\tilde{W}\tilde{\Pi}\tilde{\mathbf{x}}_k \right\|^2 + \frac{\alpha^2 c_0 \tilde{\rho}_w^2}{1 - q} \mathbb{E} \left[\|\Pi \mathbf{y}_k\|^2 \middle| \mathcal{F}_k \right], \end{aligned} \quad (60)$$

where we invoke Jensen's inequality for some $q > 0$ (to be determined later) and Lemma 2.1 for the last inequality.

For the term $\|\tilde{\Pi}\tilde{W}\tilde{\Pi}\tilde{\mathbf{x}}_k\|^2$, we have

$$\frac{1}{q} \mathbb{E} \left[\left\| \tilde{\Pi}\tilde{W}\tilde{\Pi}\tilde{\mathbf{x}}_k \right\|^2 \middle| \mathcal{F}_{k-1} \right] \leq \frac{1}{q^2} \left\| \tilde{\Pi}\tilde{W}^2\tilde{\Pi}\tilde{\mathbf{x}}_{k-1} \right\|^2 + \frac{\alpha^2 c_0 \tilde{\rho}_w^4}{q(1 - q)} \mathbb{E} \left[\|\Pi \mathbf{y}_{k-1}\|^2 \middle| \mathcal{F}_{k-1} \right]. \quad (61)$$

Choosing $q = \tilde{\rho}_w$ in (60) and (61) and repeating such procedures lead to

$$\begin{aligned} \mathbb{E} \left[\left\| \tilde{\Pi}\tilde{\mathbf{x}}_k \right\|^2 \right] &\leq \frac{1}{\tilde{\rho}_w^k} \left\| \tilde{\Pi}\tilde{W}^k\tilde{\Pi}\tilde{\mathbf{x}}_0 \right\|^2 + \frac{\alpha^2 c_0 \tilde{\rho}_w}{1 - \tilde{\rho}_w} \sum_{t=0}^{k-1} \tilde{\rho}_w^{k-t} \mathbb{E} \left[\|\Pi \mathbf{y}_t\|^2 \right] \\ &\leq c_0 \tilde{\rho}_w^k \|\Pi \mathbf{x}_0\|^2 + \frac{\alpha^2 c_0 \tilde{\rho}_w}{1 - \tilde{\rho}_w} \sum_{t=0}^{k-1} \tilde{\rho}_w^{k-t} \mathbb{E} \left[\|\Pi \mathbf{y}_t\|^2 \right] := \mathcal{R}_k^x, \end{aligned} \quad (62)$$

where we apply Lemma 2.1 for the last inequality. For the term \mathcal{R}_k^x , we have

$$\mathcal{R}_{k+1}^x = \tilde{\rho}_w \mathcal{R}_k^x + \frac{\alpha^2 c_0 \tilde{\rho}_w^2}{1 - \tilde{\rho}_w} \mathbb{E} \left[\|\Pi \mathbf{y}_k\|^2 \right] \leq \tilde{\rho}_w \mathcal{R}_k^x + \frac{\alpha^2 c_0 \tilde{\rho}_w}{1 - \tilde{\rho}_w} \mathbb{E} \left[\left\| \tilde{\Pi}\tilde{\mathbf{y}}_k \right\|^2 \right]. \quad (63)$$

Similarly, we can construct a sequence $\{\mathcal{R}_k^y\}$ that bounds the term $\mathbb{E}[\|\tilde{\Pi}\tilde{\mathbf{y}}\|^2]$ as follows.

Relation (55) guides us to rewrite $(\mathbf{z}_{k+1} - \mathbf{z}_k)$ as in (64), so that we can take advantage of the additional coefficient $(1 - \beta)$ and reduce the impact of the data heterogeneity due to $\sum_{i=1}^n \|\nabla f_i(x_{i,k})\|^2$ as mentioned in Subsection 2.1.

$$\begin{aligned} \mathbf{z}_{k+1} - \mathbf{z}_k &= -(1 - \beta)(\mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top)) + (1 - \beta)[\mathbf{g}_{k+1} - \nabla F(\mathbf{x}_{k+1})] \\ &\quad + (1 - \beta)[\nabla F(\mathbf{x}_{k+1}) - \nabla F(\mathbf{1}\bar{x}_{k+1}^\top)] + (1 - \beta)[\nabla F(\mathbf{1}\bar{x}_{k+1}^\top) - \nabla F(\mathbf{1}\bar{x}_k^\top)]. \end{aligned} \quad (64)$$

The above relation follows from (9c). Therefore, we can rewrite the update of $\tilde{\mathbf{y}}$ as follows.

$$\begin{aligned} \tilde{\mathbf{y}}_{k+1} &= \tilde{W}\tilde{\mathbf{y}}_k - (1 - \beta)\tilde{W}[\mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top)]_\# + (1 - \beta)\tilde{W}[\mathbf{g}_{k+1} - \nabla F(\mathbf{x}_{k+1})]_\# \\ &\quad + (1 - \beta)\tilde{W}[\nabla F(\mathbf{x}_{k+1}) - \nabla F(\mathbf{1}\bar{x}_{k+1}^\top)]_\# + (1 - \beta)\tilde{W}[\nabla F(\mathbf{1}\bar{x}_{k+1}^\top) - \nabla F(\mathbf{1}\bar{x}_k^\top)]_\#. \end{aligned} \quad (65)$$

Denote $\mathcal{A} := \tilde{\mathbf{y}}_k - (1-\beta)[\mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top)]_{\#} + (1-\beta)[\nabla F(\mathbf{x}_{k+1}) - \nabla F(\mathbf{1}\bar{x}_{k+1}^\top)]_{\#} + (1-\beta)[\nabla F(\mathbf{1}\bar{x}_{k+1}^\top) - \nabla F(\mathbf{1}\bar{x}_k^\top)]_{\#}$. We then have

$$\begin{aligned} \mathbb{E} \left[\left\| \tilde{\Pi} \tilde{\mathbf{y}}_{k+1} \right\|^2 \middle| \mathcal{F}_{k+1} \right] &= \left\| \tilde{\Pi} \tilde{W} \tilde{\Pi} \mathcal{A} \right\|^2 + (1-\beta)^2 \mathbb{E} \left[\left\| \tilde{\Pi} \tilde{W} \tilde{\Pi} [\mathbf{g}_{k+1} - \nabla F(\mathbf{x}_{k+1})]_{\#} \right\|^2 \middle| \mathcal{F}_{k+1} \right] \\ &\quad + 2\mathbb{E} \left[\left\langle \tilde{\Pi} \tilde{W} \tilde{\Pi} \mathcal{A}, (1-\beta) \tilde{\Pi} \tilde{W} \tilde{\Pi} [\mathbf{g}_{k+1} - \nabla F(\mathbf{x}_{k+1})]_{\#} \right\rangle \middle| \mathcal{F}_{k+1} \right] \\ &\leq \frac{1}{\tilde{\rho}_w} \left\| \tilde{\Pi} \tilde{W} \tilde{\Pi} \tilde{\mathbf{y}}_k \right\|^2 + \frac{3(1-\beta)^2 c_0 \tilde{\rho}_w^2}{1-\tilde{\rho}_w} \left[\left\| \mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top) \right\|^2 + L^2 \|\Pi \mathbf{x}_{k+1}\|^2 + \alpha^2 n L^2 \|\bar{z}_k\|^2 \right] \\ &\quad + n c_0 \tilde{\rho}_w^2 (1-\beta)^2 (2C\sigma_f^* + \sigma^2) + n c_0 \tilde{\rho}_w^2 C (1-\beta)^2 \left\{ 2[f(\bar{x}_{k+1}) - f^*] + \frac{L}{n} \|\Pi \mathbf{x}_{k+1}\|^2 \right\}, \end{aligned} \quad (66)$$

where we invoke Lemma 2.1, Assumptions 1.1 and 1.2, and (51) for the last inequality. Similarly, we can obtain the recursion for $\mathbb{E}[\|\tilde{\Pi} \tilde{W} \tilde{\Pi} \tilde{\mathbf{y}}_k\|^2 | \mathcal{F}_k]$:

$$\begin{aligned} \frac{1}{\tilde{\rho}_w} \mathbb{E} \left[\left\| \tilde{\Pi} \tilde{W} \tilde{\Pi} \tilde{\mathbf{y}}_k \right\|^2 \middle| \mathcal{F}_k \right] &\leq \frac{1}{\tilde{\rho}_w^2} \left\| \tilde{\Pi} \tilde{W}^2 \tilde{\Pi} \tilde{\mathbf{y}}_{k-1} \right\|^2 + n c_0 \tilde{\rho}_w^3 (1-\beta)^2 (2C\sigma_f^* + \sigma^2) \\ &\quad + \frac{3(1-\beta)^2 c_0 \tilde{\rho}_w^3}{1-\tilde{\rho}_w} \left[\left\| \mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top) \right\|^2 + L^2 \|\Pi \mathbf{x}_k\|^2 + \alpha^2 n L^2 \|\bar{z}_{k-1}\|^2 \right] \\ &\quad + n c_0 \tilde{\rho}_w^3 (1-\beta)^2 C \left\{ 2[f(\bar{x}_k) - f^*] + \frac{L}{n} \|\Pi \mathbf{x}_k\|^2 \right\}. \end{aligned} \quad (67)$$

Taking the full expectation in (66) and (67) and using the similar steps as in the preceding analysis lead to

$$\begin{aligned} \mathbb{E} \left[\left\| \tilde{\Pi} \tilde{\mathbf{y}}_k \right\|^2 \right] &\leq \frac{1}{\tilde{\rho}_w^k} \mathbb{E} \left[\left\| \tilde{\Pi} \tilde{W}^k \tilde{\Pi} \tilde{\mathbf{y}}_0 \right\|^2 \right] + n c_0 \tilde{\rho}_w^2 (1-\beta)^2 (2C\sigma_f^* + \sigma^2) \sum_{t=0}^{k-1} \tilde{\rho}_w^{k-1-t} \\ &\quad + \frac{3(1-\beta)^2 c_0 \tilde{\rho}_w^2}{1-\tilde{\rho}_w} \sum_{t=0}^{k-1} \tilde{\rho}_w^{k-1-t} \left\{ \mathbb{E} \left[\left\| \mathbf{z}_t - \nabla F(\mathbf{1}\bar{x}_t^\top) \right\|^2 \right] + L^2 \mathbb{E} \left[\|\Pi \mathbf{x}_{t+1}\|^2 \right] + \alpha^2 L^2 n \mathbb{E} \left[\|\bar{z}_t\|^2 \right] \right\} \\ &\quad + n c_0 \tilde{\rho}_w^2 (1-\beta)^2 C \sum_{t=1}^k \tilde{\rho}_w^{k-t} \left\{ 2\mathbb{E} [f(\bar{x}_t) - f^*] + \frac{L}{n} \mathbb{E} \left[\|\Pi \mathbf{x}_t\|^2 \right] \right\} \\ &\leq c_0 \tilde{\rho}_w^k \mathbb{E} \left[\|\Pi \mathbf{y}_0\|^2 \right] + \frac{3(1-\beta)^2 c_0 \tilde{\rho}_w^2}{1-\tilde{\rho}_w} \sum_{t=0}^{k-1} \tilde{\rho}_w^{k-1-t} \left\{ \mathbb{E} \left[\left\| \mathbf{z}_t - \nabla F(\mathbf{1}\bar{x}_t^\top) \right\|^2 \right] + L^2 \mathbb{E} \left[\|\Pi \mathbf{x}_{t+1}\|^2 \right] + \alpha^2 L^2 n \mathbb{E} \left[\|\bar{z}_t\|^2 \right] \right\} \\ &\quad + n c_0 \tilde{\rho}_w^2 (1-\beta)^2 (2C\sigma_f^* + \sigma^2) \sum_{t=0}^{k-1} \tilde{\rho}_w^{k-1-t} + n c_0 \tilde{\rho}_w^2 (1-\beta)^2 C \sum_{t=1}^k \tilde{\rho}_w^{k-t} \left\{ 2\mathbb{E} [f(\bar{x}_t) - f^*] + \frac{L}{n} \mathbb{E} \left[\|\Pi \mathbf{x}_t\|^2 \right] \right\} \\ &:= \mathcal{R}_k^y, \quad k \geq 1, \end{aligned} \quad (68)$$

where we invoke Lemma 2.1 for the last inequality. Then we can obtain the recursion for \mathcal{R}_k^y according to (68):

$$\begin{aligned} \mathcal{R}_{k+1}^y &= \tilde{\rho}_w \mathcal{R}_k^y + \frac{3(1-\beta)^2 c_0 \tilde{\rho}_w^2}{1-\tilde{\rho}_w} \left\{ \mathbb{E} \left[\left\| \mathbf{z}_k - \nabla F(\mathbf{1}\bar{x}_k^\top) \right\|^2 \right] + L^2 \|\Pi \mathbf{x}_{k+1}\|^2 + \alpha^2 L^2 n \mathbb{E} \left[\|\bar{z}_k\|^2 \right] \right\} \\ &\quad + n c_0 \tilde{\rho}_w^2 (1-\beta)^2 (2C\sigma_f^* + \sigma^2) + n c_0 \tilde{\rho}_w^2 (1-\beta)^2 C \left\{ 2\mathbb{E} [f(\bar{x}_{k+1}) - f^*] + \frac{L}{n} \mathbb{E} \left[\|\Pi \mathbf{x}_{k+1}\|^2 \right] \right\}, \quad k \geq 1. \end{aligned} \quad (69)$$

Note that (68) defines \mathcal{R}_1^y as follows

$$\begin{aligned} \mathcal{R}_1^y &= c_0 \tilde{\rho}_w \mathbb{E} \left[\|\Pi \mathbf{y}_0\|^2 \right] + \frac{3(1-\beta)^2 c_0 \tilde{\rho}_w^2}{1-\tilde{\rho}_w} \left\{ \mathbb{E} \left[\left\| \mathbf{z}_0 - \nabla F(\mathbf{1}\bar{x}_0^\top) \right\|^2 \right] + L^2 \mathbb{E} \left[\|\Pi \mathbf{x}_1\|^2 \right] + \alpha^2 L^2 n \mathbb{E} \left[\|\bar{z}_0\|^2 \right] \right\} \\ &\quad + n c_0 \tilde{\rho}_w^2 (1-\beta)^2 (2C\sigma_f^* + \sigma^2) + n c_0 \tilde{\rho}_w^2 (1-\beta)^2 C \left\{ 2\mathbb{E} [f(\bar{x}_1) - f^*] + \frac{L}{n} \mathbb{E} \left[\|\Pi \mathbf{x}_1\|^2 \right] \right\}. \end{aligned}$$

Hence, we can define $\mathcal{R}_0^y := c_0 \mathbb{E}[\|\Pi \mathbf{y}_0\|^2] + n c_0 \tilde{\rho}_w (1-\beta)^2 (2C\sigma_f^* + \sigma^2)$ so that the recursion (69) also holds for $k=0$.

Noting that $\mathbb{E}[\|\Pi\mathbf{x}_{k+1}\|^2] \leq \mathbb{E}[\|\tilde{\Pi}\tilde{\mathbf{x}}_{k+1}\|^2] \leq \mathcal{R}_{k+1}^x$ and $\mathbb{E}[\|\tilde{\Pi}\tilde{\mathbf{y}}_k\|^2] \leq \mathcal{R}_k^y$, the desired result follows by invoking (63) and Lemmas 3.3-3.6:

$$\begin{aligned}
\mathcal{R}_{k+1}^y &\leq \tilde{\rho}_w \mathcal{R}_k^y + \frac{3c_0(1-\beta)^2 L(L+C)}{1-\tilde{\rho}_w} \left(\tilde{\rho}_w \mathcal{R}_k^x + \frac{\alpha^2 c_0 \tilde{\rho}_w}{1-\tilde{\rho}_w} \mathcal{R}_k^y \right) + \frac{3c_0(1-\beta)^2}{1-\tilde{\rho}_w} \left\{ \beta \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right] \right. \\
&\quad + 2\alpha n(L+C) \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] + 2(1-\beta)L(L+C) \mathbb{E} \left[\|\Pi\mathbf{x}_k\|^2 \right] + 2n(1-\beta)^2 C \mathbb{E} \left[f(\bar{d}_k) - f^* \right] \\
&\quad + 2\alpha n(1-\beta) C \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] + n(1-\beta)^2 (2C\sigma_f^* + \sigma^2) \left. \right\} + \alpha(1-\beta)nc_0 \left(2C + \frac{3\alpha L^2(1-\beta)}{1-\tilde{\rho}_w} \right) \left\{ \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] \right. \\
&\quad + \frac{2C(1-\beta)^2}{n} \mathbb{E} \left[f(\bar{d}_k) - f^* \right] + 3(1-\beta) \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] + \frac{(1-\beta)^2 (2C\sigma_f^* + \sigma^2)}{n} + \frac{2(1-\beta)L(L+C)}{n} \mathcal{R}_k^x \left. \right\} \\
&\quad + 2nc_0(1-\beta)^2 C \left\{ \left(1 + \frac{2\alpha^2 CL}{n} \right) \mathbb{E} \left[f(\bar{d}_k) - f^* \right] - \frac{\alpha}{2} \left(1 - \frac{4\alpha^2 CL}{n(1-\beta)} \right) \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] \right. \\
&\quad + \frac{\alpha L^2}{n} \mathbb{E} \left[\|\Pi\mathbf{x}_k\|^2 \right] + \frac{\alpha^3 \beta^2 L(L+2C)}{(1-\beta)^2} \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] + \frac{\alpha^2 L(2C\sigma_f^* + \sigma^2)}{n} \left. \right\} + nc_0 \tilde{\rho}_w^2 (1-\beta)^2 (2C\sigma_f^* + \sigma^2) \\
&\quad + \frac{2nc_0(1-\beta)^2 C\alpha}{1-\beta} \left\{ 2\mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] + 2\alpha^2 L^2 \mathbb{E} \left[\|\bar{z}_k\|^2 \right] \right\} \\
&\leq \left[\tilde{\rho}_w + \frac{3\alpha^2 c_0^2 (1-\beta)^2 L(L+C)}{(1-\tilde{\rho}_w)^2} \right] \mathcal{R}_k^y + \frac{3c_0(1-\beta)^2 L}{1-\tilde{\rho}_w} \left[L+C+2(L+C)+2\alpha(L+C)^2+\alpha CL \right] \mathcal{R}_k^x \\
&\quad + \frac{3c_0(1-\beta)^2 \beta}{1-\tilde{\rho}_w} \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right] + 3\alpha n(1-\beta)c_0(L+C) \left[\frac{2(1-\beta)}{1-\tilde{\rho}_w} + 1 + \frac{4\alpha^2 CL}{3(1-\beta)} \right] \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] \\
&\quad + 2nc_0(1-\beta)^2 C \left[\frac{3(1-\beta)^2}{1-\tilde{\rho}_w} + \frac{3\alpha(1-\beta)(C+L)}{n} + 2 \right] \mathbb{E} \left[f(\bar{d}_k) - f^* \right] \\
&\quad + 4\alpha nc_0(1-\beta) \left[C + \frac{6(1-\beta)^2 C}{4(1-\tilde{\rho}_w)} + \frac{9(C+L)(1-\beta)}{4} \right] \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] \\
&\quad + nc_0(1-\beta)^2 \left[1 + \frac{3(1-\beta)^2}{1-\tilde{\rho}_w} + \frac{3\alpha(1-\beta)(L+C)}{n} + \frac{2\alpha^2 CL}{n} \right] (2C\sigma_f^* + \sigma^2) \\
&\leq \frac{1+\tilde{\rho}_w}{2} \mathcal{R}_k^y + \frac{15c_0(1-\beta)^2 L(L+C)}{1-\tilde{\rho}_w} \mathcal{R}_k^x + \frac{3c_0(1-\beta)^2 \beta}{1-\tilde{\rho}_w} \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right] \\
&\quad + 9\alpha nc_0(1-\beta)(C+L) \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] + 12nc_0(1-\beta)^2 C \mathbb{E} \left[f(\bar{d}_k) - f^* \right] \\
&\quad + 20\alpha nc_0(1-\beta)(C+L) \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] + 6nc_0(1-\beta)^2 (2C\sigma_f^* + \sigma^2),
\end{aligned}$$

where we let $1-\beta \leq 1-\tilde{\rho}_w$ and the stepsize satisfy

$$\alpha \leq \min \left\{ \frac{1}{2(L+C)}, \sqrt{\frac{1-\tilde{\rho}_w}{6c_0^2 L(L+C)}}, \sqrt{\frac{1-\beta}{4CL}} \right\}.$$

A.8 Proof of Lemma 3.8

We define the Lyapunov function \mathcal{L}_k as

$$\mathcal{L}_k := \mathbb{E} \left[f(\bar{d}_k) - f^* \right] + \alpha \mathcal{C}_1 \mathcal{R}_k^x + \alpha^3 \mathcal{C}_2 \mathcal{R}_k^y + \alpha^3 \mathcal{C}_3 \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] + \alpha^3 \mathcal{C}_4 \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right], \quad (70)$$

where \mathcal{C}_1 - \mathcal{C}_4 are positive constants to be determined later.

Combing the results of Lemmas 3.4-3.7 and (70) leads to

$$\begin{aligned}
\mathcal{L}_{k+1} \leq & \left[1 + \frac{2\alpha^2 CL}{n} + 12\alpha^3 \mathcal{C}_2 n C (1-\beta)^2 c_0 + \frac{2\alpha^3 C (1-\beta)^2 \mathcal{C}_3}{n} + 2n\alpha^3 \mathcal{C}_4 (1-\beta)^2 C \right] \mathbb{E} [f(\bar{d}_k) - f^*] \\
& + \left[\frac{\alpha L^2}{n} + \alpha \mathcal{C}_1 \tilde{\rho}_w + \frac{15\alpha^3 \mathcal{C}_2 (1-\beta)^2 c_0 L(L+C)}{1-\tilde{\rho}_w} + \frac{2\alpha^3 \mathcal{C}_3 (1-\beta) L(L+C)}{n} + 2\alpha^3 \mathcal{C}_4 (1-\beta)(L+C)L \right] \mathcal{R}_k^x \\
& + \left[\frac{\alpha^3 c_0 \tilde{\rho}_w \mathcal{C}_1}{1-\tilde{\rho}_w} + \frac{(1+\tilde{\rho}_w)\alpha^3 \mathcal{C}_2}{2} \right] \mathcal{R}_k^y \\
& + \left[\frac{\alpha^3 \beta^2 L(L+2C)}{(1-\beta)^2} + 9\alpha^4 \mathcal{C}_2 n(C+L)(1-\beta)c_0 + \frac{(1+\beta)\alpha^3 \mathcal{C}_3}{2} + 2\alpha^4 \mathcal{C}_4 n(C+L) \right] \mathbb{E} [\|\bar{z}_{k-1}\|^2] \\
& + \left[\frac{3(1-\beta)^2 \alpha^3 \mathcal{C}_2 c_0 \beta}{1-\tilde{\rho}_w} + \alpha^3 \mathcal{C}_4 \beta \right] \mathbb{E} [\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2] \\
& - \frac{\alpha}{2} \left[1 - \frac{4\alpha^2 CL}{n(1-\beta)} - 40\alpha^3 n c_0 (1-\beta)(C+L)\mathcal{C}_2 - 6(1-\beta)\alpha^2 \mathcal{C}_3 - 4n(1-\beta)\alpha^3 \mathcal{C}_4 C \right] \mathbb{E} [\|\nabla f(\bar{x}_k)\|^2] \\
& + \left[\frac{L}{n} + 6\alpha \mathcal{C}_2 n c_0 (1-\beta)^2 + \frac{\alpha \mathcal{C}_3 (1-\beta)^2}{n} + n\alpha \mathcal{C}_4 (1-\beta)^2 \right] \alpha^2 (2C\sigma_f^* + \sigma^2).
\end{aligned} \tag{71}$$

Therefore, it suffices to determine \mathcal{C}_1 - \mathcal{C}_4 and the parameters α and β such that

$$\frac{\alpha L^2}{n} + \frac{15\alpha^3 \mathcal{C}_2 (1-\beta)^2 c_0 L(L+C)}{1-\tilde{\rho}_w} + \frac{2\alpha^3 \mathcal{C}_3 (1-\beta) L(L+C)}{n} + 2\alpha^3 \mathcal{C}_4 (1-\beta)(L+C)L \leq (1-\tilde{\rho}_w)\alpha \mathcal{C}_1, \tag{72a}$$

$$\frac{\alpha^3 c_0 \tilde{\rho}_w \mathcal{C}_1}{1-\tilde{\rho}_w} + \frac{(1+\tilde{\rho}_w)\alpha^3 \mathcal{C}_2}{2} \leq \alpha^3 \mathcal{C}_2, \tag{72b}$$

$$\frac{\alpha^3 \beta^2 L(L+2C)}{(1-\beta)^2} + 9\alpha^4 \mathcal{C}_2 n(C+L)(1-\beta)c_0 + \frac{(1+\beta)\alpha^3 \mathcal{C}_3}{2} + 2\alpha^4 \mathcal{C}_4 n(C+L) \leq \alpha^3 \mathcal{C}_3, \tag{72c}$$

$$\frac{3(1-\beta)^2 \alpha^3 \mathcal{C}_2 c_0 \beta}{1-\tilde{\rho}_w} + \alpha^3 \mathcal{C}_4 \beta \leq \alpha^3 \mathcal{C}_4. \tag{72d}$$

From (72d), it is sufficient to choose $\mathcal{C}_4 = 4c_0 \mathcal{C}_2$ given that $\beta \geq \tilde{\rho}_w$. According to (72b), we can choose $\mathcal{C}_2 = 4c_0 \tilde{\rho}_w \mathcal{C}_1 / (1-\tilde{\rho}_w)^2$. Substituting the above \mathcal{C}_2 and \mathcal{C}_4 into (72a) and (72c) leads to

$$\frac{L^2}{n} + \frac{2\alpha^2 \mathcal{C}_3 (1-\beta) L(L+C)}{n} \leq \left(1 - \tilde{\rho}_w - \frac{60\alpha^2 (1-\beta)^2 c_0^2 \tilde{\rho}_w L(L+C)}{(1-\tilde{\rho}_w)^3} - \frac{32\alpha^2 c_0^2 (1-\beta)(L+C)L\tilde{\rho}_w}{(1-\tilde{\rho}_w)^2} \right) \mathcal{C}_1, \tag{73a}$$

$$\frac{\beta^2 L(L+2C)}{(1-\beta)^2} + \frac{36\alpha n(1-\beta)c_0^2 \mathcal{C}_1 \tilde{\rho}_w}{(1-\tilde{\rho}_w)^2} + \frac{8\alpha c_0^2 n(C+L)\mathcal{C}_1 \tilde{\rho}_w}{(1-\tilde{\rho}_w)^2} \leq \frac{(1-\beta)\mathcal{C}_3}{2}. \tag{73b}$$

Note that $(1-\beta) \leq (1-\tilde{\rho}_w)$. Letting the stepsize α satisfy

$$\alpha \leq \min \left\{ \sqrt{\frac{(1-\tilde{\rho}_w)^2}{240c_0^2 L(L+C)}}, \sqrt{\frac{(1-\tilde{\rho}_w)^2}{128c_0^2 L(L+C)}} \right\} \tag{74}$$

leads to

$$\mathcal{C}_1 \geq \frac{2L^2}{n(1-\tilde{\rho}_w)} + \frac{4\alpha^2 \mathcal{C}_3 (1-\beta) L(L+C)}{n(1-\tilde{\rho}_w)}. \tag{75}$$

According to (73b), (74), and (75), we can choose \mathcal{C}_1 and \mathcal{C}_3 as follows:

$$\mathcal{C}_1 = \frac{3L^2}{n(1-\tilde{\rho}_w)}, \quad \mathcal{C}_3 = \frac{4L(L+2C)}{(1-\beta)^3}. \tag{76}$$

To make such a choice feasible, it is sufficient to let the stepsize α and the parameter β satisfy

$$\alpha \leq \min \left\{ \frac{1-\tilde{\rho}_w}{48L}, \frac{1}{216(C+L)} \right\}, \quad \beta \geq \tilde{\rho}_w,$$

so that the following inequality, which is derived from (73b), holds

$$\frac{108\alpha(1-\beta)c_0CL^2}{(1-\tilde{\rho}_w)^3} + \frac{24\alpha c_0(C+L)L^2}{(1-\tilde{\rho}_w)^3} \leq \frac{L(L+2C)}{(1-\beta)^2}.$$

Therefore, we have determined the constants \mathcal{C}_1 - \mathcal{C}_4 :

$$\mathcal{C}_1 = \frac{3L^2}{n(1-\tilde{\rho}_w)}, \mathcal{C}_2 = \frac{12c_0\tilde{\rho}_wL^2}{n(1-\tilde{\rho}_w)^3}, \mathcal{C}_3 = \frac{4L(L+2C)}{(1-\beta)^3}, \mathcal{C}_4 = \frac{48c_0^2\tilde{\rho}_wL^2}{n(1-\tilde{\rho}_w)^3}. \quad (77)$$

Hence, the inequality (71) becomes

$$\begin{aligned} \mathcal{L}_{k+1} &\leq \left[1 + \frac{2\alpha^2CL}{n} + \frac{240\alpha^3c_0^2\tilde{\rho}_wCL^2(1-\beta)^2}{(1-\tilde{\rho}_w)^3} + \frac{8\alpha^3CL(L+2C)}{n(1-\beta)} \right] \mathcal{L}_k \\ &\quad - \frac{\alpha}{2} \left[1 - \frac{4\alpha^2CL}{n(1-\beta)} - \frac{672\alpha^3c_0^2\tilde{\rho}_w(1-\beta)L^2(C+L)}{(1-\tilde{\rho}_w)^3} - \frac{24\alpha^2L(L+2C)}{(1-\beta)^2} \right] \mathbb{E} \left[\|\nabla f(\bar{x}_k)\|^2 \right] \\ &\quad + \left[\frac{L}{n} + \frac{72\alpha c_0^2\tilde{\rho}_wL^2(1-\beta)^2}{(1-\tilde{\rho}_w)^3} + \frac{4\alpha L(L+2C)}{n(1-\beta)} \right] \alpha^2(2C\sigma_f^* + \sigma^2). \end{aligned} \quad (78)$$

Letting the stepsize satisfy

$$\alpha \leq \min \left\{ \frac{1-\beta}{4(L+2C)}, \left(\frac{(1-\tilde{\rho}_w)^3}{4032c_0^2(1-\beta)L^2(C+L)} \right)^{1/3}, \sqrt{\frac{(1-\beta)^2}{144L(L+2C)}}, \sqrt{\frac{1-\beta}{24CL}} \right\}$$

yields the desired result for the recursion of \mathcal{L}_k .

We next consider the term \mathcal{L}_0 in the following. Noting the definition of \mathcal{R}_0^x , \mathcal{R}_0^y in (14) and that

$$\begin{aligned} \mathbb{E} \left[\|\Pi \mathbf{y}_0\|^2 \right] &\leq (1-\beta)^2 \mathbb{E} \left[\|\mathbf{g}_0\|^2 \right] \\ &\leq (1-\beta)^2 \left\{ 2nC[f(\bar{x}_0) - f^*] + 2nC\sigma_f^* + n\sigma^2 + 2L(L+C)\|\Pi \mathbf{x}_0\|^2 + 2 \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2 \right\}, \\ \|\mathbf{z}_{-1} - \nabla F(\mathbf{1}\bar{x}_{-1}^\top)\|^2 &= \|\nabla F(\mathbf{1}\bar{x}_0^\top)\|^2 = \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2, \end{aligned} \quad (79)$$

we have

$$\begin{aligned} \mathcal{L}_0 &= f(\bar{x}_0) - f^* + \frac{3\alpha L^2}{n(1-\tilde{\rho}_w)} \mathcal{R}_0^x + \frac{12\alpha^3c_0\tilde{\rho}_wL^2}{n(1-\tilde{\rho}_w)^3} \mathcal{R}_0^y + \frac{48\alpha^3c_0^2\tilde{\rho}_wL^2}{n(1-\tilde{\rho}_w)^3} \|\mathbf{z}_{-1} - \nabla F(\mathbf{1}\bar{x}_{-1}^\top)\|^2 \\ &\leq \left[1 + \frac{6\alpha^3c_0^2L^2C(1-\beta)^2}{(1-\tilde{\rho}_w)^2} + \frac{24\alpha^3c_0^2L^2C(1-\beta)^2}{(1-\tilde{\rho}_w)^3} \right] \Delta_0 \\ &\quad + \left[c_0 + \frac{2\alpha^2c_0(1-\beta)^2L(L+C)}{1-\tilde{\rho}_w} + \frac{8\alpha^2c_0^2L(L+C)(1-\beta)^2}{(1-\tilde{\rho}_w)^2} \right] \frac{3\alpha L^2 \|\Pi \mathbf{x}_0\|^2}{n(1-\tilde{\rho}_w)} \\ &\quad + \frac{27\alpha^3c_0^2L^2(1-\beta)^2(2C\sigma_f^* + \sigma^2)}{(1-\tilde{\rho}_w)^3} + \frac{72\alpha^3c_0^2L^2 \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{n(1-\tilde{\rho}_w)^3} \\ &\leq 2\Delta_0 + \frac{9\alpha L^2 \|\Pi \mathbf{x}_0\|^2}{n(1-\tilde{\rho}_w)} + \frac{27\alpha^3c_0^2L^2(1-\beta)^2(2C\sigma_f^* + \sigma^2)}{(1-\tilde{\rho}_w)^2} + \frac{72\alpha^3c_0^2L^2 \sum_{i=1}^n \|\nabla f_i(\bar{x}_0)\|^2}{n(1-\tilde{\rho}_w)^3}. \end{aligned} \quad (80)$$

B Proofs for the PL Condition Case

B.1 Proof of Lemma 4.4

Similar to the derivation of (49), we have

$$\begin{aligned}
\mathbb{E} [f(\bar{d}_{k+1}) - f^* | \mathcal{F}_k] &\leq f(\bar{d}_k) - f^* - \alpha \left\langle \nabla f(\bar{d}_k), \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\rangle + \frac{L}{2} \mathbb{E} \left[\|\alpha \bar{g}_k\|^2 | \mathcal{F}_k \right] \\
&\leq f(\bar{d}_k) - f^* - \frac{\alpha}{2} \|\nabla f(\bar{d}_k)\|^2 - \frac{\alpha(1-\alpha L)}{2} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\|^2 \\
&\quad + \frac{\alpha}{2} \left\| \nabla f(\bar{d}_k) - \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\|^2 + \frac{\alpha^2 L}{2} \mathbb{E} \left[\left\| \bar{g}_k - \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_{i,k}) \right\|^2 | \mathcal{F}_k \right] \\
&\leq (1-\alpha\mu) [f(\bar{d}_k) - f^*] + \frac{\alpha L^2}{2n} \sum_{i=1}^n \|\bar{d}_k - x_{i,k}\|^2 + \frac{\alpha^2 L(2C\sigma_f^* + \sigma^2)}{n} + \frac{\alpha^2 L^2 C}{2n^2} \|\Pi \mathbf{x}_k\|^2 \\
&\quad + \frac{\alpha^2 LC}{n} \left\{ \left(1 + \frac{\alpha\beta L}{(1-\beta)} \right) [f(\bar{d}_k) - f^*] + \frac{\alpha\beta(1 + \alpha\beta L/(1-\beta))}{2(1-\beta)} \|\bar{z}_{k-1}\|^2 \right\},
\end{aligned} \tag{81}$$

where we invoke L -smoothness of f_i , Assumptions 1.2 and 4.3, and (47) for the last inequality.

In light of the relation $\bar{d}_k - \bar{x}_k = -\alpha\beta\bar{z}_{k-1}/(1-\beta)$, we have

$$\frac{1}{n} \sum_{i=1}^n \|\bar{d}_k - x_{i,k}\|^2 = \frac{\alpha^2\beta^2}{(1-\beta)^2} \|\bar{z}_{k-1}\|^2 + \frac{1}{n} \|\Pi \mathbf{x}_k\|^2. \tag{82}$$

Combing (81)-(82), letting $\alpha \leq (1-\beta)/L$, and taking the full expectation lead to

$$\begin{aligned}
\mathbb{E} [f(\bar{d}_{k+1}) - f^*] &\leq \left(1 - \alpha\mu + \frac{2\alpha^2 LC}{n} \right) \mathbb{E} [f(\bar{d}_k) - f^*] + \frac{\alpha L^2}{n} \mathbb{E} \left[\|\Pi \mathbf{x}_k\|^2 \right] + \frac{\alpha^2 L(2C\sigma_f^* + \sigma^2)}{n} \\
&\quad + \frac{\alpha^3 \beta L [C(1-\beta) + L\beta]}{(1-\beta)^2} \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right].
\end{aligned}$$

Letting $\alpha \leq \mu/(4LC)$ yields the desired result.

B.2 Proof of Lemma 4.5

Similar to the derivation of (71), we determine the new recursion of \mathcal{L}_k in light of Assumption 4.3 as follows:

$$\begin{aligned}
\mathcal{L}_{k+1} &\leq \left[1 - \frac{\alpha\mu}{2} + \frac{240\alpha^3 c_0^2 \tilde{\rho}_w C L^2}{1 - \tilde{\rho}_w} + \frac{8\alpha^3 C L(L+2C)}{n(1-\beta)} \right] \mathbb{E} [f(\bar{d}_k) - f^*] \\
&\quad + \left[\frac{\alpha L^2}{n} + \frac{3\alpha L^2 \tilde{\rho}_w}{n(1-\tilde{\rho}_w)} + \frac{180\alpha^3 c_0^2 \tilde{\rho}_w L^3 (L+C)}{n(1-\tilde{\rho}_w)} + \frac{8\alpha^3 L^2 (L+2C)(L+C)}{n(1-\beta)^2} + \frac{96\alpha^3 c_0^2 \tilde{\rho}_w (L+C)L^3}{n(1-\tilde{\rho}_w)^2} \right] \mathcal{R}_k^x \\
&\quad + \left[\frac{3\alpha^3 c_0 L^2 \tilde{\rho}_w}{n(1-\tilde{\rho}_w)^2} + \frac{12(1+\tilde{\rho}_w)\alpha^3 c_0 \tilde{\rho}_w L^2}{2n(1-\tilde{\rho}_w)^3} \right] \mathcal{R}_k^y \\
&\quad + \left[\frac{\alpha^3 \beta L(L+2C)}{(1-\beta)^2} + \frac{108\alpha^4 L^2 (C+L)c_0^2 \tilde{\rho}_w}{(1-\tilde{\rho}_w)^2} + \frac{4(1+\beta)\alpha^3 L(L+2C)}{2(1-\beta)^3} + \frac{96\alpha^4 c_0^2 \tilde{\rho}_w L^2 (C+L)}{(1-\tilde{\rho}_w)^3} \right] \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] \\
&\quad + \left[\frac{36\alpha^3 c_0^2 \tilde{\rho}_w \beta L^2 (1-\beta)^2}{n(1-\tilde{\rho}_w)^4} + \frac{48\alpha^3 c_0^2 \tilde{\rho}_w L^2 \beta}{n(1-\tilde{\rho}_w)^3} \right] \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right] \\
&\quad + \frac{48\alpha^3 L(L+C)}{(1-\beta)^2} \left\{ 4L\mathbb{E} [f(\bar{d}_k) - f^*] + \frac{2\alpha^2 \beta^2 L^2}{(1-\beta)^2} \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] \right\} \\
&\quad + \left[\frac{L}{n} + \frac{72\alpha c_0^2 \tilde{\rho}_w L^2 (1-\beta)^2}{(1-\tilde{\rho}_w)^3} + \frac{4\alpha L(L+2C)}{n(1-\beta)} \right] \alpha^2 (2C\sigma_f^* + \sigma^2),
\end{aligned} \tag{83}$$

where we consider $\alpha \leq 1/(28c_0^2L)$, $\beta \geq \tilde{\rho}_w$, and the estimate in (84). The estimate (84) holds due to (52)

$$\|\nabla f(\bar{x}_k)\|^2 \leq 2\|\nabla f(\bar{d}_k)\|^2 + \frac{2\alpha^2\beta^2L^2}{(1-\beta)^2}\|\bar{z}_{k-1}\|^2 \leq 4L[f(\bar{d}_k) - f^*] + \frac{2\alpha^2\beta^2L^2}{(1-\beta)^2}\|\bar{z}_{k-1}\|^2. \quad (84)$$

To derive the desired result in (28), it is then sufficient to determine α and β such that

$$\frac{240\alpha^3c_0^2\tilde{\rho}_wCL^2}{1-\tilde{\rho}_w} + \frac{8\alpha^3CL(L+2C)}{n(1-\beta)} + \frac{192\alpha^3L^2(L+C)}{(1-\beta)^2} \leq \frac{\alpha\mu}{4}, \quad (85a)$$

$$12\alpha^2c_0^2\tilde{\rho}_w^3LC + \frac{8\alpha^2(L+2C)(L+C)(1-\tilde{\rho}_w)}{3(1-\beta)^2} + \frac{32\alpha^2c_0^2\tilde{\rho}_w(L+C)L}{1-\tilde{\rho}_w} \leq \frac{2(1-\tilde{\rho}_w)}{3} - \frac{\alpha\mu}{4}, \quad (85b)$$

$$0 \leq \frac{1-\tilde{\rho}_w}{4} - \frac{\alpha\mu}{4}, \quad (85c)$$

$$\frac{27\alpha L(C+L)c_0^2\tilde{\rho}_w(1-\beta)}{L+2C} + \frac{24\alpha c_0^2\tilde{\rho}_wL(C+L)}{L+2C} + \frac{24\alpha^2(L+C)L^2\beta^2}{(L+2C)(1-\beta)} \leq \frac{1-\beta}{4} - \frac{\alpha\mu}{4}, \quad (85d)$$

$$\frac{3(1-\beta)^2}{4(1-\tilde{\rho}_w)} \leq 1-\beta - \frac{\alpha\mu}{4}. \quad (85e)$$

To ensure (85a) holds, it suffices to have

$$\frac{240\alpha^3c_0^2L(L+C)^2}{(1-\beta)^2} \leq \frac{\alpha\mu}{4}, \quad \beta \geq 1 - \frac{11}{12}(1-\tilde{\rho}_w).$$

Then, we let $\alpha \leq \sqrt{\mu(1-\beta)^2/[960c_0^2L(L+C)^2]}$.

For (85b) to hold, it suffices to let the stepsize α satisfy

$$\alpha \leq \min \left\{ \frac{4(1-\tilde{\rho}_w)}{3\mu}, \sqrt{\frac{1-\tilde{\rho}_w}{108c_0^2LC}}, \sqrt{\frac{(1-\tilde{\rho}_w)^2}{288c_0L(L+C)}}, \sqrt{\frac{(1-\beta)^2}{48(L+C)^2}} \right\}.$$

For (85c), it is sufficient to let $\alpha \leq (1-\tilde{\rho}_w)/\mu$.

For (85d), it suffices to have

$$\alpha \leq \min \left\{ \frac{1-\beta}{3\mu}, \frac{1}{486c_0^2L}, \frac{1-\beta}{432c_0L}, \sqrt{\frac{(1-\beta)^2}{432L^2}} \right\}.$$

Finally, we choose $\beta \geq (1+11\tilde{\rho}_w)/12$ such that (85e) holds and obtain the desired result.

B.3 Proof of Theorem 4.2

We first derive an upper bound for $\mathbb{E}[f(\bar{d}_k) - f^*]$. Unrolling the recursion (28) in Lemma 4.5 and rearranging the terms yield

$$\begin{aligned} \mathbb{E}[f(\bar{d}_k) - f^*] &\leq \mathcal{L}_k \leq \left(1 - \frac{\alpha\mu}{4}\right)^k \mathcal{L}_0 + \left[\frac{L}{n} + \frac{72\alpha c_0^2\tilde{\rho}_wL^2(1-\beta)^2}{(1-\tilde{\rho}_w)^3} + \frac{4\alpha L(L+2C)}{n(1-\beta)} \right] \frac{4\alpha(2C\sigma_f^* + \sigma^2)}{\mu} \\ &\leq \left(1 - \frac{\alpha\mu}{4}\right)^k \mathcal{L}_0 + \frac{4\alpha L(2C\sigma_f^* + \sigma^2)}{n\mu} + \frac{16\alpha^2L^2\mathcal{S}_1(2C\sigma_f^* + \sigma^2)}{(1-\tilde{\rho}_w)\mu}, \end{aligned} \quad (86)$$

where

$$\mathcal{S}_1 := \frac{18c_0^2(1-\beta)^2}{(1-\tilde{\rho}_w)^2} + \frac{(L+2C)(1-\tilde{\rho}_w)}{nL(1-\beta)}.$$

We next construct a new Lyapunov function \mathcal{H}_k in (87) such that the consensus error $\mathbb{E}[\|\Pi\mathbf{x}_k\|^2]$ can be bounded.

$$\mathcal{H}_k := \mathcal{R}_k^x + \alpha^2\mathcal{C}_5\mathcal{R}_k^y + \alpha^3\mathcal{C}_6\mathbb{E}\left[\|\bar{z}_{k-1}\|^2\right] + \alpha^2\mathcal{C}_7\mathbb{E}\left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2\right]. \quad (87)$$

Combing the results of Lemmas 3.5-3.7, invoking (84), and rearranging the terms yield

$$\begin{aligned}
\mathcal{H}_{k+1} \leq & \left[\tilde{\rho}_w + \frac{15\alpha^2\mathcal{C}_5(1-\beta)^2c_0L(C+L)}{1-\tilde{\rho}_w} + \frac{2\alpha^3(1-\beta)L(L+C)\mathcal{C}_6}{n} + 2\alpha^2\mathcal{C}_7(1-\beta)L(L+C) \right] \mathcal{R}_k^x \\
& + \left[\frac{\alpha^2c_0\tilde{\rho}_w}{1-\tilde{\rho}_w} + \frac{\alpha^2\mathcal{C}_5(1+\tilde{\rho}_w)}{2} \right] \mathcal{R}_k^y \\
& + \left[9\alpha^3\mathcal{C}_5nc_0(1-\beta)(C+L) + \frac{(1+\beta)\alpha^3\mathcal{C}_6}{2} + 2\alpha^3\mathcal{C}_7n(C+L) \right] \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] \\
& + \left[\frac{3\alpha^2\mathcal{C}_5(1-\beta)^2c_0\beta}{1-\tilde{\rho}_w} + \beta\alpha^2\mathcal{C}_7 \right] \mathbb{E} \left[\|\mathbf{z}_{k-1} - \nabla F(\mathbf{1}\bar{x}_{k-1}^\top)\|^2 \right] \\
& + \left[12\alpha^2\mathcal{C}_5nc_0(1-\beta)^2C + \frac{2\alpha^3\mathcal{C}_6C(1-\beta)^2}{n} + 2\alpha^2\mathcal{C}_7n(1-\beta)^2C \right] \mathcal{L}_k \\
& + \left[20\alpha^3\mathcal{C}_5nc_0(1-\beta)(C+L) + 3\alpha^3\mathcal{C}_6(1-\beta) + 2\alpha^3\mathcal{C}_7n(1-\beta)C \right] \left\{ 4L\mathcal{L}_k + \frac{2\alpha^2\beta^2L^2}{(1-\beta)^2} \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] \right\} \\
& + \left[6\mathcal{C}_5nc_0(1-\beta)^2 + \frac{\alpha\mathcal{C}_6(1-\beta)^2}{n} + \mathcal{C}_7n(1-\beta)^2 \right] \alpha^2(2C\sigma_f^* + \sigma^2).
\end{aligned} \tag{88}$$

Letting $\alpha \leq (1-\beta)/(5L)$, it is sufficient to determine \mathcal{C}_5 - \mathcal{C}_7 , and β such that

$$\tilde{\rho}_w + \frac{15\alpha^2\mathcal{C}_5(1-\beta)^2c_0L(C+L)}{1-\tilde{\rho}_w} + \frac{2\alpha^3(1-\beta)L(L+C)\mathcal{C}_6}{n} + 2\alpha^2\mathcal{C}_7(1-\beta)L(L+C) \leq \frac{5+\beta}{6}, \tag{89a}$$

$$\frac{\alpha^2c_0\tilde{\rho}_w}{1-\tilde{\rho}_w} + \frac{\alpha^2\mathcal{C}_5(1+\tilde{\rho}_w)}{2} \leq \frac{5+\beta}{6}\alpha^2\mathcal{C}_5, \tag{89b}$$

$$9\alpha^3\mathcal{C}_5nc_0(1-\beta)(C+L) + \frac{(1+\beta)\alpha^3\mathcal{C}_6}{2} + 2\alpha^3\mathcal{C}_7n(C+L) \leq \frac{5+\beta}{6}\alpha^3\mathcal{C}_6, \tag{89c}$$

$$\frac{3\alpha^2\mathcal{C}_5(1-\beta)^2c_0\beta}{1-\tilde{\rho}_w} + \alpha^2\mathcal{C}_7\beta \leq \frac{5+\beta}{6}\alpha^2\mathcal{C}_7. \tag{89d}$$

For (89d), we can choose $\mathcal{C}_7 = 18c_0\mathcal{C}_5$. For (89b) it suffices that

$$\frac{c_0\tilde{\rho}_w}{1-\tilde{\rho}_w} \leq \left(\frac{5+\beta}{6} - \frac{1+\tilde{\rho}_w}{2} \right) \mathcal{C}_5.$$

We can then choose

$$\mathcal{C}_5 := \frac{3c_0\tilde{\rho}_w}{(1-\tilde{\rho}_w)^2}, \quad \mathcal{C}_7 := \frac{54c_0^2\tilde{\rho}_w}{(1-\tilde{\rho}_w)^2}, \quad \beta \geq \tilde{\rho}_w.$$

Substituting \mathcal{C}_5 and \mathcal{C}_7 into (89c) yields

$$\frac{27c_0^2\tilde{\rho}_wnC(1-\beta)}{(1-\tilde{\rho}_w)^2} + \frac{108c_0^2\tilde{\rho}_wn(C+L)}{(1-\tilde{\rho}_w)^2} \leq \frac{(1-\beta)\mathcal{C}_6}{3}. \tag{90}$$

Therefore, we can choose

$$\mathcal{C}_6 := \frac{405nc_0^2(C+L)}{(1-\tilde{\rho}_w)^3}, \quad \beta \geq \tilde{\rho}_w. \tag{91}$$

Substituting $\mathcal{C}_5 - \mathcal{C}_7$ into (89a) and noting $\beta \geq \tilde{\rho}_w$ lead to

$$\frac{45\alpha^2(1-\beta)^2c_0^2L(C+L)\tilde{\rho}_w}{(1-\tilde{\rho}_w)^3} + \frac{910c_0^2\alpha^3(1-\beta)L(L+C)^2}{(1-\tilde{\rho}_w)^3} + \frac{108c_0^2\tilde{\rho}_w\alpha^2(1-\beta)L(L+C)}{(1-\tilde{\rho}_w)^2} \leq \frac{5(1-\tilde{\rho}_w)}{6}. \tag{92}$$

It suffices to let $\alpha \leq (1 - \tilde{\rho}_w)/[160c_0(L + C)]$ for (92) to hold. Then, the inequality (88) becomes

$$\begin{aligned} \mathcal{H}_{k+1} &\leq \frac{5 + \beta}{6} \mathcal{H}_k + \frac{72nc_0^2\alpha^2(1 - \beta)^2(2C\sigma_f^* + \sigma^2)}{(1 - \tilde{\rho}_w)^2} \left[1 + \frac{45\alpha(C + L)}{8(1 - \tilde{\rho}_w)} \right] \\ &\quad + \left[\frac{144\alpha^2nc_0^2(1 - \beta)^2(C + L)(1 + 5\alpha L/(1 - \beta))}{(1 - \tilde{\rho}_w)^2} + \frac{6370\alpha^3nc_0^2(1 - \beta)C(C + L)}{(1 - \tilde{\rho}_w)^3} \right] \mathcal{L}_k \\ &\leq \frac{5 + \beta}{6} \mathcal{H}_k + \frac{168\alpha^2(1 - \beta)^2nc_0^2(2C\sigma_f^* + \sigma^2)}{(1 - \tilde{\rho}_w)^2} + \frac{288\alpha^2c_0^2(1 - \beta)^2n(C + L)\mathcal{L}_k}{(1 - \tilde{\rho}_w)^2}, \end{aligned} \quad (93)$$

where the last inequality follows from $\alpha \leq (1 - \beta)/[234(C + L)]$.

Unrolling (93) and substituting (86) yield

$$\begin{aligned} \mathcal{H}_k &\leq \left(\frac{5 + \beta}{6} \right)^k \mathcal{H}_0 + \frac{1008\alpha^2(1 - \beta)nc_0^2(2C\sigma_f^* + \sigma^2)}{(1 - \tilde{\rho}_w)^2} \\ &\quad + \frac{288\alpha^2c_0^2(1 - \beta)^2n(C + L)\mathcal{L}_0}{(1 - \tilde{\rho}_w)^2} \left(1 - \frac{\alpha\mu}{4} \right)^k \sum_{j=0}^k \left(\frac{1 - \frac{1-\beta}{6}}{1 - \frac{\alpha\mu}{4}} \right)^j \\ &\quad + \frac{1728\alpha^2(1 - \beta)c_0^2n(C + L)}{(1 - \tilde{\rho}_w)^2} \left\{ \frac{4\alpha L(2C\sigma_f^* + \sigma^2)}{n\mu} + \frac{16\alpha^2L^2\mathcal{S}_1(2C\sigma_f^* + \sigma^2)}{(1 - \tilde{\rho}_w)\mu} \right\} \\ &\leq \left(\frac{5 + \beta}{6} \right)^k \mathcal{H}_0 + \frac{25n\mathcal{L}_0}{L + C} \left(1 - \frac{\alpha\mu}{4} \right)^k + \frac{1008\alpha^2(1 + \mathcal{S}_2)(1 - \beta)nc_0^2(2C\sigma_f^* + \sigma^2)}{(1 - \tilde{\rho}_w)^2}, \end{aligned} \quad (94)$$

where the following relations are noted:

$$\begin{aligned} \sum_{j=0}^k \left(\frac{1 - \frac{1-\beta}{6}}{1 - \frac{\alpha\mu}{4}} \right)^j &\leq \sum_{j=0}^k \left(\frac{10 + 2\beta}{11 + \beta} \right)^j \leq \frac{12}{1 - \beta}, \quad \mathcal{S}_2 := \frac{L}{n\mu} + \frac{\mathcal{S}_1L}{\mu}, \\ \alpha &\leq \min \left\{ \frac{1 - \beta}{4L}, \frac{1}{12(C + L)} \right\}. \end{aligned} \quad (95)$$

Substituting (86) and (94) into the result of Lemma 3.5 and invoking (84) yield that, for $\alpha \leq (1 - \beta)/(2L\sqrt{3})$, there holds

$$\begin{aligned} \mathbb{E} \left[\|\bar{z}_k\|^2 \right] &\leq \frac{3 + \beta}{4} \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right] + \frac{(1 - \beta)^2(2C\sigma_f^* + \sigma^2)}{n} \\ &\quad + 2(1 - \beta)(C + 6L) \left\{ \left(1 - \frac{\alpha\mu}{4} \right)^k \mathcal{L}_0 + \frac{4\alpha L(2C\sigma_f^* + \sigma^2)}{n\mu} + \frac{16\alpha^2L^2\mathcal{S}_1(2C\sigma_f^* + \sigma^2)}{(1 - \tilde{\rho}_w)\mu} \right\} \\ &\quad + \frac{2(1 - \beta)L(L + C)}{n} \left\{ \left(\frac{5 + \beta}{6} \right)^k \mathcal{H}_0 + \frac{25n\mathcal{L}_0}{L} \left(1 - \frac{\alpha\mu}{4} \right)^k + \frac{1008\alpha^2(1 + \mathcal{S}_2)(1 - \beta)nc_0^2(2C\sigma_f^* + \sigma^2)}{(1 - \tilde{\rho}_w)^2} \right\} \\ &\leq \frac{24L(L + C)\mathcal{H}_0}{n} \left(\frac{5 + \beta}{6} \right)^k + 24\mathcal{L}_0(C + L) \left(1 - \frac{\alpha\mu}{4} \right)^k + \frac{4(1 - \beta)(2C\sigma_f^* + \sigma^2)}{n} \\ &\quad + \left[\frac{6L + C}{n\mu} + \frac{4\alpha L\mathcal{S}_1(C + 6L)}{(1 - \tilde{\rho}_w)\mu} + \frac{252\alpha(1 + \mathcal{S}_2)(1 - \beta)(L + C)}{(1 - \tilde{\rho}_w)^2} \right] 32\alpha L(2C\sigma_f^* + \sigma^2). \end{aligned} \quad (96)$$

From Lemma 3.3, we have

$$\left(1 - \frac{2\alpha\beta L}{1 - \beta} \right) \mathbb{E} [f(\bar{x}_k) - f^*] \leq \mathbb{E} [f(\bar{d}_k) - f^*] + \frac{2\alpha\beta}{1 - \beta} \mathbb{E} \left[\|\bar{z}_{k-1}\|^2 \right].$$

Therefore, we have for $\alpha \leq (1 - \beta)/(4\beta L)$ that

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E} [f(x_{i,k}) - f^*] &\leq 2\mathbb{E} [f(\bar{x}_k) - f^*] + \frac{L}{n} \mathbb{E} [\|\Pi \mathbf{x}_k\|^2] \\ &\leq 4\mathcal{L}_k + \frac{4\alpha\beta}{1 - \beta} \mathbb{E} [\|\bar{z}_{k-1}\|^2] + \frac{L}{n} \mathcal{H}_k. \end{aligned} \quad (97)$$

Substituting (86), (94), and (96) into (97) and setting

$$\alpha \leq \min \left\{ \frac{1 - \beta}{32(C + L)}, \frac{1 - \beta}{4(C + 6L)} \right\}$$

yields the desired result.

References

- [1] S. A. ALGHUNAIM AND K. YUAN, *A unified and refined convergence analysis for non-convex decentralized learning*, IEEE Transactions on Signal Processing, 70 (2022), pp. 3264–3279.
- [2] L. BOTTOU, F. E. CURTIS, AND J. NOCEDAL, *Optimization methods for large-scale machine learning*, Siam Review, 60 (2018), pp. 223–311.
- [3] T. BROWN, B. MANN, N. RYDER, M. SUBBIAH, J. D. KAPLAN, P. DHARIWAL, A. NEELAKANTAN, P. SHYAM, G. SASTRY, A. ASKELL, ET AL., *Language models are few-shot learners*, Advances in neural information processing systems, 33 (2020), pp. 1877–1901.
- [4] J. CHEN AND A. H. SAYED, *On the limiting behavior of distributed optimization strategies*, in 2012 50th Annual Allerton Conference on Communication, Control, and Computing (Allerton), IEEE, 2012, pp. 1535–1542.
- [5] J. CHEN AND A. H. SAYED, *On the learning behavior of adaptive networks, Part i: Transient analysis*, IEEE Transactions on Information Theory, 61 (2015), pp. 3487–3517.
- [6] A. CHOWDHERY, S. NARANG, J. DEVLIN, M. BOSMA, G. MISHRA, A. ROBERTS, P. BARHAM, H. W. CHUNG, C. SUTTON, S. GEHRMANN, ET AL., *Palm: Scaling language modeling with pathways*, arXiv preprint arXiv:2204.02311, (2022).
- [7] J. GAO, X.-W. LIU, Y.-H. DAI, Y. HUANG, AND J. GU, *Distributed stochastic gradient tracking methods with momentum acceleration for non-convex optimization*, Computational Optimization and Applications, 84 (2023), pp. 531–572.
- [8] E. GHADIMI, H. R. FEYZMAHDAVIAN, AND M. JOHANSSON, *Global convergence of the heavy-ball method for convex optimization*, in 2015 European control conference (ECC), IEEE, 2015, pp. 310–315.
- [9] K. HUANG, X. LI, A. MILZAREK, S. PU, AND J. QIU, *Distributed random reshuffling over networks*, IEEE Transactions on Signal Processing, 71 (2023), pp. 1143–1158.
- [10] K. HUANG, X. LI, AND S. PU, *Distributed stochastic optimization under a general variance condition*, arXiv preprint arXiv:2301.12677, (2023).
- [11] K. HUANG AND S. PU, *Improving the transient times for distributed stochastic gradient methods*, IEEE Transactions on Automatic Control, (2022).
- [12] K. HUANG AND S. PU, *Cedas: A compressed decentralized stochastic gradient method with improved convergence*, 2023, <https://arxiv.org/abs/2301.05872>.
- [13] K. HUANG, L. ZHOU, AND S. PU, *Distributed random reshuffling methods with improved convergence*, 2023, <https://arxiv.org/abs/2306.12037>.
- [14] H. KARIMI, J. NUTINI, AND M. SCHMIDT, *Linear convergence of gradient and proximal-gradient methods under the polyak-łojasiewicz condition*, in Joint European conference on machine learning and knowledge discovery in databases, Springer, 2016, pp. 795–811.
- [15] A. KHALED AND P. RICHTÁRIK, *Better theory for sgd in the nonconvex world*, arXiv preprint arXiv:2002.03329, (2020).
- [16] A. KOLOSKOVA, T. LIN, AND S. U. STICH, *An improved analysis of gradient tracking for decentralized machine learning*, Advances in Neural Information Processing Systems, 34 (2021), pp. 11422–11435.
- [17] A. KOLOSKOVA, S. U. STICH, AND M. JAGGI, *Decentralized stochastic optimization and gossip algorithms with compressed communication*, arXiv preprint arXiv:1902.00340, (2019).

- [18] A. KRIZHEVSKY, G. HINTON, ET AL., *Learning multiple layers of features from tiny images*, (2009).
- [19] Y. LEI, T. HU, G. LI, AND K. TANG, *Stochastic gradient descent for nonconvex learning without bounded gradient assumptions*, IEEE transactions on neural networks and learning systems, 31 (2019), pp. 4394–4400.
- [20] X. LI AND A. MILZAREK, *A unified convergence theorem for stochastic optimization methods*, arXiv preprint arXiv:2206.03907, (2022).
- [21] X. LIAN, C. ZHANG, H. ZHANG, C.-J. HSIEH, W. ZHANG, AND J. LIU, *Can decentralized algorithms outperform centralized algorithms? a case study for decentralized parallel stochastic gradient descent*, in NIPS, 2017, pp. 5336–5346.
- [22] T. LIN, S. P. KARIMIREDDY, S. U. STICH, AND M. JAGGI, *Quasi-global momentum: Accelerating decentralized deep learning on heterogeneous data*, in International Conference On Machine Learning, Vol 139, vol. 139, JMLR-JOURNAL MACHINE LEARNING RESEARCH, 2021.
- [23] Y. LIU, Y. GAO, AND W. YIN, *An improved analysis of stochastic gradient descent with momentum*, Advances in Neural Information Processing Systems, 33 (2020), pp. 18261–18271.
- [24] Y. LU AND C. DE SA, *Optimal complexity in decentralized training*, in International Conference on Machine Learning, PMLR, 2021, pp. 7111–7123.
- [25] L. LUO AND H. YE, *An optimal stochastic algorithm for decentralized nonconvex finite-sum optimization*, arXiv preprint arXiv:2210.13931, (2022).
- [26] K. MISHCHENKO, A. KHALED RAGAB BAYOUMI, AND P. RICHTÁRIK, *Random reshuffling: Simple analysis with vast improvements*, Advances in Neural Information Processing Systems, 33 (2020).
- [27] G. MORRAL, P. BIANCHI, AND G. FORT, *Success and failure of adaptation-diffusion algorithms with decaying step size in multiagent networks*, IEEE Transactions on Signal Processing, 65 (2017), pp. 2798–2813.
- [28] A. NEDIĆ, A. OLSHEVSKY, AND M. G. RABBAT, *Network topology and communication-computation tradeoffs in decentralized optimization*, Proceedings of the IEEE, 106 (2018), pp. 953–976.
- [29] A. NEDIC AND A. OZDAGLAR, *Distributed subgradient methods for multi-agent optimization*, IEEE Transactions on Automatic Control, 54 (2009), pp. 48–61.
- [30] OPENAI, *Gpt-4 technical report*, 2023, <https://arxiv.org/abs/2303.08774>.
- [31] B. T. POLYAK, *Some methods of speeding up the convergence of iteration methods*, USSR computational mathematics and mathematical physics, 4 (1964), pp. 1–17.
- [32] S. PU AND A. GARCIA, *A flocking-based approach for distributed stochastic optimization*, Operations Research, 1 (2018), pp. 267–281.
- [33] S. PU AND A. GARCIA, *Swarming for faster convergence in stochastic optimization*, SIAM Journal on Control and Optimization, 56 (2018), pp. 2997–3020.
- [34] S. PU AND A. NEDIĆ, *Distributed stochastic gradient tracking methods*, Mathematical Programming, 187 (2021), pp. 409–457.
- [35] S. PU, A. OLSHEVSKY, AND I. C. PASCHALIDIS, *A sharp estimate on the transient time of distributed stochastic gradient descent*, 2019, <https://arxiv.org/abs/1906.02702>.
- [36] S. PU, A. OLSHEVSKY, AND I. C. PASCHALIDIS, *Asymptotic network independence in distributed stochastic optimization for machine learning: Examining distributed and centralized stochastic gradient descent*, IEEE signal processing magazine, 37 (2020), pp. 114–122.
- [37] S. PU, A. OLSHEVSKY, AND I. C. PASCHALIDIS, *A sharp estimate on the transient time of distributed stochastic gradient descent*, IEEE Transactions on Automatic Control, (2021).
- [38] K. SCAMAN, F. BACH, S. BUBECK, Y. T. LEE, AND L. MASSOULIÉ, *Optimal algorithms for smooth and strongly convex distributed optimization in networks*, in international conference on machine learning, PMLR, 2017, pp. 3027–3036.
- [39] Z. SONG, L. SHI, S. PU, AND M. YAN, *Optimal gradient tracking for decentralized optimization*, Mathematical Programming, (2023), pp. 1–53.
- [40] A. SPIRIDONOFF, A. OLSHEVSKY, AND I. C. PASCHALIDIS, *Robust asynchronous stochastic gradient-push: Asymptotically optimal and network-independent performance for strongly convex functions.*, Journal of Machine Learning Research, 21 (2020), pp. 1–47.
- [41] Y. TAKEZAWA, H. BAO, K. NIWA, R. SATO, AND M. YAMADA, *Momentum tracking: Momentum acceleration for decentralized deep learning on heterogeneous data*, arXiv preprint arXiv:2209.15505, (2022).

- [42] H. TANG, X. LIAN, M. YAN, C. ZHANG, AND J. LIU, d^2 : *Decentralized training over decentralized data*, in Proceedings of the 35th International Conference on Machine Learning, J. Dy and A. Krause, eds., vol. 80 of Proceedings of Machine Learning Research, PMLR, 10–15 Jul 2018, pp. 4848–4856, <http://proceedings.mlr.press/v80/tang18a.html>.
- [43] H. TANG, X. LIAN, M. YAN, C. ZHANG, AND J. LIU, $D2$: *Decentralized training over decentralized data*, in International Conference on Machine Learning, 2018, pp. 4848–4856.
- [44] J. WANG, Y. LU, B. YUAN, B. CHEN, P. LIANG, C. DE SA, C. RE, AND C. ZHANG, *Cocktailsgd: Fine-tuning foundation models over 500mbps networks*, in International Conference on Machine Learning, PMLR, 2023, pp. 36058–36076.
- [45] Z. WANG, J. ZHANG, T.-H. CHANG, J. LI, AND Z.-Q. LUO, *Distributed stochastic consensus optimization with momentum for nonconvex nonsmooth problems*, IEEE Transactions on Signal Processing, 69 (2021), pp. 4486–4501.
- [46] T. XIAO, X. CHEN, K. BALASUBRAMANIAN, AND S. GHADIMI, *A one-sample decentralized proximal algorithm for non-convex stochastic composite optimization*, arXiv preprint arXiv:2302.09766, (2023).
- [47] R. XIN, U. A. KHAN, AND S. KAR, *An improved convergence analysis for decentralized online stochastic non-convex optimization*, IEEE Transactions on Signal Processing, 69 (2021), pp. 1842–1858, <https://doi.org/10.1109/TSP.2021.3062553>.
- [48] Y. YAN, T. YANG, Z. LI, Q. LIN, AND Y. YANG, *A unified analysis of stochastic momentum methods for deep learning*, in IJCAI International Joint Conference on Artificial Intelligence, 2018.
- [49] H. YE AND X. CHANG, *Snap-shot decentralized stochastic gradient tracking methods*, arXiv preprint arXiv:2212.05273, (2022).
- [50] B. YUAN, Y. HE, J. DAVIS, T. ZHANG, T. DAO, B. CHEN, P. S. LIANG, C. RE, AND C. ZHANG, *Decentralized training of foundation models in heterogeneous environments*, Advances in Neural Information Processing Systems, 35 (2022), pp. 25464–25477.
- [51] K. YUAN AND S. A. ALGHUNAIM, *Removing data heterogeneity influence enhances network topology dependence of decentralized sgd*, 2021, <https://arxiv.org/abs/2105.08023>.
- [52] K. YUAN, Y. CHEN, X. HUANG, Y. ZHANG, P. PAN, Y. XU, AND W. YIN, *Decentlam: Decentralized momentum sgd for large-batch deep training*, in Proceedings of the IEEE/CVF International Conference on Computer Vision, 2021, pp. 3029–3039.
- [53] K. YUAN, X. HUANG, Y. CHEN, X. ZHANG, Y. ZHANG, AND P. PAN, *Revisiting optimal convergence rate for smooth and non-convex stochastic decentralized optimization*, Advances in Neural Information Processing Systems, 35 (2022), pp. 36382–36395.
- [54] S. ZHANG, S. ROLLER, N. GOYAL, M. ARTETXE, M. CHEN, S. CHEN, C. DEWAN, M. DIAB, X. LI, X. V. LIN, ET AL., *Opt: Open pre-trained transformer language models*, arXiv preprint arXiv:2205.01068, (2022).