

Propagation of dark solitons of DNLS equation along a large-scale background

A. M. Kamchatnov, D. V. Shaykin

Institute of Spectroscopy, Russian Academy of Sciences, Troitsk, Moscow, 108840, Russia

Moscow Institute of Physics and Technology, Institutsky lane 9, Dolgoprudny, Moscow region, 141700, Russia

Skolkovo Institute of Science and Technology, Skolkovo, Moscow, 143026, Russia

Abstract

We study dynamics of dark solitons in the theory of the DNLS equation by the method based on imposing the condition that this dynamics must be Hamiltonian. Combining this condition with Stokes' remark that relationships for harmonic linear waves and small-amplitude soliton tails satisfy the same linearized equations, so the corresponding solutions can be converted one into the other by replacement of the packet's wave number k by $i\kappa$, κ being the soliton's inverse half-width, we find the Hamiltonian and the canonical momentum of the soliton's motion. The Hamilton equations are reduced to the Newton equation whose solutions for some typical situations are compared with exact numerical solutions of the DNLS equation.

Keywords: integrable nonlinear wave equations, DNLS equation, soliton

PACS: 02.30.Ik, 05.45.Yv, 43.20.Bi

Dedicated to the memory of Noel Smyth

1. Introduction

In situations when the soliton's width is much smaller than the typical length of the background wave along which the soliton propagates, one can introduce with good enough accuracy the soliton's coordinate $x(t)$ and describe its propagation as a motion of a point-like particle through a non-uniform and varying with time surrounding. In this case, evolution of the background wave is governed by the equations of dispersionless (hydrodynamic) approximation independently of the soliton's motion. However, the soliton's motion cannot be separated from the background wave evolution: this motion causes a counterflow around the soliton and such a counterflow changes drastically the soliton's dynamics. Well-known examples of this back reaction on the soliton's motion are the formation of shelves behind KdV solitons propagating

along shallow water with uneven bottom (see, e.g., [1, 2, 3, 4]) and change of the frequency of oscillations of a dark soliton in a Bose-Einstein condensate confined in a harmonic trap (see, e.g., [5, 6]).

So far, the counterflow effects were studied by different forms of perturbation analysis. It was recently noticed [7], that if the nonlinear wave equation under consideration can be written in a Hamiltonian form and one assumes that the reduction of the wave evolution to the soliton's motion along the large-scale background wave remains Hamiltonian, then such an analysis can be considerably simplified. In fact, it is well known that completely integrable equations are Hamiltonian (see, e.g., [8, 9, 10]), but their Hamiltonian dynamics is formulated in terms of the scattering data or related variables in the inverse scattering transform method. Since a large-scale background wave can be represented as a collection of a large number of long waves associated with the continuous spectrum of the linear spectral problem, the assumption that the soliton's particle-like dynamics remains Hamiltonian seems very natural. This idea was recently applied to dynamics of KdV [7] and NLS [11] soli-

Email addresses: kamch@isan.troitsk.ru
(A. M. Kamchatnov), shaykin.dv@phystech.edu
(D. V. Shaykin)

tons where it was also generalized to some non-integrable situations. In this paper, our goal is to apply this method to propagation of the DNLS soliton.

The DNLS equation

$$i\psi_t + \frac{1}{2}\psi_{xx} - i(|\psi|^2\psi)_x = 0 \quad (1)$$

has applications to propagation of the nonlinear Alfvén waves in magnetized plasma (see, e.g., [12] and references therein) and of ultrashort optical pulses in nonlinear fibers (see, e.g., [13]). Complete integrability of this equation was established in Ref. [14]. We will not use here this fact explicitly (except for the assumption that the soliton’s dynamics is Hamiltonian) and we will obtain first the Hamilton equations which govern the soliton’s motion along a non-uniform and time-dependent background wave. Then we derive the Newton equation which is more convenient for applications and, at last, compare our analytical theory with numerical solutions of Eq. (1) for the initial conditions corresponding to propagation of solitons along the large-scale rarefaction waves or non-uniform profiles created by an external potential.

2. Basic equations

Substitution

$$\psi(x, t) = \sqrt{\rho(x, t)} \exp\left(i \int^x u(x', t) dx'\right) \quad (2)$$

and separation of real and imaginary parts transform Eq. (1) to the hydrodynamic-like form

$$\begin{aligned} \rho_t + \left[\rho \left(u - \frac{3}{2}\rho\right)\right]_x &= 0, \\ u_t + uu_x - (\rho u)_x + \left(\frac{\rho_x^2}{8\rho^2} - \frac{\rho_{xx}}{4\rho}\right)_x &= 0, \end{aligned} \quad (3)$$

for real variables ρ and u which can be interpreted as “density” and “flow velocity”, correspondingly. This system has two important limits.

First, we can consider small-amplitude waves propagating along a uniform background with constant values of $\rho = \rho_0$ and $u = u_0$. Then linearization of Eqs. (2) with respect to small deviations $\rho' = \rho - \rho_0$, $u' = u - u_0$ leads after standard calculations to the dispersion relation for the harmonic waves $\rho', u' \propto \exp[i(kx - \omega t)]$:

$$\omega(k) = k \left[u_0 - 2\rho_0 \pm \sqrt{\rho_0(\rho_0 - u_0) + k^2/4} \right]. \quad (4)$$

As we see, the uniform state (ρ_0, u_0) becomes modulationally unstable for $\rho_0 < u_0$, so we will only confine ourselves to the modulationally stable situations with $\rho_0 > u_0$.

Second, we can consider large-scale waves with typical wavelength much greater than the dispersion size $O(1)$ in our non-dimensional variables, $|\rho_x/\rho|, |u_x/u| \ll 1$. Then we can neglect the dispersive term in Eqs. (3) and arrive at the dispersionless equations

$$\rho_t + \left[\rho \left(u - \frac{3}{2}\rho\right)\right]_x = 0, \quad u_t + uu_x - (\rho u)_x = 0. \quad (5)$$

The characteristic velocities of this system are equal to

$$v_{\pm} = u - 2\rho \pm \sqrt{\rho(\rho - u)}. \quad (6)$$

Naturally, they are real for $\rho > u$ and coincide with the phase velocities $\omega/k|_{k \rightarrow 0}$ of linear waves with the dispersion relation (4) in the long wavelength limit. As follows from the condition $u < \rho$, they both are negative, $v_- < v_+ < 0$. Eqs. (5) can be cast to the Riemann diagonal form

$$\frac{\partial r_{\pm}}{\partial t} + v_{\pm} \frac{\partial r_{\pm}}{\partial x} = 0. \quad (7)$$

for the Riemann invariants

$$r_{\pm} = \frac{u}{2} - \rho \pm \sqrt{\rho(\rho - u)}, \quad (8)$$

and the velocities (6) are expressed in terms of Riemann invariants by the formulas

$$v_+ = \frac{3}{2}r_+ + \frac{1}{2}r_-, \quad v_- = \frac{1}{2}r_+ + \frac{3}{2}r_-. \quad (9)$$

If some solution of Eqs. (7) is found, then the physical variables are given by the expressions

$$\rho = \frac{1}{2}(\sqrt{-r_+} \pm \sqrt{-r_-})^2, \quad u = \pm 2\sqrt{r_+ r_-}. \quad (10)$$

These equations describe evolution of the background wave.

3. DNLS soliton

Soliton solutions of the DNLS equation (1) were found in Refs. [12, 14, 15]. Here we reproduce this solution in a convenient for us form.

If we look for the traveling wave solution of Eqs. (3) in the form $\rho = \rho(\xi)$, $u = u(\xi)$, $\xi = x - Vt$, V being the phase velocity of the wave,

then Eqs. (3) becomes a pair of ordinary differential equations which can be easily integrated to give

$$u(\xi) = \frac{A}{\rho(\xi)} + V + \frac{3}{2}\rho. \quad (11)$$

$$\rho_\xi^2(\xi) = -\mathcal{R}(\rho) = -[\rho^4 + 4V\rho^3 + 4(V^2 - A + B)\rho^2 - 4C\rho + 4A^2], \quad (12)$$

where A, B, C are the integration constants. The solution is parameterized more conveniently by the zeroes ν_i , $i = 1, 2, 3, 4$, of the polynomial $\mathcal{R}(\rho)$,

$$\mathcal{R}(\rho) = \prod_{i=1}^4 (\rho - \nu_i) = \rho^4 - s_1\rho^3 + s_2\rho^2 - s_3\rho + s_4, \quad (13)$$

where

$$\begin{aligned} s_1 &= \sum_i \nu_i = -4V, \\ s_2 &= \sum_{i<j} \nu_i\nu_j = 4(V^2 - A + B), \\ s_3 &= \sum_{i<j<k} \nu_i\nu_j\nu_k = 4C, \\ s_4 &= \nu_1\nu_2\nu_3\nu_4 = 4A^2. \end{aligned} \quad (14)$$

We assume that ν_i are ordered according to

$$\nu_1 \leq \nu_2 \leq \nu_3 \leq \nu_4. \quad (15)$$

The soliton solutions appear when two middle zeroes coincide, $\nu_2 = \nu_3 = \rho_0$, and then they are equal to the background density ρ_0 of the uniform state along which the soliton propagates. If ρ varies within the interval $\nu_1 \leq \rho \leq \nu_2 = \rho_0$, then we obtain a dark soliton; if within the interval $\nu_2 = \rho_0 \leq \rho \leq \nu_4$, then the soliton is bright. To be definite, we confine ourselves to the case of dark solitons with $\nu_1 \leq \rho \leq \nu_2 = \rho_0$. The standard calculation yields the solution of Eq. (12) in the form

$$\rho = \rho_0 - \frac{(\rho_0 - \nu_1)(\nu_4 - \rho_0)}{(\nu_4 - \nu_1) \cosh^2 \theta - (\rho_0 - \nu_1)}, \quad (16)$$

where $\theta = \frac{1}{2}\kappa\xi$ and

$$\kappa = \sqrt{(\rho_0 - \nu_1)(\nu_4 - \rho_0)} \quad (17)$$

is the inverse half-width of the soliton.

Let us express ν_1, ν_4 in terms of the values (ρ_0, u_0) of the parameters of the flow at infinity. Eq. (11) and the first Eq. (14) give

$$\nu_1 + \nu_4 = -4V - 2\rho_0, \quad \nu_1\nu_4 = (2u_0 - 3\rho_0 - 2V)^2,$$




Figure 1: Dependence of ν_1 and ν_4 on the soliton's velocity V ($\rho_0 = 1, u_0 = -0.5$).

so we get

$$\begin{aligned} \nu_1 &= -2V - \rho_0 - \sqrt{(\rho_0 - u_0)(u_0 - 2\rho_0 - 2V)}, \\ \nu_4 &= -2V - \rho_0 + \sqrt{(\rho_0 - u_0)(u_0 - 2\rho_0 - 2V)}. \end{aligned} \quad (18)$$

Substitution of these values into Eq. (17) yields

$$\kappa^2 = 4 [\rho_0(\rho_0 - u_0) - (V + 2\rho_0 - u_0)^2]. \quad (19)$$

Whence, the soliton velocity V is related with the inverse half-width κ by the formula

$$V = u_0 - 2\rho_0 \pm \sqrt{\rho_0(\rho_0 - u_0) - \frac{\kappa^2}{4}}. \quad (20)$$

Comparison with Eq. (4) shows that the soliton's velocity is related with the dispersion law of linear waves by the expression

$$V = \frac{\omega(i\kappa)}{i\kappa}. \quad (21)$$

This is a particular case of the general Stokes' remark [16] (see also [17, 18]) that both linear waves $\propto \exp[i(kx - \omega t)]$ and soliton's tails $\propto \exp[\pm\kappa(x - Vt)]$ are described by the linearized equations, so the phase velocity $\omega(k)/k$ transforms to the soliton's velocity by the replacement $k \rightarrow i\kappa$.

If we rewrite Eq. (19) in the form

$$\begin{aligned} \kappa^2 &= 4 \left[\sqrt{\rho_0(\rho_0 - u_0) - 2\rho_0 + u_0 - V} \right] \\ &\quad \times \left[\sqrt{\rho_0(\rho_0 - u_0) + 2\rho_0 - u_0 + V} \right], \end{aligned} \quad (22)$$

we find at once that the soliton's velocity V must have the value between the dispersionless characteristic velocities (5),

$$v_- < V < v_+, \quad (23)$$

that is it is always negative. The parameters ν_1 and ν_4 are always positive since positive density at the center of soliton is equal to $\rho_{min} = \nu_1$ and $\nu_4 > \nu_1$. One can easily find that at the edges of the interval (23) the parameters ν_1 and ν_4 are equal to

$$\nu_1(v_-) = \rho_0, \quad \nu_1(v_+) = \begin{cases} \rho_0, & \text{for } u_0 < 0, \\ 5\rho_0 - 4u_0 \\ -4\sqrt{\rho_0(\rho_0 - u_0)}, & \text{for } 0 < u_0 < \rho_0; \end{cases} \quad (24)$$

and

$$\nu_4(v_-) = 5\rho_0 - 4u_0 + 4\sqrt{\rho_0(\rho_0 - u_0)}, \quad \nu_4(v_+) = \begin{cases} 5\rho_0 - 4u_0 \\ -4\sqrt{\rho_0(\rho_0 - u_0)}, & \text{for } u_0 < 0, \\ \rho_0, & \text{for } 0 < u_0 < \rho_0. \end{cases} \quad (25)$$

Plots of $\nu_1(V)$ and $\nu_4(V)$ are shown in Fig. 1; $\nu_1(V)$ has vanishing minimal value at

$$V = V_m = u_0 - \frac{3}{2}\rho_0. \quad (26)$$

One can easily find that this point is located inside the interval (23), that is $V_m \leq v_+(u_0)$, for $u_0 < u_m = \frac{3}{4}\rho_0$, and $v_+(u_0)$ has its maximal value $v_+(u_m) = -\frac{1}{8}\rho_0$ at this point. Thus, for $u_0 < \frac{3}{4}\rho_0$ we should distinguish two intervals of the soliton's velocity:

$$\begin{aligned} \text{(A): } & u_0 - \frac{3}{2}\rho_0 < V < v_+, \quad V + \frac{3}{2}\rho_0 - u_0 > 0, \\ \text{(B): } & v_- < V < u_0 - \frac{3}{2}\rho_0, \quad V + \frac{3}{2}\rho_0 - u_0 < 0. \end{aligned} \quad (27)$$

If $\frac{3}{4}\rho_0 < u_0 < \rho_0$, then we get a single interval

$$\text{(C): } v_- < V < v_+, \quad V + \frac{3}{2}\rho_0 - u_0 < 0. \quad (28)$$

These different intervals of the soliton's velocity are depicted in Fig. 2; such their separation is important for calculation of the soliton's phase.

The soliton's phase in the solution $\psi = \sqrt{\rho} \exp(i\phi)$ can be found by integration of Eq. (11), that is

$$\phi = \int \left(\frac{\rho_0 \sqrt{\nu_1 \nu_4}}{\rho} + V + \frac{3}{2}\rho \right) d\xi, \quad (29)$$




Figure 2: Regions of the soliton's velocity V at different values of u_0 and fixed ρ_0 with different expressions for soliton's phase. They are confined between the boundaries (23) and separated by the lines $V_m(u_0) = u_0 - \frac{3}{2}\rho_0$ and $u_0 = \frac{3}{4}\rho_0$.

where $\rho = \rho(\xi)$ is given in Eq. (16). Quite a tedious, although straightforward, calculation yields

$$\begin{aligned} \phi_A &= u_0 \xi - \arctan \left(\frac{1}{\gamma} \sqrt{\frac{\nu_4}{\nu_1}} \tanh \frac{\kappa \xi}{2} \right) \\ &\quad - 3 \arctan \left(\frac{1}{\gamma} \tanh \frac{\kappa \xi}{2} \right), \\ \phi_{B,C} &= u_0 \xi + \arctan \left(\frac{1}{\gamma} \sqrt{\frac{\nu_4}{\nu_1}} \tanh \frac{\kappa \xi}{2} \right) \\ &\quad - 3 \arctan \left(\frac{1}{\gamma} \tanh \frac{\kappa \xi}{2} \right), \end{aligned} \quad (30)$$

where

$$\gamma = \sqrt{\frac{\nu_4 - \rho_0}{\rho_0 - \nu_1}}. \quad (31)$$

Obviously, $u_0 \xi$ is the background phase over which the soliton's phase is imposed. The jump of the phase across the soliton is equal to

$$\begin{aligned} \Delta \phi_A &= -2 \arctan \left(\frac{1}{\gamma} \sqrt{\frac{\nu_4}{\nu_1}} \right) - 6 \arctan \frac{1}{\gamma} + 2\pi, \\ \Delta \phi_{B,C} &= 2 \arctan \left(\frac{1}{\gamma} \sqrt{\frac{\nu_4}{\nu_1}} \right) - 6 \arctan \frac{1}{\gamma}. \end{aligned} \quad (32)$$

Its dependence on the soliton's velocity V is shown in Fig. 3 for positive and negative signs of the background velocity u_0 . At $V = v_+$ we get $\Delta \phi = \text{sgn}(u_0)2\pi$ for all u_0 , but $\Delta \phi \sim |u_0|$, $u_0 \rightarrow 0$, tends to zero non-uniformly almost everywhere except a decreasing interval $v_+ - |u_0| < V \leq v_+$ close to the edge $V = v_+$. So one can consider $\Delta \phi = 0$ at $u_0 = 0$ for all $V < v_+$.




Figure 3: Jump of phase across the soliton as a function of its velocity V . It equals to $\Delta\phi(v_+) = \text{sgn}(u_0)2\pi$ at the edge $V = v_+$.

Combining expressions (16) for ρ and (30) for ϕ , we arrive at the soliton solution

$$\psi_{A,B,C} = \frac{\sqrt{\nu_4} \tanh(\kappa\xi/2) \pm i\sqrt{\nu_1}\gamma}{[\gamma + i \tanh(\kappa\xi/2)]^3} \times \left(\gamma^2 + \tanh^2 \frac{\kappa\xi}{2} \right) e^{iu_0\xi}, \quad (33)$$

where “+” corresponds to the region A and “-” to the regions B,C. We will use this expression in section 5 as an initial condition in numerical solutions of Eq. (1).

4. Hamilton equations for soliton motion

We assume that soliton’s width is much smaller than the characteristic background wavelength. Therefore, we can calculate the soliton’s Hamiltonian H and the canonical momentum p under assumption that the background variables ρ, u have their values $\rho(x, t), u(x, t)$ at the location $x(t)$ of the soliton at the moment t . Then Eq. (20) can be written as one of the Hamilton equations

$$\frac{dx}{dt} = u - 2\rho + \sqrt{\rho(\rho - u) - \kappa^2/4} = \frac{\partial H}{\partial p}. \quad (34)$$

It is convenient to introduce the variable ϕ instead of κ ,

$$\kappa = 2\sqrt{\rho(\rho - u)} \sin \phi, \quad (35)$$

so that

$$\frac{dx}{dt} = \frac{\partial H}{\partial p} = u - 2\rho + \sqrt{\rho(\rho - u)} \cos \phi. \quad (36)$$

Another important relation follows from preservation of the Poincaré-Cartan integral invariant by the dispersionless flow (5) (see Refs. [7, 11]). This means that in the completely integrable equations, as the DNLS equation, the wave number k of a short wavelength packet propagating along large-scale background waves is only a function of the local values of ρ and u , $k = k(\rho, u)$, which satisfies in the DNLS case the equations [19]

$$\begin{aligned} \frac{\partial k^2}{\partial \rho} &= 4 \left[\sqrt{k^2 + 4\rho(\rho - u)} - (2\rho - u) \right], \\ \frac{\partial k^2}{\partial u} &= -2 \left[\sqrt{k^2 + 4\rho(\rho - u)} - 2\rho \right], \end{aligned} \quad (37)$$

with the solution

$$k^2 = (q + 2\rho - u)^2 - 4\rho(\rho - u), \quad (38)$$

where q is an integration constant determined by the value of k at some fixed point in the (ρ, u) -plane. Then, using the Stokes reasoning (see Eq. (21) and the text below it), we find that the inverse half-width κ is related with the local background variables by the formula

$$\kappa^2 = 4\rho(\rho - u) - (q + 2\rho - u)^2. \quad (39)$$

Its substitution into Eq. (34) with account of Eq. (36) give the integral of motion

$$u/2 - \rho + \sqrt{\rho(\rho - u)} \cos \phi = q/2, \quad (40)$$

where we assumed for definiteness that $q + 2\rho - u > 0$. Differentiation of this identity along soliton’s path yields

$$\begin{aligned} \frac{1}{2} \frac{du}{dt} - \frac{d\rho}{dt} - \sqrt{\rho(\rho - u)} \sin \phi \frac{d\phi}{dt} \\ + \frac{1}{2\sqrt{\rho(\rho - u)}} \left[(2\rho - u) \frac{d\rho}{dt} - \rho \frac{du}{dt} \right] = 0, \end{aligned}$$

where $d/dt = \partial/\partial t + V\partial/\partial x$. We eliminate ρ_t and u_t with help of Eqs. (5) and find after elementary transformations the formula

$$\frac{d\phi}{dt} = - \left(\sqrt{\rho(\rho - u)} \right)_x \sin \phi. \quad (41)$$

Now we can turn to derivation of the Hamiltonian H and the canonical momentum p for soliton’s motion. We look for the momentum in the form

$$p = \rho(\rho - u)f(\phi), \quad (42)$$

where $f(\phi)$ is unknown function. Then integration of Eq. (36) with respect to p gives

$$H = (u - 2\rho)p + [\rho(\rho - u)]^{3/2} \int \cos \phi f' d\phi. \quad (43)$$

To find $f(\phi)$, we use another Hamilton equation

$$\frac{dp}{dt} = -\frac{\partial H}{\partial x}. \quad (44)$$

Substitution of Eqs. (42) and (43) with the use of Eq. (41) leads to the equation

$$f' \sin \phi = 3 \int^\phi \cos \phi f' d\phi, \quad (45)$$

which can be easily integrated to give

$$f = C(\phi - \sin \phi \cos \phi), \quad (46)$$

where C is an integration constant which appearance reflects invariance of the Hamilton equations with respect to multiplication of p and H by the same constant factor. We choose it equal to $C = 2$ and express ϕ in terms of $\dot{x} = dx/dt = V$ and the local variables $\rho = \rho(x, t)$, $u = u(x, t)$ with help of Eq. (36). As a result, we obtain

$$p = 2\rho(\rho - u) \arccos \frac{\dot{x} + 2\rho - u}{\sqrt{\rho(\rho - u)}} - 2(\dot{x} + 2\rho - u) \sqrt{\rho(\rho - u) - (\dot{x} + 2\rho - u)^2}, \quad (47)$$

and

$$H = (u - 2\rho)p + \frac{4}{3} [\rho(\rho - u) - (\dot{x} + 2\rho - u)^2]^{3/2}. \quad (48)$$

It is worth noticing that these expressions reduce to similar expressions for the NLS soliton by means of replacements $\rho(\rho - u) \rightarrow \rho_{NLS}$, $u - 2\rho \rightarrow u_{NLS}$ (see [11]).

For practical applications, it is convenient to derive the Newton equation for the soliton's motion.

5. Newton equation

In Hamiltonian mechanics, $\dot{x} = \dot{x}(x, p)$, but it is impossible to solve explicitly Eq. (47) with respect to \dot{x} . Therefore, it is more convenient to transform the Hamilton equation (44) to the Newton equation for the soliton's acceleration \ddot{x} . Differentiation of

Eq. (47) with respect to time t gives after obvious simplifications

$$\begin{aligned} \frac{dp}{dt} = & 2 \left[(2\rho - u) \frac{d\rho}{dt} - \rho \frac{du}{dt} \right] \arccos \frac{\dot{x} + 2\rho - u}{\sqrt{\rho(\rho - u)}} \\ & - 4 \left(\dot{x} + 2 \frac{d\rho}{dt} - \frac{du}{dt} \right) \sqrt{\rho(\rho - u) - (\dot{x} + 2\rho - u)^2}. \end{aligned} \quad (49)$$

The right-hand side of Eq. (44) reads

$$\begin{aligned} -\frac{\partial H}{\partial x} \Big|_p = & -(u_x - 2\rho_x)p - \frac{\partial H^{(0)}}{\partial \rho} \Big|_{\dot{x}} \rho_x \\ & - \frac{\partial H^{(0)}}{\partial u} \Big|_{\dot{x}} u_x - \frac{\partial H^{(0)}}{\partial \dot{x}} \frac{\partial \dot{x}}{\partial x} \Big|_p, \end{aligned} \quad (50)$$

where $H^{(0)} = \frac{4}{3} [\rho(\rho - u) - (\dot{x} + 2\rho - u)^2]^{3/2}$. The derivative $\frac{\partial \dot{x}}{\partial x} \Big|_p$ is to be obtained by differentiation of Eq. (47) with respect to x at constant p and this gives

$$\begin{aligned} \frac{\partial \dot{x}}{\partial x} \Big|_p = & \frac{(\rho^2 - \rho u)_x}{2\sqrt{\rho(\rho - u) - (\dot{x} + 2\rho - u)^2}} \\ & \times \arccos \frac{\dot{x} + 2\rho - u}{\sqrt{\rho(\rho - u)}} + u_x - 2\rho_x. \end{aligned} \quad (51)$$

The other derivatives are trivial. Substitution of all the derivatives into Eq. (50) followed by equating the result to Eq. (49) and elimination of ρ_t and u_t with the use of the dispersionless equations

$$\begin{aligned} \rho_t + \left[\rho \left(u - \frac{3}{2}\rho \right) \right]_x &= 0, \\ u_t + uu_x - (\rho u)_x &= -U_x, \end{aligned} \quad (52)$$

where $U(x)$ is the potential of external forces, yield after some simplifications the Newton equation

$$\begin{aligned} 2\ddot{x} = & \frac{\rho U_x}{\sqrt{\rho(\rho - u) - (\dot{x} + 2\rho - u)^2}} \\ & \times \arccos \frac{\dot{x} + 2\rho - u}{\sqrt{\rho(\rho - u)}} + \frac{d}{dt}(2\rho - u) - U_x. \end{aligned} \quad (53)$$

If there is no external potential, then the equation

$$2\ddot{x} = \frac{d}{dt}(2\rho - u) \quad (54)$$

gives at once the integral of motion

$$2\dot{x} - 2\rho + u = \text{const}. \quad (55)$$




Figure 4: Initial distribution of ρ corresponding to interaction of a dark soliton with a rarefaction wave; $\rho_L = 2$, $\rho_R = 1$, $l = 30$, $V_0 = -2.5$.




Figure 5: Path of soliton along the rarefaction wave for initial parameters indicated in Fig. 4. Solid line corresponds to the analytical solution (59), dots to the numerical solution, and dashed line describes propagation of the soliton with constant initial velocity V_0 .

Actually, it follows easily from Eqs. (36) and (40) written for the case of a uniform background without external forces. This equation is convenient for finding the soliton's paths for motion of solitons along rarefaction waves and other large-scale waves evolving without influence of external forces. Its generalization (53) is derived under the assumption that non-uniformity of the background produced by the external potential is more essential than change of the soliton's energy caused by its deformation: in our derivation of the Hamiltonian we supposed that the soliton keeps its form in slightly non-uniform background.

To check our analytical theory, we compared it with the results of exact numerical solutions of the

DNLS equation (1) for two typical physical situations. In the first one, the soliton was propagating along a rarefaction wave produced by an initial discontinuity in the distribution of ρ (see Fig. 4). In such a rarefaction wave the Riemann invariant (see Eq. (8)) $r_+ = 0$ remains constant and $r_- = -2\rho$, so that $v_- = -3\rho$. The self-similar solution of Eqs. (7) has the form $v_- = x/t$ and this gives the distribution

$$\rho(x, t) = \begin{cases} \rho_L, & x < -3\rho_L t, \\ -\frac{x}{3t}, & -3\rho_L t < x < -3\rho_R t, \\ \rho_R, & x > -3\rho_R t, \end{cases} \quad (56)$$

We assume that the left edge of the rarefaction wave propagates to the left with velocity $v_L = -3\rho_L$ faster, than the initial soliton velocity V_0 . If the initial distance between the soliton and the discontinuity equal to l , then after time $t_1 = l/|v_L - V_0|$ the soliton starts its propagation along the rarefaction wave, so its motion is governed by Eq. (55),

$$\frac{dx}{dt} + \frac{x}{3(t + t_1)} = V_0 - \rho_L, \quad (57)$$

whose solution must satisfy the initial condition

$$x(0) = -l + V_0 t_1. \quad (58)$$

This equation can be easily solved and the solution reads

$$x(t) = \frac{3}{4}(V_0 - \rho_L)(t + t_1) - \frac{3l}{4} \frac{t_1^{2/3}}{(t + t_1)^{1/3}}. \quad (59)$$

As one can see, interaction of the soliton with the rarefaction wave leads to quite a considerable effect and our analytical theory agrees very well with the numerical solution of the DNLS equation.

As another physical situation, we consider flow of the background past an obstacle created by the potential

$$U(x) = 20 \exp \left[- \left(\frac{x}{200} \right)^2 \right]. \quad (60)$$

The flow is stationary, so $\rho = \rho(x)$, $u = u(x)$ do not depend on time t and they should be obtained by solving the stationary dispersionless Eqs. (52). An easy calculation (see, e.g., analogous problem for the NLS equation in Refs. [20, 21]) yields the solution in implicit form

$$F(u(x)) = U(x), \quad (61)$$




Figure 6: Initial distribution of ρ corresponding to interaction of a dark soliton with an obstacle created by the potential (60).




Figure 7: Path of a soliton moving past an obstacle (60). Solid line corresponds to the solution of Eq. (53), dots to the numerical solution, and dashed line describes propagation of the soliton with constant initial velocity V_0 .

that is $u(x) = F^{-1}(U(x))$, where

$$F(u) = \frac{1}{3}u\sqrt{9\rho_0^2 - 6\rho_0u_0 + u^2} - \frac{1}{6}u^2 + \frac{1}{2}u_0^2 - \rho_0u_0. \quad (62)$$

The Newton equation

$$2\ddot{x} = \frac{\rho U_x}{\sqrt{\rho(\rho - u) - (\dot{x} + 2\rho - u)^2}} \times \arccos \frac{\dot{x} + 2\rho - u}{\sqrt{\rho(\rho - u)}} + 2\rho_x - u_x - U_x \quad (63)$$

should be solved with the initial conditions

$$x(0) = l, \quad \dot{x} = V_0, \quad (64)$$

and the result is shown in Fig. 7 by a solid line, whereas dots correspond to the numerical solution

of Eq. (1) with the same initial condition in the form of the soliton (33) on the stationary background flow. Dashed line corresponds to the soliton moving with constant initial velocity V_0 . Again the analytical theory agrees very well with the numerical solution.

6. Conclusion

The problem of soliton's motion along non-uniform and time-dependent background, especially in presence of external forces, is very difficult because of back reaction of the counterflow caused by a moving soliton in the large-scale background wave. Previously, different versions of the perturbation theory were developed for solving this problem, but they were quite complicated and therefore applied to a very limited number of equations (mainly, KdV and NLS equations). We suggested in Ref. [7] a new approach. Although it is basically also perturbative, many difficulties are removed by imposing the condition that the equations of soliton's motion must be Hamiltonian. Combining this condition with Stokes' remark that some relationships for harmonic linear wave can be converted into the relationships for solitons by replacement of the packet's wave number k by $i\kappa$, κ being the soliton's inverse half-width, we reduce derivation of the Hamiltonian and the canonical momentum of the soliton's motion to a straightforward calculation. The resulting Hamilton equations can be transformed to the Newton equation where the role of the external potential can be taken into account by its inclusion into the equations of the dispersionless (hydrodynamic) flow. The effectiveness of this method was demonstrated in Ref. [11] for the NLS dark soliton, where the results of Ref. [21] were easily reproduced. In the present paper, we applied this method to the DNLS soliton which has quite a nontrivial dynamics without reflection symmetry. Validity of our approach is confirmed by its good agreement with exact numerical solutions of the DNLS equation. We believe that this approach can find many other applications.

Acknowledgments

This research is funded by the research project FFUU-2024-0003 of the Institute of Spectroscopy of the Russian Academy of Sciences (Sections 1–3) and by the RSF grant number 19-72-30028 (Section 4–6).

References

- [1] V. I. Karpman, E. M. Maslov, Zh. Eksp. Teor. Fiz. 73 (1977) 537 [Sov. Phys. JETP, 46 (1977) 281].
- [2] V. I. Karpman, E. M. Maslov, Zh. Eksp. Teor. Fiz. 75 (1978) 504 [Sov. Phys. JETP, 48 (1978) 252].
- [3] C. J. Knickerbocker, A. C. Newell, J. Fluid Mech. 98 (1980) 803.
- [4] A. C. Newell, *Solitons in Mathematics and Physics*, (SIAM, Philadelphia, 1985).
- [5] Th. Busch and J. R. Anglin, Phys. Rev. Lett. 84 (2000) 2298.
- [6] V. V. Konotop and L. P. Pitaevskii, Phys. Rev. Lett. 93 (2004) 240403.
- [7] A. M. Kamchatnov and D. V. Shaykin, Phys. Rev. E 108 (2023) 054205.
- [8] V. E. Zakharov, L. D. Faddeev, Funk. Analiz Prilozh., 5 (1971) 18 [Func. Anal. Appl. 5 (1971) 280].
- [9] C. S. Gardner, J. Math. Phys. 12 (1971) 1548.
- [10] L. A. Dickey, *Soliton Equations and Hamiltonian Systems*, (Singapore, World Scientific, 2003).
- [11] A. M. Kamchatnov, Theor. Math. Phys. (to be published, 2024); (2023), arXiv:2310.04001.
- [12] C. F. Kennel, B. Buti, T. Hada, and R. Pellat, Phys. Fluids, 31 (1988) 1949.
- [13] S. A. Akhmanov, V. A. Vysloukh, and A. S. Chirkin, Usp. Fiz. Nauk, 149 (1986) 449; [Sov. Phys. Uspekhi, 29, (1986) 642].
- [14] D. J. Kaup and A. C. Newell, J. Math. Phys. 19 (1978) 798.
- [15] A. M. Kamchatnov, J. Phys. A: Math. Gen., 23 (1990) 2945.
- [16] G. G. Stokes, *Mathematical and Physical Papers*, Vol. V, p. 162 (Cambridge University Press, Cambridge, 1905).
- [17] H. Lamb, *Hydrodynamics*, (Cambridge University Press, Cambridge, 1932).
- [18] A. M. Kamchatnov, Chaos, 30 (2020) 123148.
- [19] A. M. Kamchatnov and D. V. Shaykin, Physica D, 460 (2024) 134085.
- [20] A. M. Leszczyszyn, G. A. El, Yu. G. Gladush, and A. M. Kamchatnov, Phys. Rev. A 79 (2009) 063607.
- [21] S. K. Ivanov and A. M. Kamchatnov, Chaos 32 (2022) 113142.