Improved Space Bounds for Subset Sum

Tatiana Belova * Nikolai Chukhin † Alexander S. Kulikov ‡ Ivan Mihajlin §

Abstract

More than 40 years ago, Schroeppel and Shamir presented an algorithm that solves the Subset Sum problem for n integers in time $O^*(2^{0.5n})$ and space $O^*(2^{0.25n})$. The time upper bound remains unbeaten, but the space upper bound has been improved to $O^*(2^{0.249999n})$ in a recent breakthrough paper by Nederlof and Węgrzycki (STOC 2021). Their algorithm is a clever combination of a number of previously known techniques with a new reduction and a new algorithm for the Orthogonal Vectors problem.

In this paper, we give two new algorithms for Subset Sum. We start by presenting an Arthur–Merlin algorithm: upon receiving the verifier's randomness, the prover sends an n/4-bit long proof to the verifier who checks it in (deterministic) time and space $O^*(2^{n/4})$. An interesting consequence of this result is the following fine-grained lower bound: assuming that 4-SUM cannot be solved in time $O(n^{2-\varepsilon})$ for all $\varepsilon > 0$, Circuit SAT cannot be solved in time $O(g2^{(1-\varepsilon)n})$, for all $\varepsilon > 0$ (where n and g denote the number of inputs and the number of gates, respectively).

Then, we improve the space bound by Nederlof and Wegrzycki to $O^*(2^{0.246n})$ and also simplify

Then, we improve the space bound by Nederlof and Wegrzycki to $O^*(2^{0.246n})$ and also simplify their algorithm and its analysis. We achieve this space bound by further filtering sets of subsets using a random prime number. This allows us to reduce an instance of Subset Sum to a larger number of instances of smaller size.

^{*}St. Petersburg Department of Steklov Mathematical Institute, Russian Academy of Sciences. Email: yukikomodo@gmail.com.

[†]Neapolis University Pafos and JetBrains Research. Email: buyolitsez1951@gmail.com

[‡]JetBrains Research. Email: alexander.s.kulikov@gmail.com.

[§]JetBrains Research. Email: ivmihajlin@gmail.com.

Contents

1	Overview		
	1.1	Known Results	3
		1.1.1 Time Complexity	3
		1.1.2 Space Complexity	3
		1.1.3 Space Complexity	3
		1.1.4 Proof Complexity	4
	1.2	New Results	4
		1.2.1 New Arthur–Merlin Algorithm	4
		1.2.2 New Conditional Lower Bounds for Circuit SAT	5
		1.2.3 Improved Space Upper Bound for Subset Sum	5
2	Ger	neral Setting	6
	2.1	Modular Arithmetic	6
	2.2	Prime Numbers	6
	2.3	Probability Amplification	6
	2.4		6
	2.5	Sets and Sums	7
	2.6	Weighted Orthogonal Vectors	7
3	Nev	w Arthur-Merlin Algorithm	8
4	Imp	proved Space Upper Bound for Subset Sum	9
	4.1	Algorithm	9
	4.2	Correctness	4
	4.3	Setting the Parameters	5
	4.4	Time Complexity	6
	4.5	Space Complexity	8
	4.6	Parameter Substitution	9
	4.7	Algorithm for Weighted Orthogonal Vectors	O

1 Overview

In this paper, we study the well-known Subset Sum problem and its parameterized version, k-SUM. In Subset Sum, given a set of n integers and a target integer t the goal is to check whether there is a subset of them that sum up to t. It is common to assume that the absolute value of all input integers is at most $2^{O(n)}$ (one can achieve this using hashing techniques).

In the k-SUM problem, the goal is to check whether some k of the input integers sum to zero. We consider a slightly different formulation of k-SUM: given k sequences A_1, \ldots, A_k , each containing n integers of absolute value at most $n^{O(1)}$, check whether there exist indices i_1, \ldots, i_k such that $A_1[i_1] + \cdots + A_k[i_k] = 0$. It is common to further assume that the bit-length of all integers in the k-SUM problem is at most $k \log n + O(\log \log n)$. This can be achieved by the standard fingerprinting technique.

1.1 Known Results

1.1.1 Time Complexity

2-SUM can be solved in time O(n) using binary search (the $O(\cdot)$ notation hides polylogarithmic factors): sort A_1 ; then, for each $1 \le i_2 \le n$, check whether A_1 contains an element $-A_2[i_2]$. Another well known algorithm is based on the two pointers method that proceeds as follows. Sort A_1 and A_2 and let $i_1 = 1$ and $i_2 = n$. Then, keep repeating the following: if $i_1 > n$ or $i_2 < 1$, stop; if $A_1[i_1] + A_2[i_2] = 0$, return (i_1, i_2) ; if $A_1[i_1] + A_2[i_2] > 0$, decrement i_2 (as $A_2[i_2]$ is too large: its sum with the currently smallest element of A_1 is positive); if $A_1[i_1] + A_2[i_2] < 0$, increment i_1 (as $A_1[i_1]$ is too small: its sum with the currently largest element of A_2 is negative).

The algorithm for 2-SUM allows k-SUM to be solved in time $\widetilde{O}(n^{\lceil k/2 \rceil})$ via the following reduction. Given k arrays A_1, \ldots, A_k , split them into two halves: $A_1, \ldots, A_{\lfloor k/2 \rfloor}$ and $A_{\lceil k/2 \rceil}, \ldots, A_k$. Populate arrays B_1 and B_2 with all sums of elements from the first and second halves, respectively. Then, it remains to solve 2-SUM for B_1 and B_2 .

For 3-SUM, Chan [4] proved a slightly better upper bound $O(n^2 \log \log^{O(1)} n / \log^2 n)$. It is a major open problem whether 3-SUM can be solved in time $O(n^{2-\varepsilon})$ and the 3-SUM hypothesis, stating that this is impossible, is a popular hypothesis in the field of fine-grained complexity. It is also currently unknown whether k-SUM can be solved in time $\widetilde{O}(n^{\lceil k/2 \rceil - \varepsilon})$ for any $k \geq 3$ and $\varepsilon > 0$.

The 2-SUM algorithm described above is also at the core of the strongest known upper bound $O^*(2^{n/2})$ for Subset Sum $(O^*(\cdot))$ hides polynomial factors) presented by Horowitz and Sahni [6] fifty years ago. To get this running time, partition A into two halves of size n/2; then, populate arrays A_1 and A_2 (of size $2^{n/2}$) with sums of all subsets of the two halves, respectively; then, it remains to solve 2-SUM for A_1 and A_2 . It is an important open problem to solve Subset Sum in $O^*(2^{(1/2-\varepsilon)n})$ for a constant $\varepsilon > 0$. The fastest known algorithm shows that Subset Sum can be solved in time $2^{n/2}/\operatorname{poly}(n)$ [5].

1.1.2 Space Complexity

1.1.3 Space Complexity

The algorithms that solve k-SUM and Subset Sum via a reduction to 2-SUM have high space complexity: for k-SUM, it is $O(n^{\lceil k/2 \rceil})$, whereas for Subset Sum it is $O(2^{n/2})$. It is natural to ask whether one can lower the space complexity or whether it is possible to trade off time and space. Lincoln et al. [10] gave a positive answer for k-SUM: one can solve 3-SUM in time $O(n^2)$ and space $O(\sqrt{n})$ as well as k-SUM (for $k \geq 4$) in time $O(n^{k-2+2/k})$ and space $O(\sqrt{n})$. For Subset Sum, the well-known algorithm by Schroeppel and Shamir [14] solves it in time $O^*(2^{n/2})$ and space $O(2^{n/4})$: they reduce Subset Sum to 4-SUM and note that 4-SUM can be solved in time $\widetilde{O}(n^2)$ and space O(n) (since all pairwise sums of two sorted sequences of length n can be enumerated in time $\widetilde{O}(n^2)$ and space O(n) using a priority queue data structure). Just recently, Nederlof and Wegrzycki improved the space complexity to $O(2^{0.249999n})$ [12].

¹These two versions of k-SUM are reducible to each other. One direction is easy: take k copies of the array A. For the other direction, one can use the color coding technique [2]: for every element A, add it to A_i for random $i \in [k]$; then, if a solution for the original problem exists, it survives in the resulting instance with constant probability. This randomized reduction can also be derandomized [2].

1.1.4 Proof Complexity

It is easy to certify a yes-instance of k-SUM or Subset Sum: just write down a solution; it can be checked in deterministic time O(n). Certifying no-instances is trickier. Carmosino et al. [3] showed that, for 3-SUM, there are proofs of size $\widetilde{O}(n^{1.5})$ that can be deterministically verified in time $\widetilde{O}(n^{1.5})$. By allowing the verifier to be probabilistic, Akmal et al. [1] presented a proof system for 3-SUM where the proof's size and the verification time is $\widetilde{O}(n)$. For k-SUM, they give an upper bound of $\widetilde{O}(n^{k/3})$. The corresponding proof system is known as a Merlin–Arthur protocol: a computationally unbounded prover (Merlin) prepares a proof and sends it to a probabilistic verifier (Arthur) who needs to be able to check the proof quickly with small constant error probability. We say that a problem can be solved in Merlin–Arthur time T(n), if there exists a protocol where both the proof size and the verification time do not exceed T(n).

For Subset Sum, Nederlof [11] proved an upper bound $O^*(2^{(1/2-\varepsilon)n})$, for some $\varepsilon > 10^{-3}$, on Merlin–Arthur time. Akmal et al. [1] improved the bound to $O^*(2^{n/3})$.

1.2 New Results

Below, we give an overview of the main results of the paper.

1.2.1 New Arthur-Merlin Algorithm

Recall that in an Arthur–Merlin protocol the randomness is shared (also known as public coins) and the verifier is deterministic: upon receiving the verifier's randomness, the prover prepares a proof and sends it to the verifier who checks it deterministically with small error probability (taken over public randomness). Formally, we say that a language $L \subseteq \{0,1\}^*$ belongs to a class $\mathsf{AM}[s(n),t(n)]$, if there exists an Arthur–Merlin protocol such that, for any $x \in \{0,1\}^n$, the proof length is at most s(n), the verification time is at most t(n), and for each $x \in L$, the verifier accepts with probability at least 2/3, whereas for each $x \notin L$, the verifier rejects with probability 1.2 Such a protocol implies that L can be solved by a randomized algorithm that has running time t(n) and error probability at least $2^{-s(n)}$. It is also easily parallelizable: using $k \le 2^{s(n)}$ processors, the problem can be solved in time $\frac{2^{s(n)}t(n)}{k}$. Our first main result is a new Arthur–Merlin algorithm for Subset Sum.

Theorem 1.1. Subset
$$Sum \in AM\left[n/4, O^*(2^{n/4})\right]$$
.

As discussed above, it can be parallelized easily: to solve the problem, one can enumerate possible proofs in parallel. Also, by enumerating all possible proofs, one recovers upper bounds on time and space for Subset Sum proved by Schroeppel and Shamir in 1979. Interestingly, the resulting algorithm is very simple and does not need to use the priority queue data structure as in the algorithm by Schroeppel and Shamir.

As it is the case with the previously known algorithms for Subset Sum, our algorithm follows from an algorithm for 4-SUM.

Theorem 1.2.
$$4$$
-SUM \in AM $\left[\log_2 n, \widetilde{O}(n)\right]$.

Since 2k-SUM can be reduced to 4-SUM, this extends to 2k-SUM as follows.

Corollary 1.1. For every even integer
$$k$$
, $2k$ -SUM \in AM $\left[\frac{k}{2}\log_2 n, \widetilde{O}(n^{k/2})\right]$.

The main idea of the proof of Theorem 1.2 is the following. To find integers i_1, i_2, i_3, i_4 such that $A_1[i_1] + A_2[i_2] + A_3[i_3] + A_4[i_4] = 0$, one generates a random prime $p \le n$ and uses $(A_1[i_1] + A_2[i_2]) \mod p$ as a proof. A similar idea of using a random prime for filtering subsets was used previously by Howgrave-Graham and Joux [7].

 $^{^2}$ In the standard definition of AM protocols, there is either a two-sided error probability or zero error probability of acceptance. We choose to have a zero error probability of rejection as it allows for straightforward transformation to randomized and parallel algorithms.

1.2.2 New Conditional Lower Bounds for Circuit SAT

Another interesting consequence of Theorem 1.2 is a fine-grained lower bound for the Circuit SAT problem. Circuit SAT is a generalization of SAT where instead of a formula in CNF, one is given an arbitrary Boolean circuit with g binary gates and n inputs (and the goal, as usual, is to check whether it is satisfiable). Being a generalization of SAT, Circuit SAT cannot be solved in time $O^*(2^{(1-\varepsilon)n})$, for any constant $\varepsilon > 0$, under the Strong Exponential Time Hypothesis (that states that for every $\varepsilon > 0$, there exists k such that k-SAT cannot be solved in time $O(2^{(1-\varepsilon)n})$). Even designing a $2^{(1-\varepsilon)n}$ time algorithm solving Circuit SAT for circuits with $g \leq 8n$ gates is challenging: as shown by [8], this would imply new circuit lower bounds. We prove the following conditional lower bound for Circuit SAT.

Theorem 1.3. If, for any $\varepsilon > 0$, 4-SUM cannot be solved in time $O(n^{2-\varepsilon})$, then, for any $\varepsilon > 0$, Circuit SAT cannot be solved in time $O(g2^{(1-\varepsilon)n})$.

This theorem however does not exclude a $O(g^{O(1)}2^{(1-\varepsilon)n})$ -time algorithm for Circuit-SAT in the same manner. To get such a fine-grained lower bound, one needs to push the algorithm from Corollary 1.1 further (i.e., to trade most of the n^k time upper bound for nondeterminism).

Open Problem 1.1. Prove or disprove: 2k-SUM \in AM $\left[k \log_2 n, n^{o(k)}\right]$.

Corollary 1.2. Assume that 2k-SUM \in AM $\left[k\log_2 n, n^{o(k)}\right]$. Then, if for any $\varepsilon > 0$ and any $k \geq 1$, 2k-SUM cannot be solved in time $O(n^{k-\varepsilon})$, then, for any $\varepsilon > 0$, Circuit SAT cannot be solved in time $O(q^{O(1)}2^{(1-\varepsilon)n})$.

1.2.3 Improved Space Upper Bound for Subset Sum

Our second main result is an improved space upper bound for Subset Sum. We achieve this by improving and simplifying the algorithm by Nederlof and Wegrzycki.

Theorem 1.4. There exists a Monte Carlo algorithm with constant success probability that solves Subset Sum for instances with n integers of absolute value at most $2^{O(n)}$ in time $O^*(2^{0.5n})$ and space $O^*(2^{0.246n})$.

The algorithm by Nederlof and Węgrzycki proceeds roughly as follows. Let $I \subseteq \mathbb{Z}$, |I| = n, be an instance of Subset Sum. Assume that it is a yes-instance and that a solution $S \subseteq I$ has size n/2. Assume further that there exists a subset $M \subseteq I$ of size $\Theta(n)$ such that $|S \cap M| = |M|/2$ and M is a perfect mixer: the weights of all subsets of M are (pairwise) distinct. Partition $I \setminus M$ into two parts L and R of equal size, hence, $I = L \sqcup M \sqcup R$. We will be looking for two disjoint sets $S_1 \subseteq L \sqcup M$ and $S_2 \subseteq M \sqcup R$ such that $|S_1 \cap M| = |S_2 \cap M| = |M|/4$ and $S_1 \sqcup S_2$ is a solution. This problem can be solved by a reduction to the Weighted Orthogonal Vectors (WOV) problem that can be viewed as a hybrid of the Orthogonal Vectors and the 2-SUM problems: given two families \mathcal{L} and \mathcal{R} of weighted sets and a target integer t, find two disjoint sets whose sum of weights is equal to t. A naive such reduction would proceed similarly to the algorithm by Horowitz and Sahni [6]: let

$$\mathcal{L} = \{ (S_1 \cap M, w(S_1)) \colon S_1 \subseteq L \sqcup M, |S_1 \cap M| = |M/4| \}, \\ \mathcal{R} = \{ (S_2 \cap M, w(S_2)) \colon S_2 \subseteq M \sqcup R, |S_2 \cap M| = |M/4| \}.$$

Then, it remains to solve WOV for \mathcal{L} , \mathcal{R} and t. The bad news is that $|L \sqcup M| > n/2$, hence enumerating all $S_1 \subseteq L \sqcup M$ leads to a running time worse than $O^*(2^{n/2})$. The good news is that the solution S is represented by exponentially many pairs (S_1, S_2) : there are many ways to distribute $S \cap M$ between S_1 and S_2 . This allows one to use the representation technique: take a random prime p of the order $2^{|M|/2}$ and a random remainder $r \in \mathbb{Z}_p$; filter \mathcal{L} and \mathcal{R} to leave only subsets S_1 and S_2 such that $w(S_1) \equiv_p r$ and $w(S_2) \equiv_p t - r$. This way, one reduces the space complexity since \mathcal{L} and \mathcal{R} become about p times smaller

In this work, we further introduce the following idea to reduce the space complexity of the algorithm and to simplify it at the same time. We use another prime q to further filter \mathcal{L} and \mathcal{R} : this allows us to reduce the original instance to a larger number of instances that in turn have smaller size. The idea is best illustrated by the following toy example. Consider an instance I of size n of Subset Sum. As in the

algorithm by Horowitz and Sahni [6], partition I arbitrarily into two parts L and R of size n/2. Assume now that we are given a magic integer $q = \Theta(2^{n/4})$ that hashes all subsets of L and R almost perfectly: for each $r \in \mathbb{Z}_q$,

$$|\{A \subseteq L : w(A) \equiv_q r\}|, |\{B \subseteq R : w(B) \equiv_q t - r\}| = O(2^{n/4}).$$

This allows us to solve I in time $O^*(2^{n/2})$ and space $O^*(2^{n/4})$ without using a reduction to 4-SUM and priority queues machinery of Schroeppel and Shamir's algorithm: for each r, construct sequences of all subsets of L and R having weights r and t-r modulo q, respectively, and solve 2-SUM for the two resulting sequences of length $O(2^{n/4})$. Thus, instead of reducing I to a single instance of 2-SUM of size $2^{n/2}$, we reduce it to $2^{n/4}$ instances of size $2^{n/4}$. We show that even though in general a randomly chosen integer q does not give a uniform distribution of remainders, one can still use this idea to reduce the space.

On a technical side, we choose a mixer M more carefully. This allows us to cover various corner cases, when the size of the solution S is not n/2 and when the mixer M is far from being perfect, in the same manner, thus simplifying the algorithm by Nederlof and Węgrzycki.

2 General Setting

2.1 Modular Arithmetic

We write $a \equiv_m b$ to denote that integers a and b have the same remainder modulo m. By \mathbb{Z}_m we denote the ring of remainders modulo m. We make use of the following well known fact.

Theorem 2.1 (Chinese Remainder Theorem). For any coprime positive integers p and q and any $a \in \mathbb{Z}_p$ and $b \in \mathbb{Z}_q$, there exists a unique $x \in \mathbb{Z}_{pq}$ such that $x \equiv_p a$ and $x \equiv_q b$. Furthermore, it can be computed (using the extended Euclidean algorithm) in time $O(\log^2 p + \log^2 q)$.

2.2 Prime Numbers

Given an integer t, one can generate a uniform random prime from [t, 2t] in time $O(\log^{O(1)} t)$: to do this, one selects a random integer from [t, 2t] and checks whether it is prime [9]; the expected number of probes is $O(\log t)$ due to the law of distribution of prime numbers. We also use the following estimate: any positive integer k has at most $\log_2 k$ distinct prime divisors; hence the probability that a random prime from [t, 2t] divides k is at most $O(\frac{\log k \log t}{t})$.

2.3 Probability Amplification

We use the following standard success probability amplification trick: if a "good" event A occurs with probability at least p, then for A to occur at least once with constant probability, 1/p independent repetitions are sufficient: $(1-p)^{1/p} \leq (e^{-p})^{1/p} = e^{-1}$.

2.4 Growth Rate and Entropy

Much like $O(\cdot)$ hides constant factors, $O^*(\cdot)$ and $\widetilde{O}(\cdot)$ hide factors that grow polynomially in the input size and polylogarithmically, respectively: for example, $n^2 \cdot 1.5^n = O^*(1.5^n)$ and $\log^3 n \cdot n^2 = \widetilde{O}(n^2)$. $\Omega^*(\cdot)$, $\Theta^*(\cdot)$, $\widetilde{\Omega}(\cdot)$, and $\widetilde{\Theta}(\cdot)$ are defined similarly.

We use the following estimates for binomial coefficients: for any constant $0 \le \alpha \le 1$,

$$\Omega(n^{-1/2})2^{h(\alpha)n} \le \binom{n}{\alpha n} \le 2^{h(\alpha)n} \text{ and hence } \binom{n}{\alpha n} = \Theta^* \left(2^{h(\alpha)n}\right), \tag{1}$$

where $h(x) = -x \log_2 x - (1-x) \log_2 (1-x)$ is the binary entropy function. The function h is concave: for any $0 \le \beta \le 1$ and any $0 \le x, y \le 1$, $\beta h(x) + (1-\beta)h(y) \le h(\beta x + (1-\beta)y)$.

2.5 Sets and Sums

For a positive integer n, $[n] = \{1, 2, ..., n\}$. For disjoint sets A and B, by $A \sqcup B$ we denote their union. Let $S \subseteq \mathbb{Z}$ be a finite set of integers. By w(S) we denote the weight of S: $w(S) = \sum_{a \in S} a$. By 2^S we denote the power set of S, i.e., the set of its subsets: $2^S = \{A : A \subseteq S\}$. By $\binom{S}{k}$ we denote the set of all subsets of S of size k: $\binom{S}{k} = \{A \subseteq S : |A| = k\}$. For a family of sets $\mathcal{F} \subseteq 2^S$, by $w(\mathcal{F})$ we denote the set of their weights: $w(\mathcal{F}) = \{w(A) : A \in \mathcal{F}\}$.

Given S, one can compute $|w(2^S)|$ in time and space $O^*(2^{|S|})$ by iterating over all of its subsets. If $|w(2^S)| = 2^{(1-\varepsilon)|S|}$, we call S an ε -mixer.

Lemma 2.1. Let $S \subseteq A$ and |A| = n. For m = O(1) uniformly randomly chosen disjoint non-empty sets $A_1, \ldots, A_m \subseteq A$,

$$\Pr\left[\frac{|A_i \cap S|}{|A_i|} \approx \frac{|S|}{|A|}, \text{ for all } i \in [m]\right] = \Omega^*(1),$$

where $\frac{|A_i \cap S|}{|A_i|} \approx \frac{|S|}{|A|}$ means that $\left| |A_i \cap S| - |A_i| \frac{|S|}{|A|} \right| = O(1)$.

Proof. We will show that

$$\Pr\left[|A_i \cap S| = \left\lfloor |A_i| \frac{|S|}{|A|} \right\rfloor \ \forall i \in [m-1] \land |A_m \cap S| = \left| |A_m| \frac{|S| - \sum_{i < m} \left\lfloor A_i \frac{|S|}{|A|} \right\rfloor}{|A| - \sum_{i < m} |A_i|} \right| \right] = \Omega^*(1) \ .$$

After that it is easy to see that $\left| |A_i \cap S| - |A_i| \frac{|S|}{|A|} \right| = O(1), \ \forall i \in [m].$

Hence, we can think that all numbers $|A_i| \frac{|S|}{|A|}$ are integers.

Since m = O(1), it suffices to estimate the probability for m = 1. Let $B \subseteq A$ be a random set and

$$\gamma = \frac{\left\lfloor |B| \cdot \frac{|S|}{|A|} \right\rfloor}{|B|}.$$

Then

$$\Pr\left[|B \cap S| = \left\lfloor |B| \frac{|S|}{|A|} \right\rfloor \right] = \frac{\binom{|S|}{\gamma|B|} \binom{|A \setminus S|}{(1-\gamma)|B|}}{\binom{|A|}{|B|}} \qquad \text{(by (1))}$$
$$= \Omega^* \left(2^{\gamma|A|h(|B|/|A|) + (1-\gamma)|A|h(|B|/|A|) - |A|h(|B|/|A|)} \right) = \Omega^*(1).$$

2.6 Weighted Orthogonal Vectors

Definition 2.1. [Weighted Orthogonal Vectors, WOV] Given families of N weighted sets $\mathcal{A}, \mathcal{B} \subseteq 2^{[d]} \times \mathbb{N}$, and a target integer t, the problem is to find $(A, w_A) \in \mathcal{A}$ and $(B, w_B) \in \mathcal{B}$ such that $A \cap B = \emptyset$ and $w_A + w_B = t$.

Lemma 2.2. For any $\sigma \in [0, \frac{1}{2}]$, there is a Monte-Carlo algorithm that, given $\mathcal{A} \subseteq \binom{[d]}{\sigma d} \times \mathbb{N}$, $\mathcal{B} \subseteq \binom{[d]}{(1/2-\sigma)d} \times \mathbb{N}$ and a target integer t, solves $WOV(\mathcal{A}, \mathcal{B})$ in time $\widetilde{O}\left((|\mathcal{A}| + |\mathcal{B}|)2^{d(1-h(1/4))}\right)$ and space $\widetilde{O}\left(|\mathcal{A}| + |\mathcal{B}| + 2^d\right)$.

To prove the lemma, we utilize the algorithm from [12], but with a different parameter range. See proofs in section 4.7.

3 New Arthur-Merlin Algorithm

Theorem 1.2. 4-SUM \in AM $\left[\log_2 n, \widetilde{O}(n)\right]$.

Proof. Given arrays A_1, A_2, A_3, A_4 each consisting of n 4 log n-bit integers, our goal is to find four indices $i_1, i_2, i_3, i_4 \in [n]$ such that $A_1[i_1] + A_2[i_2] + A_3[i_3] + A_4[i_4] = 0$. Without loss of generality, assume that A_i 's are sorted and do not contain duplicates: for every $i \in [4], A_i[1] < A_i[2] < \cdots < A_i[n]$. Assume that $i_1^*, i_2^*, i_3^*, i_4^*$ is a solution and let $s = A_1[i_1^*] + A_2[i_2^*]$.

Let \mathcal{P} be a randomized algorithm that, given an integer n, returns a uniform random prime from [2..n]. Since one can check whether a given integer k is prime or not in time $O(\log^{O(1)} k)$, the expected running time of \mathcal{P} is $O(\log^{O(1)} n)$: one picks a random $k \in [2..n]$ and checks whether it is prime; if it is not, one repeats; the expected number of probes is $O(\log n)$.

We are ready to describe the protocol.

- **Stage 0: Shared randomness.** Using the shared randomness, Arthur and Merlin generate a random prime $p = \mathcal{P}(n)$.
- Stage 1: Preparing a proof. Merlin sends an integer $0 \le r < p$ to Arthur.
- Stage 2: Verifying the proof. Arthur takes all elements of A_1 and A_2 modulo p and sorts both arrays. Using the two pointers method (or binary search), he enumerates all pairs $(i_1, i_2) \in [n]^2$ such that $(A_1[i_1] + A_2[i_2]) \equiv r \mod p$. For each such pair (i_1, i_2) , he adds $A_1[i_1] + A_2[i_2]$ to a new array A_{12} . If the size of A_{12} becomes larger than $O(n \log^3 n)$, Arthur rejects immediately. Then, he does the same for A_3 and A_4 , but for the remainder -r (rather than r) and creates an array A_{34} . Finally, he solves 2-SUM for A_{12} and A_{34} (the two arrays may have different lengths, but this can be fixed easily by padding one of them with dummy elements).

Now, we analyze the protocol. It is clear that the proof size is at most $\log_2 n$ and that the running time of the verification stage is $\widetilde{O}(n)$. Also, if there is no solution for the original instance, Arthur rejects with probability 1, as he only accepts if the solution is found. Below, we show that if Merlin sends $r=s \mod p$ (recall that $s=A_1[i_1^*]+A_2[i_2^*]$), Arthur accepts with good enough probability. Consider the set $S=\{(i_1,i_2)\in [n]^2\colon A_1[i_1]+A_2[i_2]\equiv s \mod p\}$ (hence, $|A_{12}|=|S|$). It contains

Consider the set $S = \{(i_1, i_2) \in [n]^2 : A_1[i_1] + A_2[i_2] \equiv s \mod p\}$ (hence, $|A_{12}| = |S|$). It contains at most n true positives, that is, pairs (i_1, i_2) such that $A_1[i_1] + A_2[i_2] = s$: as the arrays do not contain duplicates, for every i_1 there is at most one matching i_2 . All the other pairs (i_1, i_2) in S are false positives: $A_1[i_1] + A_2[i_2] - s \neq 0$, but $A_1[i_1] + A_2[i_2] - s \equiv 0 \mod p$. Recall that $|A_1[i_1] + A_2[i_2] - s| = O(n^4)$, hence the number of prime divisors of $A_1[i_1] + A_2[i_2] - s$ is $O(\log n)$, Section 2.2. The probability that a random prime $2 \leq p \leq n$ divides $(A_1[i_1] + A_2[i_2] - s)$ is $O(\frac{\log^2 n}{n})$. Thus, the expected number of false positives is at most $n^2 \cdot O\left(\frac{\log^2 n}{n}\right) = O(n\log^2 n)$. By Markov's inequality, the probability that the number of false positives is larger than $O(n\log^3 n)$ (and hence that $|S| = O(n\log^3 n)$) is at most $1/\log n$.

Thus, with probability at least $1 - 2/\log n$, both A_{12} and A_{34} have length at most $O(n \log^3 n)$ and by solving 2-SUM for them, Arthur finds a solution for the original yes-instance.

Theorem 1.3. If, for any $\varepsilon > 0$, 4-SUM cannot be solved in time $O(n^{2-\varepsilon})$, then, for any $\varepsilon > 0$, Circuit SAT cannot be solved in time $O(q2^{(1-\varepsilon)n})$.

Proof. Assume that, for some $\varepsilon > 0$, there exists an algorithm \mathcal{A} that checks whether a given Boolean circuit with g binary gates and n inputs is satisfiable in time $O(g2^{(1-\varepsilon)n})$. Using \mathcal{A} , we will solve 4-SUM in time $O(n^{2-\varepsilon})$.

Consider the verification algorithm from Theorem 1.2 as a Turing machine M: M has three read-only input tapes (I_R, I_P, I) : the tape I_R contains random bits, the tape I_P contains a proof, and the tape I contains four input arrays. For any four arrays (A_1, A_2, A_3, A_4) of size n, if it is a yes-instance, there exists a proof of size at most $\log_2 n$ such that M accepts with probability at least 2/3, whereas for a no-instance, M rejects every proof with probability 1. The machine M compares $2 \log n$ -bit integers and moves two pointers through two arrays. Hence, the running time of M is O(n).

Given (A_1, A_2, A_3, A_4) , fix the contents of the tape I and fix the random bits of the tape I_R . This turns M into a machine M' with $\log_2 n$ input bits and running time $\widetilde{O}(n)$.

As proved by [13], a Turing machine recognizing a language $L \subseteq \{0,1\}^*$ with running time t(n) can be converted to an infinite series $\{C_n\}_{n=1}^{\infty}$ of circuits of size $O(t(n)\log t(n))$: for every n, C_n has n inputs and $O(t(n)\log t(n))$ gates and computes a function $f \colon \{0,1\}^n \to \{0,1\}$ such that $f^{-1}(1) = L \cap \{0,1\}^n$. The circuit C_n can be produced in time $O(t(n)\log t(n))$.

Applying this to the machine M', we get a circuit C with $\widetilde{O}(n)$ gates and $\log_2 n$ input bits. If A_1, A_2, A_3, A_4 is a yes-instance of 4-SUM, the circuit C is satisfiable with probability at least 2/3; otherwise it is unsatisfiable. Using the algorithm \mathcal{A} , we can check the satisfiability of C in time $\widetilde{O}(n2^{(1-\varepsilon)\log_2 n}) = \widetilde{O}(n^{2-\varepsilon})$.

Theorem 1.1. Subset $Sum \in AM\left[n/4, O^*(2^{n/4})\right]$.

Proof. We use the standard reduction from Subset SUM to 4-SUM. Given a sequence A of n integers, partition them into four parts of size n/4 and create sequences A_1, A_2, A_3, A_4 of size $2^{n/4}$ containing sums of all subsets of the corresponding parts. Use the protocol from Theorem 1.2 for the resulting instance of 4-SUM.

Corollary 1.1. For every even integer k, 2k-SUM \in AM $\left\lceil \frac{k}{2} \log_2 n, \widetilde{O}(n^{k/2}) \right\rceil$.

Proof. Reduce 2k-SUM to 4-SUM: given sequences A_1, \ldots, A_{2k} of length n, partition them into four parts each containing k/2 sequences of length n. Let B_1, B_2, B_3, B_4 be sequences of length $n^{k/2}$ containing all sums of k/2 elements of the corresponding part (where we take exactly one element from each sequence). Use the protocol from Theorem 1.2 for the resulting instance of 4-SUM.

Corollary 1.2. Assume that 2k-SUM \in AM $\left[k\log_2 n, n^{o(k)}\right]$. Then, if for any $\varepsilon > 0$ and any $k \geq 1$, 2k-SUM cannot be solved in time $O(n^{k-\varepsilon})$, then, for any $\varepsilon > 0$, Circuit SAT cannot be solved in time $O(q^{O(1)}2^{(1-\varepsilon)n})$.

Proof. Assume that Circuit SAT can be solved in time $O(g^c 2^{(1-\varepsilon)n})$ for constants $c, \varepsilon > 0$.

Similarly to the proof of Theorem 1.3, given an instance of 2k-SUM, create a circuit C with $k \log_2 n$ inputs and of size $n^{o(k)}$ that solves this instance with one-sided error probability at most 1/3.

Using the above algorithm, one can check the satisfiability of C in time

$$O(n^{c \cdot o(k)} \cdot 2^{(1-\varepsilon)k \log_2 n}) = O(n^{o(k)} n^{(1-\varepsilon)k}).$$

For large enough k, this allows to solve 2k-SUM in time $O(n^{(1-\varepsilon/2)k})$.

4 Improved Space Upper Bound for Subset Sum

4.1 Algorithm

Let (I,t) be an instance of Subset Sum (as usual, n=|I|) and assume that it is a yes-instance: we will present a randomized algorithm that is correct on no-instances with probability 1, so, as usual, the main challenge is to obtain $\Omega^*(1)$ success probability for yes-instances. Let $S\subseteq I$ be a solution: w(S)=t. It is unknown to us, but we may assume that we know its size: there are just n possibilities for |S|, so with a polynomial overhead we can enumerate all of them. Thus, assume that $|S|=\alpha n$ where $0<\alpha\le 1$ is known to us.

For a parameter $\beta = \beta(\alpha) < 0.15$ to be specified later, select randomly pairwise disjoint sets $M_L, M, M_R \in \binom{I}{\beta n}$. In time and space $O^*(2^{0.15n})$, find $\varepsilon_L, \varepsilon, \varepsilon_R$ such that M_L is an ε_L -mixer, M is an ε -mixer, and M_R is ε_R -mixer. Without loss of generality, we assume that $\varepsilon \leq \varepsilon_L, \varepsilon_R$. Consider the probability that S touches exactly half of $M_L \sqcup M \sqcup M_R$:

$$\Pr\left[\left|\left(M_L \cup M \cup M_R\right) \cap S\right| = 3\beta n/2\right] = \frac{\binom{\alpha n}{3\beta n/2}\binom{(1-\alpha)n}{3\beta n/2}}{\binom{n}{3\beta n}} = \Theta^*\left(2^{-\lambda n}\right)$$

$$\frac{\binom{\alpha n}{3\beta n/2}\binom{(1-\alpha)n}{3\beta n/2}}{\binom{n}{3\beta n}} = 2^{-\lambda n} \iff \frac{1}{n}\log\frac{\binom{n}{3\beta n}}{\binom{\alpha n}{3\beta n/2}\binom{(1-\alpha)n}{3\beta n/2}} = \lambda \iff$$

$$\iff \lambda = h(3\beta) - \alpha h\left(\frac{3\beta}{2\alpha}\right) - (1-\alpha)h\left(\frac{3\beta}{2(1-\alpha)}\right), \qquad \text{(by (1))}$$

hence,

$$\lambda = h(3\beta) - \alpha h\left(\frac{3\beta}{2\alpha}\right) - (1-\alpha)h\left(\frac{3\beta}{2(1-\alpha)}\right). \tag{2}$$

By repeating this process $2^{\lambda n}$ times, we ensure that the event $|(M_L \cup M \cup M_R) \cap S| = 3\beta n/2$ happens with probability $\Omega^*(1)$ (recall Section 2.3). Further, we may assume that

$$|M_L \cap S| = |M \cap S| = |M_R \cap S| = \beta n/2,$$

as by Lemma 2.1, $|M_L \cap S| = |M \cap S| = |M_R \cap S| = \beta n/2 + O(1)$ with probability $\Omega^*(1)$ conditioned on the event that S touches $3\beta n/2$ elements of $M_L \sqcup M \sqcup M_R$. An exact equality can be assumed since we can search for an exact value in the neighbourhood of $\beta n/2$ and the constant difference will not affect the memory and time complexity. In the following applications of Lemma 2.1, we omit O(1) factors for similar considerations.

We argue that either $S \cap M$ or $M \setminus S$ is an $(\leq \varepsilon)$ -mixer. Indeed, M is an ε -mixer and hence $|w(2^M)| = 2^{(1-\varepsilon)|M|}$. Now, $|S \cap M| = |M \setminus S| = |M|/2$ and $|w(2^{S \cap M})| \cdot |w(2^{M \setminus S})| \geq |w(2^M)|$. Hence,

$$\max\left\{|w(2^{S\cap M})|,|w(2^{M\setminus S})|\right\} \ge 2^{(1-\varepsilon)|M|/2},$$

implying that either $S \cap M$ or $M \setminus S$ is indeed a $(\leq \varepsilon)$ -mixer. In the following, we assume that it is $S \cap M$ that is $(\leq \varepsilon)$ -mixer. To assume this without loss of generality, we run the final algorithm twice — for (I,t) and (I,w(I)-t).

For any (finite and non-empty) set A, there exists $1 \le k \le |A|$, such that $|w({A \choose k})| \ge |w(2^A)|/|A|$ (by the pigeonhole principle). Since $|w({A \choose k})| = |w({A \choose |A|-k})|$, we can even assume that $1 \le k \le |A|/2$. Consider the corresponding k for the set $A = S \cap M$ and let $\mu = k/(2|S \cap M|)$ (hence, $0 \le \mu \le 0.25$). Then,

$$\left| w \left(\begin{pmatrix} S \cap M \\ 2\mu | S \cap M | \end{pmatrix} \right) \right| \ge \frac{\left| w \left(2^{S \cap M} \right) \right|}{\left| S \cap M \right|}. \tag{3}$$

Partition $I \setminus (M_L \sqcup M \sqcup M_R)$ randomly into $L_1 \sqcup L_2 \sqcup L_3 \sqcup L_4 \sqcup R_1 \sqcup R_2 \sqcup R_3 \sqcup R_4$ where the exact size of all eight parts will be specified later. Let $L = L_1 \sqcup L_2 \sqcup L_3 \sqcup L_4$ and $R = R_1 \sqcup R_2 \sqcup R_3 \sqcup R_4$. By Lemma 2.1, with probability $\Omega^*(1)$ the set S covers the same fraction of each of L_i 's and R_i 's. We denote this fraction by γ :

$$\gamma = \frac{|(L \sqcup R) \cap S|}{|L \sqcup R|} = \frac{\alpha - 3\beta/2}{1 - 3\beta}.$$
 (4)

Now, we apply the representation technique. Note that there exist many ways to partition S into $S_L \sqcup S_R$ such that

$$S_L \subseteq M_L \sqcup L \sqcup M \text{ and } S_R \subseteq M \sqcup R \sqcup M_R.$$
 (5)

Indeed, the elements from $S \cap M$ can be distributed arbitrarily between S_L and S_R . Below, we show that for a large random prime number p and a random remainder r modulo p, the probability that there exists a partition $S = S_L \sqcup S_R$ such that $w(S_L) \equiv_p r$ is $\Omega^*(1)$ (informally, at least one representative $S_L \sqcup S_R$ of the original solution S survives, even if we "hit" all such partitions with a large prime).

Lemma 4.1. With constant probability over choices of a prime number p such that $p \in [X/2, X]$, where $X \leq |w(2^{S \cap M})|$, the number of remainders $a \in \mathbb{Z}_p$ such that there exists a subset $M' \subseteq S \cap M$ with $|M'| = 2\mu |S \cap M|$ and $w(M') \equiv_p a$ is close to p:

$$\Pr\left[|\{a\in\mathbb{Z}_p\colon there\ exists\ M'\subseteq S\cap M, |M'|=2\mu|S\cap M|, w(M')\equiv_p a\}|\geq \Omega\left(p/n^2\right)\right]\geq 9/10.$$

Proof. Let $Q = M \cap S$. Then $|Q| = \frac{\beta}{2}n$. Recall that $|w(\binom{Q}{2\mu|Q|})| \ge |w(2^Q)|/|Q|$ and p is a random prime in [X/2, X].

Let $\mathcal{M} = \binom{Q}{2\mu|Q|}$, $c_i = |\{M : M \in \mathcal{M}, w(M) \equiv_p i\}|$, and $d_i = [c_i > 0]$. We need to show that $\Pr[|\{i : d_i = 1\}| \geq \Omega(p/n^2)] \geq 9/10$. The expected number of collisions is

$$\mathbb{E}\left[\sum_{i} c_i^2\right] = \sum_{M_1, M_2 \in \mathcal{M}} \Pr\left[p \text{ divides } w(M_2) - w(M_1)\right] \leq |\mathcal{M}| + O(n^2|\mathcal{M}|^2/X).$$

The last inequality follows from Section 2.2, all numbers are not greater than $2^{O(n)}$. Hence, $\sum_i c_i^2 = O(|\mathcal{M}| + n^2 |\mathcal{M}|^2 / X) = O(n^2 |\mathcal{M}|^2 / p)$ with constant probability. Then, the Cauchy–Schwartz inequality implies that

$$|\mathcal{M}| = \sum_{i} c_i = \sum_{i} c_i d_i \le \sqrt{\sum_{i} d_i^2} \sqrt{\sum_{i} c_i^2} = \sqrt{\sum_{i} d_i} \sqrt{\sum_{i} c_i^2}.$$

Hence,

$$\sum_i d_i \geq \frac{(\sum_i c_i)^2}{\sum_i c_i^2} \geq \frac{|\mathcal{M}|^2}{O(n^2|\mathcal{M}|^2/p)} \geq \Omega(p/n^2).$$

Take a random prime $2^{\frac{1-\varepsilon}{2}\beta n-1} \leq p \leq 2^{\frac{1-\varepsilon}{2}\beta n}$ and a random remainder $r \in \mathbb{Z}_p$. Lemma 4.1 ensures that with probability $\Omega^*(1)$, there exists $M' \subseteq S \cap M$ such that $|M'| = 2\mu |S \cap M| = \mu \beta n$ and $w(M') \equiv_p r$.

To find the partition $S_L \sqcup S_R$, we are going to enumerate all sets $A \subseteq M_L \sqcup L \sqcup M$ with $w(A) \equiv_p r$ and $|A \cap (M \cap S)| = \mu \beta n$ as well as all sets $B \subseteq M \sqcup R \sqcup M_R$ with $w(B) \equiv_p t - r$ and $|B \cap (M \cap S)| = (1/2 - \mu)\beta n$. Then, it suffices to solve WOV for the sets of A's and B's.

To actually reduce to WOV, we first construct set families $\mathcal{M}_L \subseteq 2^{M_L}$ and $\mathcal{M}_R \subseteq 2^{M_R}$ such that $|\mathcal{M}_L| = |w(2^{M_L})|$ and $|\mathcal{M}_R| = |w(2^{M_R})|$. That is, \mathcal{M}_L contains a single subset of M_L for each possible weight. One can construct \mathcal{M}_L (as well as \mathcal{M}_R) simply by going through all subsets of M_L and checking, for each subset, whether its sum is already in \mathcal{M}_L .

Then, construct the following families (see Figure 1).

$$Q_1 = \{ M' \cup L' : M' \in \mathcal{M}_L, L' \subseteq L_1, |L'| = \gamma |L_1| \},$$
(6)

$$Q_2 = \{ L' : L' \subseteq L_2, |L'| = \gamma |L_2| \}, \tag{7}$$

$$Q_3 = \{ L' : L' \subseteq L_3, |L'| = \gamma |L_3| \}, \tag{8}$$

$$Q_4 = \{ L' \cup M' : L' \subseteq L_4, M' \subseteq M, |L'| = \gamma |L_4|, |M'| = \mu \beta n \},$$
(9)

$$Q_1' = \{ M' \cup R' : M' \in \mathcal{M}_R, R' \subseteq R_1, |R'| = \gamma |R_1| \}, \tag{10}$$

$$Q_2' = \{R' : R' \subseteq R_2, |R'| = \gamma |R_2|\},\tag{11}$$

$$Q_3' = \{R' : R' \subseteq R_3, |R'| = \gamma |R_3|\},\tag{12}$$

$$Q_4' = \{ R' \cup M' : R' \subseteq R_4, M' \subseteq M, |R'| = \gamma |R_4|, |M'| = (1/2 - \mu)\beta n \}.$$
(13)




Figure 1: Partition of the instance I (white) and its solution S (gray) into parts.

Let \mathcal{Q}_{max} stand for \mathcal{Q}_i or \mathcal{Q}'_i with the maximum size. For $a, p \in \mathbb{Z}_{>0}$, let

$$\mathcal{X}_a = \left\{ (Q_1, Q_2, Q_3, Q_4) \colon Q_i \in \mathcal{Q}_i, \sum w(Q_i) = a \right\} \text{ and } \mathcal{X}_{a,p} = \bigcup_{a' \equiv_p a} \mathcal{X}_{a'}.$$

Then, it remains to solve WOV for $\mathcal{X}_{r,p}$ and $\mathcal{X}'_{(t-r),p}$. To do this, we use the idea used in the Arthur–Merlin protocol: we reduce to 4-SUM and use a cutoff for the number of found candidates. The following lemmas estimate the size of these set families.

Lemma 4.2. With probability $\Omega^*(1)$ over the choice of r, the size of $\mathcal{X}_{r,p}$ is at most $O^*(|\mathcal{Q}_{max}|^4/p)$ and there exists $M_1 \subseteq M \cap S$ such that $w((S \cap (L \cup M_L)) \cup M_1) \equiv_p r$.

Proof. Let A be an event of existing $M_1 \subseteq M \cap S$ such that $w((S \cap (L \cup M_L)) \cup M_1) \equiv_p r$. Note that

$$\mathbb{E}[\mathcal{X}_{r,p}] \le |\mathcal{Q}_{max}|^4/p \text{ and } \Pr[\mathcal{X}_{r,p} > x] \le \frac{|\mathcal{Q}_{max}|^4}{px}.$$

As $\Pr[A] = \Omega^*(1)$, we can take $x = O^*(\frac{|\mathcal{Q}_{max}|^4}{p})$ in a way that will make $\Pr[\mathcal{X}_{r,p} > x] \leq \frac{\Pr[A]}{2}$. Then

$$\Pr[\mathcal{X}_{r,p} \le x \land A] \ge \Pr[A] - \Pr[\mathcal{X}_{r,p} > x] \ge \frac{\Pr[A]}{2} = \Omega^*(1).$$

To further reduce the size of these families, we choose another random prime number

$$q = \Theta^* \left(\frac{2^{n/2}}{2^{\lambda n} |\mathcal{Q}_{max}|^2} \right),$$

and iterate over all possible remainders $s \in \mathbb{Z}_q$. We aim to find two families of sets as the above ones, but with weights congruent to $w(S_L)$ and $(t-w(S_L))$, respectively, modulo p and q. While iterating on s, at some point we try $s \equiv_q w(S_L)$, so we can assume that we guessed s correctly and construct the families for the fixed r and s. We now focus on the first family only, since we can then find the second one independently using the same algorithm.

For $a \in \mathbb{Z}_{>0}$, let $\mathcal{L}_a = \{(a, Q_4 \cap M) : (Q_1, Q_2, Q_3, Q_4) \in \mathcal{X}_a\}$. For $a_1, a_2, p, q \in \mathbb{Z}_{>0}$, let

$$\mathcal{X}_{a_1,p,a_2,q} = \bigcup_{\substack{a \equiv p a_1 \\ a \equiv a_2}} \mathcal{X}_a \text{ and } \mathcal{L}_{a_1,p,a_2,q} = \bigcup_{\substack{a \equiv p a_1 \\ a \equiv a_2}} \mathcal{L}_a.$$

Our goal is to find a family \mathcal{L} containing $\mathcal{L}_{w(S_L)}$ in order to guarantee that we later find the solution using the WOV algorithm. But we need \mathcal{L} to be small enough. Let $\ell = \frac{|\mathcal{Q}_{max}|^4}{pq} + \binom{|M|}{\max\{\mu, 1/2 - \mu\}|M|}$ represent an upper bound on the sum of the average size of $\mathcal{X}_{a_1,p,a_2,q}$ and the size of $\mathcal{L}_{w(S_L)}$. We want \mathcal{L} to be of size $\widetilde{O}(\ell)$. The following lemma shows that $\mathcal{X}_{w(S_L),p,w(S_L),q}$ contains not too many redundant sets.

Lemma 4.3. With probability $\Omega^*(1)$ over the choice of q, $|\mathcal{X}_{r,p,w(S_L),q} \setminus \mathcal{X}_{w(S_L)}| \leq \ell$.

Proof. We estimate the expected number of elements in $X_{r,p,w(S_L),q} \setminus \mathcal{X}_{w(S_L)}$. To do that, we calculate the sum of $\Pr_q[\sum w(Q_i) \equiv_q w(S_L)]$ over all tuples $(Q_1,Q_2,Q_3,Q_4) \in \mathcal{X}_{r,p} \setminus \mathcal{X}_{w(S_L)}$ such that $\sum w(Q_i) \neq w(S_L)$.

Since $\Pr_q[\sum w(Q_i) \equiv_q w(S_L)] = O^*(\frac{1}{q})$ (recall Section 2.2),

$$\mathbb{E}|\mathcal{X}_{r,p,w(S_L),q} \setminus \mathcal{X}_{w(S_L)}| = |\mathcal{X}_{r,p} \setminus \mathcal{X}_{w(S_L)}| \cdot O^*(1).$$

By Lemma 4.2, it is equal to $\widetilde{O}(|\mathcal{Q}_{max}|^4/(pq)) = \widetilde{O}(\ell)$.

Let us find the first $\leq 2\ell + 1$ elements of $\mathcal{X}_{r,s}$ using the algorithm from Lemma 4.4 which runs in time $\widetilde{O}(|\mathcal{Q}_{max}|^2 + \ell)$ and uses $\widetilde{O}(|\mathcal{Q}_{max}| + \ell)$ space.

Lemma 4.4. There is an algorithm that given Q_1, Q_2, Q_3, Q_4 , (r, p, s, q) and integer m, finds $\min(|\mathcal{X}_{r,p,s,q}|, m)$ elements of $\mathcal{X}_{r,p,s,q}$ using $\widetilde{O}(|\mathcal{Q}_{max}|^2 + m)$ time and $\widetilde{O}(|\mathcal{Q}_{max}| + m)$ space.

Proof. Let $v \in [0, pq-1]$ be an integer such that $v \equiv_p r$ and $v \equiv_q s$. We obtain such v in $\Omega^*(1)$ time by Theorem 2.1. Our goal is to output every tuple (Q_1, Q_2, Q_3, Q_4) such that $Q_i \in \mathcal{Q}_i$ and $\sum w(Q_i) \equiv_{pq} v$, one by one and halt the algorithm when the size of the output becomes m.

We can assume that all weights of sets from Q_i are in [0, pq-1]. Then, we want to output the tuples with total weights equal to v, v+pq, v+2pq or v+3pq. Now we run Schroeppel and Shamir's [14] algorithm four times keeping the total size of the output at most m.

Assume that $s \equiv_q w(S_L)$. If the algorithm yields at most 2ℓ elements then we know the whole $\mathcal{X}_{r,p,w(S_L),q}$, so we can now just construct \mathcal{L} from $\mathcal{X}_{r,p,s,q}$ in $\widetilde{O}(\ell)$ time and space. Otherwise, by Lemma 4.3, we have that $|\mathcal{X}_{w(S_L)}| > \ell$ with probability $\Omega^*(1)$. Since strictly more than half of the outputted tuples have weight exactly $w(S_L)$, we can determine $w(S_L)$ in $\widetilde{O}(\ell)$ time. When we know $w(S_L)$, we use Lemma 4.5 to obtain $\mathcal{L}_{w(S_L)}$ using $\widetilde{O}(|\mathcal{Q}_{max}|^2)$ time and $\widetilde{O}(|\mathcal{Q}_{max}|)$ space.

Lemma 4.5. There is an algorithm that given Q_1, Q_2, Q_3, Q_4 and a, finds \mathcal{L}_a using $\widetilde{O}(|\mathcal{Q}_{max}|^2)$ time and $\widetilde{O}(|\mathcal{Q}_{max}|)$ space.

Proof. Schroeppel and Shamir [14] use priority queues to implement a data structure D_1 for the following task: output all elements from $\{w(Q_1) + w(Q_2) \colon Q_1 \in \mathcal{Q}_1, Q_2 \in \mathcal{Q}_2\}$ in non-decreasing order using $\widetilde{O}(|\mathcal{Q}_1| \cdot |\mathcal{Q}_2|)$ time and $\widetilde{O}(|\mathcal{Q}_1| + |\mathcal{Q}_2|)$ space. Similarly, a data structure D_2 outputs all elements from $\{(w(Q_3) + w(Q_4), Q_4 \cap M) \colon Q_3 \in \mathcal{Q}_3, Q_4 \in \mathcal{Q}_4\}$ in non-increasing order (we compare elements by their first coordinate) using $\widetilde{O}(|\mathcal{Q}_3| \cdot |\mathcal{Q}_4|)$ time and $\widetilde{O}(|\mathcal{Q}_3| + |\mathcal{Q}_4|)$ space. By $\mathrm{inc}(w(\mathcal{Q}_1, \mathcal{Q}_2))$ and $\mathrm{dec}(w(\mathcal{Q}_3, \mathcal{Q}_4))$ we denote initilization methods for these data structures, by $\mathrm{pop}()$ we denote their method that gives the next element.

Algorithm 1 solves the problem using data structures D_1 and D_2 . It works in time $\widetilde{O}(|\mathcal{Q}_1| \cdot |\mathcal{Q}_2| + |\mathcal{Q}_3| \cdot |\mathcal{Q}_4|)$, uses $\widetilde{O}(|\mathcal{Q}_1| + |\mathcal{Q}_2| + |\mathcal{Q}_3| + |\mathcal{Q}_4|)$ space, and outputs $|\mathcal{L}_a|$ elements. Since the number of outputted subsets is at most $|\mathcal{Q}_4|$, even if we store the output, we still need only $\widetilde{O}(|\mathcal{Q}_{max}|^2)$ time and $\widetilde{O}(|\mathcal{Q}_{max}|)$ space.

```
Input : Q_1, Q_2, Q_3, Q_4, a

Output: \mathcal{L}_a = \{(a, Q_4 \cap M) : (Q_1, Q_2, Q_3, Q_4) \in \mathcal{X}_a\}

1 Initialize D_1 = \mathrm{inc}(w(Q_1, Q_2))

2 Initialize D_2 = \mathrm{dec}(w(Q_3, Q_4))

3 while (d_2, A) = D_2.\mathrm{pop}() do

4 | while d_1 = D_1.\mathrm{pop}() and d_1 + d_2 < a do

5 | skip d_1

6 | if d_1 + d_2 = a and not used[A] then

7 | output((a, A))

8 | used[A] = True
```

Algorithm 1: Pseudocode for Lemma 4.5.

Note that the size of \mathcal{L}_a is at most $\binom{|M|}{u|M|}$. Summarizing, we get the following.

Lemma 4.6. There is an algorithm that given Q_1, Q_2, Q_3, Q_4 and (r, p, s, q), uses $\widetilde{O}\left(|Q_{max}|^2 + \frac{|Q_{max}|^4}{pq} + \binom{|M|}{\mu|M|}\right)$ time and $\widetilde{O}\left(|Q_{max}| + \frac{|Q_{max}|^4}{pq} + \binom{|M|}{\mu|M|}\right)$ space. It returns a set $\mathcal{L} \subseteq \mathbb{Z} \times Q_4$ of size $\widetilde{O}\left(\frac{|Q_{max}|^4}{pq} + \binom{|M|}{\mu|M|}\right)$ that satisfies the following. If there exist a solution $S \subseteq I$ with w(S) = t, and a partition $S = S_L \sqcup S_R$, such that:

- $|S| = \alpha n$;
- $|M_L \cap S| = |M \cap S| = |M_R \cap S| = \beta n/2;$
- $S \cap M$ is a $(\leq \varepsilon)$ -mixer;
- $|S \cap L_i| = \gamma |L_i|$ for each $i \in [4]$, the same for R_i ;
- $S_L \subseteq M_L \sqcup L \sqcup M$, $S_R \subseteq M \sqcup R \sqcup M_R$;
- $|S_L \cap M| = \mu \beta n, |S_R \cap M| = (1/2 \mu) \beta n;$
- $w(S_L) \equiv_p r$, $w(S_L) \equiv_q s$, $w(S_R) \equiv_p t r$, $w(S_R) \equiv_q t s$;
- $|\mathcal{X}_{r,p,s,q} \setminus \mathcal{X}_{w(S_L)}| \leq \ell$,

then $\mathcal{L}_{w(S_L)} \subseteq \mathcal{L}$. Otherwise, \mathcal{L} may be any set.

Similarly, using Lemma 4.6, we find subset \mathcal{R} for the right side. We call WOV(\mathcal{L}, \mathcal{R}) and return True if it finds a solution. The WOV algorithm works in time $\widetilde{O}((|\mathcal{L}| + |\mathcal{R}|)2^{\beta n(1-h(1/4))})$ and space $\widetilde{O}(|\mathcal{L}| + |\mathcal{R}| + 2^{\beta n})$ by Lemma 2.2.

It is worth noting that for most choices of s our assumption that $s \equiv_q w(S_L)$ does not hold. Also, with probability $1 - \Omega^*(1)$, $|\mathcal{X}_{r,p,w(S_L),q} \setminus \mathcal{X}_{w(S_L)}|$ may happen to be greater than ℓ . If any of these cases occurs, we may find a set \mathcal{L} that was not intended (i.e. it does not contain $\mathcal{L}_{w(S_L)}$), or we may even not find anything at all (if after applying the algorithm from Lemma 4.4 we obtain $2\ell + 1$ elements, each of which occurs less than $\ell + 1$ times). In the second scenario, we simply continue with the next choice of s. Otherwise, we cannot determine whether we have found the correct \mathcal{L} or not. However, if an incorrect \mathcal{L} is found, the outcome of our algorithm remains unaffected. If $\mathrm{OV}(\mathcal{L}, \mathcal{R})$ finds a solution, it is a solution to the original problem; if not, we simply proceed to the next iteration of the loop in line 8.

Now we are ready to present the final Algorithm 2. We will choose the exact values for its parameters later.

```
Input : (I, t).
    Output: True, if S \subseteq I with w(S) = t exists, otherwise False.
 1 repeat 2^{\lambda n} times
        Select random disjoint subsets M_L, M, M_R \subseteq I of size \beta n
3
        foreach \mu such that \mu\beta n \in [0, \beta n/2] do
             Randomly partition I \setminus (M_L \sqcup M \sqcup M_R) into L, R with size as in (15)–(16)
 4
             Pick a prime p = \Theta(2^{(1-\varepsilon)\beta n/2}) and r \in \mathbb{Z}_p at random
 5
             Construct Q_1, Q_2, Q_3, Q_4 and Q'_1, Q'_2, Q'_3, Q'_4 as defined in (6)–(13)
 6
             Pick a random prime q = \Theta^* \left( 2^{n/2} / (2^{\lambda n} |\mathcal{Q}_{max}|^2) \right)
 7
             foreach s \in \mathbb{Z}_q do
 8
                 Construct \mathcal{L} and \mathcal{R} (as in Lemma 4.6)
 9
                  if WOV(\mathcal{L}, \mathcal{R}) (as in Lemma 2.2) then
10
                      return True
11
12 return False
```

Algorithm 2: The main algorithm.

4.2 Correctness

Lemma 4.7. The success probability of the algorithm is $\Omega^*(1)$.

Proof. To detect that I is indeed a yes-instance, an algorithm iteration relies on a number of events. The first one is $|(M_L \cup M \cup M_R) \cap S| = 3\beta n/2$, that occurs with probability $\Theta^* \left(2^{-\lambda n}\right)$. All the others occur with probability $\Omega^*(1)$, conditioned on the event that all the previous events occur. This implies that all of them occur with probability $\Omega^* \left(2^{-\lambda n}\right)$, and after repeating it $2^{\lambda n}$ times, the success probability of the resulting algorithm is $\Omega^*(1)$. Below, we recall all the considered events and provide links to lemmas proving their conditional probabilities.

```
1. |(M_L \cup M \cup M_R) \cap S| = 3\beta n/2, see Eq. (2).
```

- 2. $|M_L \cap S| = |M \cap S| = |M_R \cap S| = \frac{\beta}{2}n$, see Lemma 2.1.
- 3. $|S \cap L_i| = \gamma |L_i|$ and $|S \cap R_i| = \gamma |R_i|$, see Lemma 2.1.
- 4. p is a prime number, see Section 2.2.
- 5. $|\{a \in \mathbb{Z}_p : \exists M' \subseteq S \cap M, |M'| = 2\mu |S \cap M|, w(M') \equiv_p a\}| \geq \Omega(p/n^2)$, see Lemma 4.1.
- 6. $|\mathcal{X}_{r,p}| = O^*(|\mathcal{Q}_{max}|^4/p)$ and there exists a partition $M_1 \sqcup M_2 = S \cap M$ such that $w(S \cap (M_L \cup L \cup M_1)) \equiv_p r$ and $w(S \cap (M_2 \cup R \cup M_R)) \equiv_p t r$, see Lemma 4.2.
- 7. q is a prime number, see Section 2.2.

8. $|\mathcal{X}_{r,p,w(S_L),q} \setminus \mathcal{X}_{w(S_L)}| \leq \ell$, see Lemma 4.3.

In case any of those events does not happen, the algorithm completes the iteration without affecting the final result.

4.3 Setting the Parameters

Recall that $\alpha = \frac{|S|}{n}$ and we iterate over all possible α . We define $\beta = \beta(\alpha)$ as follows:

$$\beta(\alpha) = \begin{cases} 0.13, & \alpha \in [0.45, 0.55] \\ 0, & \text{otherwise.} \end{cases}$$
 (14)

Let us consider an iteration of the loop on line 3. To this point, we know the values of ε , ε_L , ε_R and $|M_R\rangle| = (1-3\beta)n$. Let us consider a partition $I\setminus (M_L\sqcup M\sqcup M_R) = L_1\sqcup L_2\sqcup L_3\sqcup L_4\sqcup R_1\sqcup R_2\sqcup R_3\sqcup R_4$ and estimate the size of \mathcal{Q}_i and \mathcal{Q}_i' . Recall that $\gamma = \frac{|(L\cup R)\cap S|}{|L\cup R|} = \frac{\alpha-\frac{3}{2}\beta}{1-3\beta}$. Observe that we use ε instead of ε_L and ε_R since $\varepsilon \leq \varepsilon_L, \varepsilon_R$.

$$\begin{aligned} |\mathcal{Q}_{1}| &\leq \binom{|L_{1}|}{\gamma |L_{1}|} \cdot 2^{(1-\varepsilon)\beta n} & |\mathcal{Q}'_{1}| \leq \binom{|R_{1}|}{\gamma |R_{1}|} \cdot 2^{(1-\varepsilon)\beta n} \\ |\mathcal{Q}_{2}| &= \binom{|L_{2}|}{\gamma |L_{2}|} & |\mathcal{Q}'_{2}| = \binom{|R_{2}|}{\gamma |R_{2}|} \\ |\mathcal{Q}_{3}| &= \binom{|L_{3}|}{\gamma |L_{3}|} & |\mathcal{Q}'_{3}| = \binom{|R_{3}|}{\gamma |R_{3}|} \\ |\mathcal{Q}_{4}| &= \binom{|L_{4}|}{\gamma |L_{4}|} \cdot \binom{|M|}{\mu \beta n} & |\mathcal{Q}'_{4}| = \binom{|R_{4}|}{\gamma |R_{4}|} \cdot \binom{|M|}{\left(\frac{1}{2} - \mu\right)\beta n}. \end{aligned}$$

We construct \mathcal{L} and \mathcal{R} in time depending on $|\mathcal{Q}_{max}|^2 + |\mathcal{Q}'_{max}|^2$. Note that this function is minimized when all $|Q_i|$ and $|Q'_i|$ are roughly the same. We can achieve that by a careful choice of $|L_i|$ and $|R_i|$.

We define the sizes of |L| and |R| as shown below and require that $|L_1| + |L_2| + |L_3| + |L_4| = |L|$ and $|R_1| + |R_2| + |R_3| + |R_4| = |R|.$

$$|L| = \frac{1 - 3\beta - \chi\beta}{2}n,\tag{15}$$

$$|R| = \frac{1 - 3\beta + \chi \beta}{2} n,\tag{16}$$

where $\chi = \frac{h(\mu) - h\left(\frac{1}{2} - \mu\right)}{h(\gamma)}$ is a balancing parameter needed to ensure that $|\mathcal{Q}_{max}| \approx |\mathcal{Q}'_{max}|$. Now we will show that all the parameters are chosen correctly and will define sizes of $L_1, L_2, L_3, L_4, R_1, R_2, R_3, R_4$, so we never, for example, choose the size for L_i to be negative, or repeat the algorithm a number of times smaller than one.

Lemma 4.8. λ defined in Equation (2) is a non-negative number.

Proof. Entropy is a concave function, hence

$$\alpha h\left(\frac{3\beta}{2\alpha}\right) + (1-\alpha)h\left(\frac{3\beta}{2(1-\alpha)}\right) \le h\left(\frac{3\beta}{2} + \frac{3\beta}{2}\right) = h(3\beta).$$

This implies that $\lambda \geq 0$.

Lemma 4.9. For our choice of β , for every $\alpha \in [0,1]$, $|L| \geq 0$, where

$$|L| = n \frac{h(\gamma) - 3\beta h(\gamma) - \beta h(\mu) + \beta h(1/2 - \mu)}{2h(\gamma)}.$$
 (Eq. (15))

Proof.

$$|L| \ge 0 \iff h(\gamma) - 3\beta h(\gamma) + \beta(-h(\mu) + h(1/2 - \mu)) \ge 0$$
$$h(\gamma) - 3\beta h(\gamma) + \beta(\underbrace{-h(\mu) + h(1/2 - \mu)}_{>-1}) \ge h(\gamma) - 3\beta h(\gamma) - \beta = h(\gamma)(1 - 3\beta) - \beta$$

Consider two cases:

1. $\alpha \in [0.45, 0.55]$, then $\beta = 0.13, h(\gamma) \ge h(0.45) \ge 0.99$ and

$$h(\gamma)(1-3\beta) - \beta \ge 0.99(1-3\cdot 0.13) - 0.13 \ge 0.47$$

2. $\alpha \in [0, 0.45) \cup (0.55, 1]$, then $\beta = 0$ and $|L| = \frac{1}{2}n$.

Lemma 4.10. For our choice of β , for every $\alpha \in [0,1]$ we can choose $|L_1|$, $|L_2|$, $|L_3|$, $|L_4|$, $|R_1|$, $|R_2|$, $|R_3|$, $|R_4| \ge 0$ such that $|L_1| + |L_2| + |L_3| + |L_4| = |L|$, $|R_1| + |R_2| + |R_3| + |R_4| = |R|$, and

$$\log |\mathcal{Q}_{max}| \le n \cdot \max \left\{ \frac{h(\gamma) - 3\beta h(\gamma) + 2\beta - 2\varepsilon\beta + \beta h(\mu) + \beta h(1/2 - \mu)}{8}, (1 - \varepsilon)\beta \right\}. \tag{17}$$

Proof. To make all $|Q_i|$ and $|Q_i'|$ equal $n \frac{(h(\gamma)-3\beta h(\gamma)+2\beta-2\varepsilon\beta+\beta h(\mu)+\beta h(1/2-\mu))}{8}$ we set sizes of L_i and R_i as follows:

$$|L_1| = n \frac{h(\gamma) - 3\beta h(\gamma) - 6\beta + 6\varepsilon\beta + \beta h(\mu) + \beta h\left(\frac{1}{2} - \mu\right)}{8h(\gamma)} = |R_1|$$
(18)

$$|L_2| = |L_3| = n \frac{h(\gamma) - 3\beta h(\gamma) + 2\beta - 2\varepsilon\beta + \beta h(\mu) + \beta h\left(\frac{1}{2} - \mu\right)}{8h(\gamma)} = |R_2| = |R_3|$$

$$(19)$$

$$|L_4| = n \frac{h(\gamma) - 3\beta h(\gamma) + 2\beta - 2\varepsilon\beta - 7\beta h(\mu) + \beta h\left(\frac{1}{2} - \mu\right)}{8h(\gamma)}$$
(20)

$$|R_4| = n \frac{h(\gamma) - 3\beta h(\gamma) + 2\beta - 2\varepsilon\beta + \beta h(\mu) - 7\beta h\left(\frac{1}{2} - \mu\right)}{8h(\gamma)}$$
(21)

If all $|L_i|$ and $|R_i|$ are non-negative, then the statement of the lemma holds. Assume that some values are negative. In this case we replace all $|L_i| < 0$ with $|L_i| = 0$ and scale all the $|L_i|$ in such a way that $|L_1| + |L_2| + |L_3| + |L_4| = |L|$. Then we do the same procedure with $|R_i|$. We can see that for every $|L_i| = 0$, $|\mathcal{Q}_i| \le 2^{(1-\varepsilon)\beta n}$, and for $|L_i| > 0$, $|\mathcal{Q}_i| \le n \frac{(h(\gamma) - 3\beta h(\gamma) + 2\beta - 2\varepsilon\beta + \beta h(\mu) + \beta h(1/2 - \mu))}{8}$. The same works for $|\mathcal{Q}_i'|$. Thus, the lemma holds.

We have set all the parameters and now we are ready to estimate the time and space complexity of Algorithm 2.

4.4 Time Complexity

The running time is dominated by the following parts.

- Repeating the whole algorithm $2^{\lambda n}$ times (Line 1).
- Constructing Q_1, Q_2, Q_3, Q_4 and Q'_1, Q'_2, Q'_3, Q'_4 takes $O^*(|Q_{max}| + 2^{\beta n})$ time (Line 6).
- Repeating q times (Line 8).
- Finding \mathcal{L}, \mathcal{R} using Lemma 4.6 in time $O^*\left(|\mathcal{Q}_{max}|^2 + \frac{|\mathcal{Q}_{max}|^4}{pq} + \binom{|M|}{\max\{\mu, 1/2 \mu\}|M|}\right)$ (Line 9).
- Solving WOV(\mathcal{L}, \mathcal{R}) using Lemma 2.2 in time O^* (($|\mathcal{L}| + |\mathcal{R}|$) $2^{\beta n(1-h(1/4))}$) (Line 10). The O^* here hides logarithmic factor in $|\mathcal{L}| + |\mathcal{R}|$ that grows polynomially in n.

Let us consider an iteration of the loop on line 3. Note that the values of $p, q, |Q_{max}|$ and μ are now determined. The running time of Algorithm 2 on this specific iteration is

$$O^* \left(q \cdot \left(|\mathcal{Q}_{max}|^2 + \frac{|\mathcal{Q}_{max}|^4}{pq} + \left(\frac{|\mathcal{Q}_{max}|^4}{pq} + 2^{\beta nh(\max\{\mu, 1/2 - \mu\})} \right) 2^{\beta n(1 - h(1/4))} \right) \right).$$

To bound the total running time of Algorithm 2, we can multiply the number of the iterations $\Theta^*(2^{\lambda n})$ by the upper bound on the slowest iteration.

Lemma 4.11. The total running time of the Algorithm 2 is $O^*(2^{n/2}+2^{T(\alpha,\beta)\cdot n})$, where we define $T(\alpha,\beta)$ as follows:

$$T(\alpha, \beta) = \frac{1}{2} \left(h(\gamma) - 3\beta h(\gamma) + 3\beta + 2\lambda \right). \tag{22}$$

Here, $\gamma = \gamma(\alpha)$ and $\lambda = \lambda(\alpha)$.

Proof. As it was shown the total running time does not exceed

$$O^* \left(2^{\lambda n} \cdot q \left(|\mathcal{Q}_{max}|^2 + \frac{|\mathcal{Q}_{max}|^4}{pq} + \left(\frac{|\mathcal{Q}_{max}|^4}{pq} + 2^{\beta nh(\max\{\mu, 1/2 - \mu\})} \right) 2^{\beta n(1 - h(1/4))} \right) \right) ,$$

where the values of p, q, $|Q_{max}|$ and μ correspond to the slowest iteration of the loop on line 3.

Since p and q can vary only with polynomial factors, here by them we will denote the upper and lower bounds on their values, so let $\log p = \frac{(1-\varepsilon)}{2}\beta n$ and $\log q = \frac{1}{2}n - \lambda n - 2\log|\mathcal{Q}_{max}|$. Let us estimate the logarithm of the first summand inside the parenthesis:

$$\log(2^{\lambda n} \cdot q \cdot |\mathcal{Q}_{max}|^2) = \lambda n + \underbrace{\frac{1}{2}n - \lambda n - 2\log|\mathcal{Q}_{max}|}_{\log q} + 2\log|\mathcal{Q}_{max}| = \frac{1}{2}n.$$

We estimate the last term of the sum (the fourth one) assuming that $\mu \geq \frac{1}{2} - \mu$, the other case can be shown similarly:

$$\log \left(2^{\beta nh(\mu)+\beta n(1-h(1/4))}\right) = \beta nh(\mu) + \beta n - \beta nh(1/4) \le 2\beta n.$$

Recall that we choose $\beta < 0.15$, then $2\beta n \leq 0.5n$.

The second summand is at most the third one, so it suffices to estimate the logarithm of the third summand. Recall that that $\log |\mathcal{Q}_{max}| \leq n \cdot \max\{\frac{h(\gamma) - 3\beta h(\gamma) + 2\beta - 2\varepsilon\beta + \beta h(\mu) + \beta h(1/2 - \mu)}{8}, (1 - \varepsilon)\beta\}$. If $\log |\mathcal{Q}_{max}| \leq (1 - \varepsilon)\beta n$, then the third summand is dominated by $\frac{1}{2}(9\beta - 2\beta h(\frac{1}{4}))$ and substituting $\beta = 0.13$ yields value $\approx 0.48 < 0.5$. Therefore, this case is not considered further.

$$\begin{split} &\log(2^{\lambda n}q\frac{|\mathcal{Q}_{max}|^4}{pq}2^{\beta n(1-h(1/4))})\\ &=\lambda n + 4\log|\mathcal{Q}_{max}| - \underbrace{\frac{1-\varepsilon}{2}\beta n}_{\log p} + \beta n - \beta nh(1/4)\\ &=\lambda n + n\frac{h(\gamma) - 3\beta h(\gamma) + 3\beta - \varepsilon\beta + \beta h(\mu) + \beta h(1/2 - \mu) - 2\beta h(1/4)}{2} \end{split}$$

Having $(-\varepsilon\beta) \le 0$, $h(\mu) + h(1/2 - \mu) \le 2h(1/4)$, this expression can be dominated by

$$\lambda n + n \frac{h(\gamma) - 3\beta h(\gamma) + 3\beta}{2}$$
.

So, the total running time of the algorithm is $O^*(2^{n/2} + 2^{T(\alpha,\beta)\cdot n})$, where

$$T(\alpha, \beta) = \frac{1}{2} \left(h(\gamma) - 3\beta h(\gamma) + 3\beta + 2\lambda \right). \tag{23}$$

Corollary 4.1. If for every $\alpha \in [0,1]$, $T(\alpha,\beta(\alpha)) \leq 0.5$, then Algorithm 2 has running time $O^*(2^{0.5n})$. We complete the analysis in Section 4.6.

4.5Space Complexity

The space usage is dominated by the following parts.

- Constructing Q_1, Q_2, Q_3, Q_4 and Q'_1, Q'_2, Q'_3, Q'_4 with $O^*(|Q_{max}|)$ space (Line 6).
- Finding \mathcal{L}, \mathcal{R} using Lemma 4.6 with $O^*\left(|\mathcal{Q}_{max}| + \frac{|\mathcal{Q}_{max}|^4}{pq} + \binom{|M|}{\max\{\mu, 1/2 \mu\}|M|}\right)$ space (Line 9).
- Solving WOV(\mathcal{L}, \mathcal{R}) using Lemma 2.2 with $O^*(|\mathcal{L}| + |\mathcal{R}| + 2^{\beta n})$ space (Line 10).

For a specific iteration of the loop on line 3, the space usage is at most the sum of the above expressions, where $|\mathcal{L}|$, $|\mathcal{R}| = O^*(\frac{|\mathcal{Q}_{max}|^4}{pq} + \binom{|M|}{\max\{\mu,1/2-\mu\}})|M|)$. The total space complexity of Algorithm 2 can be bounded by the space usage of an iteration, requiring the most space. Therefore, it suffices to prove that $2^{\beta n}$, $\binom{|M|}{\max\{\mu,1/2-\mu\}|M|}$, $|\mathcal{Q}_{max}|$ and $\frac{|\mathcal{Q}_{max}|^4}{pq}$ are at most $O^*(2^{0.246n})$

on each iteration.

We can see that the following lemma holds.

Lemma 4.12. If for every $\alpha \in [0,1]$, $S(\alpha,\beta(\alpha)) \leq 0.246$, then the total space usage of Algorithm 2 does not exceed O^* (2^{0.246n}).

Lemma 4.13. The total space usage of Algorithm 2 is $O^*(2^{0.246n} + 2^{S(\alpha,\beta) \cdot n})$, where $S(\alpha,\beta)$ is defined as follows:

$$S(\alpha,\beta) = \frac{1}{4} \left(3h(\gamma) - 9\beta h(\gamma) + 6\beta h(\frac{1}{4}) + 4\beta - 2 + 4\lambda \right). \tag{24}$$

Here, we interpret γ as $\gamma = \gamma(\alpha)$ and λ as $\lambda(\alpha)$.

Proof. As we have already discussed, it is sufficient to estimate the following values: $\binom{|M|}{\max\{\mu,1/2-\mu\}|M|}$, $|\mathcal{Q}_{max}|$ and $\frac{|\mathcal{Q}_{max}|^4}{pq}$, for the iteration of the loop in line 3 that requires the most space. To do that, we estimate their logarithms. The first one is way smaller than 0.246n, since

 $\beta < 0.15$. The second one is equal to $\beta nh(\mu) \leq \beta n \leq 0.246n$. Recall that that $\log |\mathcal{Q}_{max}| \leq n \cdot \max\{\frac{h(\gamma) - 3\beta h(\gamma) + 2\beta - 2\varepsilon\beta + \beta h(\mu) + \beta h(1/2 - \mu)}{8}, (1 - \varepsilon)\beta\}$. If $\log |\mathcal{Q}_{max}| \leq (1 - \varepsilon)\beta n$, then the third value is dominated by $\frac{1}{2}(11\beta - 1)$ and substituting $\beta = 0.13$ yields value $\approx 0.215 < 0.246$. Therefore, this case is not considered further.

$$\log |\mathcal{Q}_{max}| = \frac{1}{8} \left(h(\gamma) - 3\beta h(\gamma) + 2\beta - 2\varepsilon\beta + \beta h(\mu) + \beta h(\frac{1}{2} - \mu) \right) n.$$

Having $(-2\varepsilon\beta) \le 0$, $h(\mu) + h(1/2 - \mu) \le 2$ and $\beta < 0.15$, we get

$$\log |\mathcal{Q}_{max}| \le \frac{1}{8} (1+\beta)n < 0.246n.$$

It is left to estimate $\log \frac{|\mathcal{Q}_{max}|^4}{na}$.

$$\log \frac{|\mathcal{Q}_{max}|^4}{pq} = 4\log |\mathcal{Q}_{max}| - \underbrace{\frac{1-\varepsilon}{2}\beta n}_{p} - \underbrace{\frac{1}{2}n + 2\log |\mathcal{Q}_{max}| + \lambda n}_{q} =$$

$$= \lambda n + n \frac{3h(\gamma) - 9\beta h(\gamma) + 3\beta h(\mu) + 3\beta h(1/2 - \mu) + 4\beta - 4\varepsilon\beta - 2}{4\beta n + 2\beta n + 2\beta$$

Having $(-4\varepsilon\beta) \le 0$ and $h(\mu) + h(1/2 - \mu) \le 2h(1/4)$, we dominate this expression by

$$\lambda n + \frac{3h(\gamma) - 9\beta h(\gamma) + 6\beta h(1/4) + 4\beta - 2}{4}.$$

That concludes the proof.

Corollary 4.2. If for every $\alpha \in [0,1]$, $S(\alpha,\beta(\alpha)) \leq 0.246$, then the total space usage of Algorithm 2 does not exceed O^* ($2^{0.246n}$).

We complete the analysis in the following section.

4.6 Parameter Substitution

Lemma 4.14. [Maximum point for function of time complexity] The function $T(\alpha, \beta)$ (see (22)) satisfies $T(\alpha, \beta) \leq T(0.5, 0.13)$ when β is defined as in (14).

Proof of Lemma 4.14. Note that

$$T(\alpha,\beta) = \frac{1}{2} \left(h(\gamma) - 3\beta h(\gamma) + 3\beta + 2(\alpha h \left(\frac{3\beta}{2\alpha}\right) + (1-\alpha)h \left(\frac{3\beta}{2(1-\alpha)}\right) - h(3\beta))) \right),$$

$$\frac{\partial^2 T(\alpha,\beta)}{\partial \alpha^2} = \frac{18\alpha^2\beta - 2\alpha^2 - 18\alpha\beta - 9\beta^2 + 2\alpha + 6\beta}{\alpha(2\alpha - 3\beta)(1-\alpha)(-2 + 3\beta + 2\alpha)\ln(2)}.$$

Consider two cases (the case $\alpha > 0.5$ is symmetrical).

1. $0.45 \le \alpha \le 0.5$.

In this regime, we have $\beta=0.13$, the roots of the polynomials of the denominator and numerator are either <0.2 or ≥ 0.5 . By checking that $\frac{\partial T(\alpha,\beta)}{\partial \alpha}|_{a=\frac{1}{2}}=0$ and $\frac{\partial^2 T(\alpha,\beta)}{\partial \alpha^2}|_{a=\frac{1}{2}}=0$, we verify that $a=\frac{1}{2}$ is indeed the maximum and have

$$T(\alpha, 0.13) \le T(0.5, 0.13)$$
.

2. $\alpha < 0.45$.

In this case, $\beta=0$. Then all the roots of the polynomials of the denominator and numerator are either 0 or 1. By checking that $\frac{\partial^2 T(\alpha,\beta)}{\partial^2 \alpha}|_{\alpha \in [0,0.45]} < 0$ and $\frac{\partial T(\alpha,\beta)}{\partial \alpha}|_{\alpha \in [0,0.45]} > 0$, we get that

$$T(\alpha, 0) \le T(0.45, 0) \le T(0.5, 0.13)$$
.

Lemma 4.15. [Maximum point for function of space complexity] The function $S(\alpha, \beta)$ (see (24)) satisfies $S(\alpha, \beta) \leq S(0.5, 0.13)$ when β is defined as in (14).

Proof of Lemma 4.15. Recall that

$$S(\alpha,\beta) = \frac{1}{4}(3h(\gamma) - 9\beta h(\gamma) + 6\beta h\left(\frac{1}{4}\right) + 4\beta - 2 + 4(\alpha h\left(\frac{3\beta}{2\alpha}\right) + (1-\alpha)h\left(\frac{3\beta}{2(1-\alpha)}\right) - h(3\beta))).$$

Consider two cases ($\alpha > 0.5$ is symmetrical).

1. $0.45 \le \alpha \le 0.5$. Then, $\beta = 0.13$.

$$\frac{\partial^2 S(\alpha, \beta)}{\partial \alpha^2} = -\frac{3(7\alpha^2\beta - \alpha^2 - 7\alpha\beta - 3\beta^2 + \alpha + 2\beta)}{\alpha(\alpha - 1)(2\alpha - 3\beta)(3\beta + 2\alpha - 2)\ln(2)}$$

One can check that all the roots of the numerator and denominator are either < 0.2 or > 0.5. As $\frac{\partial^2 S(\alpha,\beta)}{\partial \alpha^2}|_{\alpha=\frac{1}{2}} < 0$, we conclude that S is concave on the interval [0.45,0.5]. The only thing left to do is to verify that $\frac{\partial S(\alpha,\beta)}{\partial \alpha}|_{\alpha=\frac{1}{2}} = 0$ which implies that $S(\alpha,0.13) \leq S(\frac{1}{2},0.13)$ if $0.45 \leq \alpha \leq 0.5$.

2. $\alpha < 0.45$. Then, $\beta = 0$.

$$S(\alpha, 0) = -\frac{1}{4} (3\alpha \log(\alpha) - 3\alpha \log(1 - \alpha) + 2\log(2) + 3\log(1 - \alpha))$$
$$\frac{\partial^2 S(\alpha, 0)}{\partial \alpha^2} = -\frac{3}{4\alpha(1 - \alpha)\ln(2)} < 0$$

Checking that $\frac{\partial S(\alpha,0)}{\partial \alpha}|_{\alpha \in [0,0.45]} > 0$ we get that it is maximal at $\alpha = 0.45$ and $S(\alpha,0) \leq S(0.45,0) \leq S(0.5,0.13)$. This concludes the proof.

Hence, the time and space complexity do not exceed T(0.5, 0.13) and S(0.5, 0.13), respectively (for any α and β as in (14)). Recall that $\lambda = h(3\beta) - \alpha h(\frac{3\beta}{2\alpha}) - (1-\alpha)h(\frac{3\beta}{2(1-\alpha)})$, and $\gamma = \frac{\alpha - \frac{3}{2}\beta}{1-3\beta}$, see Equation (4). By plugging $\alpha = 0.5, \beta = 0.13$, we get the following values for the parameters:

$$\begin{split} \gamma &= 0.5, \\ h(\gamma) &= 1, \\ \lambda &= 0, \\ T(0.5, 0.13) &= \frac{1}{2} \cdot (1 - 3 \cdot 0.13 + 3 \cdot 0.13) = \frac{1}{2}, \\ S(0.5, 0.13) &\leq \frac{1}{4} \cdot (3 - 9 \cdot 0.13 + 6 \cdot 0.13 \cdot 0.812 + 4 \cdot 0.13 - 2) \leq 0.246. \end{split}$$

Combining statements 4.1 and 4.2, we get that the algorithm runs in time $O^*(2^{0.5n})$ and space $O^*(2^{0.246n})$, which completes the proof of Theorem 1.4.

4.7 Algorithm for Weighted Orthogonal Vectors

Definition 2.1. [Weighted Orthogonal Vectors, WOV] Given families of N weighted sets $\mathcal{A}, \mathcal{B} \subseteq 2^{[d]} \times \mathbb{N}$, and a target integer t, the problem is to find $(A, w_A) \in \mathcal{A}$ and $(B, w_B) \in \mathcal{B}$ such that $A \cap B = \emptyset$ and $w_A + w_B = t$.

Definition 4.1 (Orthogonal Vectors, OV). Given families of vectors $\mathcal{A}, \mathcal{B} \subseteq 2^{[d]}$. Decide if there exists a pair $A \in \mathcal{A}, B \in \mathcal{B}$ such that $A \cap B = \emptyset$.

Lemma 4.16. For any $\mu \in [0, \frac{1}{2}]$, there is a Monte-Carlo algorithm that, given $\mathcal{A} \subseteq \binom{[d]}{\mu d}$ and $\mathcal{B} \subseteq \binom{[d]}{(1/2-\mu)d}$, solves $\mathrm{OV}(\mathcal{A},\mathcal{B})$ in time

$$\widetilde{O}\left((|\mathcal{A}|+|\mathcal{B}|)2^{d(1-h(1/4))}\right)$$

and space $\widetilde{O}(|\mathcal{A}| + |\mathcal{B}| + 2^d)$.

Proof. In case $\mu \in [0.2, 0.3]$, this becomes exactly Theorem 6.1 from [12] applied for $\alpha = \frac{1}{2}, \sigma = 2\mu$. The restriction $\sigma \in [0.4, 0.6]$ in that theorem appears due to the following lemma.

Lemma 4.17 (Lemma B.1 from [12]). For large enough n and $\lambda \in [0.4, 0.5]$ and $\sigma \in [0.4, 0.6]$, the following inequality holds:

$$\min_{x} \left\{ \frac{\binom{1-\lambda\sigma}{x-\lambda\sigma}_{n} + \binom{1-(1-\sigma)\lambda}{x}_{n}}{\binom{1-\lambda}{x-\lambda\sigma}_{n}} \right\} \le 2^{n(1/2+\lambda-h(\lambda/2))} n^{O(1)}.$$

Here, $\binom{\alpha}{\beta}_n$ denotes $\binom{\alpha n}{\beta n}$.

We need that condition to hold for any $\sigma \in [0,1]$ when $\lambda = 0.5$. For this reason, we adjust the statement of the lemma.

Lemma 4.18. For large enough n and $\sigma \in [0,1]$ the following inequality holds:

$$\min_{x} \left\{ \frac{\binom{1-\sigma/2}{x-\sigma/2}_{n} + \binom{1-(1-\sigma)/2}{x}_{n}}{\binom{1/2}{x-\sigma/2}_{n}} \right\} \leq 2^{n(1-h(1/4))} n^{O(1)}.$$

Proof. We will use $\mu = \frac{\sigma}{2}$, $\mu \in [0, 0.5]$ to simplify the calculations.

$$\min_{x} \left\{ \frac{\binom{1-\mu}{x-\mu}_{n} + \binom{\frac{1}{2}+\mu}{x}_{n}}{\binom{1/2}{x-\mu}_{n}} \right\} = O^{*} \left(\min_{x} \left\{ 2^{\max\{(1-\mu)h(\frac{x-\mu}{1-\mu}), (\frac{1}{2}+\mu)h(\frac{2x}{1+2\mu})\} - \frac{1}{2}h(2x-2\mu)} \right\} \right) \le C^{*} \left(\min_{x} \left\{ 2^{\max\{(1-\mu)h(\frac{x-\mu}{1-\mu}), (\frac{1}{2}+\mu)h(\frac{2x}{1+2\mu})\} - \frac{1}{2}h(2x-2\mu)} \right\} \right) \le C^{*} \left(\min_{x} \left\{ 2^{\max\{(1-\mu)h(\frac{x-\mu}{1-\mu}), (\frac{1}{2}+\mu)h(\frac{2x}{1+2\mu})\} - \frac{1}{2}h(2x-2\mu)} \right\} \right) \le C^{*} \left(\min_{x} \left\{ 2^{\max\{(1-\mu)h(\frac{x-\mu}{1-\mu}), (\frac{1}{2}+\mu)h(\frac{2x}{1+2\mu})\} - \frac{1}{2}h(2x-2\mu)} \right\} \right) \le C^{*} \left(\min_{x} \left\{ 2^{\max\{(1-\mu)h(\frac{x-\mu}{1-\mu}), (\frac{1}{2}+\mu)h(\frac{2x}{1+2\mu})\} - \frac{1}{2}h(2x-2\mu)} \right\} \right) \le C^{*} \left(\min_{x} \left\{ 2^{\min_{x} \left\{$$

(we use $x = \frac{1}{2} + \log(3)(\mu - \frac{1}{4})$ as a good enough approximation of the minimum)

$$\leq O^*(2^{\max\{(1-\mu)h(\frac{(\frac{1}{2}+\log(3)(\mu-\frac{1}{4}))-\mu}{1-\mu}),(\frac{1}{2}+\mu)h(\frac{2(\frac{1}{2}+\log(3)(\mu-\frac{1}{4}))}{1+2\mu})\}-\frac{1}{2}h(2(\frac{1}{2}+\log(3)(\mu-\frac{1}{4}))-2\mu)})$$

Let:

$$f_1 = (1 - \mu)h\left(\frac{\left(\frac{1}{2} + \log(3)\left(\mu - \frac{1}{4}\right)\right) - \mu}{1 - \mu}\right) - \frac{1}{2}h\left(2\left(\frac{1}{2} + \log(3)\left(\mu - \frac{1}{4}\right)\right) - 2\mu\right),$$

$$f_2 = \left(\frac{1}{2} + \mu\right) h\left(\frac{2\left(\frac{1}{2} + \log(3)\left(\mu - \frac{1}{4}\right)\right)}{1 + 2\mu}\right) - \frac{1}{2}h\left(2\left(\frac{1}{2} + \log(3)\left(\mu - \frac{1}{4}\right)\right) - 2\mu\right).$$

We will show that both of these functions have a maximum at $\mu = \frac{1}{4}$. Indeed,

$$\frac{\partial^2 f_1}{\partial \mu^2} = \frac{(3\log 3 - 2)^2}{(1 - \mu)(4\log 3\mu - 4\mu - \log 3 + 2)(4\log(3)\mu - \log(3) - 2)},$$

$$\frac{\partial^2 f_2}{\partial \mu^2} = -\frac{8(\log 3 - 1)^2}{(4\log(3)\mu - 4\mu - \log 3 + 2)(4\log(3)\mu - 4\mu - \log(3))} \; .$$

By analyzing the roots of the polynomials in the denominators one can show that for $0 \le \mu \le \frac{1}{2}$: $\frac{\partial^2 f_1}{\partial \mu^2} < 0$ and $\frac{\partial^2 f_2}{\partial \mu^2} < 0$, which means that there is at most one maximum for each of the functions on this interval. Now we just need to notice that

$$\frac{\partial f_1}{\partial \mu}|_{\mu=\frac{1}{2}} = \frac{\partial f_2}{\partial \mu}|_{\mu=\frac{1}{2}} = 0$$

and $f_1|_{mu=\frac{1}{2}} = f_2|_{mu=\frac{1}{2}} = 1 - h(\frac{1}{4})$. This concludes the proof.

Now we can apply the proof of Theorem 6.1 from [12] with Lemma 4.17 replaced by Lemma 4.18. That completes the proof of Lemma 4.16.

Lemma 2.2. For any $\sigma \in [0, \frac{1}{2}]$, there is a Monte-Carlo algorithm that, given $\mathcal{A} \subseteq \binom{[d]}{\sigma d} \times \mathbb{N}$, $\mathcal{B} \subseteq \binom{[d]}{(1/2-\sigma)d} \times \mathbb{N}$ and a target integer t, solves $WOV(\mathcal{A},\mathcal{B})$ in time $\widetilde{O}\left((|\mathcal{A}|+|\mathcal{B}|)2^{d(1-h(1/4))}\right)$ and space $\widetilde{O}\left(|\mathcal{A}|+|\mathcal{B}|+2^d\right)$.

Proof of Lemma 2.2. To prove the lemma we show a reduction from WOV to OV. We split \mathcal{A} and \mathcal{B} into the following groups of sets with the same sum in time $\widetilde{O}(|\mathcal{A}| + |\mathcal{B}|)$.

$$\mathcal{L}(a) = \{A \colon (A, a) \in \mathcal{A}\},\$$

$$\mathcal{R}(b) = \{B \colon (B, b) \in \mathcal{B}\}.$$

As in the algorithm for 2-SUM, we use the two pointers technique to iterate over every possible (a, b) such that a + b = t in time \widetilde{O} (number of groups). For every such pair we call $\mathrm{OV}(\mathcal{L}(a), \mathcal{R}(b))$. By Lemma 4.16, its running time and space usage depend linearly on $(|\mathcal{L}(a)| + |\mathcal{R}(b)|)$, so the total time and space complexity are at most as for $\mathrm{OV}(\mathcal{A}, \mathcal{B})$.

Acknowledgments

Research is partially supported by the grant 075-15-2022-289 for creation and development of Euler International Mathematical Institute, and by the Foundation for the Advancement of Theoretical Physics and Mathematics "BASIS".

References

- [1] Shyan Akmal, Lijie Chen, Ce Jin, Malvika Raj, and Ryan Williams. Improved Merlin–Arthur protocols for central problems in fine-grained complexity. In *ITCS*, volume 215 of *LIPIcs*, pages 3:1–3:25. Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2022.
- [2] Noga Alon, Raphael Yuster, and Uri Zwick. Color-coding. J. ACM, 42(4):844-856, 1995.
- [3] Marco L. Carmosino, Jiawei Gao, Russell Impagliazzo, Ivan Mihajlin, Ramamohan Paturi, and Stefan Schneider. Nondeterministic extensions of the strong exponential time hypothesis and consequences for non-reducibility. In *ITCS 2016*, pages 261–270. ACM, 2016.
- [4] Timothy M. Chan. More logarithmic-factor speedups for 3sum, (median, +)-convolution, and some geometric 3sum-hard problems. *ACM Trans. Algorithms*, 16(1):7:1–7:23, 2020.
- [5] Xi Chen, Yaonan Jin, Tim Randolph, and Rocco A. Servedio. Subset sum in time $2^{n/2}$ / poly(n). In APPROX/RANDOM, volume 275 of LIPIcs, pages 39:1–39:18. Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2023.
- [6] Ellis Horowitz and Sartaj Sahni. Computing partitions with applications to the knapsack problem. J. ACM, 21(2):277–292, 1974.
- [7] Nick Howgrave-Graham and Antoine Joux. New generic algorithms for hard knapsacks. In *Annual International Conference on the Theory and Applications of Cryptographic Techniques*, pages 235–256. Springer, 2010.
- [8] Hamidreza Jahanjou, Eric Miles, and Emanuele Viola. Local reductions. In *ICALP 2015*, pages 749–760. Springer, 2015.
- [9] Hendrik W. Lenstra, Jr. and Carl Pomerance. Primality testing with gaussian periods. *Journal of the European Mathematical Society*, 21(4):1229–1269, 2019.
- [10] Andrea Lincoln, Virginia Vassilevska Williams, Joshua R. Wang, and R. Ryan Williams. Deterministic time-space trade-offs for k-SUM. In *ICALP*, volume 55 of *LIPIcs*, pages 58:1–58:14. Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2016.
- [11] Jesper Nederlof. A short note on Merlin–Arthur protocols for subset sum. *Inf. Process. Lett.*, 118:15–16, 2017.
- [12] Jesper Nederlof and Karol Węgrzycki. Improving Schroeppel and Shamir's algorithm for subset sum via orthogonal vectors. In *STOC*, pages 1670–1683. ACM, 2021.
- [13] Nicholas Pippenger and Michael J. Fischer. Relations among complexity measures. J. ACM, 26(2):361-381, 1979.
- [14] Richard Schroeppel and Adi Shamir. A $T=O(2^{n/2})$, $S=O(2^{n/4})$ algorithm for certain NP-complete problems. SIAM J. Comput., 10(3):456–464, 1981.