

Quantum particle localization observables on Cauchy surfaces of Minkowski spacetime and their causal properties

Carmine De Rosa^a and Valter Moretti^b

Department of Mathematics, University of Trento, and INFN-TIFPA
via Sommarive 14, I-38123 Povo (Trento), Italy.

^acarmine.derosa@unitn.it

^bvalter.moretti@unitn.it

Abstract

We introduce and study a general notion of spatial localization on spacelike smooth Cauchy surfaces of quantum systems in Minkowski spacetime. The notion is constructed in terms of a coherent family of normalized POVMs, one for each said Cauchy surface. We prove that a family of POVMs of this type automatically satisfies a causality condition which generalizes Castrigiano's one and implies it when restricting to flat spacelike Cauchy surfaces. As a consequence no conflict with Hegerfeldt's theorem arises. We furthermore prove that such families of POVMs do exist for massive Klein-Gordon particles, since some of them are extensions of already known spatial localization observables. These are constructed out of positive definite kernels or are defined in terms of the stress-energy tensor operator. Some further features of these structures are investigated, in particular, the relation with the triple of Newton-Wigner selfadjoint operators and a modified form of Heisenberg inequality in the rest 3-spaces of Minkowski reference frames.

Contents

1	Introduction	2
1.1	The subtle issue of spatial localization of quantum relativistic systems	2
1.2	Structure of the paper	8
1.3	General definitions, notations, and conventions	9
2	Causal structures of spacetimes, Minkowski spacetime	10
2.1	Globally hyperbolic spacetimes and their causal structure	10
2.2	Minkowski spacetime geometric and causal structures	12
3	Spacelike Cauchy localization observables in \mathbb{M} and a general causality condition	15
3.1	A general notion of spatial localization in terms of POVMs	15
3.2	The general causality condition for spacelike Cauchy localization observables . .	17
3.3	The general causality condition is valid for every spacelike Cauchy localization observable	18

4	Massive KG particles, conserved currents	23
4.1	One-particle Hilbert space of real massive KG particles in \mathbb{M}	23
4.2	Conserved quantities on generic smooth Cauchy surfaces of \mathbb{M}	25
5	Spacelike Cauchy localization observables out of causal kernels of massive KG particles	26
5.1	A notion of spatial localization for massive KG particles out of causal kernels . . .	26
5.2	Properties of \mathbb{T}_Σ^g , for spacelike flat Cauchy surfaces Σ : Covariance, causality, no strict localizability, New	
5.3	A spacelike Cauchy localization observable $\mathbb{T}^g = \{\mathbb{T}_S^g\}_{S \in \mathcal{C}_M^s}$ for a massive KG particle	30
6	Spacelike Cauchy localization observables out of the stress-energy tensor of massive KG particles	
6.1	The POVM \mathbb{M}_Σ^n for massive KG particles	35
6.2	Properties of \mathbb{M}_Σ^n for spacelike flat Cauchy surfaces Σ : Covariance, causality, no strict localizability, New	
6.3	A spacelike Cauchy localization observable $\mathbb{M}^n = \{\mathbb{M}_S^n\}_{S \in \mathcal{C}_M^s}$ for massive KG particles.	40
7	Moments of \mathbb{T}_Σ^g and \mathbb{M}_Σ^n, Newton-Wigner operator, and Heisenberg inequality	42
8	Discussion	44
A	Proof of some propositions	45
B	Conditions for positive-definite kernels on \mathbb{R}^n	55

1 Introduction

1.1 The subtle issue of spatial localization of quantum relativistic systems

The study of notions of *spatial localization* at given time for a quantum relativistic particle can be traced back to the seminal paper by Newton and Wigner [30]. There, spatial localization was referred to the rest 3-space Σ , at given time, of an inertial (Minkowskian) reference frame. Later, guided by Mackey’s imprimitivity theory, Wightman established [36] an uniqueness theorem. He proved that the joint (projector-valued) spectral measure \mathbb{Q}_Σ on Σ of the triple of *Newton-Wigner selfadjoint operators* $N_\Sigma^1, N_\Sigma^2, N_\Sigma^3$ is the unique projector valued measure (PVM) which is covariant with respect to the Euclidean group of isometries of Σ and complies with some further technical hypotheses. If the notion of spatial localization is described in terms of selfadjoint observables, they must be the Newton-Wigner ones necessarily.

Unfortunately, these operators and their common spectral measure (their joint PVM) resulted to be plagued by a number of fundamental issues related to causality. The *Hegerfeldt theorem(s)* [21, 22] and the *Malament theorem* [26], with several modern reformulation mainly due to Busch [6] and Halvorson and Clifton [19], proved that some causality requirements and the request of energy positivity are definitely incompatible for quantum particles described as one-particle states of a quantum relativistic field (Wigner particles).

Malament's result and its modern extensions and reformulations are directly or indirectly related with the description of post measurement states, in terms of *Lüders-von Neuman projection postulate* or referring to a *Kraus decomposition* of the effects of the POVM describing the localization observable. This is a deep and outstanding issue [2] we shall not discuss in this paper.

Conversely, the various versions of the Hegerfeldt theorem only focus on the spatial detection probability of a quantum relativistic particle at given time. The most elementary version is like this. Let us consider the rest 3-space Σ of a Minkowski observer and a quantum relativistic free particle defined according to Wigner classification, with every mass $m \geq 0$ and every permitted spin s . Suppose that the probability to detect the particle, whose pure quantum state is represented by the normalized vector ψ , vanishes outside a *bounded spatial region* Δ at time $t = 0$. Then, the same state ψ gives rise to a *strictly positive* probability to find the particle *arbitrarily far* from Δ and at *arbitrarily small* time $t > 0$. In other words, *superluminal propagation of probability* shows up here. The hypotheses leading to Hegerfeldt's result are quite mild: the crucial one is *positivity of energy* (more precisely below boundedness) which is embodied in the Wigner's definition of particle¹. In that case, pure states are represented by unit vectors in the *one-particle Hilbert space* \mathcal{H} of the Fock space of the associated quantum field. In the proof of the theorem, the detection probability is supposed to be computed through a PVM or, more generally, by means of a POVM (see below) labelled by the sets of a given rest 3-space Σ of an inertial observer in Minkowski spacetime.

The impact on NW operators is evident. Probability distributions which vanish outside a bounded region Δ are trivially constructed when adopting a spatial localization notion in terms of a PVM on Σ . The position observable are in this case a triple of mutually compatible selfadjoint operators

$$X^k := \int_{\Sigma} x^k dP_{\Sigma}(x), \quad k = 1, 2, 3$$

in the Hilbert space \mathcal{H} . $P_{\Sigma} = P_{\Sigma}(\Delta)$ for $\Delta \in \mathcal{B}(\Sigma)$ (where henceforth $\mathcal{B}(\Sigma)$ is the family of Borel sets on Σ) is their joint spectral measure. A normalized vector in $\psi \in P_{\Sigma}(\Delta)\mathcal{H}$ immediately yields a probability distribution $\langle \psi | P(E) | \psi \rangle$ which vanishes if $E \cap \Delta = \emptyset$, fitting the hypothesis of Hegerfeldt's theorem. If we do not accept superluminal propagation of detection probability, on account of Hegerfeldt achievement, we are committed to reject any description of the spatial localization in terms of PVM, i.e., selfadjoint operators. The first victim of this reasoning is the very triple of Newton-Wigner operators $X^k = N_{\Sigma}^k$, $k = 1, 2, 3$ ².

Should we rule out spatial localization described by PVMs on the ground of the theoretical evidence of superluminal propagation of probability?

The answer needs a certain analysis. What we can control in laboratories in practice are

¹The proof of Hegerfeldt's theorem relies on properties of analytic continuations of one-parameter evolution semigroups and turns out quite involved. A quantitative analysis of the effects of the Hegerfeldt theorem in the form discussed in [21] appears in [17].

²As already found in [29], in the new picture arising from modern achievements as the present work, the Newton-Wigner observable is however recovered in terms of the *first moments* of permitted localization observables which are described in terms of POVMs.

just macroscopic objects and devices. At macroscopic level, superluminal propagation of information is forbidden. So, a better perspective to tackle this issue is wondering *whether or not the superluminal propagation of probability predicted by the Hegerfeldt theorem can be used to propagate superluminal macroscopic information*. The answer is positive, in case of states whose probability distribution at $t = 0$ vanishes outside a bounded spatial region. A corresponding ideal experiment was discussed in [29] by one of the authors of this work. The conclusion is that, to describe spatial localization on a rest 3-space Σ at some time of an inertial observer, the use of a suitable *normalized positive-operator valued measure* (POVM) is compulsory. A \mathcal{H} -POVM [7] on Σ is a map

$$\mathcal{B}(\Sigma) \ni \Delta \mapsto \mathbf{A}_\Sigma(A) \in \mathfrak{B}(\mathcal{H}) \quad (1)$$

such that $0 \leq \mathbf{A}_\Sigma(\Delta) \leq I$ for every $\Delta \in \mathcal{B}(\Sigma)$, where $\mathcal{B}(\Sigma) \ni \Delta \mapsto \langle \psi | \mathbf{A}_\Sigma(A) \psi' \rangle$ is a complex measure for every $\psi, \psi' \in \mathcal{H}$. \mathbf{A}_Σ is *normalized* if $\mathbf{A}_\Sigma(\Sigma) = I$. If $\psi \in \mathcal{H}$ is normalized as well, the meaning of $\langle \psi | \mathbf{A}_\Sigma(\Delta) \psi \rangle$ is the probability to find the system in Δ , when its state is ψ .

There are POVMs which do *not* admit probability distributions $\mathcal{B}(\Sigma) \ni E \mapsto \langle \psi | \mathbf{A}_\Sigma(E) \psi \rangle$ localized in bounded regions for any conceivable pure state $\psi \in \mathcal{H}$. Contrarily to PVMs which always permits probability distributions localized in bounded sets as viewed above. Hegerfeldt's theorem does not exclude POVMs to describe the spatial localization of relativistic quantum particles. More precisely, it permits the POVMs whose probability distributions to detect the particle is not supported in a bounded region for every pure state $\psi \in \mathcal{H}$. It is obviously necessary that probability distributions localized in bounded regions can be arbitrarily approximated by permitted distributions. It in fact happens in concrete POVMs describing spatial localization [8, 29, 9].

Mathematically speaking, at this juncture, it is convenient to give the following definition, where \mathbb{M} denotes the Minkowski spacetime and $\mathcal{C}_{\mathbb{M}}^{sf}$ is the family of all possible rest 3-spaces Σ of inertial observers at any given instant of their proper time.

Definition 1. Given a quantum system described in the Hilbert space \mathcal{H} , a **spatial localization observable** of it in \mathbb{M} is a collection $\mathbf{A} := \{\mathbf{A}_\Sigma\}_{\Sigma \in \mathcal{C}_{\mathbb{M}}^{sf}}$, where each \mathbf{A}_Σ is a normalized \mathcal{H} -POVM on $\mathcal{B}(\Sigma)$. ■

In principle, a *coherence condition* should be also imposed if one wants the probability to find the particle in Δ be independent of the rest space Σ containing it. One should require that, if $\Delta \subset \Sigma \cap \Sigma'$, then $\mathbf{A}_\Sigma(\Delta) = \mathbf{A}_{\Sigma'}(\Delta)$. However, the POVMs \mathbf{A}_Σ of all known spatial localization observables \mathbf{A} are absolutely continuous with respect to the Lebesgue measure of Σ . Therefore, since $\Delta \subset \Sigma \cap \Sigma'$ has zero measure if $\Sigma \neq \Sigma'$, this condition would not be effective because $\mathbf{A}_\Sigma(\Delta) = \mathbf{A}_{\Sigma'}(\Delta) = 0$ in any cases.

Definition 1 is nothing but Definition 18 stated in [29] for a *relativistic spatial localization observable* without the requirement of \mathcal{P}_+ -covariance.

An important remark is the following one. Consider the case $\Delta, \Delta' \in \Sigma$ and $\Delta \cap \Delta' = \emptyset$. Referring to quantum relativistic particles (with positive energy), operators $\mathbf{A}_\Sigma(\Delta)$ and $\mathbf{A}_\Sigma(\Delta')$ cannot commute, although Δ and Δ' are causally separated in the considered case. That is

due to general no-go results³ [6, 19, 11, 2]. This is a delicate issue somehow related to the *Reeh-Schlieder property* if embedding all the theoretical description in the framework of local algebras of observables. Roughly speaking, the operators $A_\Sigma(\Delta)$ cannot belong to a local algebra in the sense of *Haag-Kastler*. Whether or not this non-commutativity permits superluminal propagation of macroscopic information should be analyzed in a perspective (as in [15, 16]) where *measurement instruments* are explicitly studied. This analysis would concern post measurement states which are outside of the goals of this paper.

To get closer to the results achieved in this paper, we observe that there is a more sophisticated version of the Hegerfeldt theorem [22]: If a state $\psi \in \mathcal{H}$ produces a probability distribution which vanishes sufficiently rapidly at infinity at $t = 0$, then a more specific type of *superluminal propagation of probability* takes place again at later times.

The modern overall description of the types of violation was only recently formalized by Castrigiano. Let us quickly review the *causality condition* and the *causal time evolution condition*, introduced in [8] for localization POVMs in Minkowski spacetime \mathbb{M} , to reformulate the second form of Hegerfeldt's theorem [22].

In the following, $\mathcal{L}(\Sigma)$ is the family of Lebesgue sets of $\Sigma \in \mathcal{C}_M^{sf}$, and the considered POVMs are defined on this σ -algebra larger than the more usual⁴ $\mathcal{B}(\Sigma)$. Finally, if $\Sigma, \Sigma' \in \mathcal{C}_M^{sf}$ and $\Delta \in \mathcal{L}(\Sigma)$, we define the **region of influence of Δ on Σ'**

$$\Delta' := (J^+(\Delta) \cup J^-(\Delta)) \cap \Sigma'. \quad (2)$$

($J^\pm(\Delta)$ are the standard *causal sets* emanated from Δ as defined in Section 2.) The physical meaning of Δ' should be obvious. *It is the largest spatial region on Σ' which may have macroscopic causal relations with $\Delta \subset \Sigma$* , when assuming the causal structure of Minkowski spacetime based on the bounded propagation velocity of macroscopic physical signals.

Let us consider a spatial localization observable $A := \{A_\Sigma\}_{\Sigma \in \mathcal{C}_M^{sf}}$, where $A_\Sigma : \mathcal{L}(\Sigma) \rightarrow \mathfrak{B}(\mathcal{H})$.

Definition 2. A satisfies **Castrigiano's causality condition (CC)** [8] if

$$A_{\Sigma'}(\Delta') \geq A_\Sigma(\Delta) \quad \text{when } \Delta \in \mathcal{L}(\Sigma), \quad (3)$$

for every $\Sigma, \Sigma' \in \mathcal{C}_M^{sf}$. ■

The physical meaning of the condition above is evident: detection probability cannot propagate faster than the light speed. A weaker condition which is implied by CC but does not imply CC, is the *causal time evolution*. This condition – and not CC – was actually the specific subject of Hegerfeldt's investigation. It only considers the case where Σ and Σ' are rest spaces of a *common inertial observer* n , and thus they are related by means of the time evolution proper

³See Lemma 4 on p.13 of [11] and its proof in particular.

⁴The use of the Lebesgue σ -algebra $\mathcal{L}(\Sigma)$ in place of the Borel one $\mathcal{B}(\Sigma)$ is compulsory here. In fact, as discussed in [8], if defining the relevant POVMs on $\mathcal{B}(\Sigma)$ rather than on $\mathcal{L}(\Sigma)$ it is not guaranteed that $\Delta' \in \mathcal{B}(\Sigma')$ when $\Delta \in \mathcal{B}(\Sigma)$, making meaningless (3). It is however true that $\Delta' \in \mathcal{L}(\Sigma')$ for every $\Delta \subset \Sigma$ (even if $\Delta \notin \mathcal{B}(\Sigma)$!) as established in [8]. The use of $\mathcal{L}(\Sigma)$ is not a serious issue, even if initially dealing with POVMs defined on the Borel sets, since the standard completion procedure uniquely extends them to POVMs on $\mathcal{L}(\Sigma)$.

of n . Geometrically speaking, that is equivalent to saying that the normal vectors n_Σ and $n_{\Sigma'}$ coincide (with n).

Definition 3. A satisfies the **causal time evolution condition** condition (CT) [8] if

$$A_{\Sigma'}(\Delta') \geq A_\Sigma(\Delta) \quad \text{when } \Delta \in \mathcal{L}(\Sigma), \quad (4)$$

for every $\Sigma, \Sigma' \in \mathcal{C}_M^{sf}$ with $n_\Sigma = n_{\Sigma'}$. ■

The afore-mentioned second version of the Hegerfeldt theorem [22] (see also further formalizations in [9]) can be stated as follows, for a free quantum relativistic particle of any spin and strictly positive mass $m > 0$. (The result was actually extended [22] to more relativistic general systems in, also self-interacting, but satisfying energy positivity).

Theorem 4 ([22]). *Consider a spatial localization observable A for a massive Wigner particle described in the Hilbert space \mathcal{H} . If there is a unit vector $\psi \in \mathcal{H}$ such that the associated probability to find the particle outside a ball of radius r in a certain Σ is bounded by $K_1 e^{-K_2 r}$, for all $r > 0$ and constants $K_1 \geq 0$ and $K_2 \geq 2\frac{mc}{\hbar}$, then A fails to satisfy CT.*

Remark 5. As consequence, A fails to satisfy CC as well. ■

Castringiano proved in [8] that spatial localization observables exist which satisfy CC for spin 1/2 fermions. Another family of *causal* – namely satisfying CC – spatial localization observable $\{M_\Sigma^n\}_{\Sigma \in \mathcal{C}_M^{sf}}$ was obtained in [29] by one of the authors of this work for scalar, real, massive Klein-Gordon particles. That observable was constructed by generalizing a more specific observable introduced by D. Terno [34] and also studied in [29], where it was rigorously established that it satisfies CT. More recently, D. Castrigiano rigorously proved in [9] that the spatial localization observables for Klein-Gordon particles with causal kernels $\{T_\Sigma^g\}_{\Sigma \in \mathcal{C}_M^{sf}}$ introduced in [18, 20] satisfy CC. These three types of causal spatial localization observables also enjoy natural covariance properties with respect to the orthochronous Poincaré group in M as expected. In this sense, they are *relativistic* spatial localization observables (we shall come back to those features later in the paper).

A natural issue, discussed in the first part of this work, is whether or not we can physically admit a spatial localization observable that *violates* CC. In other words, *is it possible to transmit superluminal macroscopic information by exploiting it?*

It is not easy to re-adapt the ideal experiment presented in [29] to the case of a POVM and a state whose probability distribution *is not supported in a bounded region* but vanishes sufficiently fast at infinity to switch on the second version (Theorem 4) of Hegerfeldt's achievement, thus violating CT and CC. It seems to the author of [29] that a re-adaptation of the reasoning in [29] should involve some choice about a post-measurement state prescription or about some quantum instrument. As declared before, we do not want to pursue that route in this paper, since we do not intend to deal with more sophisticated theoretical notions than POVMs.

As a matter of fact, CC can be seen as a consequence of a more general causality condition we shall introduce in Section 3. The crucial overall idea⁵ is to extend the notion of spatial

⁵D. Castrigiano communicated to V.M. that R.F. Werner independently had a similar idea to prove CC.

localization to a broader class of 3-dimensional surfaces in Minkowski spacetime. Minkowski spacetime is in fact *globally hyperbolic*. In other words, it admits certain special 3-dimensional surfaces, called *Cauchy surfaces*, which can be used as the place where assigning initial data of causal (hyperbolic) PDEs. From the geometric perspective, each of these surfaces are met by all (inextendible) causal curves, exactly once by timelike ones. No macroscopic physical information can be transmitted from a region on a *spacelike* Cauchy surface to another separated region on the *same* Cauchy surface. Rest spaces of inertial reference frames at given times are just a special flat case of smooth Cauchy surfaces. *From a relativistic perspective, it seems natural to think of spacelike Cauchy surfaces as a generalization of the notion of space at a given time, where to localize quantum systems like particles*⁶. We therefore extend the notion of spatial localization observable to a more general notion where the possible regions Δ , where a particle can be detected, are subsets of generic spacelike Cauchy surface S . To this end, each (generally curved) Cauchy surface S is equipped with a *normalized POVM* A_S .

A *spacelike Cauchy localization observable* (Definition 21) is the family all these normalized POVMs, when S varies in the collection of all spacelike Cauchy surfaces. An important *coherence condition* is also imposed: if $\Delta \subset S \cap S'$, then $A_S(\Delta) = A_{S'}(\Delta)$. In other words, the probability to find the particle in Δ is independent of the Cauchy surface, but it is a function of Δ and the state $\psi \in \mathcal{H}$ only. Finally we also require that the probability to find a particle in a zero measure set on S vanishes.

At this juncture, a *general causality condition* GCC (Definition 26) for spacelike Cauchy localization observables can be stated, in the spirit of Definition 2. The general causality condition is just obtained by replacing the flat spacelike Cauchy surfaces Σ, Σ' in Definition 2 for generic spacelike Cauchy surfaces S, S' . This general causality condition evidently implies CC when restricting A to the POVMs A_Σ defined on the subfamily flat Cauchy surfaces Σ . A natural issue pops out here:

How is selective the general causality condition?

Quite surprisingly, the answer is that there is *no* selection at all: *every spacelike Cauchy localization observable automatically satisfies the general causality condition* (Theorem 32). As a corollary, *if a spatial localization observable $\{A_\Sigma\}_{\Sigma \in \mathcal{C}_M^{sf}}$ can be extended to a spacelike Cauchy localization observable $\{A_S\}_{S \in \mathcal{C}_M^s}$, then the former automatically satisfies CC*.

It seems quite remarkable that the result only uses a general, and very natural, notion of localization. The achieved result proves in fact the general causality relation should not be imposed as an independent postulate because it is physics that asks for it. In summary, the presented result proves that mere localizability implies causality independently of any kinematical argument.

The strategy to prove our main result stated above takes advantage of some general technical facts in Lorentzian geometry on existence of spacelike Cauchy surfaces adapted to given submanifolds with boundary [3, 4].

Other approaches exist where also considering many particles one renounces to energy positivity and defines a notion of spatial localization on generic Cauchy surfaces in terms of PVMs

⁶However, see [32] and below in the text for a different viewpoint on this issue.

[24, 25], finding similar causality conditions. A different general approach to the Born rule for the spatial localization of a particle formulated in a generic spacetime is discussed in [32]. There, a more general notion of 3-space at given time is introduced and analyzed (also relying on technical results in [27]). That notion of space is adapted (transverse) to the considered probability current and, for this reason, it is not necessarily spacelike nor a Cauchy surface.

The second part of the work is mostly devoted to prove that spacelike Cauchy localization observables *do exist*. Some of them are (uniquely determined) extensions $\{\mathbb{T}_S^g\}_{S \in \mathcal{C}_M^s}$ and $\{\mathbb{M}_S^n\}_{S \in \mathcal{C}_M^s}$ of the respective above mentioned spatial localization observables $\{\mathbb{T}_\Sigma^g\}_{\Sigma \in \mathcal{C}_M^{sf}}$ and $\{\mathbb{M}_\Sigma^n\}_{\Sigma \in \mathcal{C}_M^{sf}}$. The method to demonstrate the results in the second part uses ideas physically formulated in [18, 23, 20], made rigorous in [29, 9] and here further improved.

Some further results about the interplay of spatial localization observables, the NW operators, and the Heisenberg inequality appear in several spots of the main text and in a cumulative proposition in the last section.

In details, the achievements of this work are as follows.

- (1) **Theorem 32:** *Every spacelike Cauchy localization observable satisfies the general causality condition.*

Corollary 33: *If a spatial localization observable can be extended to a spacelike Cauchy localization observable, then it satisfies Castrigiano's causality condition.*

- (2) **Theorems 49 and 59:** *Both $\{\mathbb{T}_\Sigma^g\}_{\Sigma \in \mathcal{C}_M^{sf}}$ and $\{\mathbb{M}_\Sigma^n\}_{\Sigma \in \mathcal{C}_M^{sf}}$ uniquely extend to corresponding spacelike Cauchy localization observables $\{\mathbb{T}_S^g\}_{S \in \mathcal{C}_M^s}$ and $\{\mathbb{M}_S^n\}_{S \in \mathcal{C}_M^s}$. The former can be even defined for smooth Cauchy surfaces S which are not spacelike.*

- (3) **Theorems 44, 57, Proposition 60:** *Both $\{\mathbb{T}_\Sigma^g\}_{\Sigma \in \mathcal{C}_M^{sf}}$ and $\{\mathbb{M}_\Sigma^n\}_{\Sigma \in \mathcal{C}_M^{sf}}$ give rise to the Newton Wigner operators as their first moments on every inertial rest space Σ . The NW operators are the unique selfadjoint operators which satisfy this property. A generalized Heisenberg inequality holds in all cases.*

1.2 Structure of the paper

After a list of notations and conventions adopted in this work, Section 2 concerns a recap of the causal structure of globally hyperbolic spacetimes and Minkowski spacetime in particular. Section 3, after introducing the relevant definitions, presents our main results about causality: every spacelike Cauchy localization observable is causal. Section 4 collects some technical notions and results necessary to pass to the second set of achievements. These results are established in Sections 5 and 6: the possibility to extend some relevant spatial localization observables defined in [9] and [29] to corresponding spacelike Cauchy localization observables. In the same section, we shall discuss some features of spatial localization observables. Section 7 is devoted to discuss some general facts about Newton Wigner observables, Heisenberg inequality and spatial localization observables. After a final discussion in Section 8, the appendices contain the proofs of several technical lemmata and propositions asserted in the main text.

1.3 General definitions, notations, and conventions

Barring few changes (like the symbol for the flat Cauchy surfaces), we shall adopt the same notation as in [29].

Throughout $\mathbb{R}_+ := [0, +\infty)$, $\overline{\mathbb{R}_+} := \mathbb{R}_+ \cup \{+\infty\}$, and **smooth** means C^∞ . The light speed is $c = 1$. The normalized Planck constant is $\hbar = 1$. Furthermore we shall take advantage of the following notation:

$$\vec{x} \equiv (x^1, x^2, x^3) \in \mathbb{R}^3 \quad \text{and} \quad \nabla f = (\partial_{x^1} f, \partial_{x^2} f, \partial_{x^3} f)$$

if $f = f(\vec{x})$ is defined on a suitable open domain of \mathbb{R}^3 .

The *Lebesgue measure* (also restricted to the Borel sets) on \mathbb{R}^n will be denoted by $d^n x$, when x^1, \dots, x^n are orthonormal Cartesian coordinates on \mathbb{R}^n . We use also the notation $|B| := \int_B 1 d^n x$. The family of *Borel sets* on a topological space X will be denoted by $\mathcal{B}(X)$. The *Lebesgue σ -algebra* on \mathbb{R}^n will be indicated by $\mathcal{L}(\mathbb{R}^n)$.

We assume the following normalisation convention concerning volume n -forms in \mathbb{R}^n

$$\int_A dx^1 \wedge \dots \wedge dx^n := \int_A dx^1 \dots dx^n . \quad (5)$$

The inner product $\mathcal{H} \times \mathcal{H} \ni (x, y) \mapsto \langle x|y \rangle \in \mathbb{C}$ in a complex Hilbert space \mathcal{H} is assumed to be linear in the *right* entry.

If $A : D(A) \rightarrow \mathcal{H}$ is an operator in the complex Hilbert space \mathcal{H} with domain given by the linear subspace $D(A) \subset \mathcal{H}$, saying that A is **positive**, written $A \geq 0$, means $\langle \phi|A\phi \rangle \geq 0$ for all $\phi \in D(A)$. Furthermore, if $D(A) = D(B) \subset \mathcal{H}$ for a corresponding pair of operators A, B , then $A \geq B$ (also written $B \leq A$) means $A - B \geq 0$.

$\mathfrak{B}(\mathcal{H})$ throughout denotes the *unital C^* -algebra of bounded operators* [28] $A : \mathcal{H} \rightarrow \mathcal{H}$, where \mathcal{H} is a complex Hilbert space.

Let $\mathcal{M}(X)$ be a σ -algebra on X and \mathcal{H} a complex Hilbert space. A **\mathcal{H} -POVM** (Positive-Operator Valued Measure) on $\mathcal{M}(X)$ [7] is a map

$$A : \mathcal{M}(X) \ni \Delta \rightarrow A(\Delta) \in \mathfrak{B}(\mathcal{H})$$

such that $0 \leq A(\Delta) \leq I$ and $\Sigma(X) \ni \Delta \mapsto \langle \psi|A(\Delta)\phi \rangle \in \mathbb{C}$ is a σ -additive complex measure (with finite total variation) for every $\psi, \phi \in \mathcal{H}$. It turns out that a POVM is also σ -additive in the strong operator topology. A is **normalized** if $A(X) = I$.

A normalized \mathcal{H} -POVM A on $\mathcal{M}(X)$ is a **\mathcal{H} -PVM** (Projector Valued Measure) on $\mathcal{M}(X)$ if $A(\Delta)^2 = A(\Delta)$ (i.e., $A(\Delta)$ is an *orthogonal projector* in \mathcal{H}) for every $\Delta \in \mathcal{M}(X)$. We shall use standard notions and constructions of operator theory and spectral theory (see, e.g., [28]).

An \mathcal{H} -POVM A on $\mathcal{M}(X)$ is **absolutely continuous** with respect to a positive measure $\mu : \mathcal{M}(X) \rightarrow \overline{\mathbb{R}_+}$, written $A \ll \mu$, if $\langle \psi|A(E)\psi \rangle = 0$ for every $\psi \in \mathcal{H}$ when $\mu(E) = 0$, for $E \in \mathcal{M}(X)$.

2 Causal structures of spacetimes, Minkowski spacetime

This section is devoted to quickly introduce the basic causal structures we shall use in the rest of the paper (see, e.g., [31]). The reader who is already familiar with these notions and wants to achieve the main results soon, may temporarily skip this part, coming back to it when necessary, or for understanding a specific notation or a definition.

2.1 Globally hyperbolic spacetimes and their causal structure

A *spacetime* is the most general scenario where formulating any macroscopic physical theory according with standard notions of causality (see, e.g., [31]). It is a continuous set of *events*, a smooth manifold, equipped with geometric structures which account for causal relations. The crucial notion underpinning these notions is the Lorentzian metric tensor field on the spacetime.

Definition 6. A n -dimensional **spacetime** (M, g) is a connected n -dimensional ($n \geq 2$) smooth manifold M equipped with a **Lorentzian metric** tensor field g . It is a smooth assignment to every $T_p^*M \otimes T_p^*M$ of nondegenerate symmetric tensors g_p with constant signature $-1, +1, \dots, +1$. If $p \in M$, $v \in T_pM \setminus \{0\}$ is **spacelike**, **timelike**, **lightlike** if respectively $g_p(v, v) > 0$, $g_p(v, v) < 0$, $g_p(v, v) = 0$. **Causal vectors** are both timelike and lightlike. The zero vector $0 \in T_pM$ is spacelike *per definition*. ■

We adopt the same terminology for smooth curves $(a, b) \ni s \mapsto \gamma(s) \in M$: they are **space-like**, **timelike**, **lightlike**, **causal** according to the character of their tangent vector $\dot{\gamma}(s)$ *supposed to be uniform along γ* .

Co-vectors in T_p^*M are classified as **spacelike**, **timelike**, **lightlike**, **causal**, according to the associated elements of T_pM through the natural isomorphism $T_pM \ni v \mapsto g_p(v, \cdot) \in T_p^*M$. The indefinite inner product induced on T_p^*M by that isomorphism will be denoted by $g_p^\#$.

Definition 7. In a n -dimensional spacetime (M, g) , an m -dimensional ($m \leq n$) embedded smooth submanifold S (possibly with boundary) is **spacelike** if the tangent vectors at every point (including the ones tangent to the boundary if any) are spacelike. If $m = n - 1$ and S has no boundary, that is equivalent to saying that the co-normal vector to S is *timelike* everywhere. ■

To go on, observe that the set of timelike vectors in T_pM is made of two disjoint open cones V_p and V'_p .

Definition 8. A spacetime (M, g) is **time oriented** if there exists a continuous timelike vector field T on M . If $p \in M$,

- (a) the open cone $V_p^+ \subset T_pM$ of the pair V_p and V'_p which contains T_p is called the **future open cone at p** ; the causal vectors of $\overline{V_p^+} \setminus \{0\}$ are said to be **future-directed**;
- (b) the remaining cone V_p^- is the **past open cone at p** ; the causal vectors of $\overline{V_p^-} \setminus \{0\}$ are said to be **past-directed**.

A global continuous choice of V_p^+ for every $p \in M$ as above is a **temporal orientation** of (M, g) .

A causal co-vector is **future-directed** or **past-directed** if it is the image of a, respectively, future-directed or past-directed causal vector through $T_p M \ni v \mapsto g_p(v, \cdot) \in T_p^* M$ ■

Remark 9. (1) As M is connected, there are two possible temporal orientations or none.

- (2) If $t \in V_p^+$ and $u \in T_p M$ is causal, then u future-directed $\Leftrightarrow g_p(t, u) < 0$.
(The same result is valid for t future-directed lightlike and non-parallel to u .) ■

Definition 10. Let (M, g) be a time oriented spacetime. If $A \subset M$,

- (a) its **chronological future** $I^+(A) \subset M$ is the set of $q \in M$ such that there is a future-directed timelike smooth curve $\gamma : (a, b) \rightarrow M$ with $\gamma(t_1) \in A$ and $\gamma(t_2) = q$, for $t_1 < t_2$;
(b) its **causal future** $J^+(A) \subset M$ is the set of $q \in M$ such that either $q \in A$ or there is a future-directed causal smooth curve $\gamma : (a, b) \rightarrow M$ with $\gamma(t_1) \in A$ and $\gamma(t_2) = q$, for $t_1 < t_2$.

The **chronological past** $I^-(A)$ and the **causal past** $J^-(A)$ are defined similarly.

- (c) A is **achronal** if $A \cap I^+(A) = A \cap I^-(A) = \emptyset$,
(d) A is **acausal** if $A \cap (J^+(A) \setminus A) = A \cap (J^-(A) \setminus A) = \emptyset$. ■

Physically speaking, macroscopic physical information is transported along causal (future-directed) curves. The existence of the sets $J^\pm(p)$ (and also V_p^\pm) is the mathematical description of the finite propagation speed of physical information exiting (or entering) p .

Definition 11. Let (M, g) be a time oriented spacetime. A future-directed causal smooth curve $\gamma : (a, b) \rightarrow M$, with $a, b \in [-\infty, +\infty]$, is **future**, resp. **past**, **inextendible** if there is no $p \in M$ such that, respectively, $\gamma(s) \rightarrow p$ for $s \rightarrow b$ or $s \rightarrow a$.

γ is **inextendible** if it is both future and past inextendible. ■

We shall now focus attention on the so called *globally hyperbolic spacetimes*. This kind of spacetimes is of the utmost physical interest for many reasons, in particular because a wide family of, roughly speaking, *hyperbolic PDEs* of great physical relevance – as the Einstein equations, Klein-Gordon equations, Dirac equations – admit existence and uniqueness theorems. Cauchy data are given on special subsets called *Cauchy surfaces*. Very interestingly, the definition of globally hyperbolic spacetime and Cauchy surface is not related to PDEs, but only relies on the above geometric causal structures. *Spacelike* Cauchy surfaces are also the natural representation of instantaneous rest spaces of globally extended observers.

Definition 12. A time oriented spacetime (M, g) is said to be **globally hyperbolic** (e.g., see [3]) if it includes a **Cauchy surface**. That is a set $S \subset M$ which intersects every inextendible timelike smooth curve exactly once. ■

Remark 13. There are features of Cauchy surfaces in a n -dimensional globally hyperbolic spacetime (M, g) which deserve mention also because they are technically important for this paper.

- (a) Every Cauchy surface is in particular achronal. It is also met by every *causal* inextendible curve, but not necessarily once [31].
- (b) As it was established by Geroch, a Cauchy surface S is a closed (in M) embedded topological submanifold of co-dimension 1 of M which, in turn, is homeomorphic to $\mathbb{R} \times S$. All Cauchy surfaces are homeomorphic. [31].
- (c) If a Cauchy surface S is also a smooth submanifold of M , then its tangent vectors at each point must be either spacelike or lightlike, since S does not contain timelike curves.
- (d) As established by Bernal and Sánchez [3], every globally hyperbolic spacetime admits Cauchy surfaces which are *spacelike* smooth submanifolds of co-dimension 1 (necessarily closed in M and embedded for (b)). In turn, M is diffeomorphic to $\mathbb{R} \times S$.
The following further facts are of crucial interest to our work.

- (d1) (Proposition 5.18 in [4]) *Let S_-, S_+ be disjoint spacelike smooth Cauchy surfaces of (M, g) with $S_- \subset I^-(S_+)$. If $S \subset I^+(S_-) \cap I^-(S_+)$ is a closed (in M) connected spacelike smooth embedded 1-codimensional submanifold of M , then S is a Cauchy surface of (M, g) as well.*
- (d2) (Theorem 1.1 and Remark 4.14 of [4]) *If $K \subset M$ is a compact spacelike acausal smooth p -dimensional ($p = 1, 2, \dots, n - 1$) submanifold with boundary of (M, g) not necessarily connected, then there is a spacelike smooth Cauchy surface $S \supset K$.*
- (e) A smooth *spacelike* Cauchy surface S meets exactly once also every inextendible causal curve [3]. In particular, S is therefore acausal. ■

Remark 14. \mathcal{C}_M will henceforth denote the family of all smooth Cauchy surfaces of (M, g) . ■

2.2 Minkowski spacetime geometric and causal structures

Definition 15. Minkowski spacetime $(\mathbb{M}, \mathbb{V}, \mathbf{g})$ is a 4-dimensional real affine space whose space of translation – denoted by \mathbb{V} – is equipped with a Lorentzian inner product with signature⁷ $(-1, +1, +1, +1)$ indicated by $\mathbf{g} : \mathbb{V} \times \mathbb{V} \ni (k, p) \mapsto \mathbf{g}(k, p) =: k \cdot p \in \mathbb{R}$. ■

This definition is consistent with Definition 8 because \mathbb{M} is automatically a smooth 4-dimensional manifold with respect to the natural smooth structure induced by its affine structure. It is defined by requiring that, upon the choice of an origin $o \in \mathbb{M}$, the bijective map $I_p^o : \mathbb{M} \ni p \mapsto p - o \in \mathbb{V}$ is a diffeomorphism. $\iota_p := dI_p^o : T_p\mathbb{M} \rightarrow \mathbb{V}$ is a natural isomorphism independent of the choice of o . \mathbf{g} can therefore be exported to each tangent space $T_p\mathbb{M}$ as $g_p(u, v) := \mathbf{g}(\iota_p u, \iota_p v)$. The

⁷Differently from the choice of [8, 9]

construction defines a smooth manifold diffeomorphic to \mathbb{R}^4 endowed with a Lorentzian metric tensor field – called **Minkowski metric** – still denoted by \mathbf{g} . In the rest of the paper, for $u, v \in T_p\mathbb{M}$, we use the notation $\mathbf{g}(u, v) := u \cdot v := g_p(u, v)$.

Minkowski spacetime (\mathbb{M}, \mathbf{g}) is assumed to be *time oriented* and $dt_p(V_p^+) =: V_+ \subset V$ (which does not depend on p) is the **future open cone** in V by definition.

The **orthochronous Poincaré group** \mathcal{P}_+ is the group of time-orientation preserving isometries of (\mathbb{M}, \mathbf{g}) . \mathcal{P}_+ turns out to be the semidirect product $V \rtimes \mathcal{L}_+$ of V (the abelian group of displacements of \mathbb{M}) and the **orthochronous Lorentz group** \mathcal{L}_+ consisting of the V_+ -preserving linear isometries of (V, \mathbf{g}) . Upon the choice of an origin $o \in \mathbb{M}$ and referring to the map I_p^o above, the action of $(v, \Lambda) \in \mathcal{P}_+$ is

$$(v, \Lambda) : \mathbb{M} \ni p \mapsto o + v + \Lambda(p - o) \in \mathbb{M}. \quad (6)$$

This action also defines the structure of semidirect product $\mathcal{P}_+ = V \rtimes \mathcal{L}_+$.

A **Minkowski reference frame**, physically corresponding to an *inertial observer*, is defined by a future-directed timelike *unit* vector n . The set of these unit vectors will be denoted by $T_+ \subset V_+$. Take $o \in \mathbb{M}$ and a **Minkowski basis**, i.e., $e_0, e_1, e_2, e_3 \in V$ such that $e_0 \in T_+$ and $\mathbf{g}(e_\mu, e_\nu) = \eta_{\mu\nu}$, where $[\eta_{\mu\nu}]_{\mu, \nu=0,1,2,3} := \text{diag}(-1, 1, 1, 1)$. The global (bijective Cartesian) chart $\mathbb{M} \ni p \mapsto (x^0(p), x^1(p), x^2(p), x^3(p)) \in \mathbb{R}^4$ such that $p = o + \sum_{\mu=0}^3 x^\mu(p) e_\mu$ is a **Minkowski chart** on \mathbb{M} (with **origin** o and **axes** e_0, e_1, e_2, e_3) by definition. The vectors $\partial_{x^\mu}|_p \in T_p\mathbb{M}$ of the local bases associated to the coordinates are mapped to e_μ by ι_p and it holds both $\mathbf{g}(\partial_{x^\mu}, \partial_{x^\nu}) = \eta_{\mu\nu}$ and $\mathbf{g}^\#(dx^\mu, dx^\nu) = \eta^{\mu\nu} := \eta_{\mu\nu}$ constantly everywhere. These identities imply in particular that the metric tensor field \mathbf{g} is *globally flat*. A Minkowski chart x^0, x^1, x^2, x^3 is **adapted** to a Minkowski reference frame $n \in T_+$ if $\partial_{x^0} = n$ everywhere (where again the natural isomorphism ι_p is understood). A **rest space** Σ_n of $n \in T_+$ is any 3-dimensional plane orthogonal to n . Rest spaces are smooth spacelike 3-dimensional submanifolds. The surjective coordinate function $x^0 : \mathbb{M} \rightarrow \mathbb{R}$ of a Minkowski chart adapted to n defines a **global time coordinate** of n . The possible global time coordinates of n are defined up to an arbitrary additive constant. If a Minkowski chart x^0, x^1, x^2, x^3 is adapted to n , the **time slices**

$$\mathbb{R}_{x_0^0}^3 := \{(x_0^0, x^1, x^2, x^3) \mid (x^1, x^2, x^3) \in \mathbb{R}^3\}$$

at constant time $x^0 = x_0^0$ are the coordinate representation of the rest spaces Σ_{n, x_0^0} of n (adopting the notation used in [29]). The rest spaces of n are, in fact, bijectively labelled by the values of x^0 itself and the remaining coordinates x^1, x^2, x^3 define an admissible global chart on each submanifold Σ_{n, x_0^0} . This chart is an orthogonal Cartesian coordinate system on Σ_{n, x_0^0} with respect to the (Euclidean) metric induced by \mathbf{g} on Σ_{n, x_0^0} and the affine structure induced by the one of \mathbb{M} . So that, for instance, the *Lebesgue σ -algebra* $\mathcal{L}(\Sigma_{n, x_0^0})$ and the *Lebesgue measure* are univocally defined on each Σ_{n, x_0^0} independently of the Minkowski chart adapted to n . The Lebesgue measure on Σ_{n, x_0^0} coincides with the completion of the Borel measure canonically induced by \mathbf{g} on its embedded submanifolds of \mathbb{M} (see (10) below).

Given Minkowski chart x^0, x^1, x^2, x^3 , the **spatial components** of $k \in T_p \equiv V$ along the local basis $\partial_{x^0}|_p, \partial_{x^1}|_p, \partial_{x^2}|_p, \partial_{x^3}|_p \in T_p\mathbb{M}$ (i.e., the Minkowski basis $e_0, e_1, e_2, e_3 \in V$ associated to the

said Minkowski chart) are $\vec{k} := (k^1, k^2, k^3)$ and the **temporal component** is k^0 . Referring to the spatial components of $k, p \in \mathbb{V}$, their inner product in \mathbb{R}^3 is again denoted by the dot $\vec{k} \cdot \vec{h}$.

An important feature of Minkowski spacetime \mathbb{M} is that the structure of $J^\pm(A)$ and $I^\pm(A)$ simplify because \mathbb{M} is convex. The following result is true, whose elementary proof is left to the reader.

Proposition 16. *If $A \subset \mathbb{M}$, the following holds.*

- (a) $I^+(A)$ is made of the points $q \in \mathbb{M}$ such that there is $p \in A$ for that the $q - p \in \mathbb{V}_+$.
- (b) $J^+(A)$ is made of the points $q \in \mathbb{M}$ such that there is $p \in A$ for that $q - p \in \overline{\mathbb{V}_+}$ (so $p = q$ is admitted).

Analogous facts are true for $J^-(A)$ and $I^-(A)$.

Proposition 17 ([31]). *Minkowski spacetime is globally hyperbolic since the rest spaces of every Minkowski reference frame are (spacelike) Cauchy surfaces.*

Definition 18. A **spacelike Cauchy surface** in \mathbb{M} is a spacelike *smooth* Cauchy surface of (\mathbb{M}, \mathbf{g}) . Their family is denoted by $\mathcal{C}_\mathbb{M}^s$.

A **spacelike flat Cauchy surfaces** in \mathbb{M} is a rest space of any Minkowski reference frame. Their family will be denoted by $\mathcal{C}_\mathbb{M}^{sf}$. ■

Remark 19. In the following, S will denote a smooth Cauchy surface of \mathbb{M} , i.e., an element of $\mathcal{C}_\mathbb{M}$. However, *in case S is flat and spacelike*, i.e. $S \in \mathcal{C}_\mathbb{M}^{sf}$, we shall very often use the symbol Σ in place of S , especially when viewing them as 3-rest spaces of inertial observers according to the notation of [29]. ■

In addition to spacelike flat Cauchy surfaces, there are many other types of smooth Cauchy surfaces in \mathbb{M} . The following proposition concern their description.

Proposition 20. *Consider a given Minkowski chart x^0, x^1, x^2, x^3 and $S \in \mathcal{C}_\mathbb{M}$.*

- (a) S is determined by a smooth map $x^0 = t_S(\vec{x})$, $\vec{x} \in \mathbb{R}^3$ with

$$|\nabla t_S(\vec{x})| \leq 1 \quad \text{for every } \vec{x} \in \mathbb{R}^3. \quad \text{Here, } < \text{ replaces } \leq \text{ if } S \text{ is spacelike.} \quad (7)$$

- (b) S is diffeomorphic to \mathbb{R}^3 as the coordinates $(x^1, x^2, x^3) \in \mathbb{R}^3$ define a global chart on it.

Proof. See Appendix A. □

Let us now focus on the case where $S \in \mathcal{C}_\mathbb{M}^s$, i.e., the smooth Cauchy surface S is spacelike.

- (1) The future-directed unit normal vector and co-vector to S at $(t_S(\vec{x}), \vec{x})$ are respectively

$$n_S(\vec{x}) := \frac{\partial_{x^0} + \sum_{k=1}^3 \frac{\partial t_S}{\partial x^k} \partial_{x^k}}{\sqrt{1 - |\nabla t_S(\vec{x})|^2}}, \quad \mathbf{g}(n_S, \cdot)(\vec{x}) := \frac{-dx^0 + \sum_{k=1}^3 \frac{\partial t_S}{\partial x^k} dx^k}{\sqrt{1 - |\nabla t_S(\vec{x})|^2}}. \quad (8)$$

- (2) The Riemannian metric h^S induced on S and represented in terms of the local coordinates $\vec{x} \in \mathbb{R}^3$ is the pullback of g through the embedding function. It reads

$$h_{ab}^S := \sum_{\alpha,\beta=0}^3 \eta_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^a} \frac{\partial x^\beta}{\partial x^b} = \delta_{ab} - \frac{\partial t_S}{\partial x^a} \frac{\partial t_S}{\partial x^b} \quad (9)$$

where $a, b = 1, 2, 3$ and $x^0 = t_S(\vec{x})$.

- (c) Correspondingly, the Borel measure on S induced by the metric is equivalent to the Lebesgue measure d^3x on \mathbb{R}^3 (restricted to the Borel sets) and reads⁸

$$\nu_S(A) := \int_A \sqrt{\det h^S} d^3x = \int_A \sqrt{1 - |\nabla t_S(\vec{x})|^2} d^3x \quad \text{for every } A \in \mathcal{B}(S). \quad (10)$$

In case the smooth Cauchy surface S is generic, we can still define *non vanishing, causal* and *future-directed* normal vectors and co-vectors to S at $(t_S(\vec{x}), \vec{x})$ but, generally speaking, we cannot normalize them

$$\nu_S(\vec{x}) := \partial_{x^0} + \sum_{k=1}^3 \frac{\partial t_S}{\partial x^k} \partial_{x^k}, \quad \mathbf{g}(\nu_S, \cdot)(\vec{x}) := -dx^0 + \sum_{k=1}^3 \frac{\partial t_S}{\partial x^k} dx^k. \quad (11)$$

The induced metric turns out to be degenerate where these vectors are lightlike.

3 Spacelike Cauchy localization observables in \mathbb{M} and a general causality condition

We pass to present our generalized notion of spatial localization and a corresponding generalized causality condition. The latter extends CC stated in Definition 2. The general notion of spatial localization will be given in terms of families of POVMs on spacelike Cauchy surfaces. This notion extends the analogous Definition 1, where only flat Cauchy surfaces were considered.

3.1 A general notion of spatial localization in terms of POVMs

The POVMs we shall use will be defined on a *completion* of the Borel σ -algebra on $S \in \mathcal{C}_{\mathbb{M}}^s$. This is necessary because the *region of influence* $\Delta' \subset S'$ of a set $\Delta \in \mathcal{B}(S)$ is not necessarily in $\mathcal{B}(S')$, but it necessarily stays in the $\nu_{S'}$ completion of $\mathcal{B}(S')$ (see below). This fact was already true [8, 9, 29] when dealing with flat Cauchy surfaces Σ , where the said completion of $\mathcal{B}(\Sigma)$ was nothing but the Lebesgue σ -algebra. If $S \in \mathcal{C}_{\mathbb{M}}^s$, we denote by

$$\mathcal{M}(S) := \overline{\mathcal{B}(S)}^{\nu_S}, \quad \overline{\nu_S} \quad (12)$$

the completion σ -algebra of $\mathcal{B}(S)$ and the completion measure of ν_S with respect to the positive Borel measure ν_S (10) induced on S by the spacetime metric.

⁸We take advantage of $\det(I + cd^t) = 1 + d^t c$ for $c, d \in \mathbb{R}^n$ and I the identity of $M(n, \mathbb{R})$.

Definition 21. Consider a quantum system described in the Hilbert space \mathcal{H} . A **spacelike Cauchy localization observable** (for short **spacelike Cauchy localization**) of the system in \mathbb{M} is a family of normalized \mathcal{H} -POVMs $\mathbf{A} := \{A_S\}_{S \in \mathcal{C}_{\mathbb{M}}^s}$ where $A_S : \mathcal{M}(S) \rightarrow \mathfrak{B}(\mathcal{H})$, such that

(a) \mathbf{A} satisfies the **coherence condition**

$$A_S(\Delta) = A_{S'}(\Delta) \quad \text{if } S, S' \in \mathcal{C}_{\mathbb{M}}^s, \Delta \subset S \cap S' \text{ and } \Delta \in \mathcal{M}(S) \cap \mathcal{M}(S'); \quad (13)$$

(b) A_S is absolutely continuous with respect to $\overline{\nu_S}$, for every $S \in \mathcal{C}_{\mathbb{M}}^s$. ■

Physically speaking, requirement (b) $A_S \ll \overline{\nu_S}$ means that there is no chance to find a particle in a spatial region with zero measure. This condition will play a crucial technical role in establishing Theorem 32.

When restricting to flat Cauchy surfaces, the definition of spacelike Cauchy localization observable boils down to the definition of spatial localization observable as in Definition 1.

Remark 22. (1) Definition 21, in principle, is valid for a generic quantum system and not necessarily for a Wigner particle.

(2) A different approach [10] in defining our localization observable would concern an assignment of effects on a suitable family of acausal subsets of \mathbb{M} , without declaring that they give rise to families of POVMs on the Cauchy surfaces of $\mathcal{C}_{\mathbb{M}}^s$, but recovering this fact at a more advanced stage of the theory. This physically deeper approach would avoid to impose the coherence condition (13), since it would be encapsulated in the formalism automatically. On the other hand, this perspective would turn out technically involved when proving Theorem 32 below, in view of used mathematical technology which relies upon features of smooth spacelike Cauchy surfaces. ■

Since the examples of POVMs we shall consider later are initially defined on $\mathcal{B}(S)$, the following extension results are of relevance to our work.

Proposition 23. *Let A_S be a normalized \mathcal{H} -POVM defined on $\mathcal{B}(S)$ for $S \in \mathcal{C}_{\mathbb{M}}^s$. If $A_S \ll \nu_S$, then there exists a unique normalized \mathcal{H} -POVM \tilde{A}_S on $\mathcal{M}(S)$ which extends A_S and such that $\tilde{A}_S \ll \overline{\nu_S}$.*

Proof. See Appendix A. □

Proposition 24. *Suppose that the family of POVMs $\{A_S\}_{S \in \mathcal{C}_{\mathbb{M}}^s}$, where $A_S : \mathcal{B}(S) \rightarrow \mathfrak{B}(\mathcal{H})$, satisfies*

(a) $A_S(\Delta) = A_{S'}(\Delta)$ for every $\Delta \in \mathcal{B}(S) \cap \mathcal{B}(S')$;

(b) $A_S \ll \nu_S$.

Then, extending each POVM according to Proposition 23, we obtain a spacelike Cauchy localization observable $\{\tilde{A}_S\}_{S \in \mathcal{C}_M^s}$.

Remark 25. In the rest of the paper we shall use the same symbol A_S also for the extension to $\mathcal{M}(S)$ above denote by \tilde{A}_S . ■

In principle, it would be possible to define from scratch a notion of Cauchy localization for smooth Cauchy surfaces S which are not necessarily spacelike on the corresponding Borel σ -algebra $\mathcal{B}(S)$. As a matter of fact, we shall define such type of families of POVMs T^g in Sect.5. However Theorem 32 below needs several general results about *spacelike* smooth Cauchy surfaces. The generalisation of the results presented in this section to families of POVMs defined on generic Cauchy surfaces (also non-smooth) will be investigated elsewhere.

3.2 The general causality condition for spacelike Cauchy localization observables

We can state a natural generalization of CC.

Definition 26. If $S, S' \in \mathcal{C}_M^s$ and $\Delta \in \mathcal{M}(S)$, its **region of influence on S'** is

$$\Delta' := (J^+(\Delta) \cup J^-(\Delta)) \cap S'. \quad (14)$$

A spacelike Cauchy localization A satisfies the **general causality condition (GCC)** if, for every $S, S' \in \mathcal{C}_M^s$,

$$A_{S'}(\Delta') \geq A_S(\Delta) \quad (15)$$

when $\Delta \in \mathcal{M}(S)$ satisfies

$$\Delta' \in \mathcal{M}(S'). \quad (16)$$

■

If $S = \Sigma$ and $S' = \Sigma'$ are spacelike *flat* Cauchy surfaces of Minkowski spacetime, the completed measures $\overline{\nu_\Sigma}$ and $\overline{\nu_{\Sigma'}}$ are nothing but the Lebesgue measures on Σ and Σ' . In this situation, as already said, it turns out that [8] $\Delta' \in \mathcal{M}(\Sigma')$ whenever $\Delta \in \mathcal{M}(\Sigma)$. We do not know if this fact is general. It is however possible to prove the following fact, whose proof is inspired by analogous ideas and proofs in [10]. To this end, we observe that $S \in \mathcal{C}_M$ is a *Polish space* [12]. It can be proved in various ways, the most economic way is to use the diffeomorphism of S and \mathbb{R}^3 according to Proposition 20.

Proposition 27. Consider $S, S' \in \mathcal{C}_M^s$. If $\Delta \in \mathcal{B}(S)$, then $\Delta' \in \overline{\mathcal{B}(S')}^{\nu_{S'}} =: \mathcal{M}(S')$.

Proof. Define the following continuous function

$$\eta : S \times S' \ni (p, q) \mapsto (p, \mathbf{g}(p - q, p - q)) \in S \times \mathbb{R}.$$

The set $\Delta \times (-\infty, 0]$ is in $\mathcal{B}(S \times \mathbb{R})$ trivially. Since η is Borel measurable (as it is continuous),

$$\Delta \times \Delta' = \eta^{-1}(\Delta \times (-\infty, 0])$$

is in $\mathcal{B}(S \times S')$. At this point, we can use Proposition 8.4.4 in [12] obtaining that the projection of the above set onto S' is *universally measurable*: it stays in the σ -algebra obtained by completing $\mathcal{B}(S')$ with respect to any positive finite Borel measure on S' . To conclude we prove that Δ' must therefore belong to $\mathcal{M}(S')$. Upon the identification of S' with \mathbb{R}^3 through the global chart defined in (b) of Proposition 20, consider the Borel finite measure $\nu_n(E) = \int_E \chi_{B_n} \sqrt{1 - |\nabla t_{S'}|^2} d^3x$, where B_n is the open ball of radius $n \in \mathbb{N}$ and center the origin of \mathbb{R}^3 . $\nu_n|_{\mathcal{B}(B_n)}$ is equivalent to the \mathbb{R}^3 Lebesgue measure restricted to B_n . Since $\Delta' \cap B_n \in \overline{\mathcal{B}(S')}^{\nu_n}$, due to Lemma 61, it must be $\Delta' \cap B_n = E_n \cup N_n$ where $E_n \in \mathcal{B}(B_n) \subset \mathcal{B}(S')$ and $N_n \subset Z_n$ with $Z_n \in \mathcal{B}(B_n) \subset \mathcal{B}(S')$ and $|Z_n| = 0$, that is $\nu_{S'}(Z_n) = 0$. In summary, $\Delta' = \cup_{n \in \mathbb{N}} E_n \cup N_n = (\cup_{n \in \mathbb{N}} E_n) \cup (\cup_{n \in \mathbb{N}} N_n)$ with $\cup_{n \in \mathbb{N}} E_n \in \mathcal{B}(S')$ and $\cup_{n \in \mathbb{N}} N_n \subset \cup_{n \in \mathbb{N}} Z_n \in \mathcal{B}(S')$, where $0 \leq \nu_{S'}(\cup_{n \in \mathbb{N}} Z_n) \leq \sum_n \nu_{S'}(Z_n) = 0$, so that $\nu_{S'}(\cup_{n \in \mathbb{N}} Z_n) = 0$. By construction $\Delta' \in \overline{\mathcal{B}(S')}^{\nu_{S'}} =: \mathcal{M}(S')$. \square

3.3 The general causality condition is valid for every spacelike Cauchy localization observable

We pass to prove that GCC is actually valid for every spacelike Cauchy localization. The proof consists of some steps. The general structure of the demonstration enjoys many similarities with some proofs originally introduced in [29] and later generalized in [9]. However, to extend these constructions to generic spacelike Cauchy surfaces, we shall also take advantage of some fundamental achievements by Bernal and Sánchez [3, 4] here specialized to Cauchy surfaces of Minkowski spacetime.

Lemma 28. *Consider a Minkowski chart on \mathbb{M} and consider a pair of spacelike smooth 3-dimensional embedded manifolds S, S' respectively described as the sets*

$$S := \{(x^0 = t_S(\vec{x}), \vec{x}) \mid \vec{x} \in \mathbb{R}^3\}, \quad S' := \{(x^0 = t_{S'}(\vec{x}), \vec{x}) \mid \vec{x} \in \mathbb{R}^3\}$$

for a pair of smooth functions $t_S : \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t_{S'} : \mathbb{R}^3 \rightarrow \mathbb{R}$ (where necessarily $|\nabla t_S| < 1$ and $|\nabla t_{S'}| < 1$ everywhere). If $S \in \mathcal{C}_{\mathbb{M}}^s$ and $t_{S'}(\vec{x}) = t_S(\vec{x})$ outside a compact set in \mathbb{R}^3 , then $S' \in \mathcal{C}_{\mathbb{M}}^s$ as well.

Proof. S' is diffeomorphic to \mathbb{R}^3 through the projection map $S' \ni (x^0, t_{S'}(\vec{x})) \mapsto \vec{x} \in \mathbb{R}^3$. Therefore S' is connected and closed as it is the preimage of the closed set $\{0\}$ according to the continuous (actually smooth) map $f : \mathbb{R}^4 \ni (x^0, \vec{x}) \mapsto x^0 - t_{S'}(\vec{x}) \in \mathbb{R}$. Let us define $T = \max\{|t_S(\vec{x}) - t_{S'}(\vec{x})| \mid \vec{x} \in \mathbb{R}^3\}$. Notice that T is finite because the function is continuous and compactly supported. Since the curves tangent to the x^0 axis are timelike, we have that $S' \subset I^+(S_{-2T}) \cap I^-(S_{2T})$, where

$$S_\tau := \{(x^0 = \tau + t_S(\vec{x}), \vec{x}) \mid \vec{x} \in \mathbb{R}^3\}.$$

Similarly, $S_{-2T} \subset I^-(S_{2T})$ and also $S_{\pm 2T}$ are spacelike Cauchy surfaces because they are obtained through isometries out of the spacelike Cauchy surface S . According to (d1) in Remark 13, S' is a spacelike Cauchy surface. \square

Proposition 29. *If A is a spacelike Cauchy localization, then (16) and (15) are true when $\Delta \subset S \in \mathcal{C}_{\mathbb{M}}$ is a spacelike compact smooth 3-submanifold with boundary.*

Proof. We henceforth choose a Minkowski global chart and we describe S and S' through the global chart as in (b) of Proposition 20. In particular $\psi' : S' \ni p \mapsto \vec{x}(p) \in \mathbb{R}^3$ will be denote the said global chart on S' .

As Δ is compact, it belongs to $\mathcal{M}(S)$. Furthermore $\Delta' = (J^+(\Delta) \cup J^-(\Delta)) \cap S'$ is compact as well (Corollary A.5.4 [1]), so that it belongs to $\mathcal{M}(S')$. Next we consider a sequence of open sets $F'_n \subset S'$ such that (a) $\partial F'_n \cap \Delta' = \emptyset$, (b) $\overline{F'_{n+1}} \subset F'_n$, and (c) $\bigcap_{n \in \mathbb{N}} F'_n = \Delta'$. This sequence is recursively constructed by extracting a finite subcovering of a covering of Δ' made of coordinate open balls $\psi'^{-1}(B_{r_n}(\psi'(p_i)))$ of radius r_n , $p_i \in \Delta'$, where $r_n \rightarrow 0$ for $n \rightarrow +\infty$, and $r_{n+1} < \text{dist}(\Delta', \partial F'_n)$, where $\text{dist}(A, B) := \inf_{(x,y) \in A \times B} \text{dist}(\psi'(x), \psi'(y))$ for $A, B \subset S'$. The latter distance being the standard distance of couples of points in \mathbb{R}^3 (notice that $\text{dist}(\Delta', \partial F'_n)$ is strictly positive and finite by construction). The boundary of each $F'_n \subset S'$ is a C^0 submanifold which is also smooth up to a zero-measure subset of the boundary of the balls made of part of the intersections of the boundaries of a finite number of open balls. Each $\partial F'_n$ can be therefore slightly locally smoothed in order to transform each F'_n to a corresponding open set R'_n , such that $\overline{R'_n}$ is a compact smooth submanifold with boundary of the spacelike Cauchy surface S' and, as before, $\Delta' \cap \partial R'_n = \emptyset$, (b) $\overline{R'_{n+1}} \subset R'_n$, and (c) $\bigcap_{n \in \mathbb{N}} R'_n = \Delta'$. Let us now focus on the further family of open relatively compact sets

$$dR'_n := R'_n \setminus \overline{R'_{n+1}}.$$

$\overline{dR'_n}$ is a compact smooth submanifold with boundary of S' and $\overline{dR'_n}$ has no intesection with $J^+(\Delta) \cup J^-(\Delta)$ by construction. Since Δ is a smooth submanifold with boundary of the spacelike Cauchy surface S , we conclude that $\Delta \cup \overline{dR'_n}$ is a (non connected) spacelike and acausal compact smooth 3-submanifold with boundary of \mathbb{M} . According to (d2) in Remark 13, there is a spacelike Cauchy surface S''_n which includes $\Delta \cup \overline{dR'_n}$. Finally consider the set

$$S_n := \{(x^0 = t_{S_n}(\vec{x}), \vec{x}) \mid \vec{x} \in \mathbb{R}^3\},$$

where the map $t_{S_n} : \mathbb{R}^3 \rightarrow \mathbb{M}$ is constructed as follows:

$$t_{S_n}(\vec{x}) := \begin{cases} t_{S''_n}(\vec{x}) & \text{if } \vec{x} \in \psi'(R'_n), \\ t_{S'}(\vec{x}) & \text{if } \vec{x} \notin \psi'(R'_n). \end{cases}$$

This map is smooth by construction and $|\nabla t_{S_n}(\vec{x})| < 1$ everywhere, in particular $dx^0 - dt_{S_n} \neq 0$ everywhere, so that the S_n of t_{S_n} is a spacelike smooth submanifold. This is a spacelike compact deformation of the Cauchy surface S' as it coincides to it outside the compact $\overline{R'_n}$. According to Lemma 28, S_n is a spacelike Cauchy surface as well.

To go on, we pass to consider the two normalized \mathcal{H} -POVM A_S and $A_{S'}$. We can decompose

$$I = A_{S_n}(S_n) = A_{S_n}(\Delta) + A_{S_n}(\psi''^{-1} \circ \psi'(R'_n) \setminus \Delta) + A_{S_n}(S_n \setminus \psi''^{-1} \circ \psi'(R'_n))$$

where $\psi''_n : S''_n \rightarrow \mathbb{R}^3$ is the global chart on S''_n according to (b) of Proposition 20. Similarly,

$$I = A_{S'}(S') = A_{S'}(R'_n) + A_{S'}(S' \setminus R'_n).$$

We know that $S_n \setminus \psi''_n{}^{-1} \circ \psi'(R'_n) = S' \setminus R'_n$ because S' and S_n coincide thereon. As a consequence of the coherence condition (13), $A_{S'}(S' \setminus R'_n) = A_{S_n}(S_n \setminus \psi''_n{}^{-1} \circ \psi'(R'_n))$ so that,

$$A_{S_n}(\Delta) + A_{S_n}(\psi''_n{}^{-1} \circ \psi'(R'_n) \setminus \Delta) = A_{S'}(R'_n)$$

Using again the coherence condition with $\Delta \subset S \cap S_n$, we have obtained that

$$A_S(\Delta) + A_{S_n}(\psi''_n{}^{-1} \circ \psi'(R_n \setminus \Delta)) = A_{S'}(R'_n)$$

so that, for every $\psi \in \mathcal{H}$, we achieve the relation between positive finite measures

$$\langle \psi | A_S(\Delta) \psi \rangle + \langle \psi | A_{S_n}(\psi''_n{}^{-1} \circ \psi'(R'_n \setminus \Delta)) \psi \rangle = \langle \psi | A_{S'}(R'_n) \psi \rangle.$$

At this juncture, $\langle \psi | A_{S'}(R'_n) \psi \rangle < +\infty$, $\bigcap_{n \in \mathbb{N}} R'_n = \Delta'$, $R'_{n+1} \subset R'_n$, and external continuity yield

$$\langle \psi | A_S(\Delta) \psi \rangle + \inf_{n \in \mathbb{N}} \langle \psi | A_{S_n}(\psi''_n{}^{-1} \circ \psi'(R'_n \setminus \Delta)) \psi \rangle = \inf_{n \in \mathbb{N}} \langle \psi | A_{S'}(R'_n) \psi \rangle = \langle \psi | A_{S'}(\Delta') \psi \rangle.$$

As $\inf_{n \in \mathbb{N}} \langle \psi | A_{S_n}(\psi''_n{}^{-1} \circ \psi'(R'_n \setminus \Delta)) \psi \rangle \geq 0$, we have proved that

$$\langle \psi | A_S(\Delta) \psi \rangle \leq \langle \psi | A_{S'}(\Delta') \psi \rangle.$$

This is the thesis by exploiting a standard polarization argument and arbitrariness of ψ in a complex Hilbert space. \square

Proposition 30. *If A is a spacelike Cauchy localization, then (16) and (15) are true when $\Delta \subset S \in \mathcal{C}_M^s$ is open.*

Proof. Consider $S, S' \in \mathcal{C}_M^s$ and refer to the definition (14) of Δ' . Generally speaking, if $\Delta \subset S$ is open set, from Prop. 16 and the fact that S is a smooth submanifold (a C^0 embedded submanifold is enough actually), it is not difficult to prove that $J^\pm(\Delta)$ are open sets⁹ in \mathbb{M} . Hence $\Delta' \subset S'$ is open as well. Therefore (16) is true in the considered case.

Let us denote by $\psi : S \rightarrow \mathbb{R}^3$ the global chart on S constructed through a function t_S in a Minkowski chart x^0, \vec{x} as prescribed in (b) of Proposition 20. We shall use this identification to see the points of S as points of \mathbb{R}^3 . This identification enjoys an important property. The *completed* measures $\bar{\nu}_S$ and the standard Lebesgue measure on \mathbb{R}^3 turn out to be *equivalent* through the identification ψ as a straightforward consequence of the fact that, in (10), the map $\mathbb{R}^3 \ni \vec{x} \mapsto \sqrt{1 - |\nabla t_S(\vec{x})|^2}$ is continuous and strictly positive. The impact of this remark relies upon the following known result (Theorem 1.26 in [14]). If $A \subset \mathbb{R}^n$ is open, $\delta > 0$, there exist a

⁹This is generally false in other spacetimes!

countable collection $\{B_j\}_{j=1,2,\dots}$ of disjoint (non-empty) closed balls $B_j \subset A$ with diameter less than δ , such that

$$\left| A \setminus \bigcup_{j \in \mathbb{N}} B_j \right| = 0,$$

the bar denoting the Lebesgue measure on \mathbb{R}^3 . Passing to S , it means that (for every $\delta > 0$) there exist a countable family of pairwise disjoint compact smooth spacelike submanifolds with boundary $\Delta_j := \psi^{-1}(B_j) \subset \Delta \subset S$, such that

$$\bar{\nu}_S \left(\Delta \setminus \bigcup_{j \in \mathbb{N}} \Delta_j \right) = 0.$$

Therefore, since the positive measure $\langle \psi | \mathbf{A}_S(\cdot) \psi \rangle$ is finite and is absolutely continuous with respect to ν_S according to Prop. 23:

$$\langle \psi | \mathbf{A}_S(\Delta) \psi \rangle = \left\langle \psi \left| \mathbf{A}_S \left(\bigcup_{j \in \mathbb{N}} \Delta_j \right) \psi \right. \right\rangle, \quad (17)$$

if $\psi \in \mathcal{H}$. If we define the compact set $\Delta_N := \bigcup_{j=0}^N \Delta_j \subset \Delta$ and Δ'_N correspondingly, we can apply Proposition 29 obtaining that, for every $\psi \in \mathcal{H}$,

$$\langle \psi | \mathbf{A}_{S'}(\Delta'_N) \psi \rangle \geq \langle \psi | \mathbf{A}_{S'}(\Delta_N) \psi \rangle. \quad (18)$$

As both sequences are non-decreasing, the limit for $N \rightarrow +\infty$ exist and, using inner continuity, (17), and monotony, we find

$$\langle \psi | \mathbf{A}_{S'}(\Delta') \psi \rangle \geq \lim_{N \rightarrow +\infty} \langle \psi | \mathbf{A}_{S'}(\Delta'_N) \psi \rangle \geq \lim_{N \rightarrow +\infty} \langle \psi | \mathbf{A}_{S'}(\Delta_N) \psi \rangle = \langle \psi | \mathbf{A}_{S'}(\Delta) \psi \rangle, \quad \forall \psi \in \mathcal{H}.$$

This is the thesis by exploiting a standard polarization argument and arbitrariness of $\psi \in \mathcal{H}$ since \mathcal{H} is complex. \square

Proposition 31. *If \mathbf{A} is a spacelike Cauchy localization, then (16) and (15) are true when $\Delta \subset S \in \mathcal{C}_{\mathbb{M}}$ is compact.*

Proof. (16) is valid just because, as already observed in the proof of Proposition 29, Δ' is compact if Δ is. So we have to establish the validity of (15) only. We refer to a global chart $\psi : S \rightarrow \mathbb{R}^3$ on S constructed out of the map t_S in a Minkowski chart x^0, \vec{x} as in (b) of Proposition 20. If $\Delta \subset S$ is compact we can construct a sequence of open sets $F_n \supset \Delta$ such that $F_{n+1} \subset F_n$ and $\bigcap_{n \in \mathbb{N}} F_n = \Delta$. F_n is the finite union of coordinate balls of radius $r_n \rightarrow 0$ such that $r_{n+1} < \text{dist}(\Delta, F_n)$ where we adopted the same notation as in the proof of Prop. 29. In view of Prop 31, taking account of the external continuity of the involved measures

$$\langle \psi | \mathbf{A}_S(\Delta) \psi \rangle = \lim_{n \rightarrow +\infty} \langle \psi | \mathbf{A}_S(F_n) \psi \rangle \leq \lim_{n \rightarrow +\infty} \langle \psi | \mathbf{A}_{S'}(F'_n) \psi \rangle = \langle \psi | \mathbf{A}_{S'}(\bigcap_{n \in \mathbb{N}} F'_n) \psi \rangle.$$

The proof ends if proving that $\bigcap_{n \in \mathbb{N}} F'_n = \Delta'$. In, fact, obviously $\bigcap_{n \in \mathbb{N}} F_n \supseteq \Delta'$. On the other hand, if $p \in \bigcap_{n \in \mathbb{N}} F'_n$ then there is a causal segment from p to $q_n \in F_n \subset \overline{F_0}$ which is compact and stays in S because it is closed. As a consequence there is a subsequence $q_{n_k} \rightarrow q \in \overline{F_0}$ for $k \rightarrow +\infty$. It is easy to see that $q \in \Delta$ (otherwise q would stay at some distance from the compact Δ and this is not admitted in view of the very construction of the sets $F_{n_k} \ni q_{n_k} \rightarrow q$). The limit $p - q$ of the causal segments $p - q_n$ is still causal or 0 (as the set of causal vectors and $\{0\}$ is closed). We conclude that $p \in (J^+(\Delta) \cup J^-(\Delta)) \cap S = \Delta'$. We have established that $\bigcap_{n \in \mathbb{N}} F_n \subset \Delta'$, so that $\bigcap_{n \in \mathbb{N}} F_n = \Delta'$ concluding the proof. \square

We are in a position to prove the first main result of this work: spacelike Cauchy localizations always satisfy the Cauchy causality requirement in Definition 26. Thus, in particular, also Castrigiano's causal requirement when restricting to spacelike flat Cauchy surfaces.

Theorem 32. *Let $\mathbf{A} := \{\mathbf{A}_S\}_{S \in \mathcal{C}_M^s}$ be a spacelike Cauchy localization observable, then it satisfies the general causal condition in Definition 26.*

More generally, if $S, S' \in \mathcal{C}_M^s$, $\Delta \in \mathcal{M}(S)$, and $\psi \in \mathcal{H}$, then

$$\langle \psi | \mathbf{A}_S(\Delta) \psi \rangle \leq \sup \{ \langle \psi | \mathbf{A}_{S'}(K') \psi \rangle \mid K \subset \Delta, K \text{ compact} \}, \quad (19)$$

even if $\Delta' \notin \mathcal{M}(S')$. (Above $K' := (J^+(K) \cup J^-(K)) \cap S'$ as usual.)

Proof. The positive Borel measure $\mathcal{B}(S) \ni \Delta \mapsto \langle \psi | \mathbf{A}_S(\Delta) \psi \rangle$ is regular because S is countable union of compacts with finite measure (Theorem 2.18 in [33]). As a consequence, the completion measure $\mathcal{M}(S) \ni \Delta \mapsto \langle \psi | \mathbf{A}_S(\Delta) \psi \rangle$ is regular as well (Prop. 1.59 in [12]). If $\Delta \in \mathcal{M}(S)$, internal regularity yields

$$\langle \psi | \mathbf{A}_S(\Delta) \psi \rangle = \sup \{ \langle \psi | \mathbf{A}_S(K) \psi \rangle \mid K \subset \Delta \subset S, K \text{ compact} \}.$$

At this juncture Prop. 31 entails (19).

If we also know that $\Delta' \in \mathcal{M}(S')$, noticing that the sets K' are compact and satisfy $K' \subset \Delta'$, but they are not necessarily all compact sets in S' satisfying the latter condition, internal regularity entails

$$\sup \{ \langle \psi | \mathbf{A}_{S'}(K') \psi \rangle \mid K \subset \Delta \subset S, K \text{ compact} \} \leq \langle \psi | \mathbf{A}_{S'}(\Delta') \psi \rangle.$$

However Prop. 31 implies

$$\sup \{ \langle \psi | \mathbf{A}_S(K) \psi \rangle \mid K \subset \Delta \subset S, K \text{ compact} \} \leq \sup \{ \langle \psi | \mathbf{A}_{S'}(K') \psi \rangle \mid K \subset \Delta \subset S, K \text{ compact} \}.$$

Therefore $\langle \psi | \mathbf{A}_S(\Delta) \psi \rangle \leq \langle \psi | \mathbf{A}_{S'}(\Delta') \psi \rangle$. This is just (15) by exploiting a standard polarization argument and arbitrariness of $\psi \in \mathcal{H}$ since \mathcal{H} is complex. \square

Corollary 33. *If a spatial localization observable $\{\mathbf{A}_\Sigma\}_{\Sigma \in \mathcal{C}_M^{sf}}$ can be extended to a spacelike Cauchy localization observable $\{\mathbf{A}_S\}_{S \in \mathcal{C}_M^s}$, then the former automatically satisfies CC in Definition 2.*

Proof. Consider only spacelike flat Cauchy surfaces, noticing that there $\mathcal{M}(\Sigma) = \mathcal{L}(\Sigma)$ and that $\Delta' \in \mathcal{L}(\Sigma')$ for every $\Delta \subset \Sigma$ (even if $\Delta \notin \mathcal{B}(S)$!) as established in [8]. In that case GCC in Definition 26 boils down to CC in Definition 2. \square

Remark 34. In the proof of Theorem 32, the coherence condition (13) of Definition 21 is of utmost relevance. This is one of the reasons why we inserted that requirement already of Definition 21. However, an alternative approach would consist of removing this condition from the Definition 21 and to directly add it in the hypotheses of Theorem 32, proving again the validity of GCC. At this juncture it is worth stressing that GCC trivially implies the coherence condition 13 (since $\Delta = \Delta'$ if $\Delta \subset S \cap S'$). The conclusion is that, if using from scratch a weaker version of Definition 21, i.e., without the requirement of coherence (13), the general causal condition (15)-(16) and the coherence condition (13) would be *equivalent*¹⁰. \blacksquare

4 Massive KG particles, conserved currents

In this section, we shall introduce some basic notion and results to construct examples of spatial Cauchy localization observables. The former section is a recap about the one-particle structure of the massive real Klein-Gordon particle. The latter deals with some general properties of conserved currents and associated exact volume 3-forms.

4.1 One-particle Hilbert space of real massive KG particles in \mathbb{M}

According to Section 2.2, we fix a preferred origin $o \in \mathbb{M}$, so that the map $\mathbb{V} \ni x \mapsto o + x \in \mathbb{M}^4$ defines a one-to-one correspondence between points of \mathbb{M} and vectors in \mathbb{V} (the differential of this map being ι_p). This identification is very useful when dealing with the Fourier transformation on \mathbb{M} , where the product $p \cdot x$ enters the play. All the theory developed in this work does not depend on the choice of o . With this structure, the active action (6) of $(y, \Lambda) \in \mathcal{P}_+$ on \mathbb{M} takes the form $\mathbb{M} \ni o + x \mapsto o + y + \Lambda x \in \mathbb{M}$. Given a mass constant $m > 0$, the **future mass-shell** in \mathbb{V} is $\mathbb{V}_{m,+} := \{k \in V_+ \mid k \cdot k = -m^2\}$. For every Minkowski chart x^0, x^1, x^2, x^3 ,

$$k \in \mathbb{V}_{m,+} \quad \Leftrightarrow \quad k \in \mathbb{V} \quad \text{and} \quad k^0 = \sqrt{\vec{k}^2 + m^2} > 0. \quad (20)$$

The *one-particle Hilbert space* of a Klein-Gordon particle of mass $m > 0$ is isomorphic to $L^2(\mathbb{R}^3, d^3p)$ upon the choice of a Minkowski chart x^0, x^1, x^2, x^3 . Here \mathbb{R}^3 , is the space of momenta \vec{p} and the *momentum representation* wavefunctions $\phi = \phi(\vec{p})$ are normalized elements of $L^2(\mathbb{R}^3, d^3p)$. To have a completely covariant formulation, one takes advantage of the canonical Hilbert space isomorphism

$$F : L^2(\mathbb{V}_{m,+}, \mu_m) \ni \psi \mapsto \phi_\psi \in L^2(\mathbb{R}^3, d^3k) \quad \text{where} \quad \phi_\psi(\vec{k}) := \frac{\psi(k)}{\sqrt{k^0(\vec{k})}} \quad (21)$$

¹⁰We are grateful to D.P.L. Castrigiano for this observation.

There,

$$\mathcal{H} := L^2(\mathbb{V}_{m,+}, \mu_m) \quad \text{with} \quad d\mu_m(k) := \frac{d^3k}{k^0(\vec{k})} \quad (22)$$

is the (covariant) **one-particle Hilbert space** of a real Klein-Gordon particle with mass $m > 0$, where μ_m is the \mathcal{P}_+ -**invariant measure** on the mass shell $\mathbb{V}_{m,+}$. Covariance is here manifest because, the standard unitary representation of \mathcal{P}_+ on the one-particle space

$$\mathcal{P}_+ \ni h \mapsto U_h \in \mathfrak{B}(\mathcal{H}) \quad (23)$$

takes the *equivariant* form

$$(U_h \psi)(k) = e^{-ik \cdot y_h} \psi(\Lambda_h^{-1} k), \quad \text{for every } k \in \mathbb{V}_{m,+}, \psi \in \mathcal{H}, h = (y_h, \Lambda_h) \in \mathcal{P}_+. \quad (24)$$

For future convenience, we define the dense subspaces $\mathcal{D}(\mathcal{H}) \subset \mathcal{S}(\mathcal{H}) \subset \mathcal{H}$

$$\mathcal{D}(\mathcal{H}) := F^{-1}(C_c^\infty(\mathbb{R}^3)) \quad \text{and} \quad \mathcal{S}(\mathcal{H}) := F^{-1}(\mathcal{S}(\mathbb{R}^3)), \quad (25)$$

where $\mathcal{S}(\mathbb{R}^3)$ is the usual *Schwartz space* on \mathbb{R}^3 . It easy to prove that, in view of (24), the definition of $\mathcal{D}(\mathcal{H})$ and $\mathcal{S}(\mathcal{H})$ do not depend on the choice of the Minkowski chart used to construct F . If $\psi \in \mathcal{D}(\mathcal{H})$, or more generally $\psi \in \mathcal{S}(\mathcal{H})$, the associated (complex) **covariant wavefunction** is

$$\varphi_\psi(x) := \int_{\mathbb{V}_{m,+}} \frac{\psi(p) e^{ip \cdot x}}{(2\pi)^{3/2}} d\mu_m(p). \quad (26)$$

Notice that $\varphi_\psi \in C^\infty(\mathbb{M})$, it is also bounded with all of its derivatives, and it solves the **Klein-Gordon equation** in \mathbb{M}

$$(\square - m^2)\varphi_\psi = 0, \quad \text{where } \square := \eta^{\mu\nu} \partial_\mu \partial_\nu \text{ in every Minkowski chart.} \quad (27)$$

Furthermore, the action of U on φ_ψ is straightforward and explains the adjective "covariant":

$$\varphi_{U_h \psi}(x) = \varphi_\psi(h^{-1}x) \quad \forall h \in \mathcal{P}_+, \forall x \in \mathbb{M}. \quad (28)$$

Due to (26),(27) and Proposition 37 below applied to the bounded conserved smooth current

$$J_\mu^{\psi, \psi'} := \overline{\varphi_\psi} \partial_\mu \varphi_{\psi'} - \varphi_{\psi'} \partial_\mu \overline{\varphi_\psi},$$

we have that the scalar product of \mathcal{H} satisfies, for $\psi, \psi' \in \mathcal{S}(\mathcal{H})$,

$$\langle \psi | \psi' \rangle = \frac{i}{2} \int_S (\overline{\varphi_\psi}(x) n_S^\mu(x) \partial_\mu \varphi_{\psi'}(x) - \varphi_{\psi'}(x) n_S^\mu(x) \partial_\mu \overline{\varphi_\psi}(x)) d\nu_S(x), \quad (29)$$

for every spacelike smooth Cauchy surface S in \mathbb{M} . An analogous formula for generic smooth Cauchy surfaces can be established on account of Propositions 36 and 37 below.

Various issues concerning the possibility to directly or indirectly interpret φ_ψ as a wavefunction in classical sense somehow related to notions of spatial localization, and the failure of these expectations accumulated over the years, were discussed in [29].

4.2 Conserved quantities on generic smooth Cauchy surfaces of \mathbb{M}

From now on, we assume that \mathbb{M} is oriented. Consider a smooth vector field J in \mathbb{M} . We can associate a 3-form to it¹¹, where we henceforth take advantage of the summation convention over repeated Greek indices,

$$\omega^J(x) := \frac{1}{3!} J^\delta(x) \epsilon_{\delta\alpha\beta\gamma} dx^\alpha \wedge dx^\beta \wedge dx^\gamma. \quad (30)$$

Above $\epsilon_{\delta\alpha\beta\gamma}$ is the **Levi-Civita (pseudo)tensor** which, in Minkowski coordinates, coincides to the sign of the permutation $(\delta, \alpha, \beta, \gamma)$ of $(0, 1, 2, 3)$, or vanishes in case of repetitions.

Let S be a smooth Cauchy surface determined by the map $t_S : \mathbb{R}^3 \rightarrow \mathbb{R}$ in Minkowski coordinates x^0, x^1, x^2, x^3 where v_S is the future-directed normal vector to S as in (11). If $J \equiv (J^0, \vec{J})$ is either zero or causal and future directed depending on $x \in \mathbb{M}$, then we define

$$\int_A \omega^J := \int_A (J^0 - \vec{J} \cdot \nabla t_S) d^3x = - \int_A J \cdot v_S d^3x \in [0, +\infty], \quad \forall A \in \mathcal{B}(S) \quad (31)$$

In the last two integrals, A is interpreted as a subset of \mathbb{R}^3 according to the canonical projection $\mathbb{R}^4 \ni (x^0, \vec{x}) \mapsto \vec{x} \in \mathbb{R}^3$ and the integrals are well definite in $[0, +\infty]$ because

$$-J \cdot v_S \geq 0. \quad (32)$$

Remark 35. The integral of the form ω^J as defined by the right-most term in (31) is a standard integral in the sense of measure theory for a generic Borel set A . This fact permits to use standard arguments of measure theory. On the other hand, if A is compact, the definition agrees with the standard integral of smooth n -forms on compact sets and we can take advantage of standard results in this context as the Poincaré theorem. ■

We have a preliminary proposition whose elementary proof is left to the reader.

Proposition 36. *If J is a smooth vector field on \mathbb{M} and ω^J is defined as in (30), then the following facts are valid (where (5) has been taken into account)*

(1) J is conserved iff ω^J is closed:

$$\partial_\alpha J^\alpha(x) = 0 \quad \Leftrightarrow \quad d\omega^J(x) = 0. \quad (33)$$

(2) If $S \in \mathcal{C}_{\mathbb{M}}^s$, then

$$\int_A \omega^J = - \int_A J \cdot n_S d\nu_S. \quad (34)$$

We pass now to state a folklore statement which is actually technically very useful. The unexpectedly technical proof is in the appendix.

Proposition 37. *Let J be a smooth vector field on \mathbb{M} such that*

¹¹ ω^J can equivalently be written as the *interior product* of J and the volume 4-form of (\mathbb{M}, \mathbf{g}) .

- (1) it is **conserved**, i.e., $\partial_\alpha J^\alpha = 0$ everywhere;
- (2) if $x \in \mathbb{M}$, then either $J(x) = 0$ or $J(x)$ is causal and future-directed, i.e., $J(x) \in \overline{\mathbb{V}_+}$;
- (3) it is **bounded**, i.e., its components in a (thus every) Minkowski chart are bounded functions.

Then, if $S, S' \in \mathcal{C}_\mathbb{M}$,

$$\int_S \omega^J = \int_{S'} \omega^J \in [0, +\infty]. \quad (35)$$

If, e.g. $S' \in \mathcal{C}_\mathbb{M}^s$, then the above identity can be re-written as

$$\int_S \omega^J = - \int_{S'} J \cdot n_{S'} d\nu_{S'}. \quad (36)$$

Proof. See Appendix A □

5 Spacelike Cauchy localization observables out of causal kernels of massive KG particles

We now pass to make contact with a family of normalized POVMs, for a massive real Klein-Gordon particle, defined on spacelike flat Cauchy surfaces and introduced by Gerlach, Gromes, Petzold, and Rosenthal [18] and Henning, Wolf [20] and, much more recently, rigorously studied by Castrigiano in [9], proving in particular that these POVMs satisfy CC. Next we pass to extend these POVMs to the full family of smooth Cauchy surfaces showing that, when considering only the spacelike Cauchy surfaces, they induce corresponding spacelike Cauchy localization observables. Before, we have to recall some basic definitions.

Remark 38. We stress that the localization observable we shall construct is more general than the notion of spacelike Cauchy localization. In fact the POVMs we shall construct are also defined for smooth Cauchy surfaces which are not spacelike. This feature may have some consequences in the analysis of the structure of causal regions of \mathbb{M} in terms of orthocomplemented lattices [9]. ■

5.1 A notion of spatial localization for massive KG particles out of causal kernels

Consider a Minkowski chart x^0, x^1, x^2, x^3 . We want to introduce an $L^2(\mathbb{R}^3, d^3p)$ POVM rigorously discussed in [9] – there named POL – on the rest space at time x^0 denoted below by $\mathbb{R}_{x^0}^3$. The POVM $\mathbb{T}_{\mathbb{R}_{x^0}^3}$ defines the position in $\mathbb{R}_{x^0}^3$ of a scalar real Klein Gordon particle of mass $m > 0$ – whose pure states are defined by functions $\phi = \phi(\vec{p})$ in momentum representation with

Hilbert space $L^2(\mathbb{R}^3, d^3p)$. It satisfies the condition

$$\langle \phi | T_{\mathbb{R}_{x^0}^3}(\Delta) \phi \rangle = \frac{1}{(2\pi)^3} \int_{\Delta} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{K(\vec{k}, \vec{p})}{2\sqrt{p^0 k^0}} e^{i(\vec{p}-\vec{k}) \cdot \vec{x} - i(p^0 - k^0)x^0} \overline{\phi(\vec{k})} \phi(\vec{p}) d^3p d^3k d^3x, \Delta \in \mathcal{B}(\mathbb{R}_{x^0}^3). \quad (37)$$

Above, $\vec{x} := (x^1, x^2, x^3)$, the components k^0 and p^0 are determined by $\vec{k}, \vec{p} \in \mathbb{R}^3$ as prescribed in (20) and the vectors ϕ stay in a suitable dense subspace of $L^2(\mathbb{R}^3, d^3p)$. More specifically, referring to the discussion of Sect 11 [9]:

Definition 39. Consider a Minkowski chart x^0, x^1, x^2, x^3 on \mathbb{M} . A **POL with causal kernel**¹² on $\mathbb{R}_{x^0}^3$ is a normalized $L^2(\mathbb{R}^3, d^3p)$ -POVM $T_{\mathbb{R}_{x^0}^3}^g$ such that¹³

- (a) (37) is valid when $\phi \in L^1(\mathbb{R}^3, dk^3) \cap L^2(\mathbb{R}^3, dk^3)$;
- (b) $K := K_g$ is a **causal kernel**, i.e., it has the structure

$$K_g(\vec{k}, \vec{p}) := (k^0 + p^0)g(-k \cdot p), \quad k, p \in \mathbb{V}_{m,+} \quad (38)$$

for $g : [m^2, +\infty) \rightarrow \mathbb{R}$ which is continuous, normalized to $g(m^2) = 1$ and it is such that $K_g : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{C}$ is *positive definite*. ■

Remark 40. (1) We recall the reader that $K : X \times X \rightarrow \mathbb{C}$ is a **positive definite kernel** if

$$\sum_{i,j=1}^N \overline{c_i} c_j K(k_i, k_j) \geq 0, \quad \forall \{c_j\}_{j=1,\dots,N} \subset \mathbb{C}, \forall \{k_j\}_{j=1,\dots,N} \subset X, \forall N = 1, 2, \dots \quad (39)$$

Notice that a positive definite kernel is necessarily **Hermitian**: $K(p, q) = \overline{K(q, p)}$.

- (2) We stress that in (38), $k = (k^0(\vec{k}), \vec{k})$ and $p = (p^0(\vec{p}), \vec{p})$ in accordance with (20), so that we can see K_g either as a function on $\mathbb{R}^3 \times \mathbb{R}^3$ or on $\mathbb{V}_{m,+} \times \mathbb{V}_{m,+}$ indifferently. ■

It is possible to prove (see [9] for details) that \mathcal{P}_+ -covariant localization POVMs with causal kernel do exist. In particular, a family of functions g as in the above definition which give rise to corresponding \mathcal{P}_+ -covariant POVM with causal kernel is

$$g_r(z) := \frac{(2m^2)^r}{(m^2 + z)^r} \quad z \geq m^2, r \geq 3/2. \quad (40)$$

Finite convex combinations of functions g_r also define causal kernels. Even some pointwise limits of these convex combinations do the job as discussed in [9]:

$$g(z) = \left(\frac{(1+c)m^2}{cm^2 + z} \right)^n, \quad z \geq m^2 \quad \text{for } -1 < c \leq 1 \text{ and } \mathbb{N} \ni n > 1.$$

¹²According to [9].

¹³In [9], a further denominator $1/\sqrt{k^0 p^0}$ should take place under $k^0 + p^0$ in (38). This factor has been moved in (37) here, so that the final formulas of the POVM are actually identical. Obviously, this does not affect the positivity of that kernel.

It is convenient to re-write the POVM $\mathbb{T}_{\mathbb{R}^3_{x^0}}^g$ of Definition 39 into a different form, where (a) the dependence on a chart disappears, (b) the spacelike flat Cauchy surface normal to a reference frame remains, and (c) some covariance properties explicitly show up. A similar reformulation already appeared in [9], but we use here a slightly different approach and notation, more useful for our final goals. To this end, as in [9], we first define a current $j_g(k, p) \in \mathbb{V}_+$ by means of

$$j_g(k, p) := \frac{1}{2}(k + p)g(-k \cdot p), \quad p, k \in \mathbb{V}_{m,+}, \quad (41)$$

so that (37) can be rephrased to an equivalent form

$$\langle \phi | \mathbb{T}_{\mathbb{R}^3_{x^0}}^g(\Delta) \phi \rangle = -\frac{1}{(2\pi)^3} \int_{\Delta} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{j_g(p, k) \cdot n}{\sqrt{p^0 k^0}} e^{i(p-k) \cdot x} \overline{\phi(\vec{k})} \phi(\vec{p}) d^3 p d^3 k d\nu_{\mathbb{R}^3_{x^0}}(x), \quad \Delta \in \mathcal{B}(\mathbb{R}^3_{x^0}). \quad (42)$$

Above, the Minkowski chart x^0, x^1, x^2, x^3 is adapted to $n := \partial_{x^0}$. Furthermore, up to a spacetime displacement of the origin of the coordinates, we can always assume that $(0, 0, 0, 0)$ corresponds to the origin $o \in \mathbb{M}$ initially fixed¹⁴ (beginning of Section 5). The vector $x \in \mathbb{V}$ in the exponent in (42) is such that the point $o + x \in \mathbb{M}$ has coordinates x^0, x^1, x^2, x^3 . We might therefore write $d\mu_{\mathbb{R}^3_{x^0}}(o + x)$ rather than $d\mu_{\mathbb{R}^3_{x^0}}(x)$. However, this misuse of notation does not produce troubles because the map $\mathbb{V} \ni x \mapsto o + x \in \mathbb{M}$ is one-to-one. The factor $1/\sqrt{p^0 k^0}$ in (42) stems from the choice of the Hilbert space $L^2(\mathbb{R}^3, d^3 p)$. It can be removed by passing to covariant one-particle Hilbert space $\mathcal{H} := L^2(\mathbb{V}_{m,+}, \mu_m)$ (22). With this prescription, if $\Sigma \in \mathcal{C}_{\mathbb{M}}^{sf}$, (42) can be rephrased to

$$\begin{aligned} & \langle \psi | \mathbb{T}_{\Sigma}^g(\Delta) \psi \rangle = \\ & - \int_{\Delta} \int_{\mathbb{V}_{m,+}} \int_{\mathbb{V}_{m,+}} \frac{j_g(p, k) \cdot n_{\Sigma}}{(2\pi)^3} e^{i(p-k) \cdot x} \overline{\psi(k)} \psi(p) d\mu_m(p) \mu_m(k) d\nu_{\Sigma}(x), \quad \Delta \in \mathcal{B}(\Sigma) \end{aligned} \quad (43)$$

It is now evident that no choice of a *Minkowski chart* enters (43) and the only spot where a *Minkowski reference frame* n_{Σ} takes place is when assigning the spacelike flat Cauchy surface Σ , since n_{Σ} is the future directed unit normal vector to Σ . The vectors x entering the exponential satisfy $o + x \in S$, which, in turn, is the integration space of the external integral.

5.2 Properties of \mathbb{T}_{Σ}^g , for spacelike flat Cauchy surfaces Σ : Covariance, causality, no strict localizability, Newton-Wigner, Heisenberg inequality,

If we fix a function g as in Definition 39, the arising spatial localization observable $\{\mathbb{T}_{\Sigma}^g\}_{\Sigma \in \mathcal{C}_{\mathbb{M}}^{sf}}$ is \mathcal{P}_+ -covariant in the sense discussed in [9]: If U is the unitary representation of \mathcal{P}_+ in the one-particle space introduced in (23) and (24), it holds

$$U_h \mathbb{T}_{\Sigma}^g(\Delta) U_h^{-1} = \mathbb{T}_{h\Sigma}^g(h\Delta), \quad \forall \Sigma \in \mathcal{C}_{\mathbb{M}}^{sf}, \forall \Delta \in \mathcal{L}(\Sigma), \quad \forall h \in \mathcal{P}_+. \quad (44)$$

¹⁴Changes of the origin $o \in \mathbb{M}$ can be re-absorbed in the definition of wavefunctions through an obvious unitary transformation.

Notice that $h\Sigma$ is an analogous spacelike flat Cauchy surface normal to $n_{h\Sigma} = \Lambda_h n_\Sigma$ if $h = (y_h, \Lambda_h)$. We do not enter into the details of these properties because we shall not use them in the rest of the paper. The spatial localization observable $\{\mathbb{T}_\Sigma^g\}_{\Sigma \in \mathcal{C}_M^{sf}}$ complies with the definition of *relativistic* spatial localization observable proposed in [29] (Definition 18 therein). Denoting the POVMs extended from $\mathcal{B}(\Sigma)$ to $\mathcal{L}(\Sigma)$ with the same symbol \mathbb{T}_Σ^g , we can prove the following result.

Theorem 41 ([9]). *If $g : [m^2, +\infty) \rightarrow \mathbb{R}$ continuous, normalized to $g(m^2) = 1$ such that the kernel $K_g : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{C}$ in (38) is positive definite, then spatial localization observable $\{\mathbb{T}_\Sigma^g\}_{\Sigma \in \mathcal{C}_M^{sf}}$ satisfies CC in Definition 2.*

Proof. Theorem 56 [9]. □

Corollary 42. *If $\psi \in \mathcal{H}$, the localization probability associated to the spatial localization observable $\{\mathbb{T}_\Sigma^g\}_{\Sigma \in \mathcal{C}_M^{sf}}$ cannot be zero outside a bounded set in Σ .*

Proof. If localized states as above exist CT would fail as a consequence of the Hegerfeldt theorem 4 and thus also CC would be false. (See [29] for a discussion on this point.) □

Remark 43. In spite of this obstruction, it is possible to show [9] that probability distributions localized in bounded sets can be arbitrarily well approximated by probability distributions arising by suitable sequences of state ψ_n . ■

There is an interesting relation between the *first moment* of the POVM \mathbb{T}_Σ^g on a spacelike flat Cauchy surface Σ and the *Newton-Wigner* selfadjoint operators $N_\Sigma^1, N_\Sigma^2, N_\Sigma^3$ [29] associated to a Minkowski chart x^0, x^1, x^2, x^3 such that the slice $x^0 = 0$ coincides with Σ . Similarly to Thm 26 in [29], one sees that the following result is valid. Below $\Delta_\psi x_{\mathbb{T}_\Sigma^g}^a$ denotes the standard deviation for the coordinate x^a referred to the probability distribution $\langle \psi | \mathbb{T}_\Sigma^g(\cdot) \psi \rangle$ constructed out of the POVM \mathbb{T}_Σ^g in the state defined by the unit vector ψ .

Theorem 44. *Suppose that g in the definition of \mathbb{T}_Σ^g is real, bounded, and smooth, then the following is true.*

(a) *The a -first moment of \mathbb{T}_Σ^g is defined for every $\psi \in \mathcal{D}(\mathcal{H})$ with $\|\psi\| = 1$ and*

$$\int_\Sigma x^a \langle \psi | \mathbb{T}_\Sigma^g(d^3x) \psi \rangle = \langle \psi | N_\Sigma^a \psi \rangle, \quad \text{where } a = 1, 2, 3. \quad (45)$$

In particular, N_Σ^a is the unique selfadjoint operator in \mathcal{H} which satisfies the identity above.

(b) *The **Heisenberg inequality** turns out to be corrected as, for $\psi \in \mathcal{D}(\mathcal{H})$,*

$$\Delta_\psi x_{\mathbb{T}_\Sigma^g}^a \Delta_\psi P_a \geq \frac{\hbar}{2} \sqrt{1 + 4(\Delta_\psi P_a)^2 \langle \psi | \mathbb{K}_a^{\mathbb{T}_\Sigma^g} \psi \rangle}, \quad a = 1, 2, 3.$$

$\mathbb{K}_a^{\mathbb{T}_\Sigma^g} \in \mathfrak{B}(\mathcal{H})$ *is a selfadjoint operator, which is a (spectral) function of the four momentum observable P with the form (79), such that $\mathbb{K}_a^{\mathbb{T}_\Sigma^g} \geq 0$.*

(c) If g is of the form (40), or convex combinations of them, then (a) and (b) also hold for $\psi \in \mathcal{S}(\mathcal{H})$.

Proof. See Appendix A. □

Notice that the left-hand side of (45) does not depend on g .

Remark 45. (1) If also the identity

$$\int_{\Sigma} (x^a)^2 \langle \psi | \mathbb{T}_{\Sigma}^g(d^3x) \psi \rangle = \langle \psi | (N_{\Sigma}^a)^2 \psi \rangle \quad (\text{false!}),$$

were valid one could apply a known theorem by Naimark about the decomposition of maximally symmetric operators (here N_{Σ}^a) in terms of POVMs (see Theorem 23 in [13] and the discussion about it) obtaining $\mathbb{T}_{\Sigma}^g = \mathbb{Q}_{\Sigma}$. This is obviously false and it is also reflected by the appearance of the term $\langle \psi | \mathbb{K}_a^{\mathbb{T}_{\Sigma}^g} \psi \rangle$ in the modified Heisenberg inequality.

(2) If $U_t^{(n)}$ is the unitary time evolutor corresponding to the time evolution along n in \mathbb{M} , it is easy to see that the Heisenberg evolution $U_t^{(n)} N_{\Sigma}^a U_t^{(n)\dagger}$ of N_{Σ}^a on the right-hand side of (45) equals the integral on the left-hand side over the correspondingly temporally translated time slice Σ_t . As already observed in [29], this fact implies that the worldline $\mathbb{R} \ni t \mapsto (t, \int_{\Sigma_t} \vec{x} \langle \psi | \mathbb{T}_{\Sigma_t}^g(d^3x) \psi \rangle)$ is *timelike* (Corollary 14 in [29]) as expected by massive particles. ■

5.3 A spacelike Cauchy localization observable $\mathbb{T}^g = \{\mathbb{T}_S^g\}_{S \in \mathcal{C}_{\mathbb{M}}^s}$ for a massive KG particle

Let us focus again on the POVM $T_{g, \mathbb{R}_{x_0}^3}$ satisfying (37). The equivalent form (43) of (37) is actually already prompt to be generalized to any smooth Cauchy surface S . Heuristically, if $S \subset \mathbb{M}$ is a spacelike Cauchy surface defined by $x^0 = t_S(\vec{x})$, we expect that the current j_g also defines a normalized POVM whose expectation value on $\Delta \in \mathcal{B}(S)$ is:

$$\langle \psi | \mathbb{T}_S^g(\Delta) \psi \rangle = - \int_{\Delta} \int_{\mathcal{V}_{m,+}} \int_{\mathcal{V}_{m,+}} \frac{j_g(p, k) \cdot n_S(x)}{(2\pi)^3} e^{i(p-k) \cdot x} \overline{\psi(k)} \psi(p) d\mu_m(p) d\mu_m(k) dv_S(x), \quad \Delta \in \mathcal{B}(S) \quad (46)$$

when ψ belongs to a suitable subspace of \mathcal{H} and where n_S is the normal future-directed unit vector to S . In case S is smooth but not spacelike, we can expect the version in terms of forms be valid

$$\langle \psi | \mathbb{T}_S^g(\Delta) \psi \rangle = \int_{\Delta} \omega^{J_{g,\psi}}, \quad \Delta \in \mathcal{B}(S), \quad (47)$$

where

$$J_{g,\psi} := \frac{1}{2(2\pi)^3} \int_{\mathcal{V}_{m,+}} \int_{\mathcal{V}_{m,+}} (k+p)g(-k \cdot p) e^{i(p-k) \cdot x} \overline{\psi(k)} \psi(p) d\mu_m(p) d\mu_m(k). \quad (48)$$

As we shall see shortly, this current is conserved: $\partial_\mu J_{g,\psi}^\mu = 0$, so that $d\omega^{J_{g,\psi}} = 0$ and that it is zero or causal and future directed at each point of \mathbb{M} .

To extend the definition of \mathbb{T}_S^g to every smooth Cauchy surface of \mathbb{M} , we cannot directly follow the approach of [9] – based on a smart decompositions of $\mathbb{T}(\Delta)$ in terms of isometries and the standard PVM of the position observable on \mathbb{R}^3 – because of the appearance of various spurious terms containing $t_S(\vec{x})$ and $\nabla t_S(\vec{x})$ in the expression of $\omega^{J_{g,\psi}}$. Nevertheless, the wanted extension is feasible through a more indirect way.

There are various lemmata we shall exploit to achieve the wanted result.

Lemma 46. *Consider $g : [m^2, +\infty) \rightarrow \mathbb{R}$ continuous, normalized to $g(m^2) = 1$ such that the kernel $K_g : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{C}$ in (38) is positive definite.*

If $\psi \in \mathcal{D}(\mathcal{H})$ then the current $J_{g,\psi}$ defined in (48) is smooth, bounded, conserved, and zero or causal and future-directed at any point of \mathbb{M} .

Proof. Fix a Minkowski chart x^0, x^1, x^2, x^3 . Let us take $\phi \in C_c^\infty(\mathbb{R}^3) \subset L^2(\mathbb{R}^3, d^3k)$ and let us consider the vector field in \mathbb{M} defined in (48), now in equivalent terms of ϕ instead of ψ according to (21),

$$J_{g,\phi}^\mu(x) := \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(p^\mu + k^\mu)}{2(2\pi)^3 \sqrt{k^0 p^0}} g(-k \cdot p) e^{i(p-k) \cdot x} \overline{\phi(\vec{k})} \phi(\vec{p}) d^3p d^3k.$$

With the said hypotheses on ϕ , a direct use of the Lebesgue dominate convergence theorem permit us to pass the x^ν derivatives of every order under the integration symbol proving that J is bounded and smooth. However, the first derivative yields

$$\partial_\mu J_{g,\phi}^\mu(x) = i \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(p^\mu + k^\mu)(p_\mu - k_\mu)}{2(2\pi)^3 \sqrt{k^0 p^0}} g(-k \cdot p) e^{i(p-k) \cdot x} \overline{\phi(\vec{k})} \phi(\vec{p}) d^3p d^3k = 0$$

since $(p^\mu + k^\mu)(p_\mu - k_\mu) = -m^2 + m^2 + k^\mu p_\mu - p^\mu k_\mu = 0$. Finally, as established in [9], $J(x)$ is either zero or causal and future-directed for every $x \in \mathbb{M}$ (see Theorem 52 in [9] and (c) in its proof). \square

Lemma 47. *Consider g as in Lemma 46. If S is a smooth Cauchy surface, it holds*

$$\int_S \omega^{J_{g,\psi}} = \langle \psi | \psi \rangle. \quad (49)$$

for every $\psi \in \mathcal{D}(\mathcal{H})$.

Proof. In view of Lemma 46, since $J_{g,\psi}$ satisfies all required hypotheses, we can apply Proposition 37 for S' given by the time-slice at $x^0 = 0$, taking (31) into account and finally obtaining that

$$\int_S \omega^{J_{g,\phi}} = \int_{S'} \omega^{J_{g,\phi}} = \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(p^0 + k^0)}{2(2\pi)^3 \sqrt{k^0 p^0}} g(-k \cdot p) e^{i(\vec{p}-\vec{k}) \cdot \vec{x}} \overline{\phi(\vec{k})} \phi(\vec{p}) d^3p d^3k d^3x = \langle \phi | \phi \rangle_{L^2(\mathbb{R}^3)},$$

which is the thesis. The last identity is due to the fact that, as established in [9], the last integral is nothing but $\langle \phi | \mathbb{T}_{\mathbb{R}_0^3}(\mathbb{R}^3) \phi \rangle$ where $\mathcal{B}(\mathbb{R}^3) \ni \Delta \mapsto \mathbb{T}_{\mathbb{R}_0^3}(\Delta)$ is a normalized POVM in $\mathbb{R}_{x^0=0}^3$, because is a POVM defined as in Definition 39. \square

Lemma 48. *Let \mathcal{K} be a complex Hilbert space and $D \subset \mathcal{K}$ a dense subspace. Consider a Hermitian map $\Lambda : D \times D \rightarrow \mathbb{C}$ such that $|\Lambda(x, x)| \leq C\|x\|^2$ for some constant $C \geq 0$ and $\Lambda(x, x) \geq 0$ for every $x \in D$. Then a unique operator $A \in \mathfrak{B}(\mathcal{K})$ exists such that $\Lambda(x, x) = \langle x|Ax \rangle$ for all $x \in D$. Furthermore $A \geq 0$ and $\|A\| \leq C$.*

Proof. See Appendix A. □

We are in a position to prove that a normalized POVM induced by the operators $\mathbb{T}_S^g(\Delta)$ exists on every smooth Cauchy surface of \mathbb{M} and satisfies the natural coherence requirement (13) also in that case.

Theorem 49. *Let $S \in \mathcal{C}_{\mathbb{M}}$ and suppose that $g : [m^2, +\infty) \rightarrow \mathbb{R}$ is continuous, $g(m^2) = 1$ and the kernel in (38) is positive definite. Then,*

- (a) *there is a unique \mathcal{H} -POVM, still indicated by $\{\mathbb{T}_S^g(\Delta)\}_{\Delta \in \mathcal{B}(S)}$, which satisfies (47) when $\psi \in \mathcal{D}(\mathcal{H})$:*

$$\langle \psi | \mathbb{T}_S^g(\Delta) \psi \rangle = \int_{\Delta} \omega^{J_{g,\psi}}, \quad \Delta \in \mathcal{B}(S),$$

so that, $\mathbb{T}_S^g \ll \nu_S$ if $S \in \mathcal{C}_{\mathbb{M}}^s$ due to (34).

- (b) *\mathbb{T}_S^g is normalized: $\mathbb{T}_S^g(S) = I$;*
(c) *\mathbb{T}_S^g satisfies Definition 39 (i.e., it is a POL with causal kernel according to [9]) when $S = \Sigma$ is a time slice of a Minkowski coordinate system;*
(d) *if $\Delta \in S \cap S'$ is a Borel set, where $S' \in \mathcal{C}_{\mathbb{M}}$, then*

$$\mathbb{T}_S^g(\Delta) = \mathbb{T}_{S'}^g(\Delta).$$

The family $\mathbb{T}^g := \{\mathbb{T}_S^g\}_{S \in \mathcal{C}_{\mathbb{M}}^s}$, where we extend each POVMs to $\mathcal{M}(S)$ according to Proposition 23, is a spacelike Cauchy localization observable in the sense of Definition 21.

Proof. Item (c) is a trivial consequence of (a),(b), and [9], so we prove (a), (b), and (d). The last statement is evident in view of the items (a), (b), (d) and Def. 21

We prove (a) and (b) together. Let $x^0 = t_S(\vec{x})$ be the map which defines S by identifying it with the spatial \mathbb{R}^3 in a given Minkowski system of coordinates x^0, \vec{x} according to Proposition 20. To make easier the notation we shall omit g in \mathbb{T}_S^g . If $\Delta \in \mathcal{B}(\mathbb{R}^3)$ is bounded, so that the integral below is defined, consider the Hermitian form

$$\begin{aligned} & \Lambda_{\Delta}(\psi, \psi') \\ & := \int_{\Delta} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(p^0 + k^0) - (\vec{p} + \vec{k}) \cdot \nabla t_S(\vec{x})}{2(2\pi)^3 \sqrt{k^0 p^0}} g(-k \cdot p) e^{i(\vec{p} - \vec{k}) \cdot \vec{x} - i(p^0 - k^0)t_S(\vec{x})} \overline{\phi_{\psi}(\vec{k})} \phi_{\psi'}(\vec{p}) d^3 p d^3 k d^3 x \end{aligned}$$

for $\psi, \psi' \in \mathcal{D}(\mathcal{H})$ and $\phi_\psi, \phi_{\psi'}$ are the corresponding elements in the isomorphic Hilbert space $L^2(\mathbb{R}^3, d^3p)$. Since (use (31))

$$\Lambda(\psi, \psi) = \int_{\Delta} \omega^{J_{g,\psi}},$$

we conclude that

$$0 \leq \Lambda(\psi, \psi) \leq \langle \psi | \psi \rangle. \quad (50)$$

The former inequality arises from the last identity in (31) when remembering that $J_{g,\psi}$ is zero or causal and future directed for Lemma 46. The latter inequality in (50) is consequence of the positivity of the integrand $\omega^{J_{g,\psi}}$ in coordinates (again for (31)) and of Lemma 47. We can finally apply Lemma 48 proving that there exists $\mathsf{T}_S(\Delta) \in \mathfrak{B}(\mathcal{H})$ such that $0 \leq \mathsf{T}_S(\Delta) \leq I$ and

$$\langle \psi | \mathsf{T}_S(\Delta) \psi \rangle = \int_{\Delta} \omega^{J_{g,\psi}}. \quad (51)$$

Eq.(47) is therefore satisfied for Δ bounded. To go on we define $\mathcal{F}(S)$ as the ring of Borel sets of S which are bounded in \mathbb{R}^3 . It is clear that the σ -algebra generated by $\mathcal{F}(S)$ is $\mathcal{B}(S)$ itself. For the elements $\Delta \in \mathcal{F}(S)$ we can use the positive operator $\mathsf{T}_S(\Delta) : \mathcal{H} \rightarrow \mathcal{H}$ bounded by I defined above. We want to prove that, if $\psi \in \mathcal{H}$, the map

$$\nu_\psi : \mathcal{F}(S) \ni \Delta \mapsto \langle \psi | \mathsf{T}_S(\Delta) \psi \rangle \in [0, \|\psi\|^2] \quad \psi \in \mathcal{H}$$

is a σ -additive premeasure on $\mathcal{F}(S)$. We stress that σ -additivity is in guaranteed by the monotone convergence theorem referred to the integral in (51) – using the fact that the integrand is positive. However this argument works *only when* $\psi \in \mathcal{D}(\mathcal{H})$ since (51) is given for this type of functions. On the other hand, the very structure of (51) implies (simple) additivity of ν_ψ for $\psi \in \mathcal{H}$ just by the continuity of $\mathsf{T}_S(\Delta)$. To prove σ -additivity for the general L^2 case, consider $\Delta \in \mathcal{F}(S)$ which is the countable union of pairwise-disjoint sets $\Delta_n \in \mathcal{F}(S)$. If $\psi \in \mathcal{H}$, since all operators $\mathsf{T}_S(\Delta_n)$ are positive and additivity holds, we conclude that

$$\mathsf{T}_N := \sum_{n=0}^N \mathsf{T}_S(\Delta_n) \geq 0 \quad \text{as well as} \quad \mathsf{T}_N \leq \mathsf{T}_{N+1} \leq I.$$

The latter inequality arises from $\mathsf{T}_N = \mathsf{T}_S(\cup_{n=0}^N \Delta_n) \leq I$. A known result on increasing sequences of positive operators (Proposition 3.76 [28]) proves that there exist a bounded operator $P : \mathcal{H} \rightarrow \mathcal{H}$ such that $0 \leq P \leq I$ and

$$\langle \psi | \mathsf{T}_N \psi \rangle \rightarrow \langle \psi | P \psi \rangle \quad \text{as } N \rightarrow +\infty \text{ for all } \psi \in \mathcal{H} \quad (52)$$

However, σ -additivity for $\psi \in \mathcal{D}(\mathcal{H})$ guarantees that

$$\sum_{n=0}^N \nu_\psi(\Delta_n) = \langle \psi | \mathsf{T}_N \psi \rangle \rightarrow \langle \psi | \mathsf{T}_S(\Delta) \psi \rangle \quad \text{as } N \rightarrow +\infty \text{ for all } \psi \in \mathcal{D}(\mathcal{H}).$$

Comparing the found limits, we conclude that $\langle \psi | \mathbb{T}_S(\Delta) \psi \rangle = \langle \psi | P \psi \rangle$ for every $\psi \in \mathcal{D}(\mathcal{H})$. By continuity of P and $\mathbb{T}_S(\Delta)$, this identity extends to $\psi \in \mathcal{D}(\mathcal{H})$ which, in turn, yields $\mathbb{T}_S(\Delta) = P$ since the Hilbert space is complex. In summary, (52) can be rephrased to

$$\sum_{n \in \mathbb{N}} \langle \phi | \mathbb{T}_S(\Delta_n) \phi \rangle = \langle \phi | \mathbb{T}_S(\cup_{n \in \mathbb{N}} \Delta_n) \phi \rangle \quad \text{for all } \phi \in \mathcal{H},$$

that is the wanted σ -additivity property for ν_ϕ . As a consequence of the Carathéodory extension theorem, there is a positive σ -additive measure $\overline{\nu}_\phi : \mathcal{B}(S) \rightarrow [0, +\infty]$ which extends ν_ϕ , for every given $\phi \in L^2(\mathbb{R}^3, d^3k)$. This measure is unique because S is countable union of sets in $\mathcal{F}(S)$ thus with premeasure ν_ϕ finite. $\overline{\nu}_\phi$ is finite by inner continuity: If $\Delta_n \in \mathcal{F}(S)$ satisfy $\Delta_n \subset \Delta_{n+1}$ and $\cup_{n \in \mathbb{N}} \Delta_n = S$, we have

$$\overline{\nu}_\phi(S) = \sup_{n \in \mathbb{N}} \overline{\nu}_\phi(\Delta_n) = \sup_{n \in \mathbb{N}} \nu_\phi(\Delta_n) \leq \|\phi\|^2$$

Now we use ν_ϕ to extend the definition of $\mathbb{T}_S(\Delta)$ to the case of $\Delta \in \mathcal{B}(S)$ without the constraint $\Delta \in \mathcal{F}(S)$.

Consider $\Delta \in \mathcal{B}(S)$ and a sequence $\Delta_n \in \mathcal{F}(S)$ such that $\Delta_n \subset \Delta_{n+1} \subset \Delta$ and $\cup_{n \in \mathbb{N}} \Delta_n = \Delta$. For instance, $\Delta_n = \Delta \cap B_n(0)$, where $B_n(0)$ is the open ball of radius n centered at the origin of $\mathbb{R}^3 \equiv S$. By construction, using additivity, $0 \leq \mathbb{T}_S(\Delta_n) \leq \mathbb{T}_S(\Delta_{n+1}) \leq I$. Therefore, again for Proposition 3.76 [28], there exists a bounded everywhere defined operator

$$\mathbb{T}_S(\Delta) := s\text{-}\lim_{n \rightarrow +\infty} \mathbb{T}_S(\Delta_n) \quad (53)$$

such that $0 \leq \mathbb{T}_S(\Delta) \leq I$. With an argument strictly similar to the one used before, this operator satisfies

$$\overline{\nu}_\phi(\Delta) = \langle \phi | \mathbb{T}_S(\Delta) \phi \rangle \quad \text{for every } \phi \in \mathcal{H}.$$

This identity proves both that $\mathbb{T}_S(\Delta)$ extends the definition for $\Delta \in \mathcal{F}(S)$ and that $\mathbb{T}_S(\Delta)$ does not depend on the used sequence of sets $\Delta_n \in \mathcal{F}(S)$ to define it. Now fix $\Delta \in \mathcal{B}(S)$ with Δ unbounded. Taking advantage of a sequence $\Delta_n := \Delta \cap B_n(0)$, according to (53) and (51), we can prove that (47) is valid also if Δ unbounded for $\psi \in \mathcal{D}(\mathcal{H})$:

$$\langle \psi | \mathbb{T}_S(\Delta) \psi \rangle = \lim_{n \rightarrow +\infty} \langle \psi | \mathbb{T}_S(B_n(0) \cap \Delta) \psi \rangle = \lim_{n \rightarrow +\infty} \int_{B_n(0) \cap \Delta} \omega^{J_{g,\psi}} = \int_{\Delta} \omega^{J_{g,\psi}}.$$

In the last limit, we exploited the monotone convergence theorem (the integrand being positive). Finally, let us consider the case $\Delta = S \equiv \mathbb{R}^3$. Again, if $\psi \in \mathcal{D}(\mathcal{H})$

$$\langle \psi | \mathbb{T}_S(S) \psi \rangle = \lim_{n \rightarrow +\infty} \langle \psi | \mathbb{T}_S(B_n(0)) \psi \rangle = \lim_{n \rightarrow +\infty} \int_{B_n(0)} \omega^{J_{g,\psi}} = \int_S \omega^{J_{g,\psi}} = \langle \psi | \psi \rangle$$

The last limit, due to the monotone convergence theorem (the integrand being positive) coincides to the integral on the whole S which is $\langle \psi | \psi \rangle$ according to Lemma 47. Hence $\langle \psi | \mathbb{T}_S(S) - I \psi \rangle = 0$

for every $\psi \in \mathcal{D}(\mathcal{H})$. The standard density and continuity argument (in a complex Hilbert space) permits to conclude that $\mathbb{T}_S(S) = I$. This result also proves (b) if \mathbb{T}_S satisfies the other requirements of a POVM. To end the proof, it is sufficient to prove that $\{\mathbb{T}_S(\Delta)\}_{\Delta \in \mathcal{B}(S)}$ is a POVM in all cases. It is equivalent to prove that $\{\langle \psi | \mathbb{T}_S(\Delta) \phi \rangle\}_{\Delta \in \mathcal{B}(S)}$ is a complex measure for every choice of $\phi, \psi \in \mathcal{D}(\mathcal{H})$. The only fact to be proved is that $\mathcal{B}(\mathbb{R}^3) \ni \Delta \mapsto \langle \psi | \mathbb{T}_S(\Delta) \phi \rangle \in \mathbb{C}$ is σ -additive. This fact immediately arise by σ -additivity of the positive Borel measure $\mathcal{B}(S) \ni \Delta \mapsto \langle \chi | \mathbb{T}_S(\Delta) \chi \rangle \in [0, \|\chi\|^2]$, by choosing $\chi = \psi \pm \phi$ and $\chi = \psi \pm i\phi$ and taking advantage of the polarization identity.

The proof of (d) is trivial. If Δ is bounded, then (51) proves the assertion immediately. If Δ is unbounded, $\langle \psi | \mathbb{T}_S(\Delta) \psi \rangle$ and $\langle \psi | \mathbb{T}_{S'}(\Delta) \psi \rangle$ can be however obtained as limits of non-decreasing sequences based on bounded sets $\Delta_n \subset \Delta$ and thus they coincide as before.

The last statement of the thesis immediately arises from Proposition 24. \square

6 Spacelike Cauchy localization observables out of the stress-energy tensor of massive KG particles

In the final Sect 7 of [29], a second type of spatial localization observable was introduced by generalizing an idea by D. Terno [34] also analysed in the first part of [29]. These POVMs denoted by $M_{n,t}^{n'}$ (with $n, n' \in \mathbb{T}_+$) were constructed on spacelike flat Cauchy surfaces out of the stress energy tensor of the Klein Gordon field.

The goal of the remaining part of this section is to prove that this different notion of POVM can be defined on spacelike Cauchy surfaces of \mathbb{M} , giving rise to a Cauchy localization according to Def. 21. We expect that the construction can be generalized to any static (or stationary) globally hyperbolic spacetime referring to the Hadamard static vacuum, since the theoretical construction does not depend on the use of Fourier transform, at least at heuristic level. This conjecture will be analysed elsewhere.

6.1 The POVM M_Σ^n for massive KG particles

In Section 7 of [29], extending a notion introduced by D. Terno [34], a family of POVMs was introduced on all spacelike flat Cauchy surfaces Σ of \mathbb{M} for a common choice of a reference frame $n \in \mathbb{T}_+$. (If $n = n_\Sigma$ one obtains the very notion introduced by Terno, studied and made rigorous in the first part of [29] that is a special case of the following discussion. A rigorous proof of CT for that observable appears in [29]).

At the level of 2nd quantization of the massive real Klein-Gordon field, the considered POVM is formally defined as follows on $\Sigma \in \mathcal{C}_\mathbb{M}^{sf}$ (we readapt the notation to the choices of the present work)

$$M_\Sigma^n(\Delta) := \frac{1}{\sqrt{H_n}} P_1 \int_\Delta : \hat{T}_{\mu\nu} : (x) n^\mu n_\Sigma^\nu d\nu_\Sigma(x) P_1 \frac{1}{\sqrt{H_n}}, \quad \Delta \in \mathcal{L}(\Sigma), \quad (54)$$

where $P_1 : \mathfrak{F}_+(\mathcal{H}) \rightarrow \mathcal{H}$ is the orthogonal projector onto the one-particle space of the symmetric Fock space $\mathfrak{F}_+(\mathcal{H}_m)$ constructed upon the Minkowski vacuum state with the Hilbert space \mathcal{H}

defined as in (22) as the one-particle subspace. $:\hat{T}_{\mu\nu}:(x)$ is the *normally ordered* stress energy tensor operator. H_n is Hamiltonian operator of the quantum field in the reference frame $n \in \mathbb{T}_+$. Σ is a *flat* spacelike Cauchy surface orthogonal with constant normal unit vector $n_\Sigma \in \mathbb{T}_+$.

The overall idea at the basis of [34] and [29] is that a physical procedure to detect a particle in a region of a flat Cauchy surface may exploit the energy of the particle. However, we have many ways to synchronize a net of detectors and, as discussed in [29]: $\langle \psi | M_\Sigma^n(\Delta) \psi \rangle$ accounts for the probability to find a particle of state ψ in $\Delta \subset \Sigma$ *using a net of detectors which are (a) at rest in n but (b) synchronized on Σ* . This possibility naturally arises from the observation [29] that

$$P_1 \int_\Delta :\hat{T}_{\mu\nu}:(x) n^\mu n_\Sigma^\nu d\nu_\Sigma(x) P_1 \geq 0, \quad (55)$$

for *every* choice of $n, n_\Sigma \in \mathbb{T}_+$ and $\Delta \in \mathcal{L}(\Sigma)$, even if positivity fails¹⁵ when removing the one-particle space projectors P_1 . The rigorous definition of the normalized POVM M_Σ^n in \mathcal{H} corresponding to the formal object (54) was given in Thm 37 of [29] in terms of a kinematic deformation of the PVM $Q_\Sigma(\Delta)$ of the *Newton-Wigner position operator* [29] (see (57) below) on the spacelike flat Cauchy surface Σ :

$$\begin{aligned} M_\Sigma^n(\Delta) := & \frac{1}{2} \left(\sqrt{\frac{H_{n_\Sigma}}{H_n}} Q_{n_\Sigma}(\Delta) \sqrt{\frac{H_n}{H_{n_\Sigma}}} + \sqrt{\frac{H_n}{H_{n_\Sigma}}} Q_{n_\Sigma}(\Delta) \sqrt{\frac{H_{n_\Sigma}}{H_n}} \right) \\ & - \frac{n \cdot n_\Sigma}{2} \sqrt{\frac{H_{n_\Sigma}}{H_n}} \left(\eta^{\mu\nu} \frac{P_\mu}{H_{n_\Sigma}} Q_{n_\Sigma}(\Delta) \frac{P_\nu}{H_{n_\Sigma}} + \frac{m}{H_{n_\Sigma}} Q_{n_\Sigma}(\Delta) \frac{m}{H_{n_\Sigma}} \right) \sqrt{\frac{H_{n_\Sigma}}{H_n}}. \end{aligned} \quad (56)$$

Above $\Delta \in \mathcal{L}(\Sigma)$, $H_m := -P \cdot m$ (for $m \in \mathbb{T}_+$) is the Hamiltonian in the Minkowski reference frame m . The various everywhere-defined *bounded* composite operators H_n/H_{n_Σ} , P_ν/H_{n_Σ} etc. are defined in terms of the joint spectral measure of the four momentum operator P^μ and standard spectral calculus. The components P^μ are referred to a Minkowski chart adapted to n .

6.2 Properties of M_Σ^n for spacelike flat Cauchy surfaces Σ : Covariance, causality, no strict localizability, Newton-Wigner, Heisenberg inequality,

Though its clear from (56) that $M_\Sigma^n(\Delta) \in \mathfrak{B}(\mathcal{H})$ and that $M_\Sigma^n(\Sigma) = I$ (notice that $Q_{n_\Sigma}(\Sigma) = I$ and $P_\mu P^\mu + m^2 I = 0$), it is not evident that $M_\Sigma^n(\Delta) \geq 0$, nor the connection between (56) and (54) seems straightforward. We spend this section about these issues because the discussion will turn out useful when we shall generalize M_Σ^n to generally curved spacelike Cauchy surfaces S .

¹⁵It is interesting to observe that a similar local positivity property *does not hold* when dealing with massive Dirac particles, even if $n = n_\Sigma$ [35]: A localization POVM cannot be constructed for this type of fermions in terms of energy on a given rest space of a Minkowski reference frame according to the approach of [34, 29]. The natural physical object, useful to this goal, is instead the fermionic current operator $:\hat{J}^\mu:(x)$ [8].

From Eq.(17) in [29], we know that, if $\Sigma \in \mathcal{C}_{\mathbb{M}}^{sf}$,

$$\langle \psi | \mathbf{Q}_{\Sigma}(\Delta) \psi \rangle = \int_{\mathbb{V}_{m,+}} \overline{\psi(p)} \int_{\Delta} \int_{\mathbb{V}_{m,+}} \frac{e^{i(q-p) \cdot x}}{(2\pi)^3} \sqrt{E_{n_{\Sigma}}(p) E_{n_{\Sigma}}(q)} \psi(q) d\mu_m(q) d\nu_{\Sigma}(x) d\mu_m(p), \quad \psi \in (\mathcal{H}). \quad (57)$$

We adopted the notation

$$E_m(p) := -m \cdot p, \quad \text{for every } m \in \mathbb{T}_+ \text{ and } p \in \mathbb{V}. \quad (58)$$

($E_m(p)$ is nothing but the component p^0 of $p \in \mathbb{V}$ in any Minkowski chart adapted to m .) At this juncture, it is not difficult to see from (56) that, if $\psi \in \mathcal{D}(\mathcal{H})$ or more generally $\psi \in \mathcal{S}(\mathcal{H}) (\supset \mathcal{D}(\mathcal{H}))$ as used in [29], then

$$\begin{aligned} \langle \psi | \mathbf{M}_{\Sigma}^n(\Delta) \psi \rangle = \\ \int_{\mathbb{V}_{m,+}} \overline{\psi(p)} \int_{\Delta} \int_{\mathbb{V}_{m,+}} \frac{e^{i(q-p) \cdot x}}{(2\pi)^3} \frac{E_n(p) E_{n_{\Sigma}}(q) + E_n(q) E_{n_{\Sigma}}(p) - n \cdot n_{\Sigma}(p \cdot q + m^2)}{2\sqrt{E_n(q) E_n(p)}} \psi(q) d\mu_m(q) d\nu_{\Sigma}(x) d\mu_m(p), \end{aligned} \quad (59)$$

Conversely, since $\mathcal{D}(\mathcal{H})$ and $\mathcal{S}(\mathcal{H})$ are dense in \mathcal{H} and $\mathbf{M}_{\Sigma}^n(\Delta) \in \mathfrak{B}(\mathcal{H})$, identity (59) completely determines $\mathbf{M}_{\Sigma}^n(\Delta)$ by polarization and continuity.

First of all, we prove that (59) can be written into an equivalent form already used in [29], which eventually leads to both the requested positivity condition and the relation with (54).

Lemma 50. *If $\psi \in \mathcal{S}(\mathcal{H})$, the right-hand side of (59) can be equivalently written with the first two integrals interchanged:*

$$\int_{\Delta} \int_{\mathbb{V}_{m,+}} \int_{\mathbb{V}_{m,+}} \frac{e^{i(q-p) \cdot x}}{(2\pi)^3} \frac{E_n(p) E_{n_{\Sigma}}(q) + E_n(q) E_{n_{\Sigma}}(p) - n \cdot n_{\Sigma}(p \cdot q + m^2)}{2\sqrt{E_n(q) E_n(p)}} \overline{\psi(p)} \psi(q) d\mu_m(q) d\mu_m(p) d\nu_{\Sigma}(x) \quad (60)$$

Proof. See Appendix A. □

To go no, if $\psi \in \mathcal{D}(\mathcal{H})$ (or more generally, $\psi \in \mathcal{S}(\mathcal{H})$) and $n \in \mathbb{T}_+$, we define the solution of the Klein-Gordon equation

$$\Phi_n^{\psi}(x) := \int_{\mathbb{V}_{m,+}} \frac{\psi(p) e^{ip \cdot x}}{(2\pi)^{3/2} \sqrt{E_n(p)}} d\mu_m(p). \quad (61)$$

$\Phi_n^{\psi} = \Phi_n^{\psi}(x)$ is smooth on \mathbb{M} and bounded with all of its derivatives. More strongly $\Phi_n^{\psi} \in \mathcal{S}(\Sigma)$ for every flat $\Sigma \in \mathcal{C}_{\mathbb{M}}^{sf}$, where $\mathcal{S}(\Sigma)$ is the usual Schwartz space on $\mathbb{R}^3 \equiv \Sigma$ referring to any Minkowski chart adapted to n_{Σ} . We stress that (61) is *not* the standard covariant Klein-Gordon wavefunction φ_{ψ} (26) associated to a state $\psi \in \mathcal{H}$, since the integrand above includes a further "anomalous" factor $E_p^{-1/2}(p)$. This latter can be traced back to the factors $H_n^{1/2}$ in (55).

Definition 51. If $\psi \in \mathcal{D}(\mathcal{H})$ and $n \in \mathbb{T}_+$, the n -normalized stress energy operator is, in components of a Minkowski chart,

$$\begin{aligned} T_{\mu\nu}^\psi(x)_n &:= \frac{1}{2} \left(\partial_\mu \overline{\Phi_n^\psi(x)} \partial_\nu \Phi_n^\psi(x) + \partial_\mu \Phi_n^\psi(x) \partial_\nu \overline{\Phi_n^\psi(x)} \right) \\ &\quad - \frac{1}{2} \eta_{\mu\nu} \left(\partial^\alpha \overline{\Phi_n^\psi(x)} \partial_\alpha \Phi_n^\psi(x) + m^2 \overline{\Phi_n^\psi(x)} \Phi_n^\psi(x) \right), \end{aligned} \quad (62)$$

and the associated current

$$J_n^{\psi\mu}(x) := n^\nu T_\nu^{\psi\mu}(x)_n. \quad (63)$$

■

Lemma 52. With the said definitions, for $\psi \in \mathcal{D}(\mathcal{H})$ and $n \in \mathbb{T}_+$,

- (a) J_n^ψ is smooth, belongs to $\mathcal{S}(\Sigma)$ for every $\Sigma \in \mathcal{C}_M^{sf}$, and is bounded;
- (b) J_n^ψ is conserved;
- (c) $J_n^\psi(x)$ is either zero or causal and past-directed if $x \in \mathbb{M}$, so that

$$J_n^\psi(x) \cdot m \geq 0 \quad \forall x \in \mathbb{M}, \forall m \in \mathbb{T}_+. \quad (64)$$

Proof. (a) is a trivial consequence of the definition. (b) Follows from the Klein Gordon equation which is satisfied by Φ_n^ψ . (c) was established in Prop. 30 [29]. \square

We are prompt to prove that M_Σ^n is a normalized POVM when Σ is a spacelike flat Cauchy surface. In particular $M_\Sigma^n(\Delta) \geq 0$.

Theorem 53. If $\Sigma \subset \mathbb{M}$ is a spacelike flat Cauchy surface, the family of operators (56), when $\Delta \in \mathcal{L}(\Sigma)$, defines a normalized \mathcal{H} -POVM. Furthermore, if $\psi \in \mathcal{D}(\mathcal{H})$ (or more generally $\psi \in \mathcal{S}(\mathcal{H})$),

$$\langle \psi | M_\Sigma^n(\Delta) \psi \rangle = \int_\Delta T_{\mu\nu}^\psi(x)_n n^\mu n_\Sigma^\nu d\nu_\Sigma(x), \quad \Sigma \in \mathcal{C}_M^{sf}. \quad (65)$$

Proof. First of all $\langle \psi | M_\Sigma^n(\Delta) \psi \rangle \geq 0$ if $\psi \in \mathcal{S}(\mathcal{H})$. Indeed, from Lemma 50 and expanding (62) as prescribed in (61), we find

$$\begin{aligned} &\langle \psi | M_\Sigma^n(\Delta) \psi \rangle \\ &= \int_\Delta \int_{\mathbb{V}_{m,+}} \int_{\mathbb{V}_{m,+}} \frac{e^{i(q-p)\cdot x}}{(2\pi)^3} \frac{E_n(p)E_{n_\Sigma}(q) + E_n(q)E_{n_\Sigma}(p)n \cdot n_\Sigma(p \cdot q + m^2)}{2\sqrt{E_n(q)E_n(p)}} \overline{\psi(p)} \psi(q) d\mu_m(q) d\mu_m(p) d\nu_\Sigma(x) \\ &= \int_\Delta J_n^\psi \cdot n_\Sigma d\nu_\Sigma(x) = \int_\Delta n^\nu T_\nu^{\psi\mu}(x)_n \cdot n_{\Sigma\mu} d\nu_\Sigma(x) \geq 0 \end{aligned}$$

on account of (64) and where the integral is finite because $J_n^\psi \in \mathcal{S}(\Sigma)$. (The found identity also establishes (65).) As a consequence $M_\Sigma^n(\Delta) \geq 0$ because, as already observed, $\langle \psi | M_\Sigma^n(\Delta) \psi \rangle$ is the limit of analogous matrix elements with $\psi \in \mathcal{D}(\mathcal{H})$ or $\mathcal{S}(\mathcal{H})$. Normalization of the POVM has been already discussed beforehand, and (56) itself implies weak σ -additivity from the analogous property of Q_Σ . \square

Regarding the connection between (56) and (54), from $E_p^{-1/2}(p)$ in (61) and the expression (62) of the stress energy tensor, it is not difficult to see that (65) is nothing but the matrix element of (54) with respect a one-particle state $\psi \in \mathcal{H}$.

Referring to the spatial localization observable $\{M_\Sigma^n\}_{n \in \mathbb{T}_+, \Sigma \in \mathcal{C}_M^{sf}}$, the \mathcal{P}_+ -covariance relations analogous to (44) are valid (Thm 37 in [29])

$$U_h M_\Sigma^n(\Delta) U_h^{-1} = M_{h\Sigma}^{\Lambda_h n}(h\Delta), \quad \forall \Sigma \in \mathcal{C}_M^{sf}, \forall \Delta \in \mathcal{L}(\Sigma), \quad \forall h = (v_h, \Lambda_h) \in \mathcal{P}_+. \quad (66)$$

The validity of these relations is actually required in the spirit of the very definition of *relativistic* spatial localization observable assumed in [29] (Definition 18 therein). And, in fact, $\{M_\Sigma^n\}_{n \in \mathbb{T}_+, \Sigma \in \mathcal{C}_M^{sf}}$ satisfies that definition in a broader sense, due to the presence of the further specification¹⁶ $n \in \mathbb{T}_+$.

Let us pass to discuss causality properties.

Theorem 54 ([29]). *For a given choice of $n \in \mathbb{T}_+$, the spatial localization observable $\{M_\Sigma^n\}_{\Sigma \in \mathcal{C}_M^{sf}}$ satisfies the CC in Definition 2.*

Proof. Theorem 39 [29]. □

Corollary 55. *If $\psi \in \mathcal{H}$, the localization probability associated to the spatial localization observable $\{M_\Sigma^n\}_{\Sigma \in \mathcal{C}_M^{sf}}$ cannot be zero outside a bounded set in Σ .*

Proof. If states producing localized probability distributions as above exist CT would fail as a consequence of the Hegerfeldt theorem 4 and thus also CC would be false. (See [29]) for a discussion on this point.) □

Remark 56. In spite of this obstruction, it is possible to show [29] that probability distributions localized in bounded sets can be arbitrarily well approximated by probability distributions arising by suitable sequences of state ψ_k when $n = n_\Sigma$. However it is not difficult to generalize this result to $n \neq n_\Sigma$. ■

As already observed for the POVMs \mathbb{T}_Σ^g , there is again the interesting relation between the *first moment* of the POVM M_Σ^n on a spacelike flat Cauchy surface Σ and the *Newton-Wigner* selfadjoint operators $N_\Sigma^1, N_\Sigma^2, N_\Sigma^3$ [29] associated to a Minkowski chart x^0, x^1, x^2, x^3 such that the slice $x^0 = 0$ coincides with Σ . Generalizing Thm 26 in [29], one sees that the following theorem is true. Below, $\Delta_\psi x_{M_\Sigma^n}^a$ denotes the standard deviation for the coordinate x^a referred to the probability distribution $\langle \psi | M_\Sigma^n(\cdot) \psi \rangle$ constructed out of the POVM M_Σ^n in the state defined by the unit vector ψ .

Theorem 57. *Referring to the POVM M_Σ^n for $\Sigma \in \mathcal{C}_M^{sf}$, the following facts are true.*

¹⁶If $n = n_\Sigma$, giving rise to the *Terno spatial localization observable* studied in first part of [29], it fully satisfies that definition

(a) The a -first moment of M_Σ^n is defined for every $\psi \in \mathfrak{S}(\mathcal{H})$ with $\|\psi\| = 1$ and

$$\int_\Sigma x^a \langle \psi | M_\Sigma^n(d^3x) \psi \rangle = \langle \psi | N_\Sigma^a \psi \rangle \quad \text{for } a = 1, 2, 3. \quad (67)$$

In particular, N_Σ^a is the unique selfadjoint operator in \mathcal{H} which satisfies the identity above.

(b) The **Heisenberg inequality** turns out to be corrected as, for $\psi \in \mathfrak{S}(\mathcal{H})$,

$$\Delta_\psi x_{M_\Sigma^n}^a \Delta_\psi P_a \geq \frac{\hbar}{2} \sqrt{1 + 4(\Delta_\psi P_a)^2 \langle \psi | K_a^{M_\Sigma^n} \psi \rangle}, \quad a = 1, 2, 3$$

where $K_a^{M_\Sigma^n} \in \mathfrak{B}(\mathcal{H})$ is a selfadjoint operator which is a (spectral) function of the four momentum observable P with the form (79) and $K_a^{M_\Sigma^n} \geq 0$.

Proof. See Appendix A. □

Observe that the result in (67) does not depend on n .

Remark 58. (1) If also the identity

$$\int_\Sigma (x^a)^2 \langle \psi | M_\Sigma^n(d^3x) \psi \rangle = \langle \psi | (N_\Sigma^a)^2 \psi \rangle \quad (\text{false!}),$$

were valid one could apply a known theorem by Naimark about the decomposition of maximally symmetric operators (here N_Σ^a) in terms of POVMs (see Theorem 23 in [13] and the discussion about it) obtaining $M_\Sigma^n = Q_\Sigma$. This is obviously false and it is also reflected by the presence of the term $\langle \psi | K_a^{M_\Sigma^n} \psi \rangle$ in the modified Heisenberg inequality.

(2) If $U_t^{(n)}$ is the unitary time evolutor corresponding to the time evolution along n in the spacetime \mathbb{M} , it is easy to see that the Heisenberg evolution $U_t^{(n)} N_\Sigma^a U_t^{(n)\dagger}$ of N_Σ^a on the right-hand side of (67) equals the integral on the left-hand side over the correspondingly temporally translated time slice Σ_t . As already observed in [29], this fact implies that the worldline $\mathbb{R} \ni t \mapsto (t, \int_{\Sigma_t} \vec{x} \langle \psi | M_{\Sigma_t}^n(d^3x) \psi \rangle)$ is *timelike* (Corollary 14 in [29]) as expected by massive particles. ■

6.3 A spacelike Cauchy localization observable $M^n = \{M_S^n\}_{S \in \mathcal{C}_\mathbb{M}^s}$ for massive KG particles.

We are in a position to prove that a normalized POVM M_S^n exists on every spacelike Cauchy surface S of \mathbb{M} . We shall also obtain that the elements of the POVMs do not depend on the Cauchy surface they belong to. In other words we have a spacelike Cauchy localization M^n .

Theorem 59. Consider $S \in \mathcal{C}_\mathbb{M}^s$ and $n \in \mathbb{T}_+$. Then,

(a) there is a unique \mathcal{H} -POVM, still indicated by $\{M_S^n(\Delta)\}_{\Delta \in \mathcal{B}(S)}$, which satisfies (65) also for generic spacelike Cauchy's surface S when $\psi \in \mathcal{D}(\mathcal{H})$:

$$\langle \psi | M_S^n(\Delta) \psi \rangle = \int_{\Delta} T_{\mu\nu}^{\psi}(x)_n n^{\mu} n_S^{\nu} d\nu_S(x), \quad \Delta \in \mathcal{B}(S), \quad (68)$$

so that, $M_S^n \ll \nu_S$.

(b) M_S^n is normalized: $M_S^n(S) = I$;

(c) M_{Σ}^n satisfies (56) when $\Sigma \in \mathcal{C}_{\mathbb{M}}^{sf}$;

(d) if $\Delta \in S \cap S'$ is a Borel set, where $S \in \mathcal{C}_{\mathbb{M}}^s$, then

$$M_S^n(\Delta) = M_{S'}^n(\Delta).$$

The family $M^n := \{M_S^n\}_{S \in \mathcal{C}_{\mathbb{M}}^s}$, where we extend each POVMs to $\mathcal{M}(S)$ according to Proposition 23, is a spacelike Cauchy localization observable according to Definition 21.

Proof. Item (c) is a trivial consequence of (a) and (b), so we prove (a), (b), and (d).

Let $x^0 = t_S(\vec{x})$ be the map which defines S by identifying it with the spatial \mathbb{R}^3 in a given Minkowski system of coordinates x^0, \vec{x} according to Proposition 20. If $\Delta \in \mathcal{B}(\mathbb{R}^3)$ is bounded consider the Hermitian form

$$\Lambda(\psi, \psi') := \int_{\Delta} \omega^{J_n^{\psi\psi'}},$$

where, taking (61) into account,

$$\begin{aligned} T_{\mu\nu}^{\psi\psi'}(x)_n &:= \frac{1}{2} \left(\partial_{\mu} \overline{\Phi_n^{\psi}}(x) \partial_{\nu} \Phi_n^{\psi'}(x) + \partial_{\mu} \Phi_n^{\psi'}(x) \partial_{\nu} \overline{\Phi_n^{\psi}}(x) \right) \\ &\quad - \frac{1}{2} \eta_{\mu\nu} \left(\partial^{\alpha} \overline{\Phi_n^{\psi}}(x) \partial_{\alpha} \Phi_n^{\psi'}(x) + m^2 \overline{\Phi_n^{\psi}}(x) \Phi_n^{\psi'}(x) \right), \end{aligned} \quad (69)$$

and

$$J_n^{\psi\psi'\mu}(x) := n^{\nu} T_{\nu}^{\psi\psi'\mu}(x)_n, \quad (70)$$

As Δ is bounded, the integral is well defined. Since (use (31)) we conclude that

$$0 \leq \Lambda(\psi, \psi) \leq \langle \psi | \psi \rangle. \quad (71)$$

The former inequality arises from the last identity in (31) when observing that (see (63)) $J_n^{\psi\psi} = J_n^{\psi}$ is zero or causal and future directed as established in Lemma 52. The latter inequality in (71) is consequence of the positivity of the integrand $\omega^{J_n^{\psi}}$ in coordinates (again for (31)) and (64) which first of all imply (we write $J_n^{\psi} = J_n^{\psi\psi}$)

$$\int_{\Delta} \omega^{J_n^{\psi}} \leq \int_S \omega^{J_n^{\psi}}.$$

On the other hand

$$\int_S \omega^{J_n^\psi} = \langle \psi | \psi \rangle$$

because we can apply Proposition 36 (since J_n^ψ is also smooth, bounded, and conserved according to Lemma 52) choosing a spacelike flat surface S' , obtaining

$$\int_S \omega^{J_n^\psi} = \int_{S'} \omega^{J_n^\psi} = \langle \psi | \psi \rangle. \quad (72)$$

The last identity is due to the fact that the last integral is nothing but $\langle \psi | M_{S'}^n(\Delta) \psi \rangle = \langle \psi | \psi \rangle$ due to Theorem 53.

We can finally apply Lemma 48 proving that there exists $M_S^n(\Delta) \in \mathfrak{B}(\mathcal{H})$ such that $0 \leq M_S^n(\Delta) \leq I$ and

$$\langle \psi | M_S^n(\Delta) \psi \rangle = \int_\Delta \omega^{J_n^\psi} = \int_\Delta T_{\mu\nu}^\psi(x) n^\mu n_S^\nu d\nu_S(x), \quad (73)$$

so that (68) is satisfied for Δ bounded. From this point on, the proof is identical to the one of Theorem 49. In particular, the normalization condition (b) follows now from (72). The last statement of the thesis immediately arises from Proposition 24. \square

7 Moments of T_Σ^g and M_Σ^n , Newton-Wigner operator, and Heisenberg inequality

We address the reader to the discussion and references in [29] about the Newton-Wigner observables for a massive spinless particle.

Referring to the representation $L^2(\mathbb{R}^3, d^3p)$ (see Sect. 4.1) of the one-particle space \mathcal{H} through the Hilbert space isomorphism (21) $F : \mathcal{H} \rightarrow L^2(\mathbb{R}^3, d^3p)$, the joint PVM of the selfadjoint Newton Wigner operators N^1, N^2, N^3 for a massive spinless particle, associated to the coordinates $\vec{x} = (x^1, x^2, x^3)$ on \mathbb{R}^3 in a Minkowski frame x^0, x^1, x^2, x^3 at time $x^0 = 0$, is given by

$$(FQ(\Delta)F^{-1}\phi)(\vec{p}) := \int_\Delta \frac{e^{i(\vec{q}-\vec{p})\cdot\vec{x}}}{(2\pi)^3} \phi(\vec{q}) d^3q, \quad \phi \in \mathcal{S}(\mathbb{R}^3) \quad \text{or} \quad C_c^\infty(\mathbb{R}^3), \quad \Delta \in \mathcal{B}(\mathbb{R}^3). \quad (74)$$

It extends by continuity to the whole Hilbert space. Notice that $\mathfrak{S}(\mathcal{H}) = F(\mathcal{S}(\mathbb{R}^3))$ and $\mathcal{D}(\mathcal{H}) = F(C_c^\infty(\mathbb{R}^3))$ are invariant spaces and cores for each N^a and thereon they are unitarily equivalent to the respective differential operator $FN^a|_{\mathcal{S}(\mathbb{R}^3)}F^{-1} = i\frac{\partial}{\partial p^a}$.

Theorems 44 and 57 are subcases of the following result.

Proposition 60. *Consider a normalized POVM on \mathbb{R}^3 satisfying (with $\phi_\psi := F\psi$)*

$$\langle \psi | A(\Delta) \psi \rangle = \int_\Delta d^3x \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} d^3p d^3q \frac{e^{i(\vec{q}-\vec{p})\cdot\vec{x}}}{(2\pi)^3} \phi_\psi(\vec{p}) K_A(\vec{p}, \vec{q}) \phi_\psi(\vec{q}), \quad \psi \in \mathcal{D}(\mathcal{H}), \Delta \in \mathcal{B}(\mathbb{R}^3) \quad (75)$$

where K_A is a positive definite kernel which satisfies $K_A(\vec{p}, \vec{p}) = 1$ for every $\vec{p} \in \mathbb{R}^3$.

(a) If $K_{\mathbf{A}}$ is real and smooth, then

$$\int_{\mathbb{R}^3} x^a \langle \psi | \mathbf{A}(d^3x) \psi \rangle = \langle \psi | N^a \psi \rangle \quad \text{for } a = 1, 2, 3 \text{ and } \psi \in \mathcal{D}(\mathcal{H}), \text{ with } \|\psi\| = 1, \quad (76)$$

where N^a is the unique selfadjoint operator in \mathcal{H} which satisfies the identity above. More generally, if $\alpha := (\alpha_1, \alpha_2, \alpha_3)$ is a multi index, so that $x^\alpha := (x^1)^{\alpha_1} (x^2)^{\alpha_2} (x^3)^{\alpha_3}$,

$$\begin{aligned} & \int_{\mathbb{R}^3} (x^1)^{\alpha_1} (x^2)^{\alpha_2} (x^3)^{\alpha_3} \langle \psi | \mathbf{A}(d^3x) \psi \rangle = \langle \psi | (N^1)^{\alpha_1} (N^2)^{\alpha_2} (N^3)^{\alpha_3} \psi \rangle \\ & + \sum_{1 < |\beta|, \beta \leq \alpha} \binom{\alpha}{\beta} \int_{\mathbb{R}^3} \left(i^{|\alpha|-|\beta|} \frac{\partial^{|\alpha|-|\beta|} \overline{\phi_\psi(\vec{p})}}{\partial p^{\alpha-\beta}} \right) \left(i^{|\beta|} \frac{\partial^{|\beta|} K_{\mathbf{A}}(\vec{p}, \vec{q})}{\partial p^\beta} \right) \Big|_{\vec{q}=\vec{p}} \phi_\psi(\vec{p}) d^3p. \end{aligned} \quad (77)$$

(b) If $K_{\mathbf{A}}$ is real, smooth, and has polynomial growth with all of its derivatives of any order, then (75), (76), (77) are also valid for $\psi \in \mathcal{S}(\mathcal{H})$.

(c) If $K_{\mathbf{A}}$ is as in (b), a formula for the second moment holds for $\psi \in \mathcal{S}(\mathcal{H})$ with $\|\psi\| = 1$

$$\int_{\mathbb{R}^3} (x^a)^2 \langle \psi | \mathbf{A}(d^3x) \psi \rangle = \langle \psi | (N^a)^2 \psi \rangle + \langle \psi | \mathbf{K}_a^{\mathbf{A}} \psi \rangle \quad \text{for } a = 1, 2, 3. \quad (78)$$

$\mathbf{K}_a^{\mathbf{A}} \geq 0$ is the (generally unbounded) multiplicative selfadjoint operator

$$\left(\mathbf{K}_a^{\mathbf{A}} \psi \right) (p) := \sqrt{p^0} \frac{\partial}{\partial q_a} \frac{\partial}{\partial p_a} K_{\mathbf{A}}(\vec{q}, \vec{p}) \Big|_{\vec{p}=\vec{q}} \frac{\psi(p)}{\sqrt{p^0}}, \quad \psi \in D(\mathbf{K}_a^{\mathbf{A}}), \quad (79)$$

with $\mathbf{K}_a^{\mathbf{A}} \in \mathfrak{B}(\mathcal{H})$ if $\mathbb{R}^3 \ni \vec{p} \mapsto \frac{\partial}{\partial q_a} \frac{\partial}{\partial p_a} K_{\mathbf{A}}(\vec{q}, \vec{p}) \Big|_{\vec{p}=\vec{q}} \in \mathbb{R}$ is bounded.

(d) If $K_{\mathbf{A}}$ is as in (b), a **modified Heisenberg inequality** holds for $\psi \in \mathcal{S}(\mathcal{H})$ with $\|\psi\| = 1$ (restoring Plank's constant),

$$\Delta_\psi x_{\mathbf{A}}^a \Delta_\psi P_a \geq \frac{\hbar}{2} \sqrt{1 + 4(\Delta_\psi P_a)^2 \langle \psi | \mathbf{K}_a^{\mathbf{A}} \psi \rangle}, \quad a = 1, 2, 3. \quad (80)$$

Above, $\Delta_\psi x_{\mathbf{A}}^a$ is the standard deviation of the probability distribution $\langle \psi | \mathbf{A}(\cdot) \psi \rangle$, $\Delta_\psi P_a$ is the standard deviation of the probability distribution of the a -component of the momentum observable in the state ψ .

(e) Suppose that $K_{\mathbf{A}}$ is not necessarily real but is **finite** according to [9]. In other words, it has the form, for $N < +\infty$ suitable measurable functions u_j

$$K_{\mathbf{A}}(\vec{p}, \vec{q}) = \sum_{j=1}^N \overline{u_j(\vec{p})} u_j(\vec{q}).$$

In addition, assume that the N functions $u_j : \mathbb{R}^3 \rightarrow \mathbb{C}$ are smooth with polynomial growth with all of their derivatives of any order. Then the standard Heisenberg inequality holds in any cases for $\psi \in \mathcal{S}(\mathcal{H})$ with $\|\psi\| = 1$

$$\Delta_\psi x_A^a \Delta_\psi P_a \geq \frac{\hbar}{2}.$$

Proof. See Appendix A □

8 Discussion

The major achievement of this work (Theorem 32) is the result that, if we use POVMs to describe the probability of spatial localization of quantum systems in a sufficiently general way, then *spatial localization*, in the sense of a *spacelike Cauchy localization observable* Definition 21, *implies causality*, in the terms of our *general causal condition* GCC in Definition 26. The physical postulates at the ground of this implication, encapsulated in the notion of spacelike Cauchy localization observable, are the following ones.

- (1) It is supposed that every *spacelike* Cauchy surface can be used to localize the system. In other words, if we fill a spacelike Cauchy surface with a net of detectors, we must find somewhere the quantum system on that 3-space.
- (2) If, for a given spacelike Cauchy localization observable, i.e., for a specific type of detectors, a pair of Cauchy surface coincide in a region, then they share the same detectors therein.
- (3) There is no chance to detect the quantum system in a spatial region with zero measure on a spacelike Cauchy surface.

An observation about the need for condition (2), we named *coherence condition*, is important. As we have seen in Remark (34), condition (2) can be removed from the definition of spacelike Cauchy localization observable and inserted as a further hypothesis of Theorem 32. Within this scheme, the coherence condition would be recovered as a corollary of Theorem 32. In other words, if (1) and (3) are valid for a family of POVMs, then the *coherence condition* is *equivalent* to GCC.

We also proved, as a second achievement in double form (Theorems 49 and 59): The above general notions of localization exist at theoretical level. In fact, we presented two of them for massive Klein-Gordon particles. In the second case the localization observable was constructed in terms of physical quantities of the system (its energy). There is no evident obstruction to construct similar spacelike Cauchy localization observables for other types of particles like fermions considered in [8]. The basic ingredient to construct these observables is a conserved causal current constructed out of the state of the system. It seems plausible that spacelike Cauchy localization observables can be built up also in a more general spacetime (referring to the one-particle structure of quantum field Gaussian states) provided the spacetime is globally hyperbolic. This is because the central technology to produce the former achievement, some

technical results about Cauchy surfaces [3, 4] are at disposal in generic globally hyperbolic spacetimes. On the other hand the explicit structure of spacelike Cauchy localization observable relies upon notions, like conserved currents, which can be generalized to every globally hyperbolic spacetime. The only mathematically delicate issue which deserves attention is the fact that the region of influence Δ' of a Borel set Δ is again Borel or in a natural completion of that σ -algebra.

When coming back to flat Cauchy surfaces, i.e., rest frames of inertial observers, the resulting spatial localization observables show interesting features. Since causality is not violated by the distribution of probability of a Klein-Gordon massive particle, the no-go Hegerfeldt theorem is made harmless. There is a price to pay however: no strictly localized (in bounded spatial sets) probability distributions are permitted. Another interesting fact, already evident in [29], is that the Newton-Wigner operators insist to play some role in this much less naive picture, in spite of the fact that the Hegerfeldt no go results seemed to have ruled out them long time ago. Even if they no longer represent observables, they account for the *timelike* spacetime evolution of the first moment of a massive particle.

A widely open issue is the relation between the constructed POVMs and their decompositions in terms of Kraus operators or quantum instruments. Related to this issue is the fact that the effects of the presented POVMs do not commute even when are localized in causally separated sets. This is an urgent problem when analysing all the construction from the perspective of the local operator algebras theory. These outstanding problems will be investigated elsewhere.

Acknowledgments

V.M. is very grateful to D.P.L.Castrigiano for various remarks, suggestions, and discussions about several issues appearing in this paper. We thank N.Pinamonti and M.Sánchez for helpful discussions on some technical problems, and S. Lill and M. Reddiger for pointing out some relevant literature. This work has been written within the activities of INdAM-GNFM

A Proof of some propositions

Proof of Proposition 20. Every coordinate curve $\mathbb{R} \ni x^0 \mapsto (x^0, \vec{x}) \in \mathbb{R} \times \mathbb{R}^3 \equiv \mathbb{M}$ is smooth timelike and thus it intersects S exactly once for a corresponding value $x^0 = t_S(\vec{x})$, therefore it defines a map $\mathbb{R}^3 \ni \vec{x} \mapsto t_S(\vec{x}) \in \mathbb{R}$. In the rest of the proof write t in place of t_S for shortness. By definition $S \equiv \{(t(\vec{x}), \vec{x}) \mid \vec{x} \in \mathbb{R}^3\}$. Since S is an embedded co-dimension 1 submanifold, it must be locally described as the locus $f(x^0, x^1, x^2, x^3) = 0$ of a smooth map with $\mathbf{g}(df, df) \leq 0$ because the g -normal vectors to df are spacelike or lightlike and thus df is timelike or lightlike. It must be in particular $\partial_{x^0} f \neq 0$. As a consequence of the implicit function theorem, in a neighborhood of every $p \in S$, we can represent S as a smooth map $x^0 = t'(\vec{x})$. Therefore $t(\vec{x}) = t'(\vec{x})$ is locally smooth and thus smooth. Finally, as S is the locus of $g(x^0, x^1, x^2, x^3) = 0$ for $g(x^0, x^1, x^2, x^3) = x^0 - t(\vec{x})$, a normal co-vector to S is $v = dx^0 - \sum_{k=1}^3 \frac{\partial t}{\partial x^k} dx^k$. If S is spacelike, then its normal vector is timelike, written as $\mathbf{g}(v, v) < 0$, is equivalent to $|\nabla t| < 1$. In the other cases v can also be lightlike $\mathbf{g}(v, v) \leq 0$ which is equivalent to $|\nabla t| \leq 1$.

The final statement is now obvious, since the wanted diffeomorphism is the identity map $\mathbb{R}^3 \ni p \mapsto p \in \mathbb{R}^3$ when adopting on S the coordinates \vec{x} . \square

Proof of Proposition 23. We refer to [12] for the definition of completion of a σ -algebra and a positive measure on it. We divide the main proof into some further lemmata.

Lemma 61. *Let $\mu: \mathcal{B} \rightarrow \overline{\mathbb{R}}_+$ be a positive measure on a σ -algebra \mathcal{B} on X , $\overline{\mathcal{B}}^\mu$ be the completion of \mathcal{B} with respect to μ , and let us denote by $\overline{\mu}$ the completion measure on $\overline{\mathcal{B}}^\mu$. Then $\Delta \in \overline{\mathcal{B}}^\mu$ if and only if $\Delta = \Delta_0 \cup N$ with $\Delta_0 \in \mathcal{B}$ and $N \subset N_0 \in \mathcal{B}$ with $\mu(N_0) = 0$. Furthermore, for every decomposition of Δ as above, $\overline{\mu}(\Delta) = \mu(\Delta_0)$.*

Proof. Direct inspection. \square

Lemma 62. *Let $\mu: \mathcal{B} \rightarrow \overline{\mathbb{R}}_+$ be a positive measure on a σ -algebra \mathcal{B} on X and $\mathcal{L} := \overline{\mathcal{B}}^\mu$ be the completion of \mathcal{B} with respect to μ . Let $\nu: \mathcal{B} \rightarrow \overline{\mathbb{R}}_+$ be a positive measure such that $\nu \ll \mu$. Then the completion $\overline{\mathcal{B}}^\nu$ of \mathcal{B} respect to ν satisfies $\mathcal{L} \subset \overline{\mathcal{B}}^\nu$, and there exists a unique positive measure $\tilde{\nu}$ which extends ν on \mathcal{L} and $\tilde{\nu} \ll \overline{\mu}$, where $\overline{\mu}$ is the completion of μ on \mathcal{L} . Finally $\tilde{\nu}(X) = \nu(X)$.*

Proof. (Uniqueness) By Lemma 61, if $\Delta \in \mathcal{L}$ we can decompose $\Delta = \Delta' \cup N$ where $\Delta' \in \mathcal{B}$ and $N \subset \tilde{N} \in \mathcal{B}$ is such that $\mu(\tilde{N}) = 0$. Suppose that $\tilde{\nu}$ extends ν on \mathcal{L} . Sub-additivity and monotony yield

$$\tilde{\nu}(\Delta) \leq \tilde{\nu}(\Delta') + \tilde{\nu}(N) \leq \tilde{\nu}(\Delta') + \tilde{\nu}(\tilde{N}) = \nu(\Delta') + \nu(\tilde{N}) = \nu(\Delta')$$

where, in the last step, we used $\nu \ll \mu$. By monotony $\tilde{\nu}(\Delta) \geq \tilde{\nu}(\Delta') = \nu(\Delta')$, therefore $\tilde{\nu}(\Delta) = \nu(\Delta')$. In summary, an extension has to be absolutely continuous with respect to $\overline{\mu}$. Indeed $\overline{\mu}(\Delta) = 0$ implies $\mu(\Delta') = 0$ and therefore $\tilde{\nu}(\Delta) = \nu(\Delta') = 0$.

(Existence) Let $\overline{\mathcal{B}}^\nu$ be the completion of \mathcal{B} with respect to ν and $\overline{\nu}$ the completion of ν on $\overline{\mathcal{B}}^\nu$. Notice that $\mathcal{L} \subset \overline{\mathcal{B}}^\nu$, indeed $\mu(\tilde{N}) = 0$ implies $\nu(\tilde{N}) = 0$. Therefore $\tilde{\nu} := \overline{\nu}|_{\mathcal{L}}$ is a well defined finite measure which extends ν on \mathcal{L} . Since $X \in \mathcal{B} \cap \overline{\mathcal{B}}^\nu$, we finally have $\tilde{\nu}(X) = \nu(X)$. \square

In the following, if $\mathbf{E}: \mathcal{B} \rightarrow \mathfrak{B}(\mathcal{H})$ is a \mathcal{H} -POVM and $\psi, \phi \in \mathcal{H}$, $\mu_{\psi, \phi}^{\mathbf{E}}(\Delta) := \langle \psi | \mathbf{E}(\Delta) \phi \rangle$ for $\Delta \in \mathcal{B}$. Furthermore $\mu_{\psi}^{\mathbf{E}}(\Delta) := \mu_{\psi, \psi}^{\mathbf{E}}(\Delta)$.

Lemma 63. *Let $\mathbf{A}: \mathcal{B} \rightarrow \mathfrak{B}(\mathcal{H})$ be a normalized \mathcal{H} -POVM and $\mu: \mathcal{B} \rightarrow \overline{\mathbb{R}}_+$ be a positive measure on a σ -algebra \mathcal{B} such that, for every $\psi \in \mathcal{H}$, $\mu_{\psi}^{\mathbf{A}} \ll \mu$. Then there exists a unique normalized \mathcal{H} -POVM $\tilde{\mathbf{A}}$ on $\mathcal{L} := \overline{\mathcal{B}}^\mu$ which extends \mathbf{A} and such that for every $\mu_{\psi}^{\tilde{\mathbf{A}}} \ll \overline{\mu}$.*

Proof. Taking advantage of Lemma 62, for every $\psi \in \mathcal{H}$, we can extend $\mu_{\psi}^{\mathbf{A}}$ to $\widetilde{\mu_{\psi}^{\tilde{\mathbf{A}}}}$ on \mathcal{L} . Then, we define

$$\widetilde{\mu_{\phi, \psi}^{\tilde{\mathbf{A}}}}(\Delta) := \widetilde{\mu_{\phi+\psi}^{\mathbf{A}}}(\Delta) - \widetilde{\mu_{\phi-\psi}^{\mathbf{A}}}(\Delta) - \widetilde{\mu_{\phi-i\psi}^{\mathbf{A}}}(\Delta) + \widetilde{\mu_{\phi+i\psi}^{\mathbf{A}}}(\Delta).$$

For a fixed $\Delta \in \mathcal{L}$ - $\Delta = \Delta' \cup N$ as in the proof of Lemma 62 - the map $\mathcal{H} \times \mathcal{H} \ni (\phi, \psi) \mapsto \widetilde{\mu_{\phi, \psi}^{\tilde{\mathbf{A}}}}(\Delta)$ is sesquilinear and continuous since $\widetilde{\mu_{\phi, \psi}^{\tilde{\mathbf{A}}}}(\Delta) = \mu_{\phi, \psi}^{\mathbf{A}}(\Delta')$. As a consequence, for $\Delta \in \mathcal{L}$,

the map $\mathcal{H} \ni \psi \mapsto \widetilde{\mu_{\phi, \psi}^A}(\Delta) \in \mathbb{C}$ is a linear and continuous functional with norm bounded $\|\psi\|$. A straightforward use of the Riesz lemma gives that there is a unique operator $\tilde{A}(\Delta) : \mathcal{H} \rightarrow \mathcal{H}$ such that $\langle \psi | \tilde{A}(\Delta) \phi \rangle = \widetilde{\mu_{\phi, \psi}^A}(\Delta)$. $\tilde{A}(\Delta)$ is positive because $\mu_{\psi}^{\tilde{A}}$ is positive. By construction $\mathcal{L} \ni \Delta \mapsto \tilde{A}(\Delta) \in \mathcal{B}(\mathcal{H})$ defines a normalized POVM. Normalization arises from $\langle \phi | \tilde{A}(S) \psi \rangle = \widetilde{\mu_{\phi, \psi}^A}(S) = \mu_{\phi, \psi}^A(S) = \langle \phi | \psi \rangle$. Finally, $\mu_{\psi}^{\tilde{A}} = \widetilde{\mu_{\psi}^A} \ll \bar{\mu}$. \square

We can now conclude the proof of the main proposition. If $S \in \mathcal{C}_{\mathbb{M}}^s$, set $\mathcal{B} = \mathcal{B}(S)$, $\mathbf{A} := \mathbf{A}_S : \mathcal{B}(S) \rightarrow \mathfrak{B}(\mathcal{H})$ and $\mu = \nu_S$ in Lemma 63. The map $\mathcal{M}(S) \ni \Delta \mapsto \tilde{A}(\Delta) \in \mathfrak{B}(\mathcal{H})$ defines the wanted extension. It is unique because every extension of the considered type is completely determined by the positive measure $\widetilde{\mu_{\psi}^A}$ which is uniquely determined by μ_{ψ}^A , thus by \mathbf{A} itself. \square

Proof of Proposition 24. We have only to prove that the extended POVMs satisfy the coherence condition (13). Take $\psi \in \mathcal{H}$. If $\Delta \in \mathcal{M}(S)$ then, according to the construction in the proof of Proposition 23, there exists a decomposition $\Delta = \Delta_0 \cup N$ with $\Delta_0 \in \mathcal{B}(S)$ and $N \subset N_0 \in \mathcal{B}(S)$, and finally $\langle \psi | \mathbf{A}_S(N_0) \psi \rangle = 0$, $\langle \psi | \tilde{\mathbf{A}}_S(\Delta) \psi \rangle = \langle \psi | \mathbf{A}_S(\Delta_0) \psi \rangle$. These last two identities do not depend on the decomposition of Δ as $\Delta_0 \cup N$. Since $\Delta_0, N, N_0 \subset \Delta \subset S \cap S'$, we can repeat everything for $\langle \psi | \mathbf{A}_{S'}(\Delta) \psi \rangle$ with the same decomposition of Δ , proving that

$$\langle \psi | \mathbf{A}_{S'}(N_0) \psi \rangle = \langle \psi | \mathbf{A}_S(N_0) \psi \rangle = 0, \quad \langle \psi | \tilde{\mathbf{A}}_{S'}(\Delta) \psi \rangle = \langle \psi | \mathbf{A}_{S'}(\Delta_0) \psi \rangle = \langle \psi | \mathbf{A}_S(\Delta_0) \psi \rangle = \langle \psi | \tilde{\mathbf{A}}_S(\Delta) \psi \rangle$$

where we used the fact that \mathbf{A} satisfies the coherence condition on the Borel sets $\Delta_0, N_0 \subset S \cap S'$ by hypothesis. In particular $\langle \psi | \tilde{\mathbf{A}}_{S'}(\Delta) \psi \rangle = \langle \psi | \tilde{\mathbf{A}}_S(\Delta) \psi \rangle$. Arbitrariness of ψ proves the validity of the coherence condition for the extended POVMs ending the proof. \square

Proof of Proposition 37. Notice that (36) is nothing but (35) when exploiting (34). Evidently, it is sufficient to establish the thesis when S' is a Minkowski time slice and S is a generic smooth Cauchy surface. From now on, we consider a Minkowski coordinate frame x^0, x^1, x^3, x^4 and describe S in terms of a smooth function $x^0 = t(\vec{x})$, taking S' as the time-slice $x^0 = 0$. Define the auxiliary vector field

$$K := \frac{1}{(1+r^2)^2} \partial_{x^0}, \quad \text{where } r := \sqrt{\sum_{k=1}^3 (x^k)^2}.$$

By construction K is conserved, everywhere non-vanishing, timelike, and future-directed. Finally,

$$-K \cdot n_{S'} = \frac{1}{(1+r^2)^2} > 0 \quad \text{and} \quad - \int_{S'} K \cdot n_{S'} d\nu_{S'} = \int_{S'} \omega^K < +\infty. \quad (81)$$

Define B'_R as the open ball in $S' \equiv \mathbb{R}^3$ centered at the origin and with finite radius $R > 0$. Correspondingly $B_R \subset S$ is the open set in S defined by $x^0 = t(\vec{x})$, $\vec{x} \in B'_R$. Let $T_R \subset \mathbb{M}$ be the closed (compact) cylinder with bases $\overline{B_R}$ and $\overline{B'_R}$ and lateral surface parallel to K (i.e. to ∂_{x^0}). We are explicitly assuming that the closures of B'_R and B_R have no intersection for now and we shall treat later the case where the two surfaces intersect. As K is conserved, $d\omega^K = 0$ for

(1) Proposition 36. Furthermore ∂T_R is an orientable smooth submanifold up to zero-measure subsets, we can apply the Poincaré theorem, proving that $\int_{\partial T_R} \omega^K = \int_{T_R} d\omega^K = 0$. Since the lateral surface gives no contribution as easily arises from (30) using the fact that J is parallel to that surface, the identity boils down to

$$\int_{B'_R} \omega^K = \int_{B_R} \omega^K .$$

This identity is valid also when B'_R and B_R have intersection (just because S and S' do and B'_R passes through that intersection). In that case, as R is finite so that $t(\vec{x})$ is bounded for $\vec{x} \in B'_R \subset S' \equiv \mathbb{R}^3$, we can move S' parallelly to ∂_{x^0} till to another x^0 -slice S'' in the past of S' , in order that B_R and B''_R (the projection of B'_R onto S'') do not meet. By construction, using the above argument once again for the relevant submanifolds with orientable boundary, since now the bases do not touch each other,

$$\int_{B''_R} \omega^K = \int_{B'_R} \omega^K ,$$

and also

$$\int_{B''_R} \omega^K = \int_{B_R} \omega^K .$$

In summary

$$\int_{B'_R} \omega^K = \int_{B_R} \omega^K .$$

is valid even if B_R and B'_R have intersection.

Taking $R \rightarrow +\infty$, a direct application of the monotone convergence theorem (as the integrals can be written as Lebesgue integrals in \mathbb{R}^3 of non negative functions according to (31) proves that

$$0 \leq \int_{S'} \omega^K = \int_S \omega^K < +\infty , \quad (82)$$

where we also used (81).

To go on, define $Y := J + K$ noticing that this smooth conserved vector field is everywhere timelike and future directed. Since it is also bounded, its flow is global and defines a smooth one-parameter group of diffeomorphisms $\{\phi_s^Y\}_{s \in \mathbb{R}}$ of \mathbb{M} . Finally, the integral curves of Y are future directed and inextendible. In fact, suppose that $\gamma(s) \rightarrow q \in \mathbb{M}$ for $s \rightarrow +\infty$, i.e., the curve is not future inextendible (the same argument applies for $s \rightarrow -\infty$). If $Y^k(q) \neq 0$ for some $k = 0, 1, 2, 3$, then it happens in a connected neighborhood U of q where the sign of $Y^k = \frac{dx^k}{ds}$ is therefore fixed. There we can use $z = x^k$ to parametrize γ obtaining $s(z) = \int_0^z \frac{dx^k}{Y^k(\gamma(x^k))}$. However, since s varies till $+\infty$ when γ approaches q in U , then $Y^k(\gamma(x^k))$ must vanish as $x^k \rightarrow x^k(q)$ contrarily to the hypothesis. Hence all components of Y must vanish at q but this was excluded *a priori*.

As a consequences of these properties, the map

$$\Phi : \mathbb{R} \times S' \equiv \mathbb{R} \times \mathbb{R}^3 \ni (s, p) \mapsto \phi_s^Y(p) \in \mathbb{M}$$

turns out to be a diffeomorphism as well. In fact, Φ is injective because $\phi_s^Y(p) = \phi_{s'}^Y(p')$ implies $p = \phi_{s'-s}^Y(p')$ but this would mean that the integral curve of Y through p touches twice S' , and this is not possible unless $s = s'$ and thus $p = p'$; surjectivity of Φ follows from the fact that, if $q \in \mathbb{M}$ there is an integral line of Y through q at $s = 0$ and, it being timelike and inextendible, it must intersect S' at some q' for $s = T_q$ because S' is a Cauchy surface. Therefore $\Phi(-T_q, q') = q$; finally $d\Phi \neq 0$ everywhere, because (a) $d\Phi(s, p)e_0 = Y(\phi_s^Y(p))$ and (e_0, e_1, e_2, e_3) denoting the canonical basis of $\mathbb{R} \times \mathbb{R}^3 \equiv \mathbb{R} \times S'$ (b) $d\Phi(s, p)e_k = d\phi_s^Y(p)e_k$ defines a basis of the tangent space at $\phi_s^Y(p)$ of the embedded 3D submanifold $\phi_s^Y(S')$ ¹⁷ that is transverse to $Y(\phi_s^Y(p)) = d\phi_s^Y Y(p)$ by construction. In summary, $d\Phi(s, p)$ sends the canonical basis of $\mathbb{R}^4 = \mathbb{R} \times \mathbb{R}^3 \equiv \mathbb{R} \times S'$ to a basis of $T_{\phi_s(p)}\mathbb{M}$ and it is therefore bijective. The situation is identical to the one of the vector K , with the only difference that we have replaced K for Y and x^0 for the global parameter s . Now s, x^1, x^2, x^3 – where the latter three coordinates are taken on S' – define a global chart on \mathbb{M} . In these coordinates, the action of ϕ^Y is trivial $\phi_t^Y : \mathbb{R}^4 \ni (s, x^1, x^2, x^3) \mapsto (s + t, x^1, x^2, x^3) \in \mathbb{R}^4$. Taking advantage of the diffeomorphism Φ , one easily proves the following facts whose details are left to the reader. If $B'_R \subset S' \equiv \mathbb{R}^3$ is as before an open ball of radius $R > 0$, $\Gamma_R := \Phi(\mathbb{R} \times \overline{B'_R})$ is a smooth 4-manifold with orientable boundary embedded in \mathbb{M} . The boundary $\partial\Gamma_R$ is made of the integral lines of Y exiting $\partial B'_R$. We extract from Γ_R the compact cylinder T_R with bases given by the closures of B'_R and the closure of the smooth embedded submanifold

$$B_R := S \cap \Gamma_R = S \cap \{\phi_t(B'_R) \mid t \in \mathbb{R}\}$$

(which is not a ball in general now!). As before we assume that the closures of B_R and $B'_{R'}$ have no intersection and the case of non empty intersection can be treated exactly as before in the representation $\mathbb{R} \times \mathbb{R}^3$ where $s \in \mathbb{R}$ and $\mathbb{R}^3 \equiv S'$. Notice that $B_R \subset B'_{R'} \subset S$ if $R < R'$ by construction and also

$$\cup_{R>0} B_R = S. \quad (83)$$

This identity is valid because, for every $p \in S$, there is $R' > 0$ such that $p \in B'_{R'}$. (In fact, as before, there is an integral curve of Y passing through p and, since this curve is timelike and inextendible, there must be $p' \in S'$ such that $\phi_s^Y(p') = p$ for some s because S' is Cauchy. Since the union of the sets B'_R covers S' , then $p' \in B'_{R'}$ for some $R' > 0$ sufficiently large and so $p \in B'_{R'}$.) ∂T_R is an orientable smooth submanifold up to zero-measure sets and we can apply the Poincaré theorem as before, obtaining

$$\int_{B'_R} \omega^Y = \int_{B_R} \omega^Y,$$

where we have disregarded the contribution of the lateral surface because it is made of integral lines of Y itself and no contribution to the boundary integral arises according to (30). Taking

¹⁷Indeed, $\phi_s^Y : \mathbb{M} \rightarrow \mathbb{M}$ is a diffeomorphism and thus it transforms the embedded submanifold $S' \equiv \mathbb{R}^3$ transverse to $Y|_{S'}$ into the embedded submanifold $\phi_s^Y(S')$ transverse to $d\phi_s^Y|_{S'}Y = Y|_{\phi_s^Y(S')}$, in particular, $d\phi_s^Y$ bijectively transforms the tangent spaces of these manifolds accordingly.

the limit for $R \rightarrow +\infty$ on both sides, the monotone convergence theorem (taking positivity of integrands into account according to (2) Proposition 36 and (83)) yields

$$\int_{S'} \omega^{K+J} = \int_S \omega^{K+J}.$$

Namely,

$$\int_{S'} \omega^K + \omega^J = \int_S \omega^K + \omega^J.$$

On the ground of the validity of (82), this identity boils down to

$$0 \leq \int_{S'} \omega^J = \int_S \omega^J \leq +\infty.$$

In particular both integrals converge or diverge simultaneously. \square

Proof of Lemma 48. The Cauchy-Schwartz inequality implies $|\Lambda(x, y)| \leq C\|x\|\|y\|$ if $x, y \in D$. For a fixed $x \in D$ in $\Lambda(x, y)$, the linear map on the dense subspace $D \ni y \mapsto \Lambda(x, y) \in \mathbb{C}$ is therefore bounded and thus it uniquely extends to the whole Hilbert space to a bounded linear map. Riesz' lemma implies that $\Lambda(x, y) = \langle A_x | y \rangle$, for some $A_x \in \mathcal{K}$. $\Lambda(x, y) = \langle A_x | y \rangle$ also implies that the map $D \ni x \mapsto A_x$ is linear: $A_x := A'x$ for some $A' : D \rightarrow \mathcal{H}$. Moreover

$$\|A'x\| = \sup_{\|y\|=1} |\Lambda(x, y)| \leq C\|x\|,$$

so that A' is also bounded by C and therefore uniquely extends to the whole \mathcal{K} to an operator with the same norm. The adjoint $A := A'^* \in \mathfrak{B}(\mathcal{K})$ is the wanted operator because $\langle x | Ay \rangle = \langle A'x | y \rangle = \Lambda(x, y)$. Notice that $\|A\| = \|A'^*\| = \|A'\| \leq C$. Moreover $A \geq 0$ because $\langle x | Ax \rangle = \Lambda(x, x) \geq 0$ on D and the inequality extends to the whole Hilbert space by continuity since A is bounded and D dense. Uniqueness is now trivial, an operator $B \in \mathfrak{B}(\mathcal{K})$ with $\Lambda(x, x) = \langle x | Bx \rangle$ for $x \in D$, by polarization coincides there with A . Since D is dense, $B = A$. \square

Proof of Lemma 50. Passing to a Minkowski chart adapted to n_Σ , the right-hand side of (59) is a finite linear combination of integrals of the form

$$I := \int_{\mathbb{R}^3} d^3 p \overline{f(\vec{p})} \int_{\Delta} d^3 x \int_{\mathbb{R}^3} d^3 q \frac{e^{i(\vec{q}-\vec{p}) \cdot \vec{x}}}{(2\pi)^3} g(\vec{q}) = \int_{\mathbb{R}^3} d^3 p \overline{f(\vec{p})} \int_{\mathbb{R}^3} d^3 x \chi_{\Delta}(\vec{x}) \int_{\mathbb{R}^3} d^3 q \frac{e^{i(\vec{q}-\vec{p}) \cdot \vec{x}}}{(2\pi)^3} g(\vec{q})$$

where $f, g \in (\mathbb{R}^3)$ and the exponentials containing x^0 are embodied in these functions. Therefore, proving the thesis for I is enough for ending the proof of the theorem. We have,

$$I = \int_{\mathbb{R}^3} d^3 p \overline{f(\vec{p})} \int_{\mathbb{R}^3} d^3 x \chi_{\Delta}(\vec{x}) \frac{e^{-i\vec{p} \cdot \vec{x}}}{(2\pi)^{3/2}} \int_{\mathbb{R}^3} d^3 q \frac{e^{i\vec{q} \cdot \vec{x}}}{(2\pi)^{3/2}} g(\vec{q})$$

$$= \int_{\mathbb{R}^3} d^3 p \overline{f(\vec{p})} \int_{\mathbb{R}^3} d^3 x \frac{e^{-i\vec{p}\cdot\vec{x}}}{(2\pi)^{3/2}} \chi_{\Delta}(\vec{x}) \hat{g}(\vec{x}) = \int_{\mathbb{R}^3} d^3 x \overline{\hat{f}(\vec{x})} \chi_{\Delta}(\vec{x}) \hat{g}(\vec{x}).$$

Above, $\chi_{\Delta} \cdot \hat{g} \in L^1(\mathbb{R}^3, d^3 x) \cap L^2(\mathbb{R}^3, d^3 x)$ because χ_{Δ} is bounded and \hat{g} Schwartz, and in the last identity we exploited the Plancherel theorem in L^2 . From the very definition of Fourier transform, we finally have

$$I = \int_{\mathbb{R}^3} d^3 x \chi_{\Delta}(\vec{x}) \int_{\mathbb{R}^3} d^3 p \int_{\mathbb{R}^3} d^3 q \frac{e^{i(\vec{q}-\vec{p})\cdot\vec{x}}}{(2\pi)^3} \overline{f(\vec{p})} g(\vec{q}) = \int_{\Delta} d^3 x \int_{\mathbb{R}^3} d^3 p \int_{\mathbb{R}^3} d^3 q \frac{e^{i(\vec{q}-\vec{p})\cdot\vec{x}}}{(2\pi)^3} \overline{f(\vec{p})} g(\vec{q}).$$

Last identity concludes the proof. \square

Proof of Proposition 60 (a) and (b). We adopt throughout the easier notation $\phi := \phi_{\psi}$. We start by observing that the function $\mathbb{R}^6 \ni (\vec{p}, \vec{q}) \mapsto F(\vec{p}, \vec{q}) := \overline{\phi(\vec{p})} K_{\Lambda}(\vec{p}, \vec{q}) \phi(\vec{q})$ is $\mathcal{S}(\mathbb{R}^6)$ in both cases (a) and (b), and thus

$$\mathbb{R}^3 \ni \vec{x} \mapsto f(\vec{x}) := \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} d^3 p d^3 q \frac{e^{i(\vec{q}-\vec{p})\cdot\vec{x}}}{(2\pi)^3} F(\vec{p}, \vec{q})$$

is in $\mathcal{S}(\mathbb{R}^3)$ as it can be proved by direct inspection. Notice that, in the case (b), (75) must be valid also for $\phi \in \mathcal{S}(\mathbb{R}^3)$ since, for a fixed $\phi \in \mathcal{S}(\mathbb{R}^3)$, there is a sequence of functions $C_c^{\infty}(\mathbb{R}^3) \ni \phi_n \rightarrow \phi$ pointwise and $|\phi_n| \leq \phi$. Hence we can take the limit on both sides of (75) by using continuity of $A(\Delta)$ on the left-hand side and the dominated convergence theorem on the right-hand side.

We pass to compute the various moments of the probability measure $\langle \psi | A(\cdot) \psi \rangle$ taking advantage of the theory of Fourier transformation of distributions in $\mathcal{S}'(\mathbb{R}^n)$. If $\alpha := (\alpha_1, \alpha_2, \alpha_3)$ is a multi index, so that $x^{\alpha} := (x^1)^{\alpha_1} (x^2)^{\alpha_2} (x^3)^{\alpha_3}$, we can write

$$\int_{\mathbb{R}^3} x^{\alpha} \langle \psi | A(d^3 x) \psi \rangle = \int_{\mathbb{R}^3} x^{\alpha} f(x) \frac{d^3 x}{(2\pi)^{\frac{3}{2}}} = \left\langle \frac{1}{(2\pi)^{\frac{3}{2}}}, M^{\alpha} f \right\rangle$$

where M^{α} is the multiplicative operator with x^{α} and $f \in \mathcal{S}(\mathbb{R}^3)$ is defined above. Changing variables to $\vec{u} := \vec{q} - \vec{p}$ and $\vec{v} := \vec{q}$ we have,

$$f(\vec{x}) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int_{\mathbb{R}^3} e^{i\vec{u}\cdot\vec{x}} \left(\int_{\mathbb{R}^3} F(\vec{u} - \vec{v}, \vec{v}) d^3 v \right) d^3 u$$

with is the inverse Fourier transform, always in $\mathcal{S}(\mathbb{R}^3)$, of

$$G(\vec{u}) := \frac{1}{(2\pi)^{\frac{3}{2}}} \int_{\mathbb{R}^3} e^{-i\vec{u}\cdot\vec{x}} f(\vec{x}) d^3 x = \int_{\mathbb{R}^3} F(\vec{u} - \vec{v}, \vec{v}) d^3 v.$$

Therefore

$$\int_{\mathbb{R}^3} x^{\alpha} \langle \psi | A(d^3 x) \psi \rangle = \langle \mathcal{F}^{-1} \delta_0, i^{|\alpha|} \mathcal{F}^{-1} \partial^{\alpha} G \rangle = i^{|\alpha|} \langle \delta_0, \partial^{\alpha} G \rangle = i^{|\alpha|} \partial^{\alpha} G(0),$$

$$\partial^\alpha G(0) = \int_{\mathbb{R}^3} \frac{\partial^{|\alpha|}}{\partial p^\alpha} F(\vec{v} - \vec{u}, \vec{v})|_{\vec{u}=0} d^3 v = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \int_{\mathbb{R}^3} \frac{\partial^{|\alpha|-|\beta|} \bar{\phi}}{\partial p^{\alpha-\beta}}(\vec{p}) \frac{\partial^{|\beta|} K_A}{\partial p^\beta} \Big|_{\vec{q}=\vec{p}} \phi(\vec{p}) d^3 p.$$

To go on, defining $l(\vec{p}, \vec{q}) := (\vec{p}, \vec{p})$, since $K_A(\vec{p}, \vec{p}) = 1$, we have that $0 = \partial^\alpha (K_A \circ l) = (\partial^\alpha K_A) \circ l$ for $|\alpha| = 1$. Therefore we obtain the general expression for the α -th moment (77),

$$\int_{\mathbb{R}^3} x^\alpha \langle \psi | A(d^3 x) \psi \rangle = \langle \psi | N^\alpha \psi \rangle + \sum_{1 < |\beta|, \beta \leq \alpha} \binom{\alpha}{\beta} \int_{\mathbb{R}^3} \left(i^{|\alpha|-|\beta|} \frac{\partial^{|\alpha|-|\beta|} \bar{\phi}}{\partial p^{\alpha-\beta}} \right) (\vec{p}) \left(i^{|\beta|} \frac{\partial^{|\beta|} K_A}{\partial p^\beta} \right) \Big|_{\vec{q}=\vec{p}} \phi(\vec{p}) d^3 p.$$

In particular, for $|\alpha| = 1$, we find (76)

$$\int_{\mathbb{R}^3} x^a \langle \psi | A(d^3 x) \psi \rangle = \langle \psi | N^a \psi \rangle \quad a = 1, 2, 3.$$

Every other selfadjoint operator B in \mathcal{H} which satisfies the identity above must also satisfy (from polarization and density)

$$B|_{\mathcal{D}(\mathcal{H})} = N^a|_{\mathcal{D}(\mathcal{H})}$$

(also replacing $\mathcal{D}(\mathcal{H})$ for $\mathcal{S}(\mathcal{H})$). Since $\mathcal{D}(\mathcal{H})$ (resp. $\mathcal{S}(\mathcal{H})$) is a core for N^a :

$$N^a = \overline{N^a|_{\mathcal{D}(\mathcal{H})}} = \overline{B|_{\mathcal{D}(\mathcal{H})}} \subset \overline{B} = B.$$

Since a selfadjoint operator (here N^a) is maximally symmetric, it must hold $N^a = B$. This ends the proof of (a) and (b).

We can pass to prove (c) and (d). For $|\alpha| = 2$, the formula above for the α -th moment yields

$$\int_{\mathbb{R}^3} x^\alpha \langle \psi | A(d^3 x) \psi \rangle = \langle \psi | N^\alpha \psi \rangle + \int_{\mathbb{R}^3} \bar{\phi}(\vec{p}) \left(i^2 \frac{\partial^{|\alpha|} K_A}{\partial p^\alpha} \right) \Big|_{\vec{q}=\vec{p}} \phi(\vec{p}) d^3 p.$$

However, taking advantage of $\frac{\partial^{|\alpha|} K_A}{\partial p^\alpha} \circ l = -\frac{\partial^{|\alpha|} K_A}{\partial q^{\alpha_1} \partial p^{\alpha_2}} \circ l$ for $|\alpha| = 2$ and when only a component of α does not vanish, we can rephrase the found result as

$$\int_{\mathbb{R}^3} (x^a)^2 \langle \psi | A(d^3 x) \psi \rangle = \langle \psi | (N^a)^2 \psi \rangle + \int_{\mathbb{R}^3} d^3 p \bar{\phi}(\vec{p}) \phi(\vec{q}) \frac{\partial}{\partial q_a} \frac{\partial}{\partial p_a} K_A(\vec{q}, \vec{p}) \Big|_{\vec{p}=\vec{q}}.$$

The multiplicative operator K_a^A defined in (79), here viewed in $L^2(\mathbb{R}^3, d^3 p)$, defined by the function $\frac{\partial}{\partial p_a} \frac{\partial}{\partial q_a} K_A(\vec{p}, \vec{q}) \Big|_{\vec{p}=\vec{q}}$ is a selfadjoint operator which is a spectral function of the four momentum operator P as this function is real (see e.g. [28]). The domain of the operator trivially includes $C_c^\infty(\mathbb{R}^3)$, but also $\mathcal{S}(\mathbb{R}^3)$ when $K_A(\vec{p}, \vec{q})$ has polynomial growth with all of its derivatives. Evidently K_a^A is also bounded, and thus in $\mathfrak{B}(\mathcal{H})$, if the function $\frac{\partial}{\partial p_a} \frac{\partial}{\partial q_a} K_A(\vec{p}, \vec{q}) \Big|_{\vec{p}=\vec{q}}$ is bounded. Finally the operator K_a^A is positive just because

$$\frac{\partial}{\partial p_a} \frac{\partial}{\partial q_a} K_A(\vec{p}, \vec{q}) \Big|_{\vec{p}=\vec{q}} \geq 0.$$

This can be proved as follows. Consider a real smooth mollificator (with a common compact support) such that $\psi_n(\vec{x} - \vec{x}_0) \rightarrow \delta(\vec{x} - \vec{x}_0)$ weakly for $n \rightarrow +\infty$. As K_A is positive definite, we have

$$0 \leq \int_{\mathbb{R}^3} d^3 p \int_{\mathbb{R}^3} d^3 q \partial_{p^a} \psi_n(\vec{p} - \vec{k}) \partial_{q^a} \psi_n(\vec{q} - \vec{k}) K_A(\vec{p}, \vec{q}) \rightarrow \left. \frac{\partial}{\partial p_a} \frac{\partial}{\partial q_a} K_A(\vec{p}, \vec{q}) \right|_{\vec{p}=\vec{q}=\vec{k}}$$

uniformly as a function of \vec{k} because K_A is continuous (Proposition 4.21 [5]) so that the resulting function of \vec{k} is non-negative as well. In summary,

$$\begin{aligned} (\Delta_\psi x^a)^2 &:= \int_{\mathbb{R}^3} (x^a)^2 \langle \psi | A(d^3 x) \psi \rangle - \left(\int_{\mathbb{R}^3} x^a \langle \psi | A(d^3 x) \psi \rangle \right)^2 \\ &= \langle \psi | (N^a)^2 \psi \rangle - \langle \psi | N^a \psi \rangle^2 + \langle \psi | K_A^A \psi \rangle = (\Delta_\psi N^a)^2 + \langle \psi | K_A^A \psi \rangle. \end{aligned}$$

Multiplying both sides of the found identity with $(\Delta_\psi P_a)^2$ and taking the standard Heisenberg inequality into account we get the thesis:

$$(\Delta_\psi x_A^a \Delta_\psi P_a)^2 \geq \frac{\hbar^2}{4} + \hbar^2 (\Delta_\psi P_a)^2 \langle \psi | K_A^A \psi \rangle$$

where the second factor \hbar^2 arises when restoring the Planck constant in the exponentials $e^{i(\vec{q}-\vec{p})\cdot\vec{x}/\hbar}$ so that the representation of N^a on $\mathcal{S}(\mathbb{R}^3)$ in the $L^2(\mathbb{R}^3, d^3 p)$ space is $i\hbar \frac{\partial}{\partial p_a}$.

(e) $\mathcal{H}_N := \bigoplus_{j=1}^N \mathcal{H}$ is the complex Hilbert space of vectors of measurable complex valued functions $\Psi := (\psi_1, \dots, \psi_N)$ equipped with the Hilbert space structure arising from the Hermitian scalar product

$$\langle \Psi | \Psi' \rangle_N := \sum_{j=1}^N \int_{\mathbb{R}^3} \overline{\psi_j(\vec{p})} \psi'_j(\vec{p}) d^3 p.$$

If $\psi \in \mathcal{S}(\mathcal{H})$, define $\Phi_\psi := N^{-1/2}(\phi_\psi u_1, \dots, \phi_\psi u_N)$. Each component is in $\mathcal{S}(\mathbb{R}^3)$ in our hypotheses. Finally, using $K_A(\vec{p}, \vec{p}) = 1$

$$\langle \Phi_\psi | \Phi_\psi \rangle_N = \frac{1}{N} \sum_{j=1}^N \int_{\mathbb{R}^3} \overline{\phi_\psi(\vec{p})} u_j(\vec{p}) u_j(\vec{p}) \phi_\psi(\vec{p}) d^3 p = \frac{1}{N} \sum_{j=1}^N \int_{\mathbb{R}^3} \overline{\phi_\psi(\vec{p})} K_A(\vec{p}, \vec{p}) \phi_\psi(\vec{p}) d^3 p = \langle \psi | \psi \rangle. \quad (84)$$

With the same procedure as in the proof of (c) and (d), one sees that, for every constant $c \in \mathbb{R}$

$$\begin{aligned} \int_{\mathbb{R}^3} (x^a - c)^2 \langle \psi | A(d^3 x) \psi \rangle &= \frac{1}{N} \sum_{j=1}^N \int_{\mathbb{R}^3} \overline{\phi_\psi(\vec{p})} u_j(\vec{p}) (i\partial_{p_a} - c)^2 \phi_\psi(\vec{p}) u_j(\vec{p}) d^3 p \\ &= \langle \Phi_\psi | (i\partial_{p_a} - c)^2 \Phi_\psi \rangle_N = \langle (i\partial_{p_a} - c) \Phi_\psi | (i\partial_{p_a} - c) \Phi_\psi \rangle_N = \| (i\partial_{p_a} - c) \Phi_\psi \|^2 : \end{aligned}$$

where we have used the fact that $i\partial_{p_a} - c$, acting componentwise, is symmetric on $\oplus_{j=1}^N \mathcal{S}(\mathbb{R}^3)$. Similarly

$$\langle \psi | (P_a - bI)^2 \psi \rangle = \frac{1}{N} \sum_{j=1}^N \int_{\mathbb{R}^3} \overline{\phi_\psi(\vec{p})} u_j(\vec{p}) (p_a - b)^2 \phi_\psi(\vec{p}) u_j(\vec{p}) d^3 p = \|(p_a - d)\Phi_\psi\|^2.$$

Where the multiplicative operator $p_a - d$, acting componentwise, is symmetric on $\oplus_{j=1}^N \mathcal{S}(\mathbb{R}^3)$. Cauchy-Schwarz inequality and (84) yield

$$\begin{aligned} \|(i\partial_{p_a} - c)\Phi_\psi\| \|(p_a - d)\Phi_\psi\| &\geq | \langle (i\partial_{p_a} - c)\Phi_\psi | (p_a - d)\Phi_\psi \rangle_N | = | \langle \Phi_\psi | (i\partial_{p_a} - c)(p_a - d)\Phi_\psi \rangle_N | \\ &\geq | \text{Im} \langle \Phi_\psi | (i\partial_{p_a} - c)(p_a - d)\Phi_\psi \rangle_N | = \frac{1}{2} | \langle \Phi_\psi | [i\partial_{p_a}, p_a] \Phi_\psi \rangle_N | = \frac{1}{2} \langle \Phi_\psi | \Phi_\psi \rangle_N = \frac{1}{2} \langle \psi | \psi \rangle. \end{aligned}$$

Restoring \hbar , we have found that, for $\|\psi\| = 1$

$$\sqrt{\int_{\mathbb{R}^3} (x^a - c)^2 \langle \psi | \mathbf{A}(d^3 x) \psi \rangle} \sqrt{\langle \psi | (P_a - bI)^2 \psi \rangle} \geq \frac{\hbar}{2}$$

This is the thesis when replacing c for $\int_{\mathbb{R}^3} x^a \langle \psi | \mathbf{A}(d^3 x) \psi \rangle$ and b for $\langle \psi | P_a \psi \rangle$. \square

Proof of Theorems 44 and 57. Those theorems are easy consequences of Proposition 60. Both $M_\Sigma^n(\Delta)$ and $T_\Sigma^g(\Delta)$, for g real and smooth, have the expression (75) if Σ coincides with the slice at $x^0 = 0$ of a Minkowski chart (see (42) and (59) for $x^0 = 0$). In details,

$$K_{T^g}(\vec{q}, \vec{p}) := \frac{(q^0 + p^0)g(\vec{q} \cdot \vec{p} - q^0 p^0)}{2\sqrt{q^0 p^0}}$$

and

$$K_{M_\Sigma^n}(\vec{q}, \vec{p}) := \frac{p \cdot n q \cdot n_\Sigma + q \cdot n p \cdot n_\Sigma - n \cdot n_\Sigma (p \cdot q + m^2)}{2q \cdot n p \cdot n}.$$

These kernels are positive definite, respectively, in view of Def. 39 and Thm. 53.

The kernels $K_{M_\Sigma^n}(\vec{q}, \vec{p})$, and $K_{T_\Sigma^g}(\vec{q}, \vec{p})$ when g has the form (40) and convex combinations of these functions, have polynomial growth with all of their derivatives, so the case (b) of the proposition applies to these cases.

It is not difficult to prove in particular that the functions of \vec{p} given by $\left. \frac{\partial}{\partial q_a} \frac{\partial}{\partial p_a} K_{M_\Sigma^n}(\vec{q}, \vec{p}) \right|_{\vec{p}=\vec{q}}$ and $\left. \frac{\partial}{\partial q_a} \frac{\partial}{\partial p_a} K_{T_\Sigma^g}(\vec{q}, \vec{p}) \right|_{\vec{p}=\vec{q}}$ are bounded (in the second case for every smooth g , it is not necessary the form (40)) and therefore the corresponding operator in (79) is a positive selfadjoint operator in $\mathfrak{B}(\mathcal{H})$. \square

B Conditions for positive-definite kernels on \mathbb{R}^n

We prove here a useful property of positive definite kernels.

Proposition 64. *Let $K : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{C}$ be continuous, the following facts are equivalent.*

(a) *Every $N \times N$ matrix $[K(k_i, k_j)]_{i,j=1,\dots,N}$ for every choice of $k_1, \dots, k_N \in \mathbb{R}^n$ and $N = 1, 2, \dots$ are positive semidefinite (in particular Hermitian);*

(b) *K is positive definite according to (39);*

$$\sum_{i,j=1}^N \overline{c_i} c_j K(k_i, k_j) \geq 0, \quad \forall \{c_j\}_{j=1,\dots,N} \subset \mathbb{C}, \forall \{k_j\}_{j=1,\dots,N} \subset X, \forall N = 1, 2, \dots; \quad (39)$$

(c) *the bilinear functional induced by K on $C_c(\mathbb{R}^n)$ is positive:*

$$\int_{\mathbb{R}^n \times \mathbb{R}^n} \overline{f(p)} K(p, q) f(q) d^n p d^n q \geq 0, \quad \forall f \in C_c(\mathbb{R}^n). \quad (85)$$

Proof. Evidently (a) and (b) are equivalent, so we only prove that (b) and (c) are equivalent. We start by establishing that K positive definite implies (85). Define a compactly supported continuous function on \mathbb{R}^{2n} as $g(p, q) = \overline{f(p)} K(p, q) f(q)$ and let $Q \subset \mathbb{R}^{2n}$ be an open, $2n$ -square such that $\text{supp}(g) \subset Q$. Then, for any $N \in \mathbb{N}$ there exist a family of N^{2n} pairwise-disjoint open $2n$ -squares $Q_i^N \subset Q$, $i = 1, \dots, N^{2n}$, with common (Lebesgue) measure $|Q_i^N| = \frac{|Q|}{N^{2n}}$ and $\overline{Q} = \bigcup_{i=1}^{N^{2n}} \overline{Q_i^N}$. Choosing $x_i^N \in Q_i^N$ with $i = 1, \dots, N^{2n}$ for every given N , we can define the family of simple functions $g_N(x) = \sum_{i=1}^{N^{2n}} g(x_i^N) \chi_{Q_i^N}(x)$. Since $\text{diam}(Q_i^N) = \sqrt{2d} \frac{|Q|^{\frac{1}{2d}}}{N}$, uniform continuity of g on the compact \overline{Q} implies that for $\epsilon > 0$ there is N_ϵ such that $|g(x_i^N) - g(x)| < \epsilon$ if $x \in Q_i^N$ and $N > N_\epsilon$ and for $i = 1, \dots, N^{2n}$. Therefore, if $N > N_\epsilon$ and $x \in Q$, $|g_N(x) - g(x)| \leq \sum_{i=1}^{N^{2n}} |g(x_i^N) - g(x)| \chi_{Q_i^N}(x) < \epsilon$, so that g_N converges to g on Q uniformly and also in L^1 since $|Q| < +\infty$. The proof is over because

$$\begin{aligned} \int_{\mathbb{R}^n \times \mathbb{R}^n} \overline{f(p)} K(p, q) f(q) d^n p d^n q &= \int_Q g d^n p d^n q = \lim_{N \rightarrow +\infty} \int_Q g_N d^n p d^n q \\ &= \lim_{N \rightarrow +\infty} \sum_{i=1}^{N^{2n}} \overline{f(p_i^N)} K(p_i^N, q_i^N) f(q_i^N) \frac{|Q_N|}{N^{2n}} \geq 0 \end{aligned}$$

We eventually prove that (85) entails that the continuous kernel K is positive definite. To this end, we can choose $c_1, \dots, c_N \in \mathbb{C}$, $p_1, \dots, p_N \in \mathbb{R}^n$ and a $L \in (1, +\infty)$ parametrized family of functions

$$f_L := \sum_{k=1}^N c_k L h(L(p - p_k)) \in C_c(\mathbb{R}^n)$$

where $h : \mathbb{R}^n \rightarrow [0, +\infty)$ is continuous, compactly supported, $h(0) = 1$, and $\int_{\mathbb{R}^n} h d^n x = 1$. In this case, it is easy to prove that

$$\int_{\mathbb{R}^n \times \mathbb{R}^n} \overline{f_L(p)} g(p, q) f_L(q) d^n p d^n q \rightarrow \sum_{k,h=1}^N \overline{c_k} c_h g(p_k, p_h) \quad \text{as } L \rightarrow +\infty$$

for every continuous function $g : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{C}$. Replacing the function f for f_L in (85), the limit as $L \rightarrow +\infty$ implies (39), ending the proof. \square

Competing Interests Declaration. All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter or materials discussed in this manuscript.

Data Availability Declaration. All data generated or analyzed during this study are contained in this document.

References

- [1] Christian Bär, Nicolas Ginoux, Frank Pfäffle. *Wave Equations on Lorentzian Manifolds and Quantization*, ESI Lectures in Mathematics and Physics, EMS Publishing House (2007)
- [2] C. Beck. *Localization Local Quantum Measurement and Relativity*, Dissertation an der Fakultät für Mathematik, Informatik und Statistik der Ludwig-Maximilian-Universität München, 2020
- [3] A. N. Bernal, M. Sánchez. *On Smooth Cauchy Hypersurfaces and Geroch's Splitting Theorem*. Commun. Math. Phys. 243, 461–470 (2003)
- [4] A. N. Bernal, M. Sánchez. *Further Results on the Smoothability of Cauchy Hypersurfaces and Cauchy Time Functions* Letters in Mathematical Physics (2006) 77:183–197
- [5] H. Brezis. *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer (2010)
- [6] P. Busch. *Unsharp localization and causality in relativistic quantum theory*. J. Phys. A: Math. Gen. 32, 37 (1999), 6535
- [7] P. Busch, P. Lahti, J.-P. Pellonpää, K. Ylínen. *Quantum Measurement*, Springer 2016
- [8] D.P.L. Castrigiano. *Dirac and Weyl Fermions - The Only Causal Systems* (2017). arXiv:1711.06556

- [9] D.P.L. Castrigiano. *Causal Localizations of the Massive Scalar Boson*. Letters in Mathematical Physics (2024) 114:2
- [10] D.P.L. Castrigiano. *private communication*. January 2024.
- [11] D.P.L. Castrigiano, A.D. Leiseifer. *Causal Localizations in Relativistic Quantum Mechanics*, J. Math. Phys. 56, 072301 (2015)
- [12] D. Cohn, *Measure Theory*. Birkhäuser (1980)
- [13] N. Drago and V. Moretti. *The notion of observable and the moment problem for *-algebras and their GNS representations*, Lett. Math. Phys. 110(7), (2020), 1711-1758
- [14] L. C. Evans and R. F. Gariepy. *Measure Theory and Fine Properties of Functions* Revised Edition, CRC Press Taylor & Francis Group (2015)
- [15] C. J. Fewster and R. Verch. *Quantum Fields and Local Measurements*, Commun. Math. Phys. **378**, 851-889 (2020)
- [16] C.L. Fewster, I. Jubb, I., and M.H. Røep. *Asymptotic Measurement Schemes for Every Observable of a Quantum Field Theory*. Ann. Henri Poincaré (2022)
- [17] F. Finster, C.F. Paganini, *Incompatibility of Frequency Splitting and Spatial Localization: A Quantitative Analysis of Hegerfeldt's Theorem*. Ann. Henri Poincaré 24, 413–467 (2023).
- [18] B. Gerlach, D. Gromes, J. Petzold, P. Rosenthal. *Über kausales Verhalten nicht-lokaler Größen und Teilchenstruktur in der Feldtheorie*. Z. Phys. 208, 381–389 (1968)
- [19] H. Halvorson and R. Clifton. *No place for particles in relativistic quantum theories?* in Ontological Aspects of Quantum Field Theory, Edited By: M. Kuhlmann, H. Lyre, and A. Wayne, World Scientific November 2002
- [20] J.J. Henning, W. Wolf. *Positive definite particle densities for the positive frequency solutions of the Klein Gordon equation with arbitrary mass*. Z. Phys. 242, 12–20 (1971)
- [21] G. C. Hegerfeldt. *Remark on causality and particle localization*, Phys. Rev. D 10, 3320 (1974)
- [22] G. C. Hegerfeldt. *Violation of Causality in Relativistic Quantum Theory?*, Phys. Rev. Lett. 54, 2395 (1985)
- [23] B. Jancewicz. *Operator density current and relativistic localization problem*, J. Math. Phys 18, 2487 (1977)

- [24] M. Lienert, R. Tumulka. *Born's rule for arbitrary Cauchy surfaces*. Lett Math Phys 110, 753–804 (2020).
- [25] S. Lill, R. Tumulka. *Another Proof of Born's Rule on Arbitrary Cauchy Surfaces*. Ann. Henri Poincaré 23, 1489–1524 (2022).
- [26] D. Malament. *In defense of dogma: Why there cannot be a relativistic quantum mechanics of (localizable) particles*. In R. Clifton (Ed.), Perspectives on Quantum Reality (pp. 1–11). Dordrecht: Kluwer. (1996)
- [27] T. Miller, M. Eckstein, P. Horodecki, R. Horodecki. *Generally covariant N-particle dynamics*. J. Geom. Phys. 160, 103990 (2021).
- [28] V. Moretti. *Spectral Theory and Quantum Mechanics* 2nd edition (2017) Springer
- [29] V. Moretti. *On the Relativistic Spatial Localization for massive real scalar Klein-Gordon quantum particles*. Lett.Math.Phys. 113, 66 (2023).
- [30] T.D. Newton, E.P. Wigner. *Localized States for Elementary Systems*, Rev. Mod. Phys. 21, 400-406 (1949)
- [31] B. O'Neill. *Semi Riemannian Geometry With Applications to Relativity*, Academic Press; 1st edition (July 12, 1983)
- [32] M.Reddiger, B.Poirier *Towards a Probabilistic Foundation of Relativistic Quantum Theory: The One-Body Born Rule in Curved Spacetime*. <https://arxiv.org/abs/2012.05212>
- [33] W.Rudin. *Real and Complex Analysis* 3d edition, McGraw-Hill, (1986)
- [34] D.R. Terno. *Localization of relativistic particles and uncertainty relations*. Phys. Rev. A 89, 042111 (2014)
- [35] D. N. Vollick. *Negative energy density states for the Dirac field in flat spacetime*. Phys. Rev. D 57, 3484 (1998)
- [36] A.S. Wightman. *On the Localizability of Quantum Mechanical Systems*, Rev. Mod. Phys. 34, 845-872 (1962)