

LOGARITHMIC CONCAVITY OF BIMATROIDS

FELIX RÖHRLE AND MARTIN ULIRSCH

ABSTRACT. A bimatroid is a matroid-like generalization of the collection of regular minors of a matrix. In this article, we use the theory of Lorentzian polynomials to study the logarithmic concavity of natural sequences associated to bimatroids. Bimatroids can be used to characterize morphisms of matroids and this observation (originally due to Kung) allows us to prove a weak version of logarithmic concavity of the number of bases of a morphism of matroids. This is weaker than the original result by Eur and Huh; it nevertheless provides us with a new perspective on Mason’s log-concavity conjecture for independent sets of matroids. We finally show that for realizable bimatroids, the regular minor polynomial is a volume polynomial. Applied to morphisms of matroids, this shows that the weak basis generating polynomial of a morphism is a volume polynomial; this confirms a conjecture of Eur–Huh for morphisms of nullity ≤ 1 and gives an algebro-geometric explanation for Mason’s log-concavity conjecture in the realizable case.

CONTENTS

Introduction	1
1. Summary of the main results	2
2. Bimatroids – the basic story	5
3. Products of bimatroids and morphisms of matroids	8
4. Lorentzian polynomials and logarithmic concavity	11
5. Volume polynomials	14
References	15

INTRODUCTION

Let a_1, \dots, a_s be a sequence of non-negative real numbers. We say (in decreasing generality) that the sequence is

- *unimodal*, if there is an index t such that $a_1 \leq \dots \leq a_t \geq \dots \geq a_s$;
- *log-concave*, if we have

$$a_k^2 \geq a_{k-1} \cdot a_{k+1}$$

for all $1 < k < s$; and

- *ultra log-concave*, if the sequence $\tilde{a}_k := a_k / \binom{s}{k}$ is log-concave.

(Ultra-)log-concave and unimodal sequences are objects of central interest throughout many fields of mathematics, since showing these properties typically requires the combination of insights coming from a priori quite different fields of mathematics. Recent examples are the proofs of log-concavity for the coefficients of the characteristic polynomial of a matroid in [AHK18] and for the number of independent sets of a matroid in [ALGV18, BH20] (also see [CP24] for further refinements), both of which require an intricate combination of methods from combinatorics, algebraic geometry and, in particular, from Hodge theory. We particularly highlight the recent introduction of the theory of Lorentzian polynomials in [BH20] (also see [ALGV18, ALGV19, AGV21] for an equivalent approach), which may be thought of as a generalization of both ultra log-concave sequences and the Hodge index theorem to the realm of homogeneous polynomials.

In this article we expand on the developments in [BH20] and apply these methods to *bimatroids* in the sense of [Kun78] (or, equivalently, to *linking systems* in the sense of [Sch79]). This will give us a new perspective on the log-concavity results for morphisms of matroids proved in [EH20]. In particular, we find a more direct proof of Mason’s log-concavity conjecture for independent sets of a matroid and prove [EH20, Conjecture 5.6] on volume polynomials for morphisms of nullity ≤ 1 .

At this point, we would like to refer the reader to the excellent survey articles [Kat16, AHK17, Bak18, Huh18a, Huh18b, Huh22, Eur24], which have significantly shaped our understanding of the subject.

Acknowledgements. The authors would like to thank Jeffrey Giansiracusa, Kevin Kühn, Arne Kuhrs, Felipe Rincón, Victoria Schleis, Pedro Souza, Hendrik Süß, and Igor Pak for helpful conversations and discussions en route to this article. Central ideas for this article were conceived during the 2023 Chow Lectures at the MPI in Leipzig, given by June Huh. We thank all speakers and organizers at this occasion.

Funding. This project has received funding from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) TRR 195 *Symbolic Tools in Mathematics and their Application*, from Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) and TRR 326 *Geometry and Arithmetic of Uniformized Structures*, project number 444845124, as well as from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Sachbeihilfe *From Riemann surfaces to tropical curves (and back again)*, project number 456557832.

Notation. For a set S we write 2^S for its power set, i.e. the set of all its subsets, and we write $\binom{S}{k}$ and $\binom{S}{\leq k}$ for the set of all subsets of size k and $\leq k$ respectively. Given two sets S and S' as well as $s \in S - S'$ and $s' \in S' - S$, we abbreviate $S - \{s\} \cup \{s'\}$ by $S_{s \leftrightarrow s'}$.

1. SUMMARY OF THE MAIN RESULTS

Let E and F be finite sets and let $A \in \mathbb{K}^{E \times F}$ be a matrix over a field \mathbb{K} indexed by $E \times F$. It is well-known that the collection of linearly independent subsets of the set of column vectors of A carries the structure of a *matroid* on the ground set F .

1.1. Regular minors. A finer invariant is the collection of *regular minors* of A , i.e. the collection of $(I, J) \in \binom{E}{k} \times \binom{F}{k}$ for $k = 0, \dots, \min\{|E|, |F|\}$, such that $\det[A]_{I, J} \neq 0$. The regular minors of A define a matroid-like structure that was introduced by Kung in [Kun78], a so-called *bimatroid* (see Section 2.1 below for a precise definition). Around the same time, and apparently unbeknownst to each other, Schrijver introduced the mathematically equivalent notion of a *linking system* in [Sch79]. In this article we follow the terminology introduced in [Kun78] and give references to the corresponding results in [Sch79], whenever appropriate.

In the realizable case we consider the extended matrix $\hat{A} = [I_E | A] \in \mathbb{K}^{E \times (E \sqcup F)}$, where $I_E \in \mathbb{K}^{E \times E}$ denotes the identity matrix of size $|E|$. The bimatroid of regular minors of the matrix A admits a cryptomorphic description as the matroid of columns of \hat{A} (see Proposition 2.2 below for the general, not necessarily realizable case).

An immediate application of Lorentzian polynomials to bimatroids implies our first result.

Theorem A. *Let E and F be finite sets and A a bimatroid on the ground set $E \times F$. Set $m = |E|$ as well as $n = |F|$ and write $m \wedge n = \min\{m, n\}$. The number $R_k(A)$ of regular $k \times k$ minors is an ultra log-concave sequence, i.e. we have*

$$\frac{R_k(A)^2}{\binom{m \wedge n}{k}^2} \geq \frac{R_{k+1}(A)}{\binom{m \wedge n}{k+1}} \cdot \frac{R_{k-1}(A)}{\binom{m \wedge n}{k-1}}$$

for all $k \geq 1$.

We point out that we already have $R_k(\mathbf{A}) = 0$, whenever k is bigger than the rank of \mathbf{A} . In the realizable case, Theorem A tells us that the number $R_k(\mathbf{A})$ of regular $k \times k$ minors of a matrix $A \in \mathbb{K}^{E \times F}$ is an ultra log-concave sequence.

1.2. Regular rectangles. In Section 2.3 below we introduce a new cryptomorphic description of a bimatroid \mathbf{A} in terms of (*horizontal or vertical*) *regular rectangles*. In the realizable case, the regular rectangles of $A \in \mathbb{K}^{E \times F}$ are precisely the (not necessarily square) submatrices $[A]_{S,T}$ for $S \subseteq E$ and $T \subseteq F$ that have maximal rank. We say that a regular rectangle is *horizontal*, if $|S| \leq |T|$ and *vertical*, if $|T| \leq |S|$.

As an application of the strongest form of the Lorentzian property of the homogeneous independent set generating polynomials used in the proof of Mason's conjecture [BH20], we obtain the following result.

Theorem B. *Let E and F be finite sets and \mathbf{A} a bimatroid on the ground set $E \times F$ and write $N = |E| + |F|$. Denote*

- by $RR_k^\uparrow(\mathbf{A})$ be the number of vertical regular rectangles in \mathbf{A} with k columns and
- by $RR_k^\leftrightarrow(\mathbf{A})$ the number of horizontal regular rectangles in \mathbf{A} with k rows.

Then both sequences $RR_k^\uparrow(\mathbf{A})$ and $RR_k^\leftrightarrow(\mathbf{A})$ are ultra log-concave. To be precise, we have

$$\frac{RR_k^\uparrow(\mathbf{A})^2}{\binom{N}{k}^2} \geq \frac{RR_{k-1}^\uparrow(\mathbf{A})}{\binom{N}{k-1}} \cdot \frac{RR_{k+1}^\uparrow(\mathbf{A})}{\binom{N}{k+1}}$$

as well as

$$\frac{RR_k^\leftrightarrow(\mathbf{A})^2}{\binom{N}{k}^2} \geq \frac{RR_{k-1}^\leftrightarrow(\mathbf{A})}{\binom{N}{k-1}} \cdot \frac{RR_{k+1}^\leftrightarrow(\mathbf{A})}{\binom{N}{k+1}}$$

for all $k \geq 1$.

1.3. Logarithmic concavity of morphisms. Let F and F' be finite sets and \mathbf{M} and \mathbf{M}' matroids on F and F' respectively. Recall that a map $\phi: F \rightarrow F'$ is said to be a *morphism of matroids* if the pullback $\phi^*\mathbf{M}'$ is a quotient of \mathbf{M} (see Section 3.2 below for a reminder on this and related notions). The *nullity* of ϕ is the difference $\text{nul}(\phi) = \text{rk}(\mathbf{M}) - \text{rk}(\phi^*\mathbf{M}')$.

In [EH20] the authors define a subset $T \subseteq F$ to be a *basis* of a morphism ϕ , if T is contained in a basis of \mathbf{M} and $\phi(T)$ contains a basis of \mathbf{M}' . Denote by $\mathcal{B}_k(\phi)$ the set of bases and by $B_k(\phi)$ the number of bases of ϕ of a fixed cardinality k . Motivated by the characterization of morphisms of matroids via bimatroids developed in [Kun78], we can prove the following result:

Theorem C. *Let $\phi: \mathbf{M} \rightarrow \mathbf{M}'$ be a morphism of matroids. Then the sequence $B_k(\phi)$ is log-concave, i.e. we have*

$$B_k(\phi)^2 \geq B_{k+1}(\phi) \cdot B_{k-1}(\phi)$$

for all $k \geq 1$.

We point out that in [EH20, Theorem 1.3] the authors prove that the sequence $B_k(\phi)$ itself is ultra log-concave, so our result is a weaker version of theirs. It has been noted in [EH20] that, when \mathbf{M}' is the uniform matroid $\mathbf{U}_{0,1}$, the set of bases of ϕ is equal to the set of independent subsets of \mathbf{M} ; hence the ultra log-concavity of $B_k(\phi)$ implies the strongest version Mason's conjecture on the ultra log-concavity of the number of independent sets of a matroid \mathbf{M} . In our case, Theorem C implies the following weaker version of Mason's conjecture.

Corollary D. *Let \mathbf{M} be a matroid and denote by $I_k(\mathbf{M})$ the number of independent sets in \mathbf{M} of cardinality k . Then the sequence $I_k(\mathbf{M})$ is log-concave.*

For proofs of the ultra log-concavity of the sequence, i.e. the strongest version of Mason's conjecture, we of course refer the reader once again to [ALGV18, BH20]. Further refinements of both Mason's

conjecture and [EH20, Theorem 1.3] can be found [CP24, Theorems 1.6 and 1.16] and the latter also contains criteria for when the inequalities are equalities.

1.4. Volume polynomials and the Eur–Huh conjecture. Let X be a normal irreducible projective variety of dimension d over an algebraically closed field \mathbb{K} and consider a collection H of nef \mathbb{Q} -divisors H_1, \dots, H_n on X . Then, by [BH20, Theorem 4.6] the *volume polynomial*

$$\text{vol}_H(w) := (w_1 H_1 + \dots + w_n H_n)^d = \sum_{|\alpha|=d} \frac{d!}{\alpha!} (H_1^{\alpha_1} \dots H_n^{\alpha_n}) \cdot w^\alpha$$

is Lorentzian. The converse does not hold; i.e. not every Lorentzian polynomial is a volume polynomial. In many situations (usually involving some form of realizability) it is therefore a natural question to ask whether a given Lorentzian polynomial is a volume polynomial. We refer the reader to [EL23] for a more elaborate theory of volume polynomials associated to discrete polymatroids.

Given a bimatroid $A \in \text{BMat}^{E \times F}$ of rank r , we may consider its *homogeneous regular minor polynomial*

$$p_{\mathcal{R}(A)}(w) = \sum_{(I,J) \in \mathcal{R}(A)} \prod_{e \in I^c} w_e \cdot \prod_{f \in J} w_f,$$

which is Lorentzian, since it agrees with the basis generating polynomial of the associated extended matroid \widehat{A} . Using the construction of matroid Schubert varieties, originally due to Ardila and Boocher [AB16], we find the following result.

Theorem E. *Let A be a bimatroid which is realizable over an algebraically closed field \mathbb{K} . Then the homogeneous regular minor polynomial is a volume polynomial.*

Deciding whether a given Lorentzian polynomial is a volume polynomial tends to be a difficult question, which requires significant geometric insight. Since our proof of Theorem C via bimatroids is more direct and does not involve a limit argument (unlike the one in [EH20]), Theorem E implies the following result in the case that both M and M' as well as the morphism ϕ are realizable.

Theorem F. *Let M and M' be matroids on finite sets F and F' and suppose that $\phi: F \rightarrow F'$ defines a morphism such that M , M' , and ϕ are realizable over an algebraically closed field \mathbb{K} . Denote by r the rank of M and by $\text{nul}(\phi)$ the nullity of ϕ . Given an integer $\alpha \geq \text{nul}(\phi)$ the α -weak homogeneous basis generating polynomial*

$$p_{\mathcal{B}(\phi)}^\alpha(w) := \sum_{k \geq 0} \sum_{T \in \mathcal{B}_k(\phi)} \binom{\alpha}{r-k} w_0^{r-k} \prod_{f \in T} w_f$$

is a volume polynomial.

If we choose $\alpha = r$, Theorem F (together with [BH20, Theorem 4.6]) provides us with an algebro-geometric explanation for the validity of Theorem C in the realizable case and, in particular, for Mason’s conjecture on the log-concavity of independent sets, as stated in Corollary D.

One may view Theorem F as a weak version of [EH20, Conjecture 5.6], which states that, in the realizable case, the *homogeneous basis generating polynomial*

$$p_{\mathcal{B}(\phi)}(w) := \sum_{k \geq 0} \sum_{T \in \mathcal{B}_k(\phi)} w_0^{|F|-k} \prod_{f \in T} w_f$$

of ϕ is a volume polynomial.

Suppose that the nullity of ϕ is ≤ 1 , which means that $\mathcal{B}_k(\phi) = \emptyset$ unless $k = \text{rk } M$ or $k = \text{rk } M - 1$. In this case we may choose $\alpha = \text{nul}(\phi)$ and we have $p_{\mathcal{B}(\phi)}(w) = w_0^{|F|-r} \cdot p_{\mathcal{B}(\phi)}^{\text{nul}(\phi)}(w)$. This product is again a volume polynomial by Proposition 5.1 (i) and (ii) below. Theorem F therefore confirms the Eur–Huh conjecture for morphisms of nullity ≤ 1 .

2. BIMATROIDS – THE BASIC STORY

In this section we recall the basic theory of bimatroids, as originally introduced in [Kun78] (also see [Sch79] for an alternative approach under the name *linking systems* and [Mur95] as well as the upcoming [GRSU24] for a valuated version of this story). This part is mostly expository and is meant to introduce the notation used in the remainder of the article. In Section 2.3, we finally introduce a new cryptomorphic characterization of bimatroids in terms of *regular rectangles*.

2.1. Regular minors and extended matroid. Bimatroids are a matroidal generalization of the combinatorial structure of the set of regular minors of a given matrix.

Definition 2.1. Let E and F be finite sets and write $\binom{E}{*} \times \binom{F}{*}$ for the union $\bigcup_{d \geq 0} \binom{E}{d} \times \binom{F}{d}$. A *bimatroid* A on the ground set $E \times F$ is given by a subset $\mathcal{R}(A)$ of $\binom{E}{*} \times \binom{F}{*}$, called the set of *regular minors* of A , which fulfils the following axioms:

- (1) For $d = 0$, the pair (\emptyset, \emptyset) is a regular minor.
- (2) Let (I, J) and (I', J') be regular minors of A .
 - (a) For every $i' \in I' - I$
 - there is $i \in I - I'$ such that $(I_{i \leftrightarrow i'}, J)$ is a regular minor or
 - there is $j' \in J' - J$ such that $(I \cup \{i'\}, J \cup \{j'\})$ is a regular minor.
 - (b) For every $j \in J - J'$
 - there is $j' \in J' - J$ such that $(I, J_{j \leftrightarrow j'})$ is a regular minor, or
 - there is $i \in I - I'$ such that $(I - \{i\}, J - \{j\})$ is a regular minor.

We may write $\mathcal{R}(A) = \bigcup_{d \geq 0} \mathcal{R}_d(A)$, where $\mathcal{R}_d(A)$ denotes the set of regular minors of size $d \times d$ for $0 \leq d \leq \min\{|E|, |F|\}$. The set of bimatroids on the ground set $E \times F$ is denoted by $\text{BMat}^{E \times F}$.

There is a natural bijection between $\binom{E}{*} \times \binom{F}{*}$ and $\binom{E \sqcup F}{|E|}$, given by the association

$$(1) \quad (I, J) \longmapsto I^c \sqcup J.$$

Via this bijection we may identify the set $\mathcal{R}(A)$ of *regular minors* of A on $E \times F$ with the set of bases of the *extended matroid* \hat{A} associated to A .

Proposition 2.2. *Let E and F be finite sets. A subset of $\binom{E}{*} \times \binom{F}{*}$ is the set of regular minors of a bimatroid A if and only if under bijection (1) it corresponds to the set of bases of a matroid \hat{A} on $E \sqcup F$ such that E is a basis.*

In order to prove Proposition 2.2 it is enough to observe that the basis exchange axioms for \hat{A} are precisely equivalent to the axioms defining a bimatroid.

Example 2.3 (Realizable bimatroids). Let E and F be finite sets, let \mathbb{K} be a field and $A \in \mathbb{K}^{E \times F}$ a matrix. Then set $\mathcal{R}(A)$ of those square submatrices of A , whose determinant does not vanish, form the regular minors of a bimatroid on the ground set $E \times F$.

One way to see this is to consider the extended matrix

$$\hat{A} = [I_E | A] \in \mathbb{K}^{E \times (E \sqcup F)}$$

and to note that the set of bases among the column vectors form the bases of a matroid on $E \sqcup F$. Given an element in $\binom{E \sqcup F}{|E|}$, written as $I^c \sqcup J$ for $(I, J) \in \binom{E}{*} \times \binom{F}{*}$, we find that

$$\det [\hat{A}]_{E, I^c \sqcup J} = \det [A]_{I, J}.$$

So these bases are in natural one-to-one correspondence with the regular minors of A .

Example 2.4 (Relations). Let E and F be finite sets and let R be a *relation* between E and F , i.e. a subset of $E \times F$. Given $(I, J) \in \binom{E}{*} \times \binom{F}{*}$, a *matching* between I and J is the graph of a bijection

between I and J . The set of $(I, J) \in \binom{E}{*} \times \binom{F}{*}$, for which R contains a matching between I and J defines a bimatroid $[R] \in \text{BMat}^{E \times F}$.

One way to show this is to note that a relation between E and F may be interpreted as a Boolean matrix $A[R] \in \mathbb{B}^{E \times F}$. Then the bimatroid $[R]$ may be described as the Stiefel matroid on $E \sqcup F$ associated to the extended Boolean matrix $\hat{A}[R] = [I_E | A[R]] \in \mathbb{B}^{E \times (E \sqcup F)}$, where I_E is the tropical identity matrix (see [FR15, Section 3.1] for details). In fact, we may choose a sufficiently generic lift of the Boolean matrix $\hat{A}[R] = [I_E | A[R]]$ (which is always possible over an infinite field) and apply Example 2.3.

Note that the graph of every map $\phi: F \rightarrow E$ is a relation between E and F . So, in particular, we have an induced bimatroid $[\phi] \in \text{BMat}^{E \times F}$.

Example 2.5 (Bond bimatroids). Let F be a finite set and M a matroid on F on rank r . Choose a basis B of M . The *bond bimatroid* $\text{Bond}_B(M) \in \text{BMat}^{B \times F}$ of M with respect to the basis B is defined as follows: A pair $(I, J) \in \binom{B}{*} \times \binom{F}{*}$ is a regular minor of $\text{Bond}_B(M)$ if and only if the (not necessarily disjoint) union $I^c \cup J$ is a basis of M .

A quick way to prove this is to observe that the regular minors of $\text{Bond}_B(M)$ naturally correspond to the set of bases of the matroid on the disjoint union $B \sqcup F$, where a subset of size r is a basis if and only if its image in $B \cup F$ is a basis of M .

2.2. Relative rank. Let E and F be finite sets. A bimatroid A on $E \times F$ admits another cryptomorphic description in terms of the *relative rank function* $r_A: 2^E \times 2^F \rightarrow \mathbb{Z}_{\geq 0}$, which associates to $(S, T) \in 2^E \times 2^F$ the maximal number $d \geq 0$, for which there is a regular $d \times d$ -minor $(I, J) \in \mathcal{R}_d(A)$ with $I \subseteq S$ and $J \subseteq T$. So we always have $0 \leq r(S, T) \leq \min\{|S|, |T|\}$ for all $(S, T) \in 2^E \times 2^F$. The number $r_A(E, F)$ is called the *rank* of the bimatroid A .

The relative rank provides us with a cryptomorphic description of bimatroids (see [Kun78, Section 5] as well as [Sch79, (alternative) Definition 2.2]).

Proposition 2.6. *Let E and F be two finite sets. A function $r: 2^E \times 2^F \rightarrow \mathbb{Z}_{\geq 0}$ is the relative rank function of a bimatroid if and only if it fulfils the following properties:*

- (1) For every $(S, T) \in 2^E \times 2^F$ we have $r(S, T) \leq \min\{|S|, |T|\}$.
- (2) For every $(S, T) \in 2^E \times 2^F$ and $e \in E$ and $f \in F$, we have

$$r(S, T) \leq r(S \cup \{e\}, T) \leq r(S, T) + 1$$

as well as

$$r(S, T) \leq r(S, T \cup \{f\}) \leq r(S, T) + 1 .$$

- (3) For two pairs $(S, T), (S', T') \in 2^E \times 2^F$, we have

$$r(S, T) + r(S', T') \geq r(S \cup S', T \cap T') + r(S \cap S', T \cup T') .$$

Proof. Denote by \hat{r} the rank function of the extended matroid \hat{A} on $E \sqcup F$. Then we have

$$r(S, T) = \hat{r}(S^c \sqcup T) - |S^c|$$

for all $(S, T) \in 2^E \times 2^F$. The cryptomorphism here now follows from the cryptomorphic definition of matroids by their rank function. \square

Remark 2.7. In [Kun78, Section 5] Kung uses the normalization $r(\emptyset, \emptyset) = 0$ instead of the axiom $r(S, T) \leq \min\{|S|, |T|\}$. We believe that this is not correct, since otherwise the function $r(S, T) = |S| + |T|$ for $(S, T) \in 2^E \times 2^F$ would be the relative rank function of a bimatroid which does not match with the idea that the relative rank should be the size of the largest regular minor in (S, T) . To avoid this issue, we follow [Sch79, (alternative) Definition 2.2].

Example 2.8 (Transpose of a bimatroid). Let E and F be finite sets and A a bimatroid on $E \times F$. The transpose A^T of A is a bimatroid on the ground set $F \times E$; a pair $(J, I) \in \binom{F}{*} \times \binom{E}{*}$ is a regular minor of A^T if and only if (I, J) is a regular minor of A . The relative rank function of A^T is given by

$$r_{A^T}(T, S) = r_A(S, T)$$

for $S \times T \subseteq E \times F$. One way to see that A^T is indeed a bimatroid is to observe that the associated matroid is the dual matroid to \hat{A} . In terms of rank functions, this can be deduced from

$$\begin{aligned} \hat{r}_{A^T}(T^c \sqcup S) &= r_{A^T}(T, S) + |T^c| \\ &= r_A(S, T) + |S^c| + |S| - |E| + |T^c| \\ &= \hat{r}_A(S^c \sqcup T) + (|S| + |T^c|) - |E| \end{aligned}$$

for every $(S, T) \in 2^E \times 2^F$ and the observation that the right hand side is the rank function of the matroid dual to \hat{A} .

2.3. Regular rectangles. Let E and F be finite sets and A a bimatroid on $E \times F$. Given two subsets $S \subseteq E$ and $T \subseteq F$, we say that the pair (S, T) is a *regular rectangular minor* (for short a *regular rectangle*), if $r_A(S, T) = \min\{|S|, |T|\}$ or, equivalently, if there are $I \subseteq S$ and $J \subseteq T$ such that $|I| = |J| = \min\{|S|, |T|\}$ and $(I, J) \in \mathcal{R}(A)$. We call a pair $(S, T) \in 2^E \times 2^F$ a *vertical regular rectangle*, if $r_A(S, T) = |T| \leq |S|$, and a *horizontal regular rectangle*, if $r_A(S, T) = |S| \leq |T|$. We write $\mathcal{RR}(A)$ for the set of regular rectangles and denote the subsets of vertical and horizontal rectangles by $\mathcal{RR}^\dagger(A)$ and $\mathcal{RR}^\leftrightarrow(A)$ respectively.

Regular rectangles can be used to give another cryptomorphic description of a bimatroid A .

Proposition 2.9. *Let E and F be finite sets and A a bimatroid on the ground set $E \times F$. The association $(S, T) \mapsto S^c \sqcup T$ induces a bijection between $\mathcal{RR}^\dagger(A)$ and the independent sets of \hat{A} .*

Proposition 2.9 immediately implies the following cryptomorphic characterization of bimatroids.

Corollary 2.10. *Let E and F be finite sets. A subset \mathcal{RR}^\dagger of $2^E \times 2^F$ is the set of vertical regular rectangles of a bimatroid A on $E \times F$ if and only if the following axioms hold:*

- (1) $(\emptyset, \emptyset) \in \mathcal{RR}^\dagger$
- (2) For all $(S, T) \in \mathcal{RR}^\dagger$ we have $|T| \leq |S|$.
- (3) Given $(S, T) \in \mathcal{RR}^\dagger$ as well as $S \subseteq S' \subseteq E$ and $T' \subseteq T$, the pair (S', T') is also in \mathcal{RR}^\dagger .
- (4) Suppose we are given $(S, T), (S', T') \in \mathcal{RR}^\dagger$ with $|T| - |S| > |T'| - |S'|$ there is
 - $s' \in S' - S$ such that $(S' - \{s'\}) \times T' \in \mathcal{RR}^\dagger$ or
 - $t \in T - T'$ such that $S' \times (T' \cup \{t\}) \in \mathcal{RR}^\dagger$.

Proof. It is enough to note that all subsets of E are always independent in \hat{A} . The axioms are mere translations of the independent set axioms of \hat{A} using the bijection in Proposition 2.9. \square

In the proof of Proposition 2.9 we will make use of the following bimatroidal generalization of the Laplace expansion for matrices.

Lemma 2.11 (Laplace expansion). *Let E and F be finite sets and A a bimatroid on $E \times F$. Consider a regular minor $(I, J) \in \mathcal{R}(A)$.*

- (1) For every $j \in J$ there is $i \in I$ such that $(I - \{i\}, J - \{j\})$ is a regular minor of A .
- (2) For every $i \in I$ there is $j \in J$ such that $(I - \{i\}, J - \{j\})$ is a regular minor of A .

Proof. Part (1) immediately follows from Definition 2.1 Axiom (2) (b) applied with $(I', J') = (\emptyset, \emptyset)$. Part (2) follows with the same argument applied to A^T . \square

Proof of Proposition 2.9. Note, that the association $(S, T) \mapsto S^c \sqcup T$ defines a natural bijection

$$\{(S, T) \in 2^E \times 2^F \mid |S| \geq |T|\} \longrightarrow \binom{E \sqcup F}{\leq |E|}.$$

We now observe that the image of this map restricted to \mathcal{RR}^\dagger is precisely the set of independent subsets of \hat{A} . An independent set of \hat{A} corresponds to a pair $(S, T) \in 2^E \times 2^F$ with $|S| \geq |T|$ such that there are $I \subseteq S$ and $T \subseteq J \subseteq F$ with $(I, J) \in \mathcal{R}(A)$. If $I = S$ or $J = T$, this is already a regular rectangle. If not, we may apply the Laplacian expansion from Lemma 2.11 above (possibly several times) and find $I' \subseteq I$ such that $(I', T) \in \mathcal{R}(A)$. \square

Working with A^T instead of A we get a similar axiomatic characterization of the horizontal regular rectangles of a bimatroid. We leave the details of this reformulation to the avid reader.

3. PRODUCTS OF BIMATROIDS AND MORPHISMS OF MATROIDS

In this section we recall the construction of products of bimatroids from [Kun78, Sch79] as well as the basic terminology of morphisms of matroids (following [EH20]). We refer the reader to [Wel76, Chapter 17], to [Kun86], and the recent categorical exploration in [HP18] for more background on this notion. Section 3.3 contains the central construction for the proof of Theorem C in the next section.

3.1. Products of bimatroids. The central new feature of bimatroids is that, just like for matrices, but unlike for matroids, one can form products. In order to motivate this construction we recall the generalized *Cauchy–Binet formula*. Let E, F , and G be finite sets and \mathbb{K} be a field. Given two matrices $A \in \mathbb{K}^{E \times F}$ and $B \in \mathbb{K}^{F \times G}$ and a pair $(I, K) \in \binom{E}{*} \times \binom{G}{*}$, we have

$$\det[A \cdot B]_{I, K} = \sum_{J \in \binom{F}{|I|}} \pm \det[A]_{I, J} \cdot \det[B]_{J, K}.$$

We point out that this formula is an immediate consequence of the fact that for two linear maps $f: V' \rightarrow V''$ and $g: V \rightarrow V'$ between finite-dimensional vector spaces the induced maps on exterior products fulfil $\wedge(f \circ g) = \wedge(f) \circ \wedge(g)$.

The Cauchy–Binet formula tells us that a minor $[A \cdot B]_{I, K}$ can only be regular if there is $J \in \binom{F}{|I|}$ such that both $[A]_{I, J}$ and $[B]_{J, K}$ are regular as well.

Proposition 3.1. *Let E, F , and G be finite sets and let A and B be bimatroids on the ground sets $E \times F$ and $F \times G$ respectively. Then the set $\mathcal{R}(A \cdot B)$ of those $(I, K) \in \binom{E}{*} \times \binom{G}{*}$, for which there is $J \in \binom{F}{*}$ such that both $(I, J) \in \mathcal{R}(A)$ and $(J, K) \in \mathcal{R}(B)$, defines the set of regular minors of a bimatroid $A \cdot B$ on the ground set $E \times G$.*

The bimatroid $A \cdot B$ is called the *product* of A and B . A proof of Proposition 3.1 via rank functions, which uses the matroid intersection theorem, can be found in [Kun78, Section 6] and [Sch79, Theorem 3.5]. We complement this with an elementary verification of the regular minor axioms from Definition 2.1.

Proof of Proposition 3.1. Axiom (1) automatically holds, since $(\emptyset, \emptyset) \in \mathcal{R}(A)$ and $(\emptyset, \emptyset) \in \mathcal{R}(B)$. To verify Axiom (2) we consider $(I, K), (I', K') \in \mathcal{R}(A \cdot B)$. Note that there are $J \in \binom{F}{|I|}$ and respectively $J' \in \binom{F}{|I'|}$ such that both $(I, J), (I', J') \in \mathcal{R}(A)$ and $(J, K), (J', K') \in \mathcal{R}(B)$.

For **Part (a)** we let $i' \in I' - I$. We apply Axiom (2) Part (a) to the bimatroid A and see that we are in one of the following two cases: In the **first case** there is $i \in I - I'$ such that $(I_{i \leftrightarrow i'}, J) \in \mathcal{R}(A)$. In this case, we also have $(I_{i \leftrightarrow i'}, K) \in \mathcal{R}(A \cdot B)$ and we are done. In the **second case** there is $j' \in J' - J$ such that $(I \cup \{i'\}, J \cup \{j'\}) \in \mathcal{R}(A)$. We may now apply Axiom (2) Part (a) for the bimatroid B . If there is $k' \in K' - K$ such that $(J \cup \{j'\}, K \cup \{k'\}) \in \mathcal{R}(B)$, we immediately deduce $(I \cup \{i'\}, K \cup \{k'\}) \in \mathcal{R}(A \cdot B)$. Therefore, we now suppose that there is $j \in J - J'$ such that $(J_{j \leftrightarrow j'}, K) \in \mathcal{R}(B)$. We now apply Axiom (2) Part (b) to the bimatroid A with the two regular minors (I, J) and $(I \cup \{i'\}, J \cup \{j'\})$ and find that there

has to be $i \in I - I'$ such that $(I_{i \leftrightarrow i'}, J_{j \leftrightarrow j'}) \in \mathcal{R}(A)$. In this case it follows that $(I_{i \leftrightarrow i'}, K) \in \mathcal{R}(A \cdot B)$ and we are done.

In order to verify **Part (b)** of Axiom (2) for $A \cdot B$ we consider $k \in K - K'$. If we apply Axiom (2) Part (b) to B we find ourselves in one of the following two cases: In the **first case** there is $k' \in K' - K$ such that $(J, K_{k \leftrightarrow k'}) \in \mathcal{R}(B)$ and we immediately deduce $(I, K_{k \leftrightarrow k'}) \in \mathcal{R}(A \cdot B)$. In the **second case** there is $j \in J - J'$ such that $(J - \{j\}, K - \{k\}) \in \mathcal{R}(B)$. We may now apply Axiom (2) Part (b) for the bimatroid A . If there is $i \in I - I'$ such that $(I - \{i\}, J - \{j\}) \in \mathcal{R}(A)$ we immediately deduce $(I - \{i\}, K - \{k\}) \in \mathcal{R}(A \cdot B)$ and we are done. So let us now assume that instead there is a $j' \in J' - J$ such that $(I, J_{j \leftrightarrow j'}) \in \mathcal{R}(A)$. In this situation we apply Axiom (2) Part (a) to B with the regular minors $(J - \{j\}, K - \{k\})$ and (J, K) and find $k' \in K' - K$ such that $(J_{j \leftrightarrow j'}, K_{k \leftrightarrow k'}) \in \mathcal{R}(B)$. This implies $(I, K_{k \leftrightarrow k'}) \in \mathcal{R}(A \cdot B)$. \square

Example 3.2. Let E, F , and G be finite sets and let R be a relation between E and F and S a relation between F and G . Recall (e.g. from [HV19, Definition 0.5]) that the *product* $R \cdot S$ is defined by

$$R \cdot S = \{(e, g) \in E \times G \mid \text{there is } f \in F \text{ such that } (e, f) \in R \text{ and } (f, g) \in S\} .$$

This product is compatible with the product of bimatroids, i.e. we have

$$[R \cdot S] = [R] \cdot [S] .$$

In particular, given two maps $\phi: F \rightarrow E$ and $\psi: G \rightarrow F$, we have

$$[\phi \circ \psi] = [\phi] \cdot [\psi] .$$

Example 3.3. Let E and F be finite sets and A a bimatroid on the ground set $E \times F$. Given subsets $E' \subseteq E$ and $F' \subseteq F$, we write $i: E' \hookrightarrow E$ and $j: F' \hookrightarrow F$ for the inclusion maps. Then the *restriction* $A|_{E' \times F'}$ of A to $E' \times F'$ is defined by

$$A|_{E' \times F'} = [i]^T \cdot A \cdot [j] \in \mathbf{BMat}^{E' \times F'} .$$

According to this definition, an element $(I, J) \in \binom{E'}{*} \times \binom{F'}{*}$ is a regular minor of $A|_{E' \times F'}$ if and only if it is a regular minor of A .

The avid reader may now immediately verify that bimatroids naturally form a category \mathbf{BMat} whose objects are finite sets and in which morphisms $F \rightarrow E$ (for finite sets E and F) are bimatroids on the ground set $E \times F$ (with products as composition). Explicitly this means the following:

- The product is associative: For all bimatroids $A \in \mathbf{BMat}^{E \times F}$, $B \in \mathbf{BMat}^{F \times G}$, and $C \in \mathbf{BMat}^{G \times H}$ and finite sets E, F, G , and H we have

$$(A \cdot B) \cdot C = A \cdot (B \cdot C) .$$

- The bimatroids $I_E = [\text{id}_E]$ (for a finite set E) serve as identity morphisms: For all bimatroids $A \in \mathbf{BMat}^{E \times F}$ we have

$$I_E \cdot A = A = A \cdot I_F .$$

Remark 3.4. Taking transpose naturally endows the category \mathbf{BMat} with a dagger structure (see [HV19, Section 2.3] for details on this notion). In our situation this means the following: Given finite sets E, F , and G , we have $I_E^T = I_E$ as well as $(A^T)^T = A$ and $(A \cdot B)^T = B^T \cdot A^T$ for all bimatroids $A \in \mathbf{BMat}^{E \times F}$ and $B \in \mathbf{BMat}^{F \times G}$.

3.2. Morphisms of matroids. Let F and F' be finite sets. Recall from [EH20] that, given two matroids M and M' on F and F' respectively, a map $\phi: F \rightarrow F'$ is called a *morphism* from M to M' if one (and therefore all) of the following three equivalent conditions hold:

- For all $T_1 \subseteq T_2 \subseteq F$, we have

$$r_{M'}(\phi(T_2)) - r_{M'}(\phi(T_1)) \leq r_M(T_2) - r_M(T_1).$$

- If $T' \subseteq F'$ is a cocircuit of M' , then $\phi^{-1}(T')$ is a union of cocircuits of M .
- If $T' \subseteq F'$ is a flat of M' , then $\phi^{-1}(T')$ is flat of M .

Example 3.5 (Realizable morphisms). Let \mathbb{K} be a field. Let M and M' be matroids on finite ground sets F and F' , realized by vectors $(v_s)_{s \in F}$ and $(v'_{s'})_{s' \in F'}$ spanning vector spaces V and V' . Suppose further that we have a map $\phi: F \rightarrow F'$ and a \mathbb{K} -linear map $\Phi: V \rightarrow V'$ such that $\Phi(v_s) = v'_{\phi(s)}$ for all $s \in F$. Then ϕ defines a morphism of matroids. Morphisms of this type are called *realizable* over \mathbb{K} .

Given two matroids M and N on the same ground set F , we say that N is a *quotient* of M , if the identity map id_F is a morphism of matroids from M to N . Let $\phi: F \rightarrow F'$ be a map of finite sets. For a matroid M' on F' , the *pullback matroid* ϕ^*M' is a matroid on F , whose rank function $r: 2^F \rightarrow \mathbb{Z}_{\geq 0}$ given by

$$r_{\phi^*M'}(T) = r_{M'}(\phi(T))$$

for $T \subseteq F$. Recall that by [EH20, Lemma 2.4] a map $\phi: F \rightarrow F'$ is a morphism between matroids M and M' on F and F' respectively if and only if ϕ^*M' is a quotient of M . We call the difference of ranks

$$\text{nul}(\phi) := \text{rk}(M) - \text{rk}(\phi^*M')$$

the *nullity* of ϕ .

Following [EH20, Definition 1.1], we say that a subset $T \subseteq F$ is a *basis* of the morphism ϕ if it is contained in a basis of M and $\phi(T)$ contains a basis of M' . By [EH20, Lemma 2.4] the set $\mathcal{B}(\phi)$ of bases of ϕ is either empty, when $\phi(E)$ does not span M' , or, otherwise, equal to the set of bases of the quotient $M \rightarrow \phi^*M'$. There are two extreme cases: When ϕ is the identity morphism, then $\mathcal{B}(\phi)$ is the set of bases of M and, when M' is the uniform matroid $M' = U_{0,1}$, then $\mathcal{B}(\phi)$ is the set $\mathcal{I}(M)$ of independent sets of the matroid M . The last observation explains why Theorem C implies Corollary D.

3.3. Classifying bases of morphisms. In [Kun78, Theorem 4] Kung characterizes morphisms of matroids in terms of bimatroid multiplication. This motivates the following construction.

Consider two finite sets F and F' , two matroids M on F and M' on F' , as well as a map $\phi: F \rightarrow F'$ that is a morphism of matroids. Write N for the pullback ϕ^*M' , so that N is a quotient of M on F . By the Higgs factorization theorem (see e.g. [Whi86, Theorem 8.2.8] or [Oxl11, Section 7.3]), there is a finite set \tilde{F} containing F as well as a matroid \tilde{M} on \tilde{F} such that

$$\tilde{M}|_F = M \quad \text{and} \quad \tilde{M}/Q = N$$

where we denote the complement $\tilde{F} - F$ by Q . Here, as detailed in [Oxl11, Lemma 7.3.3], we may always choose \tilde{M} (and \tilde{F}) in such a way that Q is an independent subset of \tilde{M} and $\text{rk}(\tilde{M}) = \text{rk}(M)$. Note that in this case $|Q| = \text{nul}(\phi)$.

Now choose a finite set E of size $r = \text{rk}(M)$ that is disjoint from F and contains Q . Consider the *free extension* $\tilde{M} + (E - Q)$, a matroid on $E \sqcup F$ whose bases are all subsets $S \subseteq E \sqcup F$ of size $\text{rk} \tilde{M}$ such that $S \cap \tilde{F}$ is independent in \tilde{M} (see [Whi86, Proposition 7.3.3]). We think of $\tilde{M} + (E - Q)$ as the extended matroid of a bimatroid $\tilde{A}[\phi]$ on $E \times F$. Denote by $R_{E \leftrightarrow E}$ the relation $R_{E \leftrightarrow E} := E \times E$ and define the bimatroid

$$A[\phi] := [R_{E \leftrightarrow E}] \cdot \tilde{A}[\phi]$$

on the ground set $E \times F$.

Proposition 3.6. *Let M and M' be matroids on finite sets F and F' respectively and let $\phi: F \rightarrow F'$ be a morphism from M to M' whose image spans M' . Then the map $(I, J) \mapsto J$ defines a surjection from the regular minors of $A[\phi]$ onto the set of bases of ϕ . Given a basis T of ϕ , it has $\binom{r}{r-k} = \binom{r}{k}$ many preimages, where $r = \text{rk}(M)$ and $k = |T|$.*

Proof. Since the image of ϕ spans M' , we only need to consider the quotient $N = \phi^*M'$ of M .

Consider $(I, J) \in \mathcal{R}(A[\phi])$. By the definition of $R_{E \leftrightarrow E}$ there is $I' \subseteq E$ such that $(I', J) \in \mathcal{R}(\tilde{A}[\phi])$. Writing $Q' := (E - I') \cap Q$, this means that $Q' \sqcup J$ is a basis of \tilde{M} . This immediately implies that J is independent in \tilde{M} and contained in F , and thus it is also independent in M . Since Q is independent in \tilde{M} we may form an independent set $Q \sqcup J' \subseteq \tilde{F}$ of \tilde{M} with $J' \subseteq J$ by adding elements in J to Q as long as $|Q \sqcup J'| < |Q \sqcup J|$. This process ends, as soon as $Q \sqcup J'$ is a basis of \tilde{M} . But then J' is a basis of N by [Oxl11, Corollary 3.1.8] and this shows that J contains the basis J' of N . Hence J is a basis of the quotient $M \rightarrow N$.

Conversely, suppose we are given a basis $J \subseteq F$ of the quotient $M \rightarrow N$. Then J is independent in M and contains a basis J' of N . By [Oxl11, Corollary 3.1.8] this means that $Q \sqcup J'$ is a basis of \tilde{M} . We now form a new basis $Q' \sqcup J$ of \tilde{M} by adding elements in Q to J as long as $|Q' \sqcup J| < |Q \sqcup J'|$. This process ends, once $Q' \sqcup J$ is a basis of \tilde{M} . Now set $I' := E - Q'$ and note that $(I', J) \in \mathcal{R}(\tilde{A}[\phi])$. By the definition of $R_{E \leftrightarrow E}$ this implies that for every $I \subseteq \tilde{E}$ with $|I| = |I'|$ we have $(I, J) \in \mathcal{R}(A[\phi])$. Noting that $|E| = r$, we find that the number of choices for such an I is given by $\binom{|E|}{|E|-|J|} = \binom{r}{r-k} = \binom{r}{k}$. \square

4. LORENTZIAN POLYNOMIALS AND LOGARITHMIC CONCAVITY

We begin by recalling the definition and basic properties of *Lorentzian polynomials* from [BH20]. Let n and d be non-negative integers and write $\Delta_n^d = \{\alpha \in \mathbb{Z}_{\geq 0}^n \mid |\alpha| := \alpha_1 + \dots + \alpha_n = d\}$. Denote by $H_n^d \subseteq \mathbb{R}[w_1, \dots, w_n]$ the set of homogeneous polynomials with real coefficients of degree d in n variables w_1, \dots, w_n . Whenever convenient, we write a polynomial $p(w) \in H_n^d$ in the variables $w = (w_1, \dots, w_n)$ as

$$p(w) = \sum_{\alpha \in \Delta_n^d} a_\alpha w^\alpha$$

using multi-index notation $w^\alpha = w_1^{\alpha_1} \dots w_n^{\alpha_n}$ for $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$. Following this notational logic, we also write $\partial^\alpha p$ for

$$\partial^\alpha p = \left(\frac{\partial}{\partial w_1} \right)^{\alpha_1} \dots \left(\frac{\partial}{\partial w_n} \right)^{\alpha_n} p$$

as well as $\alpha! = \alpha_1! \dots \alpha_n!$.

Denote by P_n^d the open subset of polynomials in H_n^d , for which all coefficients a_α are positive.

Definition 4.1 ([BH20, Definition 2.1]). Set $\mathring{L}_n^0 = P_n^0$, $\mathring{L}_n^1 = P_n^1$, as well as

$$\mathring{L}_n^2 = \{p \in P_n^2 \mid \text{Hess}(p) \text{ has the Lorentzian signature } (+, -, \dots, -)\}$$

and

$$\mathring{L}_n^d = \{p \in P_n^d \mid \partial^\alpha p \in \mathring{L}_n^2 \text{ for all } \alpha \in \Delta_n^{d-2}\}.$$

Polynomials in \mathring{L}_n^d are called *strictly Lorentzian* and we define the space L_n^d of *Lorentzian polynomials* as the closure of \mathring{L}_n^d in H_n^d .

In the following, we write e_1, \dots, e_n for the standard basis vectors of \mathbb{Z}^n . Recall now, e.g. from [Mur03], that a subset $S \subseteq \mathbb{Z}_{\geq 0}^n$ is said to be *M-convex*, if the following *exchange property* holds: For any $\alpha, \beta \in S$ and any index i satisfying $\alpha_i > \beta_i$, there is an index j such that $\alpha_j < \beta_j$ and $\alpha - e_i + e_j \in S$. Observe that every *M-convex* set S is automatically contained in Δ_n^d and that, if S is also contained in $\{0, 1\}^n$, the elements in S are the indicator vectors of bases of a matroid M ; here the exchange property of S is the same as the symmetric basis exchange property of M .

Lorentzian polynomials enjoy the following useful properties:

- A homogeneous polynomial $p(x, y) = \sum_{k=0}^d a_k x^k y^{d-k}$ in two variables with non-negative coefficients is Lorentzian if and only if the sequence a_k is ultra log-concave and has no internal zeros, i.e. if $a_k a_{k'} > 0$ then $a_{k''} > 0$ for all $k < k'' < k'$ (see [BH20, Example 2.26]).
- For every $p = \sum_{\alpha \in \Delta_n^d} a_\alpha w^\alpha \in L_n^d$ the support

$$\text{supp}(p) = \{\alpha \in \Delta_n^d \mid a_\alpha \neq 0\}$$

is an M -convex set (see [BH20, Theorem 2.25]). Vice versa, given an M -convex set $S \subseteq \Delta_n^d$, the indicator polynomial

$$p_S(w) = \sum_{\alpha \in S} \frac{w^\alpha}{\alpha!}$$

is Lorentzian (see [BH20, Theorem 3.10]).

- Linear coordinate changes: Given a Lorentzian polynomial $p(w) \in L_m^d$ and a matrix $A \in \mathbb{R}_{\geq 0}^{m \times n}$, the linear coordinate change $p(Az)$ is also Lorentzian (see [BH20, Theorem 2.10]).
- Given two Lorentzian polynomials $p \in L_n^d$ and $q \in L_{n'}^{d'}$, their product $p \cdot q \in L_{n+n'}^{d+d'}$ is again a Lorentzian polynomial (see [BH20, Corollary 2.32]).
- Given a Lorentzian polynomial $p(w) = \sum_{\alpha \in \Delta_n^d} a_\alpha w^\alpha$ as well as $\kappa \in \mathbb{Z}_{\geq 0}^n$, the κ -truncations

$$p_{\leq \kappa}(w) = \sum_{\alpha \leq \kappa} a_\alpha w^\alpha \quad \text{and} \quad p_{\geq \kappa}(w) = \sum_{\alpha \geq \kappa} a_\alpha w^\alpha$$

are also Lorentzian (see [RSW23, Proposition 3.3]). Here we write $\alpha \leq \beta$ for $\alpha, \beta \in \mathbb{Z}_{\geq 0}^n$ if and only $\alpha_i \leq \beta_i$ for all $1 \leq i \leq n$.

We illustrate the interactions of these properties by giving a proof of Newton's inequalities using Lorentzian polynomials (in the spirit of [Sta89, Theorem 2]).

Example 4.2 (Newton's inequalities). Let $a = (a_1, \dots, a_n)$ be a sequence of non-negative numbers. Then the homogeneous polynomials $x + a_i y$ are of degree one and thus Lorentzian for all $1 \leq i \leq n$. Therefore also the product

$$\prod_{i=1}^n (x + a_i y) = x^n + e_1(a) x^{n-1} y + \dots + e_{n-1}(a) x y^{n-1} + e_n(a) y^n$$

is Lorentzian, where e_k denotes the k -th elementary symmetric polynomial of degree n . But this means that the sequence $e_0(a), \dots, e_n(a)$ is ultra log-concave. In other words, we have just given another proof of Newton's inequalities

$$\left(\frac{e_k(a)}{\binom{n}{k}} \right)^2 \geq \left(\frac{e_{k+1}(a)}{\binom{n}{k+1}} \right) \cdot \left(\frac{e_{k-1}(a)}{\binom{n}{k-1}} \right)$$

for $1 \leq k \leq n-1$.

We are now ready to prove Theorems A, B, and C.

Proof of Theorem A. Let A be a bimatroid on the ground set $E \times F$ and set $m = |E|$ as well as $n = |F|$. Recall that the basis generating polynomial $p_{\mathcal{B}(\hat{A})}$ of the associated matroid \hat{A} is given by

$$p_{\mathcal{B}(\hat{A})}(\underline{x}, \underline{y}) = \sum_{B \in \mathcal{B}(\hat{A})} \prod_{e \in B \cap E} x_e \prod_{f \in B \cap F} y_f.$$

By [BH20, Theorem 3.10] we have that $p_{\mathcal{B}(\hat{A})}$ is Lorentzian (it is the generating function of the M -convex set of bases of a matroid). Note that by Proposition 2.2 this polynomial is equal to the regular minor polynomial $p_{\mathcal{R}(A)}$ of A .

Now apply the coordinate transformation $x_e = x$ for all $e \in E$ and $y_f = y$ for all $f \in F$ and use [BH20, Theorem 2.10] to conclude that

$$q(x, y) = \sum_{k=0}^m R_k(A) x^{m-k} y^k$$

is Lorentzian as well. Lorentzian bivariate polynomials were characterized in [BH20, Example 2.26] and this implies that the sequence $R_k(A)$ fulfils

$$\frac{R_k(\mathbf{A})^2}{\binom{m}{k}^2} \geq \frac{R_{k+1}(\mathbf{A})}{\binom{m}{k+1}} \cdot \frac{R_{k-1}(\mathbf{A})}{\binom{m}{k-1}}$$

for all $k \geq 1$. The same argument, applied to \mathbf{A}^T shows

$$\frac{R_k(\mathbf{A})^2}{\binom{n}{k}^2} \geq \frac{R_{k+1}(\mathbf{A})}{\binom{n}{k+1}} \cdot \frac{R_{k-1}(\mathbf{A})}{\binom{n}{k-1}}$$

for all $k \geq 1$. We now combine these two inequalities and find

$$\frac{R_k(\mathbf{A})^2}{\binom{m \wedge n}{k}^2} \geq \frac{R_{k+1}(\mathbf{A})}{\binom{m \wedge n}{k+1}} \cdot \frac{R_{k-1}(\mathbf{A})}{\binom{m \wedge n}{k-1}}$$

for all $k \geq 1$. □

Proof of Theorem B. Consider the homogenized independent set generating polynomial $p_{\mathcal{I}(\hat{A})}$ of the extended matroid \hat{A} , which is Lorentzian by [BH20, Theorem 4.10] (see the explanation right after [BH20, Theorem 4.14]). Set $N = |E| + |F|$. In our situation it may be written as

$$p_{\mathcal{I}(\hat{A})}(\underline{x}, \underline{y}, z) = \sum_{S \in \mathcal{I}(\hat{A})} z^{N-|S|} \prod_{e \in S \cap E} x_e \prod_{f \in S \cap F} y_f .$$

By Proposition 2.9 above, this can be rewritten as

$$p_{\mathcal{I}(\hat{A})}(\underline{x}, \underline{y}, z) = \sum_{(I, J) \in \mathcal{RR}^\dagger(\mathbf{A})} z^{N-|I^c|-|J|} \prod_{e \in I^c} x_e \prod_{f \in J} y_f .$$

Now we apply the coordinate transformation $x_e = z$ for all $e \in E$ and $y_f = y$ for all $f \in F$ to obtain

$$q(y, z) = \sum_{k=0}^N RR_k^\dagger(\mathbf{A}) z^{N-k} y^k .$$

The polynomial q is still Lorentzian by [BH20, Theorem 2.10]. Thus, by [BH20, Example 2.26], the sequence of coefficients is ultra log-concave. This proves the first part of the theorem. The second statement is shown the same way using \mathbf{A}^T instead of \mathbf{A} . □

Proof of Theorem C. Let \mathbf{M} and \mathbf{M}' be matroids on ground sets F and F' and consider a map $\phi: F \rightarrow F'$ that defines a morphism from \mathbf{M} to \mathbf{M}' . Write $r := \text{rk}(\mathbf{M})$. When $\phi(F)$ does not span \mathbf{M}' , the set of bases of ϕ is empty and there is nothing to show. So, setting $\mathbf{N} = \phi^* \mathbf{M}'$, it is enough to consider a quotient \mathbf{N} of \mathbf{M} on F . Let $p_{\mathcal{R}(\mathbf{A}[\phi])}(w)$ be the regular minor polynomial

$$p_{\mathcal{R}(\mathbf{A}[\phi])} = \sum_{(I, J) \in \mathcal{R}(\mathbf{A}[\phi])} \prod_{e \in I^c} w_e \cdot \prod_{f \in J} w_f$$

of the bimatroid $\mathbf{A}[\phi]$ constructed in Section 3.3 above. This is Lorentzian by [BH20, Theorem 3.10], since it is the basis generating polynomial of the associated extended matroid. Setting all variables w_e with $e \in E$ equal to x we obtain the weak basis generating polynomial

$$p_{w\mathcal{B}(\phi)}(w) = \sum_{T \in \mathcal{B}(\phi)} \binom{r}{k} x^{|E|-|T|} \prod_{f \in T} w_f$$

of ϕ by Proposition 3.6. This is again Lorentzian by [BH20, Theorem 2.10]. We now set all other variables w_f for $f \in F$ equal to y and obtain the polynomial

$$q(x, y) = \sum_{k \geq 0} \binom{r}{k} B_k(\phi) x^{|E|-k} y^k .$$

This polynomial is also Lorentzian by [BH20, Theorem 2.10] and so the ultra log-concavity of the sequence $\binom{r}{k} \cdot B_k(\phi)$ follows by [BH20, Example 2.26]. But this means that $B_k(\phi)$ itself is log-concave. □

5. VOLUME POLYNOMIALS

Let X be a normal irreducible projective variety of dimension d over an algebraically closed field \mathbb{K} and consider a collection H of nef \mathbb{Q} -divisors H_1, \dots, H_n on X . Then, by [BH20, Theorem 4.6] the *volume polynomial*

$$\mathrm{vol}_H(w) := (w_1 H_1 + \dots + w_n H_n)^d = \sum_{|\alpha|=d} \frac{d!}{\alpha!} (H_1^{\alpha_1} \dots H_n^{\alpha_n}) \cdot w^\alpha$$

is Lorentzian. In general, we say that a Lorentzian polynomial $p(w) \in L_n^d$ is a *volume polynomial*, if there exists a normal irreducible projective variety of dimension d over an algebraically closed field \mathbb{K} as well as a collection H of nef \mathbb{Q} -divisors H_1, \dots, H_n on X such that $\mathrm{vol}_H(w) = p(w)$.

The class of volume polynomials satisfies the following useful properties, which are probably well-known and have been communicated to us by H. Süß.

Proposition 5.1. (i) *Given two volume polynomials $p \in L_n^d$ and $q \in L_{n'}^{d'}$, their product $p \cdot q \in L_{n+n'}^{d+d'}$ is again a volume polynomial.*

(ii) *Given a volume polynomial $p(w) \in L_m^d$ and a matrix $A \in \mathbb{Q}_{\geq 0}^{m \times n}$ with non-negative rational entries, the linear transformation $p(Az) \in L_n^d$ is also a volume polynomial.*

Proof. For Part (i) we are given two normal irreducible projective varieties X and X' as well as two collections $H = (H_1, \dots, H_n)$ and $H' = (H'_1, \dots, H'_{n'})$ of nef \mathbb{Q} -divisors on X and X' respectively. Then we may consider the collection $\tilde{H} = (H_1 \times X', \dots, H_n \times X', X \times H'_1, \dots, X \times H'_{n'})$ of nef \mathbb{Q} -divisors on the product $X \times X'$ and find

$$\mathrm{vol}_{\tilde{H}}(\tilde{w}) = \mathrm{vol}_H(w) \cdot \mathrm{vol}_{H'}(w')$$

in the variables $\tilde{w} = (w, w')$ with $w = (w_1, \dots, w_n)$ and $w' = (w'_1, \dots, w'_{n'})$.

For Part (ii) we consider a normal irreducible projective variety X as well as a collection $H = (H_1, \dots, H_m)$ of nef \mathbb{Q} -divisors on X . For a matrix $A = [a_{ij}]_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} \in \mathbb{Q}_{\geq 0}^{m \times n}$ with non-negative entries, the \mathbb{Q} -divisors $H_j^A = a_{1j} H_1 + \dots + a_{mj} H_m$ are nef for all $1 \leq j \leq n$. We write H^A for the collection of \mathbb{Q} -divisors H_j^A and find

$$\begin{aligned} \mathrm{vol}_H(Aw) &= \left(\left(\sum_{1 \leq j \leq n} a_{1j} w_j \right) H_1 + \dots + \left(\sum_{1 \leq j \leq n} a_{mj} w_j \right) H_m \right)^d \\ &= \left(w_1 \left(\sum_{1 \leq i \leq m} a_{i1} H_i \right) + \dots + w_n \left(\sum_{1 \leq i \leq m} a_{in} H_i \right) \right)^d \\ &= \mathrm{vol}_{H^A}(w) \end{aligned}$$

Therefore, $\mathrm{vol}_H(Aw)$ is again a volume polynomial. □

A central source of volume polynomials are *matroid Schubert varieties*, which have first been studied in [AB16] for realizable matroids and whose geometry motivates the development of the singular Hodge theory of matroids in [BHM⁺23]. We also refer the reader to [EL23] for a generalization of matroid Schubert varieties to the setting of (realizable) discrete polymatroids/ M -convex sets.

Example 5.2. Let K be an algebraically closed field \mathbb{K} . Let M be a realizable matroid of rank r on a ground set E , realized by a set of vectors $\{v_e\}_{e \in E}$ spanning a finite-dimensional \mathbb{K} -vector space V . Without loss of generality, V is spanned by the v_e . Consider the natural surjective linear map $\bigoplus \mathbb{K}v_e \rightarrow V$; its dual is an injective linear map $V^* \rightarrow \prod_{e \in E} \mathbb{K}v_e^*$. The closure of V^* in the multiprojective space $(\mathbb{P}^1)^E$ is called the *matroid Schubert variety* of M and will be denoted by X_M . The hyperplane divisors on every projective line define a collection of ample divisors $H = (H_e)_{e \in E}$ on X . In [AB16, Theorem 1.3 (c)] the

authors show that for every multiset $\{e_1, \dots, e_r\}$ in E with underlying set S we have

$$(H_{e_1} \cdots H_{e_r}) = \begin{cases} 1 & \text{if } S \text{ is a basis of } \mathbf{M} \\ 0 & \text{else.} \end{cases}$$

This implies that for the *basis generating polynomial* $p_{\mathbf{M}}(w) = \sum_{B \in \mathcal{B}(\mathbf{M})} \prod_{b \in B} w_b$ of \mathbf{M} we have

$$p_{\mathbf{M}}(w) = \frac{1}{d!} \text{vol}_H(w).$$

So the basis generating polynomial of a matroid which is realizable over \mathbb{K} is always a volume polynomial.

We are now ready to prove Theorems E and F.

Proof of Theorem E. Let $A \in \mathbb{K}^{E \times F}$ be a matrix. Apply [AB16, Theorem 1.3 (c)] to find that the basis generating polynomial $p_{\hat{\mathbf{A}}}$ of the matroid $\hat{\mathbf{A}}$ associated to the extended matrix $\hat{A} = [I_E | A]$ is a volume polynomial. By Proposition 2.2 this is the same as the regular minor polynomial p_A of the bimatroid \mathbf{A} associated to A . Thus Theorem E is proved. \square

Proof of Theorem F. Let \mathbf{M} and \mathbf{M}' be matroids on ground sets F and F' , realized by vectors $(v_f)_{f \in F}$ and $(v'_f)_{f \in F'}$ spanning vector spaces V and V' respectively. Suppose further that we have a map $\phi: F \rightarrow F'$ and a \mathbb{K} -linear map $\Phi: V \rightarrow V'$ such that $\Phi(v_f) = v'_{\phi(f)}$ for all $f \in F$. Then ϕ defines a morphism of matroids.

When Φ does not surject onto V' , there are no bases of ϕ and there is nothing to show. Hence, we may assume that Φ is surjective.

Denote by U the kernel of Φ and let $\alpha \geq \text{nul}(\phi)$ be an integer. Since \mathbb{K} is infinite, we may generically choose a generating set $\{u_e\}_{e \in E}$ of U with $|E| = \alpha$ such that, given a linearly independent collection of vectors $(v_f)_{f \in T}$ (for $T \subseteq F$), the images $(\Phi(v_f))_{f \in T}$ span V' if and only if for every subset $S \subseteq E$ with $|S^c| + |T| = \dim V$ the set $\{u_e, v_f\}_{e \in S^c, f \in T}$ is a basis of V .

Denote by $\tilde{\mathbf{M}}_\alpha$ the matroid associated to $\{u_e, v_f\}_{e \in E, f \in F}$. We consider the basis generating polynomial

$$p_{\mathcal{B}(\tilde{\mathbf{M}}_\alpha)}(w) = \sum_{B \in \mathcal{B}(\tilde{\mathbf{M}}_\alpha)} \prod_{e \in B \cap E} w_e \cdot \prod_{f \in B \cap F} w_f$$

of $\tilde{\mathbf{M}}_\alpha$. This is a volume polynomial by the construction of matroid Schubert varieties for realizable matroids (i.e. by [AB16, Theorem 1.3 (c)]). Write r for the rank of \mathbf{M} . Setting all variables w_e for $e \in E$ equal to w_0 and noting that $|E| = \alpha$, we find the α -weak basis generating polynomial

$$p_{\mathcal{B}(\phi)}^\alpha(w) = \sum_{k \geq 0} \binom{\alpha}{r-k} w_0^{r-k} \cdot \sum_{T \in \mathcal{B}_k(\phi)} \prod_{f \in T} w_f$$

of the morphism ϕ . This is a volume polynomial by Proposition 5.1 (ii) above. \square

REFERENCES

- [AB16] Federico Ardila and Adam Boocher, *The closure of a linear space in a product of lines*, J. Algebr. Comb. **43** (2016), no. 1, 199–235.
- [AGV21] Nima Anari, Shayan Oveis Gharan, and Cynthia Vinzant, *Log-concave polynomials. I: Entropy and a deterministic approximation algorithm for counting bases of matroids*, Duke Math. J. **170** (2021), no. 16, 3459–3504.
- [AHK17] Karim Adiprasito, June Huh, and Eric Katz, *Hodge theory of matroids*, Notices Am. Math. Soc. **64** (2017), no. 1, 26–30.
- [AHK18] ———, *Hodge theory for combinatorial geometries*, Ann. Math. (2) **188** (2018), no. 2, 381–452.
- [ALGV18] Nima Anari, Kuikui Liu, Shayan Oveis Gharan, and Cynthia Vinzant, *Log-concave polynomials III: Mason’s ultra-log-concavity conjecture for independent sets of matroids*, 2018, arXiv:1811.01600.
- [ALGV19] Nima Anari, Kuikui Liu, Shayan Oveis Gharan, and Cynthia Vinzant, *Log-concave polynomials. II: High-dimensional walks and an FPRAS for counting bases of a matroid*, Proceedings of the 51st annual ACM SIGACT symposium on theory of computing, STOC ’19, Phoenix, AZ, USA, June 23–26, 2019, New York, NY: Association for Computing Machinery (ACM), 2019, pp. 1–12.

- [Bak18] Matthew Baker, *Hodge theory in combinatorics*, Bull. Am. Math. Soc., New Ser. **55** (2018), no. 1, 57–80.
- [BH20] Petter Brändén and June Huh, *Lorentzian polynomials*, Ann. of Math. (2) **192** (2020), no. 3, 821–891.
- [BHM⁺23] Tom Braden, June Huh, Jacob P. Matherne, Nicholas Proudfoot, and Botong Wang, *Singular Hodge theory for combinatorial geometries*, 2023, arXiv:2010.06088.
- [CP24] Swee Hong Chan and Igor Pak, *Log-concave poset inequalities*, 2024, arXiv:2110.10740.
- [EH20] Christopher Eur and June Huh, *Logarithmic concavity for morphisms of matroids*, Adv. Math. **367** (2020), 19, Id/No 107094.
- [EL23] Christopher Eur and Matt Larson, *Intersection theory of polymatroids*, 2023, arXiv:2301.00831.
- [Eur24] Christopher Eur, *Essence of independence: Hodge theory of matroids since June Huh*, Bull. Am. Math. Soc., New Ser. **61** (2024), no. 1, 73–102.
- [FR15] Alex Fink and Felipe Rincón, *Stiefel tropical linear spaces*, J. Comb. Theory, Ser. A **135** (2015), 291–331.
- [GRSU24] Jeffery Giansiracusa, Felipe Rincón, Victoria Schleis, and Martin Ulirsch, *Valuated bimatroids*, 2024, in preparation.
- [HP18] Chris Heunen and Vaia Patta, *The category of matroids*, Appl. Categ. Struct. **26** (2018), no. 2, 205–237.
- [Huh18a] June Huh, *Combinatorial applications of the Hodge-Riemann relations*, Proceedings of the international congress of mathematicians 2018, Volume IV. Invited lectures, Hackensack, NJ: World Scientific; Rio de Janeiro: Sociedade Brasileira de Matemática (SBM), 2018, pp. 3093–3111.
- [Huh18b] ———, *Tropical geometry of matroids*, Current developments in mathematics 2016. Papers based on selected lectures given at the current development mathematics conference, Harvard University, Cambridge, MA, USA, November 2016, Somerville, MA: International Press, 2018, pp. 1–46.
- [Huh22] ———, *Combinatorics and Hodge theory*, Proceedings of the International Congress of Mathematicians, vol. 1, 2022, pp. 212—239.
- [HV19] Chris Heunen and Jamie Vicary, *Categories for quantum theory. An introduction*, Oxf. Grad. Texts Math., vol. 28, Oxford: Oxford University Press, 2019.
- [Kat16] Eric Katz, *Matroid theory for algebraic geometers*, Nonarchimedean and tropical geometry. Based on two Simons symposia, Island of St. John, March 31 – April 6, 2013 and Puerto Rico, February 1–7, 2015, Cham: Springer, 2016, pp. 435–517.
- [Kun78] Joseph P. S. Kung, *Bimatroids and invariants*, Adv. Math. **30** (1978), 238–249.
- [Kun86] ———, *Strong maps*, Theory of matroids, Encycl. Math. Appl. 26, 224–253, 1986.
- [Mur95] Kazuo Murota, *Finding optimal minors of valuated bimatroids*, Appl. Math. Lett. **8** (1995), no. 4, 37–41.
- [Mur03] ———, *Discrete convex analysis*, SIAM Monogr. Discrete Math. Appl., vol. 10, Philadelphia, PA: SIAM Society for Industrial and Applied Mathematics, 2003.
- [Oxl11] James G. Oxley, *Matroid theory*, 2nd ed. ed., Oxf. Grad. Texts Math., vol. 21, Oxford University Press, 2011.
- [RSW23] Julius Ross, Hendrik Süß, and Thomas Wannerer, *Dually Lorentzian polynomials*, 2023, arXiv:2304.08399.
- [Sch79] A. Schrijver, *Matroids and linking systems*, J. Comb. Theory, Ser. B **26** (1979), 349–369.
- [Sta89] Richard P. Stanley, *Log-concave and unimodal sequences in algebra, combinatorics, and geometry*, Graph theory and its applications: East and West. Proceedings of the first China-USA international conference, held in Jinan, China, June 9–20, 1986., New York: New York Academy of Sciences, 1989, pp. 500–535.
- [Wel76] Dominic J. A. Welsh, *Matroid theory*, Lond. Math. Soc. Monogr., vol. 8, Academic Press, London, 1976.
- [Whi86] Neil White (ed.), *Theory of matroids*, Encycl. Math. Appl., vol. 26, Cambridge University Press, Cambridge, 1986.

UNIVERSITÄT TÜBINGEN, FACHBEREICH MATHEMATIK, 72076 TÜBINGEN, GERMANY

Email address: roehrle@math.uni-tuebingen.de

GOETHE-UNIVERSITÄT FRANKFURT, INSTITUT FÜR MATHEMATIK, 60325 FRANKFURT AM MAIN, GERMANY

Email address: ulirsch@math.uni-frankfurt.de